YAP engages RIF1 to dampen replication stress-induced DNA damage in

human squamous cell carcinoma.

Jodie Bojko[1,2], Elodie Sins[3], Benjamin Flynn[1,3], James Scarth[4], Mikal Negasi[5], Bertram

Aschenbrenner[6], Molly Guscott[4], Alexander Howard[1,7], Emily Lay[1,8], Emma Bailey[4], Natalia

Krajic[6], Marco Franciosi[9,10], Sandra Catalán Jiménez[1,11], Kelli Gallacher[1], Ilaria Di Girolamo[3],

Reem Bagabas[3], Caterina Missero[9,10], Jun Wang[4], Sarah McClelland[4], Beate M.

Lichtenberger[6], Ute Jungwirth[1,12], Gernot Walko[1,3,*]

[1]Department of Life Sciences, University of Bath, Bath, UK.

[2]Current affiliation: School of Cellular and Molecular Medicine, University of Bristol, Bristol, UK.

[3]Centre for Oral Immunobiology and Regenerative Medicine, Institute of Dentistry, Queen Mary University of
London, London, UK.

[4]Barts Cancer Institute, Queen Mary University of London, London, UK.

[5]Translational and Clinical Research Institute, Newcastle University, Newcastle, UK.

[6]Department of Dermatology, Medical University of Vienna, Vienna, Austria.

[7]Current affiliation: Astex Pharmaceuticals, Cambridge, UK.

[8]Current affiliation: Centre for Cell Biology and Cutaneous Research, Blizard Institute, Queen Mary University of
London, London, UK.

[9]Department of Biology, University of Naples Federico II, Naples, Italy.

[10]CEINGE – Biotechnologie Avanzate Franco Salvatore S.C.A.R.L., Naples, Italy.

[11]Current affiliation: ONA Therapeutics, Madrid, Spain.

[12]Cancer Research Horizons Therapeutic Innovation, Newcastle Drug Discovery Group, Translational and Clinical
Research Institute, Newcastle University, Newcastle, UK.

[*]Correspondence: g.walko@qmul.ac.uk

Summary

Squamous cell carcinoma cells experience high levels of replication stress due to oncogene-induced cell cycle deregulation. How such cells sustain rapid proliferation despite replication stress is still not fully understood. Here, we discovered, using rapid immunoprecipitation mass spectrometry of endogenous protein (RIME) analysis, that the squamous cell carcinoma oncoprotein YAP engages with RIF1, a key regulator of DNA replication and DNA damage repair under replication stress. RIF1 is highly expressed in human squamous cell carcinoma cell lines and tissues and upregulated during tumour progression. Depletion of RIF1 in squamous cell carcinoma cells exacerbates their endogenous replication stress. Mechanistically, we show that YAP interacts with RIF1 specifically at broken replication forks, and that YAP depletion impairs DNA damage repair under replication stress. Our results thus demonstrate that YAP's oncogenic functions in squamous cancers involve both transcriptional and non-transcriptional mechanisms, the latter through interaction with RIF1 to dampen replication stress.

Keywords

Hippo signalling, YAP, TAZ, RIF1, squamous cell carcinoma, replication stress.

Introduction

Squamous cell carcinomas (SCCs) are among the most common solid tumours [1], with cutaneous SCC (cSCC) being the most prevalent type, imposing a major healthcare burden due to its high incidence [2].

Oncogenic signalling drives cancer cell proliferation by interfering with cell cycle control and upregulation of transcription, thereby causing replication stress, which occurs when DNA

replication fork progression in S phase slows or stalls [3-5]. Ineffective resolution of

replication stress leads to genomic instability or mitotic catastrophes [3, 4]. Cancer cells with

high endogenous replication stress rely on DNA damage repair (DDR) for survival [3]. Of

note, stem and progenitor cells in healthy human squamous epithelia have already high

levels of endogenous replication stress due to their high proliferation rates, and couple

activation of DDR pathways to polyploidisation and terminal differentiation as a homeostatic

mechanism [6-8]. The same mechanism operates to eliminate pre-cancerous cells [7-9]. In

SCC, this protective mechanism is not fully operational, allowing SCC cells to proliferate

despite DNA damage [10].

RIF1 is a key regulator of DNA replication and DDR [11]. By organising higher order

chromatin structures into zones of coordinated DNA replication, and through recruitment of

protein phosphatase 1 (PP1) to prevent MCM (minichromosome maintenance complex)

subunit phosphorylation, RIF1 suppresses late replicating origin firing [12-19]. Under

replication stress, RIF1 has two known functions: it protects stalled replication forks from

resection [20-22], and in DDR interacts with BRCA1 at broken replication forks [23]. While

RIFs role in non-homologous end joining (NHEJ) is well documented [24-27], its role in DNA

replication under replication stress is underexplored.

The transcriptional co-regulators Yes-associated protein (YAP) and transcriptional co-

activator with PDZ-binding motif (TAZ, also called WW Domain Containing Transcription

Regulator 1 (*WWTR1*)) are essential oncoproteins in SCC [28-30]. YAP/TAZ operate as

transcriptional co-activators and co-repressors by indirectly binding to DNA via different

transcription factors (TFs; predominantly those of the TEAD family [28]) [31]. While YAP is

essential in squamous epithelial stem cell proliferation upon injury [32, 33], overexpression

of oncogenic YAP induces rapid senescence [33]. It is likely that this oncogene-induced

senescence mechanism is linked to increased YAP-driven transcription, causing transcription-

replication conflicts and replication stress [34, 35].

How YAP/TAZ control oncogenic signalling in human SCC at the molecular level, and how

human SCC cells deal with oncogene-induced replication stress remains poorly understood.

Using a multi-omics approach we characterized transcriptional and non-transcriptional

mechanisms of YAP's oncogenic role in human SCC. We identified RIF1 as a binding partner

of YAP in human SCC cells, expanding its known interaction in early amphibian development

to human cancers. Importantly, our finding that YAP engages RIF1 under replication stress

and facilitates DDR has translational implications, since replication stress is a crucial

vulnerability of cancer cells that can be exploited therapeutically.

Materials and Methods

Complete details of the materials and methods are provided in the supplemental materials

and methods.

Results

YAP/TAZ-TEAD-mediated oncogenic signalling balances proliferation and differentiation

and drives metastatic progression of human squamous cell carcinoma.

In healthy epidermis, YAP/TAZ balance cell proliferation and terminal differentiation [32, 36].

To test if YAP/TAZ play similar roles in cSCC, we performed RNAseq analysis on SCC13 cells, a

widely used human cSCC cell line [37], following acute siRNA-mediated knockdown of

YAP/TAZ (siYAP/TAZ; Supplementary Fig. 1A, B). While we were unable to significantly

deplete TAZ (*WWTR1*), the TAZ-targeting siRNA pool prevented compensatory upregulation

of TAZ in response to YAP knockdown (Supplementary Fig. 1B) [32]. DESeq2 analysis revealed

a large number of differentially expressed genes (DEGs) upon siYAP/TAZ (977 DEGs

downregulated, 2,209 DEGs upregulated; Supplementary Table 1). Downregulated DEGs

comprised both, genes positively regulated by YAP/TAZ in cooperation with TEADs [38], and

genes regulated by YAP/TAZ in cooperation with the Myb-MuB (MMB) complex [39]

(Supplementary Fig. 1C and Supplementary Table 1). Gene set enrichment analysis (GSEA)

revealed significant enrichment of pathways associated with cell cycle regulation, DNA

replication, and mitosis among the downregulated DEGs, and of pathways associated with

keratinization, cornification, extracellular matrix (ECM) remodelling, and immunoregulation

among the upregulated DEGs (Supplementary Figs. 1D, E, and Supplementary Table 1).

YAP/TAZ knockdown also led to increased terminal differentiation which was evident from (i)

enrichment of an epidermal terminal differentiation gene signature [40] in the upregulated

DEGs and of an epidermal progenitor cell signature [40] in the downregulated DEGs

(Supplementary Fig. 2A), and (ii) upregulation of transcriptional regulators of terminal

differentiation and genes belonging to the epidermal differentiation complex (EDC) [41]

(Supplementary Fig. 2B and Supplementary Table 1).

We next used the YAP/TAZ target gene expression signature derived from our RNA-seq data

to perform gene set variation analysis (GSVA) on one of the largest bulk RNA-seq datasets

from patients with cSCC (237 patients) [42]. This revealed that genes that are positively

regulated by YAP/TAZ in SCC13 cells are expressed significantly higher in cSCC tumours (as

well as their perilesional skin tissue) that have progressed to metastatic disease, compared

to those cSCC tumours that have not (Fig. 1A). Genes that were positively regulated by

YAP/TAZ in SCC13 cells were also highly expressed in cSCC metastases. Conversely, genes

that were negatively regulated by YAP/TAZ in SCC13 cells, including those genes related to

terminal differentiation, displayed significantly reduced expression in cSCCs that had

progressed to metastatic disease, as well as in the respective metastases (Fig. 1B). This

finding provides intriguing evidence in clinical samples for the involvement of YAP/TAZ in the

metastatic progression of human cSCC.

In accordance with the detected transcriptomic changes in siYAP/TAZ SCC13 cells, we

observed that cell proliferation was strongly reduced upon expression of doxycycline-

inducible YAP-targeting shRNAs (shYAP) (Fig. 1C). This was evident from reduced levels of

Ser10-phosphorylated histone H3 (pHH3), a marker of mitotic cells, and strong upregulation

of the CDK inhibitor p21 (Fig. 1D). Consistent with p21 upregulation (Fig. 1D, Supplementary

Fig. 2C) and the reduced expression of several key regulators of DNA replication

(Supplementary Fig. 1C and Supplementary Table 1), most shYAP SCC13 cells arrested in G1,

and a much smaller proportion of cells progressed through S phase (Fig. 1E). Compromised

cell cycle progression and increased terminal differentiation led to reduced 2D clonal growth

(Fig. 1F) and 3D spheroid growth (Fig. 1G) of shYAP SCC13 cells. The effects of shTAZ on

SCC13 proliferation were less strong (Figs. 1D, F). To assess if in cSCC cells YAP/TAZ were

driving oncogenic transcription through TEAD TFs, we employed the potent allosteric pan-

TEAD inhibitor GNE-7883 [43]. GNE-7883 reduced proliferation of SCC13 cells and other

cSCC cell lines in dose-dependent manner (Figs. 2A, B). GNE-7883 treatment of SCC13 cells

phenocopied shYAP by strongly inducing p21 and the terminal differentiation marker

involucrin, and downregulating pHH3 (Figs. 2C, D). GNE-7883 also effectively blocked

proliferation of SCC13 cells growing as 3D spheroids (Figs. 2E, F). Consistent with

CDKN1A/p21 being a major effector of disrupted YAP/TAZ-TEAD signalling, we found that

SCC13 cells expressing *CDKN1A*-targeting shRNAs were able to proliferate normally upon

GNE-7883 treatment (Figs. 2G, H). We conclude that YAP/TAZ-TEAD regulate the balance

between proliferation and terminal differentiation, thereby promoting a highly proliferative

and undifferentiated cell state that is relevant for disease progression. This important

function is conserved between normal epidermal keratinocytes [32, 36] and neoplastic

keratinocytes of both BCC [44] and cSCC (this study).

YAP/TAZ-TEAD-mediated oncogenic signalling is major contributor to replication stress in

squamous cell carcinoma cells.

Reduced activation of the replication stress checkpoint (phosphorylation of CHK1 on Ser317;

[45, 46]) was evident in SCC13 cells upon shYAP but not shTAZ knockdown (Supplementary

Fig. 3A), while depletion of either YAP or TAZ decreased the levels of γH2AX. In the context

of replication stress, γH2AX marks accumulation of DNA double strand breaks (DSBs) at

collapsed replication forks [6, 23, 46] (Supplementary Fig. 3A). Furthermore, using mRNA

expression data from the DepMap project [47], we discovered a strong correlation between

the expression of a replication stress signature [48] and an SCC-specific YAP/TAZ activity

signature comprising 10 highly-correlated YAP/TAZ target genes from our RNAseq dataset in

HNSCC (Supplementary Table 1, Supplementary Figs. 4A–D). To provide additional evidence

for a causal link between oncogenic YAP signalling and replication stress, we overexpressed

wildtype YAP and hyperactive YAP5SA and YAPS127A mutants [33] in SCC cells. As shown in

Supplementary Fig. 3B, all three YAP transgenes increased YAP's transcriptional output, as

evident from strongly elevated expression of the canonical YAP/TAZ-TEAD target gene *CCN1*,

and notably reduced cell proliferation when compared to empty vector control

(Supplementary Fig. 3C). Both wildtype YAP and YAPS127A enhanced replication stress, as

evident from increased γH2AX expression and increased CHK1 Ser345 phosphorylation [45,

46], while YAP5SA overexpression caused increased p21 expression, probably indicating

oncogene-induced senescence (Supplementary Fig. 3B).

To demonstrate that transcription can be a general contributor to replication stress in SCC

cells, we transiently inhibited RNA synthesis in SCC13 cells using triptolide before assessing

γH2AX levels [49]. Triptolide treatment reduced RNA synthesis (Supplementary Fig. 3D) and

γH2AX expression to a similar extend as TEAD inhibition by GNE-7883 (Supplementary Figs.

3E, F). Together, these findings showed that YAP/TAZ-TEAD-driven oncogenic signalling is a

major contributor to the endogenous replication stress of SCC cells.

A fraction of chromatin-associated YAP interacts with RIF1.

To   understand   how   YAP   executes   its   transcriptional   roles,   we   performed   rapid

immunoprecipitation mass spectrometry of endogenous protein (RIME) analysis [50] in

SCC13 cells and normal human epidermal keratinocytes (NHK) (Fig. 3), using a knockdown-

validated YAP-specific antibody that does not recognise TAZ (Supplementary Figs. 5A, B, G).

YAP immunoprecipitation (IP) efficiency was consistently high across all YAP-RIME replicates,

with   high   YAP   peptide   coverage   (63-65%)   detected   by   mass   spectrometry   analysis

(Supplementary Fig. 5C and Supplementary Table 2). Upon stringent filtering of the mass

spectrometry   data   (see   Supplementary   Materials   and   Methods)   YAP-RIME   enabled

identification of 47 and 14 high confidence YAP-interacting proteins in SCC13 cells (Fig. 3A,

Supplementary Table 2) and NHK cells (Supplementary Fig. 5D and Supplementary Table 2),

respectively.   Significantly   enriched   gene   ontology   terms   were   all   related   to   processes

occurring in the nucleus in both interactomes (Fig. 3B, Supplementary Fig. 5E). Among the

SCC13 YAP-interacting proteins were several known regulators of YAP, including Neurofibromatosis type (NF2) [51], Angiomotin-like 1 (AMOTL1) [52], PPP1CB [53], and Prolyl 4-Hydroxylase subunit alpha 2 (P4HA2) [54] (Fig. 3A). Our YAP-RIME also identified several subunits of the BAF-type SWI/SNF chromatin remodelling complex (ARID1A, SMARCA4, SMARCC1, SMARCE1, SMARCD2) [55], and key components of the NuRD chromatin remodelling complex, Chromodomain-helicase-DNA-binding protein 4 (CHD4) and Metastasis Associated 1 Family Member 2 (MTA2). While key components of the general transcriptional machinery (POLR2A, POLR2B, GTF21, TOP1) were present in the SCC13 YAP interactome, TEAD TFs were notably absent (Fig. 3A). A closer inspection of our mass spectrometry data revealed that TEADs, as well as other known YAP-interacting transcription factors, such as JunB and TP63 [56, 57], had been identified but did not pass our stringent bioinformatics filtering criteria (Supplementary Table 2). The overall poor identification of TFs by our YAP-RIME was likely due to their insufficient crosslinking to the DNA by formaldehyde alone, as previously observed for the highly dynamic transcription factor NFκB [58].

To identify physiologically relevant interactions, the YAP-interacting proteins were compared to candidate positive and negative regulators of cell proliferation and survival identified in a prior genome-wide RNA screens in SCC13 [32]. This revealed that 22 out of the 47 high confidence YAP-interacting proteins were likely relevant for cell proliferation and/or survival (Fig. 3A, Supplementary Table 2).

Our attention was drawn to RIF1, given its recent identification as a non-transcriptional YAP interaction partner in a developmental context in *Xenopus laevis* [59]. We detected RIF1 as a YAP-chromatin-interacting protein in both the SCC13 and NHK YAP-RIME experiments (Fig.

3A, Supplementary Fig. 5D) with a much stronger enrichment in SCC13 YAP (Supplementary

Fig. 5F and Supplementary Table 2). Of note, PPP1CB, one of the three catalytic subunits of

PP1—a key effector of RIF1 [13, 16, 17, 19], was also present in the SCC13 YAP-RIME (Fig. 3A,

Supplementary Table 2). Like YAP, RIF1 featured among the positive regulators of cell

proliferation and survival of SCC13 cells identified in the aforementioned genome-wide RNA

interference screen [32] (Fig. 3A, Supplementary Table 2). As expected, YAP and RIF1 were

both localised in the nucleus of SCC13 cells (Fig. 3C) and present in chromatin fractions

(Supplementary Fig. 5G), confirming their chromatin-association. To biochemically validate

the YAP-RIF1 interaction, co-IP experiments were performed under native non-cross-linking

conditions using a different YAP-specific antibody (Supplementary Fig. 6A). Indeed, RIF1

could be immunoprecipitated together with YAP from both, whole-cell detergent extracts

and nuclei-enriched fractions (Figs. 3D, E). Further validating our RIME results, CHD4 and

ARID1A also co-immunoprecipitated with YAP under native conditions. Under these

conditions, TEAD transcription factors could also be immunoprecipitated together with YAP

(Fig. 3E). In-situ proximity ligation assays (PLA) confirmed very close YAP-RIF1 proximity in

the nucleus of SCC13 cells (Fig. 3F). To achieve a sufficiently strong knockdown of both YAP

and TAZ to control for PLA signal specificity, we combined our doxycycline-inducible YAP-

targeting shRNAs with an siRNA pool targeting YAP and TAZ (Fig. 3F, Supplementary Fig. 6B).

The YAP-RIF1 PLA signal was significantly decreased upon YAP/TAZ knockdown. Consistent

with our co-IP results, YAP was also found in very close proximity to TEAD1 in the nucleus of

SCC13 cells by PLA (Fig. 3G). The YAP-RIF1 PLA signal was completely absent when the two

primary antibodies were omitted (Supplementary Figs. 6C, D) and also largely absent from

mitotic chromosomes (Supplementary Fig. 6E). To test if the YAP-RIF1 interaction also

occurred in other types of SCC, we used the lung SCC (LUSC) cell line HARA. Co-IP was

performed with nuclei-enriched HARA cell fractions treated with Benzonase nuclease to

enzymatically digest DNA and RNA (Supplementary Figs. 6F, G), demonstrating that YAP and

RIF1 are not co-immunoprecipitated because of a DNA bridge between the two proteins. In

summary, our results show that RIF1 is a binding partner of chromatin-associated YAP in

both healthy human NHKs and cancerous cSCC and LUSC cells.

YAP/TAZ-TEAD modulate RIF1 protein levels.

Our RNAseq analysis revealed that *RIF1* mRNA expression was moderately downregulated in

response to siYAP/TAZ (Supplementary Fig. 1A and Supplementary Table 1). A similar, albeit

statistically not significant trend could be confirmed in independent siYAP/TAZ samples by

RT-qPCR analysis (Fig. 4A). These mild effects of YAP depletion on *RIF1* mRNA expression are

likely indirect and may be linked to cell cycle de-regulation. Depletion of RIF1 did not impact

on expression of most of the DEGs seen upon siYAP/TAZ (Supplementary Figs. 1B, C, and 2B,

C), ruling out a direct involvement in YAP/TAZ-TEAD-controlled transcription.

The stability of a protein within a complex often depends on other components of the

complex, as proper folding of one protein may require interactions with its partners. RIF1

protein expression was lower in siYAP/TAZ compared to siNTC SCC13 cells (Fig. 4B).

Conversely, overexpression of wildtype and oncogenic YAP variants increased RIF1 protein

levels irrespective of their effects on the cell cycle (pHH3) (Fig. 4C). Interestingly, short-term

treatment (24 h) with GNE-7883 to disrupt YAP/TAZ-TEAD binding reduced RIF1 protein

expression without affecting *RIF1* mRNA levels (Figs. 4D, E). This suggests that YAP might

modulate RIF1 protein levels in a complex with TEADs. Indeed, combined siRNA-mediated

knockdown of TEAD1, 3, and 4 (siTEAD1/3/4) in SCC13 cells reduced RIF1 protein to levels

similar to those observed in siYAP/TAZ cells (Figs. 4F, G). Interestingly, depletion of YAP

impacted chromatin-associated RIF1 levels specifically during S phase (Figs. 4H, I). Since the

data presented in Figs. 3E–G showed that YAP is in complex with RIF1 and TEADs, these

findings thus suggest that YAP-TEAD modulate protein levels of a fraction of chromatin-

associated RIF1 in S phase.

RIF1 is highly expressed in squamous cell carcinoma and upregulated during tumour

progression.

To understand if RIF1 contributes to squamous carcinogenesis, we assessed RIF1 protein

expression in a panel of human cSCC cell lines, which also included an isogenic cSCC

progression series (PM1, MET1–4), including the pre-malignant PM1 precursor lesion [60].

Increased RIF1 protein levels were notable in all cSCC cell lines when compared to NHKs (Fig.

5A). Consistent with our findings that YAP-TEAD modulate RIF1 protein levels (Fig. 4), RIF1

protein expression was highest in SCC13 cells featuring high nuclear YAP levels [32], as also

indicated by a low ratio of S127-YAP to total YAP (Fig. 5A). All cSCC lines displayed higher

levels of γH2AX compared to NHKs, indicating increased endogenous replication stress (Fig.

5A).

To investigate if high RIF1 expression was a general hallmark of cancers originating from

epidermal keratinocytes, we quantified the levels of RIF1 protein in a tissue array

comprising cSCCs of different differentiation grades, as well as different types of BCCs (Figs.

5B, C). We detected increased numbers of RIF1-expressing cells within the keratin-positive

tumour cell masses of both BCC and cSCC compared to normal skin epidermis, especially

within the more aggressive tumour types of infiltrative BCC and poorly differentiated cSCC

(Figs. 5B, C). These data are corroborated by immunohistochemistry data from the Human

Protein Atlas, depicting increased numbers of RIF1-expressing tumour cells in cSCC, HNSCC

and cervical SCC (Supplementary Figs. 7A–C). To obtain a pan-cancer overview of RIF1

expression, we used the UALCAN portal [61] to access protein expression data from the

Clinical Proteomic Tumour Analysis Consortium (CPTAC) [62]. In HNSCC, RIF1 protein

expression increased with tumour grade (Fig. 5D) and RIF1 expression was higher across

several cancer types compared to normal tissue (Supplementary Fig. 7D). The UALCAN

portal also allows assessment of the impact of somatic alterations (small mutations or copy

number alterations) in signalling pathways on the cancer proteome [61, 62]. We found that

in both HNSCC (Fig. 5E) and LUSC (Supplementary Fig. 7E) increased RIF1 protein expression

predominantly correlated with somatic Hippo pathway alterations, in contrast to alterations

in other signalling pathways, implying altered Hippo pathway activity in the regulation of

RIF1 expression. In contrast, in other cancer types, alterations in pathways other than Hippo

signalling had a strong impact on increased tumour expression of RIF1 (Supplementary Figs.

7F, G). Collectively, our multi-omics analysis confirm a close relationship between Hippo

signalling and RIF1 protein expression and point towards a role of RIF1 in squamous cell

carcinogenesis.

Depletion of RIF1 increases replication fork speed in squamous cell carcinoma cells.

Consistent with a role of RIF1 in controlling S phase progression of SCC cells, we observed

that on release from a double thymidine block, shRIF1 SCC13 cells progressed more rapidly

through S phase (Fig. 6A). Phosphorylation levels of MCM2 and MCM4 proteins, targets of

the Dbf4-dependent kinase (DDK; composed of CDC7 kinase and its activator, DBF4) [16,

18], were increased in RIF1-depleted SCC13 cells in a RIF1 dosage-dependent manner (Figs.

6B, C), particularly in early and mid S phase (2- and 4-h time points post-dTb, Fig. 6C),

indicating alterations in the dynamics of origin firing [16-19]. Accordingly, chromatin loading

of PCNA, which is required for DNA replication after the action of CDC7, was also increased

during early and mid S phase of shRIF1 SCC13 cells (Fig. 6C). Therefore, the faster

progression of SCC13 cells through S phase after RIF1 depletion likely reflects an increase in

DNA replication initiation events in early and mid S phase [16, 18]. We also examined DNA

replication dynamics in shRIF1 cells using a novel long-read sequencing method, DNAscent

[63]. We discovered that RIF knockdown increased fork speed along with led to longer

replication tracks (Figs. 6D, E, and Supplementary Figs. 8A, B).

RIF1 protects against endogenous replication stress in squamous cell carcinoma cells.

Under replication stress, RIF1 protects stalled replication forks [16, 20-22], but also

facilitates homology-mediated DDR at broken replication forks [23]. The elevated expression

of RIF1 in SCC cells could therefore indicate a role for RIF1 in enabling the cells to deal with

high levels of endogenous replication stress that are, at least in part, controlled by YAP-

TEAD-driven oncogenic mechanisms (Fig. 2). Indeed, we found a strong correlation between

RIF1 expression and a replication stress signature [48] in HNSCC cell lines (Supplementary

Figs. 4A, E). In contrast to a slowing down or stalling of replication fork progression, which is

frequently observed when cells experience replication stress [3-5], our DNAscent analysis

revealed increased replication fork speed upon RIF knockdown (Figs. 6D, E). Moreover,

replication fork stalling or pausing, as defined by a shift of DNAscent 'stall score' distribution

closer to 1.0, was not detected (Fig. 6F). Previous studies have found that increased

replication speed can lead to DNA damage, whereby faster replication forks have

insufficient time to recognize damaged DNA in need of repair [64, 65]. The only moderate

activation of CHK1 (phosphorylation on Ser317 and S345; [45, 46]) and very mild increase of

ATR-mediated S33-phosphorylation of RPA32 [46] in shRIF1 SCC13 cells, in contrast to the

strongly elevated levels of γH2AX and hyperphosphorylated pS4/8-RPA32 (indicating

phosphorylation by ATM and DNA-PK in response to generation of DNA DSBs [46, 66, 67]

(Figs. 7A, B) are corroborating such a scenario. Indeed, replication fork collapse and

concomitant activation of ATM/DNA-PK signalling were previously observed in response to

ATR inhibition in cells experiencing hydroxyurea (HU)-induced replication stress [67]. It is

worth noting that DNAScent only detects short (<20 min) replication fork stalls/pauses and

currently cannot measure longer fork stalls and fork breakage. Given that RIF1 facilitates

repair of broken replication forks [23] it is likely that loss of this additional function also

contributes to the increased accumulation of DNA damage in shRIF1 cells. Because the

replication checkpoint was not robustly activated in shRIF1 SCC13 cells (Figs. 7A, B), they

were able to progress through the cell cycle despite persistent DNA damage (Figs. 7C, D). A

small increase in the G2/M fraction in shRIF1 SCC13 cells (Figs. 7C, D) was noticeable as well

as a slightly elevated pHH3 expression in siRIF1 cells (Fig. 7E). The accumulation of DNA

damage in cultures of shRIF1 SCC13 cells became more accentuated after a few passages,

when cells had undergone multiple rounds of cell division (Fig. 7F), ultimately resulting in

reduced clonal growth (Figs. 7G, H). To confirm that the replication stress-protecting effect

of RIF1 is conserved in other types of SCC, we depleted RIF1 in the HNSCC cell line SAS,

which displays a strong correlation between RIF1 expression and replication stress

(Supplementary Fig. 4E) and carries a *YAP-MAML2* gene fusion resulting in a hyperactive

YAP variant [68]. Indeed, we observed a dosage-dependent effect of shRIF1 on cell viability

(Supplementary Figs. 9A–C) and DNA damage (Supplementary Figs. 9D, E). Decreased cell

viability (Supplementary Fig. 9F) and increased γH2AX levels (Supplementary Fig. 9G) were

also detected in the LUSC cell line HARA upon shRIF1. Together, our results point to a

scenario where SCC cells depend on elevated RIF1 expression to dampen their high levels of

endogenous replication stress.

YAP is involved in DNA damage repair at collapsed replication forks.

Our discovery of RIF1 as a YAP interaction partner in human SCC cells raised the exciting

possibility that in this cancer context, YAP could play dual roles in both transcription and

DNA replication under replication stress. To test, if YAP, like RIF1, might function at

collapsed replication forks [23], we treated SCC13 cells with 4 mM HU [23, 69] for 24 h. This

triggered strong activation of the replication stress checkpoint (phosphorylation of CHK1 on

Ser317; [45, 46]) and accumulation of DNA DSBs, marked by strongly elevated γH2AX

expression (Fig. 8A). Persisting γH2AX expression following HU washout indicated

accumulation of DNA damage at stalled and subsequently inactivated replication forks [69]

(Fig. 8A). This was further confirmed by the presence of RAD51 foci [23] (Fig. 8B) and

appearance of pan-nuclear γH2AX staining, indicating severe DNA replication stress [70],

upon 24 h HU treatment (Supplementary Fig. 10A). Indeed, some cell death (appearance of

sub-G1 fraction in EdU/PI cell cycle profiles, Supplementary Fig. 10A) was noticeable from 15

h HU treatment onwards. Interestingly, YAP protein levels were increased following washout

of HU after 24 h of treatment, suggesting a potential role of YAP in DDR upon replication

fork collapse (Fig. 8C). Indeed, while the YAP-RIF1 PLA signal remained unchanged upon

short-term (3 h) HU treatment, it increased significantly after long-term (24 h) HU treatment

(Fig. 8D). Moreover, upon long-term HU treatment, more YAP molecules were found in

close vicinity of γH2AX by in-situ PLA (Fig. 8E, Supplementary Fig. 10C), suggesting that YAP

associates with RIF1 preferentially near DNA DSBs at broken replication forks. To test if YAP

is directly involved in DDR at broken replication forks, we optimized conditions for RNA

interference experiments to only minimally perturb the cell cycle in response to siYAP/TAZ

(Fig. 8F, Supplementary Fig. 10D). For these experiments, HU concentration was reduced to

2 mM to avoid potential confounding effects of HU-induced cell death. As shown in Fig. 8F,

by 48 h post-transfection with a low concentration of siYAP/TAZ (1 nM), SCC13 cells

displayed reduced levels of YAP and only modest upregulation of p21 and downregulation of

pHH3, while γH2AX levels (as a read-out of YAP-TEAD-driven replication stress) were

strongly reduced. Treatment with 2 mM HU during the last 24 h post-siYAP/TAZ transfection

led to prominent S phase arrest, as evident from strongly reduced pHH3 levels and

replication stress checkpoint activation (phosphorylation of CHK1 on Ser345) (Fig. 8F).

Under these conditions, YAP depletion did not cause a decrease in γH2AX levels but instead

led to a significant increase in comparison to untreated siYAP/TAZ cells (Fig. 8F). Since

transcription is globally dampened under high concentrations of HU [71] and YAP was

depleted specifically during the HU treatment period, these findings suggest a novel non-

transcriptional function of YAP where it is involved in DDR at broken replication forks.

Given the dual function of YAP in promoting TEAD-mediated transcription and facilitating

DDR under severe replication stress, we examined if pharmacologically induced replication

stress could act synergistically with YAP depletion to impair SCC proliferation. Of note, HU is

sometimes used in chemoradiotherapy for HNSCC patients [72]. Indeed, we observed a

synergistic effect of shYAP and low doses of HU on SCC13 clonal colony growth (Figs. 8G, H).

This finding highlights the possibility of using pharmacological replication stress inducers in

combination with YAP-TEAD inhibitors [73] to treat SCC.

Discussion

Our findings provide novel insight into the oncogenic mechanisms of hyperactive YAP/TAZ in

SCC. While YAP/TAZ promote proliferation by regulating transcriptional programmes that

drive cell cycle progression, they also suppress commitment to terminal differentiation.

Thus, the functions of YAP/TAZ are highly conserved within the epidermal lineage, as both

normal epidermal stem cells [32, 33, 36] and neoplastic keratinocytes in BCC [44] and cSCC

(this study) show strong dependency on YAP/TAZ-TEAD signalling for cell proliferation. This

suggests that cancers originating from epidermal stem/progenitor cells 'hijack' the tissue

regenerating functions of YAP/TAZ to fuel their high rates of proliferation while remaining in

an undifferentiated cell state. The increased expression of genes that are positively

regulated by YAP/TAZ in tumour tissue of human cSCCs that underwent metastatic

progression (Fig. 1) provides strong support for this idea. Our RIME analysis revealed that

the transcriptional co-regulator functions of YAP/TAZ are likely mediated by the BAF-type

SWI/SNF and the NuRD chromatin remodelling complexes. Indeed, in HNSCC cells, BAF-type

SWI/SNF and YAP-TEAD complexes were shown to be co-dependent for chromatin binding,

whereby they create a positive-feedback mechanism to maintain open chromatin for

transcription [74]. In contrast, recruitment of the NuRD complex [75] could be involved in

the observed co-repressor functions of YAP/TAZ.

In a previous study which employed a transgenic mouse model of cSCC development,

simultaneous activation of oncogenic Kras^{G12D} and knock-down of YAP/TAZ in epidermal

stem cells was found to induce apoptosis [76]. In contrast, we found that YAP/TAZ depletion

in human SCC has cytostatic- and differentiation-promoting, rather than cytotoxic effects,

which appear to be mediated by induction of p21. It is worth noting that activating

mutations in the *RAS* proto-oncogenes are not very frequent in human cSCC [2], with some

tumours (~10%) showing copy number loss of *HRAS* [77]. Induction of apoptosis in response

to YAP/TAZ depletion might thus be specific to SCCs expressing oncogenic RAS [76].

We show that a key function of RIF1 in human squamous cancers is to regulate DNA

replication dynamics. Our findings in SCC13 cells show that in the absence of RIF1, origin

firing and replication fork speed are increased (Fig. 6). However, our DNAscent analysis

revealed that the rate of replication fork stalling/pausing in SCC13 cells was not increased by

RIF1 depletion. A previous study found that replication stress can be caused by increased

fork speed and without fork stalling when the average fork velocity increases above a ~40%

gain [64]. Although the average increase in fork speed in shRIF1 SCC13 cells was only ~25%,

we believe that this increased fork speed can generate increasing levels of replication stress

and DNA damage over a few cell division cycles, as was previously shown for cells with

elevated E2F-driven transcription [78].

Since origin firing occurs preferentially within open chromatin regions that contain

transcriptional-regulatory elements, some of which were shown to be enriched for TEAD TF

binding motifs [79], modulation of RIF1 chromatin association by YAP-TEAD (Fig. 4) could

serve to coordinate YAP-TEAD-driven transcription with DNA replication. Unfortunately,

because of the confounding effects of YAP depletion on the cell cycle (Fig. 1), it was not

possible for us to assess if YAP was involved in DNA replication under unperturbed

conditions.

An important discovery of our study was a specific role of YAP in facilitating DDR at

deactivated replication forks under severe replication stress. How YAP might work together

with RIF1 to promote DDR at broken replication forks remains elusive at this point,

particularly since the mechanism by which RIF1 enables BRCA1/RAD51-dependent repair of

broken replication forks it yet not well understood [23]. Notably, this DDR function is

attributed specifically to the long isoform of RIF1 (RIF1-long), while control of replication

dynamics is mediated by both RIF1-long and RIF1-short isoforms [80].

Our findings thus provide evidence for a non-canonical, non-transcriptional role of YAP in

SCC, whereby it engages RIF1 to dampen the high levels of endogenous replication stress in

these cancers, which are—at least in part—caused by YAP's oncogenic functions itself.

Induction of replication stress together with pharmacological YAP/TAZ-TEAD inhibition [73]

could thus potentially be used as a combination therapy strategy to treat SCC.

Funding information

This work was supported by the UK Research & Innovation (UKRI) Biotechnology and

Biological Sciences Research Council (BBSRC, grant BB/T012978/1 to Gernot Walko), the UK

Research & Innovation (UKRI) Medical Research Council (MRC, grant MR/W031442/1 to

Sarah McClelland), the Academy of Medical Sciences (grant SBF005\1005 to Gernot Walko),

the British Skin Foundation (grants 004/SG/18, 036/S/18 ('Peter Massey Skin Cancer Grant')

and 011_SG_22 to Gernot Walko), the Royal Society (grant RGS\R1\191087 to Gernot

Walko), the LEO Foundation (grant LF-AW_EMEA-21-400116 to Beate Lichtenberger), the

City of Vienna Fund for Innovative Interdisciplinary Cancer Research (grant 22059 to Beate

Lichtenberger), the Austrian Science Fund (FWF) (grant 36368-B to Beate Lichtenberger),

Cancer Research UK (CRUK, grant DRCPFA-Jun24/100005 to Sarah McClelland), and the

Italian Association for Cancer Research (AIRC, grant IG25116 to Caterina Missero). Ute

Jungwirth gratefully acknowledges funding from Cancer Research UK (Newcastle Drug

Discovery Unit Programme Grant DRCDDRPGMApr2020\100002). Gernot Walko, Jun Wang,

Elodie Sins, Benjamin Flynn, and Emma Bailey also acknowledge funding support from the

Barts Charity (UK) Strategic Award to the Barts Centre for Squamous Cancer (Centre grant 'A

Centre of Excellence for Squamous Cancer', grant G-002030). Jun Wang and Emma Bailey

further acknowledge funding support from the Cancer Research UK City of London Major

Centre Award core funding support to the Barts Cancer Institute. Jodie Bojko was recipient

of a Raoul & Catherine Hughes Scholarship through the University of Bath 'Alumni and

Friends' network. Reem Bagabas was recipient of a PhD studentship from the Saudi Arabian

Cultural Bureau.

481   Acknowledgements

We would like to thank all members of the Walko and Jungwirth labs for their inputs. We

wish to thank our colleagues from the Barts Centre for Squamous Cancer at Queen Mary

University of London for critical feedback on our draft manuscript. We thank Genomics

Birmingham Genomics Service at the University of Birmingham, for the generation of the

DNAscent data. We also acknowledge support from the bioimaging facility of the Barts

Cancer Institute (Queen Mary University of London).

489   Author contributions

Jodie Bojko: Data Curation, Formal Analysis, Investigation, Methodology, Visualization,

Writing: Original Draft Preparation; Writing: Review & Editing; Elodie Sins: Data Curation,

Investigation, Methodology; Benjamin Flynn: Data Curation, Formal Analysis, Investigation,

Methodology, Software, Validation, Visualization; James Scarth: Data Curation, Formal

Analysis, Investigation, Methodology, Visualization; Mikal Nigasi: Data Curation, Formal

Analysis, Investigation, Methodology, Visualization; Bertram Aschenbrenner: Data Curation,

Formal Analysis, Investigation, Methodology, Software, Visualization; Molly Guscott: Formal

Analysis, Investigation, Methodology, Software; <u>Alexander Howard</u>: Formal Analysis,

Investigation, Methodology; <u>Emily Lay</u>: Formal Analysis, Investigation, Methodology; <u>Emma</u>

<u>Bailey</u>: Formal Analysis, Investigation, Methodology, Software, Visualization; <u>Natalia Krajic</u>**:**

Investigation, Methodology; <u>Marco Franciosi</u>: Investigation, Methodology; <u>Sandra Catalan</u>:

Investigation, Methodology; <u>Kelli Gallacher</u>: Investigation, Methodology; <u>Ilaria di Girolamo</u>:

Data Curation, Investigation; <u>Reem Bagabas</u>: Data Curation, Investigation; <u>Caterina Missero</u>:

Funding acquisition, Resources, Supervision, Validation; <u>Jun Wang</u>: Funding acquisition,

Methodology, Resources, Supervision; <u>Sarah McClelland</u>: Funding acquisition, Resources,

Supervision, Validation; <u>Beate Lichtenberger</u>: Funding acquisition, Resources, Supervision,

Validation, Writing: Original Draft Preparation; <u>Ute Jungwirth</u>: Formal Analysis, Funding

acquisition, Resources, Supervision, Validation, Visualization, Writing: Original Draft

Preparation, Writing: Review & Editing; <u>Gernot Walko</u>: Conceptualization, Data Curation,

Formal Analysis, Funding acquisition, Methodology, Project Administration, Resources,

Supervision, Validation, Visualization, Writing: Original Draft Preparation, Writing: Review

&Editing.

Data availability

The authors declare that all data supporting the findings of this study are available within

the paper and its supplementary information files. Raw Illumina sequencing data have been

deposited in the Gene Expression Omnibus (GEO) following the MIAME guidelines, under

the accession number GSE284725. The mass spectrometry proteomics data have been

deposited to the ProteomeXchange Consortium via the PRIDE partner repository with the

dataset identifier PXD061022 and 10.6019/PXD061022. DNAscent data have been deposited

on European Nucleotide Archive (ENA), accession ID PRJEB103830.

References

Dotto GP, Rustgi AK. Squamous Cell Cancers: A Unified Perspective on Biology and Genetics. *Cancer Cell* 2016; 29: 622-637.

Winge MCG, Kellman LN, Guo K, Tang JY, Swetter SM, Aasi SZ *et al.* Advances in cutaneous squamous cell carcinoma. *Nat Rev Cancer* 2023; 23: 430-449.

da Costa A, Chowdhury D, Shapiro GI, D'Andrea AD, Konstantinopoulos PA. Targeting replication stress in cancer therapy. *Nat Rev Drug Discov* 2023; 22: 38-58.

Gaillard H, Garcia-Muse T, Aguilera A. Replication stress and cancer. *Nat Rev Cancer* 2015; 15: 276-289.

Bowry A, Kelly RDW, Petermann E. Hypertranscription and replication stress in cancer. *Trends Cancer* 2021; 7: 863-877.

Molinuevo R, Freije A, Contreras L, Sanz JR, Gandarillas A. The DNA damage response links human squamous proliferation with differentiation. *J Cell Biol* 2020; 219.

Freije A, Ceballos L, Coisy M, Barnes L, Rosa M, De Diego E *et al.* Cyclin E drives human keratinocyte growth into differentiation. *Oncogene* 2012; 31: 5180-5192.

Freije A, Molinuevo R, Ceballos L, Cagigas M, Alonso-Lecue P, Rodriguez R *et al.* Inactivation of p53 in Human Keratinocytes Leads to Squamous Differentiation and Shedding via Replication Stress and Mitotic Slippage. *Cell Rep* 2014; 9: 1349-1360.

Gandarillas A, Davies D, Blanchard JM. Normal and c-Myc-promoted human keratinocyte differentiation both occur via a novel cell cycle involving cellular growth and endoreplication. *Oncogene* 2000; 19: 3278-3289.

Alonso-Lecue P, de Pedro I, Coulon V, Molinuevo R, Lorz C, Segrelles C *et al.* Inefficient differentiation response to cell cycle stress leads to genomic instability and malignant progression of squamous carcinoma cells. *Cell Death Dis* 2017; 8: e2901.

Blasiak J, Szczepanska J, Sobczuk A, Fila M, Pawlowska E. RIF1 Links Replication Timing with Fork Reactivation and DNA Double-Strand Break Repair. *Int J Mol Sci* 2021; 22.

Foti R, Gnan S, Cornacchia D, Dileep V, Bulut-Karslioglu A, Diehl S *et al.* Nuclear Architecture Organized by Rif1 Underpins the Replication-Timing Program. *Mol Cell* 2016; 61: 260-273.

13 Gnan S, Flyamer IM, Klein KN, Castelli E, Rapp A, Maiser A *et al*. Nuclear
organisation and replication timing are coupled through RIF1-PP1 interaction. *Nat*
Commun 2021; 12: 2910.

14 Moriyama K, Yoshizawa-Sugata N, Masai H. Oligomer formation and G-quadruplex
binding by purified murine Rif1 protein, a key organizer of higher-order chromatin
architecture. *J Biol Chem* 2018; 293: 3607-3624.

15 Cornacchia D, Dileep V, Quivy JP, Foti R, Tili F, Santarella-Mellwig R *et al*. Mouse
Rif1 is a key regulator of the replication-timing programme in mammalian cells.
EMBO J 2012; 31: 3678-3690.

16 Alver RC, Chadha GS, Gillespie PJ, Blow JJ. Reversal of DDK-Mediated MCM
Phosphorylation by Rif1-PP1 Regulates Replication Initiation and Replisome Stability
Independently of ATR/Chk1. *Cell Rep* 2017; 18: 2508-2520.

17 Moiseeva TN, Yin Y, Calderon MJ, Qian C, Schamus-Haynes S, Sugitani N *et al*. An
ATR and CHK1 kinase signaling mechanism that limits origin firing during
unperturbed DNA replication. *Proc Natl Acad Sci U S A* 2019; 116: 13374-13383.

18 Yamazaki S, Ishii A, Kanoh Y, Oda M, Nishito Y, Masai H. Rif1 regulates the
replication timing domains on the human genome. *EMBO J* 2012; 31: 3667-3677.

19 Hiraga SI, Ly T, Garzon J, Horejsi Z, Ohkubo YN, Endo A *et al*. Human RIF1 and
protein phosphatase 1 stimulate DNA replication origin licensing but suppress origin
activation. *EMBO Rep* 2017; 18: 403-419.

20 Buonomo SB, Wu Y, Ferguson D, de Lange T. Mammalian Rif1 contributes to
replication stress survival and homology-directed repair. *J Cell Biol* 2009; 187: 385-
398.

21 Mukherjee C, Tripathi V, Manolika EM, Heijink AM, Ricci G, Merzouk S *et al*. RIF1
promotes replication fork protection and efficient restart to maintain genome stability.
Nat Commun 2019; 10: 3287.

22 Garzon J, Ursich S, Lopes M, Hiraga SI, Donaldson AD. Human RIF1-Protein
Phosphatase 1 Prevents Degradation and Breakage of Nascent DNA on Replication
Stalling. *Cell Rep* 2019; 27: 2558-2566 e2554.

23 Dong Q, Day M, Saito Y, Parker E, Watts LP, Kanemaki MT *et al*. The human RIF1-
Long isoform interacts with BRCA1 to promote recombinational fork repair under
DNA replication stress. *Nat Commun* 2025; 16: 5820.

24 Wang H, Zhao A, Chen L, Zhong X, Liao J, Gao M *et al*. Human RIF1 encodes an
anti-apoptotic factor required for DNA repair. *Carcinogenesis* 2009; 30: 1314-1319.

25 Eke I, Zong D, Aryankalayil MJ, Sandfort V, Bylicky MA, Rath BH *et al*. 53BP1/RIF1
signaling promotes cell survival after multifractionated radiotherapy. *Nucleic Acids*
Res 2020; 48: 1314-1326.

615
616 26 Mei Y, Peng C, Liu YB, Wang J, Zhou HH. Silencing RIF1 decreases cell growth,
617 migration and increases cisplatin sensitivity of human cervical cancer cells.
618 *Oncotarget* 2017; 8: 107044-107051.

619
620 27 Noordermeer SM, Adam S, Setiaputra D, Barazas M, Pettitt SJ, Ling AK *et al*. The
621 shieldin complex mediates 53BP1-dependent DNA repair. *Nature* 2018; 560: 117-
622 121.

623
624 28 Franklin JM, Wu Z, Guan KL. Insights into recent findings and clinical application of
625 YAP and TAZ in cancer. *Nat Rev Cancer* 2023; 23: 512-525.

626
627 29 Howard A, Bojko J, Flynn B, Bowen S, Jungwirth U, Walko G. Targeting the
628 Hippo/YAP/TAZ signalling pathway: Novel opportunities for therapeutic interventions
629 into skin cancers. *Exp Dermatol* 2022; 31: 1477-1499.

630
631 30 Maehama T, Nishio M, Otani J, Mak TW, Suzuki A. The role of Hippo-YAP signaling
632 in squamous cell carcinomas. *Cancer Sci* 2021; 112: 51-60.

633
634 31 Hillmer RE, Link BA. The Roles of Hippo Signaling Transducers Yap and Taz in
635 Chromatin Remodeling. *Cells* 2019; 8.

636
637 32 Walko G, Woodhouse S, Pisco AO, Rognoni E, Liakath-Ali K, Lichtenberger BM *et al*.
638 A genome-wide screen identifies YAP/WBP2 interplay conferring growth advantage
639 on human epidermal stem cells. *Nat Commun* 2017; 8: 14744.

640
641 33 De Rosa L, Secone Seconetti A, De Santis G, Pellacani G, Hirsch T, Rothoeft T *et al*.
642 Laminin 332-Dependent YAP Dysregulation Depletes Epidermal Stem Cells in
643 Junctional Epidermolysis Bullosa. *Cell Rep* 2019; 27: 2036-2049 e2036.

644
645 34 Lavado A, Park JY, Pare J, Finkelstein D, Pan H, Xu B *et al*. The Hippo Pathway
646 Prevents YAP/TAZ-Driven Hypertranscription and Controls Neural Progenitor
647 Number. *Dev Cell* 2018; 47: 576-591 e578.

648
649 35 Gertzmann D, Presek C, Mattes AL, Sanger M, Zoller M, Schulein-Volk C *et al*.
650 Oncogenic YAP sensitizes cells to CHK1 inhibition via CDK4/6 driven G1
651 acceleration. *EMBO Rep* 2025.

652
653 36 Yuan Y, Park J, Feng A, Awasthi P, Wang Z, Chen Q *et al*. YAP1/TAZ-TEAD
654 transcriptional networks maintain skin homeostasis by regulating cell proliferation and
655 limiting KLF4 activity. *Nat Commun* 2020; 11: 1472.

657   37   Rheinwald JG, Beckett MA. Tumorigenic keratinocyte lines requiring anchorage and
658   fibroblast support cultured from human squamous cell carcinomas. *Cancer Res*
659   1981; 41: 1657-1663.

660
661   38   Zanconato F, Forcato M, Battilana G, Azzolin L, Quaranta E, Bodega B *et al*.
662   Genome-wide association between YAP/TAZ/TEAD and AP-1 at enhancers drives
663   oncogenic growth. *Nat Cell Biol* 2015; 17: 1218-1227.

664
665   39   Pattschull G, Walz S, Grundl M, Schwab M, Ruhl E, Baluapuri A *et al*. The Myb-MuvB
666   Complex Is Required for YAP-Dependent Transcription of Mitotic Genes. *Cell Rep*
667   2019; 27: 3533-3546 e3537.

668
669   40   Lopez-Pajares V, Qu K, Zhang J, Webster DE, Barajas BC, Siprashvili Z *et al*. A
670   LncRNA-MAF:MAFB transcription factor network regulates epidermal differentiation.
671   *Dev Cell* 2015; 32: 693-706.

672
673   41   Kypriotou M, Huber M, Hohl D. The human epidermal differentiation complex:
674   cornified envelope precursors, S100 proteins and the 'fused genes' family. *Exp*
675   *Dermatol* 2012; 21: 643-649.

676
677   42   Wang J, Harwood CA, Bailey E, Bewicke-Copley F, Anene CA, Thomson J *et al*.
678   Transcriptomic analysis of cutaneous squamous cell carcinoma reveals a multigene
679   prognostic signature associated with metastasis. *J Am Acad Dermatol* 2023; 89:
680   1159-1166.

681
682   43   Hagenbeek TJ, Zbieg JR, Hafner M, Mroue R, Lacap JA, Sodir NM *et al*. An allosteric
683   pan-TEAD inhibitor blocks oncogenic YAP/TAZ signaling and overcomes KRAS
684   G12C inhibitor resistance. *Nat Cancer* 2023; 4: 812-828.

685
686   44   Yuan Y, Salinas Parra N, Chen Q, Iglesias-Bartolome R. Oncogenic Hedgehog-
687   smoothened signaling depends on YAP1-TAZ/TEAD transcription to restrain
688   differentiation in basal cell carcinoma. *J Invest Dermatol* 2021.

689
690   45   Ashley AK, Shrivastav M, Nie J, Amerin C, Troksa K, Glanzer JG *et al*. DNA-PK
691   phosphorylation of RPA32 Ser4/Ser8 regulates replication stress checkpoint
692   activation, fork restart, homologous recombination and mitotic catastrophe. *DNA*
693   *Repair (Amst)* 2014; 21: 131-139.

694
695   46   Liu S, Opiyo SO, Manthey K, Glanzer JG, Ashley AK, Amerin C *et al*. Distinct roles for
696   DNA-PK, ATM and ATR in RPA phosphorylation and checkpoint activation in
697   response to replication stress. *Nucleic Acids Res* 2012; 40: 10780-10794.

698
699   47   Dempster JM, Pacini C, Pantel S, Behan FM, Green T, Krill-Burger J *et al*. Agreement
700   between two large pan-cancer CRISPR-Cas9 gene dependency data sets. *Nat*
701   *Commun* 2019; 10: 5817.

703 48   Takahashi N, Kim S, Schultz CW, Rajapakse VN, Zhang Y, Redon CE *et al*.
704   Replication stress defines distinct molecular subtypes across cancers. *Cancer Res*
705   *Commun* 2022; 2: 503-517.

706
707 49   Kotsantis P, Silva LM, Irmscher S, Jones RM, Folkes L, Gromak N *et al*. Increased
708   global transcription activity as a mechanism of replication stress in cancer. *Nat*
709   *Commun* 2016; 7: 13087.

710
711 50   Mohammed H, Taylor C, Brown GD, Papachristou EK, Carroll JS, D'Santos CS.
712   Rapid immunoprecipitation mass spectrometry of endogenous proteins (RIME) for
713   analysis of chromatin complexes. *Nat Protoc* 2016; 11: 316-326.

714
715 51   Zhang N, Bai H, David KK, Dong J, Zheng Y, Cai J *et al*. The Merlin/NF2 tumor
716   suppressor functions through the YAP oncoprotein to regulate tissue homeostasis in
717   mammals. *Dev Cell* 2010; 19: 27-38.

718
719 52   Zhou Y, Zhang J, Li H, Huang T, Wong CC, Wu F *et al*. AMOTL1 enhances YAP1
720   stability and promotes YAP1-driven gastric oncogenesis. *Oncogene* 2020; 39: 4375-
721   4389.

722
723 53   Wang P, Bai Y, Song B, Wang Y, Liu D, Lai Y *et al*. PP1A-mediated
724   dephosphorylation positively regulates YAP2 activity. *PLoS One* 2011; 6: e24288.

725
726 54   Zhu M, Peng R, Liang X, Lan Z, Tang M, Hou P *et al*. P4HA2-induced prolyl
727   hydroxylation suppresses YAP1-mediated prostate cancer cell migration, invasion,
728   and metastasis. *Oncogene* 2021; 40: 6049-6056.

729
730 55   Mittal P, Roberts CWM. The SWI/SNF complex in cancer - biology, biomarkers and
731   therapy. *Nat Rev Clin Oncol* 2020; 17: 435-448.

732
733 56   He L, Pratt H, Gao M, Wei F, Weng Z, Struhl K. YAP and TAZ are transcriptional co-
734   activators of AP-1 proteins and STAT3 during breast cellular transformation. *Elife*
735   2021; 10.

736
737 57   Ning B, Tilston-Lunel AM, Simonetti J, Hicks-Berthet J, Matschulat A, Pfefferkorn R *et*
738   *al*. Convergence of YAP/TAZ, TEAD and TP63 activity is associated with bronchial
739   premalignant severity and progression. *J Exp Clin Cancer Res* 2023; 42: 116.

740
741 58   Nowak DE, Tian B, Brasier AR. Two-step cross-linking method for identification of
742   NF-kappaB gene network by chromatin immunoprecipitation. *Biotechniques* 2005;
743   39: 715-725.

744
59   Melendez Garcia R, Haccard O, Chesneau A, Narassimprakash H, Roger J, Perron
M *et al*. A non-transcriptional function of Yap regulates the DNA replication program
in Xenopus laevis. *Elife* 2022; 11.

60 Proby CM, Purdie KJ, Sexton CJ, Purkis P, Navsaria HA, Stables JN *et al*.
Spontaneous keratinocyte cell lines representing early and advanced stages of
malignant transformation of the epidermis. *Exp Dermatol* 2000; 9: 104-117.

752
753 61 Chandrashekar DS, Karthikeyan SK, Korla PK, Patel H, Shovon AR, Athar M *et al*.
754   UALCAN: An update to the integrated cancer data analysis platform. *Neoplasia* 2022;
755   25: 18-27.

756
757 62 Zhang Y, Chen F, Chandrashekar DS, Varambally S, Creighton CJ. Proteogenomic
758   characterization of 2002 human cancers reveals pan-cancer molecular subtypes and
759   associated pathways. *Nat Commun* 2022; 13: 2669.

760
761 63 Jones MJK, Rai SK, Pfuderer PL, Bonfim-Melo A, Pagan JK, Clarke PR *et al*. A high-
762   resolution, nanopore-based artificial intelligence assay for DNA replication stress in
763   human cancer cells. *Nat Commun* 2025; 16: 7732.

764
765 64 Maya-Mendoza A, Moudry P, Merchut-Maya JM, Lee M, Strauss R, Bartek J. High
766   speed of fork progression induces DNA replication stress and genomic instability.
767   *Nature* 2018; 559: 279-284.

768
769 65 Raso MC, Djoric N, Walser F, Hess S, Schmid FM, Burger S *et al*. Interferon-
770   stimulated gene 15 accelerates replication fork progression inducing chromosomal
771   breakage. *J Cell Biol* 2020; 219.

772
773 66 Fousek-Schuller VJ, Borgstahl GEO. The Intriguing Mystery of RPA Phosphorylation
774   in DNA Double-Strand Break Repair. *Genes (Basel)* 2024; 15.

775
776 67 Toledo LI, Altmeyer M, Rask MB, Lukas C, Larsen DH, Povlsen LK *et al*. ATR
777   prohibits replication catastrophe by preventing global exhaustion of RPA. *Cell* 2013;
778   155: 1088-1103.

779
780 68 Picco G, Chen ED, Alonso LG, Behan FM, Goncalves E, Bignell G *et al*. Functional
781   linkage of gene fusions to cancer cell fitness assessed by pharmacological and
782   CRISPR-Cas9 screening. *Nat Commun* 2019; 10: 2198.

783
784 69 Petermann E, Orta ML, Issaeva N, Schultz N, Helleday T. Hydroxyurea-stalled
785   replication forks become progressively inactivated and require two different RAD51-
786   mediated pathways for restart and repair. *Mol Cell* 2010; 37: 492-502.

787
788 70 Moeglin E, Desplancq D, Conic S, Oulad-Abdelghani M, Stoessel A, Chiper M *et al*.
789   Uniform Widespread Nuclear Phosphorylation of Histone H2AX Is an Indicator of
790   Lethal DNA Replication Stress. *Cancers (Basel)* 2019; 11.

791
792 71 Cui P, Lin Q, Xin C, Han L, An L, Wang Y *et al*. Hydroxyurea-induced global
793   transcriptional suppression in mouse ES cells. *Carcinogenesis* 2010; 31: 1661-1668.

795 72   Li Q, Tie Y, Alu A, Ma X, Shi H. Targeted therapy for head and neck cancer: signaling
796   pathways and clinical studies. *Signal Transduct Target Ther* 2023; 8: 31.

797
798 73   Sato K, Faraji F, Cervantes-Villagrana RD, Wu X, Koshizuka K, Ishikawa T *et al*.
799   Targeting YAP/TAZ-TEAD signaling as a therapeutic approach in head and neck
800   squamous cell carcinoma. *Cancer Lett* 2025: 217467.

801
802 74   Chang CY, Shipony Z, Lin SG, Kuo A, Xiong X, Loh KM *et al*. Increased ACTL6A
803   occupancy within mSWI/SNF chromatin remodelers drives human squamous cell
804   carcinoma. *Mol Cell* 2021; 81: 4964-4978 e4968.

805
806 75   Kim M, Kim T, Johnson RL, Lim DS. Transcriptional co-repressor function of the
807   hippo pathway transducers YAP and TAZ. *Cell Rep* 2015; 11: 270-282.

808
809 76   Debaugnies M, Sanchez-Danes A, Rorive S, Raphael M, Liagre M, Parent MA *et al*.
810   YAP and TAZ are essential for basal and squamous cell carcinoma initiation. *EMBO*
811   *Rep* 2018.

812
813 77   Inman GJ, Wang J, Nagano A, Alexandrov LB, Purdie KJ, Taylor RG *et al*. The
814   genomic landscape of cutaneous SCC reveals drivers and a novel azathioprine
815   associated mutational signature. *Nat Commun* 2018; 9: 3667.

816
817 78   Pennycook BR, Vesela E, Peripolli S, Singh T, Barr AR, Bertoli C *et al*. E2F-
818   dependent transcription determines replication capacity and S phase length. *Nat*
819   *Commun* 2020; 11: 3503.

820
821 79   Kumagai A, Dunphy WG. Binding of the Treslin-MTBP Complex to Specific Regions
822   of the Human Genome Promotes the Initiation of DNA Replication. *Cell Rep* 2020;
823   32: 108178.

824
80   Watts LP, Natsume T, Saito Y, Garzon J, Dong Q, Boteva L *et al*. The RIF1-long
splice variant promotes G1 phase 53BP1 nuclear bodies to protect against replication
stress. *Elife* 2020; 9.

Figure legends

Figure 1. YAP/TAZ-TEAD-mediated oncogenic transcriptional programmes control the

balance between proliferation and differentiation in squamous cell carcinoma

progression.

(**A**, **B**) Violin blots (thick line, median; thin lines, upper- and lower quartiles) show the

distribution of Gene Set Variation Analysis (GSVA) scores for differentially expressed genes

(DEGs) that were (A) downregulated or (B) upregulated in siYAP/TAZ SCC13 cells across

different cSCC patient samples [42]. Norm_no_met / Tum_no_met: perilesional normal skin

and primary cSCC tissue from patients with no metastasis over 3 years; Norm_with_met /

Tum_with_met: perilesional normal skin and primary cSCC tissue from patients with

metastasis during 3 years follow-up; Met: tissue from metastasis. Pairwise comparisons

between sample groups were conducted using a *t* test. The p-values from these comparisons

are indicated above the boxplots with brackets linking the compared groups. The

significance levels are denoted as follows: *p=0.032; **p=0.0021; ***p=0.0002;

****p<0.0001. (**C**) Photograph shows lack of medium acidification by doxycycline-induced

shYAP compared to shNTC SCC13 cells during a 4-day culture period. (**D**) Immunoblot

analysis of SCC13 cells expressing doxycycline-induced shYAP, shTAZ, or shNTC for 72 h, using

the indicated antibodies. Actin or cyclophilin B (CBP) were used as loading controls. (**E**) Cell

cycle analysis by propidium iodide (PI) labelling of asynchronous populations of shNTC and

shYAP SCC13 cells (grey profiles), and of cells synchronised at the G1/S boundary by double

thymidine block (dTb) and released into S phase and analysed at indicated time points

(T=hours post thymidine release, coloured profiles). (**F**) Colony formation assays were

quantified by measuring the percentage of well area occupied by colonies. Bars show the

means from n=3 independent experiments (performed with 3 technical replicates),

individual data points (different shapes indicate different experiments) show the means

from each experiment, error bars show SD. Exact p-values are shown, RM one way ANOVA

with Geisser-Greenhouse correction and Dunnett's multiple comparison test. (**G**) 3D

spheroid growth of shNTC and shYAP SCC13 cells. p-values are shown for multiple

comparisons at all timepoints, Ordinary two-way ANOVA and Šidák's multiple comparison

test. *p<0.033, **p<0.002, ***p<0.001. Sources of variation: shRNA, p<0.001; time,

p<0.001; shRNA x time, p<0.001.

Figure 2. **Pharmacological inhibition of YAP/TAZ-TEAD complex formation phenocopies YAP**

depletion.

(**A**) Representative proliferation dose–response curves (means±SD) of SCC13 cells treated

with different TEAD inhibitors. (**B**) Representative viability dose–response curves

(means±SD) of different cell lines from a human cSCC progression series [60] treated with

different GNE-7883 concentrations. (**C**) Immunoblot analysis of SCC13 cells treated with

vehicle control (DMSO) or GNE-7883 (2 µM) for 48 h, using the indicated antibodies. Tubulin

was used as loading control. (**D**) qRT-PCR analysis of *IVL* mRNA expression in SCC13 cells

treated with GNE-7883 (2 µM). Data shown are from n=1 experiment performed with three

technical replicates. Individual data points: mean fold change in mRNA abundance

(normalized to GAPDH) compared to DMSO control. Bars represent the means from all

replicates; error bars represent SD. (**E**) SCC13 3D spheroid growth in the presence of vehicle

(DMSO) or different concentrations of GNE-7883 (GNE). p-values are shown for multiple

comparisons at 12 days timepoint, Ordinary two-way ANOVA and Dunnett's multiple

comparison test. *p<0.033, **p<0.002, ***p<0.001. Sources of variation: GNE treatment,

p<0.001; time, p<0.001; GNE treatment x time, p<0.001. (**F**) Cell viability measured at the

endpoint of the experiment shown in (E). Each datapoint represents one spheroid. Bars

represent the means from all spheroids; error bars represent SD. Exact p-values are shown,

Kruskal-Wallis test with Dunn's multiple comparison test. (**G**) Immunoblot analysis of SCC13

cells expressing shNTC or different shP21 treated with vehicle control (DMSO) or GNE-7883

(2 µM), using the indicated antibodies. Tubulin was used as loading control. (**H**) Phase

contrast images show that p21 depletion rescues cell viability in GNE-7883 (2 μM)-treated

SCC13 cells. Bar, 10 μm.

Figure 3. RIME identifies RIF1 as an interactor of chromatin-associated YAP in squamous

cell carcinoma cells.

(**A**) Interaction network of YAP protein interactors identified by RIME in SCC13 cells. Oval

nodes: proteins that were previously identified as candidate regulators of SCC13 cell

proliferation and/or survival in a genome wide RNA-interference (RNAi) screen (hit:

'Yes')[32]; rectangular shaped nodes: proteins that were not identified in the RNAi screen

(hit: 'No'). Red connecting lines (edges) show previously identified YAP interactions

according to the STRING Homo sapiens database, and nodes with red borders indicate

known YAP interactors (manually curated from STRING and literature search); dashed red

lines show manually curated known YAP interactions that were not yet present in the

STRING database. Proteins were manually grouped based on function (gene names

displayed, generated in Cytoscape 3.9.1 using STRING app 1.7.1). (**B**) Gene ontology terms

('molecular functions' and 'biological process') enriched within the high-confidence YAP-

interacting proteins identified by RIME. (**C**) Confocal fluorescence image (z-projection) of YAP

(green) and RIF1 (red) nuclear localisation in SCC13 cells. DNA was counterstained with

Hoechst dye (blue). Scale bar, 10 μm. (**D**, **E**) Immunoblot analysis of YAP and IgG-control

immunoprecipitates and respective inputs from SCC13 (D) whole cell extracts and (E) and

nuclei-enriched fractions using the indicated antibodies. (**F**, **G**) Representative confocal

microscopy images (z-projections) and image quantifications of (F) YAP-RIF1 and (G) YAP-

TEAD1 in-situ PLA in control (siNTC&shNTC) and YAP/TAZ knockdown (siYAP/TAZ&shYAP)

cells (PLA signal, green; DNA (DAPI) blue; n >250 cells (YAP-RIF1), n >500 cells (YAP-TEAD1)

per condition in n=3 independent experiments). Scale bars, 5 µm. Datapoints show fold

decrease in mean nuclear PLA signal intensity in siYAP/TAZ&shYAP cells compared to

siNTC&shNTC cells. Lines show means from all three experiments, error bars show SD. Exact

p-values are shown, one sample two-tailed *t* test.

Figure 4. YAP/TAZ-TEAD modulate RIF1 protein stability.

(**A**) qRT-PCR analysis of *RIF1* mRNA expression in SCC13 cells transfected with an siRNA

SMARTpool targeting YAP/TAZ (siY/T) or a non-targeting control SMARTpool (siNTC) for 48 h.

Data shown are from n=3 independent experiments performed with three technical

replicates. Individual data points: mean fold change in mRNA abundance (normalized to

GAPDH) compared to siNTC in each experiment. Bars represent the means from all

experiments, error bars represent SD. Exact p-values are shown, paired *t* test. (**B–D**)

Immunoblot analysis of SCC13 cells (B) transfected with siNTC or siY/T for 48 h, (C)

expressing empty vector control (MCS) or different YAP transgenes upon doxycycline (dox)

induction, or (D) treated with GNE-7883 (2 µM) for 24 h, using the indicated antibodies.

Tubulin or cyclophilin B (CBP) were used as loading controls. Arrows, position of full-length

RIF1 bands. In (B, D) numbers below lanes represent protein ratios relative to an arbitrary

level of 1.0 set for siNTC or DMSO control, respectively. (**E**) qRT-PCR analysis of *RIF1* mRNA

expression in SCC13 cells treated with GNE-7883 for 24 h. Data shown are from n=1

experiment performed with three technical replicates. Individual data points: mean fold

change in mRNA abundance (normalized to GAPDH) compared to DMSO control. Bars

represent the means from all replicates, error bars represent SD. (**F**) Immunoblot analysis of

RIF1 protein levels in SCC13 cells transfected with siRNA SMARTpools targeting TEAD1,

TEAD3, TEAD4, or YAP/TAZ (siY/T), or an siNTC SMARTpool for 24 h. Cyclophilin B (CBP) was

used as loading control. (**G**) Quantification of RIF1 protein expression in experiment shown

in (F). Individual data points show fold change in RIF1 protein levels (normalised to tubulin)

in siTEAD1, siTEAD3, siTEAD4, siTEAD1/3/4, and siY/T compared to siNTC in n=3 independent

experiments (different shapes of data points), bars show the mean, error bars show SD. (**H**)

Immunoblot analysis of chromatin fractions from SCC13 cells expressing doxycycline-induced

shYAP or shNTC for 48 h, using the indicated antibodies. Cells were synchronised at the G1/S

boundary by double thymidine block (dTb) and released into the S phase. Histone H3 (HH3)

was used as loading control. Arrow, position of full-length RIF1 band. (**I**) Quantification of

RIF1 protein levels in experiment shown in (H). Individual data points show fold change in

RIF1 protein levels (normalised to HH3) compared to shNTC at timepoint 0 h in n=2

independent experiments (different shapes of data points).

Figure 5. Increased expression of RIF1 in squamous cell carcinoma.

(**A**) Immunoblot analysis of normal human epidermal keratinocytes (NHKs), keratinocytes

isolated from a pre-malignant dysplastic skin lesion (PM1), various cell lines isolated from

primary cutaneous SCCs (cSCC; SCC13, SCC-NR, MET1), a recurrent cSCC (MET2), and a

lymph node metastasis (MET4), using the indicated antibodies. Actin was used as loading

control. Arrow, position of full-length RIF1 band. (**B**, **C**) Quantification (B) and representative

images (C) of RIF1 expression in healthy adult human skin (HS), nodular (Nod) and (Inf)

infiltrative basal cell carcinoma (BCC), and well-differentiated (WD) and poorly differentiated

(PD) cSCC. Tissue sections were immunolabelled with antibodies against RIF1 (green), and

keratin 14 (Krt14, blue) to identify normal and neoplastic epidermal keratinocytes. White

dotted line demarcates dermal–epidermal boundary in HS. Scale bar, 50 μm. Superplots in

(B) show percentage of RIF1-positive keratinocytes in different areas of the same tumour

(coloured symbols), mean percentage of RIF1-positive keratinocytes in 3–5 different patient

samples (black datapoints), and mean percentage of RIF1-positive keratinocytes in each

sample group (lines). Error bars show SD from each sample group. *$p<0.05$, Brown-Forsythe

and Welsh ANOVA test with Dunett's multiple comparison test. (**D**, **E**) RIF1 protein

expression in HNSCC samples from the Clinical Proteomic Tumour Analysis Consortium

(CPTAC), stratified according to (D) tumour stage, or (E) presence of alterations in the Hippo

pathway or other signalling pathways. Jitter plots (overlayed onto box plots) show Z-values

(standard deviations from the median across samples for the given sample type) from

individual samples. Box plots indicate the median (middle line), 25th percentile (bottom line),

75th percentile (top line), and minimum and maximum (whiskers). Log2 spectral count ratio

values from CPTAC were first normalized within each sample profile, then normalized across

samples.   Graphs   were   downloaded   from   the   UALCAN   database

(https://ualcan.path.uab.edu/). (D) $p<0.05$ for all pairwise comparisons. In (E) exact p-values

are shown.

Figure 6. Depletion of RIF1 in squamous cell carcinoma cells affects DNA replication

dynamics.

(**A**) Representative cell cycle analysis by PI labelling of asynchronous populations of shNTC

and shRIF1 SCC13 cells (grey profiles) and of cells synchronised at the G1/S boundary by

double thymidine block (dTb) and released into S phase (T=hours post thymidine release,

coloured profiles). (**B**, **C**) Immunoblot analysis of (B) cell extracts and (C) chromatin fractions

from shNTC and shRIF1 SCC13 cells, using the indicated antibodies. In (C) cells were

synchronised at the G1/S boundary by double thymidine block (dTB) and released into the S

phase. Note that DDK-induced phosphorylation of MCM4 results in a band shift. Cyclophilin

B (CPB), tubulin, and histone H3 (HH3) were used as loading controls. shRIF1-017 was used

in experiments shown in (A, C). Arrows in (B, C), position of full-length RIF1 band. (**D, E**)

Replication fork speed (D) and replication track length (E) measured by DNAscent. Mean fork

speed (kb/min) and track length (bp) are indicated. Black bars show medians. Total number

of forks scored in n=2 independent experiments: shNTC=5632; shRIF1=6216. (**F**) Combined

distribution of stall scores from n=2 independent experiments. Mean stall cores are

indicated.

Figure 7. RIF1 protects against endogenous replication stress.

(**A, B**) Immunoblot analysis of cell extracts from SCC13 cells expressing shNTC or different

shRIF1, using the indicated antibodies. Cyclophilin B (CPB), tubulin, or GAPDH were used as

loading controls. Cells treated for 24 h with 4 mM HU served as positive controls for severe

replication stress. Positions of strongly hyperphosphorylated RPA32 bands are indicated with

asterisks. Arrows, positions of full-length RIF1 bands. (**C**) Representative cell cycle profile (PI

labelling following a 30-min EdU pulse) of unsynchronised shNTC and shRIF1 SCC13 cells.

Gates used for quantification of percentage of cells in the different cell cycle phases are

shown. (**D**) Quantification of percentage of cells in different cell cycle phases. Bars show

means from n=4 independent experiments, individual data points show cell percentages in

each experiment, error bars show SD. Exact p-values are shown, paired *t* test. (**E, F**)

Immunoblot analysis of cell extracts from SCC13 cells transfected with (E) siNTC, siYAP/TAZ

(siY/T), or siRIF1 for 48 h, or treated with cisplatin (CP, 4 µM) or hydroxyurea (HU, 4 mM) for

24 h to induce S phase arrest, or (F) expressing shNTC or different shRIF1, using the indicated

antibodies. Tubulin was used as loading control. (**G**) Representative images of colony

formation efficiency of shNTC and shRIF1 SCC13 cells. Cells in triplicate wells were stained

with crystal violet. (**H**) Percentage of well area occupied by colonies. Bars show the means

from n=3 independent experiments (performed with 3 technical replicates), individual data

points (different shapes indicate different experiments) show the means from each

experiment, error bars show SD. Exact p-values are shown, RM one way ANOVA with

Geisser-Greenhouse correction and Dunnett's multiple comparison test.

1007

Figure 8. YAP is involved in DNA damage repair at collapsed replication forks.

(**A**) Immunoblot analysis of SCC13 control cells and cells exposed to 4 mM hydroxyurea (HU)

for 24 h, using the indicated antibodies. Washout, medium removal and replacement with

fresh medium. Tubulin or cyclophilin B (CBP) were used as loading controls. (**B**)

Representative microscopy images of RAD51 (green) foci in SCC13 cells treated with 4 mM

HU for the indicated times. Dashed outlines indicate nuclear areas from DNA counterstain

with DAPI. Scale bar, 25 μm. (**C**) Immunoblot analysis of SCC13 cells exposed to 4 mM HU for

24 h, followed by HU washout, using the indicated antibodies. Tubulin was used as loading

control. (**D**, **E**) Image quantifications of (D) YAP-RIF1 and (E) YAP-γH2AX in-situ PLA in SCC13

cells treated with 4 mM HU for the indicated times. >200 cells were measured per

experimental condition in n=3 independent experiments. Superplots show nuclear PLA

signal intensity per cell (grey datapoints), mean nuclear PLA signal intensity of the respective

cell populations in each of n=3 experiments (black datapoints, different shapes indicate

different experiments), and mean nuclear PLA signal intensity from all three experiments

(lines). Error bars show SD from all three experiments. Exact p-values are shown. (D) Brown-

Forsythe and Welsh ANOVA test with Dunett's multiple comparison test, (E) unpaired *t* test.

(**F**) Immunoblot analysis of SCC13 cells treated with 2 mM HU for 24 h or left untreated

(control), 24 h after transfection with siNTC or siYAP/TAZ (siY/T), using the indicated

antibodies. Tubulin was used as loading control. (**G, H**) Synergism of YAP depletion (shYAP)

and HU treatment on clonal colony growth was quantified by measuring the percentage of

well area occupied by colonies after 10 days in culture. In (H) values have been normalized

to controls (no HU treatment) for shYAP and shNTC. Bars show the means from n=3

independent experiments (performed with 3 technical replicates), individual data points

show the means from each experiment, error bars show SD. p-values are shown for multiple

comparisons, RM two-way ANOVA (matched values are spread across a row) and Šidak's

multiple comparison test. *p<0.033, **p<0.002, ***p<0.001. Sources of variation in (G, H):

HU concentration, p<0.001; shRNA, p<0.001; HU concentration x shRNA, p<0.001.

1035

Figure 1_Bojko et al.

Figure 2_Bojko et al.

A

Intracellular Trafficking

Proteasome

Chromatin Remodelling

RNAi screen hit
No
Yes

DNA Repair

DNA Replication

Cell Cycle

Gene Expression

Nuclear Structure

Cell Signalling

B

SCC13 YAP-RIME GO MF

- Nucleosome binding
- Nucleosomal DNA binding
- Chromatin DNA binding
- Transcription coactivator activity
- RNA polymerase II-specific DNA-binding transcription factor binding

-log10(FDR)

SCC13 RIME GO BP

- Nucleosome disassembly
- Protein-DNA complex disassembly
- Chromatin remodeling
- Protein-containing complex disassembly
- Protein complex oligomerization
- Nucleosome organization
- Protein-DNA complex subunit organization

-log10(FDR)

C

YAP   RIF1   DAPI   Merge

D

Mw [kDa]   Input (1%)   YAP-IP   IgG-IP

70 —   YAP
250 —   RIF1

Whole cell lysate

E

Mw [kDa]   Input (1%)   YAP-IP   IgG-IP

70 —
50 —   YAP
250 —   RIF1
50 —   pan-TEAD
250 —   ARID1A
250 —   CHD4

Nuclear fraction

F

shNTC

shYAP/TAZ

YAP-RIF1 PLA

0.0337

Mean decrease in YAP-RIF1 PLA Signal

shYAP/TAZ / shNTC Mean nuclear PLA intensity

G

shNTC

shYAP/TAZ

YAP-TEAD1 PLA

0.0211

Mean decrease in YAP-TEAD1 PLA Signal

shYAP/TAZ / shNTC Mean nuclear PLA intensity

Figure 3_Bojko et al.

Figure 4_Bojko et al.

Figure 5_Bojko et al.

Figure 6_Bojko et al.

Figure 7_Bojko et al.

Figure 8_Bojko et al.