Benchmarking AI Models for *In Silico* Gene Perturbation of Cells

Chen Li[1,2#], Haoxiang Gao[2#], Yuli She[2#], Haiyang Bian[1,2], Qing Chen[2], Kai Liu[2*],

Lei Wei[1*] and Xuegong Zhang[1,3*]

1 MOE Key Laboratory of Bioinformatics and Bioinformatics Division of BNRIST, Department of Automation, Tsinghua University, Beijing 100084, China

2 Anew Therapeutics Pte. Ltd., Singapore 068896, Singapore

3 Center for Synthetic and Systems Biology, School of Life Sciences and School of Medicine, Tsinghua University, Beijing 100084, China

* To whom correspondence should be addressed. Email: zhangxg@tsinghua.edu.cn; weilei92@tsinghua.edu.cn; kai.liu@anewbt.com

# These authors contributed equally.

Abstract

Understanding perturbations at the single-cell level is essential for unraveling cellular mechanisms and their implications in health and disease. The growing availability of biological data has driven the development of a variety of *in silico* perturbation methods designed for single-cell analysis, which offer a means to address many inherent limitations of experimental approaches. However, these computational methods are often tailored to specific scenarios and validated on limited datasets and metrics, making their evaluation and comparison challenging. In this work, we introduce a comprehensive benchmarking framework to systematically evaluate *in silico* perturbation methods across four key scenarios: predicting effects of unseen perturbations in known cell types, predicting effects of observed perturbations in unseen cell types, zero-shot transfer to bulk RNA-seq of cell lines, and application to real-world biological cases. For each scenario, we curated diverse and abundant datasets, standardizing them into flexible formats to enable efficient analysis. Additionally, we developed multiple metrics tailored to each scenario, facilitating a thorough and comparative evaluation of these methods. Our benchmarking study assessed 10 methods, ranging from linear baselines to advanced machine learning approaches, across these scenarios. While some methods demonstrated surprising efficacy in specific contexts, significant challenges remain, particularly in zero-shot predictions and the modeling of complex biological processes. This work provides a valuable resource for evaluating and improving *in silico* perturbation methods, serving as a foundation for bridging computational predictions with experimental validation and real-world biological applications.

Introduction

Understanding responses of gene expression to gene perturbations in cells is crucial for decoding cellular functions and has important implications for many applications in biomedical research. With advancements in single-cell sequencing and gene perturbation technologies, methods such as Perturb-seq[1] and CROP-seq[2], which integrate single-cell RNA sequencing (scRNA-seq) with CRISPR-based perturbations, enable large-scale pooled assays[3, 4] to investigate gene functions and cellular responses at single-cell resolution. However, experimental approaches face significant limitations: with ~20,000 protein-coding genes and ~400 major cell types in humans[5], it is practically impossible to explore all gene perturbations across all cell types, let alone explore the vast array of gene combinations. Moreover, these experiments usually require multi-day cell incubation, but many cell types cannot survive in culture for extended periods, limiting its applicability to many biological contexts.

These constraints have spurred the development of a number of artificial intelligence (AI) models for *in silico* gene perturbation, which predict cellular states (typically represented as gene expression profiles) following gene perturbations. Dynamo[6] introduced the concept of *in silico* perturbation for single-cell datasets. It leverages the analytical Jacobian of a reconstructed vector field function to encode gene regulatory networks (GRNs). By propagating perturbations through the GRN, Dynamo predicts cell-fate outcomes following perturbation. CellOracle[7] developed an algorithm to infer causal GRNs using both scRNA-seq and single-cell assay for transposase-accessible chromatin sequencing (scATAC-seq) data, enabling GRN-based *in silico* perturbations. GEARS[8] marked a significant advancement as the first method capable of predicting transcriptional outcomes for unseen perturbations by training on Perturb-seq data, learning gene embeddings through a gene co-expression graph and perturbation embeddings through a Gene Ontology (GO) term knowledge graph. Inspired by GEARS, subsequent methods[9, 10] refined this strategy to achieve improved performance. In parallel, approaches focusing on predicting specific treatment responses in novel cell types have emerged, with the variational auto-encoder (VAE) framework widely adopted. Examples include scGen[11], CPA[12], sVAE[13], and SAMS-VAE[14]. Additionally, optimal transport theory has also been applied to *in silico* perturbation, with notable examples including CellOT[15] and CINEMA-OT[16]. Recently, single-cell foundation models have introduced a new paradigm in *in silico* perturbation. Models like scGPT[17] and scFoundation[18] leverage large-scale pretrained architectures to deliver promising performance. Overall, these *in silico* gene perturbation methods utilized advanced statistical and deep learning algorithms, showing great potential to expand our understanding of gene perturbations and cellular responses for various cell types.

However, several challenges hinder the development and comparison of *in silico* gene perturbation methods. These methods are often tailored to specific scenarios, validated on limited datasets, and assessed using inconsistent metrics. For example, GEARS aims to predict unseen perturbations in known cell types, whereas scGen and CPA focus on predicting responses in unseen cell types with known perturbations. In addition, these methods are typically evaluated on narrowly focused datasets and metrics suited to specific contexts. For

instance, Dynamo and CellOracle assessed their performance on datasets representing biological developmental trajectories, such as the hematopoietic data, while GEARS and scGPT used single-cell perturbation datasets for validation. This reliance on limited and narrowly focused datasets introduces biases into the evaluation process, complicating the assessment of their effectiveness and generalizability. Furthermore, the inconsistency in evaluation metrics across studies, ranging from qualitative metrics like trajectory direction alignment to various quantitative metrics, further complicates performance comparisons.

Some benchmarking studies have attempted to evaluate existing *in silico* perturbation methods[19-22], but they often focus on a limited subset of methods, datasets, or scenarios, leading to incomplete or biased conclusions. For example, Wu et al.[20] proposed a perturbation benchmark for various machine learning methods, but it did not include evaluations for predicting unseen perturbations. Similarly, Eltze el al.[22] conducted a study claiming that existing deep learning methods do not outperform simple linear models, but the evaluation relied on the L2 error across all genes, which may focus more on the significantly varied genes while overlooking the invariant ones. These limitations highlight the absence of a comprehensive benchmarking framework that provides standardized datasets and evaluation metrics to fairly compare methods. Just as the MNIST[23] and ImageNet[24] datasets revolutionized computer vision by enabling systematic performance comparisons, a well-designed benchmark could play a transformative role in the development of *in silico* perturbation methods.

In this work, we presented a comprehensive benchmarking framework with rich collection of data to systematically evaluate *in silico* gene perturbation method. Depending on various usage requirements, we categorized *in silico* perturbation methods into four distinct scenarios: unseen perturbation transfer, useen cell type transfer, zero-shot transfer, and cell state transition prediction. We curated and filtered the gene perturbation datasets of scPerturb, retaining 17 single-cell datasets encompassing 984,000 cells and 3,190 perturbations. We designed a comprehensive set of evaluation metrics to assess various aspects of method performance. We conducted benchmarking study to assess 10 methods across all scenarios, including linear baseline methods, specific machine learning methods and large pretrained AI models. We observed that while some methods demonstrated surprising efficacy in specific contexts, significant challenges remain, particularly in zero-shot predictions and the modeling of complex biological processes. The presented benchmarking framework and datasets offered a valuable resource for the scientific community, serving as a foundation to promote the further development and refinement of AI models for *in silico* perturbation.

Results

Framework for benchmarking *in silico* perturbation

We present a benchmarking framework designed to evaluate *in silico* gene perturbation methods across four biologically and computationally relevant scenarios (**Fig. 1a**). The unseen perturbation transfer scenario assesses the ability to predict effects of previously unobserved perturbations in known cell types. The unseen

cell type transfer scenario evaluates the prediction of known perturbations in previously unobserved cell types.

The zero-shot transfer scenario evaluates model performance in predicting perturbation effects measured

within the CMAP framework, where perturbations are assessed using bulk RNA-seq data of cell lines.

The cell state transition prediction scenario focuses on predicting effects of key genes driving cell state

transitions in specific biological processes. The first two scenarios emphasize well-defined transfer tasks with

available training data, while the latter two reflect more challenging real-world applications where *in silico*

perturbation tools often need to generalize without task-specific training.

For the unseen perturbation transfer scenario, we collected single-cell perturbation datasets from scPerturb[25]

as benchmarking data (**Fig. 1b**). After quality control, preprocessing, and reformatting, we curated 17 single-

cell datasets, resulting in 984,000 cells and 3,190 gene perturbations, including CRISPR-Cas9 knockout,

CRISPR interference (CRISPRi), and CRISPR activation (CRISPRa) (Methods). Genes were classified into

differentially expressed genes (DEGs) and non-DEGs. For DEGs, benchmarking metrics included Pearson

Fig. 1. Overview of the *in silico* perturbation benchmarking framework. (a) Four benchmarking scenarios evaluated in this study. **(b)** Datasets curated for benchmarking. **(c)** Metrics for designed for evaluation across scenarios. **(d)** Methods benchmarked in this study.

correlation coefficient (PCC) and direction accuracy (ACC), while performance for non-DEGs was assessed

using mean absolute error (MAE) (**Fig. 1c**).

The unseen cell type transfer scenario used two datasets from Replogle et al.[4], which contained the largest

number of perturbations among our curated datasets, selecting 490 shared perturbations as benchmarking data

(**Fig. 1b**). The dataset from the K562 cell line was used for training, while the RPE1 dataset served as testing

data. Evaluation metrics were consistent with those in the unseen perturbation transfer scenario (**Fig. 1c**).

In the zero-shot transfer scenario, we employed Connectivity Map (CMAP) data[26], measured at the bulk level,

to evaluate model performance (**Fig. 1b**). Single-cell RNA-seq datasets[27, 28] from six cell lines overlapping

with CMAP data were used as the input for *in silico* perturbation methods. After quality control, a total of

21,361 perturbations from six cell lines served as the ground truth (Methods). For benchmarking, we selected

the top significant genes (genes with high absolute z-scores) from the CMAP data for each perturbation and

calculated direction accuracy (ACC) of these genes (**Fig. 1c**, Methods).

For the cell state transition prediction scenario, we collected four scRNA-seq datasets[29-32] representing distinct

biological processes, including development, reprogramming, and disease progression (**Fig. 1b**). For each

dataset, there are substantial reliable evidence showing that individual genes or specific gene combinations

can drive certain cell state transitions. Qualitative evaluation included streamline plots in uniform manifold

approximation and projection (UMAP) space to visualize predictions, while a ranking transition score (RTS)

was introduced to quantitatively measure accuracy of predicted transitions in response to perturbations (**Fig.**

1c, Methods).

Considering reproducibility and broad applicability, we evaluated 10 methods in this work, spanning linear

models, gene regulatory network (GRN)-based approaches, and machine learning techniques (**Fig. 1d**,

Methods). These included linear models[22] (with and without scGPT embeddings, referred to as Linear and

Linear-GPT, respectively), correlation-based method, CellOracle[7], GEARS[8], scGen[11], CPA[12], CellOT[15],

scGPT[17], and scFoundation[18]. Given the varying design and applicability of these methods, we selected and

evaluated an appropriate subset for each specific scenario.

Benchmarking in the unseen perturbation transfer scenario

140   In the unseen perturbation transfer scenario (**Fig. 2a**), we utilized version 1.3 of the scPerturb database as our

141   benchmarking data. scPerturb contains 50 single-cell perturbation datasets for RNA and protein expression.

142   Following rigorous data filtering and standardized processing (Methods), we curated 17 scRNA-seq datasets[1-

143   4, 33-37] for analysis.

Fig. 2. Benchmarking in the unseen perturbation transfer scenario. (a), Schematic diagram of unseen perturbation transfer. **(b)** Performance of methods on 17 single-cell perturbation datasets. Each dataset is exhibited with attributes of perturbation type, perturbation number, and mean E-distance. Metrics are divided into 4 parts: DEG_PCC (Pearson correlation coefficient for differentially expressed genes), DEG_ACC (direction accuracy for differentially expressed genes), SHG_MAE (mean absolute error for the genes that are stably highly expressed), and SLG_MAE (mean absolute error for the genes that are stably lowly expressed). The top three methods for each metric on each dataset are highlighted. Color indicates ranking, and bar length represents the magnitude of the metric. For DEG_PCC, negative values are not shown. **(c)** Hierarchically-clustered heatmap of methods' performance rankings across datasets. The datasets are broadly clustered into 3 groups (G1, G2, and G3). **(d)** Scatter plot showing the mean E-distance and the number of perturbations for each dataset, accompanied by bar plots comparing the mean E-distance and perturbation number across different dataset groups. **(e)** Radar plot of DEG_PCC for 7 methods across 4 selected datasets. **(f)** Box plots comparing DEG_PCC and DEG_ACC of 7 methods on Norman2019 and Replogle2022_K562 datasets, with different levels of DEGs (top 20, 50, 100, 200, and all) selected for each metric. Center line, median; box limits, upper and lower quartiles; whiskers, 1.5× interquartile range.

We calculated the E-distance, a statistical measure of the distance between point clouds, applied to pre- and

post-perturbation cell pairs for each perturbation as an indicator of the magnitude of perturbation effect[25]. The

number of perturbations in these 17 datasets varies widely, ranging from 7 (Adamson2016_001 and

Adamson2016_005) to 1,299 (Replogle2022_K562), highlighting substantial differences in data scale.

Similarly, the mean E-distances also vary considerably across datasets, spanning from 0.1 to 48.0, likely

reflecting differences in experimental conditions. We summarized key attributes of these datasets, including

the gene perturbation technique, the number of perturbations and the mean E-distance in **Fig. 2b**.

Considering that only a subset of genes is affected by specific perturbations, we categorized the genes in the

gene expression profiles for each perturbation into three sets: Differentially expressed genes (DEG), which

exhibit significant expression differences between pre- and post-perturbation cells; (2) Stably highly

expressed genes (SHG), which are not differentially expressed and show high expression in pre-perturbation

cells; and (3) Stably lowly expressed genes (SLG), which are not differentially expressed and show low

expression in pre-perturbation cells. For the DEG set, we evaluated performance using PCC (DEG_PCC) and

ACC (DEG_ACC) values, while for the SHG and SLG set, we assessed performance using MAE values,

referred to as SHG_MAE and SLG_MAE, respectively (Methods).

To benchmark the methods, we evaluated the performance of six methods: Linear, Linear-GPT, CellOracle,

GEARS, scGPT, and scFoundation (Methods). We also included a basic approach that averages gene

expression across all cells within all known perturbations as the prediction of unseen perturbations (referred to

as KnownAverage).

The benchmarking results across the 17 datasets were summarized in **Fig. 2b**. Notably, the KnownAverage

method consistently demonstrated some of the best overall performance across all four types of metrics.

Specifically, for the DEG_PCC metric, the KnownAverage method ranked among the top three in most

datasets. The Linear and Linear-scGPT method also achieved top ranks in specific datasets, such as

Adamson2016_005, Dixit2016, and Papalexi2021_array. Meanwhile, single-cell foundation models, including

scGPT and scFoundation, outperformed other methods in datasets such as Replogle2022_K562,

Replogle2022_RPE1, and Norman2019. GEARS and CellOracle achieved top ranking in only a few datasets,

with CellOracle showing the weakest overall performance. For DEG_ACC, the overall performance trends

mirrored those of DEG_PCC, with the KnownAverage method exhibiting superior performance and ranking

top in nearly all the datasets.

For both SHG_MAE and SLG_MAE, the KnownAverage method again achieved best performance in nearly

all datasets, and Linear-scGPT consistently ranks in the top three for most cases. Although deep learning

methods such as GEARS, scGPT, and scFoundation did not consistently achieve top ranks, their metrics were

not significantly worse than those of the best-performing methods. The Linear method exhibited a notable

performance drop in specific datasets, such as Papalexi2021_array and Replogle2022_RPE1.

We performed hierarchical clustering on the heatmap of DEG_PCC rankings of models across the 17 datasets

(we did not include CellOracle because it is not compatible with all datasets.). The results revealed distinct

performance patterns, grouping the datasets into three clusters: G1, G2, and G3 (**Fig. 2c**). To better understand
these groupings, we examined the perturbation number and mean E-distance for each dataset and compared
the average attributes of the three groups (**Fig. 2d**). Group G3 exhibited the highest average perturbation
number (527.8) and mean E-distance (24.2), indicating that these datasets include a wide range of
perturbations with significant phenotypic effects pre-perturbation cells. Single-cell foundation models, such as
scGPT and scFoundation, performed best on G3 datasets, likely due to the abundance of heterogeneous
perturbations in the training data, which supports the learning capabilities of large deep learning models.

Group G2 displayed a moderate average perturbation number (87.2) but a low average E-distance (2.2),
suggesting that perturbations in these datasets induced only subtle shifts in expression profiles. Interestingly,
scFoundation achieved the best performance on G2 datasets, demonstrating its ability to capture slight
perturbation effects more effectively than other methods, including scGPT.

Group G1 showed the lowest average perturbation number (25.2). In these datasets, simple methods like
KnownAverage, Linear, and Linear-scGPT outperformed deep learning methods. This performance gap is
likely attributed to the restricted number of perturbations, which limits the capacity of deep learning models to
effectively learn perturbation effects. Additionally, some datasets in G1 include perturbed genes that belong to
the same functional group. For instance, in the Datlinger2021 dataset[2], the 20 perturbed genes were
specifically selected for their relevance to the T cell receptor pathway. This likely reduces the complexity and
heterogeneity of the perturbations, making the simpler methods more effective in modeling the observed
perturbation effects.

Based on these observations, we conclude that datasets with higher perturbation numbers and larger E-
distances, such as those in D3, are better suited for extending to genome-wide unseen perturbations. These
characteristics provide a robust foundation for fair and comprehensive comparisons of different methods.

We further selected 4 datasets from Group G3 for final benchmarking, focusing on datasets with a high
number of perturbations (>150) and substantial E-distances (>5): Replogle2022_K562[4],
Replogle2022_RPE1[4], Norman2019[38], and Sunshine2023[39]. For the DEG_PCC metric, single-cell foundation
models, including scFoundation and scGPT, outperformed all other methods, indicating their ability to capture
significant gene expression changes (**Fig. 2e**). Overall, scFoundation outperformed scGPT, particularly in the
Sunshine2023 dataset. The AverageKnown method and GEARS exhibited comparable performance, with
average PCCs of 0.734 and 0.722, respectively. The Linear method and CellOracle showed generally poor
performance across these datasets. Though Linear-scGPT outperformed the Linear method, it still did not
surpass the AverageKnown method, raising questions about whether the gene embeddings from scGPT
genuinely aids this task. For the DEG_ACC metric, the AverageKnown method, scGPT, and scFoundation
demonstrated comparable performance, with average ACCs of 0.732, 0.726, and 0.705, respectively (**Fig. S1**).
Other methods, including GEARS, Linear-scGPT, Linear, and CellOracle, showed lower ACCs of 0.692,
0.689, 0.621, and 0.601, respectively, indicating limited effectiveness in predicting the direction of gene

expression changes. Additionally, single-cell foundation models demonstrated improved performance when focusing on higher-ranked DEGs, as shown in **Fig. 2f** and **S2**, suggesting that these models are better at capturing the most significant transcriptional changes.

For the SHG_MAE and SLG_MAE metrics, the mean method, scGPT, and scFoundation consistently performed well (**Fig. S1**). CellOracle performed slightly worse on the Sunshine2023 dataset for SHG_MAE, while GEARS showed slightly lower performance on both the Sunshine2023 and Norman2019 datasets for SLG_MAE. In contrast, linear-type methods performed poorly across both SHG_MAE and SLG_MAE.

Overall, single-cell foundation models, such as scGPT and scFoundation, emerged as the best-performing methods in the unseen perturbation transfer scenario. With sufficient perturbation data for training, they are capable of accurately predicting the expression changes of DEGs while preserving the expression levels of non-DEGs. Interestingly, the simplest approach, AverageKnown, also performed remarkably well across various datasets, including large ones like Replogle2022_K562. By visualizing the average expression changes of perturbations in this dataset using a clustering heatmap, we observed that more than half of the perturbed genes were grouped into a single cluster (**Fig. S3**). As a result, the AverageKnown method can achieve good performance by simply averaging these perturbed cells. GEARS also demonstrated potential as an advanced solution for unseen perturbations but performed slightly worse than the single-cell foundation models. In contrast, Linear, Linear-scGPT, and CellOracle consistently underperformed compared to deep learning-based methods.

Benchmarking in the unseen cell type transfer scenario

In the unseen cell type transfer scenario (**Fig. 3a**), we utilized the Replogle2022_K562 and Replogle2022_RPE1 datasets from the same study[4] for benchmarking. These datasets were generated using Perturb-seq experiments conducted on two distinct cell lines: K562, a lymphoblast cell line derived from a patient with chronic myelogenous leukemia, and RPE1, a retinal pigment epithelial cell line. After standardized data processing, we identified 490 shared gene perturbations and 3,144 overlapping genes between the two datasets.

For benchmarking, all models were trained on the shared perturbations from Replogle2022_K562 and then tested on the corresponding perturbations in Replogle2022_RPE1. Metrics were calculated based on the 3,144 overlapping genes (Methods). In this scenario, we selected scGen, CPA, CellOT, scGPT, and scFoundation as the benchmarking methods. Additionally, we included a simple baseline method, DirectTransfer, which directly transfers the gene expression changes of perturbations from Replogle2022_K562 to Replogle2022_RPE1 without additional model adjustments.

Before comparing different methods, we first assessed the similarity of shared perturbations between the two datasets. For each perturbation in each dataset, we calculated the gene expression alterations of averaged gene expression profiles between perturbed cells and unperturbed cells, and then calculated the PCCs of gene expression alterations (Alter_PCCs) for each perturbation in Replogle2022_K562 and Replogle2022_RPE1. The similarity provides a critical indicator of the difficulty in unseen cell type transfer prediction. As shown in **Fig. 3b**, the distribution of Alter_PCCs exhibits a bimodal pattern: a large group of perturbations clusters around 0.05, indicating divergent effects on gene expression between the two cell lines, while a smaller group clusters around 0.45, suggesting potentially similar effects. The distributions of Alter_PCC values suggest that

Fig. 3. Benchmarking in the unseen cell type transfer scenario. (a) Schematic diagram of unseen cell type transfer. **(b)** Density plot illustrating the distribution of PCCs of gene expression alterations across 490 common perturbations. **(c)** Scatter plots of 490 common perturbations, with the x-axis and y-axis represent log(E-distance) in the K562 and RPE1 cell lines, respectively. The perturbations are grouped into four categories (P1, P2, P3, and P4) based on log(E-distance) thresholds. **(d)** Box plots showing the PCCs of gene expression alterations (Alter_PCC) across different perturbation groups. Center line, median; box limits, upper and lower quartiles; whiskers, 1.5× interquartile range. **(e)** Radar plot comparing DEG_PCC, DEG_ACC, SHG_MAE, and SLG_MAE across the 490 perturbations for 6 methods. **(f)** Radar plots comparing DEG_PCC for 6 methods across 4 perturbation groups. **(g)** GO enrichment analysis for genes in each perturbation group.

these perturbations should be categorized into distinct groups for a more comprehensive benchmarking.

We further visualized each perturbation's E-distances in Replogle2022_K562 and Replogle2022_RPE1 (**Fig.**
3c). Using a log(E-distance) threshold of 3 in each dataset, we classified the 490 shared perturbations into four
sets (Fig. 3C): P1 (above the threshold in both datasets), P2 (above in K562 but below in RPE1), P3 (below in
K562 but above in RPE1), and P4 (below in both datasets). These classifications reflect the varying effects of
perturbations, which are associated with differing Alter_PCC values. For instance, P1 comprises perturbations
that induce substantial and similar phenotypic changes in both K562 and RPE1 cell lines, with a median
Alter_PCC value of 0.46—significantly higher than those observed in the other three sets (**Fig. 3d**).

We first evaluated all methods across overall 490 shared perturbations (**Fig. 3e**). Results showed that
DirectTransfer outperforms all other methods across various metrics, including DEG_PCC, DEG_ACC,
SHG_MAE, and SLG_MAE. For example, the mean DEG_PCC of DirectTransfer was 0.606, significantly
higher than the other five methods.

Notably, DirectTransfer achieved its highest performance in the P1 group (**Fig. 3f**). CellOT and CPA
demonstrated moderate performance in DEG_PCC and DEG_ACC, and CellOT showed a decline in
SHG_MAE and SLG_MAE. In contrast, scGPT, scFoundation, and scGen demonstrated poor performance
overall. Gene ontology (GO) enrichment analysis revealed that P1 perturbations are associated with
fundamental biological processes such as mRNA processing and RNA splicing (Fig. 3G), explaining their
significant and consistent effects across K562 and RPE1 cell lines. This indicates that predicting the effects of
P1 perturbations in unseen cell types is relatively easier for current methods.

Perturbations in P2 were associated with only five genes, such as *GAB2*, which is involved in the positive
regulation of leukocyte and mast cell degranulation (**Fig. 3g**), and this suggests genes in P2 may play an
important role in K562, but have a weak effect in RPE1. For perturbations in P2, scFoundation achieved
performance close to DirectTransfer in DEG_PCC, while scGen, CPA, CellOT, and scGPT yielded extremely
low DEG_PCC values (**Fig. 3f**), indicating that these models struggle to quantitatively capture the expression
changes of DEGs.

Perturbations in P3 were enriched in some biological processes related to protein production and secretion,
such as COPII vesicle coating and protein targeting to ER (**Fig. 3g**). This likely reflects the epithelial nature of
RPE1 cells, which are involved in more processing and transport of membrane and secretory proteins. On P3
perturbations, scGPT performed comparably to DirectTransfer across all metrics, whereas other methods
showed notably poorer performance (**Fig. 3f**).

Perturbations in P4 corresponded to genes with slight effects across both K562 and RPE1 cells. These
perturbations were enriched in some biological processes that are not predominant in cells, such as
mitochondrial translation and mitochondrial gene expression (**Fig. 3g**). DirectTransfer still maintained the best
overall performance on P4 perturbations, followed by scGPT and scFoundation (**Fig. 3f**).

All the above results showed that advanced machine learning or statistical methods failed to outperform the

simplest baseline, DirectTransfer, in the unseen cell type transfer scenario. While some methods have

demonstrated effectiveness in predicting specific treatment responses in unseen cell types, such as scGen

predicting the cellular response to IFN-β stimulation in CD4 T cells[11], and CellOT predicting responses of

patients with glioblastoma to panobinostat[15], these models struggled in the context of gene perturbations.

Transformer-based models, such as scGPT and scFoundation, generally outperformed autoencoder-based

methods, such as scGen, CPA, and CellOT, but still fell significantly short of the baseline performance of

DirectTransfer. This highlights the ongoing need for new, more effective methods tailored to the unseen cell

type transfer scenario. Furthermore, when applying *in silico* perturbation methods to predict effects on unseen

cell types, the choice of gene perturbations plays a crucial role in determining the results. Perturbing certain

housekeeping genes may elicit similar cellular responses across different cell types, as seen in the P1 and P4

groups, while other genes may have varying functional importance across cell types, as in P2 and P3,

depending on their specific roles across different cell types.

Benchmarking in the zero-shot transfer scenario

In practical applications, researchers often aim to apply *in silico* perturbation methods directly to their own

datasets, where training data from matching experimental settings with perturbations are typically unavailable.

To address this scenario, we established a benchmark for evaluating model performance in transferring to bulk

RNA-seq data (**Fig. 4a**).

We used CMAP data as the testing data for benchmarking. We collected single-cell datasets[27, 28] of six cell

lines that overlap with CMAP data: A375, PC3, MCF7, HT29, HepG2, and A549. Consistency between gene

expression profiles in the single-cell data and the corresponding CMAP data was verified (**Fig. 4b**, Methods).

These single-cell datasets were processed as inputs to *in silico* methods to directly predict gene expression

changes resulting from gene perturbations without any additional training. For each perturbation, we selected

significantly varied genes in CMAP data by setting a threshold of absolute z-scores (Methods), and these

genes consist of two equal parts: up-regulated genes (genes with positive z-scores) and down-regulated genes

(genes with negative z-scores). Model performance was evaluated using direction accuracy (ACC), calculated

by comparing the predicted results to the ground-truth labels from CMAP data (Methods). In total, 21,361

perturbations across the six cell lines were included in this benchmarking analysis (**Fig. 4c**).

We included several fundamental baselines as benchmarking methods, including the Correlation method and

DirectTransfer method (Methods). The Correlation method used PCCs among genes as associated change

factors for *in silico* perturbation; when a gene is perturbed, genes with higher PCCs relative to it are more

significantly affected (Methods). For DirectTransfer, we utilized two sources: genome-scale Perturb-seq data

on K562 cells (DirectTransfer_K562), and from the L1000 platform[26], where PC3 data was selected as the

Fig. 4. Benchmarking in the zero-shot transfer scenario. (a) Schematic diagram of zero-shot transfer. **(b)** Heatmap of correlations between single-cell data and CMAP data. **(c)** Pie chart illustrating the distribution of perturbations across six cell lines in the CMAP dataset. **(d)** Expression levels of CMAP's up-regulated and down-regulated genes in single-cell data. Up-regulated genes: genes that are up-regulated after perturbations in CMAP data; down-regulated genes: genes that are down-regulated after perturbations in CMAP data. **(e)** Bar plots comparing direction accuracy (ACC) across 9 methods for the 6 cell lines. Each bar represents the median ACC, with black bars indicating the upper and lower quartiles. The range of gray bars represent the upper quartile and lower quartile of the performance of Random. **(f)** Scatter plots showing the mean ACC of up-regulated and down-regulated genes. The gray dashed lines indicate the performance of Random for up-regulated and down-regulated genes.

transfer reference as it showed a high number of perturbations (DirectTransfer_L1000). Additionally, we

benchmarked deep learning methods such as CellOracle, CPA, scGPT, and scFoundation, which were trained

on a merged data collection of scPerturb datasets we curated (Methods). To provide a comparison baseline, a

random prediction approach (referred to as Random) was included, where genes are randomly upregulated or

downregulated.

We observed a distinctive expression pattern of CMAP's significant varied genes in single-cell data: up-

regulated genes in CMAP tend to exhibit low expression levels in single-cell data, whereas down-regulated

genes in CMAP often display high expression levels in single-cell data. As shown in **Fig. 4d** and **Fig. S4**,

across all the cell lines, this trend is consistent across all analyzed cell lines. Building on this observation, we

introduced an AverageSC method for this scenario, which simply averages the expression values of all

landmark genes in single cells to generate predictions. Additionally, we found that single-cell foundation

models often predict lower gene expression values relative to the ground-truth values (**Fig. S5**). This bias will

complicate the accurate assessment of the model performance. To mitigate this issue, we proposed a strategy

for single-cell foundation models that involves using predictions for unperturbed cells as a reference

(Methods), aiming to more precisely capture perturbation effects.

We visualized the performance of each method on individual cell lines (**Fig. 4e**) and summarized the overall

accuracies for predicting both up-regulated genes and down-regulated genes for all methods (**Fig. 4f**).

CellOracle failed to predict the direction of gene expression changes across all the cell lines. Similarly, CPA

performed poorly, with results slightly worse than Random in certain cell lines, such as HepG2. The

Correlation method showed marginal improvement over Random (**Fig. 4f**). Among the DirectTransfer

methods, DirectTransfer_K562 performed at a Random level, while DirectTransfer_L1000 achieved the

highest overall accuracy, with a mean value of 0.692. This suggested that transferring gene expression changes

from the same platform, such as the L1000 platform, can provide a reliable reference. Both scGPT and

scFoundation showed better performance than Random (**Fig. 4f**). Specifically, scGPT performed better than

Random in cell line HT29, HepG2, and A549, but performed at Random levels in other cell lines. In contrast,

scFoundation outperformed Random in A375, PC3, HT29, and A549. Interestingly, scGPT demonstrated

higher accuracy for predicting up-regulated genes, while scFoundation performed better for down-regulated

genes (**Fig. 4f**). The AverageSC method achieved the best performance aside from DirectTransfer_L1000,

indicating that leveraging the inherent patterns in perturbation data can yield strong predictive results.

Overall, only DirectTransfer_L1000 and the AverageSC method achieved significantly better performance

than Random. However, DirectTransfer_L1000 relies on data from the same platform, which is typically

unavailable in practical applications, limiting its utility. Meanwhile, the AverageSC method, which merely

averages gene expression values, does not account for gene perturbation mechanisms and is therefore

impractical for understanding or modeling perturbation effects. Most other methods failed to predict the

direction of change for significant genes, highlighting the limitations of current approaches in predicting

perturbation effects in zero-shot scenarios. While single-cell foundation models demonstrated some potential

to exceed Random performance, their best accuracy on an individual cell line was approximately 0.55—only

marginally above random and still far from practical applicability.

We further evaluated the performance of scGPT with varying scales of total perturbations in the merged data

collection (Methods), using 10%, 40%, 70%, and 100% of the perturbations. Results indicated a clear scaling

effect, wich scGPT achieving progressively higher performance as the number of perturbations increased (**Fig.**

S6). We anticipated that as more single-cell perturbation data becomes available across a wider range of cell

types and genome-scale gene perturbations, the performance of single-cell foundation models is likely to

improve. However, for now, quantitatively predicting perturbation effects in a zero-shot transfer scenario

using *in silico* perturbation methods remains a significant challenge.

Benchmarking in the cell state transition prediction scenario

Fig. 5. Benchmarking in the cell state transition prediction scenario. (a), Ground-truth cell state transitions resulting from the knockdown or overexpression of specific genes. Transition directions are visualized in UMAP space. **(b-f)** Streamline plots display the predicted results from 6 methods across five cases: recovery of exhausted T cells with *PDCD1* knockdown (b), induction of ADM with *PTF1A* knockdown (c)**,** MEP transitioning to LMPP with *GATA1* knockdown (d), GMP transitioning to LMPP with *SPI1* knockdown (e), and fibroblasts reprogrammed into stem cells by four TFs: POU5F1, SOX2, KLF4, and MYC (f).

Many biological processes, such as organ development and cellular reprogramming, are driven by individual

genes or small sets of key genes. Evaluating whether computational methods can accurately predict cell state

transitions influenced by these genes is critical for assessing their applicability to researchers' own data and

for guiding experimental design.

To establish a benchmarking framework, we collected four single-cell datasets representing distinct biological

processes (Methods): Sun2022[29] for validating that the knockout of *PDCD1* can promote the recovery of

exhausted T cells, restoring their effector and memory functions[40, 41]; Ainciburu2023[30] for validating that SPI1

and *GATA1* perform as a canonical two-gene toggle-switch motif driving the differentiation of hematopoietic

cells[6, 42], where SPI1 is demonstrated to promote the granulocyte-monocyte progenitor (GMP) lineage, and

GATA1 commits to the megakaryocyte-erythrocyte progenitor (MEP) lineage; Nair2023[31] for validating

fibroblasts can be reprogrammed into stem cells by four transcription factors (TFs): POU5F1, SOX2, KLF4,

and MYC[43]; Steele2020[32] for validating the knockout of *PTF1A*[44, 45]can induce acinar cells transforming into

duct-like cells (acinar-ductal metaplasia, ADM). UMAP visualization of these datasets, along with the ground-

truth cell state transitions, was shown in **Fig. 5a**. To evaluate performance, we visualized cell transition trends

after *in silico* gene perturbations using streamline plots (**Fig. 5b-f**, Methods).

We benchmarked the methods same as in the zero-shot transfer scenario: DirectTransfer, the Correlation

method, CellOracle, CPA, scGPT, and scFoundation. It should be noted that not all methods can be applied to

all four datasets. For instance, CellOracle is restricted to *in silico* perturbations of TFs with known binding

motifs, and *PDCD1* in the Sun2022 dataset is not included in its TF list. Similarly, CPA and DirectTransfer

rely on reference data with the matching perturbations, which were unavailable for the Nair2023 dataset and

for PTF1A in the Steele2020 dataset, as these perturbed genes are not included in our filtered scPerturb

datasets.

The cell states in the Sun2022 and Steele2020 dataset exhibited discrete cell states, while cells in

Ainciburu2023 and Nair2023 form continuous trajectories. Accurately predicting transitions in datasets with

discrete cell states requires models to capture more substantial changes in cell states following gene

perturbation. As shown in the UMAP plots of both pre- and post-perturbation cells (**Fig. S7**), the predicted

gene expression changes are subtle, with cells tending to remain in their original states. In streamline plots,

most methods fail to capture the correct transition trends (**Fig. 5b-c**).

In the Ainciburu2023 dataset, both the Correlation method and CellOracle successfully predicted the

transitions from MEP to LMPP following *GATA1* knockout, and from GMP to LMPP following *SPI1*

knockout (**Fig. 5d-e**). However, other methods failed to capture these specific lineage transitions. In the

Nair2023 dataset, CellOracle provided the most accurate visualization of the transition from fibroblast-like

cells to stem-like cells in the streamline plot (**Fig. 5f**). scGPT also exhibited an accurate trend in the plot,

though it was less pronounced. In contrast, the Correlation method misrepresented the transition, erroneously

depicting fibroblast-like cells shifting to keratinocyte-like cells instead of stem-like cells.

Additionally, it is worth noting that CPA failed to predict cell state transitions across all datasets. This

limitation may be due to CPA's reliance on reference data, which can introduce batch effects, resulting in discrepancies between the predicted and actual cell states (**Fig. S7**). Among the single-cell foundation models, scGPT consistently outperformed scFoundation across all datasets.

In summary, predicting transitions between discrete cell states remains a significant challenge for current methods. For cell states within continuous trajectories, methods that leverage gene expression correlations, such as the Correlation method and CellOracle, which employed a regression model to infer relationships among genes, perform well in capturing the overall transition trends. Conversely, methods that rely on reference training data often struggle to capture accurate cell state transitions, likely due to discrepancies in cell types or experimental platforms between the reference data and the target dataset. Single-cell foundation models, which are trained on large datasets, show promising results in certain cases. For instance, scGPT demonstrates notable performance on the Nair2023 dataset. However, their performance may still be limited by the characteristics of the dataset and the complexity of the transitions being modeled. While single-cell foundation models like scGPT hold potential, they are not yet universally applicable and require further refinement to achieve broader applicability in diverse scenarios.

Discussion

Our benchmark categorized *in silico* perturbation methods into four distinct usage scenarios. We observed that no single method consistently excels across all scenarios, highlighting the need of tailoring approaches to specific applications. Single-cell foundation models demonstrated strong performance in predicting unseen perturbations while the DirectTransfer method, despite its simplicity, outperformed others in the unseen cell type transfer scenario. For the cell state transition prediction scenario, methods like the Correlation method and CellOracle proved to be the most effective. These findings underscore the importance of aligning method development with the intended usage scenario. Clearly defining the applicable scenarios for each method can not only guide their improvement but also help researchers select the most suitable approach for their specific needs.

One of the most appealing aspects of *in silico* perturbations is that researchers can easily apply them to their data of interest and conduct virtual perturbation experiments. However, due to current data limitations, most usage scenarios of *in silico* gene perturbations still rely on zero-shot predictions, where no training data is provided. Our benchmarking results reveal that existing methods barely outperform Random predictions in terms of quantitatively predicting gene expression changes in zero-shot predictions.

Although achieving reliable predictions with *in silico* perturbation methods remains a long-term challenge, advancements in this field offer hope for future progress. With the ongoing development of biological technologies, single-cell perturbation experiments are expected to cover a broader range of cell types and perturbations. The gradually accumulation of such data will likely enhance the performance of computational

methods. Rood el al.[46] proposed a concept of "perturbation cell atlas", but how to curate and build such an

atlas remains open challenges. Beyond data, innovation in model architectures is also essential. Current

transformer-based models show promise in leveraging massive datasets, but there is considerable potential for

exploring alternative architectures. For instance, recent studies have highlighted the potential of diffusion

models for performing *in silico* perturbations[47], offering a promising direction for further research and

development in this domain.

While experimental data can cover individual perturbations at the genome scale on a single cell type,

extending these experiments to cover all possible perturbation combinations remains infeasible. *In silico*

methods present a promising solution to guide the selection of perturbation combinations for experimental

validation. Several studies have attempted to benchmark *in silico* methods for predicting the effects of

perturbation combinations[8, 9, 18, 48]. However, we believe that the currently available single-cell perturbation

data involving simultaneous perturbation of multiple genes is insufficient for a comprehensive benchmarking.

With the accumulation of more extensive and diverse data in the future, we plan to incorporate this important

benchmark to address the growing need for evaluating combination perturbation predictions.

Our benchmark focuses exclusively on *in silico* methods designed for scRNA-seq data, as it is the most

abundant resource for single-cell perturbation studies and serves as the foundation for the development of

most existing methods. Single-cell perturbation methods for sequencing other modalities, such as chromatin

accessibility[49, 50] and protein abundance[51], have also been developed. We anticipate that as datasets for these

modalities continue to grow, *in silico* methods capable of predicting molecular profiles beyond RNA

expression will emerge. These advancements could enable predictions of chromatin states, protein abundance,

and other intracellular molecular features, paving the way for a deeper understanding of cellular systems and

facilitating the design of more precise and robust perturbation experiments.

To support researchers in evaluating methods with our benchmarking framework, we developed a Python

module for users to easily use our benchmark. For each usage scenario, we implemented intuitive and flexible

interfaces that allow seamless access to benchmarking datasets and straightforward computation of

benchmarking metrics. Additionally, we included examples of each *in silico* perturbation method

benchmarking in this study, offering clear guidance for training and evaluating their own models. These

resources aim to streamline the benchmarking process and promote method development. We believe this

work will serve as a foundation for advancing *in silico* perturbation methods and will contribute to progress in

the field of biomedical research.

Methods

Data curation

scPerturb data

We utilized the scPerturb scRNA-seq datasets (version 1.3, available at https://zenodo.org/records/10044268) as benchmarking data for our study. scPerturb is a comprehensive resource of single-cell perturbation datasets, encompassing a wide range of perturbations, including gene modifications, small molecules, cytokines, and more. This dataset includes a total of 50 RNA and protein datasets. To ensure consistency and relevance, we excluded datasets that do not map CRISPR perturbations to specific genes and those derived from non-human cells.

After applying these filtering steps, we retained 17 single-cell perturbation datasets for analysis, each representing cells of a specific cell type perturbed by various genes. To prepare the data for model training and evaluation, we performed the following preprocessing steps:

- Feature Selection: We identified the top 5,000 highly variable genes (HVGs) from each dataset using standard procedures. We also included all perturbed genes originally present in the dataset's feature list.

- Normalization: Raw count data was normalized using the log-normalization method, implemented through the standard preprocessing workflow of Scanpy[52].

- Control-cell pairing: For every cell subjected to perturbation, we randomly sampled a control cell from the same dataset to form a pre- and post-perturbation cell pair. These pairs served as the basis for training and evaluating models.

- Perturbation filtering based on cell thresholds: Perturbations with fewer than 100 cells were excluded to ensure robust analysis with sufficient data.

- E-distance calculation: The E-distance for each perturbation was computed to quantify the perturbation's effect. E-distance is a statistical measure of distance between two distributions, which is utilized to indicate the degree of perturbation effect in scPerturb. We utilized the scPerturb Python library to compute this metric between control cells and perturbed cells. For each perturbation, we sampled 200 cells and selected 2,000 HVGs, as instructed by the scPerturb documentation, to perform the calculation.

- sgRNA effectiveness filtering: To address variability in sgRNA efficacy, we retained only sgRNAs that demonstrated consistent perturbation effects. Specifically, if an sgRNA's E-distance exceeded 10 and was at least 10-fold greater than another sgRNA targeting the same gene, this sgRNA was included, while the underperforming sgRNA was excluded.

- Identifying differentially expressed genes (DEGs): For each perturbation, we identified DEGs by comparing perturbed cells to control cells. Genes were ranked based on adjusted p-values computed using the Wilcoxon test in Scanpy, with those having adjusted p-values < 0.05 designated as DEGs.

- Data splitting for training, testing, and validation: The filtered perturbations were divided into training,

testing, and validation sets in a 7:2:1 ratio. The training set was used to fit the models, the validation set
to select the best model, and the testing set to evaluate the final model performance.

CMAP data

CMAP data is based on L1000, a high-throughput gene expression profiling technology that converts raw
fluorescence intensity of genes into expression levels. CMAP data contain over 1.5 million gene expression
profiles derived from approximately 3,000 gene perturbations and 5,000 drug perturbations across multiple
cell lines. For each perturbation in each cell line, CMAP provides a normalized z-score vector of 978
landmark genes, representing the extent to which these genes are affected by the perturbation in the given cell
line. This dataset is a valuable resource for understanding gene expression changes induced by perturbations.

The CMAP data is organized into five distinct levels, each representing a different stage of data processing:
Level 1 records the raw fluorescence intensity measurements. Level 2 and level 3 provide deconvoluted and
inferred gene expression values, respectively. Level 4 provides robust z-scores derived from the expression
data. Level 5 aggregates the z-scores, offering a comprehensive summary of perturbation effects. In our study,
we used level 5 data as the benchmark ground truth due to its aggregated and robust representation of gene
expression changes.

To align the CMAP data with single-cell datasets, we collected single-cell data with six overlapping cell lines:
A375, PC3, MCF7, HT29, HepG2, and A549. These single-cell datasets were used as control cell inputs for
model evaluation. Each single-cell dataset underwent the following preprocessing steps: (1) 3,000 HVGs were
selected, with perturbation genes from the CMAP data added to the feature set, and (2) raw count data was
log-normalized.

For each cell line, we calculated the PCC between the single-cell data and the CMAP data. For the single-cell
data, we selected 3,000 HVGs from the single-cell data and averaged the expression levels across all cells to
obtain a mean expression vector. For the CMAP data, we extracted control samples corresponding to the
specific cell line and averaged them to a mean expression vector with a length of 978. The PCC was then
computed using the genes common to both mean expression vectors.

For each perturbation in the CMAP dataset, the level 5 data provide an aggregated z-score vector representing
the gene expression changes. Positive z-scores indicate increased expression following the perturbation (up-
regulated), while negative z-scores indicate decreased expression (down-regulated). To refine the analysis, we
selected genes with z-scores greater than 1.5 or less than –1.5 as significantly affected and excluded genes
with an expression level less than 0.1 in the single-cell datasets. From this filtered gene set, we further
sampled genes to ensure an equal representation of positive and negative z-scores, balancing the directionality
of expression changes. These genes were identified as top significant genes, and we calculated direction
accuracy using these genes to evaluate model performance.

Single-cell data with cell state transitions

We compiled four single-cell datasets encompassing five distinct biological processes for our study. To ensure
consistency and minimize batch effects, we selected data from a single donor for datasets with multiple
donors.

• Sun2022 dataset: This dataset originates from cells within the tumor microenvironment of gastric cancer.
We selected donor GC05 and retained three CD8+ T cell subsets: exhausted CD8+ T cells (Tex), effector
memory CD8+ T cells (Tem), and residual memory CD8+ T cells (Trm). The biological process of
interest involves the knockdown of *PDCD1*, which is expected to induce a transition from Tex to Tem.

• Ainciburu2023 dataset: Derived from hematopoietic cells in a human cell atlas, this dataset includes data
from the 19-year-old donor with the following cell types: lympho-myeloid primed progenitors (LMPPs),
granulocyte-monocyte progenitors (GMPs), megakaryocyte-erythrocyte progenitors (MEPs),
hematopoietic stem cells (HSCs), and plasmacytoid dendritic cells (pDCs). Two biological transitions
were investigated: knockdown of GATA1 leading to MEP-to-LMPP transition, and knockdown of SPI1
leading to GMP-to-LMPP transition.

• Nair2023 dataset: This dataset originates from a fibroblast reprogramming experiment. We included three
cell types for analysis: fibroblast-like cells, keratinocyte-like cells, and stem-like cells. The focus is on
the reprogramming of fibroblast-like cells into stem-like cells via the overexpression of canonical
reprogramming factors: POU5F1, SOX2, KLF4, and MYC.

• Steele2020 dataset: This dataset was obtained from a human pancreas cell atlas, using data from donor
N1. We selected acinar cells, ductal cells, and stellate cells as cell types of interest. The biological
process studied is Acinar-Ductal Metaplasia (ADM), a transition from acinar cells to ductal cells, which
can be induced by the knockdown of *PTF1A*.

All datasets were processed using a standardized pipeline, which included log-normalization of the count data
followed by the selection of HVGs using Scanpy with parameters set to min_mean=0.0125, max_mean=3,
and min_disp=0.4.

564 Implement of *in silico* perturbation methods

565 In this study, we evaluated 10 *in silico* perturbation methods across four distinct benchmarking scenarios.
566 Some of the methods and applicable to multiple scenarios. For each method, parameter settings were fixed
567 according to official guidelines and tailored to align with our benchmarking datasets.

568 Linear

569 The linear method is adopted from another benchmarking study[22]. The optimization objective is

$$\arg \min_{\mathbf{W}} ||\mathbf{Y}^{\text{train}} - (\mathbf{GWP}^T + \boldsymbol{b})||_2^2, \tag{1}$$

where $\mathbf{Y}^{\text{train}}$ is the gene expression matrix of perturbed cells, rows represent the gene feature set, and columns represent the perturbed genes in the training data. $\mathbf{G}$ is the gene embedding matrix, which is obtained from the top k principal components from the PCA (principal component analysis) on $\mathbf{Y}^{\text{train}}$. $\mathbf{P}$ is subset from $\mathbf{G}$ with the perturbed genes in the training data. $\mathbf{W} \in \mathbb{R}^{k \times k}$ is the fitted matrix, and $\boldsymbol{b}$ is the vector of average gene expressions of $\mathbf{Y}^{\text{train}}$: $\boldsymbol{b} = \frac{1}{N} \sum_{i=1}^{N} \mathbf{Y}_{:i}^{train}$. $\mathbf{W}$ is solved using the normal equations:

$$\mathbf{W} = (\mathbf{G}^T \mathbf{G} + \lambda \mathbf{I})^{-1} \mathbf{G}^T (\mathbf{Y}^{\text{train}} - \boldsymbol{b}) \mathbf{P} (\mathbf{P}^T \mathbf{P} + \lambda \mathbf{I})^{-1}. \tag{2}$$

Here, we set k to 512 and λ to 0.1. Then the fitted $\mathbf{W}$ is used for prediction:

$$\widehat{\mathbf{Y}} = \boldsymbol{b} + \mathbf{GWP}^T, \tag{3}$$

and $\widehat{\mathbf{Y}}$ is the predicted gene expression, and $\mathbf{P}$ subset from $\mathbf{G}$ denotes the embeddings of perturbed genes in the testing data.

Linear with scGPT embeddings

Linear-scGPT employs the same modeling approach as the linear method, with the key distinction being that the gene embedding matrix $\mathbf{G}$ is derived from the scGPT model rather than the PCA components of the training data. We followed the instruction from GitHub repository of the Linear method to obtain the scGPT gene embeddings: https://github.com/const-ae/linear_perturbation_prediction-Paper/blob/main/benchmark/src/extract_gene_embedding_scgpt.py.

Correlation

We first constructed a gene-gene correlation matrix, where each element represents the PCC between two genes. When a gene is perturbed, the expressions of other genes change according to a scaling factor determined by the corresponding PCCs. If a gene is knocked down, its expression is set directly to 0. If a gene is overexpressed, 1 is added to its original normalized expression value.

CellOracle

We followed the tutorial provided on the CellOracle website at https://morris-lab.github.io/CellOracle.documentation/tutorials/simulation.html. For each dataset, the GRN was inferred from control cells and subsequently applied to simulate expected gene expression changes under gene perturbations. Since scATAC-seq data from the same tissue as the scRNA-seq datasets is unavailable, we utilize the function 'load_human_promoter_base_GRN' to provide base GRNs for all datasets.

scGen

We followed the instructions provided on the scGen website via

600   https://scgen.readthedocs.io/en/stable/tutorials/scgen_perturbation_prediction.html. For each perturbation

601   transferred to an unseen cell type, we train a model for prediction, similar to the procedures used for CPA and

602   CellOT.

CPA

604   We followed the tutorial provided on the CPA website: https://cpa-

605   tools.readthedocs.io/en/latest/tutorials/Kang.html. While CPA was originally designed for the unseen cell type

606   transfer scenario, we adapted it for application in zero-shot transfer and cell state transition prediction

607   scenarios. In these cases, the training data was sourced from the scPerturb datasets.

CellOT

609   We adopted the codes provided on the CellOT GitHub repository: https://github.com/bunnech/cellot. Since

610   our control group data contains significantly fewer cells compared to the tutorial dataset, we reduced 'n_iters'

611   from 100,000 to 1,000 to improve computational efficiency.

GEARS

613   We followed the tutorials provided on the GEARS Github repository: https://github.com/snap-

614   stanford/GEARS. GEARS is designed for the unseen perturbation scenario. We first reformatted our data to

615   align with GEARS' input requirements. The model was then trained using its default parameter settings, with

616   the number of epochs to 20. The best-performing model, selected based on its performance on the validation

617   dataset, was used to evaluate the test dataset.

scGPT

The scGPT model is versatile in various scenarios for its ability to accept any length of genes as input. In

unseen perturbation transfer and unseen cell type transfer scenarios, we followed the guidance on scGPT

Github repository to reformat data and train models: https://github.com/bowang-

lab/scGPT/blob/main/tutorials/Tutorial_Perturbation.ipynb. Default parameter settings were used, except that

the model was trained with all genes in the feature set. Similar to GEARS, we trained the scGPT model for 20

epochs and selected the best-performing model based on validation results for testing. For zero-shot transfer

and cell state transition scenarios, the scGPT model was trained on a combined dataset comprising a total of

17 datasets, ensuring broader generalization across different biological contexts.

scFoundation

scFoundation also employs transformer-based architectures, making it suitable for *in silico* perturbation tasks,

similar to the training paradigm of scGPT. We adopted the code from scFoundation Github repository:

https://github.com/biomap-research/scFoundation. Specifically, we leveraged the encoder module of the

scFoundation model to train the perturbation model. The training and testing procedures followed the same

setup as in scGPT. Given that scFoundation incorporates more transformer layers than scGPT, we set the

learning rate to 1e-5.

Training and application of single-cell foundation models for the zero-shot transfer scenario

We curated 17 processed scPerturb datasets to serve as the training data for scGPT and scFoundation. Each

dataset was carefully processed and split into training, testing, and validation subsets. These subsets were

subsequently merged across all datasets to create comprehensive training, testing, and validation datasets. The

models were trained on the combined training set using random shuffling to ensure robust learning.

Considering that the cells in the same batch may come from different datasets and hold gene features with

different lengths, we performed the data padding with 'MASK' token, which are not involved in training

models. For both scGPT and scFoundation, we trained the model for 20 epochs and selected the best-

performing models as the final model for zero-shot transfer.

We observed that when trained single-cell foundation models are applied to novel datasets, the predicted gene

expression values tend to be lower than the ground-truth gene expression values (Fig. S5). To address this

bias, we employed a correction strategy using control cells. Specifically, let the gene expression of control cell

i be denoted as x_i, and the model's predicted output with perturbation as y_i. We then used the model to

predict the gene expression of the same control cell without perturbation, denoted as y_i'. The difference $y_i -$

y_i' was taken as the final predicted gene expression change.

Benchmarking metrics

We utilized different metrics in different scenarios to comprehensively and fairly evaluate all the *in silico*

perturbation methods. In unseen perturbation transfer and unseen cell type transfer scenarios, the single-cell

perturbation datasets from scPerturb are adopted for model training and testing. For each perturbation, we

denote $x_i \in \mathbb{R}^n$ and $y_i \in \mathbb{R}^n$ as average gene expression vectors of control cells and perturbed cells,

respectively, and n denotes the length of gene feature set.

Pearson correlation coefficient (PCC)

The PCC value is calculated by the following equation:

658
$$\text{PCC}(\Delta \boldsymbol{x}_i, \Delta \widetilde{\boldsymbol{x}}_i) = \frac{E[(\Delta \boldsymbol{x}_i - \mu_i)(\Delta \widetilde{\boldsymbol{x}}_i - \widetilde{\mu}_i)]}{\sigma_i \, \widetilde{\sigma}_i},\tag{4}$$

659   where $\Delta \boldsymbol{x}_i$ and $\Delta \widetilde{\boldsymbol{x}}_i$ represent the ground-truth and predicted changed expression vectors of perturbation i,

660   respectively. Here, μ_i and $\widetilde{\mu}_i$ denote the mean values of $\Delta \boldsymbol{x}_i$ and $\Delta \widetilde{\boldsymbol{x}}_i$, while σ_i and $\widetilde{\sigma}_i$ are their

661   corresponding standard deviations. PCC values are computed specifically for DE genes in each perturbation.

662   **Direction accuracy (ACC)**

663   We calculated ACC by the following equation:

$$\text{ACC}(\Delta \boldsymbol{x}_i, \Delta \tilde{\boldsymbol{x}}_i) = \frac{\Sigma_{i=1}^{n} \mathbb{I}\big(\text{sign}(\Delta x_i) - \text{sign}(\Delta \tilde{x}_i)\big)}{n}, \tag{5}$$

665   where $\mathbb{I}(\cdot)$ is an indicator function defined as:

$$\mathbb{I}(\text{x}) = \begin{cases} 1, & x = 0, \\ 0, & else. \end{cases} \tag{6}$$

In the unseen perturbation transfer and unseen cell type transfer scenarios, ACC values are calculated for
DEGs. While in the zero-shot transfer scenario, the ACC values are computed for the top significant genes
from the CMAP dataset.

Mean absolute error (MAE)

The MAE is calculated for the non-DEGs, including stably highly expressed genes (SHGs) and stably lowly
expressed genes (SLGs) for each perturbation, and it is defined as the following equation:

$$\text{MAE}(\Delta \boldsymbol{x}_i, \Delta \tilde{\boldsymbol{x}}_i) = \frac{\Sigma_{i=1}^{n} |\Delta x_i - \Delta \tilde{x}_i|}{n}. \tag{7}$$

Streamline plot

The model's predicted expression change vector was used as the cellular velocity for each cell. Subsequently,
the 'velocity_embedding_stream' function from the scVelo[53] package was employed to visualize the
streamlines on the UMAP plot.

Data availability

The scPerturb datasets were downloaded from https://zenodo.org/records/10044268. The CMAP data was
downloaded from Gene Expression Omnibus (GEO) with accession number GSE92742. The scRNA-seq
datasets containing 6 cell lines intersected with CMAP data were obtained from the Broad Institute's single-
cell portal (SCP542) and Sequence Archive (CNSA) with accession number CNP0003658. scRNA-seq
datasets used in the cell state transition prediction scenario were downloaded from different resources:
Sun2022 dataset (OMIX001073), Ainciburu2023 dataset (GSE180298), Nair2023 dataset (GSE242424), and
Steele2020 dataset (GSE155698).

Acknowledgements

The work is supported in part of National Natural Science Foundation of China (62250005, 62373210,
62433001, and 92470105) and The National Key R&D Program of China (2021YFF1200900).

Author contributions

C.L., H.G. and X.Z. conceived and designed the study. X.Z, K.L and Y.S. supervised the study and provided

critical revisions. C.L. and H.G. collected the data. C.L. implemented the methods, conducted the

experiments, and analyzed the results. C.L. and L.W. drafted the manuscript and prepared the figures. X.Z.,

K.L., H.G., and Y.S. reviewed the manuscript and provided critical insights. All authors contributed to

reviewing and approving the final manuscript.

695 Competing interests

696 K.L., H.G., and Y.S. are employees of Anew Therapeutics Pte. Ltd. C.L. and H.B. contributed to this work

697 while interning at Anew Therapeutics Pte. Ltd. The remaining authors declare no competing interests.

698 Reference

1. Dixit, A. et al. Perturb-Seq: Dissecting Molecular Circuits with Scalable Single-Cell RNA Profiling of
Pooled Genetic Screens. *Cell* **167**, 1853-1866 e1817 (2016).

2. Datlinger, P. et al. Pooled CRISPR screening with single-cell transcriptome readout. *Nat Methods* **14**,
297-301 (2017).

3. Adamson, B. et al. A Multiplexed Single-Cell CRISPR Screening Platform Enables Systematic
Dissection of the Unfolded Protein Response. *Cell* **167**, 1867-1882 e1821 (2016).

4. Replogle, J.M. et al. Mapping information-rich genotype-phenotype landscapes with genome-scale
Perturb-seq. *Cell* **185**, 2559-2575 e2528 (2022).

5. Hatton, I.A. et al. The human cell count and size distribution. *Proc Natl Acad Sci U S A* **120**,
e2303077120 (2023).

6. Qiu, X. et al. Mapping transcriptomic vector fields of single cells. *Cell* **185**, 690-711 e645 (2022).

7. Kamimoto, K. et al. Dissecting cell identity via network inference and in silico gene perturbation.
Nature **614**, 742-751 (2023).

8. Roohani, Y., Huang, K. & Leskovec, J. Predicting transcriptional outcomes of novel multigene
perturbations with GEARS. *Nat Biotechnol* **42**, 927-935 (2024).

9. Gao, Y. et al. Toward subtask-decomposition-based learning and benchmarking for predicting genetic
perturbation outcomes and beyond. *Nat Comput Sci* **4**, 773-785 (2024).

10. Bai, D., Ellington, C.N., Mo, S., Song, L. & Xing, E.P. AttentionPert: accurately modeling
multiplexed genetic perturbations with multi-scale effects. *Bioinformatics* **40**, i453-i461 (2024).

11. Lotfollahi, M., Wolf, F.A. & Theis, F.J. scGen predicts single-cell perturbation responses. *Nat
Methods* **16**, 715-721 (2019).

12. Lotfollahi, M. et al. Predicting cellular responses to complex perturbations in high-throughput
screens. *Mol Syst Biol* **19**, e11517 (2023).

13. Lopez, R. et al. Learning causal representations of single cells via sparse mechanism shift modeling
[C]. Proceedings of the Conference on Causal Learning and Reasoning. PMLR, 2023: 662-691.

14. Bereket, M. & Karaletsos, T. Modelling Cellular Perturbations with the Sparse Additive Mechanism
Shift Variational Autoencoder [C]. Proceedings of the 37th Conference on Neural Information
Processing Systems (NeurIPS). 2023.

15. Bunne, C. et al. Learning single-cell perturbation responses using neural optimal transport. *Nat

728   *Methods* **20**, 1759-1768 (2023).

729   16.   Dong, M. et al. Causal identification of single-cell experimental perturbation effects with CINEMA-
730   OT. *Nat Methods* **20**, 1769-1779 (2023).

731   17.   Cui, H. et al. scGPT: toward building a foundation model for single-cell multi-omics using generative
732   AI. *Nat Methods* **21**, 1470-1480 (2024).

733   18.   Hao, M. et al. Large-scale foundation model on single-cell transcriptomics. *Nat Methods* **21**, 1481-
734   1491 (2024).

735   19.   Csendes, G., Szalay, K.Z. & Szalai, B.J.b. Benchmarking a foundational cell model for post-
736   perturbation RNAseq prediction. *bioRxiv* (2024).

737   20.   Wu, Y. et al. PerturBench: Benchmarking Machine Learning Models for Cellular Perturbation
738   Analysis. *arXiv* (2024).

739   21.   Wenteler, A. et al. PertEval-scFM: Benchmarking Single-Cell Foundation Models for Perturbation
740   Effect Prediction. *bioRxiv* (2024).

741   22.   Ahlmann-Eltze, C., Huber, W. & Anders, S.J.B. Deep learning-based predictions of gene perturbation
742   effects do not yet outperform simple linear methods. *bioRxiv* (2024).

743   23.   Hall, C.C. The hazards of external version. *West J Surg Obstet Gynecol* **61**, 239-261 (1953).

744   24.   Deng, J. et al. ImageNet: A Large-Scale Hierarchical Image Database [C]. Proceedings of the IEEE-
745   Computer-Society Conference on Computer Vision and Pattern Recognition Workshops. 2009: 248-
746   255.

747   25.   Peidli, S. et al. scPerturb: harmonized single-cell perturbation data. *Nat Methods* **21**, 531-540 (2024).

748   26.   Subramanian, A. et al. A Next Generation Connectivity Map: L1000 Platform and the First 1,000,000
749   Profiles. *Cell* **171**, 1437-1452 e1417 (2017).

750   27.   Kinker, G.S. et al. Pan-cancer single-cell RNA-seq identifies recurring programs of cellular
751   heterogeneity. *Nat Genet* **52**, 1208-1218 (2020).

752   28.   Zhu, Q. et al. Single cell multi-omics reveal intra-cell-line heterogeneity across human cancer cell
753   lines. *Nat Commun* **14**, 8170 (2023).

754   29.   Sun, K. et al. scRNA-seq of gastric tumor shows complex intercellular interaction with an alternative
755   T cell exhaustion trajectory. *Nat Commun* **13**, 4943 (2022).

756   30.   Ainciburu, M. et al. Uncovering perturbations in human hematopoiesis associated with healthy aging
757   and myeloid malignancies at single-cell resolution. *Elife* **12** (2023).

758   31.   Nair, S. et al. Transcription factor stoichiometry, motif affinity and syntax regulate single-cell
759   chromatin dynamics during fibroblast reprogramming to pluripotency. *bioRxiv* (2023).

760   32.   Steele, N.G. et al. Multimodal Mapping of the Tumor and Peripheral Blood Immune Landscape in
761   Human Pancreatic Cancer. *Nat Cancer* **1**, 1097-1112 (2020).

762   33.   Datlinger, P. et al. Ultra-high-throughput single-cell RNA sequencing and perturbation screening with
763   combinatorial fluidic indexing. *Nat Methods* **18**, 635-642 (2021).

764   34.   Shifrut, E. et al. Genome-wide CRISPR Screens in Primary Human T Cells Reveal Key Regulators of
765   Immune Function. *Cell* **175**, 1958-1971 e1915 (2018).

766   35.   Tian, R. et al. Genome-wide CRISPRi/a screens in human neurons link lysosomal failure to
767   ferroptosis. *Nat Neurosci* **24**, 1020-1034 (2021).

768   36.   Tian, R. et al. CRISPR Interference-Based Platform for Multimodal Genetic Screens in Human iPSC-
769   Derived Neurons. *Neuron* **104**, 239-255 e212 (2019).

37.   Xu, Z., Sziraki, A., Lee, J., Zhou, W. & Cao, J. Dissecting key regulators of transcriptome kinetics
through scalable single-cell RNA profiling of pooled CRISPR screens. *Nat Biotechnol* **42**, 1218-1223
(2024).

38.   Norman, T.M. et al. Exploring genetic interaction manifolds constructed from rich single-cell
phenotypes. *Science* **365**, 786-793 (2019).

39.   Sunshine, S. et al. Systematic functional interrogation of SARS-CoV-2 host factors using Perturb-seq.
Nat Commun **14**, 6245 (2023).

40.   Jiang, Y., Li, Y. & Zhu, B. T-cell exhaustion in the tumor microenvironment. *Cell Death Dis* **6**, e1792
(2015).

41.   Lee, J., Ahn, E., Kissick, H.T. & Ahmed, R. Reinvigorating Exhausted T Cells by Blockade of the PD-
1 Pathway. *For Immunopathol Dis Therap* **6**, 7-17 (2015).

42.   Guo, M.H. et al. Comprehensive population-based genome sequencing provides insight into
hematopoietic regulatory mechanisms. *Proc Natl Acad Sci U S A* **114**, E327-E336 (2017).

43.   Takahashi, K. & Yamanaka, S. Induction of pluripotent stem cells from mouse embryonic and adult
fibroblast cultures by defined factors. *Cell* **126**, 663-676 (2006).

44.   Krah, N.M. et al. The acinar differentiation determinant PTF1A inhibits initiation of pancreatic ductal
adenocarcinoma. *Elife* **4** (2015).

45.   Jakubison, B.L. et al. Induced PTF1a expression in pancreatic ductal adenocarcinoma cells activates
acinar gene networks, reduces tumorigenic properties, and sensitizes cells to gemcitabine treatment.
Mol Oncol **12**, 1104-1124 (2018).

46.   Rood, J.E., Hupalowska, A. & Regev, A. Toward a foundation model of causal cell and tissue biology
with a Perturbation Cell and Tissue Atlas. *Cell* **187**, 4520-4545 (2024).

47.   He, S. et al. Squidiff: Predicting cellular development and responses to perturbations using a diffusion
model. *bioRxiv* (2024).

48.   Li, C. et al. scDirect: key transcription factor identification for directing cell state transitions based on
single-cell multi-omics data. *bioRxiv* (2024).

49.   Pierce, S.E., Granja, J.M. & Greenleaf, W.J. High-throughput single-cell chromatin accessibility
CRISPR screens enable unbiased identification of regulatory networks in cancer. *Nat Commun* **12**,
2969 (2021).

50.   Liscovitch-Brauer, N. et al. Profiling the genetic determinants of chromatin accessibility with scalable
single-cell CRISPR screens. *Nat Biotechnol* **39**, 1270-1277 (2021).

51.   Papalexi, E. et al. Characterizing the molecular regulation of inhibitory immune checkpoints with
multimodal single-cell screens. *Nat Genet* **53**, 322-331 (2021).

52.   Wolf, F.A., Angerer, P. & Theis, F.J. SCANPY: large-scale single-cell gene expression data analysis.
Genome Biol **19**, 15 (2018).

53.   Bergen, V., Lange, M., Peidli, S., Wolf, F.A. & Theis, F.J. Generalizing RNA velocity to transient cell
states through dynamical modeling. *Nat Biotechnol* **38**, 1408-1414 (2020).