Genomic signals of local adaptation in *Eleginops maclovinus* from Northern Chilean Patagonia

C. Eliza Claure [1,2,3,4], Garrett D. McKinney [5], J. Dellis Rocha [6], José M. Yáñez[7], Iván Pérez-Santos [1], Cristian B. Canales-Aguirre [1,2]

[1] Centro i~mar, Universidad de Los Lagos, Camino a Chinquihue 6 km, Puerto Montt, Chile
[2] Núcleo Milenio INVASAL, Concepción, Chile
[3] Programa de Magister en Ciencias, mención Producción, Manejo y Conservación de Recursos Naturales, Universidad de Los Lagos
[4] Programa de Doctorado en Ciencias, mención Conservación y Manejo de Recursos Naturales, Universidad de Los Lagos
[5] Washington Department of Fish and Wildlife, Seattle, Washington, USA
[6] Escuela de Obstetricia, Facultad de Ciencias para el Cuidado de la Salud, Universidad San Sebastián Sede de La Patagonia, Puerto Montt, Región de Los Lagos
[7] Facultad de Ciencias Veterinarias y Pecuarias, Universidad de Chile, Av Santa Rosa 11735, La Pintana, Santiago 8820808, Chile

*Corresponding author:
Cristian B. Canales-Aguirre
Centro i~mar, Universidad de Los Lagos
Camino a Chinquihue 6 km, Puerto Montt, Chile
cristian.canales@ulagos.cl

Abstract

Understanding the evolutionary mechanisms that shape the adaptive divergence across

spatially heterogeneous environments is a challenging task for evolutionary ecologists. The

Chilean marine Patagonia is a complex ecosystem with diverse geomorphology and

physical-chemical oceanographic conditions. There is limited research evaluating the

interactions between selective forces and environmental conditions in this area. This study

focuses on identifying the genomic signals of local adaptation of the endemic marine fish,

Eleginops maclovinus from Chilean North Patagonia. To achieve this goal, we used an

environmental marine database (temperature, salinity, oxygen, phosphate nitrate and

silicate concentration) with collected from 1995 to 2018 and 11,961 SNPs obtained from

246 individuals from 10 sampling locations across this area. We identified putative

adaptive loci using ten bioinformatic software tools, where five were based on population

genetic differentiation (PGD) and five based on the genotype-environment association

(GEA). We identified 392 adaptative loci using PGD and 2,164 associated with at least one

of the six environmental variables analyzed using GEA. A total of 131 loci were shared

between the PGD and GEA approaches, of which 37 were associated with genes involved

in the growth, metabolism and homeostasis. Then, we evaluated the variation of adaptive

loci with environmental variables using polygenic scores and found significant correlations

with temperature, salinity, and oxygen, indicating polygenic selection along environmental

gradients. This study highlights how polygenic selection drives local adaptation in

Eleginops maclovinus and underscores the value of integrating genomic and environmental

data for conservation in the Patagonian ecosystem.

Keywords*:* Patagonian blennie, SNPs, candidate genes, seascape genomics, RAD-seq.

 1   Introduction

The main aim of adaptive genomics is understanding the molecular basis of local

adaptation in species experiencing spatially heterogeneous environments. Currently,

detecting adaptative loci is possible using high-throughput sequencing technologies, which

allows thousands of single-nucleotide polymorphisms (SNPs) distributed across the

genome (Willette *et al.* 2014), to be genotyped in hundreds of individuals (Manel *et al.*

2016).

Restriction site-associated DNA sequencing (RAD-seq) method harness the high-

throughput sequencing technology to genotype individuals (Miller *et al.* 2007; Andrews *et*

al. 2016; McKinney *et al.* 2017a). Specifically, this method uses restriction enzymes that

cut at specific motifs throughout the genome to reduce their complexity (Altshuler *et al.*

2000; Miller *et al.* 2007; Baird *et al.* 2008). The use of hundreds of thousands of RAD-seq

markers for exhibiting the signatures of selection is increasingly applied to marine

organisms, offering particular advantages for complex genomes in the absence of a

reference sequence (Wang *et al.* 2015; Benestan *et al.* 2016; Cure *et al.* 2017; Dalongeville

et al. 2018; Hoey & Pinsky 2018; Li *et al.* 2019; Segovia *et al.* 2020; O'Leary *et al.* 2021).

For identifying adaptative loci, there are two predominant genome scan approaches: *(i)*

population genetic differentiation (PGD) and *(ii)* genotype-environment association (GEA).

The approach based on population genetic differentiation (PGD) identifies loci putatively

under selection by primarily comparing the genetic differentiation index (F_{ST}) and other

measures (e.g., inbreeding coefficient (F_{IS}) and locus-specific homozygosity excess) for

each locus against values predicted by a null model of neutral evolution. If natural selection

favors one allele at a particular locus in some populations, F_{ST} at that locus will be large

compared to loci in which among-population differences are the result of neutral

demographic processes (François *et al.* 2016). Contrarily, approach based on genotype-

environment association (GEA) search for associations between environmental variables

and allele frequencies across a set of populations spread over a geographic space (Rellstab

et al. 2015; Hoban *et al.* 2016). Alleles involved in local adaptation should occur at higher

frequency where they increase fitness and be at lower frequency where they decrease

fitness. Therefore, it may be possible to identify putative adaptative loci by scanning for

alleles that show greater than average genetic differentiation among populations using GEA

approach (Lewontin & Krakauer 1973; Beaumont 2005). These approaches are based on

the premise that it is possible to distinguish between the genetic signals left by adaptative

and neutral processes. However, algorithms, statistical models, and prior assumptions

considerably differ and all of them are prone to false positive associations (de Villemereuil

et al. 2014; Rellstab *et al.* 2015; de Villemereuil & Gaggiotti 2015; François *et al.* 2016).

Combining PGD and GEA outcomes may provide an efficient strategy to detect signatures

of local adaptation while reducing the number of false positives (de Villemereuil *et al.*

2014; Rellstab *et al.* 2015; Gagnaire *et al.* 2015; François *et al.* 2016).

North Patagonian ecosystems in southern Chile are constituted by extensive and

interconnected system of fjords, channels, gulf, estuaries, and bays that are affected by

physical regimes that may strongly modulate biological productivity (Pantoja *et al.* 2011;

Iriarte *et al.* 2014). It extends approximately across 140,000 km^2 from latitude ~41.5ºS

(Reloncaví Fjord) to latitude ~46.5ºS (San Rafael Lagoon) (Rodrigo 2008).

Oceanographically, Chilean Northern Patagonia is considered a transitional marine system,

where the interaction between marine and fresh waters produces strong vertical and

horizontal gradients in physical-chemical variables (Pickard 1971; Dávila *et al.* 2002;

Iriarte *et al.* 2014; Pérez-Santos *et al.* 2014; Schneider *et al.* 2014). These characteristics

make Patagonia an excellent natural laboratory to investigate how this seascape

heterogeneity results in genomics features. Thus, integrating genomic and environmental

data provides a powerful approach for uncovering the processes leading to adaptive

divergence in ecologically heterogeneous and connected marine ecosystems. So far, studies

focused on seascape genomics in Northern Patagonia are limited (Araneda *et al.* 2016;

Canales-Aguirre *et al.* 2016, 2018; Delgado *et al.* 2019; Segovia *et al.* 2020; Canales-

Aguirre *et al.* 2022).

Eleginops maclovinus (Cuvier & Valenciennes 1830), is a notothenioid fish

(Perciformes) of the monotypic family Eleginopidae (Pequeño 1989). This marine fish is

endemic to the coastal temperate and sub-Antarctic waters of South America with a

distribution range along the Atlantic and Pacific Patagonian coasts. In the Atlantic Ocean, it

is distributed from Uruguay (~35°S) to the Beagle Channel (~54°S), including the

Falkland/Malvinas Islands (Gosztonyi & Ae 1974; Eastman 1993), while in the Pacific

Ocean, from the Beagle Channel to Valparaíso (~33°S) (Pequeño 1989). This species is a

protandrous hermaphrodite with the highest fecundity (~796 eggs g^{-1}) (Brickle *et al.* 2005)

among notothenioids (80-300 eggs g^{-1}) (Kock 1992; North 2001). It is euryhaline and can

tolerate a broad range of salinity concentrations (5 and 45 PSU), however, extreme

salinities can produce dramatic metabolic responses to fuel osmoregulatory tissues (Vargas-

Chacoff *et al.* 2014, 2015a). Furthermore, juveniles of *E. maclovinus* have been described

having a large eurythermal capacity, with tolerance between 4 and 20°C (Vanella *et al.*

2012; Oyarzún *et al.* 2018; Lattuca *et al.* 2018). All these studies described above have

been conducted under controlled conditions, they are focused on one or two environmental

variables, and do not associate with genetic data. The latter leaves a gap on how loci under

selection can shape the adaptative capacity of *E. maclovinus* populations considering the

complex environmental variability in estuaries and fjords.

Studies about the geographic pattern of genetic diversity in *E. maclovinus* are

limited to three (Ceballos *et al.* 2012, 2016; Canales-Aguirre *et al.* 2022). A

phylogeographic research conducted by Ceballos et al. (2012) using mitochondrial DNA

sequences for nine locations did not find any evidence for the structuring of genetic

variation into an Atlantic and a Pacific group. Subsequently, using nine microsatellites it

was revealed a low but significant level of genetic differentiation between Pacific and

Atlantic populations and showed differences between locations in the Pacific coast

(Ceballos *et al.* 2016). However, the authors acknowledge that a single mitochondrial locus

or nine microsatellites may not be robust enough to allow them to recover significant

subdivision among geographic regions. Recently, Canales-Aguirre *et al.* (2022) used

12,050 SNPs in five locations distributed across Northern Patagonia to identify neutral and

adaptive population structure. However, the latter only included five locations, used a

public environmental database Bio-ORACLE (Tyberghein *et al.* 2012; Assis *et al.* 2018)

which is not accurate for fine-scale and fjords areas like Patagonia, and only three

approaches to identify adaptive loci. In addition, its focus was to identify natural and

adaptive loci for conservation and management purposes. To our knowledge, this is the first

study to identify the signal of local adaptation using ten different bioinformatic approaches

and six highly accurate environmental variables in populations of *E. maclovinus* along
Northern Patagonia.

In this study, we applied RAD-seq sequencing to generate SNPs genotyped in 246
individuals of *E. maclovinus* collected in North Patagonia. Here, we expanded the
sampling locations from Canales-Aguirre et al. (2022), increasing the number to ten
locations to better capture the environmental heterogeneity of their North Patagonian
seascape. Our aim was to detect SNPs putatively under selection using population genetic
differentiation (PGD) and genotype-environment association (GEA) approaches. We tested
associations between SNPs and six environmental variables (temperature, salinity
concentration of oxygen, nitrate, phosphate and silicate) obtained from oceanographic data
recorded between 1995 to 2018 in this area. We implemented a BLASTx search on putative
adaptative SNPs identified by both PGD and GEA approaches to investigate genes with
molecular functions potentially involved in local adaptation. Then, we evaluated the
variation of adaptive loci using polygenic scores where we found significant correlations
and supporting polygenic selection in North Patagonia.

2   Materials and Methods

2.1   Study area and environmental predictors

Chilean Patagonia is one of the most extended fjord regions in the world, located on the southeastern border of South America (Fig.1A), and composed by islands, peninsulas, fjords, and channels (Pantoja *et al.* 2011; Iriarte *et al.* 2014). This region has been affected by ice sheets calving into the ocean, retraction of the sea coast line and decrease in marine water temperatures since the Great Patagonian Glaciation (c. 1 Ma) during the Pleistocene (Rabassa *et al.* 2011). The Northern Patagonia is characterized by interactions between marine and fresh water, generating sharp vertical and horizontal salinity gradients (Dávila *et al.* 2002; Pérez-Santos *et al.* 2014; Schneider *et al.* 2014). The temperature shows a latitudinal gradient, which decreases southward until the Antarctic Polar Front (~55°S) (Iriarte *et al.* 2014). The oxygen concentration in the fjords is the result of horizontal advection of adjacent well-oxygenated Subantarctic Waters (5–6 mL/L) representing the major source of oxygen in the deep layers of the inner seas of Patagonia (Silva & Vargas 2014). In recent years, dissolved oxygen (DO) in the fjords of Chilean Patagonia has decreased by 1–2%, potentially driven by nutrient accumulation and rising temperatures (Linford *et al.* 2023a; b, 2024). The concentrations of inorganic nutrients show a strong seasonal signal, with high nitrate and orthophosphate concentrations during winter, and lower values during spring, presumably caused by a severe increase in primary productivity when light availability in near-surface waters increases (Iriarte *et al.* 2007; González *et al.* 2011; Torres *et al.* 2014; Jacob *et al.* 2014). The concentration of silicic acid shows latitudinal and longitudinal gradient due to surface water mixing between nitrate-rich but

Silicic acid-depleted oceanic subantarctic waters and Silicic acid-rich but nitrate depleted

continental waters (Aracena *et al.* 2011; González *et al.* 2011; Torres *et al.* 2014).

Figure 1   Geographic area and environmental variables in used.

Map of sampling locations in South America. B) Sampling site locations on North Patagonian fjord ecosystems in southern Chile REL:

Reloncaví Estuary, MAN: Manao, HOR: Hornopirén, CHE: Chepu, CHA: Chaitén, RMB: Raúl Marín Balmaceda, MEL: Melinka

Island, PUY: Puyuhuapi, CIS: Port Cisnes, AYS: Port Aysén: C) Mean surface temperature (ºC) D) Mean surface salinity (g/Kg); E)

Mean surface concentration oxygen (µM); F) Mean surface nitrate (µM); G) Mean surface phosphate(µM); H) Mean surface silicate

(µM).

We compiled an environmental database using data collected through the CIMAR Program

of the National Oceanographic Committee of Chile (CONA), from CIMAR 1 (1995) to

CIMAR 24 (2018). The database comprises 1,195 sampling stations located along the inner

sea of Chiloé Island, estuaries, channels, and fjords from the Chilean Patagonia sampled in

spring and winter (http://www.shoa.cl/n_cendhoc/productos/reporte_datos.php) (Fig. S1).

At each sampling station, environmental variables were measured from the surface to the

maximum possible depth. Considering that traditional databases lack high-resolution data

near the coasts (e.g., Bio-Oracle, World Ocean Atlas, Aqua/MODIS), we highlight the use

of information obtained by CIMAR as an effective alternative to detect the environmental

heterogeneity of Northern Patagonia. Our database compiled contains six environmental

predictors: conservative temperature[1] (ºC), absolute salinity[2] (g Kg^{-1}), oxygen (mL L^{-1}),

nitrate (µM), phosphate (µM) and silicate (µM) concentrations. We used Ocean Data View

v.5.4.0. (Schlitzer 2002) for data visualization. We estimated the mean, minimum,

maximum and range of these environmental variables in each of the ten locations for a

water column until 100 m depth using the entire period of the database (1995 – 2018)

(Table S1). These latter, considering that *E. maclovinus* inhabits in coastal and estuarine

areas at depths >40 m, and the spawning occurs in spring season (Brickle *et al.* 2005)

---

[1] Conservative temperature is a thermodynamic property of seawater. It is derived from the potential enthalpy and is recommended under the TEOS-10 standard (Thermodynamic Equation of Seawater-2010) as a replacement for potential temperature as it more accurately represents the heat content in the ocean.

[2] The Absolute Salinity is the mass fraction of dissolved non☐H_2O material in a seawater sample at its temperature and pressure and is expressed in g Kg^{-1}

2.2 Sampling and genotyping

Between November 2018 and April 2019, we sampled 246 individuals of *E. maclovinus* from ten locations throughout the North Patagonian fjord ecosystems in southern Chile (Fig.1B). Five of the ten sampling locations included in this study were previously analyzed by Canales-Aguirre *et al.* (2022). Sampling, DNA extraction, RAD-sequencing library preparation, sequencing with Illumina technology and pipeline of bioinformatic analyses were performed following procedures described by Canales-Aguirre *et al.* (2022).

Loci and samples were filtered iteratively by initially removing poor-quality data, recalculating missing data proportions, and applying increasingly stringent thresholds, following McKinney et al. (2020). The first part of the filtration process consisted in excluding SNPs or individuals based on the following criteria *(i)* minor allele frequencies (MAF) per sampling location, *(ii)* genotype rate per locus, *(iii)* genotype rate per sample. Initially, filtration criterion was MAF ≤ 0.05, genotype rate threshold of 50% per locus and sample. Subsequently, MAF≤ 0.1, genotype rate threshold of 90% per locus and sample. Due to gene duplication events described to have affected the evolution of the notothenioid genome (Xu *et al.* 2008; Chen *et al.* 2008, 2019), *HDplot* (McKinney *et al.* 2017b) was used to identify paralogs, which were removed from further analysis (parameter flags: $H < 0.6$; $|D| < 5$). We removed SNPs in Hardy-Weinberg disequilibrium (p-value < 0.05) if they deviated in three or more locations to avoid potential issues such as inbreeding, population stratification, and genotyping errors in the identified SNPs (Wigginton *et al.* 2005). For reads with more than one SNP, we only kept those with the highest F_{ST}, to reduce the influence from linked loci in the results. The resulting filtered

227 (.vcf) file was converted into the corresponding file formats for the subsequent analyses by

228 using *PGDSpider* v. 2.1.1.5. (Lischer & Excoffier 2012).

2.3 Detecting putative loci under selection

We employed ten different software programs to identify putative adaptive loci: five

based on population genetic differentiation (PGD) and five on genotype-environment

association (GEA). We combined multiple approaches to minimize false positive rates and

maximize our chances of detecting genomic signals of local adaptation (Rellstab *et al.*

2015; François *et al.* 2016). For the PGD approach, we used *fsthet* v.1.0.1 (Flanagan &

Jones 2017), *BayeScan* v.2.1 (Foll & Gaggiotti 2008), *OutFLANK* v.0.2 (Whitlock &

Lotterhos 2015), *PCAdapt* v.4.3.3 (Luu *et al.* 2017; Privé *et al.* 2020), and *Arlequin*

v.3.5.2.2 (Excoffier & Lischer 2010) (Table 1). For the GEA approach, we used six

environmental variables and five software programs: *Latent Factors Mixed Model (LFMM)*

(Frichot *et al.* 2013), *BayeScEnv* (de Villemereuil & Gaggiotti 2015), *Redundancy Analysis*

(RDA).(Hair *et al.* 1995; Zuur *et al.* 2010; Capblancq *et al.* 2018; Capblancq & Forester

2021)*, Moran Spectral Outlier Detection (MSOD)* (Wagner *et al.* 2017), and *Samβada*

(Stucki *et al.* 2017; Duruz *et al.* 2019) (Table 2). In all programs, a 5% False Discovery

Rate and respective p-values or q-values were applied to detect putative adaptive loci. We

employed *UpSet* diagrams (Lex *et al.* 2014) to visually compare the results from the ten

software programs, identifying loci flagged as outliers by at least two programs as putative

adaptive loci. For subsequent analyses, loci were classified into two categories: putative

adaptive loci (under directional selection) and putative neutral loci, excluding loci under

balancing selection.

Table 1 Description and parameters used for software with the PGD approach.

Software	Description	Parameters	References
i) *fsthet*	*fsthet* identifies loci with extreme F_{ST} values by calculating smoothed quantiles for the F_{ST}-heterozygosity distribution, using its own empirical distribution instead of a null model due to changes in population structure.	We used a confidence level of 95% to detect F_{ST} outliers	(Flanagan & Jones 2017)
ii) *BayeScan*	*BayeScan* estimates the probability of selection for each locus using a Bayesian test that distinguishes between locus-specific selection effects (α) and population-specific demography effects (β), by measuring the discordance between global and population-specific allele frequencies.	We ran it with default parameters: 20 pilot runs of 5,000 iterations followed by 50,000 iterations with an additional burn-in of 50,000 and prior odd of 10.	(Foll & Gaggiotti 2008)
iii) *OutFLANK*	OutFLANK calculates a neutral distribution of F_{ST} values and assigns q-values to each locus to detect adaptive loci influenced by selection, without assuming a specific population's demographic history, reducing false positives.	We ran it with default parameters: LeftTrimFraction = 0.05, RightTrimFraction = 0.05, Hmin = 0.1.	(Whitlock & Lotterhos 2015)
iv) *PCAdapt*	*PCAdapt* performs a Principal Component Analysis (PCA) and computes the p-values for each locus to detect putative adaptative loci.	The most appropriate number of clusters K was chosen by testing K=1:10. We detected outliers using K=2 and p-values < 0.05 associated with Mahalanobis distances calculated with the q-value package.	(Dabney *et al.* 2010; Luu *et al.* 2017)
v) *Arlequin*	*Arlequin* calculates observed heterozygosity (H_O) to create a null distribution of F_{ST} values and associated p-values for each locus.	This was executed under a hierarchical island model with 20,000 simulations and 100 demes, which implements FDIST2 methodology.	(Excoffier & Lischer 2010)

251 **Table 2** Description and parameters used for software with the GEA approach.

Software	Description	Parameters	References
i) Latent factors mixed model (*LFMM*)	*LFMM* is a hierarchical Bayesian mixed model, which uses latent factors to correct population structure	We estimated the number of ancestral populations K before the analysis testing from K 1 to 10 using the LEA v.3.2.0 R package and determined K from the cross-entropy criteria and Cattell's	(Cattell 1966; Devlin & Roeder 1999; Frichot *et al.*

		while fitting a linear regression between allele frequencies and environmental variables.	rule from the sNMF output. LFMM analyses were conducted separately for each environmental variable, with 10,000 iterations, and burning of 5,000 and five replicates.	2013, 2014; Frichot & François 2015)
ii)	*BayeScEnv*	*BayeScEnv* detects putative adaptive loci by identifying loci that show large positive F_{ST} values (outside the neutral model F_{ST} distribution) that are significantly correlated with environmental variables.	We used 20 pilot runs, with 50,000 iterations each and an additional burn-in of 50,000 iterations for each of the six standardized environmental variables.	(de Villemereuil & Gaggiotti 2015).
iii)	*Redundancy Analysis (RDA)*	*Redundancy Analysis (RDA)* is a multivariate ordination method to detect loci putatively under selection based on correlations with environmental variables.	We calculated the variance inflation factor (vif.cca function) to exclude variables with a value ≥ 5, and conducted linear regressions between allele frequencies and environmental variables at each locus. Using PCA, we produced ordination axes, identified putative adaptive loci (± 3 SD from the mean), and evaluated the correlation using the Pearson correlation coefficient.	(Hair *et al.* 1995; Zuur *et al.* 2010; Rellstab *et al.* 2015; Forester *et al.* 2018)
iv)	*Moran spectral outlier detection (MSOD)*	*MSOD* uses Moran eigenvector maps (MEM) to quantify the distribution of allele frequencies across a range of spatial scales represented by MEM spatial eigenvectors	We compared the power spectrum of each locus to the average power spectrum to identify outliers using z-scores and a cut-off of 0.05. Subsequently, we applied Moran spectral randomization (MSR) with 999 permutations using adespatial.	(Dray *et al.* 2006, 2021; Wagner & Dray 2015; Wagner *et al.* 2017)
v)	*Samβada*	*Samβada* is a spatial approach that uses multiples univariate logistic regression models to identify locus-environment associations and at the same time measures spatial autocorrelation	We coded individuals for the presence or absence of each SNP allele and measured the association with environmental parameters across sites using multiple logistic regression. P-values from these regressions were calculated via a Wald Score Test and compared to a χ^2 distribution with one degree of freedom.	(Wald 1943; Stucki *et al.* 2017; Duruz *et al.* 2019)

2.4 Genetic diversity and population structure

For the adaptative and neutral datasets, we determined, described and compared overall and population-specific genetic diversity. The expected heterozygosity (H_E), observed heterozygosity (H_O) and inbreeding coefficient (F_{IS}) were calculated for each locus and locations using *Hierfstat* v 0.04.10 R package (Goudet 2005). We assessed the genetic differentiation with pairwise F_{ST} (Weir & Cockerham 1984) comparisons between locations using the R package *StAMPP* v.1.6.2. (Pembleton *et al.* 2013) and the statistical significances were determined using 10,000 permutations.

We used two approaches to investigate the genetic structure: Bayesian clustering program *Structure* v.2.3.4. (Pritchard *et al.* 2000) and discriminant analysis of principal components (DAPC) (Jombart *et al.* 2010) using r package *adegenet* v.2.1.5. (Jombart 2008; Jombart & Ahmed 2011). *Structure* is a Bayesian method, which uses a Markov chain Monte Carlo (MCMC) algorithm based on the Gibbs sampler algorithm to identify the number of putative genetic clusters. This method assumes Hardy-Weinberg equilibrium (HWE) and linkage equilibrium among loci in sample population individuals (Pritchard *et al.* 2000). We conducted the analysis using ten replicates per K value for K=1 to K=10, with 10,000 burn-in steps and 100,000 MCMC-sampling steps. The runs were collected, sorted and merged and functions in the R package *pophelper* (Francis 2017) to produce assignment barplots. The most likely number of genetic clusters (K) was chosen using the ΔK method (Evanno *et al.* 2005). The DAPC method identifies and describes clusters of genetically related individuals from genomic datasets and allows the optimal visualization

carried out a principal component analysis (PCA), and used the principal components (PCs)

thus produced as synthetic variables for a discriminant analysis, as outlined in Jombart et al.

(2010). The optimal number of principle components was determined by cross validation

(*xvalDapc* function (Jombart *et al.* 2010)) to avoid overfitting.

We also conducted isolation by distance (IBD) analyses to test whether genetic distance

increased with geographic distance with adaptative and neutral data. We assessed the

correlation between the linearized pairwise F_{ST} matrix and the Euclidean (geographic)

distance matrix using a Mantel-test as implemented in the R-package *vegan* (Oksanen *et al.*

2016). Statistical significance was assessed with 999 permutations of the data. To test

whether the correlation between genetic and geographic distances is a result of a continuous

or distant patchy cline of genetic differentiation, local densities of distances to disentangle

both processes were plotted using 2-dimensional kernel density estimation (*kde2d* function)

of *MASS* R-package (Venables & Ripley 2013).

2.5   Additive polygenic scores (APS)

We assessed cumulative adaptive genetic variation by calculating additive polygenic

scores (APS) associated with each environmental variable (Gagnaire & Gaggiotti 2016;

Babin *et al.* 2017). For this analysis, we prioritized three biologically important

environmental variables: annual maximum temperature (°C), annual maximum salinity

(g/Kg), and annual minimum oxygen concentration (µM) (Holliday 1969; Nikinmaa &

Rees 2005; Brian *et al.* 2008; Vanella *et al.* 2012; Vargas-Chacoff *et al.* 2014). We

prioritized the maximum values of temperature and salinity considering the effect that

climate change has on them. According to the Community Climate System Model 3.0

(CCSM3) of the National Center for Atmospheric Research (NCAR), a 2.5°C increase in

the temperature of Chilean coastal areas is predicted by 2055 (Silva *et al.* 2016). Similarly,

the Center for Oceanographic Research in the Eastern South Pacific (COPAS) detected an

increase of 0.25 in salinity in the upper layers of the central-southern coast of Chile

(Schneider *et al.* 2017)We also considered the minimum oxygen concentration, as hypoxic

conditions negatively impact fish fitness (Richards 2009; Wu 2009). All these variables

have an impact of the adaptive genetic diversity of several species (Pauls *et al.* 2013;

Bernatchez 2016)..

We assessed the relationship between each of the 131 putative adaptative loci and

each environmental variable. We used genotypes (0,1,2) to determine the score for each

favored allele; if the slope is positive, and (2,1,0) if negative (Babin *et al.* 2017). We

calculated APS by summing the scores of favored alleles associated with each

environmental variable. Finally, we evaluated the relationship between individual APS and

environmental variables using three models: linear, quadratic, and null, determining the

best fit based on the lowest Akaike Information Criterion (AIC) value.

2.6   Loci annotation and Candidate Gene Identification

Loci identified as putative adaptative loci (see results section 3.5) were functionally

annotated through Blast2GO v.6.0.3 (Conesa & Götz 2008). We compared the sequences

(e.g., heat shock, immune system proteins, osmoregulation). We applied the BLASTx

algorithm with a cut-off of E-value $\leq 1 \times 10^{-5}$, homology of sequences of more than 75% and

other parameters were used as default. Subsequent analysis only considered sequences with

functional annotation (blasted, mapped and annotated).

3.1   RAD sequencing and data filtering

After quality control, 2,359,848 (511,190 RADtag) putative loci were obtained for

Stacks pipeline. Using the iterative filtering process (MAF and missing data), we identified

a total of 2,328,560 (98.67%) putative singletons and 386 putative paralog loci; the latter

were removed (1.24%). Further filtering by HWE discarded 13,215 loci (42.96%) and one

SNP per tag (the highest F_{ST}) discarded 5,582 SNPs (31.82%). A total of 11,961 SNPs

(5.01%) were retained, from 202 individuals across the 10 sampling locations (Table 3).

Table 3   Description of steps used to filter SNPs

Filtering steps	Sample size	Number of loci
SNP after STACKS	246	2,359,848
[1]MAF >= 0.05	246	1,494,046
[1]Genotyped by locus (50%)	246	167,752
[1]Genotyped by sample (50%)	236	167,752
[1]Genotyped by locus (90%)	236	54,694
[1]Genotyped by sample (90%)	202	54,694
[1]MAF >= 0.1	202	31,144
Singletons (HDPlot H=0.6; D=5)	202	30,758
Hardy Weinberg equilibrium in 100% locations	202	17,543
one SNP by tag	202	11,961

[1] Steps included in the iterative filtering process

From population genetic differentiation (PGD) and genotype-environment association

(GEA) we identified different putative adaptive loci. Out of the 11,961 loci, we identified

392 loci under positive selection by PGD, and 2,164 loci using the GEA approach. A total

of 131 loci were identified as shared between the PGD and GEA approaches (Figure 2).

Figure 2   UpSet diagram: matrix layout for all intersections of software used. A) Loci

detected by software based on Population Genetic Differentiation (PGD), B) Loci detected

by software based on Environment Genotype-Association and C) Loci shared between

PGD and GEA. The set size (horizontal bars) indicates the total number of loci detected by

that software. The intersection size (vertical bars) indicates the number of loci per set

and 313 SNPs (2.62%) under divergent selection (Figure S2). *Arlequin* identified 270 SNPs

under divergent selection (2.26%) (Figure S3). *PCAdapt* detected 15 SNPs (0.13%) under

divergent selection (Figure S4). *BayeScan* and *OutFLANK* found no outlier. For GEA

approaches, *LFMM* identified 1,785 loci under divergent selection, with the highest number

of associations to nitrate concentration (n = 684 loci) (Figure S5). *MSOD* detected 377 loci,

with the highest number of associations to temperature (n = 377 loci) (Figure S6). Finally,

RDA detected 83 loci under divergent selection, with the highest number of associations to

silicate (33 loci) (Figure S7). *Samβada* and *BayeScEnv* did not detect any correlation with

environmental variables. Overall, across GEA approaches, the nitrate concentration (n =

844), temperature (n = 748), and salinity (n = 703) were the environmental variables with

which most of the loci were correlated (Figure 3). Of the total of 2,164 that correlated with

each of the six environmental variables, 783 of the loci were significantly associated with

more than one environmental variable.

Figure 3   UpSet diagram: matrix layout for all intersections of environmental

variables.

The set size (horizontal bars) indicates the total number of loci correlated with that

environmental variable. The intersection size (vertical bars) indicates the number of loci per

set intersection, the purple dots represent the set of unique loci correlated only with that

environmental variable and no other one, while the purple dots connected by a line

represent loci shared by two, three, or more variables.

Summary statistics of genetic diversity revealed similar values for each location within

each dataset (Figure S8). Most locations showed lower values of observed heterozygosity

(H_O) than expected heterozygosity (H_E), and positive values of inbreeding coefficient F_{IS}.

For the adaptative loci dataset, H_O 0.3364 ± 0.0078 (C.I.$_{95\%}$), H_E 0.3551 ± 0,0068(C.I.$_{95\%}$),

and F_{IS} 0.0528 ± 0.0120 (C.I.$_{95\%}$). For the neutral dataset, H_O 0.3414 ± 0.0074 (C.I.$_{95\%}$), H_E

0.3485 ± 0,0063 (C.I.$_{95\%}$), and F_{IS} 0.0206 ± 0.0117 (C.I.$_{95\%}$).

Pairwise-F_{ST} comparisons showed higher levels of genetic differentiation for adaptive

loci (global F_{ST} = 0.0419 ± 0.0042 CI$_{95\%}$), than neutral loci (global F_{ST} = 0.0004 ± 0.0004

CI$_{95\%}$). For adaptative loci, all pairwise differentiation values were highly significant (p-

value ≤ 0.0004). While in neutral loci, 26 of 45 pairwise differentiation values showed

significant p-values (Table 4).

Table 4 Pairwise F_{ST} (below diagonal) and p-value (above diagonal).
383 Values between sampling site locations from North Patagonian fjord ecosystems in
384 southern Chile using A) adaptative loci and B) neutral loci. Statistical significance was
385 considered below 0.05 p-value.
386

A)	REL	MAN	HOR	CHE	CHA	RMB	MEL	PUY	CIS	AYS
REL	-	0	0	0	0	0	0	0	0	0
MAN	0.0672	-	0.0004	0	0	0	0	0	0	0
HOR	0.0495	0.0126	-	0	0	0	0	0	0.0004	0
CHE	0.0653	0.0443	0.0493	-	0	0	0	0	0	0
CHA	0.0665	0.0502	0.0503	0.0531	-	0	0	0	0	0
RMB	0.044	0.0302	0.0227	0.0502	0.0502	-	0	0	0	0
MEL	0.0755	0.0343	0.0296	0.0521	0.0539	0.0313	-	0	0	0
PUY	0.0467	0.0403	0.0266	0.0546	0.0408	0.0286	0.033	-	0	0
CIS	0.0592	0.0338	0.0210	0.0546	0.0435	0.0248	0.0355	0.0234	-	0
AYS	0.0472	0.0442	0.0199	0.0553	0.0456	0.0269	0.0432	0.0263	0.0292	-

B)	REL	MAN	HOR	CHE	CHA	RMB	MEL	PUY	CIS	AYS
REL	-	0	0	0.9997	0.7841	0.1541	0	0.0095	0.0763	0.0121
MAN	0.0018	-	0.0004	0.9909	0.3254	0	0	0	0.0008	0
HOR	0.0017	0.0011	-	0.998	0.9681	0.0018	0.0023	0	0.0174	0
CHE	-0.0018	-0.0011	-0.0013	-	1	1	0.9814	0.9893	0.9994	0.9887
CHA	-0.0394	0.0002	-0.0789	-0.0036	-	0.8519	0.8760	0.2556	0.9314	0.4809
RMB	0.0004	0.0016	0.0011	-0.0022	-0.0466	-	0.0003	0.0489	0.3379	0
MEL	0.0019	0.0019	0.0011	-0.0011	-0.524	0.0013	-	0	0.0070	0
PUY	0.0010	0.0020	0.0019	-0.0011	0.0297	0.0006	0.0020	-	0.0004	0.0002
CIS	0.0006	0.0012	0.0007	-0.0017	-0.0684	0.0002	0.0010	0.0014	-	0.0006
AYS	0.0009	0.0017	0.0015	-0.0010	0.2420	0.0012	0.0016	0.0012	0.0012	-

loci (Figure S9) and K = 7 for neutral loci (Figure S10). For adaptative loci, a visual

inspection revealed a clear separation of individuals for REL, MAN and MEL using K=2.

We observed an additional cluster which appeared to correspond to individuals from CHE

and CHA using K = 4 (Figure 4A). For neutral loci, *Structure* showed no clear geographical

genetic structure (Figure 4B). Results from DAPC analyses revealed similar patterns to

Structure (Figure 5). The adaptative dataset showed that the REL, CHA and CHE were

separate from all other locations. The first two axis explain 44.9% of the total variation. For

the neutral dataset, PET, CHA and CHE overlapped greatly with the other locations and

much less structure overall was present, although MAN seemed to be less like the other

nine locations. The first two axis explain 40.5% of the total variation (Figure 5).

Figure 4   Results from clustering analysis of the *E. maclovinus*.

Based on A) 131 putatively adaptative loci and B) 9,536 putatively neutral loci. Each

vertical line represents an individual. The colors represent the proportion of inferred

different locations. ΔR calculated for R ranging from 1 to 10, with each R repeated

tenfold.

A) adaptative loci and B) neutral loci. Axes correspond to the first two discriminant

functions and the circles represent individuals colored-coded by sampling location.

significant isolation by distance pattern (Mantel test: r $=$ 0.3458, p-value $=$ 0.0320,

Figure 6A), where the scatterplot of local densities of distances showed only one consistent

cloud of points, indicating a continuous cline of genetic differentiation. For neutral loci, we

found no detectable relationship between geographic distance and genetic distance (Mantel

test: r = -0.0435, p-value = 0.5760, Figure 6B).

Figure 6 Isolation by distance IBD among locations. A) 151 adaptive loci and B)

9,536 neutral loci. Local density of points plotted using two-dimensional kernel density

estimation. The black line is the trend estimated by Loess regression red line indicates the

linear regression. Color gradient represents density of observations (low = blue, red =

high).

The correlations between additive polygenic scores (APS) and environmental

variables were all significant ($p < 0.001$). The R^2 values ranged from 0.29 for maximum

temperature to 0.26 for both maximum salinity and minimum oxygen concentration. For all

environmental variables, APS decreased with increasing temperature, salinity, and oxygen

levels, respectively (Figure 7). The correlation between APS and environmental variables

was best represented by linear models, except for maximum temperature, which fitted

better with a quadratic model (Figure 7). The significance and R^2 values of the correlations

calculated for mean and minimum temperature, salinity, and oxygen concentration yielded

similar results, albeit with slight differences among subsets (Figure S11-S13).

Figure 7 Correlations between additive polygenic scores (APS) based on A) Annual

maximum temperature B) Annual maximum salinity and C) Annual minimum oxygen and

131 putative adaptative loci. Correlation coefficient (R^2) and p-values and AIC are

presented for each variable.

The BLASTx analysis of the contigs containing the 131 adaptative loci shared by

PGD and GEA against bony fish genomes resulted in significant hits (E-value $\leq 1 \times 10^{-5}$) for

790 fish species. The BLASTx similarity results showed that 51 of the 131 contigs

containing adaptative loci corresponded to known proteins in the non-redundant (NR)

database (E-value $1e^{-5}$), of which 36 contigs were functionally annotated. The functional

categorization of the annotated sequences involved biological process, cellular components,

and molecular function (Figure 8).

 Figure 6 Putative functional categorization and distribution according to Blast2GO.

were functionally annotated. Of these 512 sequences, three were already known to

potentially play a role in the local adaptation (Table S1). The sequences of the *SNP*

6291_214 and *44791_149* and belong to the solute carrier (SLC) family. SLC genes

encode transmembrane transporters for inorganic ions, amino acids, neurotransmitters,

sugars, purines and fatty acids, and other solute substrates (Dorwart *et al.* 2008). The SNP

42_59 is located in the vtgC (13489) gene, which is a member of the vitellogenins (Vtg)

family. In egg-laying vertebrates such as fish, estrogens activate the hepatic synthesis of

vitellogenin (Vtg) (Sumpter & Jobling 1995; Brian *et al.* 2008).

This study provides comprehensive evidence for genomic signatures of local

adaptation, of *Eleginops maclovinus* to heterogeneous environmental conditions in

Northern Patagonia. We conducted a seascape genomics study using 11,961 RAD-

sequencing markers and detected 2,164 putative adaptative loci through population genetic

differentiation (PGD) and genotype-environment association (GEA) approaches. Using 131

loci shared by PGD and GEA, we observed that the putative adaptive loci collectively

exhibit a highly significant association with various environmental variables, indicating

polygenic selection. Additionally, we found contrasting patterns of genetic variation in

neutral and adaptative loci that reinforce the possible local adaptation among high levels of

gene flow. Subsequently, we detected one gene related to thermal adaptation and two genes

that play a role in the osmoregulation. In the following, we describe how our study

contributes to understanding the role of environmental heterogeneity as a driver of spatial

patterns of putative adaptive genetic variation that could influence the local adaptation in *E.*

maclovinus. Furthermore, we discuss methodological issues about software for detecting

putative adaptative loci. Finally, we propose groups based on adaptative divergence for

creating management units.

4.1   Evidence for spatially divergent selection

Evidence suggests thermal adaptation *E. maclovinus*, which juveniles exhibit

significant eurythermal capacity, tolerating temperatures between 4 and 20°C (Vanella *et*

(Vanella *et al.* 2012), while long-term exposure to 15-20°C elevates metabolism with

tissue-specific modulation, without adverse physiological effects. Conversely, temperatures

exceeding 25°C detrimentally impact physiology, growth, food intake, and energy

substrates, mainly triglycerides (Oyarzún *et al.* 2021). These studies focus on juveniles;

thus, temperature effects on early-stage adaptation remain uncertain for this species.

Considering *E. maclovinus* spawns in spring (Brickle *et al.* 2005), higher temperatures

could expedite maturation (e.g. Scott & Pankhurst 1992; Shimizu 2003; Hokanson 2011).

Temperature affects reproductive processes' thermal inhibition (Gillet *et al.* 1996; Tveiten

et al. 2001; Okuzawa *et al.* 2003; Soria *et al.* 2008; Pankhurst & King 2010), including egg

mortality (Van Der Kraak & Pankhurst 1997; Gagliano *et al.* 2007; Janhunen *et al.* 2010),

embryonic development rate (Pauly & Pullin 1988; Rombough 1997; Smith *et al.* 2015),

and larval fish metabolism, growth, and development (Houde 1989; Blaxter 1992; Benoît *et

al.* 2000). Future research should examine temperature-induced shifts in selective mortality

in early stages of *E. maclovinus*. Additionally, temperature may influence sexual inversion

in protandrous hermaphrodites, as proposed by Baroiller et al. (1999). For instance,

elevated temperatures increased gonadal aromatase levels, promoting female proportion in

Lates calcarifer, another protandrous hermaphrodite (Athauda *et al.* 2012). This latter,

although not identified in *E. maclovinus*, can be taken as a research question for future

studies. The detection of potentially adaptive loci associated with temperature supports

molecular adaptation via gene-related heterogeneity and gene duplication events, likely

preserved in response to thermal fluctuations. Unlike most Antarctic notothenioids, *E.

maclovinus* blood shows high hemoglobin (Hb) multiplicity, with one cathodal (Hb C) and

two anodal components (Hb 1, Hb 2) (Coppola *et al.* 2010). This Hb diversity may

facilitate oxygen transport adaptation to environmental changes and metabolic demands

(Weber 1990; di Prisco & Tamburrini 1992; Feuerlein & Weber 1994; Weber *et al.* 2000;

Fago *et al.* 2001). The maintenance of multiple Hbs in response to temperature differences

in temperate waters, significantly larger than in Antarctica, is likely (Coppola *et al.* 2010; di

Prisco *et al.* 2012). Moreover, the absence of antifreeze-glycoprotein (AFGP) coding

sequence in *E. maclovinus* indicates no adaptation to near-freezing temperatures (Cheng *et*

al. 2003).

Evidence suggests salinity acts as a selective agent driving genetic differentiation in

E. maclovinus populations. Canales-Aguirre et al. (2022) identified associations between

adaptive loci and environmental variables (salinity, oxygen, primary production, current

velocity) in *E. maclovinus*. Juveniles are considered euryhalines, adapting to salinities

between 5 and 45 PSU (Vargas-Chacoff *et al.* 2014, 2015a). An inverse relationship

between the trypsin chymotrypsin−1 ratio (T/C ratio) and salinity was observed (Vargas-

Chacoff *et al.* 2015b). The T/C ratio, linked to growth and nutritional status

(Rungruangsak-Torrissen *et al.* 2006; Rungruangsak-Torrissen 2007), suggests better food

utilization in low salinity, enhancing physiological energy. It is possible that food

utilization and efficiency is better in *E. maclovinus* acclimated to low salinity, leading not

to better growth, but to increased energy available for physiological processes such.

Osmoregulation in low salinity consumes energy, affecting growth (Vargas-Chacoff *et al.*

2015b). Euryhaline fish adapt to varying osmotic challenges (Laverty & Skadhauge 2012).

Climate change, habitat degradation, and anthropogenic activities increase salinity stress

increased temperature, affecting hydrological regimes and salinity (Aguayo *et al.* 2019).

The salinity variability in microenvironments can have multiple impacts on reproduction,

including DNA damage from salt stress or oxidative stress (Dowling & Simmons 2009),

sperm motility (Beirão *et al.* 2018), incomplete egg binding (Herberg *et al.* 2018), disrupted

egg activation (Ginsburg 1963) and egg penetration issues (Yanagimachi *et al.* 2017), and

cause problems during DNA recombination (O, Wai-sum *et al.* 2006).

Our results indicate oxygen concentration exerts selective pressure on *E. maclovinus.*

Environmental oxygen variability has been crucial in fish evolution, influencing

anatomical, behavioral, and physiological adaptations for oxygen acquisition and delivery

(Nikinmaa & Rees 2005). Fish have evolved to balance oxygen delivery while avoiding

toxic reactive oxygen species (Taylor & McElwain 2010; Dong *et al.* 2011). They exhibit

diverse hemoglobin structures and functions for oxygen transport (Powers 1974; Jensen &

Weber 1985; Weber 1990; Jensen *et al.* 1998). Hemoglobin diversity in fish, including E.

maclovinus, likely results from selection pressures like low oxygen levels(Nikinmaa &

Rees 2005). Structural and functional analyses of *E. maclovinus* hemoglobin (Hb1Em)

showed higher O_2 affinity compared to *T. bernacchii,* indicating differences in temperature

sensitivity and O_2 release between sub-Antarctic and Antarctic notothenioids (Coppola *et*

al. 2012). Silva & Vargas (2014) identified hypoxic zones in Northern Patagonia's fjords

(1.1–2.0 mL L-1); we emphasize the importance of studying the ability of to adapt to

hypoxic conditions in *E. maclovinus* until now unclear.

In this study, we identified putative adaptive loci associated with the concentration of nitrate, phosphate, and silicate. Although there is no direct relationship between nutrient concentration and adaptive capacity in *E. maclovinus,* the nutrients could be a proxy for primary productivity. The productivity cycle of Patagonian channels and fjords has been divided into two seasons: a productive season characterized by a marked spring diatom bloom followed by pulsed productivity events dominated by a sequence of different phytoplankton species, and then a non-productive season that predominantly features small phytoplankton cells (Iriarte *et al.* 2007; Czypionka *et al.* 2011; Montero *et al.* 2011; Paredes & Montecino 2011). During the productive season and in the absence of persistent hydrodynamic phenomena that force nutrient replenishment in the euphotic zone, the supply of nitrate and phosphate to the well-lit surface layer would be the chief limiting factor for phytoplankton growth (Pizarro *et al.* 2000; Iriarte *et al.* 2007). The increased freshwater discharge into the Puyuhuapi channel during non-productive season coincides with increases in the concentrations of silicic acid and dissolved organic matter concentrations, as well as an increase in underwater light attenuation (Montero *et al.* 2017). A GEA analysis of Canales-Aguirre (2022) provided evidence for local adaptation associated to primary production on *E. maclovinus* likely connected to food availability (Gove *et al.* 2016; Fox *et al.* 2018). Similarly, multiple studies detected putative adaptive loci associated with chlorophyll (indicator of primary productivity), for example Boulanger *et al*. (2022) with *Mullus surmuletus,* Cayuela *et al.* (2020) with *Mallotus villosus,* Maselko et al. (2020) with *Sebastes alutus* and Diopere *et al.* (2017) with *Solea solea.*

For the Additive Polygenic Scores (APS) (Gagnaire & Gaggiotti 2016; Babin *et al.*

2017), we observed a strong correlation between polygenic scores and maximum

temperature and salinity, and minimum oxygen. This suggests that *E. maclovinus*

populations might be locally adapted to these environmental variables through a polygenic

architecture. As we highlighted in the previous section (See 4.1.), we detailed the impact of

these environmental variables on fish survival and development.

We found negative correlations between individual APS and all environmental

variables, which could indicate directional selection, where different alleles are maintained

in distinct environments with a relatively constant selection force along the environmental

gradient. Specifically, the relationship between APS and maximum temperature was best

represented by a quadratic model, suggesting stabilizing selection at intermediate

temperatures. However, caution is advised with this conclusion as the difference in AIC

between the linear and quadratic models was minimal (AIC $_{linear}$ = 1577.8, AIC $_{quadratic}$ =

1577.7).

The impact of temperature on additive genetic variation agreed with observations in

the adaptive population structure membership plots, where the REL population in the

northernmost latitudinal range with warmer temperatures separates from the other nine

populations. For salinity and oxygen concentration, the linear model better explained the

relationship with APS, indicating relatively constant selection along the salinity and oxygen

gradients. These results are consistent with the observed adaptive population structure

where the CHE and CHA populations differentiate from the rest, with the CHE population

exposed to higher oceanic salinities. Lastly, the minimum oxygen concentration is critical

for fish reproduction and development, as hypoxia can delay embryonic development and

hatching (Richards 2009; Wu 2009). However, it remains unknown how hypoxia affects the

genetic expression and reproductive development of *E. maclovinus*.

While the APS approach has potential for evaluating the cumulative effect of

potentially adaptive loci in complex environments, it has limitations. Without a reference

genome, we cannot rule out the possibility that some potentially adaptive loci are linked.

An annotated *E. maclovinus* genome would allow us to identify matches between adaptive

loci and selected genomic regions, providing insights into specific genes involved in local

adaptation (Manel *et al.* 2016). Additionally, comparing genotypic variation in potentially

adaptive loci with individual phenotypic data is necessary to establish genotype, phenotype,

and fitness associations (Barrett & Hoekstra 2011).

Our results suggest that maximum temperature, maximum salinity, and minimum

oxygen concentration may drive spatially variable selection in *E. maclovinus* populations.

Temperature, salinity, and oxygen concentration impact the environmental gradient

experienced by eggs and larvae (Holliday 1969; Blaxter 1992; Brickle *et al.* 2005; Wu

2009; Lehtonen & Kvarnemo 2015), potentially influencing local adaptation. Future

research should aim to detect molecular signatures of selection within a multilocus

quantitative genetics framework to delve deeper into the genetic architecture of polygenic

adaptive traits.

Using BLASTx to identify genes linked to adaptation through a gene ontology

approach (Haasl & Payseur 2016; Manel *et al.* 2016), we found a total of 2,425 potentially

adaptative loci with 131 loci shared by PGD and GEA, only 512 and 36 loci were

successfully annotated, respectively. In the 131 adaptative loci, only 39.69% of loci

matched known sequences in the public database, highlighting the limitation of RAD

sequencing in identifying candidate genes without a reference genome (Davey *et al.* 2013;

Andrews *et al.* 2016; Catchen *et al.* 2017; McKinney *et al.* 2017a). Both adaptive datasets

revealed loci linked to life-history traits, including thermal adaptation and osmoregulation.

Focusing on life history traits, we identified five key genes: *opa1, Stx16, FMVIA,*

foxj1b, and *Lamc3*. The gene *opa1* regulates mitochondrial fusion and fission, preventing

Cyc-c leakage and inducing apoptosis, which is linked to kidney injury (Song *et al.* 2017;

Yingjie *et al.* 2019; Herkenne *et al.* 2020; Wang *et al.* 2020; Cui *et al.* 2021; Li *et al.* 2022).

The *Stx16* is involved in the development of inducible defense structures in *Daphnia pulex*

under predation stress (Spanier *et al.* 2010). High *FMVIA* expression in the retina of striped

bass suggests a role in retinal motility (Breckler *et al.* 2000). The *foxj1b* gene is essential

for normal laterality in zebrafish embryos, affecting heart and organ positioning (Tian *et al.*

2009). Finally, *Lamc3* plays a role in the development of the parachordal chain in zebrafish

(Eve & Smith 2017).

Expanding our search to adaptive loci, we identified three key genes: *vtg3* (linked to

temperature adaptation) and *slc39a6, slc8a2a* (associated with salinity adaptation). *Vtgs* are

specialized large lipid transfer proteins produced by the liver and transported to the ovary

(Babin *et al.* 2007). The *Vtg* family includes type-I (*Vtg 1, 4, 5, 6,* and *7*), type-II (*vtg2* and

vtg8), and type-III (*vtg3*) (Yilmaz *et al.* 2018). Despite being a minor and divergent form,

vtg3 significantly contributes to zygote and early embryo development (Yilmaz *et al.*

2019). Increased temperature elevates estrogen-induced *vtg* expression in juvenile *Salmo*

trutta (Körner *et al.* 2008), suggesting a similar role for *vtg3* in *E. maclovinus* related to

thermal adaptation. The gene *slc39a6 (LIV-1)* maintains zinc homeostasis, essential for

cellular growth and differentiation (Taylor & Nicholson 2003). Low salinity increases

slc39a6 expression in tiger puffer (*Takifugu rubripes*) (Jiang *et al.* 2020), and intracellular

zinc homeostasis controls T-cell survival in zebrafish by inhibiting apoptosis (Zhao *et al.*

2019). The *SLC8* gene family (*NCX*) maintains cellular Ca2+ homeostasis (Khananshvili

2013). *Slc8a2a*, an isoform identified in zebrafish, is key for Ca2+ excretion by seawater

fish kidneys, with upregulated transcription during seawater acclimation (Liao *et al.* 2007;

Islam *et al.* 2011). These genes indicate positive selection associated with salinity in *E.*

maclovinus in Northern Patagonia.

We did not identify genes whose functions are supposed to be affected by the

concentration of oxygen or nutrients (nitrate, phosphate and silicate). However, many

outlier sequences did not have BLASTx results and may represent important proteins that

have not been adequately annotated in fish to date.

The outcomes from the ten programs varied widely in terms of the number and

identity of adaptative loci detected in this study. Such variations among methods were

expected since their algorithms and prior assumptions considerably differ. Combining

population genetic differentiation (PGD) and genotype-environment association (GEA)

approaches to detect adaptative loci of local adaptation not only reduces false-positive

discoveries, but also maximizes our chances of detecting potential signals of selection

(François *et al.* 2016). Overall, we found that GEA approach identified more adaptative loci

(2,164 loci) than PGD approach (392 loci). These outcomes are in agreement with a

simulation study demonstrating that the GEA approach has more power to detect loci under

divergent selection than PGD approach (de Villemereuil *et al.* 2014), which is not

surprising given that GEA uses multiple environmental variables (six variables in our case)

to depict signals of selection. This trend has been reported in multiple studies focused on

detecting signs of local adaptation in marine populations (Benestan *et al.* 2016; Flanagan *et*

al. 2016; Dalongeville *et al.* 2018; Bekkevold *et al.* 2020; Cayuela *et al.* 2020; Segovia *et*

al. 2020).

Inside of PGD approach, *fsthet* detected the largest number on putative adaptative

loci (n= 313 loci). This software generates smoothed quantiles for the F_{ST}–heterozygosity

relationship from the empirical distribution (Flanagan & Jones 2017). Using an empirical

distribution itself has the advantage of not assuming a particular distribution or model of

evolution. For example, *FDIST2* and *LOSITAN* used an idealized island model, which

would not be appropriate considering the geography of Northern Patagonia. Besides, this

feature is appealing because the F_{ST}–heterozygosity distribution changes based on many

potentially unknowable features of population structure (Flanagan & Jones 2017). Though,

fsthet decreased the average proportion of false positives compared with *LOSITAN*, we did

not find any studies that compared the effectiveness of *fsthet* with respect to other software

that considers the hierarchical population structure, such as *Arlequin*. The *fsthet* program is

not recommended when migration rates are low because the neutral loci saturate the full

distribution of potential F_{ST}–heterozygosity values (Flanagan & Jones 2017). In our case,

the relative migration rates showed high gene flow between locations of *E. maclovinus*

(Canales-Aguirre *et al.* 2022). The *Arlequin* identified 270 loci (200 shared with *fsthet*).

However, results of a simulation study showing that *Arlequin* consistently found more

outliers and had the highest type I and type II errors in their simulation scenarios in

compared with other methods such as *BayeScan*. Simulated data suggested that *Arlequin*

does not perform well in situations where adaptive variation contrasts with patterns of

neutral variation, likely due to the effect of hierarchical grouping based on neutral markers

(Narum & Hess 2011). The *PCAdapt* was the more conservative (n= 15 loci), whereas

BayeScan and *OutFLANK* did not detect any outliers. Population structure and the presence

of admixture individuals affect the power of identifying adaptive loci with *Bayescan* more

than *PCAdapt* (Luu *et al.* 2017). This latter, agreed with our data where we found

admixture individuals in the whole SNPs database. The presence of mixed individuals is

supported by the high relative migration rates estimated by Canales-Aguirre *et al.* (2022).

Additionally, *BayeScan* could decrease power with few populations (n <30 Aguirre-Liguori

migration and/or in cases of smaller sample size (Foll & Gaggiotti 2008; Narum & Hess

2011).

Inside the GEA approach, *LFMM* detected the largest number of potentially adaptive

loci (n= 1,785 loci). Considering that *LFMM* identified 14.92% of the loci as adaptive, it is

likely that we will find false positives among them. The main reason is that geographic and

demographic processes can lead to patterns that mimic those observed due to selection. In

fact, de Villemereuil *et al*. (2014) found high rates of false discovery in some scenarios

with complex, hierarchical structures and polygenic selection; although showed that *LFMM*

provides the best compromise between detection power and error rates in situations with

complex hierarchical neutral genetic structure and polygenic selection. However, Frichot et

al. (2013) found that *LFMM* has low rates of false positives and negatives and that it

performs slightly better than *Bayenv* in detecting weak selection. Finally, Lotterhos &

Whitlock (2015) showed that *LFMM* is quite robust to various sampling designs and

underlying demographic models. The *MSOD* detected considerably fewer potentially

adaptive loci (n=377 loci). However, the main advantage of this program is that using

Moran spectral randomization (MSR) for significance testing, avoids the inflation of type I

errors due to spatial autocorrelation, whether resulting from gene flow or the spatial

structure of the environmental factors that induce selection (Wagner & Dray 2015; Wagner

et al. 2017). Moreover, spatial outlier detection worked best for scenarios of IBD, even in

the presence of large-scale spatial genetic structure (Wagner *et al.* 2017) as in this study.

LFMM and *MSOD*. The main difference is the multivariate approach of *RDA*, while *LFMM*

and *MSOD* are univariate, meaning that they test one genetic marker at a time, and may

also test only one environmental predictor at a time (Rellstab *et al.* 2015); *RDA* identifying

covarying loci associated with the multivariate environmental variables (Legendre &

Legendre 2012). The decrease in the number of loci detected by *RDA* is because it restricts

the number of environmental variables tested based on the variance inflation factor. In the

case of univariate programs, we tested six environmental variables with four statistics for

each but restricted to one statistic per environmental variable for *RDA*, limiting the number

of possible correlations. However, simulated data showed that *RDA* has higher true-positive

and lower false-positive rates compared to *PCAdapt* and *LFMM* (Forester *et al.* 2018;

Capblancq *et al.* 2018). Furthermore, *RDA* has one important limitation: it models linear

relationships among response and predictors, meaning that nonlinear relationships will not

be detected (Capblancq & Forester 2021). Finally, we did not detect any putative adaptive

loci using the *BayeScEnv* and *Samβada* programs. Considering that *BayeScEnv* produces

fewer candidate loci than *BayeScan* (de Villemereuil & Gaggiotti 2015), it is to be expected

that if *BayeScan* did not detect any loci *BayeScEnv* either. To date, we have not identified

any studies that compare the statistical power or control of FDR of *BayeScEnv* with respect

to other programs. Therefore, we cannot conclude whether *BayeScEnv* is more or less

sensitive to certain characteristics such as hierarchical population structure, isolation by

distance, or strength of selection. Due to the recentness of the *Samβada*, we did not identify

comparative studies that allow us to evaluate the advantages and disadvantages of this with

respect to other GEA approaches.

frameworks that include more population structure, multiple selection surfaces and genetic

architectures that are more complex than the multilocus selection response modelled here.

Overall, we highlight the importance of combining PGD and GEA approaches that consider

multiple scenarios of adaptive variation, to diminish type I and II errors and avoid conclude

erroneously.

4.5 Implications for management and conservation

The 11,961 filtered loci genotyped for 202 individuals (the largest population genomic data
set reported so far for *Eleginops maclovinus*) offers high resolution for creating
management policies and decisions, with respect to patterns of adaptive genetic variation.
Although neutral loci did not reveal a clear genetic structure among locations, analysis of
131 shared adaptive loci indicated genetic differentiation separating the populations into
four groups: REL, CHE, CHA, and a fourth group comprising the remaining seven
sampling locations. Note that all neutral pairwise F_{ST} values were close to zero, similar to
showed by Canales-Aguirre *et al.* (2022) and also from fish and other marine species
(Nayfa & Zenger 2016; Asaduzzaman *et al.* 2019b; a; O'Leary *et al.* 2021). The
interpretation of extremely small F_{ST} is not straightforward (Waples & Gaggiotti 2006;
Conover *et al.* 2006; Waples *et al.* 2008), meaning it is unclear what the implications of the
patterns of weak neutral differentiation may be for demographic connectivity of
subpopulations. Nevertheless, the signals of putatively adaptive genetic differentiation
between the ten locations were significant. Our study demonstrates how the environment
shapes the distribution of adaptive genetic variation across space for *E. maclovinus*;
showing that REL, CHE and CHA retain adaptive characteristics different from the rest.
This can have important implications for prioritizing sites for protection, where the level of
adaptive genetic variation could be an indicator of the evolutionary resilience of
populations (Bonin *et al.* 2007; Sgrò *et al.* 2011). Moreover, understanding the spatial
distribution of putatively adaptive genetic variation can inform the selection of specific
sites for protection within marine reserves to maintain the adaptive potential and
evolutionary resilience of wild populations faced with environmental change (von der
Heyden 2017). The adaptive divergence reported here appears to be underpinned by a
variation in seawater temperature and oxygen concentrations in the Northern Patagonia
(Iriarte *et al.* 2014; Torres *et al.* 2014; Pérez-Santos *et al.* 2014; Silva & Vargas 2014) and

complex patterns of population structure and putatively adaptive diversity among *E.*
maclovinus locations that we showed here are not recognized in current *E. maclovinus*
management units. To date, there is no management plan for *E. maclovinus* in Chile, only
regulations for the type of fishing gear used. Future research could use neutral and
adaptative datasets to determine management units using adaptative (e.g., population
adaptative index, adaptative score) and neutral (e.g., expected heterozygosity, local
differentiation) metrics as suggested by Xuereb et al. (2021). In the Chilean Northern
Patagonia, temperature and heat content are predicted to increase and freshwater river
inputs are predicted to decrease due to climate change (Aguayo *et al.* 2019). As such, the
spatial patterns of genetic variability observed in this study can be information conservation
planning decisions by identifying for protection populations containing higher levels of
segregating polymorphisms associated with environmental conditions (e.g., temperature
and salinity) that are prone to change in the future.

Literature cited

Aguayo R, León-Muñoz J, Vargas-Baecheler J *et al.* (2019) The glass half-empty: climate

change drives lower freshwater input in the coastal system of the Chilean Northern

Patagonia. *Climatic change*, **155**, 417–435.

Aguirre-Liguori JA, Luna-Sánchez JA, Gasca-Pineda J, Eguiarte LE (2020) Evaluation of

the Minimum Sampling Design for Population Genomic and Microsatellite Studies:

An Analysis Based on Wild Maize. *Frontiers in genetics*, **11**, 870.

Altshuler D, Pollara VJ, Cowles CR *et al.* (2000) An SNP map of the human genome

generated by reduced representation shotgun sequencing. *Nature*, **407**, 513–516.

Andrews KR, Good JM, Miller MR, Luikart G, Hohenlohe PA (2016) Harnessing the

power of RADseq for ecological and evolutionary genomics. *Nature reviews.*

Genetics, **17**, 81–92.

Aracena C, Lange CB, Iriarte JL, Rebolledo L, Pantoja S (2011) Latitudinal patterns of

export production recorded in surface sediments of the Chilean Patagonian fjords

(41--55 S) as a response to water column productivity. *Continental shelf research*,

31, 340–355.

Chilean blue mussels *(Mitilus chilensis)* from different marine environments. *Ecology and evolution*, **6**, 3632–3644.

Asaduzzaman M, Igarashi Y, Wahab MA *et al.* (2019a) Population Genomics of an Anadromous Hilsa Shad *Tenualosa ilisha* Species across Its Diverse Migratory Habitats: Discrimination by Fine-Scale Local Adaptation. *Genes*, **11**.

Asaduzzaman M, Wahab MA, Rahman MJ *et al.* (2019b) Fine-scale population structure and ecotypes of anadromous Hilsa shad *(Tenualosa ilisha)* across complex aquatic ecosystems revealed by NextRAD genotyping. *Scientific reports*, **9**, 16050.

Assis J, Tyberghein L, Bosch S (2018) Bio☐ORACLE v2. 0: Extending marine data layers for bioclimatic modelling. *Global Ecology and*.

Athauda S, Anderson T, de Nys R (2012) Effect of rearing water temperature on protandrous sex inversion in cultured Asian Seabass *(Lates calcarifer)*. *General and comparative endocrinology*, **175**, 416–423.

Babin PJ, Cerdà J, Lubzens E (2007) *The Fish Oocyte: From Basic Studies to Biotechnological Applications*. Springer Science & Business Media.

Babin C, Gagnaire P-A, Pavey SA, Bernatchez L (2017) RAD-Seq Reveals Patterns of Additive Polygenic Variation Caused by Spatially-Varying Selection in the American Eel (Anguilla rostrata). *Genome biology and evolution*, **9**, 2974–2986.

Baird NA, Etter PD, Atwood TS *et al.* (2008) Rapid SNP discovery and genetic mapping using sequenced RAD markers. *PloS one*, **3**, e3376.

Baroiller J-F, Guiguen Y, Fostier A (1999) Endocrine and environmental aspects of sex

differentiation in fish. *Cellular and molecular life sciences: CMLS*, **55**, 910–931.

Barrett RDH, Hoekstra HE (2011) Molecular spandrels: tests of adaptation at the genetic

level. *Nature reviews. Genetics*, **12**, 767–780.

Beaumont MA (2005) Adaptation and speciation: what can Fst tell us? *Trends in ecology &*

evolution, **20**, 435–440.

Beirão J, Lewis JA, Wringe BF, Purchase CF (2018) A novel sperm adaptation to

evolutionary constraints on reproduction: Pre ejaculatory sperm activation in the

beach spawning capelin (Osmeridae). *Ecology and evolution*, **8**, 2343–2349.

Bekkevold D, Höjesjö J, Nielsen EE *et al.* (2020) Northern European *Salmo trutta (L.)*

populations are genetically divergent across geographical regions and

environmental gradients. *Evolutionary applications*, **13**, 400–416.

Benestan L, Quinn BK, Maaroufi H *et al.* (2016) Seascape genomics provides evidence for

thermal adaptation and current-mediated population structure in American lobster

(Homarus americanus). Molecular ecology, **25**, 5073–5092.

Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate: A practical and

powerful approach to multiple testing. *Journal of the Royal Statistical Society*, **57**,

289–300.

Benoît HP, Pepin P, Brown JA (2000) Patterns of metamorphic age and length in marine

fishes, from individuals to taxa. *Canadian journal of fisheries and aquatic sciences.*

Journal canadien des sciences halieutiques et aquatiques, **57**, 856–869.

environmental change: considerations from population genomics in fishes. *Journal*

of fish biology, **89**, 2519–2556.

Blaxter (1992) The effect of temperature on larval fishes. *Netherlands Journal of Zoology*.

Bonin A, Nicole F, Pompanon F, Miaud C, Taberlet P (2007) Population adaptive index: a

new method to help measure intraspecific genetic diversity and prioritize

populations for conservation. *Conservation biology: the journal of the Society for*

Conservation Biology, **21**, 697–708.

Boulanger E, Benestan L, Guerin P-E *et al.* (2022) Climate differently influences the

genomic patterns of two sympatric marine fish species. *The Journal of animal*

ecology, **91**, 1180–1195.

Breckler J, Au K, Cheng J, Hasson T, Burnside B (2000) Novel myosin VI isoform is

abundantly expressed in retina. *Experimental eye research*, **70**, 121–134.

Brian JV, Harris CA, Runnalls TJ *et al.* (2008) Evidence of temperature-dependent effects

on the estrogenic response of fish: implications with regard to climate change. *The*

Science of the total environment, **397**, 72–81.

Brickle P, Laptikhovsky V, Arkhipkin A (2005) Reproductive strategy of a primitive

temperate notothenioid *Eleginops maclovinus*. *Journal of fish biology*, **66**, 1044–

1059.

Canales-Aguirre CB, Ferrada-Fuentes S, Galleguillos R *et al.* (2018) High genetic diversity

and low-population differentiation in the Patagonian sprat *(Sprattus fuegensis)*

based on mitochondrial DNA. *Mitochondrial DNA. Part A, DNA mapping, sequencing, and analysis*, **29**, 1148–1155.

Canales-Aguirre CB, Ferrada-Fuentes S, Galleguillos R, Hernández CE (2016) Genetic Structure in a Small Pelagic Fish Coincides with a Marine Protected Area: Seascape Genetics in Patagonian Fjords. *PloS one*, **11**, e0160670.

Canales-Aguirre CB, Larson WA, McKinney GJ *et al.* (2022) Neutral and adaptive loci reveal fine-scale population structure in *Eleginops maclovinus* from north Patagonia. *Ecology and evolution*, **12**, e9343.

Capblancq T, Forester BR (2021) Redundancy analysis: A Swiss Army Knife for landscape genomics. *Methods in ecology and evolution / British Ecological Society*.

Capblancq T, Luu K, Blum MGB, Bazin E (2018) Evaluation of redundancy analysis to identify signatures of local adaptation. *Molecular ecology resources*, **18**, 1223–1233.

Catchen JM, Hohenlohe PA, Bernatchez L *et al.* (2017) Unbroken: RADseq remains a powerful tool for understanding the genetics of adaptation in natural populations. *Molecular ecology resources*, **17**, 362–365.

Cattell RB (1966) The Scree Test For The Number Of Factors. *Multivariate behavioral research*, **1**, 245–276.

Cayuela H, Rougemont Q, Laporte M *et al.* (2020) Shared ancestral polymorphisms and chromosomal rearrangements as potential drivers of local adaptation in a marine fish. *Molecular ecology*, **29**, 2379–2398.

Ceballos SG, Lessa EP, Licandeo R, Fernández DA (2016) Genetic relationships between

Atlantic and Pacific populations of the notothenioid fish *Eleginops maclovinus*: the

footprints of Quaternary glaciations in Patagonia. *Heredity*, **116**, 372–377.

Ceballos SG, Lessa EP, Victorio MF, Fernández DA (2012) Phylogeography of the sub-

Antarctic notothenioid fish *Eleginops maclovinus*: evidence of population

expansion. *Marine biology*, **159**, 499–505.

Chen Z, Cheng C-HC, Zhang J *et al.* (2008) Transcriptomic and genomic evolution under

constant cold in Antarctic notothenioid fish. *Proceedings of the National Academy*

of Sciences of the United States of America, **105**, 12944–12949.

Chen L, Lu Y, Li W *et al.* (2019) The genomic basis for colonizing the freezing Southern

Ocean revealed by Antarctic toothfish and Patagonian robalo genomes.

GigaScience, **8**.

Cheng C-HC, Chen L, Near TJ, Jin Y (2003) Functional antifreeze glycoprotein genes in

temperate-water New Zealand nototheniid fish infer an Antarctic evolutionary

origin. *Molecular biology and evolution*, **20**, 1897–1908.

Conesa A, Götz S (2008) Blast2GO: A comprehensive suite for functional analysis in plant

genomics. *International journal of plant genomics*, **2008**, 619832.

Conover DO, Clarke LM, Munch SB, Wagner GN (2006) Spatial and temporal scales of

adaptive divergence in marine fishes and the implications for conservation. *Journal*

of fish biology, **69**, 21–47.

Coppola D, Abbruzzetti S, Nicoletti F *et al.* (2012) ATP regulation of the ligand-binding

properties in temperate and cold-adapted haemoglobins. X-ray structure and ligand-

912  *bioSystems*, **8**, 3295–3304.

913 Coppola D, Giordano D, Vergara A *et al.* (2010) The hemoglobins of sub-Antarctic fishes

914  of the suborder Notothenioidei. *Polar science*, **4**, 295–308.

915 Cui Y, Yin K, Zheng Y *et al.* (2021) Mixed plasticizers aggravated apoptosis by NOD2-

916  RIP2-NF-κB pathway in grass carp hepatocytes. *Journal of hazardous materials*,

917  **402**, 123527.

918 Cure K, Thomas L, Hobbs J-PA, Fairclough DV, Kennington WJ (2017) Genomic

919  signatures of local adaptation reveal source-sink dynamics in a high gene flow fish

920  species. *Scientific reports*, **7**, 8618.

921 Cuvier G, Valenciennes A (1830) *Histoire naturelle des poissons*. F.-G. Levrault.

922 Czypionka T, Vargas CA, Silva N *et al.* (2011) Importance of mixotrophic nanoplankton in

923  Aysén Fjord (Southern Chile) during austral winter. *Continental shelf research*, **31**,

924  216–224.

925 Dabney A, Storey JD, Warmes GR (2010) *qvalue: Q-value estimation for false discovery*

926  *rate control*.

927 Dalongeville A, Benestan L, Mouillot D, Lobreaux S, Manel S (2018) Combining six

928  genome scan methods to detect candidate genes to salinity in the Mediterranean

929  striped red mullet *(Mullus surmuletus). BMC genomics*, **19**, 1–13.

930 Davey JW, Cezard T, Fuentes-Utrilla P *et al.* (2013) Special features of RAD Sequencing

931  data: implications for genotyping. *Molecular ecology*, **22**, 3151–3164.

relation with the salinity distribution off austral Chile (35–55°S). *Continental shelf*

research, **22**, 521–534.

Delgado ML, Górski K, Habit E, Ruzzante DE (2019) The effects of diadromy and its loss

on genomic divergence: The case of amphidromous *Galaxias maculatus*

populations. *Molecular ecology*, **28**, 5217–5231.

Devlin B, Roeder K (1999) Genomic control for association studies. *Biometrics*, **55**, 997–

1004.

Diopere E, Vandamme SG, Hablützel PI *et al.* (2017) Seascape genetics of a flatfish reveals

local selection under high levels of gene flow. *ICES journal of marine science:*

journal du conseil, **75**, 675–689.

Dong XY, Qin JG, Zhang XM (2011) Fish adaptation to oxygen variations in aquaculture

from hypoxia to hyperoxia. *African journal of fisheries and aquaculture*, **2**, 23.

Dorwart MR, Shcheynikov N, Yang D, Muallem S (2008) The solute carrier 26 family of

proteins in epithelial ion transport. *Physiology* , **23**, 104–114.

Dowling DK, Simmons LW (2009) Reactive oxygen species as universal constraints in life-

history evolution. *Proceedings of the Biological sciences / The Royal Society*, **276**,

1737–1745.

Dray S, Bauman D, Blanchet G *et al.* (2021) *adespatial: Multivariate Multiscale Spatial*

Analysis. Github.

Dray S, Legendre P, Peres-Neto PR (2006) Spatial modelling: a comprehensive framework

for principal coordinate analysis of neighbour matrices (PCNM). *Ecological*

modelling, **196**, 483–493.

Duruz S, Sevane N, Selmoni O *et al.* (2019) Rapid identification and interpretation of gene-

environment associations using the new R.SamBada landscape genomics pipeline.

Molecular ecology resources, **19**, 1355–1365.

Eastman JT (1993) *Antarctic Fish Biology: Evolution in a Unique Environment*. Academic

Press.

Evanno G, Regnaut S, Goudet J (2005) Detecting the number of clusters of individuals

using the software STRUCTURE: a simulation study. *Molecular ecology*, **14**, 2611–

2620.

Eve AMJ, Smith JC (2017) Knockdown of Laminin gamma-3 (Lamc3) impairs motoneuron

guidance in the zebrafish embryo. *Wellcome open research*, **2**, 111.

Excoffier L, Lischer HEL (2010) Arlequin suite ver 3.5: a new series of programs to

perform population genetics analyses under Linux and Windows. *Molecular*

ecology resources, **10**, 564–567.

Fago A, Forest E, Weber RE (2001) Hemoglobin and subunit multiplicity in the rainbow

trout *(Oncorhynchus mykiss)* hemoglobin system. *Fish physiology and*

biochemistry, **24**, 335–342.

Feuerlein R, Weber R (1994) Rapid and simultaneus measurement of anodic and cathodic

haemoglobins and ATP and GTP concentrations in minute quantities of fish blood.

The Journal of experimental biology, **189**, 273–277.

Flanagan SP, Jones AG (2017) Constraints on the FST Heterozygosity Outlier Approach.

The Journal of heredity, **108**, 561–573.

Flanagan SP, Rose E, Jones AG (2016) Population genomics reveals multiple drivers of

population differentiation in a sex-role-reversed pipefish. *Molecular ecology*, **25**,

5043–5072.

Foll M, Gaggiotti O (2008) A genome-scan method to identify selected loci appropriate for

both dominant and codominant markers: a Bayesian perspective. *Genetics*, **180**,

977–993.

Forester BR, Lasky JR, Wagner HH, Urban DL (2018) Comparing methods for detecting

multilocus adaptation with multivariate genotype-environment associations.

Molecular ecology, **27**, 2215–2233.

Fox MD, Williams GJ, Johnson MD *et al.* (2018) Gradients in Primary Production Predict

Trophic Strategies of Mixotrophic Corals across Spatial Scales. *Current biology:*

CB, **28**, 3355-3363.e4.

Francis RM (2017) pophelper: an R package and web app to analyse and visualize

population structure. *Molecular ecology resources*, **17**, 27–32.

François O, Martins H, Caye K, Schoville SD (2016) Controlling false discoveries in

genome scans for selection. *Molecular ecology*, **25**, 454–469.

Frichot E, François O (2015) LEA: An R package for landscape and ecological association

studies. *Methods in ecology and evolution / British Ecological Society*, **6**, 925–929.

Frichot E, Mathieu F, Trouillon T, Bouchard G, François O (2014) Fast and efficient

estimation of individual ancestry coefficients. *Genetics*, **196**, 973–983.

loci and environmental gradients using latent factor mixed models. *Molecular*

biology and evolution, **30**, 1687–1699.

Gagliano M, McCormick MI, Meekan MG (2007) Temperature-induced shifts in selective

pressure at a critical developmental transition. *Oecologia*, **152**, 219–225.

Gagnaire P-A, Broquet T, Aurelle D *et al.* (2015) Using neutral , selected , and hitchhiker

loci to assess connectivity of marine populations in the genomic era. *Evolutionary*

applications, **8**, 769–786.

Gagnaire P-A, Gaggiotti OE (2016) Detecting polygenic selection in marine populations by

combining population genomics and quantitative genetics approaches. *Current*

zoology, **62**, 603–616.

Gillet C, Breton B, Mikolajczyk T (1996) Effects of GnRHa and pimozide treatments on

the timing of ovulation and on egg quality in Arctic charr (Salvelinus alpinus) at 5

and 10 °C. *Aquatic living resources*, **9**, 257–263.

Ginsburg AS (1963) Sperm-egg association and its relationship to the activation of the egg

in salmonid fishes. *Journal of embryology and experimental morphology*, **11**, 13–

33.

González HE, Castro L, Daneri G *et al.* (2011) Seasonal plankton variability in Chilean

Patagonia fjords: Carbon flow through the pelagic food web of Aysen Fjord and

plankton dynamics in the Moraleda Channel basin. *Continental shelf research*, **31**,

225–243.

Gosztonyi AE, Ro~~ G (1974) Edad y crecimiento del "Robalo" *Eleginops maclovinus*

(Osteichthyes, Nototheniidae) en aguas de la ría Deseado y sus adyacencias.

Goudet J (2005) hierfstat, a package for r to compute and test hierarchical F-statistics.

Molecular ecology notes, **5**, 184–186.

Gove JM, McManus MA, Neuheimer AB *et al.* (2016) Near-island biological hotspots in

barren ocean basins. *Nature communications*, **7**, 10581.

Haasl RJ, Payseur BA (2016) Fifteen years of genomewide scans for selection: trends,

lessons and unaddressed genetic sources of complication. *Molecular ecology*, **25**, 5–

23.

Hair JF, Anderson RE, Tatham RL, and Black WC (1995) *Multivariate data analysis with*

readings. Prentice Hall, Englewood Cliffs, N.J.

Herberg S, Gert KR, Schleiffer A, Pauli A (2018) The Ly6/uPAR protein Bouncer is

necessary and sufficient for species-specific fertilization. *Science*, **361**, 1029–1033.

Herkenne S, Ek O, Zamberlan M *et al.* (2020) Developmental and Tumor Angiogenesis

Requires the Mitochondria-Shaping Protein Opa1. *Cell metabolism*, **31**, 987-

1003.e8.

von der Heyden S (2017) Making evolutionary history count: biodiversity planning for

coral reef fishes and the conservation of evolutionary processes. *Coral reefs* , **36**,

183–194.

Hoban S, Kelley JL, Lotterhos KE *et al.* (2016) Finding the Genomic Basis of Local

Adaptation: Pitfalls, Practical Solutions, and Future Directions. *The American*

naturalist, **188**, 379–397.

Hoey JA, Pinsky ML (2018) Genomic signatures of environmental selection despite near

panmixia in summer flounder. *Evolutionary applications*, **11**, 1732–1747.

Hokanson KEF (2011) Temperature Requirements of Some Percids and Adaptations to the

Seasonal Temperature Cycle. *Journal of the Fisheries Research Board of Canada*.

Holliday F (1969) 4 the effects of salinity on the eggs and larvae of teleosts. *Fish

Physiology*, **1**, 293–311.

Houde ED (1989) Comparative growth, mortality and energetic of marine fish larvae:

temperature an implied latitudinal effects. *Fishery bulletin* , 471–495.

Iriarte JL, González HE, Liu KK, Rivas C, Valenzuela C (2007) Spatial and temporal

variability of chlorophyll and primary productivity in surface waters of southern

Chile (41.5--43 S). *Estuarine, coastal and shelf science*, **74**, 471–480.

Iriarte JL, Pantoja S, Daneri G (2014) Oceanographic Processes in Chilean Fjords of

Patagonia: from small to large-scale studies. *Progress in Oceanography*, **129**, 1–7.

Islam Z, Kato A, Romero MF, Hirose S (2011) Identification and apical membrane

localization of an electrogenic Na^+/Ca^{2+} exchanger NCX2a likely to be involved in

renal Ca^{2+} excretion by seawater fish. *American journal of physiology. Regulatory,

integrative and comparative physiology*, **301**, R1427-39.

Jacob BG, Tapia FJ, Daneri G *et al.* (2014) Springtime size-fractionated primary

production across hydrographic and PAR-light gradients in Chilean Patagonia (41–

50°S). *Progress in oceanography*, **129**, 75–84.

Jaimanch M, Piironen J, Yeánkari N (2010) Parental effects on embryonic viability and

growth in Arctic charr *Salvelinus alpinus* at two incubation temperatures. *Journal of*

fish biology, **76**, 2558–2570.

Jensen FB, Fago A, Weber RE (1998) Hemoglobin structure and function. *Fish Physiology*,

17, 1–40.

Jensen FB, Weber RE (1985) Proton and oxygen equilibria, their anion sensitivities and

interrelationships in tench hemoglobin. *Molecular physiology*, **7**, 41–50.

Jiang J-L, Xu J, Ye L *et al.* (2020) Identification of differentially expressed genes in gills of

tiger puffer *(Takifugu rubripes)* in response to low-salinity stress. *Comparative*

biochemistry and physiology. Part B, Biochemistry & molecular biology, **243–244**,

110437.

Jombart T (2008) adegenet: a R package for the multivariate analysis of genetic markers.

Bioinformatics , **24**, 1403–1405.

Jombart T, Ahmed I (2011) adegenet 1.3-1: new tools for the analysis of genome-wide SNP

data. *Bioinformatics* , **27**, 3070–3071.

Jombart T, Devillard S, Balloux F (2010) Discriminant analysis of principal components: a

new method for the analysis of genetically structured populations. *BMC genetics*,

11, 94.

Khananshvili D (2013) The SLC8 gene family of sodium-calcium exchangers (NCX) -

structure, function, and regulation in health and disease. *Molecular aspects of*

medicine, **34**, 220–235.

Kock K-H (1992) *Antarctic Fish and Fisheries*. Cambridge University Press.

Kömer O, Konno S, Schönenberger R *et al.* (2008) Water temperature and concomitant

waterborne ethinylestradiol exposure affects the vitellogenin expression in juvenile

brown trout *(Salmo trutta). Aquatic toxicology* , **90**, 188–196.

Kültz D (2015) Physiological mechanisms used by fish to cope with salinity stress. *The*

Journal of experimental biology, **218**, 1907–1914.

Lattuca ME, Boy CC, Vanella FA, Barrantes ME, Fernández DA (2018) Thermal responses

of three native fishes from estuarine areas of the Beagle Channel, and their

implications for climate change. *Hydrobiologia*, **808**, 235–249.

Laverty G, Skadhauge E (2012) Adaptation of teleosts to very high salinity. *Comparative*

biochemistry and physiology. Part A, Molecular & integrative physiology, **163**, 1–6.

Legendre P, Legendre L (2012) *Numerical Ecology*. Elsevier.

Lehtonen TK, Kvarnemo C (2015) Density effects on fish egg survival and infections

depend on salinity. *Marine ecology progress series*, **540**, 183–191.

Lewontin RC, Krakauer J (1973) Distribution of gene frequency as a test of the theory of

the selective neutrality of polymorphisms. *Genetics*, **74**, 175–195.

Lex A, Gehlenborg N, Strobelt H, Vuillemot R, Pfister H (2014) UpSet: Visualization of

Intersecting Sets. *IEEE transactions on visualization and computer graphics*, **20**,

1983–1992.

Li Y-L, Xue D-X, Zhang B-D, Liu J-X (2019) Population Genomic Signatures of Genetic

Structure and Environmental Selection in the Catadromous Roughskin Sculpin

Trachidermus fasciatus. Genome biology and evolution, **11**, 1751–1764.

Li J, Zhang W, Zhou P *et al.* (2022) Selenium deficiency induced apoptosis via

mitochondrial pathway caused by Oxidative Stress in porcine gastric tissues.

Research in veterinary science, **144**, 142–148.

Liao B-K, Deng A-N, Chen S-C, Chou M-Y, Hwang P-P (2007) Expression and water

calcium dependence of calcium transporter isoforms in zebrafish gill

mitochondrion-rich cells. *BMC genomics*, **8**, 354.

Linford P, Pérez-Santos I, Montero P *et al.* (2023a) Oceanographic processes favoring

deoxygenation inside Patagonian fjords.

Linford P, Pérez-Santos I, Montero P *et al.* (2024) Oceanographic processes driving low-

oxygen conditions inside Patagonian fjords. *Biogeosciences*, **21**, 1433–1459.

Linford P, Pérez-Santos I, Montes I *et al.* (2023b) Recent deoxygenation of Patagonian

fjord subsurface waters connected to the Peru–Chile undercurrent and equatorial

subsurface water variability. *Global biogeochemical cycles*, **37**.

Lischer HEL, Excoffier L (2012) PGDSpider: an automated data conversion tool for

connecting population genetics and genomics programs. *Bioinformatics* , **28**, 298–

299.

Luu K, Bazin E, Blum MGB (2017) pcadapt: an R package to perform genome scans for

selection based on principal component analysis. *Molecular ecology resources*, **17**,

67–77.

Manel S, Perrier C, Pratlong M *et al.* (2016) Genomic resources and their influence on the

detection of the signal of positive selection in genome scans. *Molecular ecology*, **25**,

170–184

Maschko J, Andrews RK, Hohenlohe PA (2020) Long-lived marine species may be resilient

to environmental variability through a temporal portfolio effect. *Ecology and*

evolution, **10**, 6435–6448.

McKinney GJ, Larson WA, Seeb LW, Seeb JE (2017a) RADseq provides unprecedented

insights into molecular ecology and evolutionary genetics: comment on Breaking

RAD by Lowry et al. (2016). *Molecular ecology resources*, **17**, 356–361.

McKinney G, McPhee MV, Pascal C, Seeb JE, Seeb LW (2020) Network Analysis of

Linkage Disequilibrium Reveals Genome Architecture in Chum Salmon. *G3* , **10**,

1553–1561.

McKinney GJ, Waples RK, Seeb LW, Seeb JE (2017b) Paralogs are revealed by proportion

of heterozygotes and deviations in read ratios in genotyping□by□sequencing data

from natural populations. *Molecular Ecology*, **17**, 656–669.

Miller MR, Dunham JP, Amores A, Cresko WA (2007) Rapid and cost-effective

polymorphism identification and genotyping using restriction site associated DNA

(RAD) markers. *Genome / National Research Council Canada = Genome / Conseil*

national de recherches Canada.

Montero P, Daneri G, González HE *et al.* (2011) Seasonal variability of primary production

in a fjord ecosystem of the Chilean Patagonia: Implications for the transfer of

carbon within pelagic food webs. *Continental shelf research*, **31**, 202–215.

Montero P, Daneri G, Tapia F, Iriarte JL, Crawford D (2017) Diatom blooms and primary

production in a channel ecosystem of central Patagonia. *Latin American journal of*

aquatic research, **45**, 999–1016.

Narum SR, Hess JE (2011) Comparison of F(ST) outlier tests for SNP loci under selection.

Molecular ecology resources, **11 Suppl 1**, 184–194.

Nayfa MG, Zenger KR (2016) Unravelling the effects of gene flow and selection in highly

connected populations of the silver-lip pearl oyster (Pinctada maxima). *Marine*

genomics, **28**, 99–106.

Nikinmaa M, Rees BB (2005) Oxygen-dependent gene expression in fishes. *American*

journal of physiology. Regulatory, integrative and comparative physiology, **288**,

R1079-90.

North AW (2001) Early life history strategies of notothenioids at South Georgia. *Journal of*

fish biology, **58**, 496–505.

O, Wai-sum, Chen H, Chow PH (2006) Male genital tract antioxidant enzymes—Their

ability to preserve sperm DNA integrity. *Molecular and cellular endocrinology*,

250, 80–83.

Oksanen J, Guillaume F, Blanchet M *et al.* (2016) *vegan: Community Ecology Package*.

Okuzawa K, Kumakura N, Gen K *et al.* (2003) Effect of high water temperature on brain-

pituitary-gonad axis of the red seabream during its spawning season. *Fish*

physiology and biochemistry, **28**, 439–440.

O'Leary SJ, Hollenbeck CM, Vega RR, Portnoy DS (2021) Disentangling complex

genomic signals to understand population structure of an exploited,

estuarine‐dependent flatfish. *Ecology and evolution*.

Oyarzún R, Muñoz JLP, Pontigo JP, Morera FJ, Vargas-Chacoff L (2018) Effects of

acclimation to high environmental temperatures on intermediary metabolism and

1168 osmoregulation in the sub-Antarctic notothenioid *Eleginops maclovinus. Marine*

1169  *biology*, **165**, 1–15.

1170 Oyarzún R, Rojas JJ, Pontigo JP *et al.* (2021) Long-term effects of temperatures on the

1171  physiological response of juveniles of the eurythermal sub-antarctic notothenioid

1172  *Eleginops maclovinus. Aquaculture* , **530**, 1–11.

1173 Pankhurst NW, King HR (2010) Temperature and salmonid reproduction: implications for

1174  aquaculture. *Journal of fish biology*, **76**, 69–85.

1175 Pantoja S, Iriarte J, Daneri G (2011) Oceanography of the Chilean Patagonia. *Continental*

1176  *shelf research*, **31**, 149–153.

1177 Paredes A, Montecino V (2011) Size diversity as an expression of phytoplankton

1178  community structure and the identification of its patterns on the scale of fjords and

1179  channels. *Continental shelf research*, **31**, 272–281.

1180 Pauls SU, Nowak C, Bálint M, Pfenninger M (2013) The impact of global climate change

1181  on genetic diversity within populations and species. *Molecular ecology*, **22**, 925–

1182  946.

1183 Pauly D, Pullin RSV (1988) Hatching time in spherical, pelagic, marine fish eggs in

1184  response to temperature and egg size. *Environmental biology of fishes*, **22**, 261–271.

1185 Pembleton LW, Cogan NOI, Forster JW (2013) StAMPP: an R package for calculation of

1186  genetic differentiation and structure of mixed-ploidy level populations. *Molecular*

1187  *ecology resources*, **13**, 946–952.

Pequeño GH (1989) The geographical distribution and taxonomic arrangement of South American notothenidae fishes (Osteichthyes, Notothenidae). *Bol. Soc. Biol. Concepción, Chile*, **60**, 183–200.

Pérez-Santos I, Garcés-Vargas J, Schneider W *et al.* (2014) Double-diffusive layering and mixing in Patagonian fjords. *Progress in oceanography*, **129**, 35–49.

Pickard GL (1971) Some physical oceanographic features of inlets of Chile. *Journal of the Fisheries Research Board of Canada*, **28**, 1077–1106.

Pizarro G, Iriarte JL, Montecino V, Blanco JL, Guzmán L (2000) Distribución de la biomasa fitoplanctónica y productividad primaria máxima de fiordos y canales australes (47–50 S) en octubre 1996. *Ciencia y Tecnologia del Mar* .

Powers DA (1974) Structure, function, and molecular ecology of fish hemoglobins. *Annals of the New York Academy of Sciences*, **241**, 472–490.

di Prisco G, Maresca B, Tota B (2012) *Biology of Antarctic Fish*. Springer Science & Business Media.

di Prisco G, Tamburrini M (1992) The hemoglobins of marine and freshwater fish: the search for correlations with physiological adaptation. *Comparative biochemistry and physiology. B, Comparative biochemistry*, **102**, 661–671.

Pritchard JK, Stephens M, Donnelly P (2000) Inference of Population Structure Using Multilocus Genotype Data. *Genetics*, **155**, 945–959.

Rabassa J, Coronato A, Martínez O (2011) Late Cenozoic glaciations in Patagonia and Tierra del Fuego: an updated review. *Biological journal of the Linnean Society. Linnean Society of London*, **103**, 316–335.

Rellstab C, Gugerli F, Eckert AJ, Hancock AM, Holderegger R (2015) A practical guide to

environmental association analysis in landscape genomics. *Molecular ecology*, **24**,

4348–4370.

Richards JG (2009) Chapter 10 Metabolic and Molecular Responses of Fish to Hypoxia. In:

Fish Physiology (eds Richards JG, Farrell AP, Brauner CJ), pp. 443–485. Academic

Press.

Rodrigo C (2008) Submarine topography in the Chilean North Patagonian channels.

Progress in the oceanographic knowledge of Chilean inner waters, from Puerto

Montt to Cape Horn. Comité Oceanográfico Nacional-Pontificia Universidad

Católica de Valparaíso, Valparaíso, 19–23.

Rombough PJ (1997) The effects of temperature on embryonic and larval development. In:

Global Warming: Implications for Freshwater and Marine Fish, pp. 177–224.

Cambridge University Press.

Rungruangsak-Torrissen K (2007) Digestive efficiency, growth and qualities of muscle and

oocyte in Atlantic salmon *(Salmo Salar L.)* fed on diets with krill meal as an

alternative protein source. *Journal of food biochemistry*, **31**, 509–540.

Rungruangsak-Torrissen K, Moss R, Andresen LH, Berg A, Waagbø R (2006) Different

expressions of trypsin and chymotrypsin in relation to growth in Atlantic salmon

(Salmo salar L.). *Fish physiology and biochemistry*, **32**, 7–23.

Schlitzer R (2002) Interactive analysis and visualization of geoscience data with Ocean

Data View. *Computers & geosciences*, **28**, 1211–1218.

Schneider W, Donoso D, Garcés-Vargas J, Escribano R (2017) Water column cooling and

sea surface salinity increase in the upwelling region off central-south Chile driven

by a poleward displacement of the South Pacific High. *Progress in oceanography*,

151, 38–48.

Schneider W, Pérez-Santos I, Ross L *et al.* (2014) On the hydrography of Puyuhuapi

Channel, Chilean Patagonia. *Progress in oceanography*, **129**, 8–18.

Scott SG, Pankhurst NW (1992) Interannual variation in the reproductive cycle of the New

Zealand snapper *Pagrus auratus* (Bloch & Schneider) (Sparidae). *Journal of fish

biology*, **41**, 685–696.

Segovia NI, González-Wevar CA, Haye PA (2020) Signatures of local adaptation in the

spatial genetic structure of the ascidian *Pyura chilensis* along the southeast Pacific

coast. *Scientific reports*, **10**, 14098.

Sgrò CM, Lowe AJ, Hoffmann AA (2011) Building evolutionary resilience for conserving

biodiversity under climate change. *Evolutionary applications*, **4**, 326–337.

Shimizu A (2003) Effect of photoperiod and temperature on gonadal activity and plasma

steroid levels in a reared strain of the mummichog *(Fundulus heteroclitus)* during

different phases of its annual reproductive cycle. *General and comparative

endocrinology*, **131**, 310–324.

Silva C, Andrade I, Yáñez E *et al.* (2016) Predicting habitat suitability and geographic

distribution of anchovy (Engraulis ringens) due to climate change in the coastal

areas off Chile. *Progress in oceanography*, **146**, 159–174.

Silva N, Vargas CA (2014) Hypoxia in Chilean Patagonian Fjords. *Progress in*

oceanography, **129**, 62–74.

Smith GD, Hopkins GR, Mohammadi S *et al.* (2015) Effects of temperature on embryonic

and early larval growth and development in the rough-skinned newt *(Taricha*

granulosa). Journal of thermal biology, **51**, 89–95.

Song X-B, Liu G, Liu F *et al.* (2017) Autophagy blockade and lysosomal membrane

permeabilization contribute to lead-induced nephrotoxicity in primary rat proximal

tubular cells. *Cell death & disease*, **8**, e2863.

Soria FN, Strüssmann CA, Miranda LA (2008) High water temperatures impair the

reproductive ability of the pejerrey fish *Odontesthes bonariensis*: effects on the

hypophyseal-gonadal axis. *Physiological and biochemical zoology: PBZ*, **81**, 898–

905.

Spanier KI, Leese F, Mayer C *et al.* (2010) Predator-induced defences in *Daphnia pulex*:

selection and evaluation of internal reference genes for gene expression studies with

real-time PCR. *BMC molecular biology*, **11**, 50.

Stucki S, Orozco-terWengel P, Forester BR *et al.* (2017) High performance computation of

landscape genomic models including local indicators of spatial association.

Molecular ecology resources, **17**, 1072–1089.

Sumpter JP, Jobling S (1995) Vitellogenesis as a biomarker for estrogenic contamination of

the aquatic environment. *Environmental health perspectives*, **103 Suppl 7**, 173–178.

Taylor CT, McElwain JC (2010) Ancient atmospheres and the evolution of oxygen sensing

via the hypoxia-inducible factor in metazoans. *Physiology*, **25**, 272–279.

Taylor KM, Nicholson RI (2003) The ZIP proteins, the LIV-1 subfamily of zinc

transporters. *Biochimica et biophysica acta*, **1611**, 16–30.

Tian T, Zhao L, Zhang M, Zhao X, Meng A (2009) Both foxj1a and foxj1b are implicated

in left-right asymmetric development in zebrafish embryos. *Biochemical and*

biophysical research communications, **380**, 537–542.

Torres R, Silva N, Reid B, Frangopulos M (2014) Silicic acid enrichment of subantarctic

surface water from continental inputs along the Patagonian archipelago interior sea

(41--56 S). *Progress in oceanography*, **129**, 50–61.

Tveiten H, Solevag SE, Johnsen HK (2001) Holding temperature during the breeding

season influences final maturation and egg quality in common wolffish. *Journal of*

fish biology, **58**, 374–385.

Tyberghein L, Verbruggen H, Pauly K *et al.* (2012) Bio-ORACLE: a global environmental

dataset for marine species distribution modelling. *Global ecology and*

biogeography: a journal of macroecology, **21**, 272–281.

Van Der Kraak G, Pankhurst NW (1997) Temperature effects on the reproductive

performance of fish. In: *Global Warming: Implications for Freshwater and Marine*

Fish, pp. 159–176. Cambridge University Press.

Vanella FA, Boy CC, Fernández DA (2012) Temperature effects on growing, feeding, and

swimming energetics in the Patagonian blennie *Eleginops maclovinus* (Pisces:

Perciformes). *Polar biology*, **35**, 1861–1868.

Vargas-Chacoff L, Moneva F, Oyarzún R *et al.* (2014) Environmental salinity modified

osmoregulatory response in the sub-Antarctic notothenioid fish *Eleginops*

maclovinus. *Polar biology*, **37**, 1235–1245.

Vargas-Chacoff L, Moneva F, Oyarzún R *et al.* (2015a) Metabolic responses to salinity

changes in the subantarctic notothenioid teleost *Eleginops maclovinus*. *Polar*

biology, **39**, 1297–1308.

Vargas-Chacoff L, Saavedra E, Oyarzún R *et al.* (2015b) Effects on the metabolism,

growth, digestive capacity and osmoregulation of juvenile of Sub-Antarctic

Notothenioid fish *Eleginops maclovinus* acclimated at different salinities. *Fish*

physiology and biochemistry, **41**, 1369–1381.

Venables WN, Ripley BD (2013) *Modern Applied Statistics with S-PLUS*. Springer Science

& Business Media.

de Villemereuil P, Frichot É, Bazin É, François O, Gaggiotti OE (2014) Genome scan

methods against more complex models: when and how much should we trust them?

Molecular ecology, **23**, 2006–2019.

de Villemereuil P, Gaggiotti OE (2015) A new F_{ST} □based method to uncover local

adaptation using environmental variables (RB O'Hara, Ed,). *Methods in ecology*

and evolution / British Ecological Society, **6**, 1248–1258.

Wagner HH, Chávez-Pesqueira M, Forester BR (2017) Spatial detection of outlier loci with

Moran eigenvector maps. *Molecular ecology resources*, **17**, 1122–1135.

Wagner HH, Dray S (2015) Generating spatially constrained null models for irregularly

spaced data using Moran spectral randomization methods. *Methods in ecology and*

evolution / British Ecological Society, **6**, 1169–1178.

Wald A (1943) Tests of Statistical Hypotheses Concerning Several Parameters When the

Number of Observations is Large. *Transactions of the American Mathematical*

Society, **54**, 426–482.

Wang L, Wan ZY, Lim HS, Yue GH (2015) Genetic heterogeneity and local adaptation of

Asian seabass across Indonesian Archipelago revealed with gene-associated SNP

markers. *Fisheries research*, **170**, 205–211.

Wang Z, Wu J, Hu Z *et al.* (2020) Dexmedetomidine Alleviates Lipopolysaccharide-

Induced Acute Kidney Injury by Inhibiting p75NTR-Mediated Oxidative Stress and

Apoptosis. *Oxidative medicine and cellular longevity*, **2020**, 5454210.

Waples RS, Gaggiotti O (2006) What is a population? An empirical evaluation of some

genetic methods for identifying the number of gene pools and their degree of

connectivity. *Molecular ecology*.

Waples RS, Punt AE, Cope JM (2008) Integrating genetic data into management of marine

resources: how can we do it better? *Fish and fisheries* .

Weber (1990) Functional significance and structural basis of multiple hemoglobins with

special reference to ectothermic vertebrates. *Animal nutrition and transport*

processes.

breathing amazon catfish *Hoplosternum littorale*. *The Journal of biological chemistry*, **275**, 17297–17305.

Weir BS, Cockerham CC (1984) Estimating F-Statistics for the Analysis of Population Structure. *Evolution; international journal of organic evolution*, **38**, 1358–1370.

Whitlock MC, Lotterhos KE (2015) Reliable Detection of Loci Responsible for Local Adaptation: Inference of a Null Model through Trimming the Distribution of F(ST). *The American naturalist*, **186 Suppl 1**, S24-36.

Wigginton JE, Cutler DJ, Abecasis GR (2005) A note on exact tests of Hardy-Weinberg equilibrium. *American journal of human genetics*, **76**, 887–893.

Willette DA, Allendorf FW, Barber PH *et al.* (2014) So, you want to use next-generation sequencing in marine systems? Insight from the Pan-Pacific Advanced Studies Institute. *Bulletin of marine science*, **90**, 79–122.

Wu RSS (2009) Chapter 3 Effects of Hypoxia on Fish Reproduction and Development. In: *Fish Physiology* (eds Richards JG, Farrell AP, Brauner CJ), pp. 79–141. Academic Press.

Xu Q, Cheng C-HC, Hu P *et al.* (2008) Adaptive evolution of hepcidin genes in antarctic notothenioid fishes. *Molecular biology and evolution*, **25**, 1099–1112.

Xuereb A, D'Aloia CC, Andrello M, Bernatchez L, Fortin M-J (2021) Incorporating putatively neutral and adaptive genomic data into marine conservation planning. *Conservation biology: the journal of the Society for Conservation Biology*, **35**, 909–920.

Yanagimachi, Harumi, Matsubara (2017) Chemical and physical guidance of fish

spermatozoa into the egg through the micropyle. *Biology of blood and marrow*

transplantation: journal of the American Society for Blood and Marrow

Transplantation.

Yilmaz O, Patinote A, Nguyen T *et al.* (2019) Genome editing reveals reproductive and

developmental dependencies on specific types of vitellogenin in zebrafish *(Danio*

rerio). Molecular reproduction and development, **86**, 1168–1188.

Yilmaz O, Patinote A, Nguyen T, Bobe J (2018) Multiple vitellogenins in zebrafish *(Danio*

rerio): quantitative inventory of genes, transcripts and proteins, and relation to egg

quality. *Fish physiology and biochemistry*, **44**, 1509–1525.

Yingjie K, Haihong Y, Lingwei C *et al.* (2019) Apoptosis repressor with caspase

recruitment domain deficiency accelerates ischemia/reperfusion (I/R)-induced acute

kidney injury by suppressing inflammation and apoptosis: The role of AKT/mTOR

signaling. *Biomedicine & pharmacotherapy = Biomedecine & pharmacotherapie*,

112, 108681.

Zhao L, Tan J, Li D *et al.* (2019) SLC39A6/ZIP6 is essential for zinc homeostasis and T-

cell development in zebrafish. *Biochemical and biophysical research*

communications, **511**, 896–902.

Zuur AF, Ieno EN, Elphick CS (2010) A protocol for data exploration to avoid common

statistical problems. *Methods in ecology and evolution / British Ecological Society*,

1, 3–1