Birthdate aligns vestibular sensory neurons with central and motor partners across a sensorimotor reflex circuit for gaze stabilization

Stephanie Huang[1,2]**Emily Gershowitz**,[1]**Marie R. Greaney**,[1]**Samantha N. Davis**,[1]**David Schoppik**,[1]**Dena Goldblatt**[1,2,3,*]

[1]Depts. of Neuroscience, Otolaryngology, and the Institute for Translational Neuroscience, NYU Grossman School of Medicine
[2]Center for Neural Science, New York University
[3]Lead Contact
[*]Correspondence: goldblatt.dena@gmail.com

Developing populations of connected neurons often share spatial and/or temporal features that anticipate their
assembly. A unifying spatiotemporal motif might link sensory, central, and motor populations that comprise an
entire circuit. In the sensorimotor reflex circuit that stabilizes vertebrate gaze, central and motor partners are
paired in time (birthdate) and space (dorso-ventral). To determine if birthdate and/or dorso-ventral organiza-
tion could align the entire circuit, we measured the spatial and temporal development of the sensory circuit node:
the vestibular ganglion neurons. We discovered that progressive dorsal-to-ventral organization closely predicts
vestibular ganglion development, with additional organization along its functional (rostrocaudal) axis. With an
acute optical lesion and calcium imaging paradigm, we found that this common spatiotemporal axis anticipated
functional sensory-to-central partner matching. We propose a "first-come, first-served" model in which birth-
date organizes and assembles the sensory, central, and motor populations that comprise the gaze stabilization
circuit, a general strategy for poly-synaptic circuit assembly across embryonically-diverse neural populations.

SUMMARY STATEMENT

Circuit nodes often develop along common axes. We find that a "first-come, first-served" temporal rule underscores the
organization of a canonical sensorimotor circuit for gaze stabilization, simplifying poly-synaptic assembly.

INTRODUCTION

Individual populations that comprise neural circuits develop in both space and time. During development and matura-
tion, spatial relationships among somata and/or neuropil impose structural constraints on neural circuit assembly and
function (Balaskas et al., 2020; Bikoff et al., 2016; Cang and Feldheim, 2013; Knudsen and Brainard, 1991; Sürmeli
et al., 2011), leaving a topographic record (Erzurumlu et al., 2010; Levine et al., 2012; Penfield and Boldrey, 1937).
Progressive neurogenesis can generate similar maps in time, instructing synaptic partner matching across paired popu-
lations (Fetcho and McLean, 2010; Hamid et al., 2024; Holguera and Desplan, 2018; Kishore et al., 2020; Mark et al.,
2021; Marti et al., 2012; McLean and Fetcho, 2009; Meng et al., 2019; Pujala and Koyama, 2019; Seroka and Doe, 2019;
Wreden et al., 2017). While the alignment of individual partner populations in space and time is well-established, sen-
sorimotor circuits are almost all poly-synaptic. Further, sensorimotor circuits generally span embryologically-diverse
tissue, opening the question of whether the spatial and temporal features that pair partners might similarly organize an
entire circuit.

The vertebrate vestibulo-ocular reflex is a canonical sensorimotor circuit that uses precise sensory-to-motor partner
matching to stabilize gaze. The circuit transforms two vertical directions of head/body destabilization (nose-up/nose-
down) into compensatory eye rotation behavior (eyes-down/eyes-up). Anatomically, the circuit consists of hair cells in
the inner ear, sensory afferents in the anterior (vestibular) statoacoustic ganglion, central vestibular projection neurons,
motor neurons, and extraocular muscles (Szentágothai, 1964). Each direction of motion (up/down) is segregated into its
own specific channel comprised of directionally-selective hair cells, neurons, and muscles. The larval zebrafish, a small
model vertebrate, offers exceptional optical and genetic access to each population (Higashijima et al., 2000; Pittman

et al., 2008; Schoppik et al., 2017). Further, the external development and anatomical simplicity of larval zebrafish (Bagnall and Schoppik, 2018) affords a unique opportunity to elucidate common motifs across an entire sensorimotor circuit.

The vestibular sensory periphery and their central/motor target populations are divergently organized. Functional imaging of the vestibular ganglion established a rostrocaudal axis for nose-down/nose-up segregation, matched to hair cell position along the utricular macula (Tanimoto et al., 2022). Electron micrographs suggest that vestibular ganglion maturation proceeds in the lateral-to-medial direction (Liu et al., 2022), conversely to hair cell development (medial-to-lateral) (Baeza-Loya et al., 2025). In contrast, their central and motor neuron partners are temporally and functionally aligned along the dorsoventral axis: early-born neurons are dorsally positioned and mediate nose-up/eyes-down behavior, while their late-born, ventral counterparts perform the converse (nose-down/eyes-up) (Goldblatt et al., 2023; Greaney et al., 2016). Notably, differentiating sensory progenitors are pushed dorsally as they delaminate from the ventral otic floor (Vemaraju et al., 2012), potentially decoupling functional topography from spatiotemporal development in the vestibular ganglion. Even still, dorsal vestibular ganglion neurons include both nose-up and nose-down subtypes (Liu et al., 2022; Tanimoto et al., 2022), underscoring the point that circuit assembly cannot be solved using a "dorsal-to-dorsal" spatial map alone. Recent work uncovered that time organizes peripheral inputs to the vestibular ganglion (Liu et al., 2022) and for other peripheral sensory afferents, central targeting (Marti et al., 2012). We reasoned that, as suggested in (Liu et al., 2022; Marti et al., 2012), time may also be the "missing" variable that aligns the vestibular periphery with downstream targets, and the complex task of poly-synaptic partner matching achieved simply by assembling neurons with common temporal availability.

To determine if a common temporal motif could organize up/down channels across the entire vestibulo-ocular reflex circuit, we studied how the vestibular ganglion develops in space and time. Anatomical birthdating revealed that ganglion neurons emerged along dorsoventral and rostrocaudal gradients in time. Bipolar afferent projections to peripheral and central targets followed somatic development. With electron microscopy reconstructions, acute optical lesions, and calcium imaging, we discovered that the earliest-born ganglion neurons form functional connections with spatiotemporally-matched central cohorts. Collectively, our findings support that the vestibular ganglion develops along a shared spatiotemporal axis as its downstream central and motor counterparts, where early-born neurons preferentially contribute to a specific channel (nose-up). Our findings complete the characterization of vestibulo-ocular reflex circuit development, revealing a unified spatiotemporal pattern to circuit assembly that spans peripheral sensory, central, and motor areas.

RESULTS

Tg(-6.7Tru.Hcrtr2:GAL4-VP16) provides reliable access to the vestibular (statoacoustic) ganglion neurons that mediate vertical gaze stabilization

In zebrafish, the vestibular sensory neurons that relay utricular pitch-tilt (nose-up/nose-down) sensation are located in the anterior statoacoustic ganglion ("vestibular ganglion") (Sapède and Pujades, 2010; Taberner et al., 2017) (Figure 1A). Past *in vivo* developmental studies used a transgenic line, *Tg(-17.6isl2b:GFP)*, that labels the vestibular ganglion with a fluorescent reporter driven by the *isl2b* promoter (Dyballa et al., 2017; Pittman et al., 2008; Roberts et al., 2017; Sapède et al., 2012; Vemaraju et al., 2012; Zecca et al., 2015). Functional topography was assessed using *Tg(myo6b:jGCaMP7f);Tg(myo6b:tdTomato)* (Tanimoto et al., 2022; Toro et al., 2015). However, neither lines permit tagging subsets of neurons across development.

We instead leveraged the *Tg(-6.7Tru.Hcrtr2:GAL4-VP16)* transgenic (Lacoste et al., 2015; Schoppik et al., 2017) which labels a contiguous population whose location is consistent with the vestibular ganglion (Sapède and Pujades, 2010; Taberner et al., 2017) and that co-localizes with *Tg(-17.6isl2b:GFP)* (Figure 1A). Electron microscopy (EM) reconstructions (Liu et al., 2022) and tracing from the utricular macula (Sapède and Pujades, 2010) support that vestibular ganglion soma are unlikely to be intermingled with other neuronal types. Thus, we interpret that the contiguous *Tg(-6.7Tru.Hcrtr2:GAL4-VP16)*-labeled population is specific to vestibular ganglion neurons.

To evaluate how efficiently and reliably *Tg(-6.7Tru.Hcrtr2:GAL4-VP16)* labels the vestibular ganglion, we used two existing references: the transgenic *Tg(-17.6isl2b:GFP)* (Figure 1B) for *in vivo* comparison and a prior EM atlas of utricular-innervated vestibular ganglion neurons (Jia and Bagnall, 2022; Liu et al., 2022) (Figure 1C). We mapped the

spatial location of *Tg(-6.7Tru.Hcrtr2:GAL4-VP16)*, *Tg(-17.6isl2b:GFP)*, and EM-labeled vestibular ganglion neurons
to a common reference framework (Figure 1D-Figure 1G). Mapping was performed at 5 days post-fertilization (dpf),
when vertical vestibulo-ocular reflex circuit anatomy is stable (Schoppik et al., 2017) and tilt-evoked eye rotation be-
havior is directionally appropriate (Bianco et al., 2012; Leary et al., 2025).

We first evaluated the density of labeled neurons across all three datasets (Figure 1H). We observed a similar number of
Tg(-6.7Tru.Hcrtr2:GAL4-VP16)-labeled neurons (median, 69±15 neurons/hemisphere) as in *Tg(isl2b:GFP)* (median,
73±12 neurons/hemisphere; $p_{rank\ sum}$=0.55). However, both lines labeled fewer neurons than the EM reconstruction
(106 utricular-innervated neurons; N=1 larva; $p_{Kruskal-Wallis}$=0.39). Other estimates indicate that the vestibular ganglion
may contain as many as 100-130 neurons (Liu et al., 2022; Vemaraju et al., 2012). As neurogenesis continues through
adulthood (Schwarzer et al., 2020), the discrepancies in neuronal density could arise from transgenic biases in labeling
post-mitotic vs. all (including immature) ganglion neurons.

We next evaluated the spatial distributions of labeled neurons (Figure 1I). We compared our two *in vivo* datasets:
Tg(-6.7Tru.Hcrtr2:GAL4-VP16) to *Tg(-17.6isl2b:GFP)*. Neurons in both lines spanned approximately 50 μm in the
dorsoventral and rostrocaudal axes (dorsoventral: *hcrtr2*, 53±3 μm; *isl2b*, 51±3 μm; rostrocaudal: *hcrtr2*, 54±3 μm;
isl2b, 53±2 μm) and 60 μm in the mediolateral axis. The distribution of labeled neurons was statistically similar across
lines (dorsoventral: p_{KS}=0.86; rostrocaudal: p_{KS}=0.92; mediolateral: p_{KS}=0.99). Neurons mapped from EM recon-
structions spanned a comparable, though slightly smaller spatial extent: 47 μm dorsoventrally, 40 μm rostrocaudally,
and 40 μm mediolaterally. We interpret the minor size differences (approx. 10 μm) between *in vivo* and EM-mapped
neurons are fixation artifacts in the latter. Next, in both of our *in vivo* transgenic maps, we observed a correlation
between medial and caudal position (Figure 1G; *r*=-0.57). However, the EM reconstruction suggests that vestibular
ganglion neurons are evenly distributed across the rostrocaudal axis, with a medial bias not evident in our transgenic
maps (Figure 1G and Figure 1I). The medial bias suggests the presence of a rostro-medial domain that is not labeled
by either *Tg(-6.7Tru.Hcrtr2:GAL4-VP16)* or *Tg(-17.6isl2b:GFP)*, and which could include the additional ˜30 neurons,
whether post-mitotic or immature, that were not observed in our *in vivo* neuronal counts.

Lastly, given the tight relationships between vestibular ganglion geometry and functional organization (Liu et al., 2022;
Tanimoto et al., 2022), we considered how vestibular ganglion expression in *Tg(-6.7Tru.Hcrtr2:GAL4-VP16)* com-
pared to the *Tg(myo6b:jGCaMP7f)*;*Tg(myo6b:tdTomato)* line that was previously described (Tanimoto et al., 2022).
Tg(-6.7Tru.Hcrtr2:GAL4-VP16) and *Tg(isl2b:GFP)* appear to span a comparable spatial extent around the utricle as
Tg(myo6b:jGCaMP7f);*Tg(myo6b:tdTomato)*. However, expression may be more inconsistent along the spatial edges
(10–20 μm) and with approximately half the labeling density as *Tg(myo6b:jGcamp7f)*;*Tg(myo6b:tdTomato)* (personal
communication, M. Tanimoto). The comparable spatial extents suggests that *Tg(-6.7Tru.Hcrtr2:GAL4-VP16)* approxi-
mates, but might not comprehensively portray spatially-linked developmental patterns.

Together, our reference mapping across three datasets highlight the importance and challenges associated with aligning
variable anatomical geometry across transgenic lines and studies. We conclude that neither *Tg(-6.7Tru.Hcrtr2:GAL4-
VP16)* nor *Tg(isl2b:GFP)* comprehensively access all utricular-innervated vestibular ganglion neurons; we speculate
that the rostro-medial domain, which may contain the latest-born neurons, may be unincorporated in these transgenics.
Nevertheless, *Tg(-6.7Tru.Hcrtr2:GAL4-VP16)* provides comparable access to the vestibular ganglion as *Tg(isl2b:GFP)*,
the most comprehensively-characterized *in vivo* transgenic used to date, and to the majority of the vestibular ganglion
ganglion. The implications of potentially inconsistent labeling of the vestibular ganglion are noted where appropriate
below.

Developmental time organizes a dorsoventral map in the vestibular ganglion

To tag vestibular ganglion neurons during development, we used *Tg(-6.7Tru.Hcrtr2:GAL4-VP16)* to express a photo-
convertible reporter, Kaede (Scott et al., 2007), for permanent optical labeling of experimenter-defined temporal cohorts
(Caron et al., 2008; Goldblatt et al., 2023; Greaney et al., 2016). We tagged neuronal cohorts born at intermediate stages
between 22–72 hours post-fertilization (hpf) (Figure 2A), the timecourse over which *isl2b*+ vestibular ganglion neurons
differentiate in larval zebrafish (Haddon and Lewis, 1996; Vemaraju et al., 2012). We imaged vestibular ganglion neu-
rons at 5 dpf. Tagged neurons (i.e., born before the time of Kaede photoconversion) were readily identifiable by the
presence of red, converted Kaede; neurons born after conversion contained only green, unconverted protein (Figure 2B).
The number of tagged neurons progressively increased from 22–72 hpf, with a transient surge around 33 hpf (N=5 con-

verted larvae/timepoint; Figure 2C, Table 1). This transient phase could reflect the differentiation of neurons that are not incorporated in our genetic line, such as the rostro-medial sector (Figure 1G). Most neurons (91% of the number observed in our non-birthdated reference dataset, n=345 neurons/N=5 fish shown in Figure 1D and Figure 1H) evaluated at 5 dpf were born by 72 hpf, in line with prior reports (Vemaraju et al., 2012).

To evaluate birthdate-linked spatial organization, we compared the location of the earliest- and latest-born neurons. By the chronological midpoint (48 hpf) of our observed differentiation window (22–72 hpf), the majority of labeled ganglion neurons are already born (approx. 75%) (Table 1). Instead, the midpoint where 50% of neurons are born is 33–36 hpf (Table 1). We thus use 36 hpf as a reference midpoint to describe early- and late-born neurons. We evaluated an early-born (before 26 hpf) and late-born cohort (after 48 hpf), collectively comprising half (41%) of all vestibular ganglion neurons (Table 1). Strikingly, these cohorts were differentially organized along the dorsoventral axis (Figure 2D). We quantified the probability of observing early- and late-born soma in each spatial axis (Figure 2E). Early and late cohort position was significantly different along the dorsoventral ($p_{KS}=1.1\times10^{-16}$) and rostrocaudal axes ($p_{KS}=1.7\times10^{-4}$). Qualitatively, the earliest-born cells may be medially positioned (Figure 1B). However, statistically significant mediolateral organization was not observed ($p_{KS}=0.25$). As some medial neurons may not be labeled in *Tg(-6.7Tru.Hcrtr2:GAL4-VP16)* (Figure 1G) and the mediolateral axis was orthogonal to our imaging and mapping plane, we cannot rule out the possibility of additional mediolateral organization.

Importantly, the early-born cohort appeared caudal of center relative to our non-birthdated reference dataset (Figure 1D, Figure 1I) ($p_{KS}=1.4\times10^{-11}$) in aggregate maps (N=10 larvae; N=5 tagged at 22 hpf, and N=5 tagged at 26 hpf; Figure 2D-Figure 2E). As the vestibular ganglion is rostrocaudally organized (rostral, nose-down; caudal, nose-up) (Liu et al., 2022; Tanimoto et al., 2022), we more closely examined this caudal bias. Individual maps from N=5 larvae (Figure 2F) suggest that most neurons born before 26 hpf lie on or very near to the rostro-caudal midpoint of our (non-birthdated) reference dataset. As a secondary assessment, we re-evaluated the EM reconstruction, which had inferred neuronal age using myelination at 5 dpf (Figure 2G). We mapped myelinated (magenta; presumably earliest-born) and un-myelinated (green; late-born) neurons to our reference framework (Figure 2G). As in our data, myelinated neurons were distributed largely around the rostro-caudal midpoint, with a statistically significant caudal bias (myelinated vs. unmyelinated; Figure 2H-Figure 2I; rostrocaudal: $p_{KS}=0.01$, dorsoventral: $p_{KS}=0.93$). Though myelinated neurons were equally observed across the dorsal and ventral domains, this is not incongruent with our *in vivo* birthdating findings; myelination status at 5 dpf likely includes neurons born well-after 26 hpf. Together, *in vivo* and EM maps suggest that the earliest-born ganglion neurons may in fact be caudal biased, supporting a nose-up designation.

The spatial patterns we observed (dorsal, caudal) could reflect discrete spatiotemporal windows of neurogenesis or arise from gradients generated continuously across time. Evidence that dorsal and/or caudal neurons continue to accumulate at later developmental stages would support the latter. To differentiate these hypotheses, we evaluated the location of neurons from our *in vivo* dataset born before (Figure 3A) and after (Figure 3B) all seven of our converted timepoints (N=5 larvae/cohort; full statistics in Table 2). The dorsal bias we observed persisted until approximately 36 hpf (36 hpf vs. 33 hpf: $p_{ANOVA\ multiple\ comparisons\ test}=0.20$; Figure 3C). Qualitatively, by 48 hpf, nearly all "born after" neurons were located ventral to the midpoint of the ganglion (Figure 3B). We interpret that dorsal and ventral neurons are largely born in opposing temporal windows, offset by approximately 8–12 hours: dorsal neurons from 22–48 hpf, and ventral neurons from 30–72 hpf.

As the earliest born 30% of the vestibular ganglion may lie caudal of center (Figure 2E), we next asked if *progressive* rostrocaudal development also anticipates functional topography. We evaluated the rostro-caudal distribution of neurons at each stage relative to the spatial midpoint. The exclusive caudal bias to early-born neurons was discontinued by 30 hpf (Figure 3A), when rostral neurons first became apparent. After 33 hpf (remaining 70% of neurons), rostral and caudal neurons emerged contemporaneously (33 hpf vs. 30 hpf: $p_{ANOVA\ multiple\ comparisons\ test}=0.94$, Figure 3D, Table 2). Both caudal and rostral neurons remained unconverted after 60 hpf (Figure 3B). While our dataset contains a low number of unconverted neurons at 72 hpf, the presence of rostral neurons at this age suggests that — unlike their dorsal counterparts — ventro-rostral and ventro-caudal accumulate continuously from 33 hpf through late development.

We conclude that ganglion development progresses along a major dorsoventral and secondary rostrocaudal axis. The dorso-caudal location of the earliest-born cells may anticipate their functional (nose-up) topography. While our genetic line may not comprehensively access the rostro-medial sector, that both ventro-rostral and ventro-caudal neurons emerge contemporaneously through late development (72 hpf) suggests that spatial and functional organization are uncoupled

after 33 hpf and for the vast majority (70%) of labeled neurons. Contrastingly, the clearly delineated dorsoventral patterns we observe elevates the dorsoventral axis as a temporally-enduring predictor of vestibular ganglion accumulation.

Bipolar afferent organization follows spatial patterns of somatic development

Somatic topography anticipates axonal connectivity (Balaskas et al., 2020; Dasen, 2009; Goldblatt et al., 2023; Sürmeli et al., 2011). Vestibular ganglion neurons send bipolar afferents ventrally to peripheral hair cells in the utricular macula and dorsally to central hindbrain targets via the VIII[th] nerve (Liu et al., 2022; Sapède and Pujades, 2010) (Figures 4A and 4B). We hypothesized that the orderly positioning of early-born vestibular ganglion neurons could spatially constrain the path of their projections within each bipolar afferent bundle. We therefore analyzed the bipolar afferents of the converted neurons described previously (Figure 4C) along its width (rostrocaudal axis). Analysis in the dorsoventral axis was not possible, as projections inherently grow towards either dorsal or ventral targets. We evaluated the spatial distribution of converted axons (i.e., from neurons born before the time of conversion) at 5 dpf, reasoning that organization at 5 dpf reflects how axons become assembled into the VIII[th] nerve.

We first evaluated peripheral projections to the utricular macula (Figures 4D and 4E; N=3 hemispheres/timepoint). Converted axons from soma born before 22 hpf were only observed in the caudal macula ("born before" vs. all fluorescence, rostrocaudal axis: $p_{KS}=5.3 \times 10^{-11}$). Conversion spread rostrally by 30 hpf and spanned the full rostrocaudal extent of the macula by 36 hpf ($p_{KS}=0.50$), matching the temporal emergence of rostrally-positioned somata. Next, we evaluated projections to the VIII[th] nerve (Figures 4F and 4G; N=3 hemispheres/timepoint). In the caudal half of the axon bundle, we observed a faint but visible converted axon from soma born before 22 hpf. We observed stronger conversion in the caudal and center of the axon bundle from somata born before 26 hpf ($p_{KS}=0.016$). As with the macula, rostral spread across the VIII[th] nerve bundle was evident by 30 hpf ($p_{KS}=0.69$). Together, we conclude that rostrocaudal organization within the utricular macula and VIII[th] nerve follows neuronal birthdate: early-born soma (caudal) project caudally or center-caudally within each axon bundle, respectively.

Early-born vestibular ganglion neurons assemble with spatiotemporally- and functionally-matched central neuron cohorts

Does birthdate anticipate synaptic assembly between partner populations across the developing vestibulo-ocular reflex circuit? Vestibular ganglion neurons relay tilt sensation to central (hindbrain) projection neurons in the tangential nucleus (Goldblatt et al., 2023; Liu et al., 2022), which are necessary and sufficient for vertical (nose-up/nose-down) gaze stabilization behavior (Bianco et al., 2012; Schoppik et al., 2017). Previously, we found that both birthdate and function segregate central vestibular neurons: early-born dorsal neurons are sensitive to nose-up tilts, and late-born ventral neurons are sensitive to nose-down tilts (Goldblatt et al., 2023). This pattern mirrors the dorsoventral development we observed for their vestibular ganglion inputs and occurs over a comparable temporal window (22–48 hpf) (Goldblatt et al., 2023). Thus, we predicted that the earliest-born vestibular ganglion neurons (dorsal, caudal, nose-up) would assemble with temporally- (early-born), spatially- (dorsal), and functionally- (nose-up) matched central targets.

We first evaluated whether vestibular ganglion afferents from early-born neurons are anatomically poised to wire to temporally-matched central targets. At 5 dpf, VIII[th] nerve input from neurons born before 22 hpf exclusively targeted the early-born (dorsal) tangential nucleus (Figure 5A). Thus, early-born (dorsal) vestibular ganglion neurons may innervate commonly positioned (early-born, dorsal) central neuron partners. However, *Tg(-6.7Tru.Hcrtr2:GAL4-VP16)* labels other cranial sensory ganglia that develop comparably early and project along the VIII[th] nerve (Zecca et al., 2015).

To isolate afferents from the vestibular ganglion, we re-evaluated the EM dataset from (Jia and Bagnall, 2022; Liu et al., 2022) (Figure 1C), which had also reconstructed projections to central vestibular targets in the hindbrain. We again used myelination at 5 dpf to infer birth order. At 5 dpf, myelinated neurons preferentially localize to the lateral ganglion (Liu et al., 2022). Correspondingly, their afferents form a lateral bundle at the proximal edge of the VIII[th] nerve (Figure 5B). However, at the hindbrain entry site, the afferent bundle inverted such that myelinated afferents innervated the medial (dorsal) tangential nucleus (Figure 5B-Figure 5C). We next restricted our analysis to vestibular ganglion afferents that were synaptically coupled with the tangential nucleus, reasoning that some afferents may pass through the tangential nucleus en route to other synaptic targets. Still, tangential nucleus neurons that were synaptically coupled with myelinated ganglion afferents were preferentially dorsal and medial, aligning with its functional topography (Figure 5D) (Goldblatt et al., 2023). Together, these anatomical observations support that early-arriving sensory input from the vestibular ganglion may preferentially target the earliest-born, nose-up tangential nucleus neurons.

To test our anatomical predictions, we evaluated whether early-born neurons in the vestibular ganglion and tangential nucleus are functionally connected. We leveraged an existing acute optical lesion (Goldblatt et al., 2023; Hamling et al., 2024; Schoppik et al., 2017) and calcium imaging (Goldblatt et al., 2023; Hamling et al., 2022) paradigm (Figure 5E-Figure 5F). First, we birthdated larvae at 26 hpf to tag the earliest-born 15% of vestibular ganglion soma and visualized neurons at 5 dpf. Lesioning early-born soma was not technically feasible due to their close proximity to the dorsal VIIIth nerve bundle and local vasculature. Instead, we unilaterally lesioned somata of the 85% of ganglion neurons born after 26 hpf (Figure 5E), leaving only the early-born (dorso-caudal, nose-up) neurons available to relay tilt sensation. Somatic lesions secondarily degrade the axon projections of targeted neurons (Goldblatt et al., 2023; Hamling et al., 2024; Schoppik et al., 2017). Thus, by targeting our lesions to neuronal somata, our approach is specific to the late-born neurons and their projections. To evaluate changes in central responses to tilt sensation after ablating late-born ganglion neurons, we used Tilt-In-Place Microscopy (TIPM) (Hamling et al., 2022). Briefly, TIPM reports changes in a genetically-encoded calcium indicator, GCaMP6s (Chen et al., 2013), in central neurons after delivering nose-up and nose-down pitch tilts (Figure 5F). At 5 dpf, the vestibular ganglion is the primary source of pitch tilt information (Goldblatt et al., 2023). Therefore, loss of central neuron responses to tilt sensation should directly reflect loss of corresponding ganglion input.

We predicted that loss of late-born (ventral) vestibular ganglion inputs would eliminate pitch tilt responses in spatiotemporally-matched (late-born, ventral, nose-down) central neurons. We first imaged the tilt responses of n=172 tangential nucleus neurons from N=4 control hemispheres and n=115 neurons from N=4 hemispheres with acute, uni-lateral ganglion lesion. We observed a substantial reduction in the number of tilt-responsive neurons (median control: 89.4±6.0% responsive; lesion: 36.9±21.4% responsive; $p_{rank sum}$=0.03), driven by a near total loss to nose-down tilts (nose-down: median control, n=8±5 neurons/fish; lesion: n=1±2 neurons/fish; $p_{rank sum}$=0.03; nose-up: control, n=17±2 neurons/fish; lesion: n=16±4 neurons/fish; $p_{rank sum}$=0.09; Figure 5G). Central neurons that remained able to respond to tilt sensation were dorsally-shifted (median position: control, 25±8μm; lesion, 30±7μm, p_{KS}=0.01; Figures 5H and 5I, consistent with the loss of ventral (nose-down) neurons (Goldblatt et al., 2023).

The loss of tilt sensitivity in ventral, nose-down central neurons should consequently impair upwards eye rotation behavior. To test this prediction, we leveraged an existing assay of vestibulo-ocular reflex behavior, which measures torsional eye rotation responses (eyes-up/eyes-down) following body tilts (nose-down/nose-up) (Figure 5J) (Leary et al., 2025; Schoppik et al., 2017). We compared torsional eye rotation behavior after performing acute, bilateral lesions of either late-born ganglion neurons (N=6 fish) as before or the anterior lateral line (N=7 fish), which does not mediate vestibulo-ocular reflex behavior (control). As expected, late-born ganglion lesion significantly impaired eye rotation responses relative to control (Figure 5K-Figure 5L). Eye rotation responses were nearly entirely abolished to nose-down tilts ($p_{rank sum}$=0.002; peak angular velocity: 17.6% of mean control). We observed an additional loss of nose-up tilt responses ($p_{rank sum}$=0.001; peak angular velocity: 30.6% of mean control). These deficits are in line with the loss of functional sensitivity in the tangential nucleus.

Together, we interpret the absence of tilt responses in the ventral tangential nucleus and ensuing eye rotation deficits to be a direct consequence of loss of late-born (ventral) vestibular ganglion inputs. Correspondingly, the preservation of dorsal (nose-up) central neuron responses suggests a tight matchup in time between early-born (dorso-caudal, nose-up) ganglion neurons and their early-born (dorsal, nose-up) targets. Collectively, these experiments demonstrate early-born ganglion neurons assemble with spatiotemporally-matched central neuron targets, and that birth order is a continuous axis that links functional subtypes of vestibulo-ocular reflex neurons.

DISCUSSION

Here, to determine whether temporal organization could assemble a poly-synaptic sensorimotor circuit, we studied how time comes to assemble the vestibular ganglion and its connectivity. We discovered that development proceeds along a dorsal-to-ventral and secondarily, a caudal-to-rostral axis for a subset of early-born neurons, reminiscent of the ganglion's functional organization. Afferents from the vestibular ganglion systematically innervate their utricular and central (vestibular hindbrain) targets in a similar spatial and temporal progression. Intriguingly, these organizational motifs are shared across the entire vestibulo-ocular reflex circuit (Figure 6). Indeed, our loss-of-function experiments establish that early-born vestibular ganglion neurons preferentially assemble with early-born central partners, just as early-born central neurons assemble with early-born motor neuron targets (Goldblatt et al., 2023). Taken together, our

279 work reveals that the vertical vestibulo-ocular reflex circuit assembles sequentially in time, aligning sensory, central,
280 and motor output neurons into separate up and down channels. Our findings speak to the earliest steps in construction
281 of an ancient, conserved, and canonical sensorimotor reflex circuit.

**Limitations in genetic reagents and their implications for understanding the organization and specification of
vestibular ganglion subtypes**

Our characterization of vestibular ganglion anatomy using two transgenic lines and an existing electron microscopy
dataset reveal the challenges associated with aligning organizational patterns across studies, particularly given the small
size (50-60 μm) of the vestibular ganglion. Both transgenic reagents used in our study, *Tg(-6.7Tru.Hcrtr2:GAL4-VP16)*
and *Tg(isl2b:GFP)* likely under-sample the vestibular ganglion in neuronal number (55-70% of EM-reconstructed neu-
rons) and spatial distribution (rostro-medial access). Across transgenics, additional bias could be introduced by the
varying temporal integration of GAL4-UAS reagents compared to direct promoter-reporter fusion lines. Indeed, a
different fusion line, *Tg(myo6b:jGCaMP7f);Tg(myo6b:tdTomato)* (Tanimoto et al., 2022; Toro et al., 2015), appears
to access a similarly extended population of ganglion neurons. Further, vestibular ganglion progenitors continuously
proliferate through adulthood (Schwarzer et al., 2020), generating differentiated neurons well-past our study endpoint
(5 dpf). Intriguingly, the vestibular ganglion becomes myelinated in the lateral-to-medial direction (Liu et al., 2022),
suggesting that the youngest (unmyelinated) neurons could occupy the rostro-medial sector that was not labeled by
either *Tg(-6.7Tru.Hcrtr2:GAL4-VP16)* or *Tg(isl2b:GFP)*. Alternatively, more recent work discovered their hair cell in-
puts develop in the converse direction (medial-to-lateral) (Baeza-Loya et al., 2025), in line with our observed medial
localization to early-born ganglion neurons. Future studies could resolve these divergent observations by tracking the
migration patterns of the same vestibular ganglion neurons across development and life (Schwarzer et al., 2020), as
recent work tracked progenitor migration linked to coarse (anteroposterior) ganglion domain formation (Dyballa et al.,
2017).

The caveats in our genetic access to the vestibular ganglion merits caution in interpreting that the progressive dorsoven-
tral, and secondary rostrocaudal development we observed are the sole organizational axes. Nevertheless, the *temporal*
structure we observed may still closely reflect the developmental pressures that organize the vestibular ganglion. For
example, the dorsal-to-ventral accumulation of ganglion neurons may be closely influenced by the spatial pressures
imposed by sequential progenitor delamination: differentiated neurons are pushed dorsally by their transit-amplifying
precursors in the ventral otic floor (Vemaraju et al., 2012). Similarly, though the earliest-born 30% of ganglion neurons
likely receive nose-up utricular input based on their caudal position (Liu et al., 2022; Tanimoto et al., 2022), this early
caudal bias is abated relatively early in time (33 hpf), with the latest-born ganglion neurons distributed evenly across
the ventro-rostral and ventro-caudal ganglion. We propose that for most ganglion neurons, birthdate and functional
topography are uncoupled, and that spatial position is simply the passive byproduct of time.

Together, our considerations motivate future recordings in birthdated neurons to directly correlate the spatiotemporal
patterns we observe with the anatomical and functional geometry of the vestibular ganglion. Such experiments may be
achievable with state-of-the-art advances that overcome the incongruent axes needed to visualize the vestibular ganglion
(horizontal) and stimulate the utricular macula (vertical, gravitational) (Favre-Bulle et al., 2018; Migault et al., 2018;
Sugioka et al., 2023; Tanimoto et al., 2022), but which are outside the capacities of the present study. Prior work parsed
the statoacoustic ganglion into coarse anteroposterior (rostrocaudal) domains by sensory input modality (auditory vs.
vestibular) (Sapède and Pujades, 2010; Vemaraju et al., 2012), providing a broad framework to discover corresponding
anteroposterior patterns of progenitor delamination (Bañón and Alsina, 2023; Bell et al., 2008; Dyballa et al., 2017;
Koundakjian et al., 2007; Saini and Ladher, 2024; Vemaraju et al., 2012), molecular differentiators (Haddon et al.,
1998; Hammond and Whitfield, 2011; Laureano et al., 2022; Maier and Whitfield, 2014; Sapède and Pujades, 2010;
Whitfield et al., 2002), and VIII[th] nerve organization by sensory ganglia origin (Kimmel et al., 1990; Marti et al., 2012;
Pujol-Martí et al., 2010; Zecca et al., 2015). Here, the temporal organization we uncovered may provide molecular
access to even finer subtypes of ganglion neurons, such as by distinct input channels or functional dynamics (Liu and
Bagnall, 2023; Liu et al., 2022) — of broad importance given the general susceptibility of the vestibular periphery in
disease (Delmaghani and El-Amraoui, 2022).

Fate and connectivity in the vestibulo-ocular reflex circuit: Destined by time?

How might structural connectivity in the vestibulo-ocular reflex circuit emerge in time? Our model predicts that time reduces the complex task of matching synaptic partners across the peripheral and central nervous system to the passive assembly of first-available neurons, aligned across a common positional template. Key support comes from our finding that, as in other species (Altman and Bayer, 1981; Auclair et al., 1999; Glover, 2003; Maklad and Fritzsch, 2003; Puelles and Privat, 1977; Shaw and Alley, 1981; Vemaraju et al., 2012), across the zebrafish vestibulo-ocular reflex circuit, dorsal-to-ventral neurogenesis (Goldblatt et al., 2023; Greaney et al., 2016) proceeds over a shared temporal window (24–48 hpf) (Goldblatt et al., 2023; Greaney et al., 2016; Vemaraju et al., 2012). In spinal reflex circuits (Balaskas et al., 2020; Kishore et al., 2020; Sürmeli et al., 2011), such parallel neurogenesis imposes temporal and/or spatial constraints on axon targeting. Unlike in these circuits, sensory and central partner populations in the vestibulo-ocular reflex circuit derive from different embryological precursors with distinct environmental pressures (Haddon and Lewis, 1996), posing additional complexity to partner matching. Previously, we established that early-born, dorsally-located central vestibular neurons in the hindbrain preferentially wire to early-born, dorsal motor neurons (Goldblatt et al., 2023). Here, we demonstrate that the same spatiotemporal rules assemble vestibular ganglion neurons with their central targets. Our findings support past proposals that neuronal position – a consequence of birthdate – pre-configures connectivity in the vestibulo-ocular reflex circuit (Glover, 1996), as already established for other peripheral sensory ganglia (Pujol-Martí et al., 2010) and their central targeting (Marti et al., 2012). Vestibular ganglion axons form a 90-degree turn as they enter the hindbrain and project towards central targets (Jia and Bagnall, 2022; Liu et al., 2022); this inversion could set up the dorsoventral functional axis downstream (Goldblatt et al., 2023). A "first-come, first-served" mechanism to match temporally-available partners offers a simple means of poly-synaptic sensorimotor circuit construction.

Past work suggests that the true "first arriving" partners in the vestibulo-ocular reflex circuit are their peripheral sensory inputs. In zebrafish, hair cells in the utricular periphery become anatomically mature (Whitfield et al., 2002) and mechanically active (Tanimoto et al., 2009; 2011) as neurogenesis is only just beginning in central/motor neurons (Goldblatt et al., 2023; Greaney et al., 2016). Similar timelines have, again, been reported in zebrafish (Haddon and Lewis, 1996; Riley et al., 1997; Vemaraju et al., 2012; Zecca et al., 2015), avians (Fritzsch and Nichols, 1993), and mammals (Altman and Bayer, 1982). Together, these findings support emerging evidence (Clark et al., 2013; Goldblatt et al., 2023; 2024; Greaney et al., 2016) that the initial synaptic architecture of the vestibulo-ocular reflex circuit develops in the anterograde (sensory-to-motor) direction. Peripheral targets can influence the development of proprioceptive spinal pathways in a similar manner (Arber, 2012; Norovich et al., 2022; Poliak et al., 2016; Wenner and Frank, 1995; Wu et al., 2019). Curiously, though central axons project to motor targets soon after neurogenesis, synaptogenesis between central and motor neurons (Goldblatt et al., 2023) and between motor neurons and muscles (Clark et al., 2013) is delayed until after differentiation across all vestibulo-ocular reflex populations, and hair cell transcriptional maturation (Baeza-Loya et al., 2025) is nearly complete. We propose that the early formation of sensory-central connections instructs downstream wiring specificity. Evidence that vestibular ganglion axons project soon after differentiation, and that sensory-central synaptogenesis proceeds during the downstream synaptogenesis "delay" would strongly support an anterograde assembly model of vestibulo-ocular circuit development.

While our model predicts that temporal availability allows early-to-early born cohorts to assemble, additional forces may still refine connectivity for late-born cohorts and as circuit function matures (Leary et al., 2025). Though early input to the vestibulo-ocular reflex is exclusively nose-up (dorso-caudal), input after 30 hpf contains both up (caudal) and down (rostral) channels (Liu et al., 2022; Tanimoto et al., 2022) that may project to similar central targets, particularly to ventral neurons. This mismatch is curious, given that central projection neurons do not show strict up/down selectivity until later (Leary et al., 2025). We hypothesize that ventral central projection neurons may initially receive input from both directionally-tuned channels during development, and inappropriate sensory inputs are pruned at their central targets to enable directional selectivity in maturity. Thus, our "first-come, first-served" model is consistent with the congruent rostro-caudal (up/down) development of the ventral vestibular ganglion. Notably, sensation is dispensable for vestibulo-ocular reflex circuit development (Leary et al., 2025; Roberts et al., 2017; Ulrich et al., 2014), arguing against stimulus-driven remodeling (Hubel and Wiesel, 1964; Schmidt and Eisele, 1985) and in favor of molecular refinement (Sanes and Zipursky, 2020); Teneurin signaling has recently been shown to correct axon mis-sorting in such a manner (Spead et al., 2023). Looking ahead, measuring when and how directional selectivity emerges in central

376   neurons will be key to resolving the mechanisms that instantiate synaptic specificity.

Temporal maps: a blueprint for neural circuit construction

Our findings link time with functional circuit assembly by discovering that time can shape spatial maps over relatively
homogeneous functional scales, and across an entire poly-synaptic sensorimotor reflexive circuit. Across vertebrate and
invertebrate models, the timing of neurogenesis regulates partner availability by constraining anatomical connectivity
(Bayraktar and Doe, 2013; Berry et al., 1964; Glenwinkel et al., 2021; Goldblatt et al., 2023; Koundakjian et al.,
2007; Li et al., 2013; Meng et al., 2020; Sullivan et al., 2019), function in local ensembles (Fore et al., 2020; Huszár
et al., 2022; Younes et al., 2024), and assembly into behaviorally-separable circuits (Fetcho and McLean, 2010; Hamid
et al., 2024; Kishore et al., 2020; Liu et al., 2022; Mark et al., 2021; Marti et al., 2012; McLean and Fetcho, 2009;
McLean et al., 2007; Pujala and Koyama, 2019; Wreden et al., 2017). Here we observed that time can also organize
closely-related, binary subtypes of neurons (nose-up/nose-down) that subserve just one sensory modality (balance) and
motor behavior (eye movements). Despite the clear structural constraint that time imposes on developing sensorimotor
circuits, the extent to which individual systems may rely on passive "first-come, first-served" partner matching remains
controversial (Barsh et al., 2017; Isabella et al., 2021). Indeed, the next frontier in progressing beyond well-established
but still indirect relationships between birth order, progressive spatial development, and functional circuit assembly
continues to be limited by challenges in manipulating time itself. In other systems, transplantation experiments (Barsh
et al., 2017; Isabella et al., 2021) or genetic manipulations of temporally-restricted populations (Hamid et al., 2024;
Roome et al., 2025; Sagner et al., 2021) have laid a path towards directly linking time, neuronal fate, and topography.
For the vestibular periphery, expanded genetic atlases will be necessary for access to spatially- or temporally-restricted
subtypes within an individual sensory ganglion and/or their progenitors, though the quick regenerative capacity of the
zebrafish poses continuing constraints. Such insight may ultimately reconcile whether one rule, such as "first-come
first-served", or disparate rules may govern sensorimotor circuit assembly.

The continuity of the temporal map we discovered, from the sensory periphery to motor output, offers a path to re-
solve the extent to which the vestibulo-ocular reflex circuit is constructed holistically through shared codes or instead
constructed as local modules (Cang et al., 2008), stitched together by local signals. One hypothesis is that shared orga-
nization, whether in space or time, reflects the existence of a shared molecular code that coordinates circuit assembly.
Common, spatially-restricted patterns of chemorepulsive molecules underscore partner matching across retinal circuits
(Cheng et al., 1995; Flanagan and Vanderhaeghen, 1998; Sperry, 1963); can be spatially inverted across target popula-
tions (Triplett and Feldheim, 2012); and can even vary along spatial gradients, matching the graded functional tuning
of synaptic partners (Dombrovski et al., 2024). However, the divergent embryological origins of each vestibulo-ocular
reflex circuit populations suggests the nature of such a shared code is unlikely to be spatial, downstream of common
extrinsic induction factors (Dasen, 2009). Instead, our temporal map predicts that "molecular matching" gradients
might alternatively co-emerge in time. For example, time could establish "competency" windows in which neurons are
maximally poised to respond to local wiring specificity signals (Jefferis et al., 2001; Koundakjian et al., 2007; Petrovic
and Hummel, 2008; Rees et al., 2017; Sperry, 1963; Yogev and Shen, 2014; Zhang et al., 2024), perhaps downstream
of temporally-restricted transcription factor expression (Hamid et al., 2024; Isabella et al., 2021; Joo et al., 2013; Kur-
mangaliyev et al., 2019; Meng et al., 2020; Sullivan et al., 2019). Emerging evidence supports that such temporal codes
may be shared across related partner populations (Shin et al., 2020) and coarsely across brain regions (Sagner et al.,
2021), though whether these could act over relatively homogeneous scales remains unresolved. Future studies could
leverage the access afforded by *Tg(-6.7Tru.Hcrtr2:GAL4-VP16)* to dorsal vestibular ganglion neurons, which assemble
preferentially with spatially, temporally, and functionally-matched central targets, to test such a hypothesis. Early work
predicted that "[T]he basic connectivity scheme of the [vestibulo-ocular reflex circuit] is set up by determinate mech-
anisms of cell-cell recognition" (Glover, 2003), likely downstream of early fate specification programs (Glover, 1996;
Lunde et al., 2019). Looking ahead, it will be illuminating to resolve whether this is achieved through distinct ligand-
receptor pairs expressed in particular temporal cohorts or through just one factor, dynamically regulated or differentially
available in time.

Conclusion

423   Here, we complete our characterization of shared spatial, temporal, and functional features across a developing sen-
424   sorimotor reflex circuit. The common temporal positioning of vestibular partner populations is reminiscent of the

shared topographic maps that align sensory space/body plans with central representations in canonical (i.e., retinal, auditory, somatotopic) sensory systems. For the vestibulo-ocular reflex circuit, temporal alignment may serve as a passive constraint on functional partner matching, simplifying assembly across diverse tissues. Though additional forces may subsequently refine connectivity as the vestibulo-ocular reflex circuit matures, our findings provide a key advance towards defining the initial structural blueprints that constrain the development of poly-synaptic sensorimotor circuits.

MATERIALS AND METHODS

KEY RESOURCES TABLE

Reagent or resource	Source	Identifier
Chemicals, peptides, and recombinant proteins		
Low melting point agarose	ThermoFisher Scientific	16520
Ethyl-3-aminobenzoic acid ethyl ester (MESAB)	Sigma Aldrich	E10521
Pancuronium bromide	Sigma Aldrich	P1918
Deposited data		
Raw and analyzed data	This study	
Publicly available data		
EM connectome data	(Liu et al., 2022)	
Experimental models: Organisms/strains		
Tg(-6.7Tru.Hcrtr2:GAL4-VP16)	(Lacoste et al., 2015; Schoppik et al., 2017)	ZDB-ALT-151028-3
Tg(UAS-E1b:Kaede)	(Scott et al., 2007)	ZDB-ALT-070314-1
Tg(-17.6isl2b:GFP)	(Pittman et al., 2008)	ZDB-ALT-100322-2
Tg(UAS:GCaMP6s)	(Chen et al., 2013)	ZDB-ALT-140811-3
Software and algorithms		
Fiji/ImageJ	(Schindelin et al., 2012)	RRID: SCR_02285
Adobe Illustrator (2021)	Adobe	RRID: SCR_010279
Matlab 2020b	Mathworks	RRID: SCR_001622

RESOURCE AVAILABILITY

Lead Contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Dena Goldblatt (goldblatt.dena@gmail.com) in coordination with David Schoppik (schoppik@gmail.com).

Materials Availability

This study did not generate new unique reagents.

Data and code availability

All data and code are deposited at the Open Science Framework and are publicly available at 10.17605/OSF.IO/KP839

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Fish care

All protocols and procedures involving zebrafish were approved by the New York University Grossman School of Medicine Institutional Animal Care & Use Committee (IACUC). All larvae were raised at 28.5° C at a density of 20-50 larvae in 25-40 ml of buffered E3 (1mM HEPES added). Larvae used for birthdating experiments were raised in constant darkness; all other fish were raised on a standard 14/10h light/dark cycle. Larvae for experiments were between 1-5 days post-fertilization (dpf).

Transgenic lines

Experiments were conducted on the *mifta^-/-* background to remove pigment. Validation of statoacoustic ganglion labeling by *Tg(-6.7Tru.Hcrtr2:GAL4-VP16)* (Lacoste et al., 2015; Schoppik et al., 2017) was performed using *Tg(-17.6isl2b:GFP)* (Pittman et al., 2008). All other experiments used *Tg(-6.7Tru.Hcrtr2:GAL4-VP16)* (Lacoste et al., 2015; Schoppik et al., 2017) larvae to drive expression of the following UAS reporters: *Tg(UAS:E1b-Kaede)* (Scott et al., 2007) for birthdating only experiments and *Tg(UAS:GCaMP6s)* (Thiele et al., 2014);*Tg(US:E1b-Kaede)* for acute lesion/calcium imaging experiments. Larvae used were selected for brightness of fluorescence relative to siblings. Mendelian ratios were observed, supporting that selected larvae were homozygous for a given allele.

METHOD DETAILS

Confocal imaging

Larvae were anesthetized in 0.2 mg/mL ethyl-3-aminobenzoic acid ethyl ester (MESAB, Sigma-Aldrich E10521, St. Louis, MO) prior to imaging except where noted. Larvae were mounted dorsal side-up (axial view) or lateral side-up (sagittal view) in 2% low-melting point agarose (Thermo Fisher Scientific 16520) in E3. Birthdating images of the vestibular ganglion were collected on a Zeiss LSM800 confocal microscope with a 20x water-immersion objective (Zeiss W Plan-Apochromat 20x/1.0). All imaging windows were 160x160μm. Anatomy stacks spanned approximately 60μm in depth (mediolateral axis), sampled every micron. For two-photon imaging of the tangential nucleus, stimulus imaging windows were 297x86μm. Anatomy stacks of the tangential nucleus spanned 50–60μm, sampled every micron (dorsoventral axis). All anatomy stacks were collected between 4–5 dpf. Raw image stacks were analyzed using Fiji/ImageJ (Schindelin et al., 2012).

Optical tagging of neurons by birthdate

All experiments used larvae from *Tg(-6.7Tru.Hcrtr2:GAL4-VP16)*;*Tg(UAS:E1b-Kaede)*. Neurons were optically tagged by their time of terminal differentiation using whole-embryo Kaede photoconversions (Caron et al., 2008) as described in (Goldblatt et al., 2023). Briefly, embryos were photoconverted for five minutes on a custom-built apparatus with a 405 nm LED bulb, measured at 9 mW power using a 9.5 mm probe (Thorlabs S130C). Photoconversions were performed at experimenter-defined timepoints between 22–72 hpf. Embryos used for experiments were converted at only one timepoint. Larvae were subsequently raised in darkness to prevent background photoconversion until the time of imaging (5 dpf). Neurons born before the time of conversion were identified by the presence of converted (red) Kaede. For each experiment, basal conversion was estimated by imaging control larvae raised in darkness until confocal imaging. n=5 hemispheres from N=5 separate larvae were analyzed for each timepoint. The 60 hpf timepoint was only analyzed with respect to "born after" neurons. Only one hemisphere was analyzed for each larva.

Representations of vestibular ganglion neurons in example images and maps

The following image adjustments were performed to representative images and spatial registration data. First, to correlate the anatomy of mapped images with previous representations of vestibular ganglion topography (Liu et al., 2022), spatial maps were rotated 3.81 degrees anticlockwise. Next, the midpoints in the dorsoventral (Y) and rostrocaudal (X) axes and circumscribed boundaries shown in spatial maps and quantification plots were defined using the mean range of the n=345 neurons from N=5 *Tg(-6.7Tru.Hcrtr2:VP16-GAL4)* mapped control larvae. All representative images of photoconverted neurons ("born before") are shown after applying a threshold to subtract background and/or bleedthrough conversion (e.g., Figure 2B and Figure 5E). The thresholding method is described below ("Correction of background and/or bleedthrough Kaede conversion"). In image stacks, medial cells often appeared dimmer due to the light scattering properties of brain tissue. To aid visualization of all cells in representative maximum intensity projection images (e.g., Figure 1B, Figure 2B, and Figure 5E), brightness, contrast, and/or gamma were separately adjusted for medial and lateral planes.

Imaging and analysis of bipolar afferents

Bipolar afferent development was evaluated in the same larvae used for somatic birthdating experiments. As neuronal soma had brighter fluorescence than their bipolar afferents, to avoid oversaturation we acquired separate image stacks to evaluate projections to the VIII[th] nerve and utricular macula.

Acute lesion and calcium imaging experiments

Experiments used sighted larvae from *Tg(-6.7Tru.Hcrtr2:GAL4-VP16);Tg(UAS:E1b-Kaede);Tg(UAS:GCaMP6s)*. Larvae were birthdated at 26 hpf using whole-embryo conversions, then imaged on a confocal microscope at 4 dpf to identify neurons born before photoconversion. At 5 dpf, acute, uni-lateral lesion and calcium imaging experiments were performed using a two-photon microscope. Lesions were performed as originally validated in (Schoppik et al., 2017) and used in (Goldblatt et al., 2023; Hamling et al., 2024). To visualize the statoacoustic ganglion, larvae were mounted lateral side-up (sagittal view). Lesions were targeted to the soma of individual neurons born after the time of photoconversion (green, unconverted Kaede only) in one hemisphere only. Lesions were performed using a pulsed infrared laser (SpectraPhysics Spirit W) at 1040 mm (400 fs pulse duration, 4 pulses per cell over 10 ms) at 25–75 nJ per pulse. Anatomy stacks of the statoacoustic ganglion were acquired before and immediately after lesions to confirm the lesion extent (individual soma and secondarily, degraded axon projections). Larvae were left to recover for ten minutes and re-mounted axially for calcium imaging of the tangential nucleus. Calcium imaging after pitch-tilt stimulation was performed and analyzed as described in (Goldblatt et al., 2023; 2024; Hamling et al., 2022). Briefly, tonic 19° up/down tilts were delivered to larvae using a galvonometer and calcium activity was imaged immediately following tilt presentation. Control experiments used non-lesioned larvae to minimize potential off-target effects from commissural-projecting neurons.

Acute lesion and eye tilt behavior experiments

Lesions were performed as described above. Larvae were from *Tg(-6.7Tru.Hcrtr2:GAL4-VP16);Tg(UAS:E1b-Kaede)*. Larvae were birthdated at 26 hpf using whole-embryo conversions. At 5 dpf, bi-lateral lesions were performed: late-born (green, unconverted Kaede only) vestibular ganglion neurons or the anterior lateral line ganglion neurons (sham lesion control). Larvae were left to recover for ten minutes prior to eye tilt behavior experiments.

Torsional eye rotations were measured in response to step tilts delivered using an apparatus similar in design to (Bianco et al., 2012). All experiments took place in complete darkness. Lesioned larvae (vestibular ganglion or sham) were immobilized completely in 2% low-melting temperature agarose, and the left eye freed. The agarose was then pinned (0.1 mm stainless minutien pins, FST) to a 5 mm2 piece of Sylgard 184 (Dow Corning) which was itself pinned to Sylgard 184 at the bottom of a 10 mm2 optical glass cuvette (Azzota, via Amazon). The cuvette was filled with approx. 1 mL of E3 and placed in a custom holder on a 5-axis (X, Y, Z, pitch, roll) manipulator (ThorLabs MT3 and GN2). The fish was aligned with the optical axes of two orthogonally placed cameras such that both the left utricle and two eyes were level with the horizon (front camera) and centered about the axis of rotation. The eye-monitoring camera (Guppy Pro 2 F-031, Allied Vision Technologies) used a 5x objective (Olympus MPLN, 0.1 NA) and custom image-forming optics to create a 100x100 pixel image of the left eye of the fish (6 µm/pixel), acquired at 200 Hz. A stepper motor (Oriental Motors AR98MA-N5-3) was used to rotate the platform holding the cameras and fish. An experiment consisted of 50 cycles of four steps each. Steps were $\pm15°$ towards and away from the horizon. Each step followed a trapezoidal velocity profile peaking at 35°/sec, peak acceleration 150°/sec2. The platform velocity and acceleration were measured using integrated circuits (IDG500, Invensense and ADXL335, Analog Devices) mounted together on a breakout board (Sparkfun SEN-09268).

STATISTICAL ANALYSES AND QUANTIFICATION

Validation of vestibular ganglion labeling in *Tg(-6.7Tru.Hcrtr2:GAL4-VP16)*

For initial validation that *Tg(-6.7Tru.Hcrtr2:GAL4-VP16)* labels neurons within the anatomical constraints of the vestibular ganglion, *Tg(-6.7Tru.Hcrtr2:GAL4-VP16)* expression was compared directly against *Tg(-17.6isl2b:GFP)* in double transgenic embryos. *Tg(-6.7Tru.Hcrtr2:GAL4-VP16)* was labeled with a red UAS reporter (photoconverted *Tg(UAS-E1b:Kaede)*) and compared with co-localized *Tg(-17.6isl2b:GFP)* expression. After validating co-localization, all subsequent quantification of vestibular ganglion density and cell position (Figure 1D-Figure 1I) was performed in singly-labeled transgenic larvae. We evaluated n=220 neurons from N=3 *Tg(-17.6isl2b:GFP)* and n=345 neurons from N=5 *Tg(-6.7Tru.Hcrtr2:GAL4-VP16)* larvae.

Spatial registration of vestibular ganglion soma

Neurons were manually registered to a common coordinate framework in Adobe Illustrator (2021). The maximum intensity projections (MIPs) from all imaged larvae were aligned using stereotyped and clearly-identifiable anatomical

landmarks: the saccule, the utricle, the edge of the otic capsule, and the VIII[th] nerve branch. To control for poten-
tial bias, MIP alignment was verified by two independent observers (D.G. and S.H.). To control for differences in
digital magnification across microscopes and larvae, all MIPs were uniformly scaled to a standardized resolution (5.0
pixels/µm).

Vestibular ganglion neurons labeled by *Tg(-6.7Tru.Hcrtr2:GAL4-VP16)* were approximately 10 µm in diameter. To
simplify localization of neurons in the mediolateral (Z) axis, each anatomical image stack was subdivided into five, 12
µm thick planes (Figure 1E, Figure 1G), such that each neuron would appear in no more than two mediolateral planes.
Subdivisions were assessed for consistency using anatomical landmarks. The caudal-most neurons occupied the first
mediolateral plane (z1), and the rostral-most neurons occupied the last mediolateral plane (z5). The mediolateral mid-
point (z3) was qualitatively aligned with the dorsoventral (Y) and rostrocaudal (X) midpoints. Each mediolateral plane
was aligned with its MIP. Neurons were manually localized to a mediolateral subdivision based on the plane in which the
soma fluorescence was brightest (soma center). Reference neurons, represented as circles approximately the diameter
of a neuron, were centered over the soma in the appropriate subdivision. The mediolateral subdivision and the rostro-
caudal (X) and dorsoventral (Y) Illustrator coordinates were manually recorded for each neuron. For standardization,
XY coordinates were normalized by subtraction of the upper-most and left-most coordinates. Standardized coordinates
were then used to recreate a spatial map of neurons imaged across all fish in Matlab. Cell counts were concurrently
performed in Fiji/ImageJ (Schindelin et al., 2012) using the Cell Counter plugin.

Representations, alignment, and analysis of electron microscopy data

Electron microscopy (EM) images of the vestibular ganglion, its afferent projections, and the tangential nucleus were
reconstructed using Catmaid as described in (Liu et al., 2022). Catmaid reconstructions were exported as a catalog SVG
and imported into Illustrator for alignment. Spatial scales were preserved during image import and alignment with ref-
erence anatomical images from our *in vivo* transgenic maps. Given the minor size distortions (approx. 10 µm) between
the EM reconstructions and our *in vivo* maps, presumably a fixation artifact, the EM images were not quantitatively
aligned with transgenic maps. Instead, we generated a separate reference framework as described above. Reconstructed
images of the tangential nucleus, including analyses of tangential nucleus neurons that received synaptic input from
utricular-innervated ganglion neurons, was based on previously-published analyses, available in the supplementary ma-
terial (supplementary data 1 and 2) in (Liu et al., 2022).

Spatial registration and quantification of converted bipolar afferent projections

Maximum intensity projections were aligned using anatomical landmarks (otic capsule, utricle, etc.) into a hyperstack
in Fiji/ImageJ (Schindelin et al., 2012). A 15 µm or 35 µm-sized rectangular ROI was drawn around the extent of the
macula and VIII[th] nerve axon projections, respectively. The fluorescence gray value along the rostrocaudal axis of the
ROI was extracted using the "Plot XY Profile" function for each channel (unconverted green, converted red), and the
X profile (rostrocaudal axis) was obtained. As neurons continue to generate new, unconverted Kaede fluorescence after
the time of conversion, unconverted (green) fluorescence comprises projections from neurons born both before and after
the time of conversion. Fluorescence values were imported into Matlab. To account for variation in Kaede fluorescence
across larvae due to transgenic expression, fluorescence values were standardized by the maximum raw fluorescence
observed for each fish. Probability distributions were generated, and the median/standard deviation were calculated.
N=3 hemispheres/timepoint (N=1 hemisphere per larva) were quantified.

Correction of background and/or bleedthrough Kaede conversion

Basal Kaede conversion was estimated by imaging a total of N=18 unconverted (green only, sibling controls) larvae
raised in darkness until imaging at 5 dpf. Control larvae had comparable fluorescence as experimental larvae and were
imaged alongside experimental larvae using identical imaging settings. The amount of fluorescence observed in the red
(converted Kaede) channel was measured using Fiji/ImageJ (Schindelin et al., 2012). As controls were never exposed
to light until the time of imaging, any red fluorescence was either basal conversion or unconverted (green) fluorescence
bleedthrough. The average red Kaede fluorescence value across the vestibular ganglion was used as a "threshold" and
subtracted from all birthdated images. This ensured that cells designated as born based on the presence of red Kaede
were not artifacts of basal or bleedthrough fluorescence.

Statistical analysis of spatial organization

For each timepoint, data from all hemispheres analyzed (N=5/timepoint) were pooled given variability in the number of vestibular ganglion neurons labeled across transgenic larve (median: 69±15 neurons/fish across N=5 control hemispheres; see Figure 1I). The number of "born before" and "born after" neurons mapped per fish is listed in Table 1. Probability distributions of soma position were generated using the median and standard deviation from pooled and bootstrapped data (n=100 iterations). Spatial organization was evaluated individually in each spatial axis using a two-tailed Kolmogorov-Smirnov (KS) test given that most distributions were bimodal. All control distributions used the n=345 neurons within the boundaries of the vestibular ganglion mapped from the N=5 *Tg(-6.7Tru.Hcrtr2:GAL4-VP16)* control larvae described above.

Quantification and analysis of acute lesion and calcium imaging experiments

Calcium responses in the tangential nucleus were extracted and analyzed as described in (Goldblatt et al., 2023; 2024). In Fiji/ImageJ, regions of interest (ROI) were drawn around each tangential nucleus soma and raw fluorescence values were extracted using Matlab. Calcium responses were normalized using the mean fluorescence value of a baseline period, and the first second of the calcium response was used for analysis. Responses were analyzed with respect to two primary metrics: (1) nose-up/nose-down identity, defined using a directionality index that compared the calcium responses to up/down tilt stimulation; (2) spatial organization, mapped in Illustrator using the same methodology as described above and in (Goldblatt et al., 2023; 2024). In cell position maps, plotted neuron centers were scaled to match the strength of their fluorescence responses, evaluated using the maximum value of the $\Delta F/F$ fluorescent response.

All identifiable neurons within the tangential nucleus were analyzed. We analyzed the responses of n=140 tangential nucleus neurons from N=4 control, never-lesioned larvae (median: 36±7 neurons/larva) and n=110 neurons from N=4 larvae following acute, uni-lateral vestibular ganglion lesion (median: 28±5 neurons/larva). Qualitatively, we observed that tangential nucleus neurons in larvae after vestibular ganglion lesions had lower basal fluorescence, which may account for variation in the number of identifiable neurons per larva. All Wilcoxon rank sum tests for differences of medians were performed on a per-larva basis. Statistical analysis of neuron position were performed on aggregate data from all larvae to account for spatial biases in transgenic expression.

Analysis of eye tilt behavior data

The image was processed on-line by custom pattern matching software to derive an estimate of torsional angle (LabView 2014, National Instruments), and data was analyzed using custom MATLAB scripts (Mathworks, Natick MA). The eye's response across the experiment was first centered to remove any offset introduced by the pattern-matching algorithm. Data were then interpolated with a cubic spline to correct for occasional transient slowdowns (i.e. missed frames). The eye's velocity was estimated by differentiating the position trace; high-frequency noise was minimized using a 4-pole low-pass Butterworth filter (cutoff = 3 Hz). Each step response was evaluated manually; trials with rapid deviations in eye position indicative of horizontal saccades or gross failure of the pattern matching algorithm were excluded from analysis. The response to each step for a given fish was defined as the mean across all responses to that step across cycles. The gain was estimated by measuring the peak eye velocity occurring over the first second after the start of the step.

ACKNOWLEDGMENTS

Research was supported by the National Institute on Deafness and Communication Disorders of the National Institutes of Health under award numbers R01DC017489 and the National Institute of Neurological Disorders and Stroke under award numbers F99NS129179 and T32NS086750. The authors would like to thank Hannah Gelnaw for assistance with fish care, Martha Bagnall for invaluable insights and assistance with EM data, Jeremy Dasen, Claude Desplan, Katherine Nagel, Dan Sanes, and Masashi Tanimoto along with the members of the Schoppik and Nagel labs for their valuable feedback and discussions.

AUTHOR CONTRIBUTIONS

Conceptualization: SH, DG and DS, Methodology: SH, DG and DS, Investigation: SH, MRG, EG, and SND, Visualization: SH, Writing: SH, Editing: DG and DS, Funding Acquisition: DG and DS, Supervision: DG.

AUTHOR COMPETING INTERESTS

The authors declare no competing interests.

STATEMENT ON ARTIFICIAL INTELLIGENCE USE

No artificial intelligence tools were used in the preparation of this manuscript, including code generation, image production, data collection, analysis, and writing.

REFERENCES

J. Altman and S. A. Bayer. Development of the brain stem in the rat. V. Thymidine-radiographic study of the time of origin of neurons in the midbrain tegmentum. *The Journal of Comparative Neurology*, 198(4):677–716, Jun 1981. doi: 10.1002/cne.901980409. URL http://dx.doi.org/10.1002/cne.901980409.

J. Altman and S. A. Bayer. *Development of the Cranial Nerve Ganglia and Related Nuclei in the Rat*. Springer Berlin Heidelberg, 1982. ISBN 9783642684791. doi: 10.1007/978-3-642-68479-1. URL http://dx.doi.org/10.1007/978-3-642-68479-1.

S. Arber. Motor circuits in action: Specification, connectivity, and function. *Neuron*, 74(6):975–989, jun 2012. doi: 10.1016/j.neuron.2012.05.011. URL https://doi.org/10.1016/j.neuron.2012.05.011.

F. Auclair, R. Marchand, and J. C. Glover. Regional patterning of reticulospinal and vestibulospinal neurons in the hindbrain of mouse and rat embryos. *The Journal of Comparative Neurology*, 411(2):288–300, Aug. 1999. ISSN 1096-9861. doi: 10.1002/(sici)1096-9861(19990823)411: 2⟨288::aid-cne9⟩3.0.co;2-u. URL http://dx.doi.org/10.1002/(SICI)1096-9861(19990823)411:2<288::AID-CNE9>3.0.CO;2-U.

S. Baeza-Loya, J. T. Bùi, and D. W. Raible. Activity and retinoic acid drive hair cell spatial patterning in the zebrafish utricle. *bioRxiv*, Nov. 2025. doi: 10.1101/2025.11.17.687290. URL http://dx.doi.org/10.1101/2025.11.17.687290.

M. W. Bagnall and D. Schoppik. Development of vestibular behaviors in zebrafish. *Current Opinion in Neurobiology*, 53:83–89, Dec. 2018. doi: 10.1016/j.conb.2018.06.004. URL https://doi.org/10.1016/j.conb.2018.06.004.

N. Balaskas, D. Ng, and N. Zampieri. The positional logic of sensory-motor reflex circuit assembly. *Neuroscience*, 450:142–150, Dec. 2020. ISSN 0306-4522. doi: 10.1016/j.neuroscience.2020.04.038. URL http://dx.doi.org/10.1016/j.neuroscience.2020.04.038.

G. R. Barsh, A. J. Isabella, and C. B. Moens. Vagus motor neuron topographic map determined by parallel mechanisms of hox5 expression and time of axon initiation. *Current Biology*, 27(24):3812–3825.e3, Dec. 2017. doi: 10.1016/j.cub.2017.11.022. URL https://doi.org/10.1016/j.cub.2017.11.022.

O. A. Bayraktar and C. Q. Doe. Combinatorial temporal patterning in progenitors expands neural diversity. *Nature*, 498(7455):449–455, June 2013. doi: 10.1038/nature12266. URL https://doi.org/10.1038/nature12266.

A. Bañón and B. Alsina. Pioneer statoacoustic neurons guide neuroblast behaviour during otic ganglion assembly. *Development*, 150(21), Nov. 2023. ISSN 1477-9129. doi: 10.1242/dev.201824. URL http://dx.doi.org/10.1242/dev.201824.

D. Bell, A. Streit, I. Gorospe, I. Varela-Nieto, B. Alsina, and F. Giraldez. Spatial and temporal segregation of auditory and vestibular neurons in the otic placode. *Developmental Biology*, 322(1):109–120, Oct. 2008. ISSN 0012-1606. doi: 10.1016/j.ydbio.2008.07.011. URL http://dx.doi.org/10.1016/j.ydbio.2008.07.011.

M. Berry, A. W. Rogers, and J. T. Eayrs. Pattern of cell migration during cortical histogenesis. *Nature*, 203(4945):591–593, Aug. 1964. doi: 10.1038/203591b0. URL https://doi.org/10.1038/203591b0.

I. H. Bianco, L.-H. Ma, D. Schoppik, D. N. Robson, M. B. Orger, J. C. Beck, J. M. Li, A. F. Schier, F. Engert, and R. Baker. The tangential nucleus controls a gravito-inertial vestibulo-ocular reflex. *Current Biology*, 22(14):1285–1295, Jul 2012. doi: 10.1016/j.cub.2012.05.026. URL http://dx.doi.org/10.1016/j.cub.2012.05.026.

J. B. Bikoff, M. I. Gabitto, A. F. Rivard, E. Drobac, T. A. Machado, A. Miri, S. Brenner-Morton, E. Famojure, C. Diaz, F. J. Alvarez, G. Z. Mentis, and T. M. Jessell. Spinal inhibitory interneuron diversity delineates variant motor microcircuits. *Cell*, 165(1):207–219, mar 2016. doi: 10.1016/j.cell.2016.01.027. URL https://doi.org/10.1016/j.cell.2016.01.027.

J. Cang and D. A. Feldheim. Developmental mechanisms of topographic map formation and alignment. *Annual Review of Neuroscience*, 36(1):51–77, July 2013. ISSN 1545-4126. doi: 10.1146/annurev-neuro-062012-170341. URL http://dx.doi.org/10.1146/annurev-neuro-062012-170341.

J. Cang, C. M. Niell, X. Liu, C. Pfeiffenberger, D. A. Feldheim, and M. P. Stryker. Selective disruption of one cartesian axis of cortical maps and receptive fields by deficiency in ephrin-as and structured activity. *Neuron*, 57(4):511–523, Feb. 2008. ISSN 0896-6273. doi: 10.1016/j.neuron.2007.12.025. URL http://dx.doi.org/10.1016/j.neuron.2007.12.025.

S. J. C. Caron, D. Prober, M. Choy, and A. F. Schier. *In vivo* birthdating by BAPTISM reveals that trigeminal sensory neuron diversity depends on early neurogenesis. *Development*, 135(19):3259–3269, Aug 2008. doi: 10.1242/dev.023200. URL http://dx.doi.org/10.1242/dev.023200.

T.-W. Chen, T. J. Wardill, Y. Sun, S. R. Pulver, S. L. Renninger, A. Baohan, E. R. Schreiter, R. A. Kerr, M. B. Orger, V. Jayaraman, L. L. Looger, K. Svoboda, and D. S. Kim. Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature*, 499(7458):295–300, July 2013. doi: 10.1038/nature12354. URL https://doi.org/10.1038/nature12354.

H.-J. Cheng, M. Nakamoto, A. D. Bergemann, and J. G. Flanagan. Complementary gradients in expression and binding of elf-1 and mek4 in development of the topographic retinotectal projection map. *Cell*, 82(3):371–381, Aug. 1995. ISSN 0092-8674. doi: 10.1016/0092-8674(95)90426-3. URL http://dx.doi.org/10.1016/0092-8674(95)90426-3.

C. Clark, O. Austen, I. Poparic, and S. Guthrie. α2-chimaerin regulates a key axon guidance transition during development of the oculomotor projection. *Journal of Neuroscience*, 33(42):16540–16551, Oct 2013. doi: 10.1523/jneurosci.1869-13.2013. URL http://dx.doi.org/10.1523/JNEUROSCI.1869-13.2013.

J. S. Dasen. *Chapter 4 Transcriptional Networks in the Early Development of Sensory–Motor Circuits*, page 119–148. Elsevier, 2009. ISBN 9780123744692. doi: 10.1016/s0070-2153(09)01204-6. URL http://dx.doi.org/10.1016/S0070-2153(09)01204-6.

S. Delmaghani and A. El-Amraoui. The genetic and phenotypic landscapes of usher syndrome: from disease mechanisms to a new classification. *Human Genetics*, 141(3–4):709–735, Mar. 2022. ISSN 1432-1203. doi: 10.1007/s00439-022-02448-7. URL http://dx.doi.org/10.1007/s00439-022-02448-7.

M. Dombrovski, Y. Zang, G. Frighetto, A. Vaccari, H. Jang, P. S. Mirshahidi, F. Xie, P. Sanfilippo, B. W. Hina, A. Rehan, R. H. Hussein, P. S. Mirshahidi, C. Lee, M. A. Frye, C. R. von Reyn, Y. Z. Kurmangaliyev, G. M. Card, and S. L. Zipursky. Gradients of recognition molecules shape synaptic specificity of a visuomotor transformation. *bioRxiv*, Sept. 2024. doi: 10.1101/2024.09.04.610846. URL http://dx.doi.org/10.1101/2024.09.04.610846.

S. Dyballa, T. Savy, P. Germann, K. Mikula, M. Remesikova, R. Špir, A. Zecca, N. Peyriéras, and C. Pujades. Distribution of neurosensory progenitor pools during inner ear morphogenesis unveiled by cell lineage reconstruction. *eLife*, 6, jan 2017. doi: 10.7554/elife.22268. URL https://doi.org/10.7554/elife.22268.

R. S. Erzurumlu, Y. Murakami, and F. M. Rijli. Mapping the face in the somatosensory brainstem. *Nature Reviews Neuroscience*, 11(4):252–263, Feb. 2010. doi: 10.1038/nrn2804. URL https://doi.org/10.1038/nrn2804.

I. A. Favre-Bulle, G. Vanwalleghem, M. A. Taylor, H. Rubinsztein-Dunlop, and E. K. Scott. Cellular-resolution imaging of vestibular processing across the larval zebrafish brain. *Current Biology*, 28(23):3711–3722.e3, Dec. 2018. doi: 10.1016/j.cub.2018.09.060. URL https://doi.org/10.1016/j.cub.2018.09.060.

J. R. Fetcho and D. L. McLean. Some principles of organization of spinal neurons underlying locomotion in zebrafish and their implications. *Annals of the New York Academy of Sciences*, 1198(1):94–104, Jun 2010. doi: 10.1111/j.1749-6632.2010.05539.x. URL http://dx.doi.org/10.1111/j.1749-6632.2010.05539.x.

J. G. Flanagan and P. Vanderhaeghen. The ephrins and eph receptors in neural development. *Annual Review of Neuroscience*, 21(1):309–345, Mar. 1998. ISSN 1545-4126. doi: 10.1146/annurev.neuro.21.1.309. URL http://dx.doi.org/10.1146/annurev.neuro.21.1.309.

S. Fore, F. Acuña-Hinrichsen, K. A. Mutlu, E. M. Bartoszek, B. Serneels, N. G. Faturos, K. T. P. Chau, M. I. Cosacak, C. D. Verdugo, F. Palumbo, C. Ringers, N. Jurisch-Yaksi, C. Kizil, and E. Yaksi. Functional properties of habenular neurons are determined by developmental stage and sequential neurogenesis. *Science Advances*, 6(36), Sept. 2020. doi: 10.1126/sciadv.aaz3173. URL https://doi.org/10.1126/sciadv.aaz3173.

B. Fritzsch and D. Nichols. Dii reveals a prenatal arrival of efferents at the differentiating otocyst of mice. *Hearing Research*, 65(1–2):51–60, Feb. 1993. ISSN 0378-5955. doi: 10.1016/0378-5955(93)90200-k. URL http://dx.doi.org/10.1016/0378-5955(93)90200-k.

L. Glenwinkel, S. R. Taylor, K. Langebeck-Jensen, L. Pereira, M. B. Reilly, M. Basavaraju, I. Rafi, E. Yemini, R. Pocock, N. Sestan, M. Hammarlund, D. M. Miller, and O. Hobert. In silico analysis of the transcriptional regulatory logic of neuronal identity specification throughout the c. elegans nervous system. *eLife*, 10, June 2021. doi: 10.7554/elife.64906. URL https://doi.org/10.7554/elife.64906.

J. C. Glover. Development of second order vestibular projections in the chicken embryoa. *Annals of the New York Academy of Sciences*, 781(1):13–20, June 1996. ISSN 1749-6632. doi: 10.1111/j.1749-6632.1996.tb15689.x. URL http://dx.doi.org/10.1111/j.1749-6632.1996.tb15689.x.

J. C. Glover. The development of vestibulo-ocular circuitry in the chicken embryo. *Journal of Physiology-Paris*, 97(1):17–25, Jan 2003. doi: 10.1016/j.jphysparis.2003.10.003. URL http://dx.doi.org/10.1016/j.jphysparis.2003.10.003.

D. Goldblatt, S. Huang, M. R. Greaney, K. R. Hamling, V. Voleti, C. Perez-Campos, K. B. Patel, W. Li, E. M. Hillman, M. W. Bagnall, and D. Schoppik. Neuronal birthdate reveals topography in a vestibular brainstem circuit for gaze stabilization. *Current Biology*, 33(7):1265–1281.e7, Apr. 2023. doi: 10.1016/j.cub.2023.02.048. URL https://doi.org/10.1016/j.cub.2023.02.048.

D. Goldblatt, B. Rosti, K. R. Hamling, P. Leary, H. Panchal, M. Li, H. Gelnaw, S. Huang, C. Quainoo, and D. Schoppik. Motor neurons are dispensable for the assembly of a sensorimotor circuit for gaze stabilization. *eLife*, Apr. 2024. doi: 10.7554/elife.96893.1. URL http://dx.doi.org/10.7554/eLife.96893.1.

M. R. Greaney, A. E. Privorotskiy, K. P. D'Elia, and D. Schoppik. Extraocular motoneuron pools develop along a dorsoventral axis in zebrafish, *Danio rerio*. *Journal of Comparative Neurology*, 525(1):65–78, jun 2016. doi: 10.1002/cne.24042. URL https://doi.org/10.1002/cne.24042.

C. Haddon and J. Lewis. Early ear development in the embryo of the zebrafish, danio rerio. *The Journal of Comparative Neurology*, 365(1):113–128, Jan. 1996. ISSN 1096-9861. doi: 10.1002/(sici)1096-9861(19960129)365:1⟨113::aid-cne9⟩3.0.co;2-6. URL http://dx.doi.org/10.1002/(SICI)1096-9861(19960129)365:1<113::AID-CNE9>3.0.CO;2-6.

C. Haddon, Y.-J. Jiang, L. Smithers, and J. Lewis. Delta-notch signalling and the patterning of sensory cell differentiation in the zebrafish ear: evidence from the mind bomb mutant. *Development*, 125(23):4637–4644, Dec. 1998. ISSN 1477-9129. doi: 10.1242/dev.125.23.4637. URL http://dx.doi.org/10.1242/dev.125.23.4637.

A. Hamid, H. Gattuso, A. N. Caglar, M. Pillai, T. Steele, A. Gonzalez, K. Nagel, and M. H. Syed. The conserved rna-binding protein imp is required for the specification and function of olfactory navigation circuitry in drosophila. *Current Biology*, 34(3):473–488.e6, Feb. 2024. ISSN 0960-9822. doi: 10.1016/j.cub.2023.12.020. URL http://dx.doi.org/10.1016/j.cub.2023.12.020.

K. R. Hamling, Y. Zhu, F. Auer, and D. Schoppik. Tilt in place microscopy: a simple, low-cost solution to image neural responses to body rotations. *The Journal of Neuroscience*, 43(6):936–948, Dec. 2022. ISSN 1529-2401. doi: 10.1523/jneurosci.1736-22.2022. URL http://dx.doi.org/10.1523/JNEUROSCI.1736-22.2022.

K. R. Hamling, K. Harmon, Y. Kimura, S.-i. Higashijima, and D. Schoppik. The vestibulospinal nucleus is a locus of balance development. *The Journal of Neuroscience*, 44(30):e2315232024, May 2024. ISSN 1529-2401. doi: 10.1523/jneurosci.2315-23.2024. URL http://dx.doi.org/10.1523/jneurosci.2315-23.2024.

K. L. Hammond and T. T. Whitfield. Fgf and hh signalling act on a symmetrical pre-pattern to specify anterior and posterior identity in the zebrafish otic placode and vesicle. *Development*, 138(18):3977–3987, Sept. 2011. ISSN 0950-1991. doi: 10.1242/dev.066639. URL http://dx.doi.org/10.1242/dev.066639.

S. Higashijima, Y. Hotta, and H. Okamoto. Visualization of cranial motor neurons in live transgenic zebrafish expressing green fluorescent protein under the control of the *Islet*-1 promoter/enhancer. *Journal of Neuroscience*, 20(1):206–218, Jan 2000. URL http://www.ncbi.nlm.nih.gov/pubmed/10627598.

I. Holguera and C. Desplan. Neuronal specification in space and time. *Science*, 362(6411):176–180, Oct. 2018. ISSN 1095-9203. doi: 10.1126/science.aas9435. URL http://dx.doi.org/10.1126/science.aas9435.

D. H. Hubel and T. N. Wiesel. Effects of monocular deprivation in kittens. *Naunyn-Schmiedebergs Archiv für Experimentelle Pathologie und Pharmakologie*, 248(6):492–497, 1964. ISSN 1432-1912. doi: 10.1007/bf00348878. URL http://dx.doi.org/10.1007/BF00348878.

R. Huszár, Y. Zhang, H. Blockus, and G. Buzsáki. Preconfigured dynamics in the hippocampus are guided by embryonic birthdate and rate of neurogenesis. *Nature Neuroscience*, Aug. 2022. doi: 10.1038/s41593-022-01138-x. URL https://doi.org/10.1038/s41593-022-01138-x.

A. J. Isabella, J. A. Stonick, J. Dubrulle, and C. B. Moens. Intrinsic positional memory guides target-specific axon regeneration in the zebrafish vagus nerve. *Development*, 148(18), Sept. 2021. doi: 10.1242/dev.199706. URL https://doi.org/10.1242/dev.199706.

G. S. X. E. Jefferis, E. C. Marin, R. F. Stocker, and L. Luo. Target neuron prespecification in the olfactory map of drosophila. *Nature*, 414(6860):204–208, Nov. 2001. doi: 10.1038/35102574. URL https://doi.org/10.1038/35102574.

Y. Jia and M. W. Bagnall. Monosynaptic targets of utricular afferents in the larval zebrafish. *Frontiers in Neurology*, 13, July 2022. doi: 10.3389/fneur.2022.937054. URL https://doi.org/10.3389/fneur.2022.937054.

W. Joo, L. Sweeney, L. Liang, and L. Luo. Linking cell fate, trajectory choice, and target selection: Genetic analysis of sema-2b in olfactory axon targeting. *Neuron*, 78(4):673–686, May 2013. ISSN 0896-6273. doi: 10.1016/j.neuron.2013.03.022. URL http://dx.doi.org/10.1016/j.neuron.2013.03.022.

C. B. Kimmel, K. Hatta, and W. K. Metcalfe. Early axonal contacts during development of an identified dendrite in the brain of the zebrafish. *Neuron*, 4(4):535–545, Apr. 1990. ISSN 0896-6273. doi: 10.1016/0896-6273(90)90111-r. URL http://dx.doi.org/10.1016/0896-6273(90)90111-R.

S. Kishore, E. B. Cadoff, M. A. Agha, and D. L. McLean. Orderly compartmental mapping of premotor inhibition in the developing zebrafish spinal cord. *Science*, 370(6515):431–436, Oct. 2020. doi: 10.1126/science.abb4608. URL https://doi.org/10.1126/science.abb4608.

E. I. Knudsen and M. S. Brainard. Visual instruction of the neural map of auditory space in the developing optic tectum. *Science*, 253(5015):85–87, July 1991. doi: 10.1126/science.2063209. URL https://doi.org/10.1126/science.2063209.

E. J. Koundakjian, J. L. Appler, and L. V. Goodrich. Auditory neurons make stereotyped wiring decisions before maturation of their targets. *Journal of Neuroscience*, 27(51):14078–14088, Dec. 2007. doi: 10.1523/jneurosci.3765-07.2007. URL https://doi.org/10.1523/jneurosci.3765-07.2007.

Y. Z. Kurmangaliyev, J. Yoo, S. A. LoCascio, and S. L. Zipursky. Modular transcriptional programs separately define axon and dendrite connectivity. *eLife*, 8, Nov. 2019. ISSN 2050-084X. doi: 10.7554/elife.50822. URL http://dx.doi.org/10.7554/eLife.50822.

A. M. Lacoste, D. Schoppik, D. N. Robson, M. Haesemeyer, R. Portugues, J. M. Li, O. Randlett, C. L. Wee, F. Engert, and A. F. Schier. A convergent and essential interneuron pathway for mauthner-cell-mediated escapes. *Current Biology*, 25(11):1526–1534, jun 2015. doi: 10.1016/j.cub.2015.04.025. URL https://doi.org/10.1016/j.cub.2015.04.025.

A. S. Laureano, K. Flaherty, A.-M. Hinman, A. Jadali, T. Nakamura, S.-i. Higashijima, H. E. Sabaawy, and K. Y. Kwan. shox2 is required for vestibular statoacoustic neuron development. *Biology Open*, 11(12), Dec. 2022. ISSN 2046-6390. doi: 10.1242/bio.059599. URL http://dx.doi.org/10.1242/bio.059599.

P. Leary, C. Bellegarda, C. Quainoo, D. Goldblatt, B. Rosti, and D. Schoppik. Sensation is dispensable for the maturation of the vestibulo-ocular reflex. *Science*, 387(6729):85–90, Jan. 2025. ISSN 1095-9203. doi: 10.1126/science.adr9982. URL http://dx.doi.org/10.1126/science.adr9982.

A. J. Levine, K. A. Lewallen, and S. L. Pfaff. Spatial organization of cortical and spinal neurons controlling motor behavior. *Current Opinion in Neurobiology*, 22(5):812–821, Oct. 2012. doi: 10.1016/j.conb.2012.07.002. URL https://doi.org/10.1016/j.conb.2012.07.002.

X. Li, T. Erclik, C. Bertet, Z. Chen, R. Voutev, S. Venkatesh, J. Morante, A. Celik, and C. Desplan. Temporal patterning of drosophila medulla neuroblasts controls neural fates. *Nature*, 498(7455):456–462, June 2013. doi: 10.1038/nature12319. URL https://doi.org/10.1038/nature12319.

Z. Liu and M. W. Bagnall. Organization of vestibular circuits for postural control in zebrafish. *Current Opinion in Neurobiology*, 82:102776, Oct. 2023. ISSN 0959-4388. doi: 10.1016/j.conb.2023.102776. URL http://dx.doi.org/10.1016/j.conb.2023.102776.

Z. Liu, D. G. C. Hildebrand, J. L. Morgan, Y. Jia, N. Slimmon, and M. W. Bagnall. Organization of the gravity-sensing system in zebrafish. *Nature Communications*, 13(1), Aug. 2022. doi: 10.1038/s41467-022-32824-w. URL https://doi.org/10.1038/s41467-022-32824-w.

A. Lunde, B. W. Okaty, S. M. Dymecki, and J. C. Glover. Molecular profiling defines evolutionarily conserved transcription factor signatures of major vestibulospinal neuron groups. *eneuro*, 6(1):ENEURO.0475–18.2019, Jan. 2019. ISSN 2373-2822. doi: 10.1523/eneuro.0475-18.2019. URL http://dx.doi.org/10.1523/ENEURO.0475-18.2019.

E. C. Maier and T. T. Whitfield. Ra and fgf signalling are required in the zebrafish otic vesicle to pattern and maintain ventral otic identities. *PLoS Genetics*, 10(12):e1004858, Dec. 2014. ISSN 1553-7404. doi: 10.1371/journal.pgen.1004858. URL http://dx.doi.org/10.1371/journal.pgen.1004858.

A. Maklad and B. Fritzsch. Development of vestibular afferent projections into the hindbrain and their central targets. *Brain Research Bulletin*, 60(5–6):497–510, June 2003. ISSN 0361-9230. doi: 10.1016/s0361-9230(03)00054-6. URL http://dx.doi.org/10.1016/s0361-9230(03)00054-6.

B. Mark, S.-L. Lai, A. A. Zarin, L. Manning, H. Q. Pollington, A. Litwin-Kumar, A. Cardona, J. W. Truman, and C. Q. Doe. A developmental framework linking neurogenesis and circuit formation in the drosophila CNS. *eLife*, 10, May 2021. doi: 10.7554/elife.67510. URL https://doi.org/10.7554/elife.67510.

J. P. Marti, A. Zecca, J. P. Baudoin, A. Faucherre, K. Asakawa, K. Kawakami, and H. L. Schier. Neuronal birth order identifies a dimorphic sensorineural map. *Journal of Neuroscience*, 32(9):2976–2987, Feb. 2012. doi: 10.1523/jneurosci.5157-11.2012. URL https://doi.org/10.1523/jneurosci.5157-11.2012.

D. L. McLean and J. R. Fetcho. Spinal interneurons differentiate sequentially from those driving the fastest swimming movements in larval zebrafish to those driving the slowest ones. *Journal of Neuroscience*, 29(43):13566–13577, Oct. 2009. doi: 10.1523/jneurosci.3277-09.2009. URL https://doi.org/10.1523/jneurosci.3277-09.2009.

D. L. McLean, J. Fan, S. ichi Higashijima, M. E. Hale, and J. R. Fetcho. A topographic map of recruitment in spinal cord. *Nature*, 446(7131):71–75, Mar. 2007. doi: 10.1038/nature05588. URL https://doi.org/10.1038/nature05588.

J. L. Meng, Z. D. Marshall, M. Lobb-Rabe, and E. S. Heckscher. How prolonged expression of hunchback, a temporal transcription factor, re-wires locomotor circuits. *eLife*, 8, Sept. 2019. doi: 10.7554/elife.46089. URL https://doi.org/10.7554/elife.46089.

J. L. Meng, Y. Wang, R. A. Carrillo, and E. S. Heckscher. Temporal transcription factors determine circuit membership by permanently altering motor neuron-to-muscle synaptic partnerships. *eLife*, 9, May 2020. doi: 10.7554/elife.56898. URL https://doi.org/10.7554/elife.56898.

G. Migault, T. L. van der Plas, H. Trentesaux, T. Panier, R. Candelier, R. Proville, B. Englitz, G. Debrégeas, and V. Bormuth. Whole-brain calcium imaging during physiological vestibular stimulation in larval zebrafish. *Current Biology*, 28(23):3723–3735.e6, Dec. 2018. doi: 10.1016/j.cub.2018.10.017. URL https://doi.org/10.1016/j.cub.2018.10.017.

A. L. Norovich, S. Brenner-Morton, and T. M. Jessell. Muscle-derived cues are required to specify proprioceptor pool identity. *bioRxiv*, Aug. 2022. doi: 10.1101/2022.08.18.504408. URL http://dx.doi.org/10.1101/2022.08.18.504408.

W. Penfield and E. Boldrey. Somatic motor and sensory representation in the cerebral cortex of man as studied by electrical stimulation. *Brain*, 60(4):389–443, 1937. doi: 10.1093/brain/60.4.389. URL https://doi.org/10.1093/brain/60.4.389.

M. Petrovic and T. Hummel. Temporal identity in axonal target layer recognition. *Nature*, 456(7223):800–803, Nov. 2008. ISSN 1476-4687. doi: 10.1038/nature07407. URL http://dx.doi.org/10.1038/nature07407.

A. J. Pittman, M.-Y. Law, and C.-B. Chien. Pathfinding in a large vertebrate axon tract: isotypic interactions guide retinotectal axons at multiple choice points. *Development*, 135(17):2865–2871, sep 2008. doi: 10.1242/dev.025049. URL https://doi.org/10.1242/dev.025049.

S. Poliak, A. L. Norovich, M. Yamagata, J. R. Sanes, and T. M. Jessell. Muscle-type identity of proprioceptors specified by spatially restricted signals from limb mesenchyme. *Cell*, 164(3):512–525, Jan. 2016. ISSN 0092-8674. doi: 10.1016/j.cell.2015.12.049. URL http://dx.doi.org/10.1016/j.cell.2015.12.049.

L. Puelles and A. Privat. Do oculomotor neuroblasts migrate across the midline in the fetal rat brain? *Anatomy and Embryology*, 150(2):187–206, 1977. ISSN 1432-0568. doi: 10.1007/bf00316650. URL http://dx.doi.org/10.1007/BF00316650.

A. Pujala and M. Koyama. Chronology-based architecture of descending circuits that underlie the development of locomotor repertoire after birth. *eLife*, 8, Feb. 2019. doi: 10.7554/elife.42135. URL https://doi.org/10.7554/elife.42135.

J. Pujol-Martí, J. Baudoin, A. Faucherre, K. Kawakami, and H. López-Schier. Progressive neurogenesis defines lateralis somatotopy. *Developmental Dynamics*, 239(7):1919–1930, June 2010. ISSN 1097-0177. doi: 10.1002/dvdy.22320. URL http://dx.doi.org/10.1002/dvdy.22320.

C. L. Rees, K. Moradi, and G. A. Ascoli. Weighing the evidence in peters' rule: Does neuronal morphology predict connectivity? *Trends in Neurosciences*, 40(2):63–71, Feb. 2017. ISSN 0166-2236. doi: 10.1016/j.tins.2016.11.007. URL http://dx.doi.org/10.1016/j.tins.2016.11.007.

B. B. Riley, C. Zhu, C. Janetopoulos, and K. J. Aufderheide. A critical period of ear development controlled by distinct populations of ciliated cells in the zebrafish. *Developmental Biology*, 191(2):191–201, Nov. 1997. ISSN 0012-1606. doi: 10.1006/dbio.1997.8736. URL http://dx.doi.org/10.1006/dbio.1997.8736.

R. Roberts, J. Elsner, and M. W. Bagnall. Delayed otolith development does not impair vestibular circuit formation in zebrafish. *Journal of the Association for Research in Otolaryngology*, 18(3):415–425, Mar. 2017. doi: 10.1007/s10162-017-0617-9. URL https://doi.org/10.1007/s10162-017-0617-9.

R. B. Roome, A. Yadav, L. Flores, A. Puarr, D. Nardini, A. Richardson, R. R. Waclaw, R. M. Arkell, V. Menon, J. E. Johnson, and A. J. Levine. Ontogeny of the spinal cord dorsal horn. *bioRxiv*, Mar. 2025. doi: 10.1101/2025.03.14.643370. URL http://dx.doi.org/10.1101/2025.03.14.643370.

A. Sagner, I. Zhang, T. Watson, J. Lazaro, M. Melchionda, and J. Briscoe. A shared transcriptional code orchestrates temporal patterning of the central nervous system. *PLOS Biology*, 19(11):e3001450, Nov. 2021. doi: 10.1371/journal.pbio.3001450. URL https://doi.org/10.1371/journal.pbio.3001450.

S. S. Saini and R. K. Ladher. Timing of delamination of inner ear neurons specifies their topography and target innervation. *bioRxiv*, Nov. 2024. doi: 10.1101/2024.11.25.625134. URL http://dx.doi.org/10.1101/2024.11.25.625134.

J. R. Sanes and S. L. Zipursky. Synaptic specificity, recognition molecules, and assembly of neural circuits. *Cell*, 181(3):536–556, Apr. 2020. ISSN 0092-8674. doi: 10.1016/j.cell.2020.04.008. URL http://dx.doi.org/10.1016/j.cell.2020.04.008.

D. Sapède and C. Pujades. Hedgehog signaling governs the development of otic sensory epithelium and its associated innervation in zebrafish. *The Journal of Neuroscience*, 30(10):3612–3623, Mar. 2010. ISSN 1529-2401. doi: 10.1523/jneurosci.5109-09.2010. URL http://dx.doi.org/10.1523/JNEUROSCI.5109-09.2010.

D. Sapède, S. Dyballa, and C. Pujades. Cell lineage analysis reveals three different progenitor pools for neurosensory elements in the otic vesicle. *The Journal of Neuroscience*, 32(46):16424–16434, Nov. 2012. ISSN 1529-2401. doi: 10.1523/jneurosci.3686-12.2012. URL http://dx.doi.org/10.1523/JNEUROSCI.3686-12.2012.

J. Schindelin, I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S. Preibisch, C. Rueden, S. Saalfeld, B. Schmid, J.-Y. Tinevez, D. J. White, V. Hartenstein, K. Eliceiri, P. Tomancak, and A. Cardona. Fiji: an open-source platform for biological-image analysis. *Nature Methods*, 9(7):676–682, Jun 2012. doi: 10.1038/nmeth.2019. URL http://dx.doi.org/10.1038/nmeth.2019.

J. Schmidt and L. Eisele. Stroboscopic illumination and dark rearing block the sharpening of the regenerated retinotectal map in goldfish. *Neuroscience*, 14(2):535–546, Feb. 1985. ISSN 0306-4522. doi: 10.1016/0306-4522(85)90308-2. URL http://dx.doi.org/10.1016/0306-4522(85)90308-2.

D. Schoppik, I. H. Bianco, D. A. Prober, A. D. Douglass, D. N. Robson, J. M. Li, J. S. Greenwood, E. Soucy, F. Engert, and A. F. Schier. Gaze-stabilizing central vestibular neurons project asymmetrically to extraocular motoneuron pools. *The Journal of Neuroscience*, 37(47): 11353–11365, sep 2017. doi: 10.1523/jneurosci.1711-17.2017. URL https://doi.org/10.1523/jneurosci.1711-17.2017.

S. Schwarzer, N. Asokan, O. Bludau, J. Chae, V. Kuscha, J. Kaslin, and S. Hans. Neurogenesis in the inner ear: the zebrafish statoacoustic ganglion provides new neurons from a neurod/nestin-positive progenitor pool well into adulthood. *Development*, 147(7), Apr. 2020. ISSN 0950-1991. doi: 10.1242/dev.176750. URL http://dx.doi.org/10.1242/dev.176750.

E. K. Scott, L. Mason, A. B. Arrenberg, L. Ziv, N. J. Gosse, T. Xiao, N. C. Chi, K. Asakawa, K. Kawakami, and H. Baier. Targeting neural circuitry in zebrafish using GAL4 enhancer trapping. *Nature Methods*, mar 2007. doi: 10.1038/nmeth1033. URL https://doi.org/10.1038/nmeth1033.

A. Q. Seroka and C. Q. Doe. The hunchback temporal transcription factor determines motor neuron axon and dendrite targeting in *Drosophila*. *Development*, Jan. 2019. doi: 10.1242/dev.175570. URL https://doi.org/10.1242/dev.175570.

M. D. Shaw and K. E. Alley. Generation of the ocular motor nuclei and their cell types in the rabbit. *The Journal of Comparative Neurology*, 200 (1):69–82, Jul 1981. doi: 10.1002/cne.902000106. URL http://dx.doi.org/10.1002/cne.902000106.

M. M. Shin, C. Catela, and J. Dasen. Intrinsic control of neuronal diversity and synaptic specificity in a proprioceptive circuit. *eLife*, 9, Aug. 2020. ISSN 2050-084X. doi: 10.7554/elife.56374. URL http://dx.doi.org/10.7554/eLife.56374.

O. Spead, T. Moreland, C. J. Weaver, I. D. Costa, B. Hegarty, K. L. Kramer, and F. E. Poulain. Teneurin trans-axonal signaling prunes topographically missorted axons. *Cell Reports*, 42(3):112192, Mar. 2023. ISSN 2211-1247. doi: 10.1016/j.celrep.2023.112192. URL http://dx.doi.org/10.1016/j.celrep.2023.112192.

R. W. Sperry. Chemoaffinity in the orderly growth of never fiber patterns and connections. *Proceedings of the National Academy of Sciences*, 50 (4):703–710, Oct. 1963. doi: 10.1073/pnas.50.4.703. URL https://doi.org/10.1073/pnas.50.4.703.

T. Sugioka, M. Tanimoto, and S.-i. Higashijima. Biomechanics and neural circuits for vestibular-induced fine postural control in larval zebrafish. *Nature Communications*, 14(1), Mar. 2023. ISSN 2041-1723. doi: 10.1038/s41467-023-36682-y. URL http://dx.doi.org/10.1038/s41467-023-36682-y.

L. F. Sullivan, T. L. Warren, and C. Q. Doe. Temporal identity establishes columnar neuron morphology, connectivity, and function in a drosophila navigation circuit. *eLife*, 8, Feb. 2019. doi: 10.7554/elife.43482. URL https://doi.org/10.7554/elife.43482.

G. Sürmeli, T. Akay, G. C. Ippolito, P. W. Tucker, and T. M. Jessell. Patterns of spinal sensory-motor connectivity prescribed by a dorsoventral positional template. *Cell*, 147(3):653–665, Oct. 2011. doi: 10.1016/j.cell.2011.10.012. URL https://doi.org/10.1016/j.cell.2011.10.012.

J. Szentágothai. Pathways and synaptic articulation patterns connecting vestibular receptors and oculomotor nuclei. In *The Oculomotor System*, pages 205–223. Hoeber Medical Division, Harper and Row, 1964.

L. Taberner, A. Bañón, and B. Alsina. Anatomical map of the cranial vasculature and sensory ganglia. *Journal of Anatomy*, 232(3):431–439, Dec. 2017. ISSN 1469-7580. doi: 10.1111/joa.12762. URL http://dx.doi.org/10.1111/joa.12762.

M. Tanimoto, Y. Ota, K. Horikawa, and Y. Oda. Auditory input to cns is acquired coincidentally with development of inner ear after formation of functional afferent pathway in zebrafish. *The Journal of Neuroscience*, 29(9):2762–2767, Mar. 2009. ISSN 1529-2401. doi: 10.1523/jneurosci.5530-08.2009. URL http://dx.doi.org/10.1523/JNEUROSCI.5530-08.2009.

M. Tanimoto, Y. Ota, M. Inoue, and Y. Oda. Origin of inner ear hair cells: Morphological and functional differentiation from ciliary cells into hair cells in zebrafish inner ear. *The Journal of Neuroscience*, 31(10):3784–3794, Mar. 2011. ISSN 1529-2401. doi: 10.1523/jneurosci.5554-10.2011. URL http://dx.doi.org/10.1523/JNEUROSCI.5554-10.2011.

M. Tanimoto, I. Watakabe, and S.-i. Higashijima. Tiltable objective microscope visualizes selectivity for head motion direction and dynamics in zebrafish vestibular system. *Nature Communications*, 13(1), Dec. 2022. ISSN 2041-1723. doi: 10.1038/s41467-022-35190-9. URL http://dx.doi.org/10.1038/s41467-022-35190-9.

T. R. Thiele, J. C. Donovan, and H. Baier. Descending control of swim posture by a midbrain nucleus in zebrafish. *Neuron*, 83(3):679–691, Aug. 2014. doi: 10.1016/j.neuron.2014.04.018. URL https://doi.org/10.1016/j.neuron.2014.04.018.

C. Toro, J. G. Trapani, I. Pacentine, R. Maeda, L. Sheets, W. Mo, and T. Nicolson. Dopamine modulates the activity of sensory hair cells. *The Journal of Neuroscience*, 35(50):16494–16503, Dec. 2015. ISSN 1529-2401. doi: 10.1523/jneurosci.1691-15.2015. URL http://dx.doi.org/10.1523/JNEUROSCI.1691-15.2015.

J. W. Triplett and D. A. Feldheim. Eph and ephrin signaling in the formation of topographic maps. *Seminars in Cell & Developmental Biology*, 23 (1):7–15, Feb. 2012. ISSN 1084-9521. doi: 10.1016/j.semcdb.2011.10.026. URL http://dx.doi.org/10.1016/j.semcdb.2011.10.026.

F. Ulrich, C. Grove, J. Torres-Vázquez, and R. Baker. Development of functional hindbrain oculomotor circuitry independent of both vascularization and neuronal activity in larval zebrafish. *Current Neurobiology*, 7:62–73, 2014. doi: 10.4172/0975-9042.000122. URL http://dx.doi.org/10.4172/0975-9042.000122.

S. Vemaraju, H. Kantarci, M. S. Padanad, and B. B. Riley. A spatial and temporal gradient of fgf differentially regulates distinct stages of neural

development in the zebrafish inner ear. *PLoS Genetics*, 8(11):e1003068, Nov. 2012. ISSN 1553-7404. doi: 10.1371/journal.pgen.1003068. URL http://dx.doi.org/10.1371/journal.pgen.1003068.

P. Wenner and E. Frank. Peripheral target specification of synaptic connectivity of muscle spindle sensory neurons with spinal motoneurons. *The Journal of Neuroscience*, 15(12):8191–8198, December 1995. ISSN 1529-2401. doi: 10.1523/jneurosci.15-12-08191.1995. URL http://dx.doi.org/10.1523/JNEUROSCI.15-12-08191.1995.

T. T. Whitfield, B. B. Riley, M. Chiang, and B. Phillips. Development of the zebrafish inner ear. *Developmental Dynamics*, 223(4):427–458, Mar. 2002. ISSN 1097-0177. doi: 10.1002/dvdy.10073. URL http://dx.doi.org/10.1002/dvdy.10073.

C. C. Wreden, J. L. Meng, W. Feng, W. Chi, Z. D. Marshall, and E. S. Heckscher. Temporal cohorts of lineage-related neurons perform analogous functions in distinct sensorimotor circuits. *Current Biology*, 27(10):1521–1528.e4, May 2017. doi: 10.1016/j.cub.2017.04.024. URL https://doi.org/10.1016/j.cub.2017.04.024.

D. Wu, I. Schieren, Y. Qian, C. Zhang, T. M. Jessell, and J. C. de Nooij. A role for sensory end organ-derived signals in regulating muscle spindle proprioceptor phenotype. *The Journal of Neuroscience*, 39(22):4252–4267, Mar. 2019. ISSN 1529-2401. doi: 10.1523/jneurosci.2671-18. 2019. URL http://dx.doi.org/10.1523/JNEUROSCI.2671-18.2019.

S. Yogev and K. Shen. Cellular and molecular mechanisms of synaptic specificity. *Annual Review of Cell and Developmental Biology*, 30 (1):417–437, Oct. 2014. ISSN 1530-8995. doi: 10.1146/annurev-cellbio-100913-012953. URL http://dx.doi.org/10.1146/annurev-cellbio-100913-012953.

M. A. Younes, M. Rima, J.-P. Coutanceau, N. Peysson, F. Hoseinkhani, E. Rodrigues, E. Bullier, P. Legendre, J.-M. Mangin, and E. Hong. Birth order specified recruitment of motor circuits during spontaneous neural activity in zebrafish embryo. *bioRxiv*, Aug. 2024. doi: 10.1101/2024. 08.27.609675. URL http://dx.doi.org/10.1101/2024.08.27.609675.

A. Zecca, S. Dyballa, A. Voltes, R. Bradley, and C. Pujades. The order and place of neuronal differentiation establish the topography of sensory projections and the entry points within the hindbrain. *Journal of Neuroscience*, 35(19):7475–7486, may 2015. doi: 10.1523/jneurosci.3743-14. 2015. URL https://doi.org/10.1523/jneurosci.3743-14.2015.

I. Zhang, G. L. Boezio, J. Cornwall-Scoones, T. Frith, M. Jiang, M. Howell, R. Lovell-Badge, A. Sagner, J. Briscoe, and M. J. Delás. The cis-regulatory logic integrating spatial and temporal patterning in the vertebrate neural tube. *bioRxiv*, Apr. 2024. doi: 10.1101/2024.04.17.589864. URL http://dx.doi.org/10.1101/2024.04.17.589864.

Figure 1: *Tg(-6.7Tru.Hcrtr2:GAL4-VP16)* **and** *Tg(-17.6isl2b:GFP)* **provide comparable but partially incomplete genetic access to vestibular sensory neurons in the anterior statoacoustic ganglion.**

(A) Schematic of information flow (top and left) in the vestibulo-ocular reflex circuit. Bipolar sensory neurons in the vestibular (anterior) ganglion are highlighted as the population of focus. Right inset shows transmitted light and fluorescent micrograph image of the vestibular ganglion, labeled in green by *Tg(-6.7Tru.Hcrtr2:GAL4-VP16);Tg(UAS-E1b:Kaede)*. White arrows point to bipolar afferents to the utricular macula (mac, bottom) and VIII[th] nerve (top); u, utricle; s, saccule; aSAG, anterior statoacoustic ganglion. **(B)** Visualization of vestibular ganglion neurons labeled by *Tg(-17.6isl2b:GFP)* (left; blue) and *Tg(-6.7Tru.Hcrtr2:GAL4-VP16);Tg(UAS-E1b:Kaede)* (middle; red) in the same larva (sagittal view). Black dashed circle outlines the extent of the vestibular ganglion. **(C)** Electron microscopy (EM) reconstruction from (Liu et al., 2022) of all utricular-innervated vestibular ganglion neurons in one larvae at 5 dpf (sagittal view). Dashed lines bound the ganglion. **(D)** Soma position of mapped neurons within the boundaries of the vestibular ganglion, in larvae that expressed either *Tg(-17.6isl2b:GFP)* (n=220 neurons from N=3 larvae), *Tg(-6.7Tru.Hcrtr2:GAL4-VP16);Tg(UAS-E1b:Kaede)* (n=345 neurons from N=5 larvae; a subset of n=220 randomly selected neurons are shown), or EM reconstructed (n=106 neurons from N=1 larva). The vertical and horizontal dashed lines over blue/red show the rostrocaudal and dorsoventral midpoints based on the full spatial extent of vestibular ganglion neurons observed in the n=345 neurons from N=5 *Tg(-6.7Tru.Hcrtr2:VP16-GAL4)* larvae (Methods). Dashed lines and midpoints over grey are based on the EM-specific bounding box shown in Figure 1C. **(E)** Representative images of the vestibular ganglion in *Tg(-6.7Tru.Hcrtr2:GAL4-VP16)* (top) and *Tg(-17.6isl2b:GFP)* (bottom) across five mediolateral subdivisions (orthogonal axis to Figure 1D). Black dashed lines outline ganglion neuron somata (circle) and the VIII[th] nerve (rectangle). **(F)** Maximum intensity projection of the EM-reconstructed ganglion neurons in an axial view, emphasizing the additional medial density. **(G)** Soma position of the neurons mapped in Figure 1D, projected in the orthogonal (mediolateral) axis (left). Maps from genetically-labeled neurons (left) localize neurons to one of the five mediolateral subdivisions shown in Figure 1E; right, EM reconstruction from Figure 1F. **(H)** Number of neurons counted from each fish (dot). Solid bar shows mean across larvae for isl2b/hcrtr2. **(I)** Probability of soma position in each spatial axis for all neurons mapped. Solid and shaded lines shown the mean and standard deviation, respectively using bootstrapped estimates of variance (Methods). Short vertical lines indicate the median position of neurons from each dataset. Grey/black vertical ticks delineate the spatial extent of the EM/transgenic volumes; black vertical lines are 10% probability. All scale bars are 20μm. "isl2b" refers to *Tg(-17.6isl2b:GFP)*. "hcrtr2" refers to *Tg(-6.7Tru.Hcrtr2:GAL4-VP16)*.

Figure 2: Early- and late-born cohorts of vestibular ganglion neurons are dorsoventrally opposed.
(A) Timeline of birthdating and imaging experiments. Time of differentiation is estimated by the presence of red and green Kaede (born before the time of photoconversion) or green Kaede only (born after photoconversion). Timeline shows all experimenter-defined conversion points; each larvae is converted only once. Yellow line highlights 26 hpf as the example timepoint shown in Figure 2B. **(B)** Example vestibular ganglion neurons (1) born before (left; magenta arrows) or after (middle; green arrows) 26 hpf (right: merge). Neurons are visualized by *Tg(-6.7Tru.Hcrtr2:GAL4-VP16);Tg(UAS-E1b:Kaede)* fish imaged at 5 dpf. The "born before 26 hpf" image shown was thresholded to remove background fluorescence (Methods). Top: Sagittal maximum intensity projection used for mapping. Bottom: The same image in an orthogonal plane highlighting the mediolateral location of early-born neurons. **(C)** Percent of neurons in each fish that were born before each timepoint. Black lines indicate the mean number across individual fish (circles). N=5 fish for each age. Magenta and green shaded rectangles highlight the number of neurons born at the timepoints selected in Figure 2D. **(D)** Location of neurons born before 26 hpf (left; "earliest-born" cohort) or after 48 hpf (right; "latest-born" cohort). Number of mapped neurons: early, n=88 neurons from N=10 fish (22 hpf and 26 hpf timepoints combined); late, n=58 neurons from N=5 fish. Dashed horizontal and vertical reference lines indicate the rostrocaudal and dorsoventral midpoints of the vestibular ganglion shown in Figure 1D. **(E)** Probability of soma position for the neurons shown in Figure 2D. Short vertical axis lines indicate the median position for control (black), early-born (magenta), or late-born (green) neurons. Solid and shaded lines shown the mean and standard deviation, respectively using bootstrapped estimates of variance (Methods). **(F)** Location of neurons from N=5 randomly selected individual fish from the N=10 larvae aggregated in Figure 2D. **(G)** EM reconstruction of myelinated (presumptive early-born neurons; magenta) and unmyelinated neurons (green) from one fish at 5 dpf. Data is reconstructed from (Liu et al., 2022) contains the same neurons shown in Figure 1C. **(H)** EM-reconstructed neurons, mapped to the same sagittal view framework as in Figure 2D and Figure 2F, to show the comparative localization of myelinated neurons at 5 dpf. **(I)** Probability of soma position for myelinated and unmyelinated neurons from the EM reconstruction. All scale bars, 20μm. Full statistics are shown in Table 2.

Figure 3: Position along the dorsoventral axis predicts late-born somatic accumulation within the vestibular ganglion.
(A) Soma position of neurons born before each timepoint (N=5 fish/timepoint). Dashed horizontal and vertical reference lines indicate the rostrocaudal and dorsoventral midpoints shown in Figure 1D). **(B)** Soma position of neurons born after each timepoint (N=5 fish/timepoint). **(C)** Dorsoventral position of neurons born before (magenta) or after (green) each timepoint, shown as median/standard deviation across aggregated data. Horizontal dashed line shows the midpoint. **(D)** Rostrocaudal position of neurons born before/after each timepoint, shown as the median/standard deviation across aggregated data. Full statistics are shown in Table 2.

Figure 4: The development of bipolar afferents to local utricular targets and the VIIIth nerve mirrors somatic topography.
(A) Circuit schematic. Black dashed lines outline the circuit elements of focus. **(B)** Representative image of the vestibular ganglion, visualized using *Tg(-6.7Tru.Hcrtr2:GAL4-VP16);Tg(UAS-E1b:Kaede)*. Rectangles outline bipolar afferents to the utricular macula (bottom) and VIIIth nerve (top). **(C)** Timeline of birthdating experiments. Larvae were converted at only one timepoint (22, 26, 30, or 36 hpf) and imaged at 5 dpf. **(D)** Vestibular ganglion projections terminating at the utricular macula at four converted timepoints (left, converted Kaede; middle, unconverted Kaede; right, merge). As neurons continue to generate new, uncoverted Kaede after the time of conversion, green projections may originate from both "born before" and "born after" neurons. Black dashed vertical line shows the approximate rostrocaudal extent and midpoint of vestibular ganglion projections based on total unconverted fluorescence (all projections). Data from one example fish/timepoint shown. **(E)** Probability of observing converted Kaede (magenta) along the rostrocaudal axis. Each distribution corresponds with the timepoint shown to the left (Figure 4D). Green shows the probability of observing unconverted Kaede. N=3 larvae quantified per timepoint. **(F)** Projections to the VIIIth nerve from converted vestibular ganglion somata. Dashed boxes outline the area of the projection bundle quantified in Figure 4G. Dashed circular outline shows somata that were born before the converted timepoint and were above thresholded background levels (Methods). Non-converted, below-threshold soma fluorescence is an artifact of adjusting image brightness to visualize the fainter projection bundle. **(G)** Zoom on VIIIth nerve projections of one fish at one timepoint. Black arrow highlights a faint but visibly converted projection converted at 22 hpf. Black dashed lines show the rostrocaudal extent and midpoint. **(H)** Probability of observing converted (magenta) or unconverted (green) Kaede fluorescence along the rostrocaudal axis for each timepoint shown in Figure 4G. All scale bars, 20 μm. Stars are p<0.05; n.s., not significant.

Figure 5: Early-born vestibular ganglion neurons assemble with spatiotemporally-matched central partners in the hindbrain. (A) VIII[th] nerve axons and tangential nucleus somata born before (left; magenta, 1) and after 22 hpf (middle; green, 2). Black arrows point to VIII[th] nerve innervation. Axons and neurons are visualized using *Tg(-6.7Tru.Hcrtr2:GAL4-VP16);Tg(UAS-E1b:Kaede)*. **(B)** EM reconstruction from (Liu et al., 2022) of VIII[th] nerve projections to the tangential nucleus at 5 dpf. Birth order is inferred from myelination (magenta, myelinated; green, unmyelinated). Black dashed box shows afferent organization at the tangential nucleus. Afferents from all n=106 reconstructed ganglion neurons are shown. **(C)** Pseudo-sagittal (orthogonal to Figure 5B) projection of VIII[th] nerve projections highlighting the dorso-medial entry point of myelinated ganglion afferents. Left image shows all ganglion afferents; right, only afferents with synaptic connections onto the tangential nucleus. **(D)** EM reconstruction of the tangential nucleus (n=17 neurons) in the same view as Figure 5C. Left: neurons are pseudocolored by whether they receive synaptic input from myelinated (magenta) or unmyelinated (green) vestibular ganglion afferents. Right: neurons pseudocolored according to inferred directionality (nose-up, orange; nose-down, blue, (Liu et al., 2022)) of their vestibular ganglion inputs. Semi-transparent neurons (ventral) receive input from semicircular canal afferents but not utricular afferents, and may be included in the functional maps shown in Figure 5H. **(E)** Top: Timeline of birthdating, lesion, and tilt experiments. Larvae are converted at 26 hpf. At 5 dpf, unconverted vestibular ganglion neurons (green 1) are lesioned. Central projection neurons expressing GCaMP6s are imaged following tilt stimulation. Bottom: Vestibular ganglion neurons born after (left) or before (middle/right) 26 hpf, pre- (middle) and post- (right) lesion in a thresholded image (Methods). Circle and dashed lines from Figure 1D. **(F)** Schematic of stimulation and calcium imaging paradigm. Shaded bars show imaging periods for baseline (grey), return from nose-down tilt (blue), and return from nose-up tilt (orange). Periods between shaded bars show time when the larvae is tilted away from the imaging plane. Inset shows timecourse of step to horizontal. **(G)** The number of tilt-responsive nose-up and nose-down neurons in a given larvae (circles) after lesion of vestibular ganglion neurons born after 26 hpf, quantified as a percentage of the mean number of up/down neurons measured across N=4 control larvae. **(H)** Location of tangential nucleus neurons in control and lesioned larvae. Each circle is an individual cell, scaled by the magnitude of tilt-evoked fluorescence change. n=172 neurons from N=4 control hemispheres and n=115 neurons from N=4 lesioned hemispheres. **(I)** Distribution of central neuron somata in the dorsoventral axis in control (grey) and lesioned (red) larvae shown in Figure 5H. **(J)** Top: Vestibulo-ocular reflex trial schematic. Bottom: Body tilt kinematics. **(K)** Average eye velocity following nose-up and nose-down body tilts in control (grey, N=6) and lesioned (red, N=7) larvae. **(L)** Peak mean eye velocity after late-born ganglion lesion, shown as a percentage of the mean observed in control larvae. Scale bars are 20 μm.

Figure 6: Birth order unifies vestibulo-ocular reflex circuit organization. **(A)** Functional topography for each vestibulo-ocular reflex circuit population, adapted from(Goldblatt et al., 2023; Greaney et al., 2016; Liu et al., 2022; Tanimoto et al., 2022). Orange and blue represent nose-up and nose-down domains for each nucleus (circles) and their axonal projections (horizontal lines). **(B)** Birth order-related organization of each circuit population, adapted from the present study (yellow box) and (Goldblatt et al., 2023; Greaney et al., 2016). Magenta and green represent early- and late-born cohorts, respectively. **(C)** Timeline of key developmental events for each circuit population in zebrafish. Data is from this study (yellow box) and (Bianco et al., 2012; Clark et al., 2013; Goldblatt et al., 2023; Greaney et al., 2016; Leary et al., 2025). Up/down function in the vestibular ganglion is inferred based on correlations between anatomical position (this study) and functional topography(Liu et al., 2022; Tanimoto et al., 2022). The timecourse of sensory-central synaptogenesis is yet to be determined in zebrafish.

Age converted	# cells/fish, born before	approx. % of control, born before	# cells/fish born after
22	7±2	10%	59±7
26	11±2	16%	49±7
30	21±2	30%	44±5
33	42±2	61%	36±5
36	40±5	58%	22±4
48	52±8	75%	12±2
60	n/a		3±1
72	63±5	91%	0.5±0.5
Non-birthdated reference dataset	69±15		

Table 1: Number of statoacoustic ganglion neurons converted across developmental timepoints. Data shown as mean/std number of neurons born before or after each timepoint from N=5 hemispheres/timepoint. "Non-birthdated reference dataset" refers to the median number of neurons per fish observed from a reference dataset of the n=345 neurons vestibular ganglion neurons mapped from N=5 larvae from *Tg(-6.7Tru.Hcrtr2:VP16-GAL4);Tg(UAS-E1b:Kaede)* (Figure 1D, Figure 1G).

Age converted	Median position born before (µm)	Median position born after (µm)	p_{KS} born before vs. born after	$p_{ANOVA, mult\ compare}$ born before vs. prev. age	$p_{ANOVA, mult\ compare}$ born after vs. prev. age
Dorsoventral					
22	17.3±7.2	33.5±12.7	9.9E-10		
26	23.4±9.4	34.7±13.3	7.6E-07	0.50	0.99
30	20.7±11.3	36.1±12.5	2.3E-12	0.99	0.80
33	27.1±13.1	40.7±12.5	9.0E-09	5.2E-05	0.08
36	25.0±12.0	38.9±13.6	2.2E-11	0.20	0.97
48	28.2±11.8	41.9±9.9	7.2E-09	0.18	0.34
60	n/a	45.5±12.7	n/a		0.99
72	34.5±13.9	53.3±2.0	0.04		0.99
Rostrocaudal					
22	42.7±7.1	34.9±13.9	7.6E-06		
26	42.2±6.9	35.7±16.2	2.2E-08	0.99	0.99
30	38.2±11.5	35.3±15.4	0.08	0.10	0.99
33	35.3±11.7	42.4±15.3	6.6E-07	0.94	0.48
36	36.6±13.2	39.4±17.1	1.1E-04	0.99	0.99
48	37.2±15.1	48.2±20.6	7.9E-04	0.97	0.99
60	n/a	51.3±7.3	n/a		0.66
72	38.2±14.1	21.8±17.1	0.28		0.24
Mediolateral					
22	36.0±7.7	36.0±10.5	0.01		
26	3.0±9.1	36.0±11.6	0.03	0.99	0.99
30	36.0±10.6	36.0±12.3	0.99	0.04	0.99
33	36.0±10.7	36.0±10.8	0.005	0.99	0.48
36	36.0±10.7	36.0±10.9	0.06	0.99	0.99
48	36.0±9.7	36.0±10.5	0.52	0.97	0.99
60		36.0±5.6			0.66
72	36.0±10.1	30±8.5	0.73		0.24

Table 2: Statistical comparisons of temporal cohort somatic organization. Median (standard deviation) positions are shown as microns from the dorsal- or rostral-most edges (0 µm) of the vestibular ganglion. Data is shown as an aggregate of all neurons mapped across N=5 hemispheres/timepoint.