1   Systematic functional assessment of anti-phage systems in their native host
2
3

4   Ellie David[1]*, Clarisse Plantady[1,2,3]*, Sophiane Poissonier[1], Josie Elliott[3], Elodie Kenck[3], Justine
5   Le Boulch[1], Arnaud Gutierrez[1]#, Anne Chevallereau[1,3]#
6
7   [1] Université Paris Cité, CNRS, INSERM, Institut Cochin, Paris 75014, France.
8   [2] Phaxiam Therapeutics, 60 Avenue Rockefeller, Bâtiment Bioserra, 69008 Lyon, France
9   [3] Molecular Microbiology and Structural Biochemistry (MMSB), CNRS UMR 5086, Université
10   Claude Bernard Lyon 1, Lyon, France
11
12
*, # equal contribution
Correspondance to: anne.chevallereau@cnrs.fr

Summary

Bacterial resistance to bacteriophages (phages) relies on two primary strategies: preventing
phage attachment and blocking post-attachment steps. These post-attachment mechanisms
are mediated by diverse defence systems, including DNA-degrading systems such as
Restriction-Modification (RM) and CRISPR-Cas, as along with abortive infection systems that
induce cell death or dormancy. Computational analyses suggest that bacterial genomes
encode multiple defence systems, which may act synergistically to enhance phage resistance.
However, the regulation, interactions, and ecological roles of these systems in native hosts
remain poorly understood. This study explored the role of eight predicted defence systems
in the clinical isolate NILS69 of *E. coli* by testing its susceptibility to 93 phages. Infectivity and
adsorption assays using mutants defective in these systems revealed that only PD-T4-3 and
RM systems restricted phages able to adsorb. The RM system acted via a predicted Type IV
endonuclease and was also able to limit plasmid conjugation if the plasmid was transferred
from a donor strain lacking a methylase, which is the hallmark of Type I, II or III RM systems.
Other defence systems showed no detectable activity, likely due to phage specificity,
environmental regulation, or cofactor requirements. These findings underscore the need for
further studies to investigate the regulation and ecological roles of bacterial defence systems
in their native host contexts.

Introduction

The balance between resistance and susceptibility of bacteria to their viruses (bacteriophages,
or phages) is governed by two primary strategies. The first strategy consists in preventing the
attachment of phages to the bacterial cell surface. This can occur through the loss,
modification, or masking of phage receptors, effectively rendering the host inaccessible to
infection. The second blocks post-adsorption steps in the phage lifecycle, a process mediated
by specialized defence systems. In recent years, more than a hundred novel defence systems
have been identified, revealing a remarkable diversity of mechanisms. These systems include
those that degrade invading phage DNA, such as Restriction-Modification (RM) systems and

CRISPR-Cas systems. Others inhibit phage gene expression or replication, exemplified by mechanisms like Viperins and chemical-based defences (1–3). Additionally, some defence systems rely on extreme strategies, including triggering programmed cell death or inducing dormancy, to stop phage proliferation. These mechanisms are known as abortive infection systems (4). The variety of such defences reflects the constant and dynamic evolutionary struggle between bacteria and their viral predators, as highlighted in recent literature (5).

Computational analyses based on remote sequence homology searches have predicted that the bacterial defensome (i.e., the set of defence genes present in the same genome) is composed of 5-6 defence systems in average, although this number is highly variable, even between related strains (6). These findings have led to the view of the defensome as an integrated bacterial immune system, where individual defence systems are involved in a complex network of interactions. Supporting this view, synergistic associations of defence systems have been reported recently (7–12). Carrying multiple defence systems in a single genome is thought to have two main positive effects on bacterial fitness: it protects against a larger diversity of phages (11,13), as individual defence systems may have a narrow specificity and restrict only few phage genera (13), and it limits the emergence of phages that escape bacterial defences (7,11,14,15). However, we still know very little about how defence systems interact with one another, how they are regulated, and how they impact the ecology and evolution of phage-bacteria interactions. One reason for this knowledge gap is that research necessarily needed an initial effort focused on scaling up the discovery and characterisation of defence systems, which rely on novel informatic tools and synthetic biology. Consequently, most defence systems were studied from their heterologous expression in model organisms such as *Escherichia coli* and *Bacillus subtilis* (16–18) . Comparatively, fewer defence systems have been studied in their original host, which hinders our understanding of the activity, regulation, fitness costs and benefits of defence genes (and their combinations) in their native condition. Recent studies have interrogated whether and how the composition of the defensome impacts the range of phages capable of infecting a host. Intuitively, one might expect a negative correlation between the number of phages infecting a host and the number and diversity of host defences. However, experimental findings have yielded contrasting results. A recent study evaluating interactions between clinical isolates of *Pseudomonas aeruginosa* and its phages found a positive correlation between the number of defence systems encoded in the isolates and their resistance to the tested phages (13). In contrast, a large-scale screen of phage-bacteria interactions in *Escherichia coli* found that, within this cohort, phage resistance was more likely determined by adsorption factors whereas a very low correlation was observed between the number of defence systems and the level of phage susceptibility (19). This suggests that, in this dataset, defence systems played only a marginal role in shaping the range of phages capable of infecting a host. Additionally, this study observed that the host range of specialist and generalist phages tended to overlap, generating an interaction matrix with a nested pattern.

In another study, the investigation of interactions between natural, longitudinally sampled, populations of *Vibrio* and their phages generated a modular interaction matrix: a specific subset of phages infected a specific subset of related hosts and thus, the host range of phages from two distinct subsets displayed very little overlap (20). In this case, the lack of interactions between phages and bacteria that belong to distinct modules is due to lack of adsorption. Conversely, the lack of interaction between phages and bacteria that belong to the same module is not explained by adsorption factors but correlates with the presence of defence

systems (20). In sum, the contribution of the defensome in determining phage-bacteria ecological interactions remains unclear. In addition, the functionality and role of defence systems in their native host have been overlooked, and therefore, little is known about their levels of expression, their regulation, and their interactions with other defence systems.

In this study, we aimed to complement ecological interaction matrices by providing a genetic approach to assess the relative contribution of defence systems and adsorption factors in shaping phage host range. Specifically, we employed a systematic genetic strategy to directly evaluate the activity of predicted defence systems in their native bacterial hosts. This approach allowed us to quantify the respective contributions of individual defence systems in restricting the host range of phages or modulating their infectivity levels. By integrating these direct genetic assessments with ecological interaction data, we sought to bridge the gap between theoretical predictions and empirical observations, offering a more nuanced understanding of the interplay between phage resistance mechanisms and host susceptibility.

Results

Phage sensitivity of *E. coli* clinical isolate NILS69 is mainly explained by adsorption factors

To test the role of defence systems in shaping phage susceptibility, we chose the clinical *E. coli* B2 phylogroup, uropathogenic isolate NILS69 as a model system. This strain is part of a collection of *E. coli* Natural Isolates with Low Subculture passages (NILS), developed to provide the community with *E. coli* natural isolates that cover the diversity of phylogroups while remaining as close as possible to the original isolated strain (21).

We initially evaluated phage infectivity in wildtype (WT) NILS69 by screening the recently published Antonina Guelin (AG) collection, which gathers virulent dsDNA phages belonging to 19 different genera (19), through solid and liquid infection succeptibility assays (Supplementary Figure 1A,B). Ten phages out of 93 tested were able to affect the growth of NILS69 in solid infection assay amongst which six affected NILS69 growth in liquid (Supplementary Table 1).

To determine whether the infectivity of certain phages could be revealed under different culture conditions, we varied the parameters of the solid infection assay, including incubation temperature (30°C, 42°C) and salt concentration (by adding 30 mM $MgSO_4$ and 15 mM $CaCl_2$ to the growth medium), but none of these conditions changed the range of infecting phages (Supplementary Figure 1A).

We next characterized the infectivity of these 10 phages in more details. In solid medium, they form plaques with low efficiency in efficiency of plating assays (EOP ranging from 10^{-5} to 10^{-1}), except for phage NIC06_P2 (Figure 1A). For some of these phages, infection on solid medium did not result in the formation of individual plaques but rather in a large lysis area. Therefore, here and for all EOP assays, we scored EOP as the highest dilution factor allowing the detection of phage lysis activity, divided by the PFU counted on the production strain. This lysis area, occurring at high phage concentration, is likely the reflection of the phage population overwhelming the host defence mechanisms. When introduced at a multiplicity of infection (MOI) of 1 in liquid medium, they show different impact on bacterial growth, which is either unaffected (536_P1, P6, P7, P9), delayed (LF73_P4, DIJ07_P1, AN24_P4) or strongly

suppressed (LF73_P1, LF73_P3, NIC06_P2) (Figure 1B). The effects of phages on bacterial
growth were overall consistent between liquid and solid assays: the phages with highest EOP
(i.e., producing as many plaques on NILS69 as on the production strain) cause the strongest
reduction of bacterial growth in liquid, whereas phages with low EOP do not affect bacterial
growth. The only exception is phage LF73_P4, which has little effect in liquid infection assay
despite having a relatively high EOP.
142
Next, time-resolved adsorption assays were performed for each of the 93 phages on NILS69
to test whether the host range defined by our initial screening was due to a defect in phage
adsorption. We confirmed that the 83 phages defined as non-infective by our screen failed to
bind to NILS69 (Supplementary Figure 2). Therefore, the susceptibility of NILS69 to these 93
phages appears to be exclusively determined by its surface composition. Previous genome
analysis indicated that NILS69 carries capsule loci while also exhibiting a rough LPS
phenotype. This combination of capsular structures and the absence of full-length O-antigen
may contribute to increased resistance to phage adsorption.
151
152 **Restriction Modification system and PD-T4-3 reduce the virulence of infecting phages**
153
We next wondered whether intracellular defences could impact the infectivity levels of the 10
phages that infect NILS69. NILS69 carries 6 defence systems as predicted by PADLOC (22)
and DefenseFinder(6) (Figure 1C and Supplementary Table 2), as well as 7 additional putative
systems predicted by PADLOC. Individual deletion mutants of PD-T4-3, PD-T7-1, PsyrTA,
SoFIC and Thoeris were constructed (Figure 1D). Regarding the predicted type II RM system,
computational tools identified a HNH enconuclease associated with two methyltransferases
(red arrows in Figure 1E). Manual inspection of the neighbouring region revealed two
additional HNH endonucleases (referred to as HNH endonuclease 1 and HNH endonuclease
3, yellow arrows in Figure 1E) encoded in direct vicinity of the type II RM, which were not
detected by prediction algorithms. A BLAST search against the REBASE database (23)
indicated that HNH endonuclease 1 and HNH endonuclease 3 are similar to two putative type
IV endonucleases (Osp6506McrB2P and ObaORFAP with e-values of 4E-40 and 7E-20
respectively), although these were predicted based on structural prediction and not verified
experimentally. As these two endonucleases might have an effect on phages, we knocked out
a large 10.6-kb region, that we refer to as RM island, encompassing the 3 HNH endonucleases
as well as the two methyltransferases. Out of the seven putative defence systems predicted
by PADLOC (22), we found PDC-SO4 and PDC-M63AB of particular interest as the first may
be similar to a Type II Toxin Antitoxin system (according to search against TADB3.0 (24)) and
the later consists In two genes that are encoded within a prophage,Ich are known to be
hotspots of defence systems (6,25–27). For these reasons, we included deletion mutants of
these 2 putative defence systems for downstream analyses.
The infectivities of the 10 phages were measured on each mutant through EOP (PFU on
mutant NILS69/ PFU on NILS69) and liquid assays (Figure 1D and Supplementary Figure 3).
Increased EOP were observed for all phages, except for NIC06_P2 and 536_P9, on mutants
lacking the RM island and PD-T4-3, indicating that these defence systems protect NILS69
against these phages, while the other systems do not seem to contribute. Liquid infection
assays yielded similar results, where the growth of the mutants lacking the RM island and PD-
T4-3 was effectively suppressed by phages. While 536_P9 did not formed more plaques on

any mutant in solid assays, this phage could effectively suppress the growth of the RM island
mutant in liquid, indicating that RM is also active against this phage (Supplementary Figure
3).
We next wondered if the phages that still induce weak lysis of NILS69 despite being restricted
by the RM system carry escape mutations or epigenetic modifications. To test this, we
collected phages by scraping from the lysis area formed at high phage concentration by
phages 536_P6 and 536_P7 growing on NILS69 and used them to reinfect both NILS69 and
the production strain 536. We predicted that, if the recovered phages were escaping NILS69
defences because of mutations or genome modifications, they would lyse NILS69 and 536
with similar efficacy. However, the recovered phages formed plaques on 536 but not on
NILS69, indicating that they were not carrying escape mutations or modifications
(Supplementary Figure 4), but were rather the product of a large phage population
overwhelming the defence capacity of NILS69.
The RM island and PD-T4-3 target a non-overlapping set of phages, with the RM island
restricting 7 phages (536_P1, 536_P6, 536_P7, 536_P9, AN24_P4, DIJ07_P1, LF73_P4) and
PD-T4-3 specifically inhibiting LF73_P1 and LF73_P3. To investigate potential negative or
positive interactions between defence systems, we generated four double mutants and four
triple mutants and tested their phage sensitivity using solid assays. Among these mutants,
only the double mutant lacking both PD-T4-3 and the RM island exhibited increased sensitivity
to phages, which was comparable to the additive effect observed in the corresponding single
mutants (Figure 1D). We used these mutants lacking multiple defence systems to screen the
AG phage collection again and confirmed that the lack of infectivity of the 83 phages is not
due to defence activity (Supplementary Figure 1A).
Altogether, our data indicate that the RM island and PD-T4-3 are the only active defence
systems in NILS69 under these conditions and against this collection of phages, and that their
combined activity provides an additive protection against a broader range of phages.
208
209

Experimental validation of the activity of an HNH endonuclease

Next, we aimed to identify which endonuclease within the RM island mediates phage
protection, hypothesizing that the type II RM system detected by DefenseFinder and PADLOC
(HNH endonuclease 2), which shows 100% identity to several restriction enzymes in REBASE,
is likely responsible for this phenotype. We constructed two mutants with deletions in different
portions of the RM island and measured phage EOP (Figure 1E). Only the mutants lacking
HNH endonuclease 3 (ΔRM_right) exhibited a phage sensitivity profile comparable to the
mutant lacking the entire RM island. These findings indicate that the type II endonuclease
detected by DefenseFinder and PADLOC is not responsible for phage protection. A sequence
homology search using BLASTP against the Standard database revealed that HNH
endonuclease 3 has 100% identity to an HNH endonuclease found in other *E. coli* isolates
(Uniprot: A0A3K0QCZ9). A structure-based homology search using AlphaFold3 (28) and
FoldSeek (29) identified a match to the Type IV methyl-directed restriction enzyme EcoKMcrA
encoded by *E. coli* K-12 MG1655 (E-value: 2.48E-04, run in Nov. 2024). Our data indicate that
the previously untested HNH endonuclease 3 has an antiphage activity and shares structural
features with a Type IV restriction enzyme (Figure 1E, Supplementary Figure 5).

Evaluation of the activity of NILS69 defence systems in the *E. coli* laboratory K-12

The apparent lack of activity of most of predicted defence systems might be linked to their target specificity. In other words, they might be active against other phages than those we tested. To explore this possibility, we attempted to cloned each individual defence system under its native upstream regulatory sequence into a low copy plasmid and subsequently transformed the constructs in *E. coli* K-12. Despite repeated attempts, we were unable to clone Thoeris and the RM island. The K-12 strain has a distinct phage sensitivity profile compared to NILS69, being sensitive to 15 of the 93 phages tested (Supplementary Figure 1A). Comparing the infectivity of the 15 phages on K-12 and on clones complemented with individual defence systems confirmed that PD-T4-3 is active and protects against 4 phages (CLB_P2, LF73_P1, NIC06_P2 and T4LD) which all belong to the *Tevenvirinae* subfamily (Figure 1F). Previous work indicated that PD-T4-3 targets phages from the *Tequatrovirus* genus (T2, T4 and T6)(18). Our data suggest that the specificity of PD-T4-3 is not limited to *Tequatrovirus* as it targets phage from *Dhakavirus* genus (CLB_P2) but PD-T4-3 likely does not target all genera from the *Tevenvirinae* subfamily since phages from *Mosigvirus* (LF82_P8) do not seem to be affected. Interestingly, phage NIC06_P2, which is unaffected by PD-T4-3 in NILS69, becomes sensitive to this defence system when overexpressed in K-12. This suggests that NIC06_P2 may possess a counter-defence strategy that is no longer effective under this condition of defence overexpression (Figure 1F). This assay also revealed that PD-T7-1 protects against the phage PDP351_P2 (*Teseptimavirus*) and has a partial protective activity as it reduces the plaque size of phage T7. None of the other defence system had a detectable protective activity in these conditions (Figure 1F). Surprisingly, the EOP of phages LM40_P1, LM40_P2 and LM40_P3 were higher when the strain expressed PsyrTA. Because we also noticed that this strain had a growth defect, we tested whehter the effect observed on LM40 phages was due to bacterial density. We show that phages LM40_P1, LM40_P2 and LM40_P3 have higher infectivity on low density bacterial lawns, and that the effect we see is not specific to PsyrTA (Supplementary Figure 6).

Restriction modification system of NILS69 limits plasmid transmission

We hypothesized that some defence systems in NILS69 may target other types of mobile genetic elements in addition to phages. Therefore, we assessed their capacity to limit plasmid transmission. We assessed the efficiency of transfer of a pRCS30, a 157kb IncC plasmid carrying multiple antibiotic resistance genes including an extended-spectrum β-lactamase gene CTX-M14(30), from its original host strain *E. coli* 513 to NILS69 and its derivative mutants.

The conjugation efficiency, measured as the ratio of transconjugants to recipients, was approximately 10^{-7} in NILS69, peaking at 1 hour. Similar efficiency was observed in all mutants except for the ΔRM strain, which exhibited a 100-fold increase (~10^{-5}), suggesting that the RM island restricts plasmid horizontal transmission. (Figure 2A). To test which endonuclease from the island was mediating this effect, we measured conjugation efficiency in the ΔRM_right and ΔRM_left mutant backgrounds and found that the same HNH endonuclease 3 was responsible for protection against both phages and plasmids (Figure 2B).

Because this endonuclease shares some similarity to a putative Type IV endonuclease, we aimed to verify if it was indeed targeting methylated invading DNA or not. To test this, we recovered transconjugants in NILS69, ΔRM_right and ΔRM_left backgrounds and measured the efficiency of a secondary transfer of pRCS30 in NILS69 and ΔRM island mutants. Plasmids from the donor strains NILS69 and ΔRM_left should be methylated, as these strains encode the two methyltransferases in the RM island, while plasmids from the donor ΔRM_right strain are expected to be unmethylated (Figure 1E). First, we observed that the transfer of pRCS30 into NILS69 is 1000-fold higher when the plasmid had previously replicated in the same background compared to its original host *E. coli* 513 (Figure 2C). Second, we confirmed that this increased conjugation efficiency is due to the presence of methyltransferases in the donor strains, as their absence prevents this increase in transfer efficiency (Figure 2C). Our data therefore indicate that NILS69 encodes an HNH endonuclease that restricts plasmid and phage infection and which activity is likely inhibited by methylation.

Finally, we tested whether plasmid vertical transmission could be affected by the two defence systems that are active in NILS69: PD-T4-3 and RM. To this aim we transformed the low copy plasmid pZS into NILS69 WT, ΔRM or ΔPD-T4-3. Plasmid-carrying bacterial clones were initially selected in the presence of antibiotics and subsequently transferred on non-selective media to measure the percentage of plasmid loss in each background (Figure 2D). While a fraction of the population lost the plasmid, this did not depend of the bacterial genetic background, therefore RM and PD-T4-3 had no effect on the vertical transmission of this plasmid.

Discussion

The role of defence systems in shaping phage-bacteria interactions is still unclear. In particular, the benefits provided by the accumulation of defence systems in a single genome remain an open question. In this study we used a direct genetic approach to investigate the impact of defence systems on the vulnerability of the *E. coli* clinical isolate NILS69 to a diverse collection of phages. Confirming a previous study, we experimentally show that the inability of phage to produce a successful infection is mainly explained by the lack of adsorption. Out of the six predicted defence systems, only two have a measurable impact on the infectivity of the phages that can bind the host; Restriction Modification and PD-T4-3. In addition, we tested two putative systems (i.e., never previously verified experimentally) predicted by PADLOC(22,31), namely PDC-M63AB and PDC-S04, but could not find evidence of defence activity in our model system. These findings raise questions about the activity of other predicted defence systems. We can foresee several reasons for the apparent inactivity of these systems. First, the limited diversity of the phage collection, which consists of virulent, double-stranded DNA phages, may introduce bias. Previous work has showed that certain defence systems are specific and target a narrow range of phages (13,18) and it may be that we have just not tested the phages that are sensitive to these defence systems.

Curiously, phage NIC06_P2 can be targeted by PD-T4-3 when expressed in *trans in E. coli* K-12, but is not targeted in the native context of NILS69.

Therefore, a second possibility is that some of these defence systems may be strongly regulated and expressed in different environmental conditions or may require additional cofactor to act. Although some regulators have been identified (11,32,33), whether defence systems are expressed and differentially regulated depending on the environment remains an unexplored area. Future studies aiming at measuring the impact of the environmental context

on defence systems activity and their consequences on phage activity will be key to advance our understanding on the ecological impact of defence systems.

A third explanation to the apparent inactivity of defence systems is given by phage-encoded counter-defence proteins. Many counter-defences have been identified recently in phages or plasmids and it clearly appears that many more remains to be uncovered (34,35). A recent effort was made to catalogue and detect all known counter-defence proteins from genomes (36). In this study, we observed that the phage NIC06_P2, which has the highest EOP on NILS69 despite an apparent low adsorption efficiency (Figure 1A, Supplementary Figure 2), is unaffected by the defence systems tested since it lyses NILS69 as efficiently as all the derivative mutants. Interestingly, this phage encodes a predicted anti-restriction nuclease (Arn) protein, which, in phage T4, inhibits the modification-dependent endonuclease McrBC (Supplementary Table 1). Phage LF73_P4 also encodes a predicted anti-RM protein, Dam, which may partially protect the phage against the RM island. In addition, NIC06_P2 is unaffected by the defence system PD-T4-3, while closely related phages are (LF73_P1 and LF73_P3). PD-T4-3 is a single gene system that encodes a GIY-YIG nuclease domain but its mechanism of action is unknown. Our data suggest that NIC06_P2 may escape PD-T4-3 in NILS69, but not in K-12 where this system is likely overexpressed (due to plasmid-based expression or to a lack of repression that might be at play in the native host).

Interrogating the effect of defence systems on other mobile genetic elements, we found that an HNH endonuclease effectively limits both phage and plasmid horizontal transmission. Interestingly, this endonuclease shares structural similarity to a Type IV methylcytosine-targeting restriction enzyme but our data indicate that the methyltransferase that are encoded directly downstream on the opposite strand limit the endonuclease activity, suggesting a Type I, II or III mode of action. This association of an HNH endonuclease with two methyltransferases might be conserved as our preliminary analysis detected this 3-gene association in *E. coli*, *Enterobacter bugandensis*, *Klebsiella pneumoniae* and *Vibrio alginolyticus*. Further molecular investigation of this endonuclease and associated methyltransferases will help to clarify their mechanism of action.

In conclusion, we conducted a comprehensive functional assessment of defence systems in a clinical *E. coli* isolate. Our study indicates that, for this strain and in the conditions tested, the main phage barrier is the bacterial envelope, while defence systems play a more marginal role and further highlights the need to study defence systems in their native hosts, in variable environmental conditions to improve our understanding of their roles in the ecology of phage-bacteria interactions.

Material and Method

Bacterial strains, phages, plasmids and growth conditiond

Bacterial strains, phages, plasmids and primers used in this study are listed in Supplementary Table 3. The medium used for all experiments was Lysogenic Broth (LB) prepared according to the Luria formulation. Unless otherwise specified, bacterial cultures were inoculated in LB from frozen stocks and propagated at 37°C with agitation. When needed, antibiotics were added at the following concentrations: ampicillin 100 μg/mL, kanamycin 100 μg/mL, chloramphenicol 30 μg/mL, streptomycin 30 μg/mL, and rifampicin 100 μg/mL. Solid or soft media were prepared by adding 15 g/L or 7 g/L of agar, respectively.

Phage efficiency scoring

Phage efficiency of plating (EOP) against NILS69 and K-12-MG1655 derivative was tested by spotting on soft agar plate 3µL of 10-fold serially diluted phage suspensions. Soft agar was inoculated to a final concentration of ≈10^{-7} CFU/mL from exponential phase growing cultures and then incubated overnight at 37°C. EOP was calculated by dividing the Plaque-Forming Units (PFU) or the highest dilution factor allowing the detection of phage lysis activity obtained on NILS69 by the number of PFU obtained on the phage production strain.. All the bacterial growth kinetic have been done in microtiter plate using "Tecan Infinite M Nano" microplate reader. Wells were inoculated with 10^5 CFU/mL from overnight culture with or without phages at a multiplicity of infection of 1. OD600nm readings were taken every 5 minutes over an 9h incubation period at 37°C with constant agitation

The screening of E. coli NILS69 against the Antonia Guelin phage collection was performed in microtiter plate using 200 µL culture inoculated with 10^5 CFU/mL from an overnight culture and phages at a 100x dilution from stocks. OD readings were taken every 5 minutes over a 9h incubation period at 37°C with constant agitation.

Adsorption assays

Adsorption assays on NILS69 were conducted according to a previously described protocol (37). Overnight bacterial cultures incubated at 37°C were transferred into a deep-well plate. Phages were introduced at MOI 0.001 in a final volume of 1mL, with each well representing a unique phage/bacteria combination. Adsorption was performed at 37°C for 30 minutes without agitation. To monitor adsorption kinetics, 150 µL samples were collected at 10, 20, and 30-minute intervals, treated with 50 µL of chloroform, and the supernatants were titrated using a spot assay. This included a negative control with phages only (no bacteria) and a positive control with phages and their reference bacterial strain, both of which were also titrated.

Deletion mutants of NILS69

Deletion mutants of the defence systems in NILS69 were constructed using site-specific recombination mediated by the **λ**-Red system, as described by Datsenko and Wanner (38). The primers listed in Supplementary Table 3 were used to amplify resistance cassettes via PCR, generating linear DNA fragments containing homology regions for recombination. This linear DNA was introduced into bacteria expressing the **λ**-Red system by electroporation, and recombinant bacteria were selected on the appropriate antibiotic-containing media. When necessary, the resistance marker was removed using FRT recombination mediated by the expression of the Flp recombinase, as described in the same study (38). All generated mutants were sequenced (Illumina, NextSeq) to confirm the deletion. The reference genome of NILS69 can be accessed on NCBI (accession: DABQWS000000000.1)

Subcloning of defence systems and expression in E. coli K-12 MG1655

The low-copy number plasmid pZS_Cat was used to subclone NILS69 defence systems and expressed them in the lab strain K-12 MG1655. Defence systems were amplified using the PCR primer listed in Supplementary Table 3, and were cloned in the pZS_Cat vector using commonly used restriction digestion and ligation. Constructions were verified by Sanger sequencing. The primers were designed in order to include a minimum of 200 nucleotide

409 upstream the start codon, in order to include any native promotor and regulatory sequences.
410 Only 5 out 8 defence systems could be cloned using this method.
411

Plasmid Vertical Transmission

The low-copy-number plasmid pZS_sfGFP was used to assess the plasmid vertical
transmission ability of NILS69 and its derivatives. This vector, derived from the pZ vector series
described by Lutz and Bujard (39) includes a low-copy-number replication origin, a kanamycin
resistance gene, and allows for the constitutive expression of superfolder-GFP (sfGFP) under
the control of an unregulated tetpromoter.
Vertical transmission of pZS_sfGFP was monitored as follows: overnight cultures were diluted
to approximately 1–10 bacteria in media containing kanamycin and incubated for 24 hours
(T1). The cultures were then diluted again to approximately 1–10 bacteria in media without
antibiotic selection and incubated for an additional 24 hours (T2). At T1 and T2, the colony-
forming units (CFU) were estimated by plating the cultures on solid media without selection.
To estimate the proportion of bacteria retaining the plasmid, ≈100 CFUs were replicated onto
plates containing kanamycin.

Plasmid Horizontal Acquisition

The IncC family conjugative plasmid pRCS30 was used to assess the ability of NILS69 and its
derivatives to acquire horizontally transferred DNA. This 157 kb plasmid, isolated from a
clinical *E. coli* strain, carries various antibiotic resistance genes, including aph(6)-Id, which
confers resistance to streptomycin. Since the spontaneous mutation frequency leading to
streptomycin resistance in NILS69 is below 10^{-9}, resistance to streptomycin was used as a
marker to track the transfer of the pRCS30 plasmid. To evaluate plasmid acquisition, recipient
NILS69 and its derivatives, grown to an OD_{600} of 0.2, were mixed at a 1:1 ratio with the
pRCS30 donor strain and incubated statically. All recipient cells carried the stable high-copy-
number plasmid pZS_sfGFP-kanR, which constitutively expressed GFP and conferred
resistance to kanamycin. After 1 hour and 24 hours, the mixed cultures were vortexed
vigorously for 1 minute, and bacteria were plated on LB agar containing kanamycin to
estimate the total count of recipient cells (kanR). Additionally, the cultures were plated on LB
agar containing both streptomycin and kanamycin to determine the number of recipient cells
that acquired the pRCS30 plasmid (kanR strepR). Conjugation efficiency was calculated as the
ratio of kanR strepR / kanR.
Conjugation assays across NILS derivatives were performed by first selecting spontaneous
mutants resistant to rifampicin (rifR), by plating the relevant strain on LB agar containing 100
µg/mL of rifampicin. These mutants were subsequently used as recipients during conjugation
assays with NILS69 derivatives carrying the pRCS30 plasmid. Here, the conjugation efficiency
was calculated as the ratio of rifR strepR / rifR.

Bioinformatic and data analysis tools

452 Defence systems from NILS69 genome were predicted using webservers of DefenseFinder
453 (6,40,41)   (https://defensefinder.mdmlab.fr/)   and   PADLOC   (22,31)

(https://padloc.otago.ac.nz/padloc/). Toxin Antitoxin systems were predicted with TADB 3.0 (https://bioinfo-mml.sjtu.edu.cn/TADB3/)

Search for sequence and structural homologies of the HNH endonucleases were done using BLAST(42) (https://blast.ncbi.nlm.nih.gov/Blast.cgi?PAGE=Proteins) against the Rebase database(23) (https://rebase.neb.com/rebase/rebase.html), AlphaFold3(28) (https://alphafold.ebi.ac.uk/) and FoldSeek(29) (https://search.foldseek.com/search).

Graphs and data analyses were generated with Prism (GraphPad).

Data accessibility

All raw data generated during these study have been deposited on https://github.com/MMSB-MEEP-lab

Acknowledgments

We acknowledge Dr Luce Landraud for proving the plasmid pRCS30 and Prof. Erick Denamur for sharing the strain NILS69. The Antonina Guelin phage collection and associated production strains were a kind gift from Dr. Laurent Debarbieux and Dr. Aude Bernheim. AG is supported by the following fundings ANR-21-CE35-0003, Emergence en recherche 2020 de l'Idex Université Paris Cité RM99J20IDXA8 and Emergence ville de Paris 2020-DAE78-EMERGENCE. AC is supported by the ATIP-Avenir program and IdEx Université Paris Cité ANR 18 IDEX 0001.

References

1. Bernheim A, Millman A, Ofir G, Meitav G, Avraham C, Shomar H, et al. Prokaryotic viperins produce diverse antiviral molecules. Nature. 16 sept 2020;1-5.

2. Kronheim S, Daniel-Ivad M, Duan Z, Hwang S, Wong AI, Mantel I, et al. A chemical defence against phage infection. Nature. déc 2018;564(7735):283-6.

3. Kever L, Hardy A, Luthe T, Hünnefeld M, Gätgens C, Milke L, et al. Aminoglycoside Antibiotics Inhibit Phage Infection by Blocking an Early Step of the Infection Cycle. mBio. 28 juin 2022;13(3):e0078322.

4. Rousset F, Sorek R. The evolutionary success of regulated cell death in bacterial immunity. Curr Opin Microbiol. août 2023;74:102312.

5. Georjon H, Bernheim A. The highly diverse antiphage defence systems of bacteria. Nat Rev Microbiol. oct 2023;21(10):686-700.

6. Tesson F, Hervé A, Mordret E, Touchon M, d'Humières C, Cury J, et al. Systematic and quantitative view of the antiviral arsenal of prokaryotes. Nat Commun. 10 mai 2022;13(1):2561.

7. Maestri A, Pons BJ, Pursey E, Chong CE, Gandon S, Custodio R, et al. The bacterial defense system MADS interacts with CRISPR-Cas to limit phage infection and escape. Cell Host Microbe. 14 août 2024;32(8):1412-1426.e11.

8. Wu Y, Garushyants SK, van den Hurk A, Aparicio-Maldonado C, Kushwaha SK, King CM, et al. Bacterial defense systems exhibit synergistic anti-phage activity. Cell Host Microbe. 10 avr 2024;32(4):557-572.e6.

9. Williams MC, Reker AE, Margolis SR, Liao J, Wiedmann M, Rojas ER, et al. Restriction endonuclease cleavage of phage DNA enables resuscitation from Cas13-induced bacterial dormancy. Nat Microbiol. mars 2023;8(3):400-9.

10. Birkholz N, Jackson SA, Fagerlund RD, Fineran PC. A mobile restriction-modification system provides phage defence and resolves an epigenetic conflict with an antagonistic endonuclease. Nucleic Acids Res. 8 avr 2022;50(6):3348-61.

11. Picton DM, Luyten YA, Morgan RD, Nelson A, Smith DL, Dryden DTF, et al. The phage defence island of a multidrug resistant plasmid uses both BREX and type IV restriction for complementary protection from viruses. 2021;18.

12. Dupuis MÈ, Villion M, Magadán AH, Moineau S. CRISPR-Cas and restriction-modification systems are compatible and increase phage resistance. Nat Commun. 2013;4:2087.

13. Costa AR, van den Berg DF, Esser JQ, Muralidharan A, van den Bossche H, Bonilla BE, et al. Accumulation of defense systems in phage-resistant strains of Pseudomonas aeruginosa. Sci Adv. 23 févr 2024;10(8):eadj0341.

14. Maguin P, Varble A, Modell JW, Marraffini LA. Cleavage of viral DNA by restriction endonucleases stimulates the type II CRISPR-Cas immune response. Mol Cell. 3 mars 2022;82(5):907-919.e7.

15. Srikant S, Guegler CK, Laub MT. The evolution of a counter-defense mechanism in a virus constrains its host range. eLife. 4 août 2022;11:e79549.

16. Doron S, Melamed S, Ofir G, Leavitt A, Lopatina A, Keren M, et al. Systematic discovery of antiphage defense systems in the microbial pangenome. Science [Internet]. 2 mars 2018 [cité 8 déc 2020];359(6379). Disponible sur: http://science.sciencemag.org/content/359/6379/eaar4120

17. Gao L, Altae-Tran H, Böhning F, Makarova KS, Segel M, Schmid-Burgk JL, et al. Diverse enzymatic activities mediate antiviral immunity in prokaryotes. Science. 28 août 2020;369(6507):1077-84.

18. Vassallo CN, Doering CR, Littlehale ML, Teodoro GIC, Laub MT. A functional selection reveals previously undetected anti-phage defence systems in the E. coli pangenome. Nat Microbiol. oct 2022;7(10):1568-79.

19. Gaborieau B, Vaysset H, Tesson F, Charachon I, Dib N, Bernier J, et al. Prediction of strain level phage-host interactions across the Escherichia genus using only genomic information. Nat Microbiol. nov 2024;9(11):2847-61.

20. Piel D, Bruto M, Labreuche Y, Blanquart F, Goudenège D, Barcia-Cruz R, et al. Phage-host coevolution in natural populations. Nat Microbiol. juill 2022;7(7):1075-86.

21. Bleibtreu A, Clermont O, Darlu P, Glodt J, Branger C, Picard B, et al. The *rpoS* Gene Is Predominantly Inactivated during Laboratory Storage and Undergoes Source-Sink Evolution in Escherichia coli Species. J Bacteriol. 15 déc 2014;196(24):4276-84.

22. Payne LJ, Todeschini TC, Wu Y, Perry BJ, Ronson CW, Fineran PC, et al. Identification and
classification of antiviral defence systems in bacteria and archaea with PADLOC reveals new
system types. Nucleic Acids Res. 8 nov 2021;49(19):10868-78.

23. Roberts RJ, Vincze T, Posfai J, Macelis D. REBASE--a database for DNA restriction and modification:
enzymes, genes and genomes. Nucleic Acids Res. janv 2010;38(Database issue):D234-236.

24. Guan J, Chen Y, Goh YX, Wang M, Tai C, Deng Z, et al. TADB 3.0: an updated database of bacterial
toxin–antitoxin loci and associated mobile genetic elements. Nucleic Acids Res. 5 janv
2024;52(D1):D784-90.

25. Rousset F, Depardieu F, Miele S, Dowding J, Laval AL, Lieberman E, et al. Phages and their
satellites encode hotspots of antiviral systems. Cell Host Microbe. 11 mai 2022;30(5):740-753.e5.

26. Hochhauser D, Millman A, Sorek R. The defense island repertoire of the Escherichia coli pan-
genome. PLoS Genet. avr 2023;19(4):e1010694.

27. Patel PH, Maxwell KL. Prophages provide a rich source of antiphage defense systems. Curr Opin
Microbiol. juin 2023;73:102321.

28. Abramson J, Adler J, Dunger J, Evans R, Green T, Pritzel A, et al. Accurate structure prediction of
biomolecular interactions with AlphaFold 3. Nature. 13 juin 2024;630(8016):493-500.

29. van Kempen M, Kim SS, Tumescheit C, Mirdita M, Lee J, Gilchrist CLM, et al. Fast and accurate
protein structure search with Foldseek. Nat Biotechnol. févr 2024;42(2):243-6.

30. Branger C, Ledda A, Billard-Pomares T, Doublet B, Fouteau S, Barbe V, et al. Extended-spectrum β-
lactamase-encoding genes are spreading on a wide range of Escherichia coli plasmids existing
prior to the use of third-generation cephalosporins. Microb Genomics. sept 2018;4(9):e000203.

31. Payne LJ, Meaden S, Mestre MR, Palmer C, Toro N, Fineran PC, et al. PADLOC: a web server for
the identification of antiviral defence systems in microbial genomes. Nucleic Acids Res. 5 juill
2022;50(W1):W541-50.

32. Gao LA, Wilkinson ME, Strecker J, Makarova KS, Macrae RK, Koonin EV, et al. Prokaryotic innate
immunity through pattern recognition of conserved viral proteins. Science. 12 août
2022;377(6607):eabm4096.

33. Picton DM, Harling-Lee JD, Duffner SJ, Went SC, Morgan RD, Hinton JCD, et al. A widespread
family of WYL-domain transcriptional regulators co-localizes with diverse phage defence systems
and islands. Nucleic Acids Res. 20 mai 2022;50(9):5191-207.

34. Mayo-Muñoz D, Pinilla-Redondo R, Camara-Wilpert S, Birkholz N, Fineran PC. Inhibitors of
bacterial immune systems: discovery, mechanisms and applications. Nat Rev Genet. avr
2024;25(4):237-54.

35. Samuel B, Mittelman K, Croitoru SY, Ben Haim M, Burstein D. Diverse anti-defence systems are
encoded in the leading region of plasmids. Nature. 7 nov 2024;635(8037):186-92.

36. Tesson F, Huiting E, Wei L, Ren J, Johnson M, Planel R, et al. Exploring the diversity of anti-defense
systems across prokaryotes, phages and mobile genetic elements. Nucleic Acids Res. 9 déc
2024;gkae1171.

37. Kropinski AM. Measurement of the rate of attachment of bacteriophage to cells. Methods Mol
Biol Clifton NJ. 2009;501:151-5.

38. Datsenko KA, Wanner BL. One-step inactivation of chromosomal genes in Escherichia coli K-12
using PCR products. Proc Natl Acad Sci U S A. 6 juin 2000;97(12):6640-5.

39. Lutz R. Independent and tight regulation of transcriptional units in Escherichia coli via the
LacR/O, the TetR/O and AraC/I1-I2 regulatory elements. Nucleic Acids Res. 15 mars
1997;25(6):1203-10.

40. Tesson F, Planel R, Egorov AA, Georjon H, Vaysset H, Brancotte B, et al. A Comprehensive
Resource for Exploring Antiphage Defense: DefenseFinder Webservice,Wiki and Databases. Peer
Community J. 25 sept 2024;4:e91.

41. Néron B, Denise R, Coluzzi C, Touchon M, Rocha EPC, Abby SS. MacSyFinder v2: Improved
modelling and search engine to identify molecular systems in genomes. Peer Community J. 24
mars 2023;3:e28.

42. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. J Mol Biol.
5 oct 1990;215(3):403-10.

586
587

590
591
Figure 1. Activity of defence systems encoded in NILS69 against ten phages
A,B. Phage infectivity on NILS69 measured through Efficiency of Plating (EOP) assays,
indicated as a ratio of the phage activity on NILS69 on the activity on phage production
strain.(A) or liquid infection assays using a multiplicity of infection (MOI) of 1 (B).

C. Genomic map of NILS69 indicating predicted defence systems (red), putative defence
systems assessed in this study (pink) and putative defence systems not tested in this study
(black). Location of prophages is indicated by yellow boxes.
D. Phage infectivity on NIL69 defence mutants measured through EOP assays
E. Schematic of the RM island locus in NILS69 and derivative mutants. Red arrows indicate
genes that were flagged as defence genes by DefenseFinder and PADLOC, yellow arrows
indicate genes that were identified as putative endonuclease by manual inspection and black
arrows indicate genes encoding hypothetical proteins (upper panel). Phage infectivity was
measured through EOP assays (PFU on mutant NILS69/PFU on WT NILS69) on indicated RM
mutants (lower panel).
F. Phage infectivity on K-12 expressing NILS69 defence systems in *trans* from pZS plasmids.
The EOP was calculated as (PFU on K-12 pZS_defence plasmid)/(PFU on K-12 pZS_empty
plasmid). Pictures of phage T7 PFU obtained on MG1655 pZS_empty or K-12 _pZS_PD-T7-1.
All experiments were performed in triplicate, except for the EOP in panel A which were
replicated 18 times. Boxes in panel A indicate minimal to maximal values, the line indicates
the mean value and individual data points are shown. Error bars in panel E indicate standard
error of mean.
613

614
615

Figure 2. NILS69 RM system limits plasmid horizontal transmission

A,B. Conjugation efficiency of the 157kb-plasmid pRCS30 into NILS69 and defence mutant derivatives. The plasmid was transferred from its native host (*E. coli* 513). Conjugation efficiency is indicated as the ratio of transconjugant Colony Forming Units (CFU) on recipient CFU after 1h or 24h of contact between the donor and the recipient strains.

C. Conjugation efficiency of pRCS30 from NILS69 to NILS69.

D. Vertical transmission of pZS_GFP plasmid in NILS69 and the ΔRM and ΔPD-T4-3 mutants. All experiments were perfomed in triplicates, except in Panel A, conjugation in NILS69 and ΔRM were performed six times. Individual data points are indicated, lines indicate mean values.

626
627
628

Electronic Supplementary material

A　　　　　　　　　　　　　　　　　**B**

Supplementary Figure 1. Screening of the Antonina Guelin phage collection on NILS69 and K-12 in solid (A) and liquid (B) infection assays.

"nt" stands for "not tested", the corresponding phages could not be amplified on their production strain. Green and orange squares indicate detectable phage infection (i.e.

635　plaques). Orange square indicate cases where only faint lysis could be observed. Grey
636　indicates no lysis

638
639
640 Supplementary Figure 2. Adorption assays of 93 phages from the Antonina Guelin on
641 NILS69 and on their control strains (phage production strains)
642

643
644
645
646   Supplementary Figure 3. Growth kinetic of NILS69 and mutant dertivative in the presence
647   of the infective phages at a MOI of 1
648
649

650
651
Supplementary Figure 4. Assessment of escape phage mutants
Phages 536_P6 and 536_P7 lysing NILS69 or NILS69ΔRM were harvested from the plate and
purified through chloroform extraction (Day 1). These phages were subjected to 10-fold serial
dilutions and spotted on lawn of strain 536, NILS69 or NILS69ΔRM (Day 2).

A   **B**

656
657
Supplementary Figure 5: Predicted structure of HNH endonuclease 3

A. HNH endonuclease 3 encoded in NILS69 has 100% identity with the amino acid sequence

of a HNH endonuclease (Uniprot A0A3K0QCZ9), which has a predicted structure in AlphaFold

(average pLDDT score of 80,56)

B. Structural alignment of NILS69_HNH endonuclease and K-12 MG1655_EcoKMrcA (e-value

2.48e-4; sequence identity 20.6%)

664
665

666
Supplementary Figure 6: Impact of bacterial lawn density on LM40_P phage infectivity in
the K-12 MG1655 strain expressing PsyrTA and its parental strain.
A, B. The bacterial lawn was inoculated directly from cultures at OD600 = 0.6 (1x).
C, D. The parental and PsyrTA strains were inoculated at dilutions of 1/10 or 10x, respectively,
from cultures at OD600 = 0.6. For each infectivity test, 3 µL of 10-fold serially diluted phage
was spotted. The T4 phage was used as a positive control for lysis.
673
674
675 **Supplementary Table 1.** List of phages infecting NILS69
676
677 **Supplementary Table 2.** List of predicted defence systems and toxin antitoxin systems
678 encoded in NILS69
679
680 **Supplementary Table 3.** Lists of plasmids, strains and primers used in this study