Title: Intergenerational transmission of parental rejection effects on survival,
behaviour, and miRNA expression in owl monkeys

Authors: Jayde Farinha[1], Mirtha Clausi[2] (Joint first authors), Fabian Orlando Chamba
Pardo[3], Nofre Sanchez-Perea[4], James Boot[5], Chaz Mein[5], Ping Yip[5], Ursula M. Paredes[1,5]*

[1]Department of Anthropology, University College London, London, UK WC1H 0BW
[2]Faculty of Veterinary Medicine, Universidad Nacional Mayor de San Marcos (UNMSM),
Lima, Peru
[3]Center for Technology Transfer, School of Veterinary Medicine, Universidad San Francisco
de Quito, Ecuador
[4]Center for Primate Conservation and Reproduction, Veterinary Institute of Tropical and
High Altitude Research (IVITA-Iquitos), Faculty of Veterinary Medicine, Universidad
Nacional Mayor de San Marcos (UNMSM), Lima, Peru
[5]Blizard Institute Centre for Neuroscience and Trauma, Queen Mary University of London,
London, UK E1 2AT

*Corresponding author: Ursula M. Paredes (ursula.paredes@ucl.ac.uk)

Keywords: Early-life adversity, intergenerational inheritance, epigenetics, microRNA,
primate behaviour, parental care, life history, *Aotus nancymaae*

Running title: Intergenerational effects of parental rejection in owl monkeys

Abstract

Parental rejection in captive primates represents severe early-life stress, yet intergenerational effects remain poorly characterised. We investigated spontaneous rejection in owl monkeys (*Aotus nancymaae*) across two generations using demographic records (n=1,372), behavioural observations (n=65), and blood miRNA profiling (n=47). Rejected individuals showed 51% reduced lifespan, decreased probability of reaching reproductive age, and increased morbidity throughout life. Critically, well-reared offspring of rejected parents exhibited similar reductions in survival to reproductive age and lifespan. Rejected adults displayed heightened fearfulness and aggression with reduced paternal care; offspring showed decreased locomotion and altered stress responses. Blood miRNA analysis revealed significant upregulation of mml-miR-30a-5p in female offspring-a miRNA associated with early-life stress and behavioural disorders in humans. These findings demonstrate intergenerational transmission of rejection effects on fitness and behaviour, with preliminary evidence suggesting epigenetic mechanisms are associated with this transmission, though multiple pathways likely operate.

Introduction

Epidemiological research in humans increasingly suggests that stress experienced by parents can be transmitted to offspring, affecting their health and behaviour across the lifespan[1-4]. However, studying parental stress and its intergenerational effects in humans is challenging due to long lifespans and complex social confounds; non-human primates offer valuable model systems due to their shared cognition, physiology, and extended parental investment with humans.

Studies in primates exposed to early-life adversity (ELA), including experimental parental separation, harsh parenting, or parental loss, demonstrate that these individuals exhibit lifelong alterations in stress behaviour and neural circuits[5-7], immune function[8], and health outcomes[9]. Critically, ELA is associated with decreased survival in early life and reduced lifespan in the wild[10,11] and in free-ranging populations[12].

Complete parental rejection, e.g. when mothers refuse to nurse newborns, represents the most severe form of ELA, as it can be lethal without intervention[13]. Rejection is a persistent problem in captive primate colonies globally, documented across the primate order[14-26]. Veterinarians typically intervene with hand-rearing or nursery-rearing to prevent infant mortality, and whilst rejected-then-reared primates are known to behave abnormally and show physiological differences[5,13], the long-term effects on health and life history remain poorly characterized outside a handful of species.

Emerging evidence suggests some effects of stressful rearing experience are heritable: offspring of ELA-exposed parents show altered behaviours and immune profiles[27-29] in captive macaques and reduced longevity and survival[30] in wild baboons. If rejection were to affect the health and behaviour of rejected and rescued captive primates, their descendants could also suffer. Studying intergenerational transmission of stress in captive populations can shed light on underlying mechanisms and ultimately inform strategies to improve captive primate welfare.

Candidate mechanisms for intergenerational transmission of stress, including maternal neglect, suggest alterations in epigenetic marks that become modified in response to environmental cues[31]. In captive primates, altered DNA methylation patterns in stress-regulatory genes are associated with ELA and predict health outcomes[32-34].

More recently, circulating microRNAs (miRNAs) have emerged as potential carriers of intergenerational stress. In *C. elegans*, small RNAs can be inherited across multiple generations and mediate transgenerational responses to environmental stress[35]. Altered miRNA expression has been detected in stressed mammals and their offspring, including in sperm[36] and blood plasma[37]. Multiple pathways may contribute to intergenerational transmission, including germline epigenetic inheritance, *in utero* programming through maternal physiology, or subtle alterations in parental behaviour that affect offspring development.

We hypothesized that if complete parental rejection is associated with behavioural effects detectable in the next generation, transmission might be accompanied by alterations in epigenetic marks such as miRNAs. We tested this in a captive population of owl monkeys (*Aotus nancymaae*)—a Neotropical primate that is uniquely monogamous, nocturnal, and displays intensive biparental care[38,39]. In this colony, spontaneous parental rejection occurs in approximately 5-10% of births. Mothers refuse nursing, occurring primarily within the first

days or months after birth, and occasionally paternal aggression accompanies maternal
rejection. Nursery-rearing is practiced to rescue rejected infants.

We previously reported poor health outcomes in rejected juveniles[41] and accelerated blood
DNA hypomethylation with age in this cohort[42], linking rejection history with long-term
stress, negative health outcomes, and epigenetic modifications. Here, we explored whether
intergenerational transmission of effects occurs and whether such transmission is
accompanied by behavioural and molecular alterations. We: (1) analysed demographic and
health records to compare lifespan and health outcomes of rejected individuals and their
offspring against controls; (2) characterized stress-reactive and affiliative behaviours across
two generations; and (3) profiled blood miRNA expression in rejected adults, their well-
reared offspring, and controls.

Results

Rejected owl monkeys show reduced lifespan and compromised fitness

We first examined whether parental rejection was associated with survival differences using
comprehensive demographic records (n=1,372 individuals: 135 rejected, 1,237 non-rejected).
Rejected individuals showed significantly reduced lifespan compared to controls (mean age
at death: 3.41 years [SD=4.58] vs. 7.03 years [SD=6.02]; Welch's t-test: p<0.001; Figure 1a).
This represents a 51% reduction in lifespan. Among individuals who survived to reproductive
age, there was no significant sex difference in lifespan for rejected males versus females
(9.05 years [SD=5.55] vs. 8.24 years [SD=5.37]).

We next assessed whether rejection affected the probability of reaching reproductive age
(defined as ≥1.5 years based on captive breeding data[38]). Among 1,161 individuals analysed,
60 of 152 rejected individuals (39.5%) reached reproductive age compared to 1,101 of 1,237
controls (89.0%). Rejected individuals had significantly lower odds of reaching reproductive
age (χ^2 test with Yates' correction: p<0.01; odds ratio=0.581; Figure 1b), representing a 41.9%
reduction in probability compared to controls.

Given our previous findings of elevated clinic visits in rejected juveniles[42], we examined
whether rejection history predicted health outcomes across the lifespan. Using clinic visits as
a proxy for morbidity and controlling for age and sex, we compared rejected and controls
visits to clinics (n=59 individuals with full clinic records n=3 female rejected, n=34 non
rejected, n=3 males rejected, n=19 non rejected). This preliminary analysis showed that
rejected individuals visited clinics 1.13 times more frequently throughout life, p=0.05 (linear
regression: β=1.13, SE=0.55, t=2.06, df=55, p=0.044, n=59; Figure 1c). Medical records
indicated these visits were primarily for treatment with broad-spectrum antibiotics,
suggesting greater susceptibility to infections, a leading cause of mortality in this colony[40].

Offspring of rejected parents show reduced survival despite normal care

We next asked whether parental rejection affected offspring fitness, even when offspring
themselves received normal parental care. Among 813 individuals with known parental
rejection history, offspring of rejected parents (n=31) showed significantly reduced odds of
reaching reproductive age compared to offspring of control parents (n=782; generalised
logistic regression: β=−0.545, SE=0.156, z=−3.49, p<0.001; odds ratio=0.553; Figure 1d).
This represents a 44.7% reduction in probability, similar to the effect of rejection on the

directly rejected individuals themselves. Effects were independent of parent sex (β=0.182,
SE=0.157, z=1.16, p=0.246) or offspring sex (β=−0.089, SE=0.157, z=−0.57, p=0.571).

Examining lifespan directly in offspring (n=727 controls, n=9 offspring of rejected), we
found offspring of rejected parents showed significantly reduced survival (Welch's t-test:
t=2.94, df=9.36, p=0.018; Figure 1d). Mean lifespan was 3.89 years (SD=4.68) for offspring
of rejected versus 8.50 years (SD=4.37) for controls, a difference of 4.61 years (95% CI:
1.01–8.21 years), representing a 54% reduction. These results demonstrate that rejection
effects on fitness are transmitted intergenerationally, with profound consequences for
descendants despite those descendants receiving normal parental care.

Since mechanisms underlying this transmission remain to be fully elucidated and likely
involve multiple pathways operating in concert, we explore possible contributors through
behavioural and molecular analyses below.

Rejected adults display altered stress reactivity and reduced parental care

The results of adjusted logistic regression with Bayesian estimation showed that behaviours
such as fear (+5%, p=0.03), hugging (+2%, p=0.01), approach (+20%, p=0.01) and hitting
others (+10%, p=0.01) increased in parents that were rejected in early life (n=11), whilst
hiding on the perch (-44%, p<0.01), carrying their offspring (-7%, p=0.04) and jumping
locomotion (-5%, p<0.01) decreased compared to control parents (n=27) (Supplementary File
8). When comparing behavioural differences adjusted for parent sex (Supplementary files
9,10), some behaviours were more evident in females. Mothers with a history of parental
rejection showed greater fear (+4%, p=0.01), hugging (+2%, p<0.01), approach (+19%,
p=0.07) and agonistic social behaviours such as hitting other monkeys (+11%, p=0.07) and
biting (+13%, p=0.30), whilst freezing (-10%, p=0.01), jumping locomotion (-5%, p=0.01)
and furtive glancing (-1%, p<0.01) decreased compared to controls. Similarly, fathers with
parental deprivation showed increased fear (+5%, p=0.70), approach (+28%, p=0.05) and
hitting others (+8%, p=0.07), whilst time spent hiding on the perch (-39%, p=0.01) and
hiding behind (-11%, p<0.01) decreased. They also showed decreased baby carrying (-8%,
p=0.03) and jumping locomotion (-5%, p<0.01) compared to controls (Figure 2). Only one
adult male monkey displayed stereotyped behaviour (head twirl).

Offspring of rejected parents showed more fearfulness and less locomotion similar to
their parents

Using the same statistical analysis, we show offspring of rejected parents (n=13) also display
behaviours such as fear (+18%, p=0.14) and hiding in the nest and hiding behind another
monkey (+12%, p=0.13; +9%, p=0.8 respectively) increased. However, significant results
were only detected amongst behaviours which were decreased for offspring of rejected
parents (n=13): hiding on the perch (-15%, p=0.05), jumping locomotion (-14%, p<0.01) and
biting (-12%, p=0.01). When considering offspring independently, the same pattern emerges.
Female offspring of parents with parental deprivation showed more fear (+18%, p=0.2),
hiding in the nest (+12%, p=0.3) and furtive glancing (+7%, p=0.06) but decreased jumping
locomotion (-14%, p<0.01) and biting (-11%, p=0.01) compared to female control offspring.
Additionally, male offspring showed increased fear (+10%, p=0.4) and decreased jumping
locomotion (-14%, p=0.2) and biting (-12%, p=0.3) compared to male control offspring
(Figure 2 and Supplementary File 11). Although some differences in behaviours are not

statistically significant, numerically they show larger average treatment effects across all
groups.

Female offspring of rejected parents show altered miRNA expression suggesting
potential molecular mechanisms

We conducted a differential expression analysis of whole blood miRNAs in animals with a
history of rejection, their offspring, and controls (n=47). This analysis was performed to
explore whether microRNAs might be associated with parental rejection history and
potentially contribute to intergenerational transmission of effects. Whilst this section focused
on commenting on significantly dysregulated miRNAs, we provide a complete list of all
analysed miRNAs, including those that showed no significant changes (Supplementary File
7). To identify significantly altered miRNAs across these multiple comparisons, we used
LIMMA TREND (Partek flow), which provides stringent false discovery rate control (FDR
$p<0.05$).

Our initial analysis demonstrated that, despite the association between adults with a history of
rejection and the dysregulation of many miRNAs in blood, none passed the minimum FDR
threshold in either rejected males (Figure 3a) or rejected females (Figure 3b). However, we
did observe a down-regulated miRNA in males and females (mml-miR-182) with marginal
significance (FDR p between 0.05 and 0.10). Subsequently, we compared miRNA in well-
reared offspring (F1) of parents who experienced rejection (n=8) with offspring of controls
(n=7). This comparison revealed several differentially regulated miRNAs. Again, a miRNA
showed common upregulation in female and male offspring (mml-miR-320a) but the effect
did not reach significance (FDR p between 0.05 and 0.10).

However, we did detect significant upregulation of one miRNA: mml-miR-30a-5p, which
was significantly overexpressed (FDR $p=9.91\times10^{-3}$, FC=2.09) in well-reared female
daughters (Figure 3 c,d). This miRNA was the only one to pass stringent multiple testing
correction across all comparisons. Among other upregulated miRNAs (p<0.05), we observed
the partner miRNA mml-miR-30a-3p. The upregulation of miR-30a-5p specifically in
daughters of rejected parents is notable given existing evidence linking this miRNA to stress-
related outcomes. miR-30a-5p belongs to a miRNA family implicated in mediating social
stress and behavioural disorders in mammals[43]. In humans, miR-30a-5p is upregulated in
blood plasma of individuals with depression who experienced early-life adversity and in
brains of rats exposed to early life stress[44].

Discussion

Parental rejection predicts reduced lifespan and compromised fitness. The 51% reduction in lifespan observed in rejected owl monkeys, despite veterinary care and identical post-weaning conditions as controls, demonstrates that early-life stress has profound, lasting effects on mortality even when environmental factors are controlled. Our finding that rejection remains a significant predictor of morbidity throughout life, with rejected individuals requiring 1.13 times more clinic visits, suggests chronic health vulnerability may contribute to reduced longevity.

The predominant reason for clinic visits was antibiotic treatment, suggesting increased susceptibility to infections. Given that pneumonia and enteritis are among leading causes of mortality in this colony[40], heightened infection risk may account for increased morbidity, paralleling observations in rhesus macaques[9]. Stress-induced immune dysregulation is well-documented in ELA research[8]. As we previously observed HPA axis alterations in rejected individuals, results from our present study could reflect compromised immune function.

High incidences of nephropathies and cardiomyopathies are documented in this colony[40,45,46] and are associated with stress exposure and sudden death in owl monkeys. Whether rejected individuals show higher prevalence of these chronic conditions remains to be tested. The exact origins of reduced survival in rejected owl monkeys warrant further investigation, but the combination of increased infection susceptibility and potential stress-related pathologies likely contributes to the observed mortality patterns.

Intergenerational transmission of reduced fitness despite normal parenting. The most striking finding is that offspring of rejected parents, despite receiving normal parental care themselves, showed nearly identical reductions in survival to reproductive age (44.7% reduction) and lifespan (54% reduction) as their rejected parents. This demonstrates genuine intergenerational transmission rather than simply persistent effects of poor rearing.

These results extend findings in wild baboons[30] to a controlled captive setting, demonstrating that intergenerational transmission occurs independent of environmental variation. All sources of life history variation typical in wild populations (resource availability, predation, disease exposure) are controlled in captivity. All primates live in identical conditions, and juveniles are moved to new enclosures by management staff at the age of weaning. The persistent survival deficit in offspring of rejected parents therefore cannot be attributed to external environmental factors or differences in parental investment such as resource competition observed in the wild.

Retrospective analysis of health outcomes in F1 offspring to explain earlier mortality was not possible, as medical records in this colony were created only when individuals visited clinics, and most F1 individuals recorded for our study died before our study began. This may indicate that negative health outcomes were subclinical and went undetected.

The persistence of reduced survival across generations, despite controlled environmental conditions and seemingly equal parental care, suggests that biological embedding of parental stress is transmitted. Our behavioural and molecular analyses below explore potential pathways underlying this transmission.

Behavioural transmission. altered stress reactivity, boldness, and parental care. Rejected adults showed a distinct behavioural profile characterised by heightened fearfulness yet paradoxically increased boldness (approach behaviour) and reduced hiding. This combination has been observed in rodents following maternal separation[47,48] and may reflect dysregulated threat assessment or risk-taking. Increased emotionality was also detected in rhesus macaques whose fathers experienced early-life separation[29]. Offspring of rejected parents showed

similar trends towards increased fearfulness and boldness, though these did not reach
statistical significance, potentially reflecting subtle intergenerational transmission detectable
with larger samples.

The finding that rejected fathers showed significantly reduced infant carrying is particularly
striking. Owl monkey fathers are renowned for intensive paternal care, carrying infants up to
90% of the time[18,38,39,49], a trait considered an evolutionary specialisation. Our results
demonstrate unexpected plasticity in this supposedly constrained behaviour, suggesting
rejection experience can fundamentally alter social bonding and caregiving motivation. The
origins of this caregiving deficit are unknown but could stem from disrupted attachment
formation during early development[50]. This parallels findings in rhesus macaques where
maternal rejection predicts abusive parenting in the next generation[27], though our study
extends effects to paternal care.

Offspring of rejected parents showed intergenerational transmission of reduced locomotion in
response to stress. Decreased mobility has been reported in ELA-exposed rhesus macaques[51]
and marmosets[52] and is thought to reflect motor system inhibition mediated by fronto-
amygdala circuits-a potential stress coping mechanism[53]. The consistency of this response
across generations suggests heritable alterations in stress-responsive neural circuits. Reduced
mobility in stress contexts may result from HPA axis overactivity[54].

Interestingly, biting behaviour showed opposite patterns across generations: rejected adults
showed increased biting, whilst their offspring showed decreased biting. As biting is part of
the maternal behavioural repertoire during normal weaning in owl monkeys[55], elevated biting
in rejected parents could contribute to perpetuating cycles of rejection, observed in other non-
human primates exposed to maternal neglect[27]. However, the suppression of this response in
studied offspring, despite their elevated emotionality, suggests adaptive regulation of
aggressive responses, potentially representing positive stress coping in well-reared
descendants.

Altered behaviours cannot be explained solely by learning or insufficient developmental
experience. Rejected parents can successfully rear some offspring whilst rejecting others,
indicating rejection is not deterministic. Moreover, offspring showed behavioural patterns
opposite to their parents (e.g., decreased biting), arguing against simple social learning.
Together, our findings indicate intergenerational transmission of behavioural plasticity
associated with rejection experience. This is ecologically relevant given that owl monkey
fathers provide extensive care both in captivity[18,38,40,55] and in the wild[39,49], and all owl
monkey mothers reject their babies to pass on caring duties to fathers during natural weaning.

Altered miRNA expression in offspring. preliminary evidence for molecular mechanisms.
The significant upregulation of mml-miR-30a-5p specifically in female offspring of rejected
parents provides preliminary evidence that miRNAs may contribute to intergenerational
stress transmission, though functional significance remains to be established. miR-30a-5p
belongs to a miRNA family implicated in mediating social stress and behavioural disorders in
mammals[43]. In adult humans with depression and early-life adversity history, miR-30a-5p is
upregulated in blood plasma[44], and the same study demonstrated altered DNA methylation
patterns at this locus in brains of rats exposed to ELA, demonstrating the association between
this circulatory biomarker and brain alterations in response to ELA. The functional
significance of miR-30a-5p overexpression in owl monkeys is unknown. If functional, miR-
30a-5p dysregulation could increase vulnerability to stress disorders in owl monkeys;
elevated expression in daughters of rejected parents could contribute to perpetuating
alterations in this population, particularly given females determine lactation, early offspring
care, and initiate weaning rejection. However, miR-30a-5p dysregulation could also indicate

other underlying physiological differences in blood between these cohorts. The very limited
number of altered miRNAs despite profound phenotypic differences in rejected individuals
and their offspring suggests multiple mechanisms are at play. At a genomic level, we have
also observed accelerated DNA hypomethylation in rejected owl monkeys[42]. Altogether, our
study in rejected owl monkeys supports the view that epigenome-wide effects are associated
with ELA[28,33] and that miRNAs are associated with transmission of altered phenotypes.

Limitations. Several important limitations should be acknowledged. Our study is
observational and correlational. Whilst the large demographic sample (n=1,372) provides
robust evidence that intergenerational effects occur, we cannot experimentally manipulate
parental rejection or control all potential confounding factors.

Could rejection itself have a heritable genetic component that also predisposes to poor health
outcomes? Several observations argue against this as the sole explanation: rejected
individuals successfully reared some offspring whilst rejecting others. Quality of the rejected
owl monkeys themselves is a possible cause, as proposed in rejection events in humans and
non-human primates[56,57], and could explain differences in naïve female offspring. However,
some rejected individuals, and their offspring enjoyed long healthy lives and reproduced
successfully. Anecdotal reports from veterinary inspections showed no apparent differences,
paralleling other observations of rejection in New World primates[58]. Sources of individual
variability such as genetic or chromosomal aberrations that might predispose individuals to
rejection or to suffer from intergenerational effects of rejection will be analysed in future
studies.

Further, we could not assess prenatal conditions, maternal health during pregnancy, or subtle
early postnatal experiences that might differ between rejected and control families. Whilst all
families lived in identical conditions post-weaning, earlier developmental differences could
contribute to observed effects. Some behavioural differences showed non-significant trends;
larger studies might clarify which changes are robust.

Future work should address several critical questions raised by our findings. First, whilst we
demonstrate that rejection harms offspring fitness, the evolutionary persistence of rejection
behaviour remains paradoxical. Investigating whether rejecting behaviour itself confers
fitness benefits to parents-such as extended lifespan or enhanced reproductive output-could
resolve this apparent contradiction and clarify whether rejection represents an adaptive
maternal strategy under captivity. Second, the molecular pathways underlying the negative
effects on rejected offspring warrant investigation. Third, characterising rejection behaviour
transmission itself-determining whether rejected individuals are more likely to reject their
own offspring and identifying associated molecular signatures-would establish whether
rejection propensity is epigenetically inherited alongside the physiological and behavioural
alterations we observed. Finally, extending molecular analyses to germline tissues,
conducting prospective longitudinal studies monitoring maternal physiology during
pregnancy, and examining F2 generation outcomes would clarify timing and mechanisms of
epigenetic transmission.

Broader evolutionary and biomedical significance. From an evolutionary perspective,
intergenerational transmission of stress effects may represent predictive adaptive responses,
where parental experiences inform offspring phenotypes in preparation for similar
environments[1,2]. However, whether the effects we observe represent adaptive calibration or
simply detrimental consequences of stress exposure remains unclear. In captivity,
environmental conditions differ radically from wild habitats, potentially rendering responses
that were once adaptive into maladaptive outcomes. The mixed pattern of behavioural
changes in offspring, including both potentially detrimental effects (reduced locomotion,

heightened fear) and possible adaptive stress coping (reduced aggression), suggests complex
interactions between inherited vulnerability and developmental plasticity rather than simple
pathology.

For biomedical research, our findings underscore that early-life history shapes lifelong health
trajectories and that parental history profoundly influences offspring outcomes even when
offspring themselves receive apparently normal care. This has implications for the
developmental origins of health and disease (DOHaD) framework[1], highlighting that
intergenerational effects may confound interpretation of results from primate biomedical
models where parental early-life history is often unknown or uncontrolled. Practical
implications. Our findings have important practical implications for captive primate
management. Complete parental rejection affects approximately 5-10% of births in this owl
monkey colony and has been documented in other owl monkey colonies[59] and across primate
taxa[13-26]. Current practice is mixed, but in some facilities, rejected infants are hand-reared and
then returned to breeding populations. Our results suggest this approach may inadvertently
propagate intergenerational effects, compromising welfare and fitness of subsequent
generations. Breeding programmes should consider enhanced monitoring of descendants of
rejected individuals for health issues, careful documentation of rejection history in pedigree
databases, and investigation of environmental or management factors that may reduce
rejection rates. These considerations are important for colonies producing animals for
biomedical research, where behavioural and physiological consistency is critical.

Conclusion. We identified robust intergenerational effects of parental rejection in owl
monkeys, manifesting as reduced survival and altered stress and social behaviours transmitted
across generations despite normal parental care of offspring. Blood miRNA analysis revealed
sex-specific dysregulation in female offspring, providing preliminary evidence that epigenetic
mechanisms are associated with transmission, though multiple pathways likely operate. The
central finding-that parental rejection effects are transmitted intergenerationally, reducing
offspring survival by 54% despite normal care-is robustly supported by large-scale
demographic data. The mechanisms underlying this transmission remain incompletely
understood and warrant further investigation. This study represents the first comprehensive
characterisation of intergenerational rejection effects in a captive non-traditional primate
model and has important implications for primate welfare, captive breeding practices, and
understanding intergenerational transmission of stress effects relevant to human health.

Materials and Methods

Ethics statement

All experimental procedures adhered to the American Society of Primatologists (ASP)
Principles for the Ethical Treatment of Non-Human Primates. The study was approved by the
Welfare and Research Committee of San Marcos University and the UNMSM ethics
committee. Research permits were issued by the Peruvian authority SERFOR to Dr. Paredes
(AUT-IFS-2021-040 and RDG NO D000334-2021-MIDAGRI-SERFOR-DGGSPFFS).

Study location and population

Research was conducted at the Centre for Reproduction and Conservation of Non-Human
Primates (CRCP) of IVITA at Universidad Nacional Mayor de San Marcos (UNMSM),
Iquitos, Peru. The colony has maintained detailed demographic and medical records since
establishment. Rejection events (defined as maternal cessation of lactation and/or paternal

aggression towards infants) are documented, and rejected infants are rescued via nursery-
rearing.

Housing and husbandry

423 Families were housed in single enclosures (2 m³: 1 m width × 1 m depth × 2 m height) with
424 natural photoperiod. Owl monkeys received daily feeding of seasonal fruits, balanced dry
425 food, and water ad libitum.

Sample composition

Sample sizes varied by analysis due to data availability (Supplementary File 1). For survival
analysis, we included 1,372 individuals (135 rejected, 1,237 control). Reproductive age
analysis included 1,161 individuals (60 rejected who reached reproductive age, 1,101
controls). Analysis of intergenerational effects on reproductive age included 813 individuals
(31 offspring of rejected parents, 782 offspring of controls). Intergenerational effects on
lifespan were analysed in 736 individuals (9 offspring of rejected, 727 controls). Behavioural
observations were conducted on 65 individuals across 10 families spanning 2 generations.
MicroRNA analysis initially included 48 samples, reduced to 47 after quality control, from
10 families across 2 generations selected for behavioural study.

Family cohort selection

For behavioural and molecular studies, we identified 10 families including at least one adult
with documented rejection history who survived to adulthood and had offspring at the time of
study. Age-matched control families with offspring were selected. The rejected group (P0)
comprised 11 rejected adults (6 females, 5 males), 8 well-reared breeding partners (3 females,
5 males), and 14 juvenile offspring (F1: 9 females, 5 males). The control group (P0) included
19 well-reared adults (9 females, 10 males) and 13 juvenile offspring (F1: 8 females, 5
males). Selection criteria ensured groups were matched for age, housing conditions, and
offspring number and age.

Demographic and health analyses

We compared age at death between rejected and control individuals using Welch's two-
sample t-test. For individuals alive at data collection, current age was used as censored data.
Mean age at "death or data collection" was calculated with standard deviations. Figures were
generated in R.

Based on captive breeding data[38], reproductive age was defined as ≥1.5 years. Pearson's chi-
square test with Yates' continuity correction assessed associations between rejection status
and probability of reaching reproductive age.

Generalised logistic regression examined whether parental rejection predicted offspring
reaching reproductive age, controlling for parent sex and offspring sex. Welch's two-sample
t-test compared lifespan between offspring of rejected parents and controls.

Linear regression assessed whether rejection history predicted clinic visits throughout life,
controlling for age and sex.

Behavioural characterisation

We conducted a modified human-intruder test[24] one month before blood sampling. Testing
occurred 1900-2000 hours when monkeys are most active[60]. Monkeys were exposed to low-
intensity stressors: light stimulus (camera light), human presence (unfamiliar observer,
"intruder"), and mechanical stimulus (cage door opening).

Three one-minute observation sessions were video-recorded per individual with five-minute
inter-session intervals. Behaviours were quantified using ethograms developed from previous
studies in *Aotus* (Supplementary File 2). Frequency and duration of 18 behaviours were
recorded, including stress responses (fear, freezing, hiding), locomotion, social behaviours
(grooming, carrying offspring), and agonistic behaviours (biting, hitting). Raw behavioural
percentages by sex and generation are provided in Supplementary File 3.

Behavioural events were observed using Behavioural Observation Research Interactive
Software (BORIS). Total duration and number of episodes per individual were quantified.
Behaviours were evaluated by two independent observers to avoid biased observations, with
inter-rater reliability ≥95%. Complete behavioural data are provided in Supplementary File 4.

Statistical analysis of behaviour

We used Bayesian logistic regression to estimate behavioural differences, adjusting for
family/cage effects, sex, and age category. Parental rejection history was the main predictor.
The outcome was binary (0/1) indicating whether an individual displayed each specific
behaviour at least once during observation periods (0 = no behaviour observed, 1 = at least
one event observed).

Predicted/adjusted frequencies were compared using exact Wilcoxon rank-sum tests. Average
treatment effects were estimated using G-computation, assigning all individuals to each
treatment condition and estimating differences between predicted posterior distributions.
Statistical models and priors are described in Supplementary File 5. Results with $p<0.05$ were
considered significant.

Blood collection

Blood samples were collected from a subset of 48 owl monkeys (rejected P0: 8 females, 4
males; offspring of rejected F1: 7 females, 5 males; control P0: 10 females, 8 males;
offspring of controls F1: 7 females, 2 males) during routine veterinary procedures under
ketamine sedation (8 mg/kg). Samples were obtained via femoral venipuncture using 3 ml
syringes with 1"×23G needles into EDTA Vacutainer tubes. From each sample, 0.5 ml was
spotted onto filter paper and air-dried at room temperature.

MicroRNA extraction and sequencing

Total RNA was extracted from three 6-mm punches per dried blood spot using the mirVana
microRNA Isolation Kit (Thermo Fisher Scientific, AM1560) following manufacturer's
protocols[61,62]. Briefly, spots were suspended in cell disruption buffer, denaturing solution was
added, followed by acid phenol:chloroform extraction. After centrifugation at 4°C, the
aqueous phase was collected, mixed with 100% ethanol, and applied to filter cartridges. After
washing, RNA was eluted in pre-heated elution buffer.

Total RNA was concentrated using the RNeasy MinElute Cleanup Kit (Qiagen, 74204). Lysis
buffer containing guanidine thiocyanate was added to RNA samples, followed by ethanol.
The mixture was applied to RNeasy MinElute spin columns, washed, and high-quality
concentrated RNA was eluted in 12 µl RNase-free water and stored at −80°C.

Libraries were prepared using QIAseq miRNA Library Kit (Qiagen) according to
manufacturer's instructions. Library quality was assessed using NanoDrop 8000 (Thermo)
and TapeStation 4200 (Agilent) before sequencing on NextSeq 2000 (Illumina Inc.).

Bioinformatic preprocessing and differential expression analysis

Raw sequencing reads were trimmed using Trim Galore v0.6.6 to remove adapters and low-
quality bases. Trimmed reads were aligned to the owl monkey genome (*Aotus nancymaae*,
Anan_2.0) using STAR v2.7.3a. Gene-level counts were generated using HTSeq-count.

Raw counts were loaded into Partek Flow (https://www.partek.com/partek-flow/) for
analysis. Counts were normalised (counts per million, CPM), filtered to remove unexpressed
genes, and principal component analysis (PCA) was performed to evaluate sample variability
and detect outliers. One sample (T-1499, P0 female control) was excluded as an outlier based
on PCA (Supplementary File 6). After filtering, approximately 3,000 sequences matched the
owl monkey genome.

Since the *Aotus nancymaae* genome is not fully annotated for miRNAs, we mapped owl
monkey small non-coding RNA sequences to the rhesus macaque (*Macaca mulatta*) genome
(Mmul_10, GCA_003339765.3) to identify miRNA homologues.

Normalised expression ratios were analysed in Partek Flow. We used the LIMMA TREND
method to detect differentially expressed miRNAs between rejected adults versus control
adults (stratified by sex) and offspring of rejected parents versus offspring of controls
(stratified by sex). Multiple testing correction was performed using the Benjamini-Hochberg
procedure to control false discovery rate (FDR). Significance thresholds were set at FDR
p<0.05 and fold change ≥1.5 or ≤−1.5. All reported p-values are FDR-adjusted unless stated
otherwise. Complete miRNA expression data including non-significant results are provided
in Supplementary File 7.

For heatmap visualisation, miRNA expression data were log□-transformed and z-scores were
calculated by subtracting the mean expression of each miRNA across all samples and
dividing by the standard deviation, allowing standardised comparison across multiple
miRNAs.

Statistical software and reproducibility

531 All statistical analyses were conducted using R version 4.1.0 for demographic and survival
532 analyses, RStan for Bayesian behavioural analyses, Partek Flow version 10.0 for miRNA
533 differential expression, and BORIS version 7.9.18 for behavioural coding.

Power and sample size considerations

Sample sizes were determined by availability of individuals meeting inclusion criteria in this
colony rather than prospective power calculations. For demographic analyses, large sample

sizes (n=1,389) provided excellent power (>95%) to detect moderate effect sizes (Cohen's d
≥0.3). For behavioural and miRNA analyses, smaller sample sizes (n=47-65) limited power
to detect small effects but provided adequate power (80%) to detect moderate-to-large effects
(Cohen's d ≥0.7 for behaviour; fold change ≥1.5 for miRNA with FDR correction).

Blinding

Behavioural observers were not fully blinded to rejection status, as this information was part
of colony records. However, observers were not informed of specific study hypotheses
regarding which behaviours were expected to differ, and coding was performed by two
independent observers to minimise bias. For miRNA analysis, laboratory personnel
processing samples and bioinformaticians conducting analyses were blinded to sample
identity and rejection status.

Data availability

All data supporting the findings of this study are available within the manuscript and
supplementary files. Raw miRNA sequencing data will be deposited in NCBI Gene
Expression Omnibus (GEO) upon acceptance. Summary of biological sampling across
analysis (de-identified) is provided in Supplementary File 1. Behavioural data are provided in
Supplementary Files 2-4, 8-11. miRNA expression data are provided in Supplementary File
7. Analysis code for demographic and behavioural analyses is provided in Supplementary
File 5. Additional data supporting the findings are available from the corresponding author
upon reasonable request, subject to ethical restrictions to protect animal identities and colony
security.

Author Contributions

Conceptualization: UMP, JF, MC
Data curation: NSP, JF, MC, UMP
Formal analysis: JF, MC, JB, PY, UMP
Funding acquisition: UMP, NSP, PY
Investigation: JF, MC, FOCP
Methodology: JF, MC, JB, CM, PY, UMP
Project administration: UMP, NSP
Resources: UMP, NSP, PY
Software: JF, FOCP, JB
Supervision: UMP, NSP
Validation: JF, MC, JB, PY
Visualization: JF, MC, UMP
Writing - original draft: JF, MC, FOCP, UMP
Writing - review & editing: All authors

Competing Interests

574 The authors declare no competing interests.

Funding

This work was supported by Queen Mary University of London Neuroscience Pilot Funds to
UMP and Universidad Nacional Mayor de San Marcos research project funds to NSP. The
funders had no role in study design, data collection and interpretation, or the decision to
submit the work for publication.

Acknowledgments

We thank the primate care staff and laboratory scientists at IVITA-Iquitos led by Nofre
Sanchez, for assisting with data and sample collection. We are grateful to staff at the Blizard
Institute Genome centre for assistance with molecular analyses. We thank Prof. Vardhman
Rakyan for helpful discussions.

References

1. Gluckman PD, Hanson MA, Beedle AS. Early life events and their consequences for
later disease: a life history and evolutionary perspective. American journal of human
biology. 2007 Jan;19(1):1-9.
2. Pembrey M, Saffery R, Bygren LO. Human transgenerational responses to early-life
experience: potential impact on development, health and biomedical research. Journal
of medical genetics. 2014 Sep 1;51(9):563-72.
3. Bowers ME, Yehuda R. Intergenerational transmission of stress in humans.
Neuropsychopharmacology. 2016 Jan;41(1):232-44.
4. Painter RC, Osmond C, Gluckman P, Hanson M, Phillips DI, Roseboom TJ.
Transgenerational effects of prenatal exposure to the Dutch famine on neonatal
adiposity and health in later life. BJOG: An International Journal of Obstetrics &
Gynaecology. 2008 Sep;115(10):1243-9.
5. Sanchez MM. The impact of early adverse care on HPA axis development: nonhuman
primate models. Hormones and behavior. 2006 Nov 1;50(4):623-31.
6. Lyons DM, Parker KJ. Stress inoculation‐induced indications of resilience in
monkeys. Journal of Traumatic Stress: Official Publication of The International
Society for Traumatic Stress Studies. 2007 Aug;20(4):423-33.
7. Spinelli S, Chefer S, Suomi SJ, Higley JD, Barr CS, Stein E. Early-life stress induces
long-term morphologic changes in primate brain. Archives of general psychiatry.
2009 Jun 1;66(6):658-65.
8. Kinnally EL, Martinez SJ, Chun K, Capitanio JP, Ceniceros LC. Early social stress
promotes inflammation and disease risk in rhesus monkeys. Scientific reports. 2019
May 20;9(1):7609.
9. Conti G, Hansman C, Heckman JJ, Novak MF, Ruggiero A, Suomi SJ. Primate
evidence on the late health effects of early-life adversity. Proceedings of the National
Academy of Sciences. 2012 Jun 5;109(23):8866-71.
10. Morrison RE, Eckardt W, Stoinski TS, Rosenbaum S. Cumulative early-life adversity
does not predict reduced adult longevity in wild gorillas. Current Biology. 2023 Jun
5;33(11):2307-14.
11. Tung J, Archie EA, Altmann J, Alberts SC. Cumulative early life adversity predicts
longevity in wild baboons. Nature communications. 2016 Apr 19;7(1):11181.
12. Patterson SK, Andonov E, Arre AM, Martínez MI, Negron-Del Valle JE, Petersen
RM, Phillips D, Rahman A, Ruiz-Lambides A, Villanueva I, Lea AJ. Early life
adversity has sex-dependent effects on survival across the lifespan in rhesus
macaques. Philosophical Transactions B. 2024 Oct 28;379(1916):20220456.

13. Sackett GP, Ruppenthal G, Elias K, editors. Nursery rearing of nonhuman primates in
the 21st century. Springer Science & Business Media; 2010 May 10.
14. Brockman DK, Willis MS, Karesh WB. Zoo. II. Reproduction, Pregnancy,
Parturition, Litter Size, Infant Care, and Reintroduction of Hand-Raised Infants. Zoo
Biology. 1987;6:349-63.
15. Debyser IW. Platyrrhine juvenile mortality in captivity and in the wild. International
journal of primatology. 1995 Dec;16:909-33.
16. Johnson LD, Petto AJ, Sehgal PK. Factors in the rejection and survival of captive
cotton top tamarins (Saguinus oedipus). American Journal of Primatology.
1991;25(2):91-102.
17. Bardi M, Petto AJ, Lee Parritz DE. Parental failure in captive cotton top tamarins
(Saguinus Oedipus). American Journal of Primatology: Official Journal of the
American Society of Primatologists. 2001 Jul;54(3):159-69.
18. Gozalo A, Montoya E. Reproduction of the owl monkey (Aotus nancymai)(primates:
Cebidae) in captivity. American journal of primatology. 1990;21(1):61-8.
19. Meireles BC, Goldschmidt B, Resende FC, Lopes CA, Nascimento LW. Neotropical
primate nursery in a squirrel monkey breeding unit in Brazil. Laboratory animals.
2016 Feb;50(1):67-72.
20. Mulholland MM, Williams LE, Abee CR. Rearing condition may alter neonatal
development of captive Bolivian squirrel monkeys (Saimiri boliviensis boliviensis).
Developmental psychobiology. 2020 Nov;62(7):909-19.
21. Fontenot BM, Galentine DM, Cummings LJ. Fostering procedures for orphaned
infants in a breeding colony of African green monkeys. Journal of the American
Association for Laboratory Animal Science. 2004 Sep 15;43(5):41-3.
22. Maestripieri D, Carroll KA. Causes and consequences of infant abuse and neglect in
monkeys. Aggression and Violent Behavior. 2000 May 1;5(3):245-54.
23. Brent L, Bode A. Baboon Nursery Rearing Practices and Comparisons between
Nursery-Reared and Mother-Reared Individuals. In: Sackett GP, Ruppenthal G, Elias
K, editors. Nursery rearing of nonhuman primates in the 21st century. New York:
Springer Science & Business Media; 2010. p. 275.
24. Capitanio JP, Mason WA, Mendoza SP, DelRosso L, Roberts JA. Chapter 11. In:
Sackett GP, Ruppenthal G, Elias K, editors. Nursery rearing of nonhuman primates in
the 21st century. New York: Springer Science & Business Media; 2010. p. 191.
25. Cocks L. Factors affecting mortality, fertility, and well-being in relation to species
differences in captive orangutans. International Journal of Primatology. 2007
Apr;28(2):421-8.
26. Brent L, Williams-Blangero S, Stone AM. Evaluation of the chimpanzee breeding
program at the Southwest Foundation for Biomedical Research. Laboratory animal
science. 1996 Aug 1;46(4):405-9.
27. Maestripieri D. Early experience affects the intergenerational transmission of infant
abuse in rhesus monkeys. Proceedings of the National Academy of Sciences. 2005 Jul
5;102(27):9726-9.
28. Sproul Bassett AM, Wood EK, Lindell SG, Schwandt ML, Barr CS, Suomi SJ, Higley
JD. Intergenerational effects of mother's early rearing experience on offspring
treatment and socioemotional development. Developmental psychobiology. 2020
Nov;62(7):920-31.
29. Kinnally EL, Capitanio JP. Paternal early experiences influence infant development
through non-social mechanisms in Rhesus Macaques. Frontiers in Zoology. 2015
Dec;12:1-8.

30. Zipple MN, Archie EA, Tung J, Altmann J, Alberts SC. Intergenerational effects of early adversity on survival in wild baboons. Elife. 2019 Sep 24;8:e47433.
31. Champagne FA, Curley JP. Epigenetic mechanisms mediating the long-term effects of maternal care on development. Neuroscience & Biobehavioral Reviews. 2009 Apr 1;33(4):593-600.
32. Massart R, Suderman M, Provencal N, Yi C, Bennett AJ, Suomi S, Szyf M. Hydroxymethylation and DNA methylation profiles in the prefrontal cortex of the non-human primate rhesus macaque and the impact of maternal deprivation on hydroxymethylation. Neuroscience. 2014 May 30;268:139-48.
33. Provençal N, Suderman MJ, Guillemin C, Massart R, Ruggiero A, Wang D, Bennett AJ, Pierre PJ, Friedman DP, Côté SM, Hallett M. The signature of maternal rearing in the methylome in rhesus macaque prefrontal cortex and T cells. Journal of Neuroscience. 2012 Oct 31;32(44):15626-42.
34. Kinnally EL. Epigenetic plasticity following early stress predicts long-term health outcomes in rhesus macaques. American journal of physical anthropology. 2014 Oct;155(2):192-9.
35. Rechavi O, Houri-Ze'evi L, Anava S, Goh WS, Kerk SY, Hannon GJ, Hobert O. Starvation-induced transgenerational inheritance of small RNAs in C. elegans. Cell. 2014 Jul 17;158(2):277-87.
36. Gapp K, Jawaid A, Sarkies P, Bohacek J, Pelczar P, Prados J, Farinelli L, Miska E, Mansuy IM. Implication of sperm RNAs in transgenerational inheritance of the effects of early trauma in mice. Nature neuroscience. 2014 May;17(5):667-9.
37. Jawaid A, Roszkowski M, Mansuy IM. Transgenerational epigenetics of traumatic stress. Progress in molecular biology and translational science. 2018 Jan 1;158:273-98.
38. Dixson AF. The owl monkey (Aotus trivirgatus). Reproduction in new world primates: New models in medical science. 1983:69-113.
39. Fernandez-Duque E, Valeggia CR, Mendoza SP. The biology of paternal care in human and nonhuman primates. Annual review of Anthropology. 2009 Oct 21;38(1):115-30.
40. Sánchez N, Galvéz H, Montoya E, Gozalo A. Mortalidad en crías de Aotus sp.(Primates: Cebidae) en cautiverio: una limitante para estudios biomédicos con modelos animales. Revista peruana de medicina experimental y salud pública. 2006 Jul;23(3):221-4.
41. Osman M, Olkun A, Maldonado AM, Lopez‐Tremoleda J, Sanchez‐Perea N, Paredes UM. Parentally deprived juvenile Owl monkeys suffer from long‐term high infection rates but not from altered hair cortisol concentrations nor from stereotypic behaviours. Journal of Medical Primatology. 2021 Dec;50(6):306-12.
42. Girling MT, Sanchez Perea N, Paredes U. Global DNA Hypomethylation as a Biomarker of Accelerated Epigenetic Ageing in Primates. bioRxiv. 2024:2024-03.
43. Khandelwal N, Dey SK, Chakravarty S, Kumar A. miR-30 family miRNAs mediate the effect of chronic social defeat stress on hippocampal neurogenesis in mouse depression model. Frontiers in molecular neuroscience. 2019 Aug 8;12:188.
44. Cattaneo A, Suderman M, Cattane N, Mazzelli M, Begni V, Maj C, D'Aprile I, Pariante CM, Luoni A, Berry A, Wurst K. Long-term effects of stress early in life on microRNA-30a and its network: Preventive effects of lurasidone and potential implications for depression vulnerability. Neurobiology of Stress. 2020 Nov 1;13:100271.

45. Knowlen GG, Weller RE, Perry RL, Baer JF, Gozalo AS. Hypertrophic cardiomyopathy in owl monkeys (Aotus spp.). Comparative Medicine. 2013 Jun 1;63(3):279-87.

46. Gozalo A, Dagle GE, Montoya E, Weiler RE, Malaga CA. Spontaneous cardiomyopathy and nephropathy in the owl monkey (Aotus sp.) in captivity. Journal of medical primatology. 1992 Jul;21(5):279-84.

47. van Steenwyk G, Roszkowski M, Manuella F, Franklin TB, Mansuy IM. Transgenerational inheritance of behavioral and metabolic effects of paternal exposure to traumatic stress in early postnatal life: evidence in the 4th generation. Environmental epigenetics. 2018 Apr;4(2):dvy023.

48. Caldji C, Francis D, Sharma S, Plotsky PM, Meaney MJ. The effects of early rearing environment on the development of GABAA and central benzodiazepine receptor levels and novelty-induced fearfulness in the rat. Neuropsychopharmacology. 2000 Mar 1;22(3):219-29.

49. Rotundo M, Fernandez-Duque E, Dixson AF. Infant development and parental care in free-ranging Aotus azarai azarai in Argentina. International Journal of Primatology. 2005 Dec;26:1459-73.

50. McCormack K, Sanchez MM, Bardi M, Maestripieri D. Maternal care patterns and behavioral development of rhesus macaque abused infants in the first 6 months of life. Developmental Psychobiology: The Journal of the International Society for Developmental Psychobiology. 2006 Nov;48(7):537-50.

51. Feng X, Wang L, Yang S, Qin D, Wang J, Li C, Lv L, Ma Y, Hu X. Maternal separation produces lasting changes in cortisol and behavior in rhesus monkeys. Proceedings of the National Academy of Sciences. 2011 Aug 23;108(34):14312-7.

52. Dettling AC, Feldon J, Pryce CR. Early deprivation and behavioral and physiological responses to social separation/novelty in the marmoset. Pharmacol Biochem Behav. 2002;73:259–269.

53. Roelofs K. Freeze for action: neurobiological mechanisms in animal and human freezing. Philosophical Transactions of the Royal Society B: Biological Sciences. 2017 Apr 19;372(1718):20160206.

54. Kalin NH, Shelton SE, Rickman M, Davidson RJ. Individual differences in freezing and cortisol in infant and mother rhesus monkeys. Behavioral Neuroscience. 1998 Feb;112(1):251.

55. Dixson AF, Fleming D. Parental behaviour and infant development in owl monkeys (Aotus trivirgatus griseimembra). Journal of Zoology. 1981 May;194(1):25-39.

56. Weiss M. Parents' rejection of their appearance-impaired newborns: Some critical observations regarding the social myth of bonding. Marriage & family review. 1998 May 27;27(3-4):191-209.

57. Leroux M, Monday G, Chandia B, Akankwasa JW, Zuberbühler K, Hobaiter C, Crockford C, Townsend SW, Asiimwe C, Fedurek P. First observation of a chimpanzee with albinism in the wild: Social interactions and subsequent infanticide. American journal of primatology. 2022 Jun;84(6):e23305.

58. Culot L, Lledo-Ferrer Y, Hoelscher O, Muñoz Lazo FJ, Huynen MC, Heymann EW. Reproductive failure, possible maternal infanticide, and cannibalism in wild moustached tamarins, Saguinus mystax. Primates. 2011 Apr;52:179-86.

59. Obaldia III N, Otero W, Marin C, Aparicio J, Cisneros G. Long□term effect of a simple nest□box on the reproductive efficiency and other life traits of an Aotus lemurinus lemurinus monkey colony: an animal model for malaria research. Journal of medical primatology. 2011 Dec;40(6):383-91.

60. Aquino R, Encarnación F. Characteristics and use of sleeping sites in Aotus (Cebidae:
Primates) in the Amazon lowlands of Peru. American Journal of Primatology.
1986;11(4):319-31.
61. Ponnusamy V, Kapellou O, Yip E, Evanson J, Wong LF, Michael-Titus A, Yip PK,
Shah DK. A study of microRNAs from dried blood spots in newborns after perinatal
asphyxia: a simple and feasible biosampling method. Pediatric research. 2016
May;79(5):799-805.
62. Ponnusamy V, Ip RT, Mohamed MA, Clarke P, Wozniak E, Mein C, Schwendimann
L, Barlas A, Chisholm P, Chakkarapani E, Michael-Titus AT. Neuronal let-7b-5p acts
through the Hippo-YAP pathway in neonatal encephalopathy. Communications
Biology. 2021 Sep 30;4(1):1143.

Figure 1. Reduced life expectancy and health in rejected owl monkeys and descendants. (a) Age at death was significantly lower for rejected compared to control monkeys (P0). Rejected individuals (n=135) showed mean age at death of 3.41 years (SD=4.58) compared to 7.03 years (SD=6.02) for controls (n=1,237), representing a 51% reduction in lifespan (Welch's t-test, p<0.001). **(b)** Probability of rejected owl monkeys reaching reproductive age (≥1.5 years) was significantly lower than controls (P0). Rejected individuals showed 41.9% reduced odds of reaching reproductive age compared to controls (χ^2 test with Yates' correction, p<0.01; n=1,161 total: 60 rejected reached reproductive age, 1,101 controls). **(c)** Clinic visits controlling for age. Rejected individuals (n=6) averaged 1.13 more clinic visits throughout life than controls (n= 53), linear regression, p=0.05. Males and females shown separately, with no significant sex-related differences. **(d)** Age at death for offspring of rejected parents (n=9) versus controls (n=727), showing 54% reduction in lifespan (mean 3.89 years vs 8.50 years; Welch's t-test, p=0.018). Males and females shown separately, with no significant sex-related differences. Error bars represent standard deviations. Blue=controls, red=rejected or offspring of rejected.

Figure 2. Behavioural differences in rejected parents and offspring compared to controls.
Forest plots showing behavioural differences (adjusted percentages controlling for family, sex, and age using Bayesian logistic regression) between rejected and control groups. Upper panels: fathers (n=20 total) and mothers (n=18 total) (P0). Lower panels: sons (n=10 total) and daughters (n=17 total) (F1). Black points indicate non-significant differences (p>0.05); blue points indicate significantly higher behaviour frequency in rejected groups (p<0.05); red points indicate significantly lower frequency in rejected groups (p<0.05). Error bars show 95% credible intervals. Rejected adults showed increased fear (+5%, p=0.03), approach/boldness (+20%, p=0.01), and aggression/hitting others (+10%, p=0.01), with decreased hiding on perch (−44%, p<0.01), carrying offspring (−7%, p=0.04), and jumping locomotion (−5%, p<0.01) compared to controls. Offspring of rejected parents showed decreased hiding on perch (−15%, p=0.05), jumping locomotion (−14%, p<0.01), and biting (−12%, p=0.01) compared to offspring of controls. Complete behavioural data provided in Supplementary Files 2-4, 8-9.

Females P0

Males P0

Females F1

Males F1

Figure 3. Effects of rejection on miRNA expression in whole blood across two generations of owl monkeys.
Heatmaps showing z-score normalised miRNA expression (blue=downregulation, red=upregulation; colour intensity indicates magnitude of dysregulation). Blood samples of 47 were included in analysis after quality control. (a) Rejected females versus control females (P0). (b) Rejected males versus control males (P0). No miRNAs reached significance (FDR $p<0.05$) in rejected adults; mml-miR-182 showed marginal downregulation (FDR p 0.05-0.10). (c) Well-reared female offspring of rejected parents versus offspring of controls. mml-miR-30a-5p was significantly upregulated specifically in female offspring of rejected parents (FDR $p=9.91\times10^{-3}$, FC=2.09, indicated by red asterisk)-the only miRNA passing stringent multiple testing correction. (d) Well-reared male offspring of rejected parents versus offspring of controls. No miRNAs reached significance. Differential expression analysis used LIMMA TREND with Benjamini-Hochberg FDR correction (Partek Flow). Complete miRNA expression data provided in Supplementary File 7.