The brain cortical similarity network: Development and sensitivity to early life stress in a

rat model

Rachel L. Smith[1,2], Stephen J. Sawiak[3,4], Lena Dorfschmidt[1], Ethan G. Dutcher[5], Jolyon A.

Jones[3,5], Joel D. Hahn[6], Olaf Sporns[7,8], Larry W. Swanson[6], Paul A. Taylor[9], Daniel R. Glen[9],

Jeffrey W. Dalley[1,3,5], Francis J. McMahon[2], Armin Raznahan[2], Petra E. Vértes[1,*], Edward T.

Bullmore[1,*]

1 Department of Psychiatry, University of Cambridge, Cambridge, CB2 0SZ, UK

2 Human Genetics Branch, National Institute of Mental Health, Bethesda, MD, USA 20892

3 Behavioural and Clinical Neuroscience Institute, University of Cambridge, Downing Site, Cambridge, UK

4 Department of Physiology, Development and Neuroscience, University of Cambridge, Cambridge, CB2

3EL, UK

5 Department of Psychology, University of Cambridge, Cambridge, CB2 3EB, UK

6 Department of Biological Sciences, University of Southern California, Los Angeles, CA, USA 90089

7 Indiana University Network Science Institute, Indiana University, Bloomington, IN, USA 47405

8 Department of Psychological and Brain Sciences, Indiana University, Bloomington, IN, USA 47405

9 Scientific and Statistical Computing Core, National Institute of Mental Health, NIH, Bethesda, MD, USA

20892

* P.E.V. contributed equally to this work with E.T.B.

Corresponding author: Rachel Smith (rls83@cam.ac.uk)

Abstract

Understanding how early life experiences shape brain network development is a key challenge in the neuroscience of mental health disorders. To address this, we used magnetic resonance imaging (MRI) similarity network analysis to study the effects of stress in the rat, an important animal model in neuropsychiatry. We measured magnetization transfer ratio (MTR) at each of 53 distinct cortical areas and estimated a cortical similarity network for each individual scan, in two independent experimental datasets. We first characterized normative network development in rats scanned repeatedly between postnatal days 20 (weanling) and 290 (mid-adulthood) (N=47), and then contrasted these findings with a cohort exposed to early life stress in the form of repeated maternal separation (RMS, N=40). The normative rat cortical similarity network exhibited biologically meaningful organization, aligning with prior cytoarchitectonic and tract-tracing data, and displayed complex topological features, including rich club organization. During postnatal and adolescent development, brain regions became more similar, including an early phase of fronto-hippocampal convergence. Early increases in inter-areal similarity were reversed in a later phase of fronto-hippocampal divergence in mid-adulthood. RMS exposure altered inter-areal similarity, especially between frontal and parahippocampal regions, that were also most active developmentally and in aging. Our results reveal how normative cortical network changes in the developing brain are influenced by early life stress. These findings suggest a new translational framework for elucidating how environmental risk factors lead to atypical development of cortical networks.

Keywords

Introduction

Altered trajectories of brain development are a major risk factor for psychiatric and

neurodevelopmental disorders (Sánchez et al., 2001; Paus et al., 2008; Lupien et al., 2009).

However, the biological mechanisms that govern brain development—and how they are shaped

by life experiences—remain incompletely understood. Tools that capture dynamic, systems-level

changes in neuroarchitecture can advance our insight into these processes. Network models of

macroscale brain organization, particularly those constructed using noninvasive neuroimaging,

provide a key framework for characterizing distributed brain structure and function (Bullmore and

Sporns, 2009, 2012; van den Heuvel and Sporns, 2011, 2013; Fornito et al., 2015). By quantifying

inter-regional relationships, these models reveal the organization and maturation of brain systems

across development and in response to experience.

Structural similarity analysis is an emerging approach for constructing individualized brain

networks using magnetic resonance imaging (MRI) (Li et al., 2017; Seidlitz et al., 2018; Sebenius

et al., 2023). These methods estimate inter-regional similarity using MRI-derived morphometric

features (Sebenius et al., 2024), resulting in networks in which nodes represent cortical areas and

edges reflect the strength of structural similarity between them. A recent implementation,

morphometric inverse divergence (MIND), quantifies similarity between voxel- or vertex-level

distributions of MRI features (Sebenius et al., 2023). MIND and related networks are reproducible

(Seidlitz et al., 2018; Sebenius et al., 2023), developmentally sensitive (Fenchel et al., 2020; Ruan

et al., 2023; Sebenius et al., 2023; Wu et al., 2023; Dorfschmidt et al., 2024), and responsive to

environmental exposures (Tian et al., 2021; González-García et al., 2023; Xiao et al., 2023;

Hettwer et al., 2024). MIND networks also show clinical relevance (Mahjoub et al., 2018; Homan

et al., 2019; Li et al., 2019; Lisowska and Rekik, 2019; Zhang et al., 2020; Cao et al., 2023),

significant heritability (Sebenius et al., 2023), and associations with gene expression (Morgan et

al., 2019; Seidlitz et al., 2020).

Despite these advantages, structural similarity networks have been almost exclusively studied in

humans, limiting their utility for testing causal mechanisms. Animal models are essential for

addressing this gap, enabling experimental manipulation and brain tissue access. Rats in

particular offer distinct advantages, including suitability for high-throughput experiments and

ethologically rich, human-relevant behaviors (Bryda, 2013; Ellenbroek and Youn, 2016). Despite

this, network representations of the rat brain remain limited. Existing data, primarily derived from

expertly curated tract-tracing studies (Swanson et al., 2017, 2019, 2020, 2022, 2024a, 2024b),

suggest that the rat connectome exhibits complex topology (Swanson et al., 2024b) akin to that

in humans (Bullmore and Sporns, 2009) and mice (Rubinov et al., 2015). However, these data

represent the composite rat brain and thus do not support individual-level analyses.

Here, we extend the MIND framework to rats, enabling construction of individual-level cortical

similarity networks from in vivo MRI and providing a tool to examine how early life environments

shape brain organization. We focus on magnetization transfer ratio (MTR), a proxy for cortical

myelination (Mancini et al., 2020)—a feature that is both developmentally dynamic (Hamano et

al., 1998; Downes and Mullins, 2014; Mengler et al., 2014) and environmentally sensitive in rats

(Bass et al., 1970; Krigman and Hogan, 1976; Breton et al., 2021; Long et al., 2021; Han et al.,

2022; Abraham et al., 2023), but underexplored in rat MRI studies (Bai et al., 1996).

We applied this method to two rat neuroimaging datasets: a longitudinal normative developmental

cohort of males scanned across postnatal development (**Fig. 1A**), and an experimental stress

cohort of males and females exposed to repeated maternal separation (RMS) as a model of early

life stress (**Fig. 1B**). We demonstrate that MIND networks reflect biologically meaningful features

of cortical organization that track age- and experience-related changes in network architecture.

These findings offer a novel approach for linking early life experiences to brain network

development.

Materials and Methods

Experimental design

Pre-existing structural MRI data from two independent cohorts, published in (Jones et al., 2024)

and (Dutcher et al., 2023), were used to assess network-level changes that occur during

development and in response to stress:

1.   Male normative developmental cohort (Jones et al., 2024) (**Fig. 1A**)

Male Lister Hooded rats (N=47) were kept on a reverse light/dark cycle with red light on from

7:30am - 7:30pm and white light for the other half of the daily cycle. Rats underwent MRI scanning

and weaning on postnatal day (PND) 20 or 21 (N=40; here referred to as PND 20). Rats were

scanned again at PND 35 (N=38), PND 63 (N=42), and once between PND 212-244 (N=43; here

referred to as PND 230). **Figure 1** shows the number of observations per animal in the male

normative developmental cohort (hence referred to as the normative developmental cohort), with

32 animals (68% of total) having scans at each of the four timepoints from PND 20 (post-weaning)

to PND 230 (aging adult). Experiments were carried out in accordance with the (U.K Animals)

Scientific Procedures Act (1986) under UK Home Office project licenses (PPL 70/7587 & PPL

70/8072) and were approved by the University of Cambridge Ethics Committee.

2.   Experimental stress cohort (Dutcher et al., 2023) (**Fig. 1B**)

Pregnant Lister Hooded rats (N=14) were purchased from Envigo (Blackthorn, UK). Litters were

delivered by spontaneous partum on gestational days 22-24. Within three days of birth, litter size

was adjusted to 4-6 pups, with each litter consisting of two female and two male pups (except for

one litter with four males and two females). If two litters were born within 24 hours of one another

(the case for 10 litters in total), pups were mixed between the litters. After litter size adjustment,

litters were allocated alternately by birth time to either the repeated maternal separation condition

(RMS; N=30 pups: 14 female, 16 male) or the control condition (N=28 pups: 14 female, 14 male).

PND 0 was defined as the day of delivery. Body weight was measured weekly starting at PND

20. Lights were on from 21:00 to 09:00. Experiments were conducted on Project License

PA9FBFA9F, in accordance with the UK Animals (Scientific Procedures) Act 1986 Amendment

Regulations 2012, the EU legislation on the protection of animals used for scientific purposes

(Directive 2010/63/EU), and the GSK Policy on the Care, Welfare and Treatment of Animal,

following ethical review by the University of Cambridge Animal Welfare and Ethical Review Body

(AWERB).

From PND 5-19 (inclusive), pups from RMS litters were separated from their dam for 6 hours a

day, starting between 11:00am-12:30pm. During separation, dams remained in their home cages

while pups were taken to a different room and placed together inside a ventilated cabinet. One

centimeter of bedding was provided and the temperature at the surface of the bedding was kept

between 30 °C and 35 °C through warming of the air and use of an electric heat pad. Control pups

were subject only to normal animal facility rearing. Following PND 20, pups from both groups

were weaned and housed in same-sex pairs at PND 20 and left undisturbed until early adulthood

except for weighing and once-weekly cage changes. All animals then underwent MRI scanning

at PND 63.

Figure 1.

A l Normative developmental cohort

B l Experimental stress cohort

Two independent experimental cohorts to assess reliability and validity of rat MRI similarity

networks as measures of developmental and stress-related changes in cortical microstructural

networks. A) Top: Study design for the normative developmental cohort: N=47 male Lister Hooded rats

were reared normally and had brain MRI scanning at PND 20 (N=40), PND 35 (N=38), PND 63 (N=42),

and within a few days of PND 230 (N=43). All MRI timepoints are indicated by a star; PND 63 is outlined in

black, as it marks the timepoint used to construct the normative network. Bottom: Scans per subject. Each

tick on the y-axis represents a subject, while the x-axis shows age. Each point represents a scan; repeated

scans of the same rat are connected by a line. **B)** Study design for the experimental stress cohorts. N=21

Lister Hooded pups were stressed by repeated maternal separation (RMS), i.e., for 1 hour/day every day

from PND 0 to PND 20, pre-weaning pups were separated from their dam. A control group of N=19 pups

was reared normally. All animals completed MRI scanning at PND 63 as young adults. As with panel A,

MRI timepoints are indicated by stars, and the primary comparison timepoint (PND 63) is outlined in black.

Figure created in https://BioRender.com.

MRI acquisition

For both cohorts, high-resolution MRI was performed on a 9.4T horizontal bore MRI system

(Bruker BioSpec 94/20; Bruker Ltd.). Images were acquired under isoflurane anesthesia using

the manufacturer-supplied rat brain array coil with the rat in a prone position. Structural images

were obtained based on a 3D multi-gradient echo sequence (TR/TE 25/2.4 ms with 6 echo images

spaced by 2.1 ms, flip angle 6° with RF spoiling of 117°). The field of view was 30.72 × 25.6 ×

20.48 mm3 with a matrix of 192 × 160 × 160 yielding isotropic resolution of 160 μm with a total

scan time of 6 min 36 sec with zero-filling acceleration (25% in the readout direction; 20% in each

phase encoding direction). Magnetization transfer pulses (10 μT, 2 kHz off-resonance) were

applied within each repetition to enhance gray-white matter contrast. Post-reconstruction, images

from each echo were averaged after weighting each by its mean signal.

Throughout all scanning procedures, rats were anesthetized with isoflurane (1-2% in 1 L/min O2:

air 1:4). Respiratory rate, oxygen saturation and pulse rate (SA Instruments; Stony Brook, NY)

were measured with anesthetic dose rates adjusted to ensure readings remained within a

physiological range. Body temperature was measured and regulated with a rectal probe and

heated water system to 36-37°C.

Image registration

Structural MRI image preprocessing was performed using the AFNI software package version

AFNI_24.2.03 (Cox, 1996). Briefly, magnetization transfer (MT) images for each rat were first

deobliqued, spatially oriented, and translated to have spatial overlap with the Waxholm Space

reference template (WHS) (Kleven et al., 2023). This space was selected due to its alignment

with gold-standard histological rat brain atlases (Paxinos and Watson, 2006; Swanson, 2018) and

the atlas's parcellation granularity (N=222 regions). The @animal_warper command (Jung et al.,

2021) was then used to nonlinearly align all MT images to the WHS template. Because some

input datasets were notably smaller than the WHS standard template, the -init_scale option was

added and used to increase the search space of the registration algorithm, scaled to the relative

size ratio of the input scan to the template. Quality control image outputs for each scan (produced

by @animal_warper) were then manually reviewed; for any images that were not successfully

registered, @animal_warper was run again with the -init_scale parameter altered to better

approximate the size ratio. All image registration scripts are available with this publication.

Magnetization transfer ratio calculation

Magnetization transfer ratio (MTR) was calculated on a per-voxel basis for each scan in native

space. First, native scans were scaled according to the receiver gain (RG) parameter used in the

scan acquisition protocols. The scaled MT and proton density (PD)-weighted scans in native

space were then used to calculate MTR according to the equation (PD - MT) / PD. Both steps

were executed using 3dcalc in AFNI (Cox, 1996).

Manual evaluation of MTR data quality resulted in the exclusion of 7 scans, typically because of

motion artifacts and particularly at the earlier timepoints. Example MT images and their quality

control images are shown in **Figure S1**. The analyzable MRI datasets available following

preprocessing and quality control of the two cohorts are summarized in **Table 1**.

	postnatal days (PND)	*N*
Normative developmental cohort		
	20	40
	35	38
	63	41
	230	43
Experimental stress cohort		
Repeated maternal separation (RMS)	63	21
Normal maternal rearing (control)	63	19

Table 1: Number of MRI scans included in the study analyses following quality control. *N* represents

the number of unique individuals scanned at each timepoint within each cohort.

Morphometric INverse Divergence (MIND) network calculation

MIND networks estimate structural similarity from MRI data (Sebenius et al., 2023). Briefly,

cortical regions are represented by a distribution of structural MRI features sampled at many

points within the region, in this case, at each voxel. The MIND similarity between each pair of

regions is then calculated using the Kullback-Leibler (KL) divergence between their feature

distributions.

<u>Input data generation</u>: Each MTR image was aligned with the WHS atlas in the native space of

the respective MT scan (the scan with the highest contrast), so that each voxel in the MTR scan

was labeled with a region of interest based on voxel assignment output from the registration

pipeline. For each scan, a two-column CSV was generated, in which the first column "Label" was

the region of interest, and the second column ("MTR") was the value for the corresponding voxel

in the MTR scan. For cortical MIND calculation, the input CSV was filtered to contain only voxels

belonging to regions under the "Cerebral cortex" hierarchical level of the WHS atlas (which

excludes olfactory bulb regions).

<u>MIND network construction</u>: MIND networks were constructed for each scan by calculating the

KL divergence between pairwise combinations of regional MTR profiles using the MIND toolkit

(https://github.com/isebenius/MIND). The MIND algorithm used can be sensitive to differences in

number of datapoints. To balance this, we estimated KL divergence for each pair of regions by

estimating the same number of voxels (5000) from each region, regardless of its size.

<u>MIND network phenotypes</u>: In network neuroscience, topography refers to the spatial layout of

brain regions, while topology refers to how those regions are interconnected. Here, we focus on

the topological structure of the normative cortical MIND network.

Edge weight and nodal strength (or weighted degree) were considered features of interest for

downstream analyses. Edge weights were calculated as $w = 1 / (1 + KL)$, per the MIND toolkit

(Sebenius et al., 2023). Nodal strength was calculated as the sum of all edge weights for a given

region; hubs were defined as the 10 regions with the highest nodal strength. The rich club

coefficient (Φ), a network measure that quantifies the tendency of hubs to be more densely

interconnected compared to lower-strength nodes (van den Heuvel and Sporns, 2011; Fornito et

al., 2016a), was calculated for the normative network as follows:

Equation 1.

$$\phi_w(s) = \frac{W_{>s}}{W_{\max,>s}}$$

where

● $W_{>s}$ = sum of edge weights among nodes with strength greater than *s* (i.e., hubs)

● $W_{\max,>s}$ = sum of the strongest edges in the network

Edge distance calculation

The midline of the WHS atlas was determined using the AFNI function 3dcalc to separate the left

and right hemispheres. To approximate the center of each WHS region of interest, the AFNI

function 3dCM -Icent (Cox, 1996) was then used. The Euclidean distance between pairwise

regional centers was then calculated as their edge length.

Rat atlas mapping

We mapped the WHS atlas into Brain Maps 4.0 (BM4) atlas space (Swanson, 2018), Zilles atlas

space (Zilles, 2012), and Allen Mouse Brain Atlas space (AMBA) (Lein et al., 2007) to compare

the MIND similarity network to cortical tract-tracing data (Swanson et al., 2017, 2024b), cortical

type (García-Cabezas et al., 2023), and mouse spatial transcriptomic expression (Yao et al.,

2023), respectively. We did so first by aligning cortical regions based on nomenclature. However,

not all regions maintained consistent nomenclature across atlases, so we also visually inspected

anatomical   alignment   of   regions   using   the   WHS   EBRAINS   online   resource

(https://www.ebrains.eu/tools/rat-brain),   BM4   atlas   maps   (https://sites.google.com/view/the-

neurome-project/brain-maps), Zilles atlas cortical maps in stereotaxic coordinates (available for

download   at   https://link.springer.com/book/10.1007/978-3-642-70573-1),   and   AMBA   online

interactive   atlas   viewer   (http://atlas.brain-map.org/atlas?atlas=1).   Briefly,   we   identified   the

position of each region in the reference (BM4, Zilles, or AMBA) atlas maps, then panned through

the WHS atlas using the EBRAINS tool to approximate the same coronal slice and identify what

region most closely aligned with the reference anatomical position. We also considered relative

positioning of surrounding regions to determine this alignment. All atlas mappings are provided

as resources in **Tables S2A-C**.

In this work, the BM4 atlas was used as the reference space for the tract-tracing comparison, the

Zilles atlas was used as the reference space for the cortical type comparison, and the AMBA atlas

provided reference space for the mouse transcriptomics comparison. If multiple WHS atlas

subdivisions comprised a single reference atlas region, the median across these subdivisions was

taken as the MIND edge weight for that region.

Tract-tracing Jaccard index calculation

To convert the cortical tract-tracing matrix (Swanson et al., 2017, 2024b) into a similarity network,

the Jaccard index between each pairwise combination of regions was calculated. The set for a

given region *a* was defined as the ordinal tract-tracing weight with each other region. The Jaccard

index *J* between two regions *a, b* was then calculated as the intersection of their sets divided by

the union as follows:

Equation 2.

$$J(a,b) = \frac{|A \cap B|}{|A \cup B|}$$

Cortex type network thresholding

WHS cortical regions were categorized by their cortical type using the data presented in (García-

Cabezas et al., 2023) and grouped according to whether they are part of the archicortical

allocortex or mesocortex (agranular or dysgranular subdivisions). Paleocortical and eulaminate

regions were excluded from this analysis due to the very small number of constituent regions

defined by the atlas. Each MIND edge was then defined as 'intra-class' or 'inter-class' based on

whether both regions were part of the same cortex type or not, respectively. To assess the extent

to which top-weighted MIND edges consisted of two regions within the same cortex type, the

normative MIND network was thresholded across densities (comprising 0-10% of top-weighted

edges), and the percentage of intra-class edges was calculated.

Statistical analysis

<u>Null network generation</u>: Ten thousand null networks were generated to assess whether

normative MIND network alignment with tract-tracing similarity and cortical type were greater than

would be expected by chance. To do so, all MIND network edges were classified into three evenly

sized bins based on distance: proximal, intermediate, and distal. Edge weights within each bin

were then reshuffled to generate a null network that preserved distance structure. The tract-

tracing and cortical type analyses were repeated for each null network to generate a null

distribution for comparison of each analysis.

<u>Developmental slope</u>: Changes that occurred in edge weight (*w*) and nodal strength (*s*) during

normative development (Δw_{dev}; Δs_{dev}) and aging (Δw_{age}; Δs_{age}) were quantified by calculating the

linear gradient or slope of age-related change in each period. Early development was defined as

PND 20 to PND 35, as the highest proportion of change occurred here, and aging was defined as

PND 63 to PND 230. For each region, a linear mixed effects model was fit with the normalized

strength as the dependent variable, continuous age and total brain volume as fixed effects, and

each individual rat as a random effect:

Equation 3A.

$$\text{normalized strength} \sim \beta_1 \, \text{age} + \beta_2 \, \text{TBV} + (1|\text{subject})$$

Edge-level analyses were run using coarse-grained systems labels for interpretability. In this

case, ROI-level edges were also included as a random effect:

Equation 3B.

$$\text{normalized weight} \sim \beta_1 \text{ age} + \beta_2 \text{ TBV} + (1|\text{subject}) + (1|\text{ROI edge})$$

The coefficient for age, β_1, was estimated as the slope for that region within the given period. The

effect size (Δw; Δs) was used to plot and compare between early development and aging epochs.

<u>RMS-control effect size</u>: Case-control analysis at PND 63 was used to identify changes that

occurred in response to RMS and were measurable in young adulthood. For each region, the

following model was used:

Equation 4A.

$$\text{normalized strength} \sim \beta_1 \text{ group} + \beta_2 \text{ sex} + \beta_3 \text{ age} + \beta_4 \text{ TBV}$$

As with development, case-control edge effects were determined at the systems-level, with ROI-

level edges included as a random effect:

Equation 4B.

$$\text{normalized weight} \sim \beta_1 \text{ group} + \beta_2 \text{ sex} + \beta_3 \text{ age} + \beta_4 \text{ TBV} + (1|\text{ROI edge})$$

The effect size associated with the group term was estimated as the actual case-control effect

size for a given region or edge. Then, for each region and edge, 1000 permutations were run in

which the group label was randomly sampled, and the case-control effect size was estimated

under the null hypothesis. The Z-score for each actual effect size in this permutation distribution

was calculated; any region with absolute value Z-score > 1.96 (P=0.05) was considered

significant.

Relating developmental changes and case-control stress effects: To characterize the relationship

between RMS case-control effects and normative developmental changes, Pearson's correlation

was run on Δw_{dev} and Δw_{age} vs edge-level PND 63 case-control effect size. To assess the extent

to which the strength of this relationship was greater than expected under the null hypothesis,

10000 permutations were run, in which the edge assignment of RMS effect size was randomly

resampled and again correlated with Δw_{dev} and Δw_{age}. The Z-score of the actual Pearson's r was

determined, and the P-value was calculated as 1 - (proportion of permuted correlations that were

smaller than the observed correlation).

Across all statistical analyses, P values were corrected for multiple comparisons using the

Benjamini-Hochberg (BH) procedure (Benjamini and Hochberg, 1995).

Rat brain plot construction

Flatmap rendering: A flatmap is a two-dimensional representation of the brain that facilitates

visualization of spatial relationships between regions (Herrick, 1948; Swanson, 2000). Flatmaps

are particularly useful for displaying cortical organization and topographic continuity, avoiding

distortions introduced by sulcal and gyral folding in three-dimensional renderings (Swanson,

2000). While flatmapping techniques have been used elsewhere in comparative neuroanatomy

and cortical cartography (Herrick, 1948; Nauta and Karten, 1970), Swanson and colleagues were

the first to systematically adapt this approach to the rat brain (Swanson, 1992; Hahn et al., 2021;

Hahn and Duckworth, 2023). In this study, rat brain flatmap plots were derived from the work of

(Hahn et al., 2021; Hahn and Duckworth, 2023).

The SVG file provided by (Hahn and Duckworth, 2023) was first converted to well-known text

(WKT) using code from https://github.com/davidmcclure/svg-to-wkt. The WKT was then parsed

into geometry objects in R using the package sf version 1.0.17 (Pebesma, 2018). Flatmaps

represent a practical and underutilized tool for visualizing architectonic variation and similarity in

the rat cortex, particularly in connectomic studies. To facilitate broader use, we provide all flatmap

construction resources.

Anatomical rendering: Rat brain maps were also visualized using an anatomical rendering that

preserved 3-dimensional brain structure. Nifti files representing the right and left hemispheres of

the WHS (generated in Edge distance calculation) were loaded into R using the package RNifti

version 1.7.0 (Clayden et al., 2024). Each region was converted to a point cloud based on voxel

intensity index using the R package lidR version 4.1.2 (Roussel et al., 2020). Outlying voxels

(noise) were removed using the isolated voxel filter (resolution = 4, N = 25) (Roussel et al., 2020).

Point clouds were converted to geometry objects in R using the package sf version 1.0.17

(Pebesma, 2018).

Results

Defining and describing the normative network

We calculated each individual rat cortical network as the {53×53} matrix of edge weights (*w*),

representing all pairwise MIND similarity values between 53 brain regions. To estimate the

normative cortical microstructural network, we computed the median of each edge weight across

the 41 male adult rat brains (postnatal day [PND] 63; young adulthood) from the normative cohort

(**Fig. 1A**; **Fig. 2A**; **Table 1**). The 53 regions were grouped into 15 coarse-grained cortical systems,

as defined by the Waxholm Space Atlas (Kleven et al., 2023), for edge-level analyses and

interpretability of results (**Fig. 1B**,**C**).

Figure 2.

A I Adult normative cortical microstructural network

B I Cortical system topography   **C I Cortical system anatomy**

Structure of the normative rat cortical microstructural network. A) Heatmap representation of the

normative rat connectome, defined as the median edge weight across rats in the normative cohort, at

postnatal day 63 (PND 63). Rows and columns are ordered by cortical systems, as indicated by annotation

bars. Tile color indicates strength of MIND similarity (edge weight, w). **B)** A flatmap rendering of cortical

systems derived from (Hahn et al., 2021) and (Hahn and Duckworth, 2023). Colors correspond to cortical

systems as indicated below. **C)** Anatomical positions of regions of interest within each cortical system. Each

point represents the center for a given cortical region of interest (N=53), as defined by the WHS; lines

represent intra-system edges. For coronal and axial plans, only right hemisphere nodes are shown.

Normative network topology

The normative network distributions of nodal strength (i.e., the sum of all edge weights connecting

it to the rest of the network, s; **Fig. 3A**) and edge weights were left-skewed (**Fig. 3B**), indicating

a generally high MTR-based anatomical similarity across the cortex. Hubs, defined as regions

with the highest strength (Fornito et al., 2016b), and rich-club organization, which examines

whether hubs have stronger similarity with each other than expected by chance (Fornito et al.,

2016a), are functionally relevant features of brain networks described in humans and other

organisms (Bullmore and Sporns, 2009; Towlson et al., 2013; Rubinov et al., 2015; Swanson et

al., 2024b). We found this principle held true in rat brain MIND networks. The hubs, defined as

the 10 nodes with the highest strength, were largely hippocampal and piriform regions (**Fig. 3C**).

The rich club coefficient (Φ), calculated using **Equation 1**, was 0.95, compared to a median

coefficient of 0.45 across 10,000 distance-corrected null networks (**Fig. 3D**,**E**), indicating a

significantly greater-than-random strength of connectivity amongst cortical hubs (Z=73;

P_{perm}=1.0*10^{-4}). This evidence for a rich club organization in MIND networks was consistent with

prior reports of rich clubs in the meta-analytic connectome from rat tract-tracing data (Swanson

et al., 2024b) and in other species and modalities of brain networks (van den Heuvel and Sporns,

2011; Towlson et al., 2013).

Figure 3.

A | **Nodal strength (*s*)**

B | **Strength distribution**

C | **Top 10 hubs**

D | **Rich club connectivity**

E | **Rich club coefficient**

Graph-based metrics of the normative network. A) Maps of normative nodal strength. Left: Topographic

(flatmap) cortical rendering (right hemisphere only). Right: Anatomical MRI rendering. Network hubs, or top

10 nodes by strength, are labeled. CA1=cornu ammonis 1; PIR1=piriform cortex, layer 1; CA3=cornu

ammonis 3; DG=dentate gyrus; S1-tr=primary somatosensory area, trunk representation; DI=dysgranular

insular cortex; PIR2=piriform cortex, layer 2; PIR3=piriform cortex, layer 3; PER35=perirhinal area 35;

Au1=primary auditory area. **B)** Nodal strength distribution of the adult normative network. Each point on the

x-axis shows the strength of a given region (defined as the sum of all edge weights for that region), while

the y-axis shows smoothed density of nodes for each strength bin. **C)** Ten regions with the highest nodal

strength, i.e., "hubs". The x-axis represents the median strength (± standard deviation) of each region; each

point is colored by cortical system as in **Figure 2**. The y-axis shows each region of interest (same

abbreviations as panel A). **D)** Anatomical representation of the network rich club. Each point represents a

hub (from panel C), and the interconnectivity among all hubs is indicated by connecting lines. The width

and color of each line indicates edge weight; i.e., the strength of similiarty between the two hubs. **E)**

Normative MIND network rich club coefficient versus 10000 distance-corrected null networks. The x-axis

shows the rich club coefficient, while the y-axis indicates density distribution. The rich club coefficient of the

normative network, defined as the sum of all hub inter-connections (N=90) divided by the sum of the top

N=90 edges, is shown in by the position of the maroon line (Φ=0.95). The black curve shows the distribution

of rich club coefficients across distance-corrected null networks (median Φ=0.45).

Network validation and reliability

Research in humans has shown that structural similarity networks reflect key neurobiological

features, including axonal connectivity and cytoarchitectonic similarity (Sebenius et al., 2023,

2024). Since this study is the first to describe structural similarity networks in rats, we aimed to

establish these relationships within the normative network reported here. Confirming the

neurobiological relevance of this network enhanced the interpretability and significance of

downstream findings.

We validated the normative MIND network against four key predictions: that MIND similarity was

greater between regions that (i) were spatially closer to each other, (ii) were more architectonically

similar to each other, (iii) had more similar gene expression profiles, and (iv) had more similar

profiles of axonal projections in prior tract-tracing data. All four of these predictions were verified

as detailed below.

Similarity–distance relationship: First, we found that morphometric similarity decreased with

increasing distance between regions such that the Euclidean distance between node centers was

negatively correlated with edge weight (i.e., MIND similarity; *r*=-0.66; *P*<0.001; **Fig. 4A**). This

result aligned with expectations, as brain regions that are in closer spatial proximity tend to exhibit

greater structural similarity (Alexander-Bloch et al., 2013).

Figure 4.

MIND-based similarity is related to metrics of cortical organization. A) Scatter plot of the distance

between two nodes (x-axis) and their MIND similarity, or edge-weight between them (y-axis). Distance is

defined as the Euclidean distance between region of interest centers, as determined using the Waxholm

Space atlas. **B)** Topographic (flatmap) rendering of cortical types (left hemisphere only). Regions are

colored by cortex type, as defined by (García-Cabezas et al., 2023). **C)** Proportion of intra-class edges

across network density thresholds. The x-axis shows the proportion of top-weighted edges considered,

while the y-axis indicates the percentage of those top-weighted edges that consists of two regions from the

same cortex class. The maroon curve shows the percentage of within-class edges in the normative MIND

network, while the gray line shows the mean percentage (± standard deviation) of within-class edges across

10000 permuted null networks. **D)** Correlation between similarity of tract-tracing connection profiles

between pairwise combinations of regions and strength of MIND similarity. The Jaccard index for edge *ij*

was defined as the intersection of tract-tracing connections (i.e., where *ik == jk*) divided by the union of all

connections containing *i*, *j*, or both (**Equation 2**). **E)** Heatmap representations of the median PND 63 control

network from the experimental stress cohort (top left of the diagonal) and the median PND 63 network from

the normative developmental cohort (bottom right of the diagonal). Panel legend is the same as **Figure 2A**.

F) The relationship between edge weights in the median PND 63 network from the normative development

cohort (x-axis) and median PND 63 control network from the experimental stress cohort (y-axis; *r*=0.86;

P<0.001). Each point represents an edge; the line of best fit is shown in maroon.

<u>Cytoarchitectonic relationships to similarity</u>: Second, if MIND similarity truly reflects

cytoarchitectonic similarity, we would expect regions of the same cortical type to exhibit higher

MIND edge weights. The rat neocortex is structured along gradients of laminar differentiation,

which expanded from allocortical areas (archicortical [hippocampal] and paleocortical [olfactory])

according to the dual origin theory (García-Cabezas et al., 2023) (**Fig. 4B**). In rodents, these

cortical gradients extend through mesocortical (agranular and dysgranular) into isocortical

(eulaminate) areas (García-Cabezas et al., 2023). To test the relationship between MIND

similarity and cytoarchitectonics, we co-registered our neuroimaging data with a cytoarchitectonic

atlas of the rat brain (García-Cabezas et al., 2023) (**Fig. 4B**), and compared MIND edge weights

between areas of the same cortex class ("intra-class" edges) versus areas of different cortex

classes ("inter-class" edges), removing isocortical regions due to small sample size (N=5 regions).

Intra-class edges between two allocortical areas ("allo-allo") had significantly higher MIND

similarity than edges between allocortical and mesocortical ("allo-meso") areas (false discovery

rate, FDR < 0.05; (**Fig. S2**). However, meso intra-class edges were not significantly more similar

than allo-meso. To probe this relationship further, we calculated the percentage of intra-class

edges (allo-allo and meso-meso) at various binary network densities, thresholding across a range

of sparse connection densities (1% to 10% of all possible pairwise edges). We then compared

these to distance-corrected null networks thresholded across the same range (**Fig. 4C**). Across

all thresholds, the empirical rat network demonstrated a higher percentage of intra-class edges

than the null networks (**Fig. 4C**), indicating a higher-than-chance density of cytoarchitectonically

similar edges among edges with the highest MIND similarity.

<u>Mouse transcriptional relationships to similarity</u>: Third, we predicted that regions with more similar

patterns of gene expression would also exhibit higher MIND similarity. However, a spatial

transcriptomic atlas of the rat brain is not yet available. To overcome this limitation, we leveraged

homology between the rat and mouse brain to access spatially comprehensive transcriptomic

data from the Allen Mouse Brain Atlas (AMBA; N=437 genes) (Yao et al., 2023). While we

acknowledge the challenges of cross-species comparisons, our goal was to assess the general

consistency between edge-wise MIND similarity and transcriptional similarity, rather than

expecting a precise one-to-one correspondence.

We tested and verified that regions with higher MIND similarity also demonstrated higher

transcriptomic similarity, calculated as the pairwise regional correlation between gene expression

profiles (**Fig. S3**). These findings support our general hypothesis that MIND similarity reflects

broad patterns of transcriptomic similarity.

<u>Tract-tracing relationships to similarity</u>: Fourth, we hypothesized that MIND similarity would reflect

patterns of axonal connectivity. To test this, we converted meta-analytic rat tract-tracing data

(Swanson et al., 2024b) to a similarity matrix by calculating the Jaccard index of the log10-

transformed ordinal connection weight between pairwise regions (**Equation 2**; **Fig. S4A**). This

transformation yielded a continuous measure representing the similarity of axonal connectivity

profiles between regions (i.e., the extent to which two regions share common connectivity

patterns).

When the tract-tracing Jaccard index was correlated with MIND edge weights, we found a positive

relationship (ρ=0.25; *P*<0.001; **Fig. 4D**). To assess the significance of this correlation, we

compared it to a null distribution generated from 10000 distance-corrected null networks. The

observed Spearman correlation was significantly greater than chance (*Z*=6.04; P_{perm}=1.0*10^{-4};

Fig. S4B), suggesting a meaningful relationship between anatomical connectivity profiles and

MIND similarity.

<u>Cross-cohort reproducibility</u>: Finally, to assess the reliability of MIND network analysis, we directly

compared the median adult MIND network for the normative developmental cohort (N=41, PND

63; **Fig. 1A**) to the median adult MIND network for an independent cohort of rats: namely, the

normally reared (control) group in the experimental stress cohort (N=19, PND 63; **Fig. 1B**; **Fig.**

4E). Edge weights were highly correlated between the normative adult and independent control

MIND networks (*r*=0.86; *P*<0.001; **Fig. 4F**), as was nodal strength (*r*=0.91; *P*<0.001). These

results demonstrate a high level of replicability of the cortical microstructural network estimated

by identically implemented MIND analyses of MTR data collected using the same sequences in

two independent cohorts of rats.

Normative developmental changes in the rat cortical microstructural network

Having established convergent validity and inter-sample replicability of MIND networks, we next

harnessed this analytic approach to model network level reorganization of the rat brain over

development. This strategy, applied to a longitudinal MRI dataset spanning PND 20, 35, 63 and

230 (N=162 total scans, **Fig. 1A**, **Methods**, **Table 1**), revealed developmental changes in (i) each

region's morphometric similarity with the rest of the brain (strength, *s*), and (ii) the morphometric

similarity (edge strength, *w*) between each unique pair of regions. Visual inspection of the median

MIND matrices for each of the four time-points indicated that there are age-related changes in the

cortical pattern of inter-areal similarity. For example, frontal cortical areas (frontal association

cortex, orbitofrontal cortex, mediofrontal cortex, and motor areas) become more similar to the rest

of the cortex during early development, and then strikingly less similar during later aging (**Fig.**

5A). Likewise, nodal strength showed a general tendency to increase during development and

decrease during aging (**Fig. 5B**).

Figure 5.

A | Normative network at each timepoint

B | Strength distribution by PND

C | Early life strength changes (Δs$_{dev}$)

E | Δs correlation

D | Age-related strength changes (Δs$_{age}$)

F | Δw correlation

G | Early life edge changes (Δw$_{dev}$)

H | Age-related edge changes (Δw$_{age}$)

The normative rat cortical connectome generally increases in similarity in early development and

decreases in similarity in aging. A) Heatmap representation of the connectome throughout development

in the normative cohort (median edge weight, at each timepoint). Rows and columns are ordered by

decreasing-to-increasing nodal strength within broader cortical system (in the same order as **Figure 2A**).

Tile color indicates strength of MIND similarity (edge weight, w). **B)** The normative strength distribution

through development, defined as the median strength per ROI at each timepoint. Each point represents a

region of interest; boxplots show the overall strength (s) distribution at each timepoint. The normative

strength distribution at PND 20 is significantly lower than the distribution at PND 63 (Dunn test P=0.019);

the nodal strength distribution at PND 230 is significantly lower than the distributions at PND 35 (Dunn test

P=0.027) and PND 63 (Dunn test P=0.003) (Krustal-Wallis P=0.002). **C)** Anatomical patterning of nodal

strength changes in early development (Δs_{dev}). Left: Volumetric rendering; Right: Flatmap rendering (left

hemisphere/top half only; (Hahn et al., 2021; Hahn and Duckworth, 2023)). Brown indicates decrease in

strength, teal indicates increase in strength. Δs_{dev} was considered significant if |t-value| of the age term in

the mixed effects model was greater than 2. **D)** Anatomical patterning of nodal strength changes in aging

(Δs_{age}; flatmap rendering right hemisphere/bottom half). Figure key same as panel C. **E)** Pearson correlation

between Δs_{dev} (x-axis) and Δs_{age} (y-axis; r=-0.39; P=0.004). Each point represents a region of interest; the

line of best fit is shown in maroon. **F)** Pearson correlation between Δw_{dev} (x-axis) and Δw_{age} (y-axis; r=-0.55;

P<0.001). Each point represents an edge; the line of best fit is shown in maroon. **G)** A circle plot of

significantly changed system-level edges in early development (Δw_{dev}; |t|> 3.3). Each datapoint in the circle

represents a broader cortical system, colored and labeled by system. The color and size of the curves

connecting two points show the change in edge weight (brown indicates decreasing similarity; teal indicates

increasing similarity). **H)** A circle plot of significantly changed edges in aging (Δw_{age}). Figure key same as

panel G.

We quantified the regional rate of change in early development as the linear gradient or slope of

age-related change in nodal strength for each cortical area between PND 20 (weanling) to PND

35 (adolescence), Δs_{dev}**,** accounting for inter-subject variation (**Equation 3A**). Similarly, to

calculate changes in aging, we applied the same model to estimate the age-related change in

strength for each cortical area from PND 63 (young adult) to PND 230 (mid adult) timepoints,

Δs_{age}.

During development, regions in most cortical systems significantly increased in nodal strength,

especially the hippocampal formation and motor cortex (Δs_{dev} t-values=7.19 and 5.22,

respectively; **Fig. 5C**). The parahippocampal region decreased in similarity with the rest of the

brain (Δs_{dev} t=-1.34). In contrast to these developmental changes, most cortical systems

decreased in nodal strength during aging, especially areas of the frontal cortex, including the

orbitofrontal (Δs_{age} t=-11.4), motor (Δs_{age} t=-6.13), and mediofrontal (Δs_{age} t=-5.94; **Fig. 5D**)

cortices. Early life and aging effects on nodal strength were negatively correlated (r=-0.39;

P=0.004; **Fig. 5E**), indicating that those regions showing the strongest strength increases in

development also tended to show the most rapid strength decreases in aging. Ten fronto-

hippocampal regions (Frontal: dorsolateral orbital area, frontal association cortex, frontal

association area 3, primary motor area, secondary motor area; Hippocampal: dentate gyrus,

perirhinal area 35, cornu ammonis 1) both significantly increased in strength in early development

and significantly decreased in strength in aging. Only one region, the parasubiculum, significantly

decreased in early development and increased in aging.

We also calculated changes in early development and aging for each inter-areal similarity or edge

in the MIND networks (Δw_{dev} and Δw_{age}, respectively; **Equation 3B**). In early development, the

hippocampal formation demonstrated notable increases in similarity with frontal regions, including

the orbitofrontal (Δw_{dev} t=17.6), motor (Δw_{dev} t=15.5), frontal association (Δw_{dev} t=11.7), and

mediofrontal (Δw_{dev} t=7.97) cortices (**Fig. 5G**). In contrast, parahippocampal regions largely

decreased in similarity with other cortical systems, including the somatosensory (Δw_{dev} t=-13.4)

and piriform (Δw_{dev} t=-11.0) cortices. In aging, the most salient changes in edge weight involved

orbitofrontal regions, which significantly diverged from every other cortical system, most notably

the somatosensory (Δw_{age} t=-30.1), hippocampal (Δw_{age} t=25.7), parahippocampal (Δw_{age} t=22.5)

regions (**Fig. 5H**). As with strength, edge weight effects during development and aging were

negatively correlated (r=-0.55; P<0.001; **Fig. 5F**), demonstrating that frontal and hippocampal

systems with the most rapid increases in pairwise similarity during early development showed the

fastest decreases in similarity, or increases in dissimilarity, during aging.

Early life environmental stressors perturb adult cortical similarity networks

We assessed the impact of early life stress—as modeled by repeated maternal separation

(RMS)—on the nodal strength of the young adult cortical microstructural network by running

multiple case-control comparisons for each cortical area at the PND 63 timepoint (**Equation 4A**;

Fig. 1B). Exposure to RMS was associated with strength decreases in most brain regions (N=40

of 53 total) (**Fig. 6A**). Four regions had significant differences following permutation testing: the

lateral entorhinal cortex, perirhinal area 36, and cingulate area 1 decreased in strength in RMS

(permutation Z-scores=-3.07, -2.31, and -2.18, respectively), while the frontal association cortex

demonstrated increased strength following RMS (Z_{perm}=2.08; **Fig. 6B**)

Figure 6.

A | Anatomical patterning of RMS strength effects

RMS < Control   RMS > Control

−4   −2   0   2   4

B | Significant RMS effects

Control
RMS

Frontal association

Cingulate area 1

Perirhinal area 36

Lateral entorhinal

−2   −1   0   1   2
Normalized strength (s)

C | Experimental cohort MIND networks

RMS PND 63 median (N=21)

Control PND 63 median (N=19)

0.9
0.8
0.7
0.6
0.5

D | Effects of RMS on edge weights

Piriform
Frontal association
Auditory
Orbito-frontal
Temporal association
Medio-frontal
Hippo-campal
Motor
Parahippo-campal
Posterior parietal
Insular
Somato-sensory
Retro-splenial
Cingulate
Visual

RMS < Control   RMS > Control

−4   0   4

Impact of early life stress on nodal strength and edge weights. A) Anatomical distribution of the post-

natal day 63 (young adulthood) RMS-control effect sizes across brain slices. For each region of interest, a

linear model was run of the normalized strength on group + age + sex + TBV, and the group statistic was

extracted as the actual effect size. Then, 1000 permutations were run in which the group assignments were

shuffled, the same linear model was run, and the group statistic was extracted as the permuted effect size.

The Z-score of the actual effect size was calculated as its position in the permuted distribution. Regions

with positive Z-scores (shown in red) demonstrated increased strength in RMS subjects in young adulthood;

regions with negative Z-scores (shown in blue) demonstrated decreased strength. The flatmap rendering

was derived from (Hahn et al., 2021; Hahn and Duckworth, 2023). **B)** Boxplots showing the nodal strength

distribution by group for the four regions with significant case-control differences at PND 63 ($|Z_{perm}| > 1.96$).

The x-axis shows nodal strength, corrected for covariates and normalized for visualization purposes. Each

point represents a subject, while the box-and-whiskers plot shows the overall distribution by group. Blue

indicates control, red indicates RMS. **C)** The median PND 63 MIND networks in control subjects (top) vs

RMS subjects (bottom). Figure legend is the same as Fig. 2A. **D)** A circle plot of significant RMS-control

edge weight differences in young adulthood (PND 63; $|t| > 3.3$). Node order and system annotation are the

same as **Figure 5** panels G and H. Curves connecting two points indicate a significant case-control

difference in edge weight; colored by effect size (with red indicating higher weight in RMS and blue

indicating higher weight in control); line width indicates absolute value effect size.

Qualitatively, visual inspection of the median MIND matrices in control (N=19) and RMS-exposed

rats (N=21) indicated that the early life stressor induced changes in the cortical pattern of inter-

areal similarity, especially in frontal and hippocampal systems (**Fig. 6C**). Indeed, edge-level case-

control analyses (**Equation 4B**) showed that RMS-exposed rats had strongly decreased similarity

between parahippocampal cortex and several other cortical areas (including the orbitofrontal

cortex (t=-6.89), mediofrontal cortex (t=-5.89), and motor cortex (t=-4.69; **Fig. 6D**). This contrasts

with markedly increased similarity between the frontal association cortex and other areas,

including the somatosensory (t=5.40) and insular cortices (t=5.17). Together, these results

indicate vulnerability in network structure to environmental stress, especially in frontal and

parahippocampal regions most sensitive to developmental and aging changes in nodal strength.

Effects of early life stress are nested within normative developmental changes in the

cortical microstructural network

Finally, we tested more formally for potential congruence between stress effects and normative

network changes in development and aging. Strikingly, variation in the effects of RMS on inter-

areal edge weights in the rat cortical microstructural network was positively correlated with

variation in the edge weight changes over normative development (Δw_{dev}; r=0.18, P_{perm}=0.03;

Z_{perm}=1.90; **Fig. S5** (left)). Specifically, those edges showing greatest similarity increases in

normative development also tended to show greater similarity increases following RMS.

Furthermore, the effects of RMS on edge weight were significantly and negatively correlated with

normative age-related changes in edge weight (Δw_{dev}; r=-0.19, P_{perm}=0.02; Z_{perm}=-2.03; **Fig. S5**

(right)), indicating that the edges with the greatest similarity increases following RMS were also

those that decreased in similarity the most in normative aging. Taken together, these results are

consistent with a model in which early life stress accelerates normative brain reorganization

during development; and areas which are most developmentally dynamic and vulnerable to stress

are also the most susceptible in aging.

Discussion

We have pioneered and validated MIND similarity network analysis as a novel approach to

inferring myelo-architectonic similarity between all cortical areas in an individual rat's brain. Using

this methodological advance, we studied N=47 rat cortical networks to investigate normative

dynamics in cortical similarity across development and aging. These changes, which primarily

involved frontal and hippocampal systems, were examined through MIND analysis of myelin-

sensitive MTR data collected up to four times over each rat's lifespan. In a second experiment,

we tested the hypothesis that early life stress exposure was associated with subsequent

abnormalities of cortical network organization. Repeated maternal separation in the first 20 days

of postnatal life affected the adult similarity of cortical areas, especially frontal and hippocampal

systems that are normatively most dynamic in adolescent development and early brain aging.

Structural MRI similarity analysis is being increasingly used as a measure of brain network

organization in various methodological and experimental contexts ((Sebenius et al., 2024) and

references therein). In general, the interpretation of MIND similarity rests on two key assumptions:

(i) that the correlation or inverse divergence between MRI features in two cortical areas reflects

their cyto-architectonic or myelo-architectonic similarity; and (ii) that cortical areas which are more

architectonically similar are more likely to be axonally inter-connected (Bazinet et al., 2023;

Sebenius et al., 2024). We can therefore think of a structural MRI similarity network as primarily

a map of cortical patterning—an architectome—which is in turn a partial proxy for the map of

axonal wiring—a connectome. To validate rat brain MIND similarity analysis, we tested both these

key assumptions. The results confirmed that cortical areas belonging to the same architectonic

class, or spatially adjacent to each other, tended to have higher MIND similarity, and that higher

MIND similarity between cortical areas was associated with stronger evidence of similarity in

axonal connectivity, based on a prior meta-analysis of tract-tracing studies (Swanson et al.,

2024b). Since the MRI data we used for this analysis were collected using a magnetization

transfer (MTR) sequence that is known to be sensitive to cortical myelination and neuropil density

(Grossman et al., 1994), we can therefore generally interpret MTR-derived MIND (dis-)similarity

as indicative of architectonic differentiation and myelination of the cortex of an individual rat.

To characterize the dynamics of such differentiation across the lifespan, we measured changes

in similarity in early life, defined as PND 20 to 35, and in later life, defined as PND 63 to 230.

Though the data on developmental patterns of rat brain myelination are sparse, a histological

study demonstrated that the rat brain begins myelinating around PND 10, and most areas are

fully myelinated by PND 24 (Downes and Mullins, 2014). Thus, our "early" developmental data

likely reflect the end of early life myelinating processes (e.g., late adolescence) and do not capture

earlier peaks in cortical myelination. We show that increasing MIND similarity in fronto-

hippocampal circuitry in late adolescent development ("early life") is coupled to rapidly decreasing

similarity of these systems in mid-adulthood aging. These data thus support the "last in, first out"

hypothesis of development and aging, in which plastic regions that mature last in early

development (namely, frontal areas related to decision-making and executive function) are more

vulnerable to age-related decline, as previously described in humans (Douaud et al., 2014; Fjell

et al., 2014; Duan et al., 2024). We demonstrate here that rat brain microarchitecture is subject

to the same phenomenon.

Developmentally sensitive fronto-(para)hippocampal circuitry also showed targeted disruptions in

young adult rats who had been exposed to repeated maternal separation. This result aligns with

studies in humans, in which later-developing plastic circuitry also shows increased vulnerability

to disease processes during early development and aging, such as schizophrenia and

Alzheimer's disease, respectively (Douaud et al., 2014; Fjell et al., 2014). The congruence

between developmentally dynamic and environmentally sensitive regions was reinforced

quantitatively, as edges that increased in similarity in development tended to show higher

similarity following RMS (**Fig. S5**). These results are consistent with an accelerated development

hypothesis of early life stress (Callaghan and Tottenham, 2016). Edges that showed higher

similarity in young adulthood following early life stress also demonstrated more rapid divergence

in normative aging. This provides further support for the concept that specific fronto-hippocampal

circuits are developmentally dynamic, vulnerable to age-related decline, and susceptible to

environmental stress (Douaud et al., 2014; Duan et al., 2024). The alignment between human

and rat data underscores the potential of this model system for experimental investigation into the

mechanisms linking early life stress, network development, and aging.

An increase in similarity between two areas (in this case, frontal and hippocampal) does not

necessarily indicate the formation of new axonal connections but likely represents coordinated

changes in (i) myelination of fibers, and/or (ii) microstructural properties, such as synaptic or

dendritic remodeling. The first hypothesis is supported by rat histological data showing that,

between PND 24 and PND 37, myelination occurs exclusively in the fornix and mammillothalamic

tract—pathways that traverse the hippocampus (Downes and Mullins, 2014). It has been argued

theoretically that regions related to memory and learning develop as the adolescent rat ventures

out and requires spatial recognition, a skill less essential earlier in life (Downes and Mullins, 2014).

Consistent with this, a study on rat hippocampal myelination reported the first appearance of

myelinated fibers at PND 17, a significant increase to near-adult levels by PND 25, and full

maturation by PND 60 (Meier et al., 2004).

In addition to myelination, changes in MIND similarity may also arise from coordinated

microstructural reorganization. Extensive cross-modality research highlights hippocampal

plasticity in rodents, especially changes in synaptic density in response to environmental

enrichment or deprivation (Lerch et al., 2011; Stein et al., 2016; Ohline and Abraham, 2019;

Bramati et al., 2023; Dayananda et al., 2023). Similarly, the frontal cortex undergoes

environmentally-sensitive synaptogenesis in early development—continuing into adulthood in rats

(Zeiss, 2021)—and synaptic pruning throughout the lifespan (Kolk and Rakic, 2022). These

shifting neuropil profiles suggest that cortical regions without direct physical connections may

exhibit high MIND similarity due to convergent synaptic architectures—or increasingly divergent

profiles with other regions. Our cross-validation data support this concept. In the tract-tracing

(Swanson et al., 2024b) Jaccard comparison, a subset of edges showed low tract-tracing

similarity but high MIND similarity, predominantly involving hippocampal regions. Excluding

hippocampal edges strengthened the correlation between MIND similarity and tract-tracing

(ρ=0.41; *P*<0.001; **Fig. S4C**). We postulate that the observed high MIND similarity, despite low

axonal connection similarity, may reflect convergent plastic reorganization of the microstructural

properties of these regions. Notably, the hypotheses of cortical fiber myelination and synaptic

density as contributors to MIND similarity are likely interconnected, as evidence suggests neural

activity can induce myelinogenesis (Demerens et al., 1996). Future studies incorporating

histological measures of synaptic density and activity-dependent myelination in the rat brain could

help clarify the relative contributions of these mechanisms to MIND similarity.

A key limitation of this study is that the normative developmental cohort included only male

animals; accordingly, the normative network defined here represents male development only. To

more fully characterize normative development, future work should explicitly investigate sex

differences in network structure and dynamics. It would also be of interest to reapply these

methods using varying parcellation schemes, as different definitions of brain regions—both

cortical and subcortical—may offer complementary or more detailed insights. Future work to

characterize MRI-derived rat brain network architecture could include additional morphometric

features, such as DTI metrics, within a multivariate MIND analysis. This may provide a broader

view of cortex-wide morphological co-variation and more directly reflect axonal connections. As

additional rat brain resources become available (for instance, a brain-wide spatial transcriptomic

atlas and consensus nomenclature/parcellations), our ability to biologically annotate the rat

structural similarity network will continue to improve.

We provide the normative MTR-derived rat cortical microstructural network and all associated

code to support further investigation and understanding of the complex organization of cortical

networks in this key model system. Our results demonstrate the biological validity and replicability

of the MIND similarity analysis and demonstrate its sensitivity to developmental and

environmental stress-related changes in cortical network configuration. We emphasize the

importance and vulnerability of key frontal and hippocampal circuitry in dynamic processes of

normative development and atypical developmental trajectories triggered by early life adversity.

Data and Code Availability: All data were made available through previous publications (Dutcher

et al., 2023; Jones et al., 2024). All visualizations were rendered using the R package ggplot

version 3.5.1 (Wickham, 2016). Code for data preprocessing (including the rat MRI preprocessing

pipeline),   data   analysis,   and   figure   construction   is   available   at

https://github.com/rlsmith1/rat_MRI_similarity_networks.

Author Contributions: *Designed research*: R.L.S., E.G.D., J.A.J., J.W.D., F.J.M., A.R., P.E.V.,

E.T.B., *Performed research*: R.L.S., S.J.S., L.D., E.G.D., J.A.J., *Contributed unpublished*

reagents/analytic tools: S.J.S., J.D.H., O.S., L.W.S., P.A.T., D.R.G., *Analyzed data*: R.L.S., *Wrote*

the paper: R.L.S., S.J.S., L.D., E.G.D., J.A.J., J.D.H., O.S., L.W.S., P.A.T., D.R.G., J.W.D., F.J.M.,

A.R., P.E.V., E.T.B.

Declaration of Competing Interests: E.T.B has consulted for SR One, GSK, Sosei Heptares,

Boehringer Ingelheim, Novartis, and Monument Therapeutics. P.E.V. has consulted for LinusBio.

Acknowledgments

R.L.S completed this work as a PhD candidate in the NIH Oxford-Cambridge Scholars Program.

R.L.S., P.A.T., D.R.G., A.R., and F.J.M NIMH Intramural Research Program of the NIH/HHS, USA

as follows: R.L.S and F.J.M (1ZIAMH002810), A.R. (1ZIAMH002949), P.A.T. and D.R.G

(ZICMH002888). L.D. and E.G.D. were supported by the Gates Cambridge Scholarship. P.E.V.

was supported by MQ: Transforming Mental Health (MQF17_24). This work received

computational support from the NIP HPC Biowulf cluster (http://hpc.nih.gov). All research from

the Department of Psychiatry at the University of Cambridge is made possible by the mental

health theme of the NIHR Cambridge Biomedical Research Centre (BRC-1215-20014) and the

NIHR East of England Applied Research Centre. E.T.B. was also supported by an NIHR Senior

References

Abraham M, Peterburs J, Mundorf A (2023) Oligodendrocytes matter: a review of animal studies
on early adversity. J Neural Transm (Vienna) 130:1177–1185.

Alexander-Bloch A, Giedd JN, Bullmore E (2013) Imaging structural co-variance between
human brain regions. Nat Rev Neurosci 14:322–336.

Bai LS, Kinosada Y, Okuda Y, Ning M, Nakagawa T (1996) Changes of magnetization transfer
ratio according to rat brain development. Nihon Igaku Hoshasen Gakkai Zasshi 56:955–
960.

Bass NH, Netsky MG, Young E (1970) Effect of neonatal malnutrition on developing cerebrum.
II. Microchemical and histologic study of myelin formation in the rat. Arch Neurol 23:303–
313.

Bazinet V, Hansen JY, Vos de Wael R, Bernhardt BC, van den Heuvel MP, Misic B (2023)
Assortative mixing in micro-architecturally annotated brain connectomes. Nat Commun
14:2850.

Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate: a practical and powerful
approach to multiple testing. Journal of the royal statistical society series b-
methodological 57:289–300.

Bramati G, Stauffer P, Nigri M, Wolfer DP, Amrein I (2023) Environmental enrichment improves
hippocampus-dependent spatial learning in female C57BL/6 mice in novel IntelliCage
sweet reward-based behavioral tests. Front Behav Neurosci 17:1256744.

Breton JM, Barraza M, Hu KY, Frias SJ, Long KLP, Kaufer D (2021) Juvenile exposure to acute
traumatic stress leads to long-lasting alterations in grey matter myelination in adult
female but not male rats. Neurobiol Stress 14:100319.

Bryda EC (2013) The Mighty Mouse: the impact of rodents on advances in biomedical research.
Mo Med 110:207–211.

Bullmore E, Sporns O (2009) Complex brain networks: graph theoretical analysis of structural
and functional systems. Nat Rev Neurosci 10:186–198.

Bullmore E, Sporns O (2012) The economy of brain network organization. Nat Rev Neurosci
13:336–349.

Callaghan BL, Tottenham N (2016) The Stress Acceleration Hypothesis: effects of early-life
adversity on emotion circuits and behavior. Current Opinion in Behavioral Sciences
7:76–81.

Cao H, Wei P, Huang Y, Wang N, Guo L-A, Fan X, Wang Z, Ren L, Piao Y, Lu J, Shan Y, He X,
Zhao G (2023) The alteration of cortical microstructure similarity in drug-resistant
epilepsy correlated with mTOR pathway genes. EBioMedicine 97:104847.

Clayden J, Cox B, Jenkinson M, Hall M, Reynolds R, Fissell K, Gailly J-L, Adler M (2024) RNifti:
Fast R and C++ Access to NIfTI Images. Available at: https://cran.r-
project.org/web/packages/RNifti/index.html [Accessed January 30, 2025].

Cox RW (1996) AFNI: Software for Analysis and Visualization of Functional Magnetic
872   Resonance Neuroimages. Comput Biomed Res 29:162–173.

873   Dayananda KK, Ahmed S, Wang D, Polis B, Islam R, Kaffman A (2023) Early life stress impairs
874   synaptic pruning in the developing hippocampus. Brain Behav Immun 107:16–31.

875   Demerens C, Stankoff B, Logak M, Anglade P, Allinquant B, Couraud F, Zalc B, Lubetzki C
876   (1996) Induction of myelination in the central nervous system by electrical activity. Proc
877   Natl Acad Sci U S A 93:9887–9892.

878   Dorfschmidt L, Váša F, White SR, Romero-García R, Kitzbichler MG, Alexander-Bloch A,
879   Cieslak M, Mehta K, Satterthwaite TD, NSPN Consortium, Bethlehem RAI, Seidlitz J,
880   Vértes PE, Bullmore ET (2024) Human adolescent brain similarity development is
881   different for paralimbic versus neocortical zones. Proc Natl Acad Sci U S A
882   121:e2314074121.

883   Douaud G, Groves AR, Tamnes CK, Westlye LT, Duff EP, Engvig A, Walhovd KB, James A,
884   Gass A, Monsch AU, Matthews PM, Fjell AM, Smith SM, Johansen-Berg H (2014) A
885   common brain network links development, aging, and vulnerability to disease. Proc Natl
886   Acad Sci U S A 111:17648–17653.

887   Downes N, Mullins P (2014) The development of myelin in the brain of the juvenile rat. Toxicol
888   Pathol 42:913–922.

889   Duan H et al. (2024) Population clustering of structural brain aging and its association with brain
890   development. Available at:
891   https://www.medrxiv.org/content/10.1101/2024.01.09.24301030v2 [Accessed April 8,
892   2024].

893   Dutcher EG, Lopez-Cruz L, Pama EAC, Lynall M-E, Bevers ICR, Jones JA, Khan S, Sawiak SJ,
894   Milton AL, Clatworthy MR, Robbins TW, Bullmore ET, Dalley JW (2023) Early-life stress
895   biases responding to negative feedback and increases amygdala volume and
896   vulnerability to later-life stress. Transl Psychiatry 13:1–11.

897   Ellenbroek B, Youn J (2016) Rodent models in neuroscience research: is it a rat race? Dis
898   Model Mech 9:1079–1087.

899   Fenchel D et al. (2020) Development of microstructural and morphological cortical profiles in the
900   neonatal brain. Cereb Cortex 30:5767–5779.

901   Fjell AM, McEvoy L, Holland D, Dale AM, Walhovd KB (2014) What is normal in normal aging?
902   Effects of aging, amyloid and Alzheimer's disease on the cerebral cortex and the
903   hippocampus. Prog Neurobiol 117:20–40.

904   Fornito A, Zalesky A, Breakspear M (2015) The connectomics of brain disorders. Nat Rev
905   Neurosci 16:159–172.

906   Fornito A, Zalesky A, Bullmore ET eds. (2016a) Chapter 6 - Components, Cores, and Clubs. In:
907   Fundamentals of Brain Network Analysis, pp 163–206. San Diego: Academic Press.

908   Fornito A, Zalesky A, Bullmore ET eds. (2016b) Chapter 5 - Centrality and Hubs. In:
909   Fundamentals of Brain Network Analysis, pp 137–161. San Diego: Academic Press.

García-Cabezas MÁ, Hacker JL, Zikopoulos B (2023) Homology of neocortical areas in rats and
primates based on cortical type analysis: an update of the Hypothesis on the Dual Origin
of the Neocortex. Brain Struct Funct 228:1069–1093.

González-García N, Buimer EEL, Moreno-López L, Sallie SN, Váša F, Lim S, Romero-Garcia R,
Scheuplein M, Whitaker KJ, Jones PB, Dolan RJ, NSPN consortium, Fonagy P, Goodyer
I, Bullmore ET, van Harmelen A-L (2023) Resilient functioning is associated with altered
structural brain network topology in adolescents exposed to childhood adversity. Dev
Psychopathol 35:2253–2263.

Grossman RI, Gomori JM, Ramer KN, Lexa FJ, Schnall MD (1994) Magnetization transfer:
theory and clinical applications in neuroradiology. Radiographics 14:279–290.

Hahn JD, Duckworth C (2023) A brain flatmap data visualization tool for mouse, rat, and human.
J Comp Neurol 531:1008–1016.

Hahn JD, Swanson LW, Bowman I, Foster NN, Zingg B, Bienkowski MS, Hintiryan H, Dong H-W
(2021) An open access mouse brain flatmap and upgraded rat and human brain
flatmaps based on current reference atlases. J Comp Neurol 529:576–594.

Hamano K, Takeya T, Iwasaki N, Nakayama J, Ohto T, Okada Y (1998) A quantitative study of
the progress of myelination in the rat central nervous system, using the
immunohistochemical method for proteolipid protein. Brain Res Dev Brain Res 108:287–
293.

Han W, Pan Y, Han Z, Cheng L, Jiang L (2022) Advanced maternal age impairs myelination in
offspring rats. Front Pediatr 10:850213.

Herrick CJ (1948) The brain of the tiger salamander, Ambystoma tigrinum. Chicago: Univ. of
Chicago Press.

Hettwer MD, Dorfschmidt L, Puhlmann LMC, Jacob LM, Paquola C, Bethlehem RAI, NSPN
Consortium, Bullmore ET, Eickhoff SB, Valk SL (2024) Longitudinal variation in resilient
psychosocial functioning is associated with ongoing cortical myelination and functional
reorganization during adolescence. Nat Commun 15:6283.

Homan P, Argyelan M, DeRosse P, Szeszko PR, Gallego JA, Hanna L, Robinson DG, Kane JM,
Lencz T, Malhotra AK (2019) Structural similarity networks predict clinical outcome in
early-phase psychosis. Neuropsychopharmacology 44:915–922.

Jones JA, Belin-Rauscent A, Jupp B, Fouyssac M, Sawiak SJ, Zuhlsdorff K, Zhukovsky P,
Hebdon L, Velazquez Sanchez C, Robbins TW, Everitt BJ, Belin D, Dalley JW (2024)
Neurobehavioral Precursors of Compulsive Cocaine Seeking in Dual Frontostriatal
Circuits. Biological Psychiatry Global Open Science 4:194–202.

Jung B, Taylor PA, Seidlitz J, Sponheim C, Perkins P, Ungerleider LG, Glen D, Messinger A
(2021) A comprehensive macaque fMRI pipeline and hierarchical atlas. Neuroimage
235:117997.

Kleven H, Bjerke IE, Clascá F, Groenewegen HJ, Bjaalie JG, Leergaard TB (2023) Waxholm
Space atlas of the rat brain: a 3D atlas supporting data analysis and integration. Nat
Methods 20:1822–1829.

Kolk SM, Rakic P (2022) Development of prefrontal cortex. Neuropsychopharmacology 47:41–
57.

Krigman MR, Hogan EL (1976) Undernutrition in the developing rat: effect upon myelination.
Brain Res 107:239–255.

Lein ES et al. (2007) Genome-wide atlas of gene expression in the adult mouse brain. Nature
445:168–176.

Lerch JP, Yiu AP, Martinez-Canabal A, Pekar T, Bohbot VD, Frankland PW, Henkelman RM,
Josselyn SA, Sled JG (2011) Maze training in mice induces MRI-detectable brain shape
changes specific to the type of learning. Neuroimage 54:2086–2095.

Li W, Yang C, Shi F, Wu S, Wang Q, Nie Y, Zhang X (2017) Construction of individual
morphological brain networks with multiple morphometric features. Front Neuroanat
11:34.

Li X, Wu K, Zhang Y, Kong L, Bertisch H, DeLisi LE (2019) Altered topological characteristics of
morphological brain network relate to language impairment in high genetic risk subjects
and schizophrenia patients. Schizophr Res 208:338–343.

Lisowska A, Rekik I (2019) Joint pairing and structured mapping of convolutional brain
morphological multiplexes for early dementia diagnosis. Brain Connect 9:22–36.

Long KLP, Chao LL, Kazama Y, An A, Hu KY, Peretz L, Muller DCY, Roan VD, Misra R, Toth
CE, Breton JM, Casazza W, Mostafavi S, Huber BR, Woodward SH, Neylan TC, Kaufer
D (2021) Regional gray matter oligodendrocyte- and myelin-related measures are
associated with differential susceptibility to stress-induced behavior in rats and humans.
Transl Psychiatry 11:1–15.

Lupien SJ, McEwen BS, Gunnar MR, Heim C (2009) Effects of stress throughout the lifespan on
the brain, behaviour and cognition. Nat Rev Neurosci 10:434–445.

Mahjoub I, Mahjoub MA, Rekik I, Alzheimer's Disease Neuroimaging Initiative (2018) Brain
multiplexes reveal morphological connectional biomarkers fingerprinting late brain
dementia states. Sci Rep 8:4103.

Mancini M, Karakuzu A, Cohen-Adad J, Cercignani M, Nichols TE, Stikov N (2020) An
interactive meta-analysis of MRI biomarkers of myelin. Elife 9 Available at:
https://elifesciences.org/articles/61523 [Accessed September 26, 2024].

Meier S, Bräuer AU, Heimrich B, Nitsch R, Savaskan NE (2004) Myelination in the hippocampus
during development and following lesion. Cell Mol Life Sci 61:1082–1094.

Mengler L, Khmelinskii A, Diedenhofen M, Po C, Staring M, Lelieveldt BPF, Hoehn M (2014)
Brain maturation of the adolescent rat cortex and striatum: changes in volume and
myelination. Neuroimage 84:35–44.

Morgan SE, Seidlitz J, Whitaker KJ, Romero-Garcia R, Clifton NE, Scarpazza C, van
Amelsvoort T, Marcelis M, van Os J, Donohoe G, Mothersill D, Corvin A, Pocklington A,
Raznahan A, McGuire P, Vértes PE, Bullmore ET (2019) Cortical patterning of abnormal
morphometric similarity in psychosis is associated with brain expression of

schizophrenia-related genes. Proc Natl Acad Sci U S A 116:9604–9609.

Nauta WJH, Karten HJ (1970) A General Profile of the Vertebrate Brain, with Sidelights on the
Ancestry of Cerebral Cortex. In: Neuroanatomy (Nauta WJH, ed), pp 520–539. Boston,
MA: Birkhäuser.

Ohline SM, Abraham WC (2019) Environmental enrichment effects on synaptic and cellular
physiology of hippocampal neurons. Neuropharmacology 145:3–12.

Paus T, Keshavan M, Giedd JN (2008) Why do many psychiatric disorders emerge during
adolescence? Nat Rev Neurosci 9:947–957.

Paxinos G, Watson C (2006) The Rat Brain in Stereotaxic Coordinates: Hard Cover Edition.
Elsevier.

Pebesma E (2018) Simple features for R: Standardized support for spatial vector data. R J
10:439.

Roussel J-R, Auty D, Coops NC, Tompalski P, Goodbody TRH, Meador AS, Bourdon J-F, de
Boissieu F, Achim A (2020) lidR: An R package for analysis of Airborne Laser Scanning
(ALS) data. Remote Sens Environ 251:112061.

Ruan J, Wang N, Li J, Wang J, Zou Q, Lv Y, Zhang H, Wang J (2023) Single-subject cortical
morphological brain networks across the adult lifespan. Hum Brain Mapp 44:5429–5449.

Rubinov M, Ypma RJ, Watson C, Bullmore ET (2015) Wiring cost and topological participation
of the mouse brain connectome. Proceedings of the National Academy of Sciences
112:10032–10037.

Sánchez MM, Ladd CO, Plotsky PM (2001) Early adverse experience as a developmental risk
factor for later psychopathology: evidence from rodent and primate models. Dev
Psychopathol 13:419–449.

Sebenius I, Dorfschmidt L, Seidlitz J, Alexander-Bloch A, Morgan SE, Bullmore E (2024)
Structural MRI of brain similarity networks. Nat Rev Neurosci:1–18.

Sebenius I, Seidlitz J, Warrier V, Bethlehem RAI, Alexander-Bloch A, Mallard TT, Garcia RR,
Bullmore ET, Morgan SE (2023) Robust estimation of cortical similarity networks from
brain MRI. Nat Neurosci 26:1461–1471.

Seidlitz J et al. (2020) Transcriptomic and cellular decoding of regional brain vulnerability to
neurogenetic disorders. Nat Commun 11:3358.

Seidlitz J, Váša F, Shinn M, Romero-Garcia R, Whitaker KJ, Vértes PE, Wagstyl K, Kirkpatrick
Reardon P, Clasen L, Liu S, Messinger A, Leopold DA, Fonagy P, Dolan RJ, Jones PB,
Goodyer IM, Raznahan A, Bullmore ET (2018) Morphometric Similarity Networks Detect
Microscale Cortical Organization and Predict Inter-Individual Cognitive Variation. Neuron
97:231-247.e7.

Stein LR, O'Dell KA, Funatsu M, Zorumski CF, Izumi Y (2016) Short-term environmental
enrichment enhances synaptic plasticity in hippocampal slices from aged rats.
Neuroscience 329:294–305.

Swanson LW (1992) Brain maps: structure of the rat brain, 1st ed.

Swanson LW (2000) 3 - A History of Neuroanatomical Mapping. In: Brain Mapping: The
1029   Systems (Toga AW, Mazziotta JC, eds), pp 77–109. San Diego: Academic Press.

1030   Swanson LW (2018) Brain maps 4.0—Structure of the rat brain: An open access atlas with
1031   global nervous system nomenclature ontology and flatmaps. J Comp Neurol 526:935–
1032   943.

1033   Swanson LW, Hahn JD, Sporns O (2017) Organizing principles for the cerebral cortex network
1034   of commissural and association connections. Proceedings of the National Academy of
1035   Sciences 114:E9692–E9701.

1036   Swanson LW, Hahn JD, Sporns O (2020) Structure–function subsystem models of female and
1037   male forebrain networks integrating cognition, affect, behavior, and bodily functions.
1038   Proceedings of the National Academy of Sciences 117:31470–31481.

1039   Swanson LW, Hahn JD, Sporns O (2022) Structure–function subsystem model and
1040   computational lesions of the central nervous system's rostral sector (forebrain and
1041   midbrain). Proceedings of the National Academy of Sciences 119:e2210931119.

1042   Swanson LW, Hahn JD, Sporns O (2024a) Network architecture of intrinsic connectivity in a
1043   mammalian spinal cord (the central nervous system's caudal sector). Proc Natl Acad Sci
1044   U S A 121:e2320953121.

1045   Swanson LW, Hahn JD, Sporns O (2024b) Neural network architecture of a mammalian brain.
1046   Proc Natl Acad Sci U S A 121:e2413422121.

1047   Swanson LW, Sporns O, Hahn JD (2019) The network architecture of rat intrinsic interbrain
1048   (diencephalic) macroconnections. Proc Natl Acad Sci U S A 116:26991–27000.

1049   Tian T, Li J, Zhang G, Wang J, Liu D, Wan C, Fang J, Wu D, Zhou Y, Zhu W (2021) Effects of
1050   childhood trauma experience and BDNF Val66Met polymorphism on brain plasticity
1051   relate to emotion regulation. Behav Brain Res 398:112949.

1052   Towlson EK, Vértes PE, Ahnert SE, Schafer WR, Bullmore ET (2013) The rich club of the C.
1053   elegans neuronal connectome. J Neurosci 33:6380–6387.

1054   van den Heuvel MP, Sporns O (2011) Rich-club organization of the human connectome. J
1055   Neurosci 31:15775–15786.

1056   van den Heuvel MP, Sporns O (2013) Network hubs in the human brain. Trends Cogn Sci
1057   17:683–696.

1058   Wickham H (2016) Ggplot2: Elegant graphics for data analysis, 2nd ed. Cham, Switzerland:
1059   Springer International Publishing.

1060   Wu X, Palaniyappan L, Yu G, Zhang K, Seidlitz J, Liu Z, Kong X, Schumann G, Feng J,
1061   Sahakian BJ, Robbins TW, Bullmore E, Zhang J (2023) Morphometric dis-similarity
1062   between cortical and subcortical areas underlies cognitive function and psychiatric
1063   symptomatology: a preadolescence study from ABCD. Mol Psychiatry 28:1146–1158.

1064   Xiao Y, Chen F, Lei W, Ke J, Dai Y, Qi R, Lu G, Zhong Y (2023) Transcriptional signal and cell

1065   specificity of genes related to cortical structural differences of post-traumatic stress
1066   disorder. J Psychiatr Res 160:28–37.

1067   Yao Z et al. (2023) A high-resolution transcriptomic and spatial atlas of cell types in the whole
1068   mouse brain. Nature 624:317–332.

1069   Zeiss CJ (2021) Comparative milestones in rodent and human postnatal central nervous system
1070   development. Toxicol Pathol 49:1368–1373.

1071   Zhang W, Lei D, Keedy SK, Ivleva EI, Eum S, Yao L, Tamminga CA, Clementz BA, Keshavan
1072   MS, Pearlson GD, Gershon ES, Bishop JR, Gong Q, Lui S, Sweeney JA (2020) Brain
1073   gray matter network organization in psychotic disorders. Neuropsychopharmacology
1074   45:666–674.

1075   Zilles K (2012) The Cortex of the Rat: A Stereotaxic Atlas. Springer Science & Business Media.