1   A continuation of the assembly of a G-quadruplex repair complex by the

2   FANCJ DNA helicase

5   Laura Campbell[1,#a], Kaitlin Lowran[1,#b], Emma Cismas[1], Colin G. Wu[1*]

7   [1] Department of Chemistry, Oakland University, Rochester, Michigan, United States of

8   America

9   [#a] Current Address: Department of Biochemistry, University of Wisconsin-Madison,

10   Madison, Wisconsin, United States of America

11   [#b] Current Address: Institute of Environmental Health Sciences, Wayne State University,

12   Detroit, Michigan, United States of America

14   * Corresponding author

15   E-mail: colinwu@oakland.edu (CGW)

Abstract

Nucleic acid sequences that are rich in guanines can form G-quadruplex (G4) structures, which can impede DNA replication or repair. The FANCJ helicase plays a role in maintaining genomic stability by facilitating DNA replication through regions of DNA that form G4 structures. This activity has been associated with an AKKQ motif in FANCJ and it is thought that this motif retains the ability to also target 8-oxoguanine-modified G4 (8oxoG4) structures. We hypothesize that the molecular recognition of FANCJ AKKQ to 8oxoG4s is dependent upon the position of DNA damage, sequence composition, and substrate stability. We previously demonstrated that reducing the loop length between guanine tetrads increased the affinity of FANCJ AKKQ when 8-oxoguanines (8oxoGs) were located at the first and fifth positions of the G4 structure. In this work, we use fluorescence spectroscopy to measure the affinities of a FANCJ AKKQ peptide for G4s formed by $(GGGT)_4$, $(GGGTT)_4$, and $(TTAGGG)_4$. G4 conformations and thermal stability were determined via circular dichroism spectroscopy. We substituted 8oxoGs at the first (8oxo1), third (8oxo3), and fifth (8oxo5) positions of each G4 sequence and found that the most destabilized structures were $(GGGT)_4$ 8oxo1, $(GGGTT)_4$ 8oxo1, and $(TTAGGG)_4$ 8oxo5 in KCl. The most destabilized structures in NaCl were $(GGGT)_4$ 8oxo1, $(GGGTT)_4$ 8oxo5, and $(TTAGGG)_4$ 8oxo5. The binding affinities of FANCJ AKKQ differed for the distinct damage positions for $(GGGT)_4$ in KCl and for $(TTAGGG)_4$ in NaCl. These findings

36 suggest that FANCJ AKKQ targets G4s and 8oxoG4s in a manner dependent upon

37 sequence composition and damage position.

Introduction

G-quadruplexes

G-quadruplexes (G4s) are stable secondary DNA structures found in guanine-rich

sequences. These structures arise via intramolecular Hoogsteen hydrogen bonding

reactions in single-stranded (ss) DNA that contains four stretches of three or more

consecutive guanines [1]. Monovalent cations further stabilize these interactions by

centrally coordinating the G4 structure. The two most biologically relevant cations are

potassium and sodium, with potassium ions generally having a greater impact on

stability [2-4].

The general sequence for G4s has been accepted as $(GGGN_{1-7})_4$, with N

representing any DNA base(s) [5]. This has led to an estimate of over 400,000 G4 motifs

being identified in the human genome [6]. Depending on solution conditions and

intervening loop sequence, G4s can adopt three main spatial conformations: parallel,

antiparallel, and hybrid (Fig 1) [4, 7]. These conformations are classified based on how

the intervening sequence, N_{1-7}, links the guanine tetrads [8, 9]. Circular dichroism (CD)

spectroscopy can be used to observe the optical properties of these conformations [10].

The respective spatial arrangements are important as certain G4-binding proteins have

preference or specificity for specific conformations; for example, the Pif1 helicase was

shown to preferentially unfold antiparallel G4s [11-15].

G4s are abundant in regions associated with genome regulation [5, 16, 17]. Due to

this accumulation, G4s are thought to play a role in translation regulation, splicing

modulation, and telomere capping [18- 20]. However, despite this apparent involvement,

an accumulation of G4s can be toxic to cells as they stall DNA repair, replication, and

RNA transcription [16]. The regulation of G4 formation has consequently emerged as a

novel therapeutic target for diseases. G4 stabilizers such as CX-5461 take advantage of

this fact by inducing replication fork blockage and single-stranded gaps or breaks [21].

An additional problem present in G4s is the susceptibility of guanine to reactive

oxygen species. Guanine has the lowest oxidation potential of all DNA base pairs and is

the most susceptible to oxidative damage from reactive oxygen species [22]. Consecutive

runs or stacking of guanines further lowers their oxidation potential [23]. When guanine

undergoes oxidative damage, it is commonly known as 8-oxoguanine (8oxoG). Due to

the ability of 8oxoG to pair with both adenine and cytosine, 8oxoGs can cause guanine-

to-thymine transversion mutations if left unrepaired [24, 25]. While 8oxoG-modified G4s

(8oxoG4s) can still form stable structures, the damage can cause a decrease in stability,

ultimately leaving the DNA vulnerable to cellular nucleases and causing untimely

shortening [26, 27]. It was also shown, via NMR, that the incorporation of 8oxoG into the

human telomeric G4 changed the G4 conformation by converting guanine glycosidic angles from *anti* to *syn* conformations [26].

FANCJ

Fanconi anemia complementation group J (FANCJ) is an iron-sulfur helicase that participates in the human DNA cross-link repair pathway [28]. Mutations in *FANCJ* can lead to an increased risk of cancer and the development of Fanconi anemia [28-30]. Previous studies have shown that FANCJ coordinates the repair of DNA intermediates arising from the nucleotide excision repair pathway [31-33]. Specifically, FANCJ binds to G4-rich regions and supports DNA replication by unwinding G4s [31, 34]. When a polymerase is stalled by G4 obstructions, FANCJ is able to bind the G4 and unfold the structure [35]. FANCJ can then recruit another polymerase, such as the translesion polymerase REV1, to continue with replication (Fig 2) [36].

An unrelated RNA helicase and G4 binding protein, RHAU, targets G4s and 8oxoG4s through an AKKQ motif. In this motif, the two lysine residues are readily available for DNA binding [11, 37]. FANCJ has also been identified to contain a similar AKKQ motif at K141/K142 [33]. This motif (Fig 3) allows FANCJ to bind and anchor to G4s and 8oxoG4s [36]. In this work, we examine the interactions of FANCJ AKKQ with G4s and 8oxoG4s of varying intervening sequence lengths and position of damage. These interactions are related to G4 conformation and thermal stability to further characterize

93   the recognition of G4 and 8oxoG4 sequences by FANCJ AKKQ. Previous data is included

94   from our study of (GGGT)$_4$ and (TTAGGG)$_4$; we expand upon this work here by

95   introducing a third sequence, (GGGTT)$_4$, with 8oxoGs systematically substituted at the

96   first, third, and fifth guanines of each G4 sequence.

Results

G-quadruplex conformation and thermal stability

CD spectroscopy was used to determine the conformation and thermal stability of

twelve G4 sequences in KCl and NaCl solutions. G4 structures have distinct optical

properties that give unique CD intensity signals (Fig 4A) [9, 10]. Moreover, CD

spectroscopy was used to determine the melting temperature (T_m) of G4 structures [38].

As temperature increases, the G4 structure unfolds, and the intensity of spectral peaks

and troughs decreases (Fig 4B). The change in intensity can be analyzed as a function of

temperature to determine the T_m, which is a direct indicator of thermal stability (Fig 4C).

The three undamaged G4 sequences were chosen based on intervening sequence

length and biological relevance. (GGGT)$_4$ is an HIV-1 integrase aptamer and contains a

single thymine base for the loop [7, 39]. G4 sequences that contain only single-residue

loops are shown to adopt a compact, parallel conformation, with thymine having the

most stability [7, 40]. (TTAGGG)$_4$ is commonly referred to as the human telomeric

sequence due to its repetition at the ends of human chromosomes and its protective role

in maintaining telomere length [19]. (GGGTT)$_4$ is an artificial sequence designed to have

a loop of intermediate length.

An 8oxoG-modified base was substituted for the first (8oxo1), third (8oxo3), or

fifth (8oxo5) guanine position of each G4 sequence (Fig 5). These positions were chosen

as previous studies for the human telomeric sequence showed that 8oxo5 was the most

thermally destabilized, 8oxo1 was the least destabilized, and 8oxo3 had an intermediate

level of destabilization [27]. Furthermore, this places an 8oxoG in each of the three tetrads

of a G4 structure.

We previously studied the G4 conformations and melting temperatures of the

(GGGT)$_4$ and (TTAGGG)$_4$ sequences with oxidative damage induced at the 8oxo1 and

8oxo5 positions [36]. This data was performed in both KCl- and NaCl-containing buffers

and is included in this manuscript for reference. In this study, we introduce damage to

the 8oxo3 position of the (GGGT)$_4$ and (TTAGGG)$_4$ sequences and study all three

positions of oxidative damage in the (GGGTT)$_4$ sequence.

All (GGGT)$_4$ sequences formed a parallel conformation in KCl (Fig 6A). Based on

the overlaying CD spectra traces, 8oxoG-modified (GGGT)$_4$ sequences did not show

significant structural changes. Due to the limitation of the machine, 95 °C was the

maximum temperature reached. As evident in Fig 6B, (GGGT)$_4$ is not fully unfolded at 95

°C and the T$_m$ can be assumed to be > 88.4 ± 8.4 °C (Table 1). This assumption is in

agreement with previous studies [7]. Each T_m value for the $(GGGT)_4$ 8oxoG4s showed a

decrease from the native sequence, with $(GGGT)_4$ 8oxo1 having the highest decrease in

thermal stability of ΔT_m = -19.0 ± 8.4 °C and $(GGGT)_4$ 8oxo3 having the lowest change of

ΔT_m = -8.1 ± 8.4 °C.

$(GGGTT)_4$ also formed a parallel conformation in KCl (Fig 6C). $(GGGTT)_4$ 8oxo1

and $(GGGTT)_4$ 8oxo5 sequences similarly formed parallel conformations, yet with slight

structural changes, as evident by differences in the CD spectra traces from 300 to 280 nm.

Conversely, the $(GGGTT)_4$ 8oxo3 sequence changed $(GGGTT)_4$ to an antiparallel

sequence. The T_m of $(GGGTT)_4$ was determined to be 83.5 ± 0.7 °C (Fig 6D; Table 1). Similar

to the $(GGGT)_4$ sequences in KCl, $(GGGTT)_4$ 8oxo1 had the highest decrease in thermal

stability of ΔT_m = -31.5 ± 1.6 °C while $(GGGTT)_4$ 8oxo3 had the lowest change of ΔT_m = -

26.3 ± 0.7 °C.

$(TTAGGG)_4$ formed a hybrid conformation in KCl (Fig 6E). $(TTAGGG)_4$ 8oxo1 and

$(TTAGGG)_4$ 8oxo5 sequences also formed hybrid conformations, although with

significant structural changes. $(TTAGGG)_4$ 8oxo1 had a significant right shift in the 295

nm peak and a double trough at 260 and 240 nm. $(TTAGGG)_4$ 8oxo5 experienced

depression in its peak at 295 nm and an increase in its shoulder at 260 nm and trough at

240 nm. Similar to $(GGGTT)_4$ 8oxo3, the $(TTAGGG)_4$ 8oxo3 sequence changed to the

antiparallel conformation. $(TTAGGG)_4$ had a T_m of 62.5 ± 0.5 °C (Fig 6F; Table 1).

Concurrent with previous literature, $(TTAGGG)_4$ 8oxo5 had the highest decrease in

thermal stability of $\Delta T_m = -23.6 \pm 0.8$ °C while (TTAGGG)$_4$ 8oxo1 had the lowest change in

thermal stability of $\Delta T_m = -8.0 \pm 0.7$ °C [27].

Table 1. Summary of G4 conformation and melting temperature in 150 mM KCl

Name	Conformation	T_m (°C)	ΔT_m (°C)
(GGGT)$_4$	Parallel	> 88.4 ± 8.4	-
(GGGT)$_4$ 8oxo1	Parallel	69.4 ± 0.6	-19.0 ± 8.4
(GGGT)$_4$ 8oxo3	Parallel	80.3 ± 0.4	-8.1 ± 8.4
(GGGT)$_4$ 8oxo5	Parallel	71.6 ± 0.4	-16.8 ± 8.4
(GGGTT)$_4$	Parallel	83.5 ± 0.7	-
(GGGTT)$_4$ 8oxo1	Parallel	52.0 ± 1.4	-31.5 ± 1.6
(GGGTT)$_4$ 8oxo3	Antiparallel	57.2 ± 0.3	-26.3 ± 0.7
(GGGTT)$_4$ 8oxo5	Parallel	55.9 ± 0.3	-27.6 ± 0.8
(TTAGGG)$_4$	Hybrid	62.5 ± 0.5	-
(TTAGGG)$_4$ 8oxo1	Hybrid-esque	54.5 ± 0.5	-8.0 ± 0.7
(TTAGGG)$_4$ 8oxo3	Antiparallel	49.5 ± 0.3	-13.0 ± 0.6
(TTAGGG)$_4$ 8oxo5	Hybrid-esque	38.9 ± 0.6	-23.6 ± 0.8

As in KCl, all (GGGT)$_4$ sequences formed parallel conformations in NaCl (Fig 7A).

However, all three (GGGT)$_4$ 8oxoG4 sequences experienced a slight change in structure,

as indicated by the differences in traces in the 300 to 280 nm region. (GGGT)$_4$ was

determined to have a T_m of 70.3 ± 0.7 °C (Fig 7B; Table 2). Similar to (GGGT)$_4$ in KCl,

(GGGT)$_4$ 8oxo1 had the highest decrease in thermal stability of ΔT_m = -27.8 ± 0.9 °C while

(GGGT)$_4$ 8oxo3 had the lowest change of ΔT_m = -13.9 ± 0.8 °C.

The native (GGGTT)$_4$ also formed a parallel conformation in NaCl (Fig 7C).

However, all (GGGTT)$_4$ 8oxoG4 sequences experienced a conformational change.

(GGGTT)$_4$ 8oxo1 and (GGGTT)$_4$ 8oxo3 formed antiparallel conformations, while

(GGGTT)$_4$ 8oxo5 formed a hybrid conformation. (GGGTT)$_4$ had a T_m of 44.8 ± 1.2 °C (Fig

7D; Table 2). (GGGTT)$_4$ 8oxo1 and (GGGTT)$_4$ 8oxo3 did not show a significant change in

thermal stability, with (GGGTT)$_4$ 8oxo1 even having an increase of ΔT_m = 2.4 ± 1.2 °C.

(GGGTT)$_4$ 8oxo5 had a significant change of ΔT_m = -17.3 ± 1.4 °C. The (GGGTT)$_4$ 8oxo5

thermal melt was taken over -5 to 75 °C because it did not appear fully folded at 15 °C.

(TTAGGG)$_4$ and its 8oxoG4 sequences formed antiparallel conformations in NaCl

(Fig 7E). All (TTAGGG)$_4$ 8oxoG4 sequences showed slight differences in intensities and

peak shifts from the spectra of the native sequence, indicating slight structural

differences. The T_m of (TTAGGG)$_4$ was determined to be 52.1 ± 0.3 °C (Fig 7F; Table 2).

Concurrent with previous literature and the KCl data above, (TTAGGG)$_4$ 8oxo5 had the

highest decrease in thermal stability of ΔT_m = -18.8 ± 0.5 °C while (TTAGGG)$_4$ 8oxo1 had

the lowest of ΔT_m = -4.3 ± 0.7 °C [27].

Table 2. Summary of G4 conformation and melting temperature in 150 mM NaCl

Name	Conformation	T_m (°C)	$\Delta\ T_m$ (°C)
(GGGT)$_4$	Parallel	70.3 ± 0.7	-
(GGGT)$_4$ 8oxo1	Parallel	42.5 ± 0.5	-27.8 ± 0.9
(GGGT)$_4$ 8oxo3	Parallel	56.4 ± 0.4	-13.9 ± 0.8
(GGGT)$_4$ 8oxo5	Parallel	44.0 ± 0.3	-26.3 ± 0.8
(GGGTT)$_4$	Parallel	44.8 ± 1.2	-
(GGGTT)$_4$ 8oxo1	Antiparallel	47.2 ± 0.3	2.4 ± 1.2
(GGGTT)$_4$ 8oxo3	Antiparallel	44.6 ± 0.2	-0.2 ± 1.2
(GGGTT)$_4$ 8oxo5	Hybrid-esque	27.5 ± 0.9	-17.3 ± 1.4
(TTAGGG)$_4$	Antiparallel	52.1 ± 0.3	-
(TTAGGG)$_4$ 8oxo1	Antiparallel	47.8 ± 0.6	-4.3 ± 0.7
(TTAGGG)$_4$ 8oxo3	Antiparallel	42.2 ± 0.7	-9.9 ± 0.8
(TTAGGG)$_4$ 8oxo5	Antiparallel	33.3 ± 0.4	-18.8 ± 0.5

FANCJ AKKQ binding

179   Fluorescence spectroscopy titrations were used to determine the interaction

180   strength between AKKQ and G4 sequences. The AKKQ sequence examined was

181   modified to contain a tryptophan residue, W, at the C-terminal end of the sequence (Fig

182   8A). Due to its aromatic properties, tryptophan produces a fluorescence peak from 300 to

450 nm. When bound to DNA, this signal is quenched, and the binding strength can be determined by the decrease in fluorescence intensity (Fig 8B). This decrease in the integrated peak was plotted against increasing DNA concentration, and the resulting curve fit to a 1:1 equilibrium binding interaction model using a nonlinear least squared analysis (Fig 8C). The resulting K value represents an equilibrium K for forming the protein-DNA complex.

Previously, we demonstrated that shortening the intervening loop length increased the affinity of FANCJ AKKQ when 8oxoGs were located at the 8oxo1 and 8oxo5 positions of the G4 structure [36]; however, our study was limited to the $(GGGT)_4$ and $(TTAGGG)_4$ sequences in KCl-containing buffer. We include this previous data for comparison. In this study, we additionally study the $(GGGT)_4$ sequence with oxidative damage induced at the 8oxo1, 8oxo3, and 8oxo5 positions of all three G4 sequences. We also perform these binding analyses in NaCl-containing buffer.

Binding interactions between FANCJ AKKQ and the G4 sequences were first analyzed in buffer containing 150 mM KCl (Fig 9). Binding affinities are summarized in Table 3. For the $(GGGT)_4$ sequences, AKKQ had the strongest interaction with $(GGGT)_4$ 8oxo1 ($K = 7.7 \pm 0.7 \times 10^5$ M^{-1}), followed by $(GGGT)_4$ 8oxo5 ($K = 4.4 \pm 0.6 \times 10^5$ M^{-1}), then $(GGGT)_4$ 8oxo3 ($K = 3.2 \pm 0.1 \times 10^5$ M^{-1}), and lastly the undamaged native sequence ($K = 2.7 \pm 0.4 \times 10^5$ M^{-1}). AKKQ bound strongest to the native $(GGGTT)_4$ sequence ($K = 6.0 \pm 0.9 \times 10^5$ M^{-1}) and weakest to the respective $(GGGTT)_4$ 8oxoG4s ($K = 8-9 \times 10^5$ M^{-1}). No

significant changes in interaction were observed between the 8oxo1, 8oxo3, and 8oxo5

damage positions for the (GGGTT)$_4$ sequence. AKKQ displayed no preference for the

8oxoG4 damage positions of (TTAGGG)$_4$ and bound with similar affinity to the native

sequence. Each K value for the (TTAGGG)$_4$ sequences was within error of 1.3-1.4 x 10^6 M^{-1}.

Table 3. Summary of FANCJ AKKQ-G4 binding by fluorescence spectroscopy in 150 mM KCl

Name	K (M^{-1})	R^2
(GGGT)$_4$	2.7 ± 0.4 x 10^5	0.997
(GGGT)$_4$ 8oxo1	7.7 ± 0.7 x 10^5	0.998
(GGGT)$_4$ 8oxo3	3.2 ± 0.1 x 10^5	0.997
(GGGT)$_4$ 8oxo5	4.4 ± 0.6 x 10^5	0.997
(GGGTT)$_4$	6.0 ± 0.9 x 10^5	0.997
(GGGTT)$_4$ 8oxo1	8.2 ± 1.0 x 10^5	0.997
(GGGTT)$_4$ 8oxo3	9.6 ± 1.0 x 10^5	0.996
(GGGTT)$_4$ 8oxo5	9.3 ± 0.8 x 10^5	0.997
(TTAGGG)$_4$	1.6 ± 0.7 x 10^6	0.996
(TTAGGG)$_4$ 8oxo1	1.3 ± 0.4 x 10^6	0.997
(TTAGGG)$_4$ 8oxo3	1.2 ± 0.1 x 10^6	0.996

(TTAGGG)$_4$ 8oxo5	$1.4 \pm 0.5 \times 10^6$	0.997

These binding experiments were repeated in buffer containing 150 mM NaCl (Fig 10). Binding affinities are summarized in Table 4. As in KCl, FANCJ AKKQ showed a preference for specific 8oxoG damage positions in the (GGGT)$_4$ sequences. AKKQ bound strongest to (GGGT)$_4$ 8oxo5 and (GGGT)$_4$ 8oxo1 (K= $3.4 \pm 0.5 \times 10^5$ M^{-1} for both), followed by (GGGT)$_4$ 8oxo3 (K = $2.9 \pm 0.5 \times 10^5$ M^{-1}), and lastly the native sequence (K = $2.3 \pm 0.3 \times 10^5$ M^{-1}). AKKQ bound with similar affinity to the native (GGGTT)$_4$ sequence, (GGGTT)$_4$ 8oxo5, and (GGGTT)$_4$ 8oxo3 (K = 4.9-5.6 $\times 10^5$ M^{-1}), with the weakest affinity for (GGGTT)$_4$ 8oxo1 (K = $3.9 \pm 1.3 \times 10^5$ M^{-1}). For the (TTAGGG)$_4$ sequences, AKKQ bound strongest to (TTAGGG)$_4$ 8oxo5 (K = $1.5 \pm 0.1 \times 10^6$ M^{-1}), followed by the native sequence and (TTAGGG)$_4$ 8oxo1 (K = 6.5×10^5 M^{-5} for both), and lastly (TTAGGG)$_4$ 8oxo3 (K = $5.1 \pm 1.1 \times 10^5$ M^{-1}).

Table 4. Summary of FANCJ AKKQ-G4 binding by fluorescence spectroscopy in 150 mM NaCl

Name	K (M^{-1})	R^2
(GGGT)$_4$	$2.3 \pm 0.3 \times 10^5$	0.9997
(GGGT)$_4$ 8oxo1	$3.4 \pm 0.5 \times 10^5$	0.9996
(GGGT)$_4$ 8oxo3	$2.9 \pm 0.5 \times 10^5$	0.9994
(GGGT)$_4$ 8oxo5	$3.4 \pm 0.5 \times 10^5$	0.9997

(GGGTT)$_4$	$4.9 \pm 2.0 \times 10^5$	0.9995
(GGGTT)$_4$ 8oxo1	$3.9 \pm 1.3 \times 10^5$	0.9993
(GGGTT)$_4$ 8oxo3	$5.2 \pm 1.7 \times 10^5$	0.9994
(GGGTT)$_4$ 8oxo5	$5.6 \pm 1.1 \times 10^5$	0.9992
(TTAGGG)$_4$	$6.5 \pm 0.6 \times 10^5$	0.9991
(TTAGGG)$_4$ 8oxo1	$6.5 \pm 1.4 \times 10^5$	0.9996
(TTAGGG)$_4$ 8oxo3	$5.1 \pm 1.1 \times 10^5$	0.9998
(TTAGGG)$_4$ 8oxo5	$1.1 \pm 0.1 \times 10^6$	0.9974

Discussion

G-quadruplexes

The CD data for the human telomeric sequence is concurrent with previous studies in that (TTAGGG)$_4$ and its respective 8oxoG4 sequences form hybrid and antiparallel conformations in KCl and antiparallel conformations in NaCl [26, 27, 41]. In both KCl and NaCl, the (TTAGGG)$_4$ 8oxo5 sequence showed the most thermal instability, while (TTAGGG)$_4$ 8oxo1 showed the least [27]. This pattern was also observed with the (GGGTT)$_4$ 8oxoG-modified sequences in NaCl. While (GGGTT)$_4$ formed a parallel conformation, its 8oxoG4 analogs formed antiparallel or hybrid conformations. Conversely, a different pattern of stability was observed in the other substrates. (GGGT)$_4$

sequences in KCl and NaCl and (GGGTT)$_4$ sequences in KCl formed predominately

parallel conformations, with only (GGGTT)$_4$ 8oxo3 forming an antiparallel conformation.

Of these sequences, the 8oxo1 analogs had the most thermal instability, while the 8oxo3

analogs had the least.

Our results indicate that the effect of an 8oxoG on G4 thermal stability is

dependent upon the location of damage and the G4 structure. Sequences forming

compact parallel conformations were most destabilized by damage to the first guanine,

which resides in the bottom tetrad of the G4 structure, and were least destabilized by

damage to the third guanine, which resides in the top tetrad. Sequences forming the more

flexible antiparallel or hybrid conformations were most destabilized by damage to the

fifth guanine, which resides in the middle tetrad, and were least destabilized by damage

to the first guanine. A similar destabilizing effect had already been observed in the hybrid

conformation of the human telomeric G4 sequence when an 8oxoG was substituted at the

fifth guanine position [26].

Potassium and sodium ions are the most physiologically relevant monovalent

cations and are commonly studied in relation to G4 conformation and stability. While the

(GGGT)$_4$ sequences formed parallel conformations in both KCl and NaCl, most (GGGTT)$_4$

and (TTAGGG)$_4$ sequences folded into other conformations. The thermal stability for all

sequences was lower in NaCl than in KCl, suggesting that sodium ions are less effective

in stabilizing G4 structures. With only sodium ions present, G4 sequences cannot adopt

the rigid conformations that occur in the presence of potassium ions and thus form less

stable conformations. These findings support previous studies, suggesting that while

both ions can stabilize G4s, potassium ions are better suited due to their optimal size and

ionic strength [42].

It is of note that in KCl, there was a correlation between the length of the

intervening sequence and the thermal stability of the native sequences. (GGGT)$_4$ had the

highest melting temperature while (TTAGGG)$_4$ had the lowest. This indicates that as

intervening sequence length increases, thermal stability decreases. This observation was

present yet less evident in NaCl. Although (GGGT)$_4$ maintained the highest melting

temperature, (GGGTT)$_4$ had the lowest. These observations are in agreement with

previous studies [4, 40].

264 FANCJ AKKQ and G-quadruplex binding

Our results indicate that FANCJ AKKQ has the ability to target both G4s and

8oxoG4s; however, this recognition depends on the sequence composition and position

of damage. For the (GGGT)$_4$ sequences, which formed compact parallel conformations,

AKKQ bound strongest to the 8oxoG4 sequences rather than the native sequence. The

observed increase in binding affinity across the 8oxoG damage positions correlates with

a decrease in thermal stability. (GGGT)$_4$ 8oxo1 had the greatest decrease in thermal

stability and the strongest interaction with AKKQ. Conversely, (GGGT)$_4$ 8oxo3 had the

lowest decrease in thermal stability and the weakest interaction with AKKQ. As the CD

spectra displayed no change in structural stability (Fig 6A), this implies that AKKQ is

targeting the 8oxoG position rather than the intervening loop.

For the $(GGGTT)_4$ sequences, which formed predominantly parallel

conformations, AKKQ preferentially bound to the 8oxoG4 sequences with no significant

changes in affinity between each damage position. Assuming that AKKQ is targeting the

intervening sequence, the preference for the 8oxoG4s can be attributed to the

destabilization of the G4 structure. Though $(GGGTT)_4$ 8oxo1 and 8oxo5 still formed

parallel conformations, the CD spectra indicated destabilization in the 280 to 300 nm

range (Fig 6C). This destabilization may have led to the intervening sequence being more

exposed than in an undamaged parallel conformation, thus facilitating better access for

AKKQ.

AKKQ bound with similar affinity to all $(TTAGGG)_4$ sequences, which form the

less rigid hybrid or antiparallel conformations and thus expose the DNA loop. This

implies that AKKQ preferentially targets the intervening sequence and is not affected by

oxidative damage in $(TTAGGG)_4$. It is of note that in KCl, FANCJ AKKQ bound to the

$(TTAGGG)_4$ sequences with an order of magnitude higher than the other G4 sequences.

However, upon changing the salt to NaCl, AKKQ bound equally to all sequences except

for the $(TTAGGG)_4$ 8oxo5 sequence, to which it bound an order of magnitude stronger.

One possible explanation for the observed differences in binding preferences is that high sodium concentrations affect the electrostatic interactions between the peptide and the G4s. Sodium ions are known to provide less stabilization to G4s compared to potassium [2-4]. We anticipated the FANCJ AKKQ peptide to adopt an alpha helical structure due to previous NMR studies of the structurally and sequentially similar AKKQ motif of the G4-binding protein RHAU [11]. Although our CD studies (Supporting information) did not detect any alpha helical structure in either salt, the addition of trifluoroethanol revealed a helical-like structure in both salt conditions. However, the spectra obtained in each salt were not superimposable. This suggests that sodium ions stabilize the peptide to a different extent than potassium ions.

As FANCJ is a DNA repair helicase, it was anticipated that FANCJ AKKQ would bind preferentially to 8oxoGs to facilitate their repair. However, a preference for the 8oxoG4 sequences was only observed for the $(GGGT)_4$ sequences, suggesting that FANCJ AKKQ may instead target the intervening sequence. This is further supported by the positive correlation between binding interaction and intervening sequence length in the undamaged G4 structures. A proposed interaction model for AKKQ and 8oxoG4s is illustrated in Fig 11.

Previous work has demonstrated that FANCJ collaborates with the translesion polymerase REV1, which unwinds G4 structures, for the successful replication of G4-rich DNA strands [43]. It is possible that FANCJ and REV1 are responsible for repairing

distinct types of G4 DNA. FANCJ appears to prefer G4 structures with exposed loops,

while REV1 may prefer more compact G4 structures. Future work should aim to

characterize the interaction of REV1 with G4s and 8oxoG4s. Furthermore, G4 damage

positions were limited to the first, third, and fifth guanine in this study; it should be

observed whether the trends identified in this study persist with systematic substitution

of 8oxoGs for each G4 damage position.

Materials and Methods

Buffers and reagents

319   All buffers and reagents were prepared with reagent-grade chemicals and Type I

320 water purified with a Smart2Pure 6 UV/UF system (ThermoFisher, Waltham, MA).

321 Solutions were sterilized through a 0.22 μm PES filter.

Peptide and DNA oligos

FANCJ AKKQ peptide (129-PEKTTLAAKLSAKKQASIW-147) was synthesized

by Genscript (Piscataway, NJ), and the powder was dissolved into 20 mM HEPES pH 7.5,

150 mM KCl or NaCl, 5 mM TCEP, and 5% (v/v) glycerol. Protein concentration was

determined using a NanoDrop One UV-Vis microvolume spectrophotometer

(ThermoFisher) and the molar extinction coefficients below. Peptide samples were

aliquoted, frozen, and stored at -20 °C. Oligodeoxyribonucleotides were purchased from

Integrated DNA Technologies (IDT, Coralville, IA) and concentrations were determined

similarly to the above stated. DNA substrates were stored at 4 °C. A list of sequences used

is described in Table 5.

Table 5. DNA Sequences.

Name	Sequence (5′→3′)	Molar Extinction Coefficient ($M^{-1}cm^{-1}$)	Molecular Weight (g/mol)
(GGGT)4	GGGTGGGTGGGTGGGT	157200	5105.3
(GGGT)4 8oxo1	T/i8oxodG/GGTGGGTGGGTGGGT	159100	5425.5
(GGGT)4 8oxo3	GG/i8oxodG/TGGGTGGGTGGGT	151600	5121.3
(GGGT)4 8oxo5	GGGTG/i8oxodG/GTGGGTGGGT	151600	5121.3
(GGGTT)4	GGGTTGGGTTGGGTTGGGTT	189600	6322.1
(GGGTT)4 8oxo1	/i8oxodG/GGTTGGGTTGGGTTGGGTT	184000	6338.1

(GGGTT)₄ 8oxo3	GG/i8oxodG/TTGGGTGGGTTGGGTT	184000	6338.1
(GGGTT)₄ 8oxo5	GGGTTG/i8oxodG/GTTGGGTTGGGTT	184000	6338.1
(TTAGGG)₄	TTAGGGTTAGGGTTAGGGTTAGGG	244600	7575.0
(TTAGGG)₄ 8oxo1	TTA/i8oxodG/GGTTAGGGTTAGGGTTA GGG	239000	7590.9
(TTAGGG)₄ 8oxo3	TTAGG/i8oxodG/TTAGGGTTAGGGTTA GGG	239000	7590.9
(TTAGGG)₄ 8oxo5	TTAGGGTTAG/i8oxodG/GTTAGGGTTA GGG	239000	7590.9

CD spectroscopy

CD measurements were performed on a JASCO J-810 spectropolarimeter

equipped with a PTC-423S peltier system and Koolance Liquid Cooling System. G4 DNA

samples were dialyzed into 20 mM boric acid, 150 mM KCl or NaCl. CD spectra scans

were taken from 220 nm to 320 nm at 25 °C with DNA samples of 0.3 mg/mL. Five traces

for each DNA sample were taken and averaged. A reference scan of buffer alone was

subtracted from the averaged scan. Temperature melt scans were taken from 15 °C to 95

°C at 262 nm (for parallel conformations) or 295 nm (for antiparallel and hybrid

conformation). Samples were heated to 95 °C and allowed to cool overnight prior to the

first melt. Three trials for each DNA sample were taken and averaged.

Melting temperature was determined using SigmaPlot as an analysis program and

the model described below by Greenfield for a two-state transition of a monomer from a

folded to an unfolded state [44]. This analysis is based on the Gibbs-Helmholtz equation

where ΔG represents Gibbs free energy, ΔH represents enthalpy, ΔC_P represents the

change in heat capacity, K represents the folding constant, and y represents the fraction

of folded DNA at any given temperature, T. θ_t is the observed ellipticity at any

temperature, θ_F is the observed ellipticity of the folded DNA, and θ_U is the observed

ellipticity of the unfolded DNA. The melting temperature, T_m, of the DNA can then be

calculated as the temperature where y = 0.5.

352
$$\Delta G = \Delta H \left(1 - \frac{T}{T_m}\right) - \Delta C_p \left((T_m - T) + T \left(\ln\left(\frac{T}{T_m}\right)\right)\right)$$

353
$$K = e^{\left(-\frac{\Delta G}{RT}\right)}$$

354
$$y = \frac{K}{1 + K}$$

355
$$[\theta]_t = \left(([\theta]_F - [\theta]_U) * y\right) + [\theta]_U$$

Fluorescence spectroscopy

Fluorescence spectroscopy measurements were performed on a Cary Eclipse

Fluorescence Spectrophotometer (Agilent Technologies, Santa Clara, CA) equipped with

a PCB 1500 water peltier system. Spectra were taken from 300 nm to 450 nm with the

excitation at 280 nm and temperature at 25 °C. A scan of buffer alone, 20 mM HEPES pH

7.5, 150 mM KCl or NaCl, 5 mM TCEP, and 5% (v/v) glycerol, was used as a baseline.

FANCJ AKKQ peptide (5 μM) was titrated with additions of a DNA substrate. The

mixture was allowed to equilibrate for 3 minutes before taking each spectra scan.

Observed fluorescence quenching (ΔF) was calculated using the following

equation, with F_0 representing total fluorescence from the FANCJ peptide alone and F_i

representing the total fluorescence after the *i*th addition of DNA:

367
$$\Delta F = \frac{F_0 - F_i}{F_0}$$

Binding isotherms were calculated by plotting ΔF versus total DNA concentration.

The resulting curve was fit to a 1:1 interaction model using Scientist 2.0 software

(Micromath, St. Louis, MO) to calculate an equilibrium K. The equilibrium K was

calculated using the following formulas, where A was the amplitude of fluorescence

quenching and K was the equilibrium association constant. D_f and D_t were the free and

total concentration of DNA, while P_f and P_t described the free and total peptide

concentration. Fluorescence titrations were performed in triplicate. The reported K values

were determined from an average of three independent data-sets.

$$\Delta F_{obs} = A\left(\frac{KD_f}{1+KD_f}\right)$$
$$D_t = D_f(1+KP_f)$$
$$P_t = P_f(1+KD_f)$$

Acknowledgements

378   Figs 1, 3-5, 8A, and 11 were created with BioRender.com.

Author contributions

380   CGW designed experiments. LC, KL, and CGW acquired data and interpreted results.

381   LC, KL, EC, and CGW prepared the manuscript draft. All authors have read and agreed

382   to the published version of the manuscript.

Financial disclosure statement

384   This work was supported by NSF grant 2142839 and AHA grant 19AIREA34460026 to

385   CGW.

Competing interests

387   Authors have no conflict of interest to disclose.

References

1. Bochman ML, Paeschke K, Zakian VA. DNA secondary structures: stability and function of G-quadruplex structures. Nature Reviews Genetics. 2012;13(11):770-80.

2. Nový J, Böhm S, Králová J, Král V, Urbanová M. Formation and temperature stability of G-quadruplex structures studied by electronic and vibrational circular dichroism spectroscopy combined with ab initio calculations. Biopolymers. 2008;89(2):144-52.

3. Chaires JB. Human telomeric G-quadruplex: thermodynamic and kinetic studies of telomeric quadruplex stability. The FEBS Journal. 2010;277(5):1098-106.

4. Guédin A, Gros J, Alberti P, Mergny J-L. How long is too long? Effects of loop size on G-quadruplex stability Nucleic Acids Research. 2010;38(21):7858-68.

5. Huppert JL, Balasubramanian S. Prevalence of quadruplexes in the human genome. Nucleic Acids Research. 2005;33(9):2908-16.

6. Lombardi EP, Holmes A, Verga D, Teulade-Fichou M-P, Nicolas A, Londoño-Vallejo A. Thermodynamically stable and genetically unstable G-quadruplexes are depleted in genomes across species Nucleic Acids Research. 2019;47(12):6098-113.

7.   Hazel P, Huppert J, Balasubramanian S, Neidle S. Loop-length-dependent folding

of G-quadruplexes. Journal of the American Chemical Society. 2004;126(50):16405-

15.

8.   Hao F, Ma Y, Guan Y. Effects of Central Loop Length and Metal Ions on the

Thermal Stability of G-Quadruplexes. Molecules. 2019;24(10).

9.   Burge S, Parkinson GN, Hazel P, Todd AK, Neidle S. Quadruplex DNA:

sequence, topology and structure. Nucleic Acids Research. 2006;34(19):5402-15.

10. Villar-Guerra RD, Trent JO, Chaires JB. G-Quadruplex Secondary Structure

Obtained from Circular Dichroism Spectroscopy. Angewandte Chemie.

2018;57(24):7171-5.

11. Heddi B, Cheong VV, Martadinata H, Phan AT. Insights into G-quadruplex

specific recognition by the DEAH-box helicase RHAU: Solution structure of a

peptide–quadruplex complex. Proceedings of the National Academy of Sciences

of the United States of America. 2015;112(31):9608-13.

12. Budhathoki JB, Maleki P, Roy WA, Janscak P, Yodh JG, Balci H. A Comparative

Study of G-Quadruplex Unfolding and DNA Reeling Activities of Human RECQ5

Helicase. Biophysical Journal. 2016;110(12):2585-96.

13. Tippana R, Hwang H, Opresko PL, Bohr VA, Myong S. Single-molecule imaging

reveals a common mechanism shared by G-quadruplex–resolving helicases.

Proceedings of the National Academy of Sciences of the United States of America. 2016;113(30):8448-53.

14. Zheng K-W, Zhang J-Y, He Y-d, Gong J-Y, Wen C-J, Chen J-N, et al. Detection of genomic G-quadruplexes in living cells using a small artificial protein. Nucleic Acids Research. 2020;48(20):11706-20.

15. Wang L, Wang Q-M, Wang Y-R, Xi X-G, Hou X-M. DNA-unwinding activity of Saccharomyces cerevisiae Pif1 is modulated by thermal stability, folding conformation, and loop lengths of G-quadruplex DNA. Journal of Biological Chemistry. 2018;293(48):18504-13.

16. Rhodes D, Lipps HJ. G-quadruplexes and their regulatory roles in biology Nucleic Acids Research. 2015;43(18):8627-37.

17. Huppert JL, Balasubramanian S. G-quadruplexes in promoters throughout the human genome. Nucleic Acids Research. 2006;35(2):406-13.

18. Beaudoin J-D, Perreault J-P. 5'-UTR G-quadruplex structures acting as translational repressors. Nucleic Acids Research. 2010;38(20):7022-36.

19. Smith JS, Chen Q, Yatsunyk LA, Nicoludis JM, Garcia MS, Kranaster R, et al. Rudimentary G-quadruplex-based telomere capping in Saccharomyces cerevisiae. Nature Structural & Molecular Biology. 2011;18(4):478-85.

20. Georgakopoulos-Soares I, Parada GE, Wong HY, Medhi R, Furlan G, Munita R, et

al. Alternative splicing modulation by G-quadruplexes. Nature Communications.

2022;13(1).

21. Xu H, Di Antonio M, McKinney S, Mathew V, Ho B, O'Neil NJ, et al. CX-5461 is a

DNA G-quadruplex stabilizer with selective lethality in BRCA1/2 deficient

tumours. Nature Communications. 2017;8.

22. Steenken S, Jovanovic S. How Easily Oxidizable Is DNA? One-Electron Reduction

Potentials of Adenosine and Guanosine Radicals in Aqueous Solution. Journal of

the American Chemical Society. 1997;119(3):617-8.

23. Oikawa S. Sequence-specific DNA damage by reactive oxygen species:

Implications for carcinogenesis and aging. Environmental Health and Preventive

Medicine. 2005;10(2):65-71.

24. Neeley WL, Essigmann JM. Mechanisms of Formation, Genotoxicity, and

Mutation of Guanine Oxidation Products. Chemical Research in Toxicology.

2006;19(4):491-505.

25. Ohno M, Sakumi K, Fukumura R, Furuichi M, Iwasaki Y, Hokama M, et al. 8-

oxoguanine causes spontaneous de novo germline mutations in mice. Scientific

Reports. 2014;4.

26. Bielskutė S, Plavec J, Podbevšek P. Impact of Oxidative Lesions on the Human Telomeric G-Quadruplex. Journal of the American Chemical Society. 2019;141(6):2594-603.

27. Vorlícková M, Tomasko M, Sagi AJ, Bednarova K, Sagi J. 8-oxoguanine in a quadruplex of the human telomere DNA sequence. The FEBS Journal. 2012;279(1):29-39.

28. Jr RMB, Cantor SB. Molecular and cellular functions of the FANCJ DNA helicase defective in cancer and in Fanconi anemia. Frontiers in Genetics. 2014;5.

29. Palovcak A, Liu W, Yuan F, Zhang Y. Maintenance of genome stability by Fanconi anemia proteins. Cell & Bioscience. 2017;7.

30. Datta A, Robert M. Brosh J. Holding All the Cards—How Fanconi Anemia Proteins Deal with Replication Stress and Preserve Genomic Stability. Genes. 2019;10(2).

31. Wu Y, Shin-ya K, Jr RMB. FANCJ helicase defective in Fanconia anemia and breast cancer unwinds G-quadruplex DNA to defend genomic stability. Molecular and Cellular Biology. 2008;28(12):4116-28.

32. Cantor SB, Xie J. Assessing the link between BACH1/FANCJ and MLH1 in DNA crosslink repair. Environmental and Molecular Mutagenesis. 2010;51(6):500-7.

33. Sarkies P, Murat P, Phillips LG, Patel KJ, Balasubramanian S, Sale JE. FANCJ coordinates two pathways that maintain epigenetic stability at G-quadruplex DNA. Nucleic Acids Research. 2012;40(4):1485-98.

34. Bosch PC, Segura-Bayona S, Koole W, Heteren JTv, Dewar JM, Tijsterman M, et al.

FANCJ promotes DNA synthesis through G-quadruplex structures. The EMBO

Journal. 2014;33(21):2521-33.

35. Wu CG, Spies M. G-quadruplex recognition and remodeling by the FANCJ

helicase. Nucleic Acids Research. 2016;44(18):8742-53.

36. Lowran K, Campbell L, Popp P, Wu CG. Assembly of a G-Quadruplex Repair

Complex by the FANCJ DNA Helicase and the REV1 Polymerase. Genes.

2019;11(1).

37. Lattmann S, Giri B, Vaughn JP, Akman SA, Nagamine Y. Role of the amino

terminal RHAU-specific motif in the recognition and resolution of guanine

quadruplex-RNA by the DEAH-box RNA helicase RHAU. Nucleic Acids

Research. 2010;38(18):6219-33.

38. Rachwal PA, Fox KR. Quadruplex melting. Methods. 2007;43(4):291-301.

39. Do NQ, Lim KW, Teo MH, Heddi B, Phan AT. Stacking of G-quadruplexes: NMR

structure of a G-rich oligonucleotide with potential anti-HIV and anticancer

activity. Nucleic Acids Research. 2011;39(21):9448-57.

40. Guédin A, Cian AD, Gros J, Lacroix L, Mergny J-L. Sequence effects in single-base

loops for quadruplexes. Biochimie. 2008;90(5):686-96.

41. Lin C, Yang D. Human Telomeric G-Quadruplex Structures and G-Quadruplex-

Interactive Compounds. Methods in Molecular Biology. 2018:171-96.

42. Hardin CC, Watson T, Corregan M, Bailey C. Cation-dependent transition

between the quadruplex and Watson-Crick hairpin forms of d(CGCG3GCG).

Biochemistry. 1992;31(3):833–41.

43. Eddy S, Ketkar A, Zafar MK, Maddukuri L, Choi J-Y, Eoff RL. Human Rev1

polymerase   disrupts   G-quadruplex   DNA.   Nucleic   Acids   Research.

2014;42(5):3272-85.

44. Greenfield NJ. Using circular dichroism collected as a function of temperature to

determine the thermodynamics of protein unfolding and binding interactions.

Nature Protocols. 2006;1(6):2527-35.

Fig 1. G-quadruplex formation and conformations.

(A) Formation of a G4 structure. (B) Graphic representation of three different intra-strand G4 conformations.

Fig 2. Molecular models of FANCJ.

(A) FANCJ possesses a G4 recognition site capable of binding G4s, and an AKKQ motif similar in structure and sequence to the AKKQ binding motif found in G4-binding protein RHAU [11, 37]. (B) The current model for FANCJ-supported DNA replication in G4-rich sequences. Upon reaching a G4 structure, the DNA polymerase stalls. FANCJ binds to the G4 structure to unwind it and another polymerase is recruited to continue DNA replication.

523   **Fig 3. Map of FANCJ**.

524   The AKKQ sequence (amino acids 129-147) has been highlighted.

Fig 4. CD spectroscopy cartoons.

(A) Graphic representation of CD spectra for parallel, antiparallel, and hybrid G4 conformations. Parallel conformations have a maximum at 262 nm and a minimum at 240 nm. Antiparallel conformations give two peak signals near 295 nm and 240 nm, and a minimum at 260 nm. Hybrid conformations show a peak at 295 nm, a shoulder around 260 nm, and a minimum at 240 nm. (B) Graphic representation of the CD spectra of a parallel G4 conformation as a function of temperature. (C) Graphic representation of a thermal melt graph of two G4s.

Fig 5. Graphic representation of a parallel G4 conformation.

An 8oxoG-modified base has been substituted at the (A) first guanine

(8oxo1), (B) third guanine (8oxo3), and (C) fifth guanine (8oxo5).

Fig 6. Experiments performed in the presence of 150 mM KCl.

(A) CD spectra of $(GGGT)_4$, $(GGGT)_4$ 8oxo1, $(GGGT)_4$ 8oxo3, and $(GGGT)_4$

8oxo5 taken from 320 to 220 nm. (B) Temperature melt of same sequences

seen in 6A. Melt taken at 262 nm from 15 to 95 °C. (C) CD spectra of

$(GGGTT)_4$, $(GGGTT)_4$ 8oxo1, $(GGGTT)_4$ 8oxo3, and $(GGGTT)_4$ 8oxo5 taken

545   from 320 to 220 nm. (D) Temperature melt of same sequences seen in 6C.

546   Melt taken at 262 nm for all sequences except (GGGTT)$_4$ 8oxo3, which was

547   taken at 295 nm. (E) CD spectra of (TTAGGG)$_4$, (TTAGGG)$_4$ 8oxo1,

548   (TTAGGG)$_4$ 8oxo3, and (TTAGGG)$_4$ 8oxo5 taken from 320 to 220 nm. (F)

549   Temperature melt of the same sequences seen in 6E. Melt taken at 295 nm.

Fig 7. Experiments performed in the presence of 150 mM NaCl.

(A) CD spectra of (GGGT)$_4$, (GGGT)$_4$ 8oxo1, (GGGT)$_4$ 8oxo3, and (GGGT)$_4$

8oxo5 taken from 320 to 220 nm. (B) Temperature melt of same sequences

as seen in 7A. Melt taken at 262 nm from 15 to 95 °C. (C) CD spectra of

(GGGTT)$_4$, (GGGTT)$_4$ 8oxo1, (GGGTT)$_4$ 8oxo3, and (GGGTT)$_4$ 8oxo5 taken

from 320 to 220 nm. (D) Temperature melt of same sequences as seen in 7C.

Melt taken at 262 nm for all sequences except (GGGTT)$_4$ 8oxo3, which was

taken at 295 nm. (E) CD spectra of (TTAGGG)$_4$, (TTAGGG)$_4$ 8oxo1,

(TTAGGG)$_4$ 8oxo3, and (TTAGGG)$_4$ 8oxo5 taken from 320 to 220 nm. (F)

Temperature melt of the same sequences as seen in 7E. Melt taken at 295

nm.

Fig 8. Overview of fluorescence spectroscopy analyses.

(A) Molecular model of fluorescence spectroscopy titrations. AKKQ is

equipped with a tryptophan residue that fluoresces. When bound to DNA,

the residue is quenched and fluorescence intensity decreases. (B) Titration

of AKKQ with $(GGGT)_4$. As DNA concentration increases, the peak

intensity decreases. (C) Analysis of fluorescence titration. The decrease in

integrated peak is plotted as a function of DNA concentration. The resulting

isotherm is fit to a 1:1 binding interaction model to generate an equilibrium

K value.

Fig 9. Summary of FANCJ AKKQ-G4 binding in KCl by fluorescence

spectroscopy.

Trials performed in 20 mM HEPES pH 7.5, 150 mM KCl, 5 mM TCEP, and

5% (v/v) glycerol. (A) FANCJ AKKQ equilibrium binding to (GGGT)$_4$ and

its respective 8oxoG4 sequences. (B) FANCJ AKKQ binding to (GGGTT)$_4$

and its respective 8oxoG4 sequences. (C) FANCJ AKKQ binding to

(TTAGGG)$_4$ and its respective 8oxoG4 sequences.

Fig 10. Summary of FANCJ AKKQ-G4 binding in NaCl by fluorescence

spectroscopy.

Trials performed in 20 mM HEPES pH 7.5, 150 mM NaCl, 5 mM TCEP, and

5% (v/v) glycerol. (A) FANCJ AKKQ binding to (GGGT)$_4$ and respective

8oxoG4 sequences. (B) FANCJ AKKQ binding to (GGGTT)$_4$ and respective

8oxoG4 sequences. (C) FANCJ AKKQ binding to (TTAGGG)$_4$ and

respective 8oxoG4 sequences.

Fig 11. Proposed binding interaction model between FANCJ AKKQ and

8oxoG4 sequences.

(A) For G4 sequences that form compact structures with short intervening

sequences, such as (GGGT)$_4$, AKKQ preferentially targets the 8oxoG site

and promotes repair. (B) For G4 sequences that expose the intervening

sequence, as with antiparallel or hybrid conformations, AKKQ

preferentially targets the intervening sequence and ignores the 8oxoG.