The Septin Cytoskeleton excludes the Cdc42 GAP Rga4 from growing sites to

promote bipolarity in fission yeast

Justin McDuffie, Goeun Chang, Maitreyi Das*

Biology Department, Boston College, Chestnut Hill, MA, 02467

*Corresponding author: Maitreyi.Das@bc.edu

Running title: Septins restrict Rga4 localization

Keywords: Septin, Cell cycle, Cdc42, GAP, Rga4, Cell polarity

Summary: Cell-cycle-dependent Rga4 localization pattern is controlled by the septin

cytoskeleton to promote bipolar growth. In interphase, the septin cytoskeleton at the cell

sides restricts Rga4 mobility and localization away from the cell ends to promote proper

Cdc42 activation dynamics.

ABSTRACT

The conserved GTPase Cdc42 regulates polarity in most eukaryotes and is

spatiotemporally regulated to confine growth to discreet areas throughout the cell cycle.

Cdc42 activity stops at the growing ends of fission yeast during mitosis and resumes

after division when the ends remove the inactivator/GAP Rga4. In interphase, Rga4

appear punctate at the cell sides restricted from the growing ends. During mitosis, Rga4

appears homogeneous and localizes to the ends. We show that this cell-cycle-

dependent Rga4 localization is septin cytoskeleton-dependent. Septins forms linear

filaments along the sides during interphase. In mitosis, the septins recede from the

sides and form a ring at the division site. In septin mutants, Rga4 is mobile,

homogeneously distributed and localizes to the ends, regardless of cell-cycle stage.

Accordingly, septin mutants show delayed Cdc42 reactivation after division and are

monopolar with disrupted Cdc42 activation dynamics. Our data describe an unusual

form of polarity regulation where a septin-dependent physical barrier restricts GAP

localization for proper Cdc42 dynamics.

INTRODUCTION

Establishment and maintenance of cell shape is fundamental to its function. Most cells

exhibit consistent cell shape after every round of cell division and this ensures their

ability to maintain function. Consistent cell shape is not just observed in multicellular

organisms that form highly organized tissues and organelles, but even in unicellular

organisms such a yeast (Das and Verde, 2013; Galliot and Ghila, 2010; Guillot and

Lecuit, 2013; Mitchison and Nurse, 1985). In most eukaryotes, polarized cell growth

stops during division and resumes immediately after to establish consistent cell shape

(Lloyd, 2013; McCusker and Kellogg, 2012). It is not well understood how polarized

growth resumes promptly after each round of cell division. To understand this process,

we use the fission yeast model system *Schizosaccharomyces pombe*. These cells stop

polarized growth from the cell ends during division and resume growth once the cells

complete cytokinesis (Mitchison and Nurse, 1985). However, we have previously shown

that resumption of polarized growth from the cell ends is independent of cytokinesis and

is instead cell-cycle-dependent and occurs at a fixed time after mitosis (Rich-Robinson

et al., 2021).

Like in many eukaryotic cells, the major regulator of polarized growth in fission yeast is

the small GTPase Cdc42 (Das and Verde, 2013; Estravis et al., 2012; Etienne-

Manneville, 2004; Johnson, 1999). Cdc42 is activated when Guanine nucleotide

exchange factors (GEFs) help it bind GTP (Bos et al., 2007; Sprang, 2001). GTP-bound

Cdc42 binds and activates downstream targets to promote polarized growth (Bos et al.,

2007). Cdc42 is inactivated when it hydrolyses the GTP to GDP with the help of the

GTPase activating proteins (GAPs) (Bos et al., 2007). Cdc42 in fission yeast has two

GEFs, Scd1 and Gef1, and three GAPs, Rga4, Rga6 and Rga3 (Chang et al., 1994;

Coll et al., 2003; Das et al., 2007; Gallo Castro and Martin, 2018; Murray and Johnson,

2001; Revilla-Guarinos et al., 2016; Tatebe et al., 2008). Our lab previously showed that

during cell division, the GAP Rga4 undergoes a change in its localization pattern (Rich-

Robinson et al., 2021). Rga4 localizes to the cell sides in a punctate manner during

interphase. It becomes diffuse along the cell cortex and localizes to the cell tips in

dividing cells, where it prevents Cdc42 activation. Once cells complete division, Rga4

resumes its punctate localization away from the cell ends and growth at the ends

resume. It is not known how Rga4 localization is regulated in a cell-cycle-dependent

manner.

Rga4 has a coiled-coil region that is required for its localization to the cell cortex (Das et

al., 2007). Cells lacking *rga4* show increased width, since in these mutants, the zone of

Cdc42 activation and consequent growth spreads over a larger area at the cell ends

(Das et al., 2007; Tatebe et al., 2008). It is not known how Rga4 localization is limited to

the cell sides and away from the ends. A recent report shows that Rga4 is removed from

the cell ends as a result of membrane flow that occurs due to extensile growth at these

ends (Gerganova et al., 2021; Rutkowski et al., 2024). During mitosis, Rga4 appears

diffuse along the membrane and appears at the cell ends (Rich-Robinson et al., 2021).

It is not known what prevents the diffuse Rga4 localization at the cell surface restricting

it to the sides.

In several eukaryotic cells the septin cytoskeleton acts as a diffusion barrier along the

membrane (Bridges and Gladfelter, 2015; Caudron and Barral, 2009; Kinoshita, 2003).

S. pombe expresses seven total septin proteins (Spn1-7p), however, only four are

involved in the regular cell cycle (Spn1-4p) (An et al., 2004; Berlin et al., 2003; Onishi et

al., 2010). The septin cytoskeleton is a heterotetramer consisting of 4 peptides Spn1-4

(An et al., 2004; Berlin et al., 2003). These peptides form heterotetramer that organize

in a palindromic manner to form larger filaments (Kinoshita, 2003; Spiliotis and

McMurray, 2020; Woods and Gladfelter, 2021). The septin cytoskeleton plays a role in

cytokinesis where it forms a ring at the division site during cytokinesis. The septin ring

acts as the landmark where proteins required for cell separation during cytokinesis are

delivered (Berlin et al., 2003; Maddox et al., 2007; Martin-Cuadrado et al., 2005; Singh

et al., 2024; Spiliotis and McMurray, 2020). It is not clear if the septin cytoskeleton plays

a role during interphase. Cells lacking the septin cytoskeleton show cell separation

defects but remain polarized (An et al., 2004; Berlin et al., 2003; Martin-Cuadrado et al.,

2005).

Here we show that the septin cytoskeleton acts as a barrier to prevent Rga4 diffusion

along the cell cortex and restrict it to the cell sides. In cells lacking septin filaments,

Rga4 localization is diffuse along the cortex and also appears at the cell ends, similar to

that observed in mitotic cells. In the presence of septin filaments, the Rga4 puncta at

the cell sides are mostly immobile. During interphase, the septin cytoskeleton forms

small filaments at the cortex that align along the long axis of the cell. The septin levels

along the cortex decrease as the cells enter mitosis, and eventually, a septin ring forms

at the end of anaphase B. Rga4 localization appears diffuse as the cells enter mitosis,

and throughout cytokinesis. Correspondingly, active Cdc42 is lost from the cell ends

during mitosis. Using the GFP-GBP trap method, we show that the septin cytoskeleton

does not interact with Rga4 and likely restricts its diffusion by acting as a barrier. The

septin mutants show a delay in reactivation of Cdc42 at the cell ends as the cells

complete division. Moreover, these mutants are monopolar compared to controls,

indicating that the presence of Rga4 at the cells ends disrupts bipolarity. Together, these

findings show that the septin cytoskeleton forms a diffusion barrier along the cortex to

restrict Rga4 localization along the cell sides. During mitosis, septin localization along

the cortex is lost, thus allowing diffusion of Rga4 along the membrane and its

localization to the cell ends. Rga4 at the ends in the absence of septin disrupt Cdc42

activation dynamics resulting in monopolar cells. These findings show that the septin

cytoskeleton maintains cell polarity in a cell-cycle-dependent manner.

RESULTS

The septin cytoskeleton restricts Rga4 to the cell sides during interphase

To investigate if the septin cytoskeleton acts as a diffusion barrier for Rga4, we analyzed Rga4-GFP localization in *spn1*$^+$ and *spn1Δ* cells. In the *spn1*$^+$ cells, Rga4-GFP along the cell sides appears punctate in interphase cells and diffuse in dividing cells. In contrast, *spn1Δ* cells Rga4-GFP show the diffuse pattern, regardless of the cell cycle stage (Fig.1A). We quantified the level of Rga4 heterogeneity along the cell sides by measuring the coefficient of variance of Rga4-GFP intensity. As reported earlier, we find that the coefficient of variance of Rga4-GFP is highest in the cells in G2 phase of the cell cycle and the lowest in the mitotic phase in *spn1*$^+$ cells (Fig 1, B) (Rich-Robinson et al., 2021). This indicates that Rga4-GFP is more homogeneous along the cell sides in mitotic cells and form puncta during G2 or interphase. In *spn1Δ* cells, the coefficient of variance in G2 cells is low and is similar to that in mitotic cells (Fig 1, C).

In mitotic, cells Rga4 appears more cytoplasmic and is also increasingly observed at the cell ends. We asked if septins restrict cytoplasmic Rga4 and prevents its localization to the cell ends. Rga4-GFP along the cells ends appear faint even in z-projected images due to the curvature of the cell and weak expression/signal. To overcome this, we took rapid timelapse movies with 5 second intervals over 5 mins of Rga4-GFP expressing *spn1+* and *spn1Δ* cells. We measured Rga4-GFP intensity in the cytoplasm and the cell ends of time-projected images of these movies (Fig1, D). We find that compared to *spn1*$^+$ cells cytoplasmic Rga4-GFP levels increase in *spn1Δ* cells (Fig1, E). Similarly, *spn1Δ* cells show enhanced Rga4-GFP levels at the cell ends compared to *spn1*$^+$ cells (Fig1, F). In control cells, Rga4-GFP shows elevated levels at the non-growing cell ends

138   compared to growing ends. We also see higher levels of Rga4-GFP at the non-growing

139   cell ends of *spn1Δ* cells but these levels are much higher than that in *spn1+* cells.

140   Moreover, the growing ends in *spn1Δ* cells show higher Rga4-GFP levels compared to

141   growing ends of *spn1+* cells. This indicates that in the absence of septins, Rga4-GFP

142   levels at the cell ends are elevated.

Septins inhibit Rga4-GFP mobility during Interphase

Next, we asked whether the increased homogeneity of Rga4-GFP along the plasma membrane is the result of increased Rga4-GFP mobility or diffusion along the membrane. The above mentioned 5-minute timelapse movies of Rga4-GFP localization were analyzed and we found that, in *spn1*[+] cells in interphase, Rga4-GFP formed a number of bright, stationary puncta along the plasma membrane that remained immobile throughout the course of timelapse imaging (Fig. 2A). In contrast, *spn1*[+] cells in mitosis, and *spn1Δ* cells at all stages of the cell-cycle showed limited Rga4-GFP puncta formation, and lacked immobile Rga4-GFP puncta. In these cells Rga4-GFP appeared to diffuse more rapidly along the plasma membrane. To quantify this observed change in Rga4 mobility, fluorescence recovery after photobleaching (FRAP) microscopy was performed on 10x10-pixel ROIs of Rga4-GFP along the cell sides of *spn1*[+] and *spn1Δ* cells in interphase and mitosis (Fig. 2B). Cell-cycle-progression was marked by the fluorescently tagged spindle pole body marker Sad1-mCherry. Fluorescence recovery curves and calculated recovery half-lives of Rga4-GFP in the photobleached ROIs showed that Rga4-GFP was much less mobile along the plasma membrane in interphase cells (Fig. 2C, D). In contrast, Rga4-GFP is much more mobile along the plasma membrane in *spn1*[+] cells during mitosis. Similarly, to mitotic *spn1*[+] cells, Rga4-GFP is much more mobile along the plasma membrane in *spn1Δ* cells, regardless of cell-cycle-stage. This indicates that in the absence of septin filaments, the diffusion of Rga4-GFP is significantly less restricted than when they are present, leading to increased protein mobility. This increased mobility along the plasma membrane may

also explain the increased localization of Rga4-GFP to the cytoplasm and the cell ends

in the septin mutants.

A previous report has indicated that the GAP Rga6 interacts with the septin filaments

and is required for proper septin localization at the cortex in interphase cells (Zheng et

al., 2022). Thus, we asked if the septin filaments also regulate Rga6 distribution along

the cortex. Similar to Rga4-GFP, Rga6-GFP also appears punctate along the cell sides

(Supplementary Fig. S1A). However, in *spn1Δ* mutants, Rga6-GFP localization

remained the same and the coefficient of variance along the cortex was not significantly

distinct between *spn1⁺* and *spn1Δ* cells, and throughout the cell cycle stages

(Supplementary Fig. S1B). This indicates that the septin cytoskeleton specifically

disrupts Rga4 distribution along the cortex.

Septins form linear filaments in interphase and a ring structure as Anaphase B ends

Given that Rga4-GFP localization in *spn1Δ* cells regardless of the cell cycle stage is similar to that of mitotic *spn1+* cells, we asked if this is due to changes in septin localization throughout the cell cycle. While extensive work has revealed septin localization at the division site during cytokinesis, it is unclear how septin proteins localize during interphase. We find that in interphase cells, the septin proteins appear as short filaments at the plasma membrane along the long axis of the cell. Spn1-tdTomato and Spn3-GFP co-localize as short linear filaments along the cell cortex (Fig 3A). The other septin protein Spn2 also appears as short filaments along the long axis of the cell as determined by Spn2-GFP (Fig. 3B). Spn4 on the other hand appears more punctate but along the cell sides as determined by Spn4-tdTomato (Fig.3C).

The septin filament distribution appears biased towards the growing cell ends. To confirm this, we investigated Spn3-GFP localization in *tea1+* and *tea1Δ* mutants. The *tea1Δ* mutants are monopolar and only grow from one end (Mata and Nurse, 1997; Verde et al., 1995). We find that in *tea1Δ* cells Spn3-GFP filaments appear closer to the growing cell ends (Supplementary Fig. S2A, red arrowheads). In *tea1Δ* cells Spn3-GFP levels are higher at the growing ends compared to the *tea1+* cells (Supplementary Fig. S2B). This is likely due to the fact that in *tea1+* cells Spn3-GFP is distributed between the two growing ends of bipolar cells while in *tea1Δ* cells all of the Spn3-GFP accumulates to the single growing end.

Since, the septin cytoskeleton appears to restrict Rga4 mobility, we asked of the septin

proteins themselves were mobile along the cell cortex. To test for mobility, we performed

FRAP analysis of Spn1-tdTomato along the cell sides in both interphase and dividing

cells. Regardless of the cell cycle stage, we do not see any recovery of the fluorescent

signal after photobleaching of Spn1-tdTomato (Supplementary Fig. S2C,D). This

indicates that the septin filaments observed along the cell sides are stable and

immobile.

Next, we investigated how septin localization changes throughout the cell cycle. We

imaged cells expressing Spn4-tdTomato along with Myo2-GFP and Sid4-GFP over

timelapse (Fig. 3D). While the Sid4-GFP labels the spindle pole body and determines

the onset of mitosis when the spindle pole body splits into two, Myo2-GFP labels the

cytokinetic actomyosin ring (Sparks et al., 1999; Wei et al., 2017). We find that Spn4-

tdTomato intensity along the cell cortex starts to drop as the cell enters mitosis, until it

plateaus (Fig.3E). The Spn4-tdTomato eventually forms a septin ring just prior to mitotic

exit. We quantified Spn4-tdTomato levels along the sides of cells in different mitotic

stages, as determined by the distance between spindle pole bodies. We find that Spn4-

tdTomato levels along the cell sides is highest in interphase cells with a single spindle

pole body (distance, 0µm, Fig. 3F). As the spindle pole body distance increases, Spn4-

tdTomato levels along the cell sides drop. Together these observations indicate that, in

interphase cells, septins form short linear filaments at the cortex along the long axis of

the cell. These filaments begin to drop when the cell enters mitosis, however the septin

ring forms after a delay close to the end of Anaphase B.

The septin cytoskeleton and Rga4 show distinct structures at the cortex

Since Rga4 distribution pattern along the cortex is septin-dependent, we investigated

whether the septin cytoskeleton colocalizes with Rga4. We observed Rga4-GFP puncta

along the cell sides in cells expressing Spn1-tdTomato. While both these proteins

localize to the cell sides, we did not see any significant colocalization pattern between

them (Fig.4A, Top plane, inset). In most cells, Rga4-GFP organizes like a corset at the

cell middle as has been reported before, while Spn1-tdTomato appears on either side of

Rga4-GFP closer to the growing ends (Fig.4A, red arrowhead). In some cells, Rga4-

GFP puncta appeared on either side of Spn1-tdTomato filaments (Fig.4A, middle plane,

inset). We asked if the dynamic Rga4-GFP is trapped by the septin via physical

interaction, thereby preventing its mobility along the cortex.

If the septin cytoskeleton physically interacts with Rga4 and entraps it, mislocalizing the

septin should lead to Rga4-GFP mislocalization. To test this, we used the GFP trap

method with a GFP-binding protein (GBP) (Wang et al., 2016). We used the

mitochondrial protein Tom20 tagged with GBP and RFP to entrap Spn3-GFP (Fig.4B).

We find that in cells expressing Tom20-GBP-RFP, Spn3-GFP is no longer observed at

the cortex and rather appears along the mitochondria colocalizing with Tom20. We

asked if mislocalizing one of the components of the septin cytoskeleton can mislocalize

the other components. We expressed Spn1-TdTomato and Spn3-GFP in cells

expressing Tom20-GBP. We find that Spn3-GFP with Tom20-GBP at the mitochondria

also entraps Spn1-tdTomato at that site (Fig.4C). This indicates that the septin

cytoskeleton is mislocalized to the mitochondria with Tom20-GBP.

Next, we asked if Rga4-GFP can be mislocalized to the mitochondria with Tom20-GBP.

We find that in cells expressing Tom20-GBP, Rga4-GFP appears away from the cortex

and along the mitochondria (Fig.4D). However, in these cells Spn1-tdTomato remained

at the cortex and did not colocalize with the mitochondria and Rga4-GFP. Moreover, in

Tom20-GBP expressing cells, Spn3-GFP at the mitochondria did not entrap Rga4-

mCherry (Fig.4D). In these cells, Rga4-mCherry remained at the cortex along the cell

sides. These observations suggest that the septin cytoskeleton does not regulate Rga4

distribution via physical interaction.

It is noteworthy that in the cells where the septin cytoskeleton localized to the

mitochondria and away from the cortex, Rga4 at the cell sides appeared less punctate

and more diffuse, similar to that observed in $spn1\Delta$ cells. To confirm this, we quantified

the coefficient of variance of Rga4-mCherry signal along the cell sides in control cells

with that of cells co-expressing Tom20-GBP and Spn3-GFP. We find that the coefficient

of variance of Rga4-mCherry is significantly reduced in cells where Spn3-GFP is

mislocalized to the Tom20-GBP containing mitochondria (Fig.4F). This indicates that the

absence of the septin cytoskeleton from the cell cortex is sufficient to alter Rga4

distribution pattern even in cells where the septin proteins are expressed. These

observations further support the hypothesis that the septin cytoskeleton acts as a

physical barrier that prevents Rga4 mobility along the cell sides.

267   **As septins recede from the cell sides, Rga4 spreads out along the membrane and**

268   **Cdc42 activity diminishes at the cell ends**

Our data show that Rga4 localization at the cortex is mostly punctate and immobile in

interphase and diffused and mobile in dividing cells. In addition, we find that the septins

show short filaments along the cell cortex in interphase and this localization is lost once

the cell enters mitosis. Based on this, we propose that Rga4 diffuses along the cortex

when the septins are lost from this region. To verify this, we observed Rga4-GFP and

Spn1-tdTomato localization along the cortex in cells at different cell cycle stages

(Supplementary Fig.S2E). As expected, we find that in early and late G2 cells Rga4-

GFP appears punctate along the cell sides. Spn1-tdTomato also localizes to this region.

In early and late mitotic cells, Rga4-GFP appears more diffuse and Spn1-tdTomato

levels along the sides is lower. Given the low levels of Rga4, it was not possible to

image Rga4-GFP through timelapse of different cell cycle stages along with Spn1-

tdTomato. The Spn1-tdTomato ring appears in post-mitotic cells while Rga4-GFP shows

diffuse localization in mitotic cells as the septin localization along the cortex decreases

(Supplementary Fig.S2E, early and late mitosis). This suggests that the removal of

septins from the cell cortex, and not the formation of the septin ring, is sufficient to

enable diffuse Rga4 localization.

Next, we asked if Rga4 mobility and its ability to localize to the cell ends leads to Cdc42

inactivation at the cell ends in dividing cells. We find that active Cdc42 at the cell ends is

not immediately lost upon the onset of mitosis, as observed in CRIB-mCherry

expressing cells with Sad1-mCherry as spindle pole body markers (Supplementary

Fig.S2F). As the Rga4-GFP appears diffuse in these cells during mitosis, we observe

loss of CRIB-mCherry from the ends. This suggests that as the septins gradually

decrease from the cell sides, Rga4 mobility along the cortex is enhanced, resulting in

eventual loss of active Cdc42 from the cell ends.

Cdc42 is active at the cell ends in interphase and at the division site in diving cells

(Merla and Johnson, 2000; Wei et al., 2016). Our previous data show that Cdc42

activation is lost at the ends during mitosis and is activated at the division site when the

cell forms the cytokinetic actomyosin ring. Moreover, delaying Cdc42 activation at the

division site did not change the timing of Cdc42 inactivation at the cell ends (Wei et al.,

2016). This indicates that Cdc42 inactivation at the ends is independent of its activation

at the division site. We analyzed the timing of CRIB-mCherry loss from the cell ends

with respect to mitotic progression, as indicated by Sad1-mCherry and changes in

Septin Spn3-GFP localization (Fig. 5A). We find that CRIB-mCherry is lost from the cell

ends about 10-20 minutes after the onset of mitosis. In contrast the septin ring forms at

the end of Anaphase B (Fig.5C). This indicates that as the cells enter mitosis, septin

levels at the cell sides drop, allowing Rga4 mobility, which coincides with the loss of

active Cdc42 from the cell ends.

Our previous report showed that Cdc42 is reactivated at the cell ends as cytokinesis

completes to enable old end growth (Rich-Robinson et al., 2021). We asked if Cdc42

reactivation at the ends is delayed in *spn1Δ* mutants. Indeed, we find that in *spn1+* cells

Cdc42 is reactivated at the cell ends about 10 minutes after the completion of

actomyosin ring constriction as indicated by the probe CRIB-3xGFP and ring marker

Rlc1-tdTomaton (Fig.5B,D). In *spn1Δ* mutants, Cdc42 reactivation at the ends occurs

later, about 15 minutes after the completion of actomyosin ring constriction (Fig.5B,D).

This suggests that Cdc42 activity at the cell ends is disrupted in the absence of the

septin cytoskeleton.

Septins promote proper cell dimensions and bipolarity

Our data suggest that loss of septins from the cell cortex results in Rga4-mediated

inactivation of Cdc42 at the cell ends during mitosis. However, in *rga4Δ* mutants, we do

not see any significant change in Cdc42 reactivation at the cell ends during mitosis

(data not shown). However, these results were not unexpected, since we have

previously shown that precocious Cdc42 activity and consequent polarized growth at

the cell ends requires both the loss of *rga4*, and the activation of the MOR/Orb6

pathway (Gupta et al., 2014; Ray et al., 2010; Rich-Robinson et al., 2021). Thus, we

asked if septin-mediated Rga4 localization plays an important role at the cell ends

during interphase.

Our data show that in *spn1Δ* cells, the ends show increased Rga4-GFP levels, even at

the growing cell ends (Fig.1F). Thus, we asked of the presence of Rga4 at the cell ends

disrupts Cdc42 dynamics at these ends. We find that *spn1Δ* cells are more monopolar

than *spn1⁺* cells (Fig.6A,B). In addition, *spn1Δ* cells also showed an increase in cell

width (Fig. 6A,C). We asked if the increase in cell width was due to an increase in active

Cdc42 levels or its distribution at the growing cell ends. We find that compared to *spn1⁺*

cells, *spn1Δ* mutants show increased CRIB-GFP levels at the growing ends (Fig.6D,E).

In addition, the *spn1Δ* mutants also show an increase in the zone of distribution of

CRIB-GFP at the growing ends (Fig.6D,F). An analysis of CRIB-GFP fraction at the

ends of *spn1⁺* cells, appears symmetric between the two ends as the cells increase in

size beyond 9µm (Fig.6G). However, in *spn1Δ* cells, the ends show increased

asymmetry of the CRIB-GFP fraction between the two cell ends, and symmetry is

partially achieved only at a much higher cell length, beyond 11µm, just before the cells

start to divide (Fig.6G). Next, we analyzed the localization patterns of the Cdc42 GEF

Scd1 and its scaffold protein Scd2. We find that compared to *spn1⁺* cells, *spn1Δ*

mutants show increased levels of Scd1-mNeonGreen and Scd2-GFP levels at the

growing cell ends (Fig. 6 H-K). These data suggest that in the absence of the septin

cytoskeleton, Cdc42 activation dynamics between the two ends is disrupted and

accumulates only at one end, resulting in monopolarity and increased cell width. Thus,

we analyzed Cdc42 activation dynamics at the cell ends using the CRIB-3xGFP probe

in *spn1⁺* and *spn1Δ* cells. As reported earlier, CRIB-3xGFP shows anticorrelated

oscillations between the two cell ends in control cells (Das et al., 2012). However, in

spn1Δ cells, CRIB-3xGFP activity fluctuated only at the growing end, while the non-

growing end barely showed any signal. This indicates that the septin cytoskeleton is

required for proper Cdc42 activation dynamics between the two cell ends.

Discussion

Our previous report showed that Cdc42 activation at the cell ends is cell-cycle-dependent and is turned off during cell division (Rich-Robinson et al., 2021). Cdc42 reactivation at the cell ends depends on the activation of the MOR/Orb6 kinase pathway and absence of Rga4 from the cell ends (Gupta et al., 2014; Rich-Robinson et al., 2021). Here we show that cell-cycle-dependent Rga4 localization is dependent on the septin cytoskeleton. The septin cytoskeleton at the cell sides prevents lateral diffusion of Rga4 along the membrane and keeps it away from the cell ends where Cdc42 is activated for polarized growth. Once the cell enters mitosis, the septin cytoskeleton gradually drops from the cell sides and this corresponds with Rga4 diffused localization to the cell ends where Cdc42 is gradually inactivated.

After division in fission yeast, cells grow in a monopolar manner from old end and transition to bipolar growth when the new end takes off (NETO) (Mitchison and Nurse, 1985). Bipolar growth occurs when the old and new ends are able to compete with each other Cdc42 activation (Das et al., 2012). The zone of growth at the cell ends is determined by the Cdc42 activation dynamics and the GAP Rga4 (Das et al., 2012; Das et al., 2007). Rga4 localizes to the cell sides like a corset and prevents the overflow of Cdc42 activation from the ends to these regions. This helps to maintain the cell width and proper polarity. During cell division, Rga4 localizes to the ingressing septum and prevent excessive Cdc42 activation at that site (Campbell et al., 2022). While Rga4 is not present at the growing cell ends, in pre-NETO cells, Rga4 localizes to the non-growing ends. Recent reports show that Rga4 does not localize to the growing cell ends

due to plasma membrane flows as a result of growth (Gerganova et al., 2021;

Rutkowski et al., 2024). Thus, non-growing ends show Rga4 given the lack of

membrane flow at these ends. We find that in cells lacking the septin cytoskeleton,

Rga4 levels at both the cell ends increase. This increase is due to the increased

mobility of Rga4 along the membrane. However, the levels at the non-growing ends are

higher than that at the growing ends, which again can be explained by the effect of

growth-dependent membrane flows. In these mutants, the cell ends receive increased

levels of Rga4 due to diffusion, while the growth dependent membrane flow removes

Rga4. This explains why the Rga4 levels at these ends are not as high as that observed

at the cell sides but sufficient to disrupt Cdc42 activation.

Cdc42 activation dynamics at the cell ends depend on positive and time-delayed

negative feedbacks (Das et al., 2012; Harrell et al., 2024; Howell et al., 2012). These

feedbacks ensure that the two cell ends can sequentially activate Cdc42 to promote

bipolar growth. Previous reports from our lab have shown that the GEF Gef1 is required

for establishment of Cdc42 activation at the new end and thus promote bipolar growth

(Hercyk and Das, 2019; Hercyk et al., 2019). Any disruption in the Cdc42 activation

pattern results in its accumulation at one end at the expense of the other (Das et al.,

2012; Harrell et al., 2024). We also observe this in the septin mutants, where Cdc42

dynamics is disrupted, resulting in monopolar growth. The low levels of Rga4 at the cell

ends can alter the dynamics of Cdc42 activation and inactivation such that it

accumulates at one end.

Our data suggest that the septin cytoskeleton does not restrict Rga4 diffusion via

protein-protein interaction. This is in agreement with previous reports that show that the

septin cytoskeleton forms a physical barrier on the membrane (Spiliotis and McMurray,

2020). The septin filaments at the cell sides are organized along the long axis of the cell

while Rga4 on the membrane is organized in small puncta. The Rga4 puncta are mostly

distinct from the septin filaments. It is unclear how the septin filaments block Rga4

mobility given that these proteins do not colocalize. One possibility is that the septins

organize the lipid compartments on the plasma membrane as has been reported in

other systems, and this regulates Rga4 localization (Caudron and Barral, 2009). Indeed,

Rga4 protein contains a coiled-coil domain that is required for membrane localization

(Das et al., 2007). It is possible that the septin cytoskeleton regulates the plasma

membrane organization, which indirectly restricts Rga4 diffusion. We are currently

investigating whether membrane interactions mediate septin-dependent Rga4

localization.

The septin cytoskeleton in fission yeast has been investigated mostly in the context of

cytokinesis (An et al., 2004; Berlin et al., 2003; Martin-Cuadrado et al., 2005; Munoz et

al., 2014; Singh et al., 2024; Wang et al., 2015). In budding yeast, the septins organize

as a collar around the bud neck and forms a ring during cytokinesis (Bridges and

Gladfelter, 2015; Woods and Gladfelter, 2021). The Septin collar in budding yeast has

been shown to restrict GAP localization near the bud neck to promote bud emergence

for daughter cell formation (Okada et al., 2013). This indicates that restricting GAP

localization by septin filaments is a conserved phenomenon. In fission yeast, the septin

ring forms as anaphase B ends, when the actomyosin ring starts to constrict. However,

the septin filament at the cell sides begin to disappear at the onset of mitosis, well

before the septin ring forms. The septin proteins have membrane-interacting domains

and are involved in lipid metabolism (Mela and Momany, 2022; Spiliotis and McMurray,

2020; Woods and Gladfelter, 2021). It is possible that a mitotic signal disables the

septin-membrane interaction resulting in its loss from the cell sides. Further

investigation will determine what leads to the loss of septins from the cell sides in

mitosis and its subsequent ring formation.

Our findings here show a role for the septin cytoskeleton in regulating polarized cell

growth in a cell-cycle-dependent manner. While most cell cycle dependent regulation is

mediated via kinase-substrate phosphorylation (Kaldis, 1999; Malumbres, 2014;

McCusker and Kellogg, 2012), here we show a regulatory system where a physical

barrier by the septins blocks Rga4 mobility, thereby regulating Cdc42 activation

dynamics. While the septin rings have been mostly identified in fungal cells, it is

possible that septins in other eukaryotes also undergo cell-cycle-dependent changes in

their organization (Delic et al., 2024; Marquardt et al., 2021). In several eukaryotes,

septins have been shown to act as diffusion barriers and maintain asymmetry in the cell

(Caudron and Barral, 2009). Thus cell-cycle-dependent changes in septin organization

may alter their ability to act as barriers. Further, investigation will shed light on the

significance of cell-cycle-dependent septin organization and its role in polarity

maintenance.

Methods and Materials

Cell Culture

The *S. pombe* strains used in this study are listed in Table 1. Cells were cultured in

yeast extract with supplements (YES – Sunrise Science Products) medium and grown

exponentially at 25°C, unless specified otherwise. Standard techniques were used for

genetic manipulation and analysis (Moreno et al., 1991). Cells were grown exponentially

for at least three rounds of eight generations each before imaging.

Microscopy

Imaging was performed at room temperature (23–25°C). We used a spinning disk

confocal microscope system with a Nikon Eclipse inverted microscope with a 100×/1.49

NA lens, a CSU-22 spinning disk system (Yokogawa Electric Corporation) and a

Photometrics EM-CCD camera (Photometrics Technology Evolve with excelon Serial

No: A13B107000). Images were acquired with MetaMorph (Molecular Devices,

Sunnyvale, CA).

A Nikon Ti2 Eclipse wide field microscope was also used with a 100x/1.49 NA objective,

an ORCA-FusionBT digital camera (Hamamatsu Model: C15440-20UP Serial No:

500428, Hamamatsu, Hamamatsu City, Japan). Images were acquired using Nikon NIS

Elements (Nikon, Melville, NY). Fluorophores were excited using an AURA Light Engine

system (Lumencor, Beaverton, OR).

Additionally, microscopy was performed with a 3i spinning disk confocal using a Zeiss

AxioObserver microscope with integrated Yokogawa spinning disk (Yokogawa CSU-X1

A1 spinning disk scanner) and a 100x/1.49 NA objective. Images were acquired with a

Teledyne Photometrics Prime 95b back-illuminated sCMOS camera (Serial No:

A20D203014, Tucson, AZ). Images were acquired using SlideBook (3i Intelligent

Imaging innovations, Denver, CO).

Fluorescence Intensity Acquisition & Quantification

Image analysis was performed on a Macintosh computer using the public domain NIH

Image program (developed at the U.S. National Institutes of Health and available on the

Internet at http://rsb.info.nih.gov/nih-image/). Images were acquired at either single,

mid-section focal planes, or as Z-stacks with 0.2-0.4 µm focal plane intervals. Z-stack

images were superimposed using sum projection for analysis. Mean intensity

measurements were used for analysis, along with integrated density values where

specified. Prior to intensity acquisition, a region of interest devoid of cells was used for

background subtraction. Cell side coefficient of variation measurements were acquired

using the segmented line tool on both cell sides. Plot profiles of these lines were

generated, and the coefficient of variation was determined via ratio of the standard

deviation of intensity values along these lines, divided by each position's average

intensity along said line. Cell end intensities were measured using freehand ROIs of

sum projected Z-stacks.

Fluorescence recovery after photobleaching

Fluorescence recovery after photobleaching (FRAP) was performed as described

before (Onwubiko et al., 2019). Fluorescently tagged proteins along the cell cortex were

first focused at the medial z-plane and an ROI was selected for bleaching. The ROI was

imaged 10 times pre-bleaching and then bleached for 10 s and images were then

acquired every 3 s for 50–100 images. A non-bleached spot on the cortex for the same

cell was analyzed to account for loss of signal due to photobleaching over time. A cell-

free region of the image was used for background subtraction. Intensity values for each

ROI were corrected for loss of signal over time and background signal. The intensity

values were then normalized against the mean pre-bleach value to determine the

recovery fraction over time. At least ten cells were analyzed for each genotype for each

experiment in carried out triplicate. The average corrected and normalized intensity

values were then plotted as fraction of recovery over time. The fraction of recovery over

time was plotted   and smoothened using GraphPad *Prism*. The half-life (t1/2) was then

computed as time for recovery of 50% of the signal.

Supplemental information titles and legends

Document S1. Figures S1–S3 and Table S1

Data Availability

The data generated are available in the published article and its online supplemental

material.

Acknowledgements

We thank Kathy Gould, Fulvia Verde, Pilar Perez and Chuanhai Fu for providing strains

and Bret Judson at the Boston College Imaging Core for imaging support. This work is

supported by the National Institutes of Health grant R01GM136847 to M.D.

Author contributions

Conceptualization: M.D. and J.M.; Methodology: M.D. and J.M.; Validation: M.D. and J.M.;

Formal analysis: M.D. and J.M.; Investigation: M.D., J.M., G.C.; Resources: M.D.; Data

curation: J.M. and G.C.; Writing- original draft: M.D. and J.M. ; Writing - review & editing:

M.D.; Supervision: M.D.; Project administration: M.D.; Funding acquisition: M.D.

Declaration of interests

The authors declare no competing or financial interests.

References

An, H., J.L. Morrell, J.L. Jennings, A.J. Link, and K.L. Gould. 2004. Requirements of fission yeast septins for complex formation, localization, and function. *Molecular biology of the cell*. 15:5551-5564.

Berlin, A., A. Paoletti, and F. Chang. 2003. Mid2p stabilizes septin rings during cytokinesis in fission yeast. *The Journal of cell biology*. 160:1083-1092.

Bos, J.L., H. Rehmann, and A. Wittinghofer. 2007. GEFs and GAPs: critical elements in the control of small G proteins. *Cell*. 129:865-877.

Bridges, A.A., and A.S. Gladfelter. 2015. Septin Form and Function at the Cell Cortex. *The Journal of biological chemistry*. 290:17173-17180.

Campbell, B.F., B.S. Hercyk, A.R. Williams, E. San Miguel, H.G. Young, and M.E. Das. 2022. Cdc42 GTPase activating proteins Rga4 and Rga6 coordinate septum synthesis and membrane trafficking at the division plane during cytokinesis. *Traffic*. 23:478-495.

Caudron, F., and Y. Barral. 2009. Septins and the lateral compartmentalization of eukaryotic membranes. *Developmental cell*. 16:493-506.

Chang, E.C., M. Barr, Y. Wang, V. Jung, H.P. Xu, and M.H. Wigler. 1994. Cooperative interaction of S. pombe proteins required for mating and morphogenesis. *Cell*. 79:131-141.

Coll, P.M., Y. Trillo, A. Ametzazurra, and P. Perez. 2003. Gef1p, a new guanine nucleotide exchange factor for Cdc42p, regulates polarity in Schizosaccharomyces pombe. *Molecular biology of the cell*. 14:313-323.

Das, M., T. Drake, D.J. Wiley, P. Buchwald, D. Vavylonis, and F. Verde. 2012. Oscillatory Dynamics of Cdc42 GTPase in the Control of Polarized Growth. *Science*. 337:239-243.

Das, M., and F. Verde. 2013. Role of Cdc42 dynamics in the control of fission yeast cell
polarization. *Biochemical Society transactions*. 41:1745-1749.

Das, M., D.J. Wiley, S. Medina, H.A. Vincent, M. Larrea, A. Oriolo, and F. Verde. 2007.
Regulation of cell diameter, For3p localization, and cell symmetry by fission yeast
Rho-GAP Rga4p. *Molecular biology of the cell*. 18:2090-2101.

Delic, S., B. Shuman, S. Lee, S. Bahmanyar, M. Momany, and M. Onishi. 2024. The
evolutionary origins and ancestral features of septins. *Front Cell Dev Biol*.
12:1406966.

Estravis, M., S. Rincon, and P. Perez. 2012. Cdc42 regulation of polarized traffic in
fission yeast. *Communicative & integrative biology*. 5:370-373.

Etienne-Manneville, S. 2004. Cdc42--the centre of polarity. *Journal of cell science*.
117:1291-1300.

Galliot, B., and L. Ghila. 2010. Cell plasticity in homeostasis and regeneration. *Mol
Reprod Dev*. 77:837-855.

Gallo Castro, D., and S.G. Martin. 2018. Differential GAP requirement for Cdc42-GTP
polarization during proliferation and sexual reproduction. *The Journal of cell
biology*. 217:4215-4229.

Gerganova, V., I. Lamas, D.M. Rutkowski, A. Vjestica, D.G. Castro, V. Vincenzetti, D.
Vavylonis, and S.G. Martin. 2021. Cell patterning by secretion-induced plasma
membrane flows. *Sci Adv*. 7:eabg6718.

Guillot, C., and T. Lecuit. 2013. Mechanics of epithelial tissue homeostasis and
morphogenesis. *Science*. 340:1185-1189.

Gupta, S., M. Govindaraghavan, and D. McCollum. 2014. Cross talk between NDR
kinase pathways coordinates cytokinesis with cell separation in
Schizosaccharomyces pombe. *Eukaryotic cell*. 13:1104-1112.

Harrell, M.A., Z. Liu, B.F. Campbell, O. Chinsen, T. Hong, and M. Das. 2024. Arp2/3-
dependent endocytosis ensures Cdc42 oscillations by removing Pak1-mediated
negative feedback. *The Journal of cell biology*. 223.

Hercyk, B.S., and M.E. Das. 2019. F-BAR Cdc15 Promotes Cdc42 Activation During
Cytokinesis and Cell Polarization in Schizosaccharomyces pombe. *Genetics*.
213:1341-1356.

Hercyk, B.S., J. Rich-Robinson, A.S. Mitoubsi, M.A. Harrell, and M.E. Das. 2019. A
novel interplay between GEFs orchestrates Cdc42 activity during cell polarity and
cytokinesis in fission yeast. *Journal of cell science*. 132.

Howell, A.S., M. Jin, C.F. Wu, T.R. Zyla, T.C. Elston, and D.J. Lew. 2012. Negative
feedback enhances robustness in the yeast polarity establishment circuit. *Cell*.
149:322-333.

Johnson, D.I. 1999. Cdc42: An essential Rho-type GTPase controlling eukaryotic cell
polarity. *Microbiology and molecular biology reviews : MMBR*. 63:54-105.

Kaldis, P. 1999. The cdk-activating kinase (CAK): from yeast to mammals. *Cellular and
molecular life sciences : CMLS*. 55:284-296.

Kinoshita, M. 2003. The septins. *Genome Biol*. 4:236.

Lloyd, A.C. 2013. The regulation of cell size. *Cell*. 154:1194-1205.

Maddox, A.S., L. Lewellyn, A. Desai, and K. Oegema. 2007. Anillin and the septins
promote asymmetric ingression of the cytokinetic furrow. *Developmental cell*.
12:827-835.

Malumbres, M. 2014. Cyclin-dependent kinases. *Genome Biol*. 15:122.

Marquardt, J., X. Chen, and E. Bi. 2021. Septin Assembly and Remodeling at the Cell
Division Site During the Cell Cycle. *Front Cell Dev Biol*. 9:793920.

Martin-Cuadrado, A.B., J.L. Morrell, M. Konomi, H. An, C. Petit, M. Osumi, M.
Balasubramanian, K.L. Gould, F. Del Rey, and C.R. de Aldana. 2005. Role of
septins and the exocyst complex in the function of hydrolytic enzymes
responsible for fission yeast cell separation. *Molecular biology of the cell*.
16:4867-4881.

Mata, J., and P. Nurse. 1997. tea1 and the microtubular cytoskeleton are important for
generating global spatial order within the fission yeast cell. *Cell*. 89:939-949.

McCusker, D., and D.R. Kellogg. 2012. Plasma membrane growth during the cell cycle:
unsolved mysteries and recent progress. *Current opinion in cell biology*. 24:845-
851.

Mela, A., and M. Momany. 2022. Septins coordinate cell wall integrity and lipid
metabolism in a sphingolipid-dependent process. *Journal of cell science*. 135.

Merla, A., and D.I. Johnson. 2000. The Cdc42p GTPase is targeted to the site of cell
division in the fission yeast Schizosaccharomyces pombe. *European journal of*
cell biology. 79:469-477.

Mitchison, J.M., and P. Nurse. 1985. Growth in cell length in the fission yeast
Schizosaccharomyces pombe. *Journal of cell science*. 75:357-376.

Moreno, S., A. Klar, and P. Nurse. 1991. Molecular genetic analysis of fission yeast
Schizosaccharomyces pombe. *Methods in enzymology*. 194:795-823.

Munoz, S., E. Manjon, and Y. Sanchez. 2014. The putative exchange factor Gef3p
interacts with Rho3p GTPase and the septin ring during cytokinesis in fission
yeast. *The Journal of biological chemistry*. 289:21995-22007.

Murray, J.M., and D.I. Johnson. 2001. The Cdc42p GTPase and its regulators Nrf1p and
Scd1p are involved in endocytic trafficking in the fission yeast
Schizosaccharomyces pombe. *The Journal of biological chemistry*. 276:3004-
3009.

Okada, S., M. Leda, J. Hanna, N.S. Savage, E. Bi, and A.B. Goryachev. 2013. Daughter
cell identity emerges from the interplay of Cdc42, septins, and exocytosis.
Developmental cell. 26:148-161.

Onishi, M., T. Koga, A. Hirata, T. Nakamura, H. Asakawa, C. Shimoda, J. Bahler, J.Q.
Wu, K. Takegawa, H. Tachikawa, J.R. Pringle, and Y. Fukui. 2010. Role of septins
in the orientation of forespore membrane extension during sporulation in fission
yeast. *Molecular and cellular biology*. 30:2057-2074.

Onwubiko, U.N., P.J. Mlynarczyk, B. Wei, J. Habiyaremye, A. Clack, S.M. Abel, and
M.E. Das. 2019. A Cdc42 GEF, Gef1, through endocytosis organizes F-BAR
Cdc15 along the actomyosin ring and promotes concentric furrowing. *Journal of*
cell science. 132.

Ray, S., K. Kume, S. Gupta, W. Ge, M. Balasubramanian, D. Hirata, and D. McCollum.
2010. The mitosis-to-interphase transition is coordinated by cross talk between
the SIN and MOR pathways in Schizosaccharomyces pombe. *The Journal of cell*
biology. 190:793-805.

Revilla-Guarinos, M.T., R. Martin-Garcia, M.A. Villar-Tajadura, M. Estravis, P.M. Coll,
and P. Perez. 2016. Rga6 is a Fission Yeast Rho GAP Involved in Cdc42
Regulation of Polarized Growth. *Molecular biology of the cell*.

Rich-Robinson, J., A. Russell, E. Mancini, and M. Das. 2021. Cdc42 reactivation at
growth sites is regulated by local cell-cycle-dependent loss of its GTPase-
activating protein Rga4 in fission yeast. *Journal of cell science*. 134.

Rutkowski, D.M., V. Vincenzetti, D. Vavylonis, and S.G. Martin. 2024. Cdc42 mobility
and membrane flows regulate fission yeast cell shape and survival. *Nat*
Commun. 15:8363.

Singh, D., Y. Liu, Y.H. Zhu, S. Zhang, S. Naegele, and J.Q. Wu. 2024. Septins function
in exocytosis via physical interactions with the exocyst complex in fission yeast
cytokinesis. *bioRxiv*.

Sparks, C.A., M. Morphew, and D. McCollum. 1999. Sid2p, a spindle pole body kinase
that regulates the onset of cytokinesis. *The Journal of cell biology*. 146:777-790.

Spiliotis, E.T., and M.A. McMurray. 2020. Masters of asymmetry - lessons and
perspectives from 50 years of septins. *Molecular biology of the cell*. 31:2289-
2297.

Sprang, S. 2001. GEFs: master regulators of G-protein activation. *Trends in biochemical*
sciences. 26:266-267.

Tatebe, H., K. Nakano, R. Maximo, and K. Shiozaki. 2008. Pom1 DYRK regulates
localization of the Rga4 GAP to ensure bipolar activation of Cdc42 in fission
yeast. *Current biology : CB*. 18:322-330.

Verde, F., J. Mata, and P. Nurse. 1995. Fission yeast cell morphogenesis: identification
of new genes and analysis of their role during the cell cycle. *The Journal of cell*
biology. 131:1529-1538.

Wang, N., I.J. Lee, G. Rask, and J.Q. Wu. 2016. Roles of the TRAPP-II Complex and
the Exocyst in Membrane Deposition during Fission Yeast Cytokinesis. *PLoS*
biology. 14:e1002437.

Wang, N., M. Wang, Y.H. Zhu, T.W. Grosel, D. Sun, D.S. Kudryashov, and J.Q. Wu.
2015. The Rho-GEF Gef3 interacts with the septin complex and activates the
GTPase Rho4 during fission yeast cytokinesis. *Molecular biology of the cell*.
26:238-255.

Wei, B., B.S. Hercyk, J. Habiyaremye, and M. Das. 2017. Spatiotemporal Analysis of
Cytokinetic Events in Fission Yeast. *J Vis Exp*:e55109.

Wei, B., B.S. Hercyk, N. Mattson, A. Mohammadi, J. Rich, E. DeBruyne, M.M. Clark,
and M. Das. 2016. Unique Spatiotemporal Activation Pattern of Cdc42 by Gef1
and Scd1 Promotes Different Events during Cytokinesis. *Molecular biology of the*
cell. 27:1235-1245.

Woods, B.L., and A.S. Gladfelter. 2021. The state of the septin cytoskeleton from

676   assembly to function. *Current opinion in cell biology*. 68:105-112.

677   Zheng, S., B. Zheng, Z. Liu, X. Ma, X. Liu, X. Yao, W. Wei, and C. Fu. 2022. The Cdc42

678   GTPase-activating protein Rga6 promotes the cortical localization of septin.

679   *Journal of cell science*. 135.

Figure Legends

Figure 1. The septin cytoskeleton restricts Rga4 to the cell sides during interphase. A. Middle plane images of Rga4-GFP in *spn1*⁺ and *spn1Δ* cells. Red brackets highlight Rga4 distribution along cell sides in interphase cells. Red asterisks highlight dividing cells. **B-C**. Quantification of the coefficient of variation of Rga4 distribution through the cell-cycle in *spn1*⁺ (B) and *spn1Δ* (C) cells. [N=3, *n*=40-50 cells each, *$P≤0.05$, *ns*=not significant,]. **D.** Rga4-GFP increasingly localizes to the cell ends and cytoplasm in *spn1Δ* cells during interphase. Images consist of superimposed middle-plane images taken at 10 second intervals over 5 minute timelapse. Red boxes highlight increased localization of Rga4 to the cell ends. **E.** Quantification of Rga4-GFP intensity within the cytoplasm of interphase cells in *spn1*⁺ and *spn1Δ* conditions measured by averaging three unique regions within the cytoplasm of each cell [**$P≤0.01$, Student's T test]. **F.** Quantification of Rga4-GFP localization to growing and non-growing cell ends of interphase cells in *spn1*⁺ and *spn1Δ* conditions. [N=3, *n*=24-28 cells each, ****$P≤0.0001$, ***$P≤0.001$, **$P≤0.01$, *$P≤0.05$, *ns*=not significant, one-way ANOVA, scale bar=5µm].

Figure 2. Septins inhibit Rga4-GFP mobility along the plasma membrane during Interphase. A. Montage of Rga4-GFP in *spn1*⁺ cells during interphase, during mitosis, and in *spn1Δ* cells during interphase, taken over 5 minute timelapse. Red arrowhead marks an Rga4-GFP puncta. **B.** Montage of Rga4-GFP recovery after photobleaching during interphase and mitosis in *spn1*⁺ and *spn1Δ* cells taken in 15 second intervals. Red circles indicate the photobleached region of interest (ROI). Red lightning bolt

indicates the timepoint of photobleaching. **C.** FRAP recovery rates of Rga4-GFP signal

intensity within the ROI over time in *spn1⁺* and *spn1Δ* cells during interphase and

mitosis. **D.** $T_{1/2}$ fluorescence recovery times of listed conditions. [N=3, *n*=10 cells each,

**$P≤0.01$, *$P≤0.05$, *ns*=not significant, one-way ANOVA, scale bar=5µm].

Figure 3. Septins form linear filaments along the cell sides in interphase and a

ring at the division site as Anaphase B ends. A. Spn3-GFP and Spn1-tdT colocalize

in linear filaments along the cell cortex during interphase, and within a ring during

division. Red and white arrowheads highlight filaments. **B.** Spn2-GFP show linear

filament like organization along the cell sides during interphase and a ring at the division

site during mitosis. Red arrowhead marks a filament. **C.** Spn4-tdTomato localize along

the cell sides during interphase in a punctate pattern and form a ring at the division site

during mitosis. Red arrowhead marks punctate organization. **D.** Montage of timelapse of

Spn4-tdTomato and Moyo2-GFP Sid4-GFP expressing cells during mitosis. Red asterisk

marks onset of mitosis, the green asterisk marks onset of septin ring formation and

purple asterisk marks end of mitosis. **E.** Quantification over time of Spn4-TdTomato

levels along the cell side of the cell shown in D. F. Quantification of Spn4-tdTomato

along the cell sides with increasing spindle pole body distances. [N=3, n=>20, *$P≤0.05$,

one-way ANOVA, scale bar=5µm or 2µm as indicated]

Figure 4. The septin cytoskeleton does not colocalize or interact with Rga4. A.

Rga4-GFP and Spn1-tdTomato show distinct localization pattern along the sides of

interphase cells. Red arrowhead zones with Rga4-GFP but no Spn1-tdTomato along the

cell side. Middle plane inset shows partial colocalization of Rga4-GFP with Spn1-

tdTomato. Top plane inset shows distinct localization patterns of these proteins. **B.**

Spn3-GFP mislocalizes to Tom20-GBP-RFP at the mitochondria. **C.** Spn3-GFP and

Spn1-tdT simultaneously mislocalize to the mitochondria in Tom20-GBP expressing

cells. **D.** Rga4-GFP mislocalizes to the mitochondria, but not Spn1-tdT in Tom20-GBP

expressing cells. **E.** Spn3-GFP mislocalizes to the mitochondria, but not Rga4-mCh in

Tom20-GBP expressing cells. **F.** Quantification of the coefficient of variation of Rga4-

mCherry distribution along cell sides in interphase cells, Spn3-GFP and Tom20-GBP

expressing cells as shown in E. [N=3, *n*=20-26 cells each, *P≤0.05, student's t-test,

scale bar=5µm].

Figure 5. As septins recede from the cell sides, Rga4 spreads out along the

membrane and Cdc42 activity diminishes at the cell ends. A. A representative

timelapse of Spn3-GFP, CRIB-mCherry and Sad1-mCherry expressing cells starting at

the onset of mitosis. B. A representative timelapse of CRIB-3xGFP and Rlc1tdTomato in

spn1+ and *spn1Δ* cells. Red asterisk marks onset of stable CRIB-3xGFP localization at

the cell ends after the completion of ring constriction marked with a purple asterisk. **C.**

Frequency plot of the timing of loss of active Cdc42 (CRIB-mCherry) from the cell ends,

septin ring formation and end of Anaphase B with respect to the onset of mitosis.

(n=10). **D.** Frequency plot of the timing of the onset of stable Cdc42 activity at the cell

ends as indicated by CRIB-3xGFP after the completion of ring constriction in *spn1+* and

spn1Δ cells (n>30). Scale bar=5µm.

Figure 6. The septin cytoskeleton promotes bipolar growth. A. Representative

images of calcofluor white-stained *spn1⁺* and *spn1Δ* cells. Red asterisks highlight

bipolar (*spn1⁺*) and monopolar (*spn1Δ*) growth. **B.** Quantification of monopolarity index

in *spn1⁺* and *spn1Δ* cells. **C.** Quantification of cell width in *spn1⁺* and *spn1Δ* cells. [N=3,

n=65-100 cells each, *P≤0.05, student's t-test]. **D.** Representative Maximum Z-

Projection images of *spn1⁺* and *spn1Δ* cells expressing CRIB-GFP. Red arrowheads

highlight Cdc42 activity at the ends of growing cells. **E.** Quantification of CRIB-GFP

intensity at growing cell ends in *spn1⁺* and *spn1Δ* cells. **F.** Quantification of the length of

CRIB-GFP cap at growing cell ends in *spn1⁺* and *spn1Δ* cells. [N=3, *n*=65-100 cells

each, *P≤0.05, **P≤0.01, student's t-test]. **G.** Heat map of the fraction of CRIB-GFP

intensity at each cell end over the total intensity with increasing cell size in *spn1⁺* and

spn1Δ cells. **H, I.** Quantification of Scd1-mNG and Scd2-GFP at the ends of *spn1⁺* and

spn1Δ cells [N=3, *n*=65-100 cells each, *P≤0.05, **P≤0.01, student's t-test]. **J,K.**

Representative images of Scd1-mNG and Scd2-GFP (Red arrowheads) at the ends of

spn1⁺ and *spn1Δ* cells. scale bar=5µm

Figure 1

A

spn1⁺   *spn1Δ*

B Rga4-GFP heterogeneity along cell sides in *spn1+* cells

C Rga4-GFP heterogeneity along cell sides in *spn1Δ* cells

D

spn1⁺

spn1Δ

E Rga4-GFP Cytoplasmic Intensity

F Rga4-GFP Intensity at Cell Ends

Figure 2

Figure 3.

Figure 4.

Figure 5.

Figure 6.