Septins mediate cell-cycle-dependent exclusion of Cdc42 GAP Rga4 from growth

sites in fission yeast

Justin McDuffie, Goeun Chang, Maitreyi Das*

Biology Department, Boston College, Chestnut Hill, MA, 02467

*Corresponding author: Maitreyi.Das@bc.edu

Running title: Septins restrict Rga4 localization

Keywords: Septin, Cell cycle, Cdc42, GAP, Rga4, Cell polarity

Summary: Cell-cycle-dependent Rga4 localization pattern is controlled by the septin

cytoskeleton to promote bipolar growth. In interphase, the septin cytoskeleton at the cell

sides restricts Rga4 mobility and localization away from the cell ends to promote proper

Cdc42 activation dynamics.

ABSTRACT

The polarized growth promoter Cdc42 is inactivated at the growing cell ends during

mitosis and reactivated after division. In fission yeast, the Cdc42 inactivator/GAP Rga4

localizes to the cell ends during mitosis and is restricted to the sides in interphase. We

show that this cell-cycle-dependent Rga4 localization is septin-dependent. Septins form

linear filaments along the sides during interphase and recede to form a medial ring

during mitosis. In septin mutants, Rga4 is mobile, homogeneously distributed, and

increasingly localizes to the cell ends and cytoplasm, similar to mitotic cells. Accordingly,

septin mutants are monopolar with disrupted Cdc42 activation dynamics. This regulation

appears to be indirect, via septin-dependent lipid organization on the cortex. We find

that the septin cytoskeleton prevents excessive phosphatidylinositol 4,5-bisphosphate

$(PI(4,5)P_2)$ and its upstream kinase Its3 at the plasma membrane. Limiting $PI(4,5)P_2$

levels enhances Rga4 puncta at the cortex. Our data describe an unusual form of

polarity regulation where the septin cytoskeleton restricts GAP localization via

membrane lipid organization.

INTRODUCTION

Establishment and maintenance of cell shape is fundamental to its function. Most cells exhibit a consistent cell shape after every round of cell division, and this ensures their ability to maintain function. Consistent cell shape is not just observed in multicellular organisms that form highly organized tissues and organelles, but even in unicellular organisms such a yeast (Das and Verde, 2013; Galliot and Ghila, 2010; Guillot and Lecuit, 2013; Mitchison and Nurse, 1985). In most eukaryotes, polarized cell growth stops during division and resumes immediately after, to establish consistent cell shape (Lloyd, 2013; McCusker and Kellogg, 2012). It is not well understood how polarized growth resumes promptly after each round of cell division. To understand this process, we use the fission yeast model system *Schizosaccharomyces pombe.* These cells stop polarized growth from the cell ends during division and resume growth once the cells complete cytokinesis (Mitchison and Nurse, 1985). However, we have previously shown that resumption of polarized growth from the cell ends is independent of cytokinesis and is instead cell-cycle-dependent and occurs at a fixed time after mitosis (Rich-Robinson et al., 2021).

Like in many eukaryotic cells, the major regulator of polarized growth in fission yeast is the small GTPase Cdc42 (Das and Verde, 2013; Estravis et al., 2012; Etienne-Manneville, 2004; Johnson, 1999). Cdc42 is activated when Guanine nucleotide exchange factors (GEFs) help it bind GTP (Bos et al., 2007; Sprang, 2001). GTP-bound Cdc42 binds and activates downstream targets to promote polarized growth (Bos et al., 2007). Cdc42 is inactivated when it hydrolyses the GTP to GDP with the help of the GTPase-activating proteins (GAPs) (Bos et al., 2007). Cdc42 in fission yeast has two

GEFs, Scd1 and Gef1, and three GAPs, Rga4, Rga6, and Rga3 (Chang et al., 1994;

Coll et al., 2003; Das et al., 2007; Gallo Castro and Martin, 2018; Murray and Johnson,

2001; Revilla-Guarinos et al., 2016; Tatebe et al., 2008). Our lab previously showed that

during cell division, the GAP Rga4 changes its localization pattern (Rich-Robinson et

al., 2021). Rga4 localizes to the cell sides in a punctate manner during interphase. It

becomes diffuse along the cell cortex and localizes to the cell tips in dividing cells,

where it prevents Cdc42 activation. Once cells complete division, Rga4 resumes its

punctate localization away from the cell ends, and growth at the ends resumes. It is not

known how Rga4 localization is regulated in a cell-cycle-dependent manner.

Rga4 has a coiled-coil region that is required for its localization to the cell cortex (Das et

al., 2007). Cells lacking *rga4* show increased width, since in these mutants, the zone of

Cdc42 activation and consequent growth spreads over a larger area at the cell ends

(Das et al., 2007; Tatebe et al., 2008). It is not known how Rga4 localization is limited to

the cell sides and away from the ends. A recent report shows that Rga4 is removed from

the cell ends as a result of membrane flow that occurs due to extensile growth at these

ends (Gerganova et al., 2021; Rutkowski et al., 2024). During mitosis, Rga4 is diffuse

along the membrane and appears at the cell ends (Rich-Robinson et al., 2021). It is not

known what prevents the diffuse Rga4 localization at the cell surface during interphase,

restricting it to the sides.

In several eukaryotic cells, the septin cytoskeleton acts as a diffusion barrier along the

membrane (Bridges and Gladfelter, 2015; Caudron and Barral, 2009; Kinoshita, 2003).

S. pombe expresses seven total septin proteins (Spn1-7p), however, only four are

involved in the regular cell cycle (Spn1-4p) (An et al., 2004; Berlin et al., 2003; Onishi et

al., 2010). The septin cytoskeleton is a hetero-octamer consisting of 2 sets of 4 peptides

Spn1-4 (An et al., 2004; Berlin et al., 2003). These peptides form hetero-octamers that

organize in a palindromic manner to form larger filaments (Kinoshita, 2003; Spiliotis and

McMurray, 2020; Woods and Gladfelter, 2021). The septin cytoskeleton plays a role in

cytokinesis, where it forms a ring at the division site. The septin ring acts as the

landmark where proteins required for cell separation during cytokinesis are delivered

(Berlin et al., 2003; Maddox et al., 2007; Martin-Cuadrado et al., 2005; Singh et al.,

2024; Spiliotis and McMurray, 2020). It is not clear if the septin cytoskeleton plays a role

during interphase. Cells lacking the septin cytoskeleton show cell separation defects but

remain polarized (An et al., 2004; Berlin et al., 2003; Martin-Cuadrado et al., 2005).

Here, we show that the septin cytoskeleton prevents Rga4 diffusion along the cell cortex

and restricts it to the cell sides. In cells lacking septin filaments, Rga4 at the cell sides is

highly mobile, appears diffuse along the cortex, and increasingly localizes to the cell

ends, similar to that observed in mitotic cells. Correspondingly, in cells lacking septins,

the Cdc42 activation dynamics are disrupted, resulting in monopolar cells. During

interphase, the septin cytoskeleton forms small filaments at the cortex that align along

the long axis of the cell. The septin levels along the cortex decrease as the cells enter

mitosis, and eventually, a septin ring forms at the end of anaphase B. Around this time,

Rga4 localization appears diffuse at the cortex, and active Cdc42 is lost from the cell

ends. We find that the mechanism for septin-dependent Rga4 localization is not via

direct protein-protein interaction as determined by acceptor photobleach FRET and the

GFP-GBP trap method. Instead, we find that upon loss of septin filaments, the lipid

organization on the plasma membrane is disrupted with enhanced $PI(4,5)P_2$ levels and

decreased PI(4)P levels at the cortex. Moreover, in these mutants, the localization of the

PI(4)P kinase Its3 is also disrupted. Correspondingly, in mutants lacking Its3 kinase

function, Rga4 shows enhanced cortical localization. These findings together suggest a

role for the septin cytoskeleton in maintaining proper lipid organization at the cortex,

thereby restricting Rga4 localization to the cell sides. These findings show that the

septin cytoskeleton maintains cell polarity in a cell-cycle-dependent manner.

RESULTS

The septin cytoskeleton restricts Rga4 to the cell sides during interphase

To investigate if the septin cytoskeleton restricts Rga4 localization, we analyzed Rga4-

GFP in $spn1^+$ and $spn1\Delta$ cells. In the $spn1^+$ cells, Rga4-GFP along the cell sides

appears punctate in interphase cells and diffuse in dividing cells. In contrast, $spn1\Delta$

cells show a more diffuse Rga4-GFP localization pattern, regardless of the cell cycle

stage (Fig.1A). We quantified the level of Rga4 heterogeneity along the cell sides by

measuring the coefficient of variance of Rga4-GFP intensity. As reported earlier, we find

that the coefficient of variance of Rga4-GFP is highest in the cells in the G2 phase of

the cell cycle and the lowest in the mitotic phase in $spn1^+$ cells (Fig. 1B) (Rich-Robinson

et al., 2021). This indicates that Rga4-GFP is more homogeneous along the cell sides in

mitotic cells and forms puncta during G2 or interphase. In $spn1\Delta$ cells, the coefficient of

variance in G2 cells is low and is similar to that in mitotic cells (Fig. 1C).

In mitotic cells, Rga4 appears more cytoplasmic and is also increasingly observed at the

cell ends. We asked if septins restrict cytoplasmic Rga4 and prevent its localization to

the cell ends. Rga4-GFP along the cell ends appears faint, even in Z-projected images,

due to the curvature of the cell and weak expression/signal. To overcome this, we took

rapid time-lapse movies with 5-second intervals over 5 mins of Rga4-GFP expressing

$spn1+$ and $spn1\Delta$ cells. We measured Rga4-GFP intensity in the cytoplasm, and the cell

ends of time-projected images of these movies (Fig.1D). We find that compared to

$spn1^+$ cells, cytoplasmic Rga4-GFP levels increase in $spn1\Delta$ cells (Fig.1E). Similarly,

$spn1\Delta$ cells show enhanced Rga4-GFP levels at the cell ends compared to $spn1^+$ cells

(Fig.1F). In control cells, Rga4-GFP shows elevated levels at the non-growing cell ends

compared to growing ends (Gerganova et al., 2021). We also see higher levels of Rga4-

GFP at the non-growing cell ends of *spn1Δ* cells, but these levels are much higher than

those in *spn1+* cells. Moreover, the growing ends in *spn1Δ* cells show higher Rga4-GFP

levels compared to growing ends of *spn1$^+$* cells. This indicates that in the absence of

septins, Rga4-GFP levels at the cell ends are elevated.

Septins inhibit Rga4-GFP mobility during Interphase

Next, we asked whether the increased homogeneity of Rga4-GFP along the plasma

membrane is the result of increased Rga4-GFP mobility or diffusion along the

membrane. The above-mentioned 5-minute time-lapse movies of Rga4-GFP localization

were analyzed, and we found that, in $spn1^+$ cells in interphase, Rga4-GFP formed a

number of bright, stationary puncta along the plasma membrane that remained

immobile throughout the course of time-lapse imaging (Fig. 1G). In contrast, $spn1^+$ cells

in mitosis, and $spn1\Delta$ cells at all stages of the cell-cycle showed limited Rga4-GFP

puncta formation, and lacked immobile Rga4-GFP puncta. In these cells, Rga4-GFP

appeared to diffuse more rapidly along the plasma membrane. To quantify this observed

change in Rga4 mobility, fluorescence recovery after photobleaching (FRAP)

microscopy was performed on 10x10-pixel ROIs of Rga4-GFP along the cell sides of

$spn1^+$ and $spn1\Delta$ cells in interphase and mitosis (Fig. 1H). Cell-cycle progression was

marked by the fluorescently tagged spindle pole body marker Sad1-mCherry and the

fluorescently tagged Myo2 light chain actomyosin ring marker Rlc1-tdTomato.

Fluorescence recovery curves and calculated recovery half-lives of Rga4-GFP in the

photobleached ROIs showed that Rga4-GFP was much less mobile along the plasma

membrane in interphase $spn1^+$ cells (Fig. 1I, J). In contrast, Rga4-GFP is more mobile

along the plasma membrane in $spn1^+$ cells during mitosis. Similar to mitotic $spn1^+$ cells,

Rga4-GFP is more mobile along the plasma membrane in $spn1\Delta$ cells, regardless of

cell-cycle stage. This indicates that in the presence of septin filaments, the diffusion of

Rga4-GFP is significantly restricted, leading to decreased protein mobility. This

decreased mobility along the plasma membrane may also explain the decreased

localization of Rga4-GFP to the cytoplasm and the cell ends in the presence of septins.

A previous report has indicated that the GAP Rga6 interacts with the septin filaments

and is required for proper septin localization at the cortex in interphase cells (Zheng et

al., 2022). Thus, we asked if the septin filaments also regulate Rga6 distribution along

the cortex. Similar to Rga4-GFP, Rga6-GFP appears punctate along the cell sides

(Supplementary Fig. S1A). However, in *spn1Δ* mutants, Rga6-GFP localization

remained the same, and the coefficient of variance along the cortex was not significantly

distinct between *spn1+* and *spn1Δ* cells, and throughout the cell cycle stages

(Supplementary Fig. S1B). This indicates that the septin cytoskeleton specifically

disrupts Rga4 distribution along the cortex.

Septins promote bipolar growth, separate from their role in facilitating proper cytokinesis and cell separation

Next, we asked if, in addition to the changes in Rga4 localization, the septin mutants displayed polarity defects. We observe that the growth patterns of *spn1Δ* mutants are altered compared to their *spn1+* counterparts. Notably, cells lacking the core septin *spn1* showed a number of septation defects as a result of failure to successfully separate after division (Fig. 2A, C), as has been reported before (An et al., 2004; Perez et al., 2015; Tasto et al., 2003; Wang et al., 2015; Wu et al., 2010; Zheng et al., 2018; Zheng et al., 2024). These septin mutants also show a significant change in their polarity, with 54% of the cells showing a monopolar growth pattern, as compared to *spn1+* cells (18%). The monopolar pattern in *spn1Δ* has been reported earlier, and this has been attributed to the cytokinesis defect observed in these cells, possibly making the new ends incapable of growth (An et al., 2004; Bohnert and Gould, 2012) (Fig. 2A, B). Similarly, we find that in the *spn4Δ* mutants, the cells show cytokinesis defects, as indicated by the high septation index and increased monopolarity (Fig. 2D- F). Both Spn1 and Spn4 are the core septin proteins and are necessary to form the septin oligomer (An et al., 2004). Previous reports have shown that Spn2 and Spn3 are not necessary to form the septin oligomer and their loss do not lead to cytokinetic defects (An et al., 2004). However, we find that even in *spn2Δ* and *spn3Δ* mutants, while the septation index is not elevated, they continue to show increased monopolarity similar to that observed in *spn1Δ* and *spn4Δ* mutants (Fig D-F). This suggests that the polarity defect observed in the septin mutants is independent of cytokinetic defects. Taken

185   together, these data show that the absence of any of these four septin monomers

186   results in polarity defects, even when cytokinetic and septation defects are not present.

Septins promote proper Cdc42 oscillatory dynamics

Our data show that in the absence of *spn1*, Rga4 localizes to the cell ends, and these cells are mostly monopolar. Given that Rga4 is a GAP for Cdc42, we asked if the septin cytoskeleton impacts Cdc42 activation levels at the cell ends. We find that neither the loss of *spn1*, or *rga4*, or both, result in significant changes to the average levels of Cdc42 activity at the growing ends of cells, as measured using the active Cdc42 bio-marker, CRIB-3xGFP (Fig. 3A, B).

As reported previously, an analysis of CRIB-3xGFP fraction at the ends of *spn1*[+] cells appears symmetric between the two ends as the cells increase in size beyond 9µm (Fig. 3C) (Das et al., 2012). However, in *spn1Δ* cells, the ends show increased asymmetry of the CRIB-3xGFP fraction between the two cell ends, and symmetry is partially achieved only at a much higher cell length, beyond 11µm, just before the cells start to divide (Fig. 3C). These data suggest that in the absence of the septin cytoskeleton, Cdc42 activation dynamics between the two ends is disrupted and accumulates only at one end, resulting in monopolarity. Next, we analyzed Cdc42 oscillatory dynamics at the cell ends using the CRIB-3xGFP probe in *spn1*[+] and *spn1Δ* cells. As reported earlier, CRIB-3xGFP shows anticorrelated oscillations between the two cell ends in control cells (Das et al., 2012). However, in *spn1Δ* cells, CRIB-3xGFP activity fluctuated only at the growing end, while the non-growing end barely showed any signal (Fig. 3D- G). While some activation of Cdc42 was transiently observed at the non-growing end, it was swiftly deactivated before establishing a competitive draw for the Cdc42 activators at the growing cell end.

We have previously shown that precocious Cdc42 activity and consequent polarized

growth at the cell ends require both the loss of *rga4* and the activation of the MOR/Orb6

pathway (Gupta et al., 2014; Ray et al., 2010; Rich-Robinson et al., 2021). Cdc42 is

reactivated at the cell ends as cytokinesis completes to enable old end growth (Rich-

Robinson et al., 2021). We asked if Cdc42 reactivation at the ends is delayed in *spn1Δ*

mutants. Indeed, we find that in *spn1+* cells, Cdc42 is reactivated at the cell ends about

10 minutes after the completion of actomyosin ring constriction as indicated by the

probe CRIB-3xGFP and ring marker Rlc1-tdTomato (Fig. 3H, I). In *spn1Δ* mutants,

Cdc42 reactivation at the ends occurs later, about 15 minutes after the completion of

actomyosin ring constriction (Fig. 3H, I). This suggests that Cdc42 activity at the cell

ends is disrupted in the absence of the septin cytoskeleton. These findings indicate that

the septin cytoskeleton regulates cell-cycle-dependent Rga4 localization, along with

proper Cdc42 activation dynamics at the cell ends.

The septin cytoskeleton shows distinct localization patterns during interphase

and mitosis.

Given that the septin cytoskeleton promotes proper cell polarization regardless of a

cytokinetic defect, we next investigated how septins organize within the cell in

interphase as the cell grows. Previous reports have shown that in interphase, the septin

cytoskeleton forms small filaments along the plasma membrane (An et al., 2004;

Gregory et al., 2025; Zheng et al., 2018). However, epitope tagging of individual septin

proteins, such as Spn1 with tdTomato, shows some functional deficiencies (Gregory et

al., 2025). To determine the proper organization of the septin proteins, we first

investigated the localization patterns of the individual septin proteins during interphase.

We find that Spn1-mEGFP shows very faint cortical puncta during interphase, as has

been reported earlier (Fig.4A) (Gregory et al., 2025). With a brighter fluorophore,

tdTomato, Spn1 appears to localize to the plasma membrane as short linear filaments

along the long axis of the cell (Fig. 4A). Similar short filaments were also visible with

Spn2-GFP, Spn3-GFP, and Spn4-tdTomato (Fig. 4A).

We next asked if these interphase filaments were dependent on the core septin protein

Spn1. We find that in the absence of *spn1*, Spn2-GFP, Spn3-GFP, and Spn4-tdTomato

are cytoplasmic and fail to localize to the plasma membrane (Fig.4B). These proteins

also fail to form the septin ring in dividing cells, as has been reported before. This

indicates that the core septin protein Spn1 is required for septin filament formation in

both interphase and mitosis. However, the filaments are distinctly organized during

these two phases.

Next, we asked if the septin proteins colocalize to these filaments on the plasma

membrane in interphase. Previous reports suggest that Spn1 proteins tolerate tagging

with monomeric fluorophores better than dimeric tags (Gregory et al., 2025). Thus, we

first investigated Spn1-tagRFP and Spn3-GFP colocalization in interphase. We find that

both Spn3-GFP and Spn1-tagRFP colocalize to the linear filaments along the long axis

of the cell (Fig. 4D). We next verified this interaction using strains co-expressing Spn1-

tdTomato and Spn3-GFP. Spn1-tdTomato and Spn3-GFP also co-localize as short linear

filaments along the cell cortex (Fig. 4D).

We also find that septin filament distribution appears biased towards the growing cell

ends. To confirm this, we investigated Spn3-GFP localization in *tea1+* and *tea1Δ*

mutants. The *tea1Δ* mutants are monopolar and only grow from one end (Mata and

Nurse, 1997; Verde et al., 1995). We find that in *tea1Δ* cells, Spn3-GFP filaments

appear closer to the growing cell ends (Supplementary Fig. S2A, red arrowheads). In

tea1Δ cells, Spn3-GFP levels are higher at the growing ends compared to the *tea1+*

cells (Supplementary Fig. S2B). This is likely due to the fact that in *tea1+* cells, Spn3-

GFP is distributed between the two growing ends of bipolar cells, while in *tea1Δ* cells, all

of the Spn3-GFP accumulates at the single growing end.

Since the septin cytoskeleton appears to restrict Rga4 mobility, we asked whether the

septin proteins themselves were mobile along the cell cortex. To test for mobility, we

performed FRAP analysis of Spn1-tdTomato along the cell sides in both interphase and

dividing cells, and analyzed the fluorescence recovery over a 5-minute timespan.

Regardless of the cell-cycle stage, we barely see any recovery of the fluorescent signal

5 minutes post-photobleaching of Spn1-tdTomato (Supplementary Fig. S3A & B). This

indicates that the septin filaments observed along the cell sides are stable and relatively

immobile.

Septin filaments recede from the cell sides as the cell enters mitosis

Our data show that Rga4 localization at the cortex is mostly punctate and immobile in

interphase and diffused and mobile in dividing cells. In addition, we find that the septins

show short filaments along the cell cortex in interphase, and this localization is lost once

the cell enters mitosis and a septin ring forms at the division site. Based on this, we

propose that Rga4 is more dynamic at the cortex when the septins are lost from this

region. To verify this hypothesis, we observed Rga4-GFP and Spn1-tdTomato

localization along the cortex in cells at different cell-cycle stages (Supplementary

Fig.S4A). As expected, we find that Rga4-GFP appears punctate along the cell sides in

early and late G2 cells. Spn1-tdTomato also localizes to this region. In early and late

mitotic cells, Rga4-GFP appears more diffuse, and Spn1-tdTomato levels along the

sides diminish. Given the low expression levels of Rga4, it was not possible to image

Rga4-GFP by time-lapse microscopy throughout the different cell-cycle stages along

with Spn1-tdTomato. The fully matured Spn1-tdTomato ring appears in post-mitotic cells

while Rga4-GFP shows diffuse localization in mitotic cells as the septin localization

along the cortex decreases (Supplementary Fig.S4A, early and late mitosis). This

suggests that the removal of septins from the cell cortex, and not the formation of the

septin ring, is sufficient to enable diffuse Rga4 localization.

Cdc42 is active at the cell ends in interphase and at the division site in dividing cells

(Merla and Johnson, 2000; Wei et al., 2016). Our previous data show that Cdc42

activation is lost at the ends during mitosis and is activated at the division site when the

cell forms the cytokinetic actomyosin ring. Moreover, delaying Cdc42 activation at the

division site did not change the timing of Cdc42 inactivation at the cell ends (Wei et al.,

2016). This indicates that Cdc42 inactivation at the ends is independent of its activation

at the division site. We analyzed the timing of CRIB-mCherry loss from the cell ends

with respect to mitotic progression, as indicated by Sad1-mCherry and changes in

septin Spn3-GFP localization (Fig. 5A). We find that CRIB-mCherry is lost from the cell

ends about 10-20 minutes after the onset of mitosis. In contrast, the septin ring forms at

the end of Anaphase B (Fig. 5B). This suggests that as the cells enter mitosis, septin

levels at the cell sides drop, Rga4 localization changes, and this coincides with the loss

of active Cdc42 from the cell ends.

Next, we investigated how septin localization changes throughout the cell cycle. We first

imaged cells expressing Spn4-tdTomato along with Myo2-GFP and Sid4-GFP over time-

lapse (Fig. 5C). We analyzed Spn4-tdTomato as this is one of the core septin proteins,

along with Spn1, required for timely cytokinesis and proper cell separation. While Sid4-

GFP labels the spindle pole body and helps visualize the onset of mitosis when the

spindle pole body splits into two, Myo2-GFP labels the cytokinetic actomyosin ring

(Sparks et al., 1999; Wei et al., 2017). We find that Spn4-tdTomato intensity along the

cell cortex drops as the cell enters mitosis, and eventually plateaus (Fig.5D). The Spn4-

tdTomato eventually forms a septin ring just prior to mitotic exit, 18 minutes after spindle

pole body separation. We quantified Spn4-tdTomato levels along the sides of cells in

different mitotic stages, as determined by the distance between spindle pole bodies. We

find that Spn4-tdTomato levels along the cell sides are highest in interphase cells with a

single spindle pole body (distance, 0μm, Fig. 5E). As the spindle pole body distance

increases, Spn4-tdTomato levels along the cell sides drop.

We further looked at Spn3-GFP localization throughout the cell cycle, as its localization

is more easily visualized through fluorescence microscopy. Similar to Spn4-tdTomato,

Spn3-GFP distribution along the cell sides also begins to change as cells enter mitosis.

Septins, which once shared a filamentous localization pattern along the cell sides during

interphase, increasingly disappear from these sides upon the onset of mitosis, starting

approximately 14 minutes after spindle pole body separation (Fig. 5F, G). At this same

time, Spn3-GFP localization to the division plane of the cell increases as the septin ring

begins to coalesce and mature (Fig. 5H). This maturation appears to occur

approximately 20 minutes after spindle pole body separation, as the spindle poles relax,

signaling the end of Anaphase B. Together, these data indicate that septins form short

linear filaments at the cortex along the long axis of the cell during interphase. These

filaments begin to disappear when the cell enters mitosis, but coalesce into a ring at the

division plane only at the end of Anaphase B.

The septin cytoskeleton and Rga4 show distinct structures at the cortex

Since the Rga4 distribution pattern along the cortex is septin-dependent, we

investigated whether the septin cytoskeleton colocalizes with Rga4. We observed Rga4-

GFP puncta along the cell sides in cells expressing Spn1-tdTomato. While both these

proteins localize to the cell sides, we did not see any significant colocalization pattern

between them (Fig.6A, Top Plane, inset). In most cells, Rga4-GFP organizes like a

corset at the cell middle as has been reported before (Das et al., 2007; Rich-Robinson

et al., 2021; Tatebe et al., 2008), while Spn1-tdTomato appears on either side of Rga4-

GFP closer to the growing ends (Supplementary Fig.S4A). In some cells, Rga4-GFP

puncta appeared on either side of Spn1-tdTomato filaments (Fig.6A, Middle Plane,

inset). To further verify this, we also analyzed cells co-expressing Rga4-GFP and Spn1-

tagRFP. We find that Spn1-tagRFP localizes on either side of the Rga4-GFP corset-like

organization in interphase cells (Fig. 6B). We asked if the dynamic Rga4-GFP is trapped

by septins via physical interaction, thereby preventing its mobility along the cortex.

If the septin cytoskeleton physically interacts with Rga4 and entraps it, this interaction

should be detectable by measuring a change in Förster Resonance Energy Transfer

(FRET) efficiency (Bajar et al., 2022; Bajar et al., 2016; McDonald et al., 2017). To test

this, we utilized an Acceptor photobleach FRET method wherein FRET efficiency was

measured between Rga4 and septin cytoskeletal proteins expressing GFP and RFP,

respectively, prior to and following complete photobleaching of the red fluorescent

acceptor (Liu et al., 2019). If interaction occurs between two proteins, bleaching of the

red-tagged acceptor fluorophore should result in a nominal increase in GFP intensity,

adjusted for normal photobleaching during the imaging process. If no interaction occurs,

no increase in GFP intensity should be detectable following acceptor photobleaching.

Indeed, when we look for interaction between two septin monomers in the cytoskeleton,

Spn3-GFP and Spn1-tagRFP, we find a distinct increase in GFP intensity and FRET

efficiency following acceptor photobleaching, indicating that there is interaction between

these proteins (Supplementary Fig. S5A, B). We used this same approach to detect

Spn1 and Rga4 interaction. We first identified regions of Rga4-GFP and Spn1-tagRFP

overlap in the cell. Next, we bleached Spn1-tagRFP and monitored the intensity of

Rga4-GFP. We fail to see an increase in FRET efficiency, and rather, Rga4-GFP

intensity decreases to a degree consistent with regular photobleaching (Supplementary

Fig. S5C, D). This absence of FRET efficiency increase between these two proteins

suggests a lack of direct interaction between them.

Further, if the septin cytoskeleton physically interacts with and entraps Rga4,

mislocalizing either the septin cytoskeleton or Rga4 should lead to mislocalization of the

other. To test this, we used the GFP trap method with a GFP-binding protein (GBP)

(Wang et al., 2016). We used the mitochondrial protein Tom20 tagged with GBP and

RFP to entrap Spn3-GFP (Fig.6C). We find that in cells expressing Tom20-GBP-RFP,

Spn3-GFP is no longer observed at the cortex, but rather appears along the

mitochondria, colocalizing with Tom20. We asked if mislocalizing one of the components

of the septin cytoskeleton could mislocalize the other components. We expressed Spn1-

TdTomato and Spn3-GFP in cells expressing Tom20-GBP. We find that Spn3-GFP with

Tom20-GBP at the mitochondria also entraps Spn1-tdTomato at that site (Fig.6D). This

indicates that mislocalizing one of the septin components to the mitochondria is

sufficient to mislocalize other proteins in the complex.

Next, we asked if Rga4-GFP can be mislocalized to the mitochondria with Tom20-GBP.

We find that in cells expressing Tom20-GBP, Rga4-GFP appears away from the cortex

and along the mitochondria (Fig.6E). However, in these cells, Spn1-tdTomato remained

at the cortex and did not colocalize with the mitochondria and Rga4-GFP. Moreover, in

Tom20-GBP expressing cells, Spn3-GFP at the mitochondria did not entrap Rga4-

mCherry (Fig.6F). In these cells, Rga4-mCherry remained at the cortex along the cell

sides. These observations, combined with our Acceptor Photobleach FRET data,

suggest that the septin cytoskeleton does not regulate Rga4 distribution via physical

interaction.

It is noteworthy that in the cells where the septin cytoskeleton localized to the

mitochondria and away from the cortex, Rga4-mCherry at the cell sides appeared less

punctate and more diffuse, similar to that observed in *spn1∆* cells. To confirm this, we

quantified the coefficient of variance of the Rga4-mCherry signal along the cell sides in

control cells with those of cells co-expressing Tom20-GBP and Spn3-GFP. We find that

the coefficient of variance of Rga4-mCherry is significantly reduced in cells where Spn3-

GFP is mislocalized to the Tom20-GBP containing mitochondria (Fig.6F). This indicates

that the absence of the septin cytoskeleton from the cell cortex is sufficient to alter Rga4

distribution pattern, even in cells where the septin proteins are expressed. These

observations further support the hypothesis that the septin cytoskeleton at the cortex in

interphase cells prevents Rga4 mobility along the cell sides.

Septins promote the proper organization of membrane lipids

Our data show that the septin cytoskeleton does not physically interact with Rga4 to

regulate its cortical dynamics and localization. Recent reports indicate that the septin

cytoskeleton regulates membrane lipid organization in the budding yeast model system

(El Alaoui et al., 2025; Pacheco et al., 2023). Similarly, septins in *Drosophila* modulate

membrane $PI(4,5)P_2$ resynthesis via regulation of the PIP5 kinase (Kumari et al., 2022).

The septin cytoskeleton contains membrane-interacting domains, including an

amphipathic helix that senses membrane curvature and specific interactions with

membrane $PI(4,5)P_2$ lipid (Beber et al., 2019; Bertin et al., 2010; Cannon et al., 2019;

Lobato-Marquez et al., 2021; Woods et al., 2021; Zhang et al., 1999). Rga4 also

contains membrane-interacting coiled-coil domains necessary for its membrane

localization (Das et al., 2007). Thus, we asked if the septin cytoskeleton regulates Rga4

localization via membrane lipid organization. The ability of septin filaments to organize

and exclude anionic and neutral lipids, respectively, has been shown to occur in

budding yeast (El Alaoui et al., 2025). To study whether septins organize specific lipids

along the *S. pombe* plasma membrane, we analyzed the localization of the

fluorescently-tagged $PI(4,5)P_2$ probe GFP-2x-PH$_{Plc}$ in *spn1$^+$* and *spn1Δ* cells (Willet et

al., 2023). In *spn1$^+$* cells, the $PI(4,5)P_2$ biomarker, GFP-2x-PH$_{Plc}$, is distributed primarily

along the cell sides and non-growing cell ends, but is diminished at growing cell ends.

However, in *spn1Δ* cells, GFP-2x-PH$_{Plc}$ levels were increased all over the cortex, and

particularly enhanced at the growing cell ends (Fig. 7A, B). It is possible that our

observed increase in GFP-2x-PH$_{Plc}$ at the cell ends of *spn1Δ* cells is the result of

enhanced lipid binding capabilities due to the absence of the septin cytoskeleton, rather

than an actual increase in the PI(4,5)P$_2$ levels. If, on the other hand, the PI(4,5)P$_2$ levels

indeed increase, we would expect to see a decrease in the levels of its precursor PI(4)P.

To test this, we analyzed the localization of the PI(4)P probe GFP-P4C-SidC in *spn1*$^+$

and *spn1Δ* cells. In *spn1*$^+$ cells, GFP-P4C-SidC localizes along the cell cortex, mainly

concentrating at the growing cell ends (Snider et al., 2020). We find that, in *spn1Δ* cells,

the overall PI(4)P levels at the cortex are reduced, including those at the cell ends (Fig.

7C, D). This suggests that the septin cytoskeleton regulates the PI(4,5)P$_2$ levels and its

organization at the plasma membrane. We asked if the disruption of the PI(4,5)P$_2$

organization is due to changes in the kinase Its3 that converts PI(4)P to the PI(4,5)P$_2$

(Deng et al., 2005). We find that, in *spn1*$^+$ cells, Its3-mNG is mainly localized along the

cell sides and mostly excluded from the growing cell ends (Fig. 7E). In *spn1Δ* mutants,

Its3-mNG levels at the cell ends are enhanced, in agreement with our data with the

PI(4,5)P$_2$ probe (Fig. 7E, F). Together, these data suggest that, in the absence of the

septin cytoskeleton, the membrane lipid organization is disrupted, with excessive

PI(4,5)P$_2$ levels.

Decreased PI(4,5)P$_2$ levels at the cell membrane enhance Rga4-GFP membrane

binding.

Our data show that the loss of the septin cytoskeleton via *spn1Δ* results in increased

PI(4,5)P$_2$ levels at the cortex (Fig. 7A, B), while simultaneously leading to decreased

distinct Rga4-GFP puncta formation and increased Rga4-GFP cortical spread, mobility,

and cytoplasmic levels (Fig. 1D, E, G-J).

We asked if a subsequent decrease in PI(4,5)P$_2$ at the cell cortex would result in an

opposite effect with regard to Rga4-GFP localization. To test this, we analyzed Rga4

localization in an *its3-1* temperature-sensitive mutant. When grown at the restrictive

temperature of 36°C, *its3-1* expressing cells fail to convert PI(4)P to PI(4,5)P$_2$ (Deng et

al., 2005). At this high temperature, even in control cells, Rga4-GFP shows aberrant

localization [data not shown], making characterization of Rga4 localization under these

conditions difficult. We therefore examined Rga4 localization to the plasma membrane

in the *its3-1* mutant at the semi-permissible temperature of 32°C over a 2-hour

incubation time.

When observing Rga4-GFP localization in *its3-1* mutants grown under semi-restrictive

conditions (32°C) for 2 hours, we find that Rga4-GFP more readily forms distinct puncta

at the cell membrane (Fig. 8A, B). Furthermore, we see that with this increase in

membrane puncta formation in the *its3-1* mutants, we subsequently observe an

increase in the membrane-to-cytoplasmic ratio of Rga4-GFP in these cells, with a shift

in the ratio favoring plasma membrane binding (Fig. 8 A-Middle Plane, C & D). These

findings indicate that increased PI(4,5)P$_2$ at the cell membrane resulting from disruption

of the septin cytoskeleton in turn leads to increased Rga4-GFP mobility along the

membrane and increased cytoplasmic exchange of Rga4. However, when the amount

of PI(4,5)P$_2$ at the cortex is reduced, Rga4-GFP more readily binds and forms distinct

puncta on the membrane, with a subsequent decrease in the amount of Rga4 localized

to the cytoplasm. Taken together, these results indicate that the septin cytoskeleton

regulates PI(4,5)P$_2$ organization at the plasma membrane, which in turn regulates Rga4

localization and polarity.

Discussion

Our previous report showed that Cdc42 activation at the cell ends is cell-cycle-

dependent and is turned off during cell division (Rich-Robinson et al., 2021). Cdc42

reactivation at the cell ends depends on the activation of the MOR/Orb6 kinase pathway

and absence of Rga4 from the cell ends (Gupta et al., 2014; Rich-Robinson et al.,

2021). Here we show that cell-cycle-dependent Rga4 localization is dependent on the

septin cytoskeleton. The septin cytoskeleton at the cell sides prevents lateral diffusion of

Rga4 along the membrane and keeps it away from the cell ends, where Cdc42 is

activated for polarized growth. Once the cell enters mitosis, the septin cytoskeleton

gradually drops from the cell sides to eventually form a medial ring, and this

corresponds with Rga4 diffused localization to the cell ends, where Cdc42 is gradually

inactivated (Fig. 8E).

After division in fission yeast, cells grow in a monopolar manner from their previously

established old end and transition to bipolar growth when the new end takes off (NETO)

(Mitchison and Nurse, 1985). Bipolar growth occurs when the old and new ends can

compete with each other for Cdc42 activation (Das et al., 2012). The zone of growth at

the cell ends is determined by the Cdc42 activation dynamics and the GAP Rga4 (Das

et al., 2012; Das et al., 2007). Rga4 localizes to the cell sides like a corset and prevents

the overflow of Cdc42 activation from the ends to these regions. This helps to maintain

the cell width and proper polarity. During cell division, Rga4 localizes to the ingressing

septum and prevents excessive Cdc42 activation at that site (Campbell et al., 2022).

While Rga4 is mostly absent at the growing cell ends, in pre-NETO cells, Rga4 localizes

to the non-growing ends. Recent reports show that Rga4 does not localize to the

growing cell ends due to plasma membrane flows as a result of growth (Gerganova et

al., 2021; Rutkowski et al., 2024). We find that in cells lacking the septin cytoskeleton,

Rga4 levels at both cell ends increase due to increased mobility along the membrane

and increased cytoplasmic exchange. However, the levels at the non-growing ends are

higher than those at the growing ends, likely due to the effect of growth-dependent

membrane flows. In these mutants, the cell ends receive increased levels of Rga4 due

to diffusion, while the growth-dependent membrane flow removes Rga4.

Cdc42 activation dynamics at the cell ends depend on positive and time-delayed

negative feedbacks (Das et al., 2012; Harrell et al., 2024; Howell et al., 2012). These

feedbacks ensure that the two cell ends can sequentially activate Cdc42 to promote

bipolar growth. Previous reports from our lab have shown that the GEF Gef1 is required

for the establishment of Cdc42 activation at the new end and thus promotes bipolar

growth (Hercyk and Das, 2019; Hercyk et al., 2019). Any disruption in the Cdc42

activation pattern results in its accumulation at one end at the expense of the other (Das

et al., 2012; Harrell et al., 2024). We also observe this in the septin mutants, where

Cdc42 dynamics are disrupted, resulting in monopolar growth. In *spn1*$^+$ cells, Rga4 can

accumulate at the non-growing ends to some degree due to the lack of displacement

from growth-dependent membrane flow. However, these levels are below a threshold

and thus do not prevent subsequent Cdc42 activation at that site, allowing for bipolar

growth. In septin mutants, however, the overall increased Rga4 accumulation at the cell

ends, particularly at the non-growing end of monopolar cells, likely reaches a threshold.

Once this threshold has been surpassed, relatively transient and low levels of Cdc42

activity at this end are not sufficient to effectively compete with the growing end for

proper oscillatory dynamics and bipolar growth.

Rga4 contains coiled-coil domains that are required for its membrane localization (Das

et al., 2007). In interphase cells, Rga4 mostly appears to form immobile puncta along

the cell sides. During mitosis, these puncta disappear, and Rga4 displays faster

dynamics with increased levels in the cytoplasm and diffuse localization along the

cortex, including the cell ends. Our data indicate that the septin cytoskeleton maintains

Rga4 puncta in interphase and restricts its localization to the cell sides. However, this

regulation is not mediated via protein-protein interaction. Due to the geometry of septin

filament organization during interphase along the long axis of the cell, it is also unlikely

that the septin cytoskeleton restricts Rga4 diffusion along the plasma membrane via

physical obstruction. Our data show that, in the fission yeast system, the septin

cytoskeleton plays a role in affecting plasma membrane lipid organization during

interphase. The presence of septins appears to promote the proper localization and

amounts of PI(4)P and PI(4,5)P_2 along the plasma membrane of interphase cells. This

localization coincides with, and is likely downstream of, the localization of the PI(4)P

kinase Its3, which is also reliant on the septin cytoskeleton for its proper localization

during interphase. In keeping with this observation, we find that in cells with diminished

PI(4,5)P_2 levels, Rga4 membrane puncta and levels are enhanced. Thus, we posit that

the septin cytoskeleton limits membrane PI(4,5)P_2 levels to promote Rga4 localization

to immobile puncta along the cell sides. In the absence of the septin cytoskeleton,

membrane PI(4,5)P_2 levels are enhanced, and Rga4 dynamics and exchange with the

cytoplasm increase (Fig. 8F).

The septin cytoskeleton in fission yeast has been investigated mostly in the context of

cytokinesis (An et al., 2004; Berlin et al., 2003; Martin-Cuadrado et al., 2005; Munoz et

al., 2014; Singh et al., 2024; Wang et al., 2015). In budding yeast, the septins organize

as a collar around the bud neck and form a ring during cytokinesis (Bridges and

Gladfelter, 2015; Woods and Gladfelter, 2021). The septin collar in budding yeast has

been shown to restrict GAP localization near the bud neck to promote bud emergence

for daughter cell formation (Okada et al., 2013). This indicates that restricting GAP

localization by septin filaments is a conserved phenomenon. Here, we show that the

septin cytoskeleton forms short linear filaments along the long axis of the cell during

interphase. While the core septins Spn1 and Spn4 are critical for the medial ring

formation during cytokinesis, all 4 septin proteins Spn1-4 are required for their role in

cell polarity. This indicates that the nature of the filaments formed in interphase is

distinct from that during mitosis. In fission yeast, the septin ring forms as anaphase B

ends, when the actomyosin ring starts to constrict. However, the septin filaments at the

cell sides begin to disappear at the onset of mitosis, well before the septin ring forms.

The septin proteins have membrane-interacting domains and are involved in lipid

metabolism (Mela and Momany, 2022; Spiliotis and McMurray, 2020; Woods and

Gladfelter, 2021). A mitotic signal may disable the septin-membrane interaction,

resulting in its loss from the cell sides. In budding yeast, the LKB1-like kinase Elm1 has

been shown to promote septin hourglass assembly and stability during bud emergence

(Marquardt et al., 2020). In mitosis, Elm1 regulates the Nim1/PAR-1-related kinase Gin4

kinase to enable septin double ring formation (Marquardt et al., 2024). However, the

Gin4 homolog in fission yeast, Cdr2, does not regulate septin ring formation (Morrell et

al., 2004). Further investigation will determine what leads to the loss of septins from the

cell sides in mitosis and its subsequent ring formation after anaphase B.

Our findings here show a role for the septin cytoskeleton in regulating polarized cell

growth in a cell-cycle-dependent manner. While most cell-cycle-dependent regulation is

mediated via kinase-substrate phosphorylation (Kaldis, 1999; Malumbres, 2014;

McCusker and Kellogg, 2012), here we show a novel regulatory system where Rga4

localization and dynamics are indirectly controlled by regulating the distribution of

specific lipids along the plasma membrane via septin localization during interphase.

While the septin rings have been mostly identified in fungal cells, it is possible that

septins in other eukaryotes also undergo cell-cycle-dependent changes in their

organization (Delic et al., 2024; Marquardt et al., 2021). In several eukaryotes, septins

have been shown to act as diffusion barriers and maintain asymmetry in the cell

(Caudron and Barral, 2009; Spiliotis and McMurray, 2020). Cell-cycle-dependent

changes in septin organization may alter their ability to act as barriers to the movements

of both proteins and membrane lipids.

Methods and Materials

Strains and cell culture

The *S. pombe* strains used in this study are listed in Table 1. All strains are isogenic to

the original strain PN567. Cells were cultured in yeast extract with supplements (YES –

Sunrise Science Products) medium and grown exponentially at 25°C, unless specified

otherwise. Standard techniques were used for genetic manipulation and analysis

(Moreno et al., 1991). Cells were grown exponentially for at least three rounds of eight

generations each before imaging.

Microscopy

Imaging was performed at room temperature (23–25°C). We used a spinning disk

confocal microscope system with a Nikon Eclipse inverted microscope with a 100×/1.49

NA lens, a CSU-22 spinning disk system (Yokogawa Electric Corporation), and a

Photometrics EM-CCD camera (Photometrics Technology Evolve with excelon Serial

No: A13B107000). Images were acquired with MetaMorph (Molecular Devices,

Sunnyvale, CA).

A Nikon Ti2 Eclipse wide-field microscope was also used with a 100x/1.49 NA objective,

an ORCA-FusionBT digital camera (Hamamatsu Model: C15440-20UP Serial No:

500428, Hamamatsu, Hamamatsu City, Japan). Images were acquired using Nikon NIS

Elements (Nikon, Melville, NY). Fluorophores were excited using an AURA Light Engine

system (Lumencor, Beaverton, OR).

Additionally, microscopy was performed with a 3i spinning disk confocal using a Zeiss

AxioObserver microscope with integrated Yokogawa spinning disk (Yokogawa CSU-X1

A1 spinning disk scanner) and a 100x/1.49 NA objective. Images were acquired with a

Teledyne Photometrics Prime 95b back-illuminated sCMOS camera (Serial No:

A20D203014, Tucson, AZ). Images were acquired using SlideBook (3i Intelligent

Imaging innovations, Denver, CO).

Fluorescence Intensity Acquisition & Quantification

Image analysis was performed on a Windows computer using the public domain NIH

ImageJ program (developed at the U.S. National Institutes of Health and available on

the Internet at http://rsb.info.nih.gov/nih-image/). Images were acquired at either single,

mid-section focal planes, or as Z-stacks with 0.2-0.4 µm focal plane intervals. Z-stack

images were superimposed using SUM-projection for analysis. Mean intensity

measurements were used for analysis, along with integrated density values where

specified. Prior to intensity acquisition, a region of interest devoid of cells was used for

background subtraction.

Analysis of protein localization heterogeneity along the cell sides

Heterogeneity of protein localization along the cell sides was analyzed by measuring the

coefficient of variance of protein distribution along said sides. Line profiles of each cell

side were drawn using ImageJ, and the standard deviations and averages of

fluorescence intensity along these line profiles were measured. The coefficient of

variance was obtained by dividing the measured standard deviation by the obtained

average, and then multiplying the result by 100%.

Fluorescence Recovery After Photobleaching

Fluorescence recovery after photobleaching (FRAP) was performed as described

before (Onwubiko et al., 2019). Fluorescently tagged proteins along the cell cortex were

first focused at the medial z-plane, and an ROI was selected for bleaching. The ROI

was imaged 10 times pre-bleaching and then bleached for 10 s, and images were then

acquired every 3 s for 50–100 images. A non-bleached spot on the cortex for the same

cell was analyzed to account for loss of signal due to photobleaching over time. A cell-

free region of the image was used for background subtraction. Intensity values for each

ROI were corrected for loss of signal over time and background signal. The intensity

values were then normalized against the mean pre-bleach value to determine the

recovery fraction over time. At least ten cells were analyzed for each genotype for each

experiment, and carried out in triplicate. The average corrected and normalized intensity

values were then plotted as a fraction of recovery over time. The fraction of recovery

over time was plotted and smoothed using GraphPad *Prism*. The half-life ($t_{1/2}$) was then

computed as the time for recovery of 50% of the signal.

Acceptor-photobleach Förster Resonance Energy Transfer (FRET)

FRET imaging was performed using a 3i spinning disk confocal using a Zeiss

AxioObserver microscope with integrated Yokogawa spinning disk (Yokogawa CSU-X1

A1 spinning disk scanner) and a 100x/1.49 NA objective. Images were acquired with a

Teledyne Photometrics Prime 95b back-illuminated sCMOS camera (Serial No:

A20D203014, Tucson, AZ). Images were acquired using SlideBook (3i Intelligent

Imaging innovations, Denver, CO). An acceptor photobleaching method was employed

for FRET imaging in live cells. The Spn1-tagRFP fluorophore was bleached throughout

the cells using 40% 561 nm laser power for 3 cycles. Middle plane images in the donor

GFP channel were acquired pre- and post-bleach. FRET percentages were calculated

by first correcting for background and photobleaching between acquisitions pre- and

post-bleach, and subsequently calculating the percentage change in GFP donor

fluorescence at areas of donor and acceptor fluorophore colocalization along the cell

cortex using ROIs of a consistent size.

Supplemental information titles and legends

Document S1. Figures S1–S5 and Table S1

Data Availability

The data generated are available in the published article and its online supplemental material.

Acknowledgements

We thank Kathy Gould, Fulvia Verde, Pilar Perez, Chuanhai Fu, Jian-Qiu Wu, and Damian Brunner for providing strains and Bret Judson at the Boston College Imaging Core for imaging support. This work is supported by the National Institutes of Health grant R01GM136847 to M.D.

Author contributions

Conceptualization: M.D. and J.M.; Methodology: M.D. and J.M.; Validation: M.D. and J.M.; Formal analysis: M.D. and J.M.; Investigation: M.D., J.M., G.C.; Resources: M.D.; Data curation: J.M. and G.C.; Writing- original draft: M.D. and J.M.; Writing - review & editing: M.D.; Supervision: M.D.; Project administration: M.D.; Funding acquisition: M.D.

Declaration of interests

The authors declare no competing or financial interests.

References

An, H., J.L. Morrell, J.L. Jennings, A.J. Link, and K.L. Gould. 2004. Requirements of fission yeast septins for complex formation, localization, and function. *Molecular biology of the cell*. 15:5551-5564.

Bajar, B.T., X. Guan, A. Lam, M.Z. Lin, R. Yasuda, T. Laviv, and J. Chu. 2022. FRET Imaging of Rho GTPase Activity with Red Fluorescent Protein-Based FRET Pairs. *Methods Mol Biol*. 2438:31-43.

Bajar, B.T., E.S. Wang, S. Zhang, M.Z. Lin, and J. Chu. 2016. A Guide to Fluorescent Protein FRET Pairs. *Sensors (Basel)*. 16.

Beber, A., M. Alqabandi, C. Prevost, F. Viars, D. Levy, P. Bassereau, A. Bertin, and S. Mangenot. 2019. Septin-based readout of PI(4,5)P2 incorporation into membranes of giant unilamellar vesicles. *Cytoskeleton*. 76:92-103.

Berlin, A., A. Paoletti, and F. Chang. 2003. Mid2p stabilizes septin rings during cytokinesis in fission yeast. *The Journal of cell biology*. 160:1083-1092.

Bertin, A., M.A. McMurray, L. Thai, G. Garcia, 3rd, V. Votin, P. Grob, T. Allyn, J. Thorner, and E. Nogales. 2010. Phosphatidylinositol-4,5-bisphosphate promotes budding yeast septin filament assembly and organization. *J Mol Biol*. 404:711-731.

Bohnert, K.A., and K.L. Gould. 2012. Cytokinesis-based constraints on polarized cell growth in fission yeast. *PLoS Genet*. 8:e1003004.

Bos, J.L., H. Rehmann, and A. Wittinghofer. 2007. GEFs and GAPs: critical elements in the control of small G proteins. *Cell*. 129:865-877.

Bridges, A.A., and A.S. Gladfelter. 2015. Septin Form and Function at the Cell Cortex. *The Journal of biological chemistry*. 290:17173-17180.

Campbell, B.F., B.S. Hercyk, A.R. Williams, E. San Miguel, H.G. Young, and M.E. Das. 2022. Cdc42 GTPase activating proteins Rga4 and Rga6 coordinate septum

synthesis and membrane trafficking at the division plane during cytokinesis.

Traffic. 23:478-495.

Cannon, K.S., B.L. Woods, J.M. Crutchley, and A.S. Gladfelter. 2019. An amphipathic

helix enables septins to sense micrometer-scale membrane curvature. *The*

Journal of cell biology. 218:1128-1137.

Caudron, F., and Y. Barral. 2009. Septins and the lateral compartmentalization of

eukaryotic membranes. *Developmental cell*. 16:493-506.

Chang, E.C., M. Barr, Y. Wang, V. Jung, H.P. Xu, and M.H. Wigler. 1994. Cooperative

interaction of S. pombe proteins required for mating and morphogenesis. *Cell*.

79:131-141.

Coll, P.M., Y. Trillo, A. Ametzazurra, and P. Perez. 2003. Gef1p, a new guanine

nucleotide exchange factor for Cdc42p, regulates polarity in

Schizosaccharomyces pombe. *Molecular biology of the cell*. 14:313-323.

Das, M., T. Drake, D.J. Wiley, P. Buchwald, D. Vavylonis, and F. Verde. 2012. Oscillatory

Dynamics of Cdc42 GTPase in the Control of Polarized Growth. *Science*.

337:239-243.

Das, M., and F. Verde. 2013. Role of Cdc42 dynamics in the control of fission yeast cell

polarization. *Biochemical Society transactions*. 41:1745-1749.

Das, M., D.J. Wiley, S. Medina, H.A. Vincent, M. Larrea, A. Oriolo, and F. Verde. 2007.

Regulation of cell diameter, For3p localization, and cell symmetry by fission yeast

Rho-GAP Rga4p. *Molecular biology of the cell*. 18:2090-2101.

Delic, S., B. Shuman, S. Lee, S. Bahmanyar, M. Momany, and M. Onishi. 2024. The

evolutionary origins and ancestral features of septins. *Front Cell Dev Biol*.

12:1406966.

Deng, L., R. Sugiura, K. Ohta, K. Tada, M. Suzuki, M. Hirata, S. Nakamura, H. Shuntoh,

and T. Kuno. 2005. Phosphatidylinositol-4-phosphate 5-kinase regulates fission

yeast cell integrity through a phospholipase C-mediated protein kinase C-
independent pathway. *The Journal of biological chemistry*. 280:27561-27568.

El Alaoui, F., I. Al-Akiki, S. Ibanes, S. Lyonnais, D. Sanchez-Fuentes, R. Desgarceaux,
C. Cazevieille, M.P. Blanchard, A. Parmeggiani, A. Carretero-Genevrier, S. Piatti,
and L. Picas. 2025. Septin assemblies promote the lipid organization of
membranes. *Structure*. 33:451-464 e455.

Estravis, M., S. Rincon, and P. Perez. 2012. Cdc42 regulation of polarized traffic in
fission yeast. *Communicative & integrative biology*. 5:370-373.

Etienne-Manneville, S. 2004. Cdc42--the centre of polarity. *Journal of cell science*.
117:1291-1300.

Galliot, B., and L. Ghila. 2010. Cell plasticity in homeostasis and regeneration. *Mol
Reprod Dev*. 77:837-855.

Gallo Castro, D., and S.G. Martin. 2018. Differential GAP requirement for Cdc42-GTP
polarization during proliferation and sexual reproduction. *The Journal of cell
biology*. 217:4215-4229.

Gerganova, V., I. Lamas, D.M. Rutkowski, A. Vjestica, D.G. Castro, V. Vincenzetti, D.
Vavylonis, and S.G. Martin. 2021. Cell patterning by secretion-induced plasma
membrane flows. *Sci Adv*. 7:eabg6718.

Gregory, J.R., I.M.A. Llaneza, A.H. Osmani, H.E. Gosselin, S.A. Sabzian, and J.Q. Wu.
2025. Localization and function of septins are susceptible to epitope tagging.
Molecular biology of the cell. 36:ar148.

Guillot, C., and T. Lecuit. 2013. Mechanics of epithelial tissue homeostasis and
morphogenesis. *Science*. 340:1185-1189.

Gupta, S., M. Govindaraghavan, and D. McCollum. 2014. Cross talk between NDR
kinase pathways coordinates cytokinesis with cell separation in
Schizosaccharomyces pombe. *Eukaryotic cell*. 13:1104-1112.

Harrell, M.A., Z. Liu, B.F. Campbell, O. Chinsen, T. Hong, and M. Das. 2024. Arp2/3-
dependent endocytosis ensures Cdc42 oscillations by removing Pak1-mediated
negative feedback. *The Journal of cell biology*. 223.

Hercyk, B.S., and M.E. Das. 2019. F-BAR Cdc15 Promotes Cdc42 Activation During
Cytokinesis and Cell Polarization in Schizosaccharomyces pombe. *Genetics*.
213:1341-1356.

Hercyk, B.S., J. Rich-Robinson, A.S. Mitoubsi, M.A. Harrell, and M.E. Das. 2019. A
novel interplay between GEFs orchestrates Cdc42 activity during cell polarity and
cytokinesis in fission yeast. *Journal of cell science*. 132.

Howell, A.S., M. Jin, C.F. Wu, T.R. Zyla, T.C. Elston, and D.J. Lew. 2012. Negative
feedback enhances robustness in the yeast polarity establishment circuit. *Cell*.
149:322-333.

Johnson, D.I. 1999. Cdc42: An essential Rho-type GTPase controlling eukaryotic cell
polarity. *Microbiology and molecular biology reviews : MMBR*. 63:54-105.

Kaldis, P. 1999. The cdk-activating kinase (CAK): from yeast to mammals. *Cellular and
molecular life sciences : CMLS*. 55:284-296.

Kinoshita, M. 2003. The septins. *Genome Biol*. 4:236.

Kumari, A., A. Ghosh, S. Kolay, and P. Raghu. 2022. Septins tune lipid kinase activity
and PI(4,5)P(2) turnover during G-protein-coupled PLC signalling in vivo. *Life Sci
Alliance*. 5.

Liu, M., M.B. Heimlicher, M. Bachler, C.C. Ibeneche-Nnewihe, E.L. Florin, D. Brunner,
and A. Hoenger. 2019. Glucose starvation triggers filamentous septin assemblies
in an S. pombe septin-2 deletion mutant. *Biol Open*. 8.

Lloyd, A.C. 2013. The regulation of cell size. *Cell*. 154:1194-1205.

Lobato-Marquez, D., J. Xu, G.O. Guler, A. Ojiakor, M. Pilhofer, and S. Mostowy. 2021.
757   Mechanistic insight into bacterial entrapment by septin cage reconstitution. *Nat*
758   *Commun*. 12:4511.

759   Maddox, A.S., L. Lewellyn, A. Desai, and K. Oegema. 2007. Anillin and the septins
760   promote asymmetric ingression of the cytokinetic furrow. *Developmental cell*.
761   12:827-835.

762   Malumbres, M. 2014. Cyclin-dependent kinases. *Genome Biol*. 15:122.

763   Marquardt, J., X. Chen, and E. Bi. 2021. Septin Assembly and Remodeling at the Cell
764   Division Site During the Cell Cycle. *Front Cell Dev Biol*. 9:793920.

765   Marquardt, J., X. Chen, and E. Bi. 2024. Reciprocal regulation by Elm1 and Gin4
766   controls septin hourglass assembly and remodeling. *The Journal of cell biology*.
767   223.

768   Marquardt, J., L.L. Yao, H. Okada, T. Svitkina, and E. Bi. 2020. The LKB1-like Kinase
769   Elm1 Controls Septin Hourglass Assembly and Stability by Regulating Filament
770   Pairing. *Current biology : CB*. 30:2386-2394 e2384.

771   Martin-Cuadrado, A.B., J.L. Morrell, M. Konomi, H. An, C. Petit, M. Osumi, M.
772   Balasubramanian, K.L. Gould, F. Del Rey, and C.R. de Aldana. 2005. Role of
773   septins and the exocyst complex in the function of hydrolytic enzymes
774   responsible for fission yeast cell separation. *Molecular biology of the cell*.
775   16:4867-4881.

776   Mata, J., and P. Nurse. 1997. tea1 and the microtubular cytoskeleton are important for
777   generating global spatial order within the fission yeast cell. *Cell*. 89:939-949.

778   McCusker, D., and D.R. Kellogg. 2012. Plasma membrane growth during the cell cycle:
779   unsolved mysteries and recent progress. *Current opinion in cell biology*. 24:845-
780   851.

McDonald, N.A., A.L. Lind, S.E. Smith, R. Li, and K.L. Gould. 2017. Nanoscale
architecture of the Schizosaccharomyces pombe contractile ring. *Elife*. 6.

Mela, A., and M. Momany. 2022. Septins coordinate cell wall integrity and lipid
metabolism in a sphingolipid-dependent process. *Journal of cell science*. 135.

Merla, A., and D.I. Johnson. 2000. The Cdc42p GTPase is targeted to the site of cell
division in the fission yeast Schizosaccharomyces pombe. *European journal of*
cell biology. 79:469-477.

Mitchison, J.M., and P. Nurse. 1985. Growth in cell length in the fission yeast
Schizosaccharomyces pombe. *Journal of cell science*. 75:357-376.

Moreno, S., A. Klar, and P. Nurse. 1991. Molecular genetic analysis of fission yeast
Schizosaccharomyces pombe. *Methods in enzymology*. 194:795-823.

Morrell, J.L., C.B. Nichols, and K.L. Gould. 2004. The GIN4 family kinase, Cdr2p, acts
independently of septins in fission yeast. *Journal of cell science*. 117:5293-5302.

Munoz, S., E. Manjon, and Y. Sanchez. 2014. The putative exchange factor Gef3p
interacts with Rho3p GTPase and the septin ring during cytokinesis in fission
yeast. *The Journal of biological chemistry*. 289:21995-22007.

Murray, J.M., and D.I. Johnson. 2001. The Cdc42p GTPase and its regulators Nrf1p and
Scd1p are involved in endocytic trafficking in the fission yeast
Schizosaccharomyces pombe. *The Journal of biological chemistry*. 276:3004-
3009.

Okada, S., M. Leda, J. Hanna, N.S. Savage, E. Bi, and A.B. Goryachev. 2013. Daughter
cell identity emerges from the interplay of Cdc42, septins, and exocytosis.
Developmental cell. 26:148-161.

Onishi, M., T. Koga, A. Hirata, T. Nakamura, H. Asakawa, C. Shimoda, J. Bahler, J.Q.
Wu, K. Takegawa, H. Tachikawa, J.R. Pringle, and Y. Fukui. 2010. Role of septins

in the orientation of forespore membrane extension during sporulation in fission
yeast. *Molecular and cellular biology*. 30:2057-2074.

Onwubiko, U.N., P.J. Mlynarczyk, B. Wei, J. Habiyaremye, A. Clack, S.M. Abel, and
M.E. Das. 2019. A Cdc42 GEF, Gef1, through endocytosis organizes F-BAR
Cdc15 along the actomyosin ring and promotes concentric furrowing. *Journal of*
cell science. 132.

Pacheco, J., A.C. Cassidy, J.P. Zewe, R.C. Wills, and G.R.V. Hammond. 2023.
PI(4,5)P2 diffuses freely in the plasma membrane even within high-density
effector protein complexes. *J Cell Biol*. 222.

Perez, P., E. Portales, and B. Santos. 2015. Rho4 interaction with exocyst and septins
regulates cell separation in fission yeast. *Microbiology (Reading)*. 161:948-959.

Ray, S., K. Kume, S. Gupta, W. Ge, M. Balasubramanian, D. Hirata, and D. McCollum.
2010. The mitosis-to-interphase transition is coordinated by cross talk between
the SIN and MOR pathways in Schizosaccharomyces pombe. *The Journal of cell*
biology. 190:793-805.

Revilla-Guarinos, M.T., R. Martin-Garcia, M.A. Villar-Tajadura, M. Estravis, P.M. Coll,
and P. Perez. 2016. Rga6 is a Fission Yeast Rho GAP Involved in Cdc42
Regulation of Polarized Growth. *Molecular biology of the cell*.

Rich-Robinson, J., A. Russell, E. Mancini, and M. Das. 2021. Cdc42 reactivation at
growth sites is regulated by local cell-cycle-dependent loss of its GTPase-
activating protein Rga4 in fission yeast. *Journal of cell science*. 134.

Rutkowski, D.M., V. Vincenzetti, D. Vavylonis, and S.G. Martin. 2024. Cdc42 mobility
and membrane flows regulate fission yeast cell shape and survival. *Nat*
Commun. 15:8363.

Singh, D., Y. Liu, Y.H. Zhu, S. Zhang, S. Naegele, and J.Q. Wu. 2024. Septins function
in exocytosis via physical interactions with the exocyst complex in fission yeast
cytokinesis. *bioRxiv*.

Snider, C.E., A.H. Willet, H.T. Brown, J.S. Chen, J.M. Evers, and K.L. Gould. 2020.
Fission yeast Opy1 is an endogenous PI(4,5)P(2) sensor that binds to the
phosphatidylinositol 4-phosphate 5-kinase Its3. *Journal of cell science*. 133.

Sparks, C.A., M. Morphew, and D. McCollum. 1999. Sid2p, a spindle pole body kinase
that regulates the onset of cytokinesis. *The Journal of cell biology*. 146:777-790.

Spiliotis, E.T., and M.A. McMurray. 2020. Masters of asymmetry - lessons and
perspectives from 50 years of septins. *Molecular biology of the cell*. 31:2289-
2297.

Sprang, S. 2001. GEFs: master regulators of G-protein activation. *Trends in biochemical*
sciences. 26:266-267.

Tasto, J.J., J.L. Morrell, and K.L. Gould. 2003. An anillin homologue, Mid2p, acts during
fission yeast cytokinesis to organize the septin ring and promote cell separation.
J Cell Biol. 160:1093-1103.

Tatebe, H., K. Nakano, R. Maximo, and K. Shiozaki. 2008. Pom1 DYRK regulates
localization of the Rga4 GAP to ensure bipolar activation of Cdc42 in fission
yeast. *Current biology : CB*. 18:322-330.

Verde, F., J. Mata, and P. Nurse. 1995. Fission yeast cell morphogenesis: identification
of new genes and analysis of their role during the cell cycle. *The Journal of cell*
biology. 131:1529-1538.

Wang, N., I.J. Lee, G. Rask, and J.Q. Wu. 2016. Roles of the TRAPP-II Complex and
the Exocyst in Membrane Deposition during Fission Yeast Cytokinesis. *PLoS*
biology. 14:e1002437.

Wang, N., M. Wang, Y.H. Zhu, T.W. Grosel, D. Sun, D.S. Kudryashov, and J.Q. Wu.
2015. The Rho-GEF Gef3 interacts with the septin complex and activates the
GTPase Rho4 during fission yeast cytokinesis. *Molecular biology of the cell*.
26:238-255.

Wei, B., B.S. Hercyk, J. Habiyaremye, and M. Das. 2017. Spatiotemporal Analysis of
Cytokinetic Events in Fission Yeast. *J Vis Exp*:e55109.

Wei, B., B.S. Hercyk, N. Mattson, A. Mohammadi, J. Rich, E. DeBruyne, M.M. Clark,
and M. Das. 2016. Unique Spatiotemporal Activation Pattern of Cdc42 by Gef1
and Scd1 Promotes Different Events during Cytokinesis. *Molecular biology of the*
cell. 27:1235-1245.

Willet, A.H., L.A. Turner, J.S. Park, L. Ren, C.E. Snider, and K.L. Gould. 2023.
Characterization of Pik1 function in fission yeast reveals its conserved role in lipid
synthesis and not cytokinesis. *Journal of cell science*. 136.

Woods, B.L., K.S. Cannon, E.J.D. Vogt, J.M. Crutchley, and A.S. Gladfelter. 2021.
Interplay of septin amphipathic helices in sensing membrane-curvature and
filament bundling. *Molecular biology of the cell*. 32:br5.

Woods, B.L., and A.S. Gladfelter. 2021. The state of the septin cytoskeleton from
assembly to function. *Current opinion in cell biology*. 68:105-112.

Wu, J.Q., Y. Ye, N. Wang, T.D. Pollard, and J.R. Pringle. 2010. Cooperation between the
septins and the actomyosin ring and role of a cell-integrity pathway during cell
division in fission yeast. *Genetics*. 186:897-915.

Zhang, J., C. Kong, H. Xie, P.S. McPherson, S. Grinstein, and W.S. Trimble. 1999.
Phosphatidylinositol polyphosphate binding to the mammalian septin H5 is
modulated by GTP. *Current biology : CB*. 9:1458-1467.

Zheng, S., F. Dong, F. Rasul, X. Yao, Q.W. Jin, F. Zheng, and C. Fu. 2018. Septins
regulate the equatorial dynamics of the separation initiation network kinase Sid2p
and glucan synthases to ensure proper cytokinesis. *FEBS J*. 285:2468-2480.

Zheng, S., B. Zheng, and C. Fu. 2024. The Roles of Septins in Regulating Fission Yeast
Cytokinesis. *J Fungi (Basel)*. 10.

Zheng, S., B. Zheng, Z. Liu, X. Ma, X. Liu, X. Yao, W. Wei, and C. Fu. 2022. The Cdc42

GTPase-activating protein Rga6 promotes the cortical localization of septin.

Journal of cell science. 135.

 Figure Legends

Figure 1. The septin cytoskeleton restricts Rga4 localization and dynamics at the

cell sides during interphase. A. Middle plane images of Rga4-GFP in *spn1$^+$* and

spn1Δ cells. Red brackets highlight Rga4 distribution along cell sides in interphase

cells. Red asterisks highlight dividing cells. **B-C**. Quantification of the coefficient of

variation of Rga4 distribution through the cell-cycle in *spn1$^+$* (B) and *spn1Δ* (C) cells.

[N=3, *n*=40-50 cells each, *$P{\leq}0.05$, *ns*=not significant,]. **D.** Rga4-GFP increasingly

localizes to the cell ends and cytoplasm in *spn1Δ* cells during interphase. Images

consist of superimposed middle-plane images taken at 10-second intervals over a 5-

minute time-lapse. Red boxes highlight increased localization of Rga4 to the cell ends.

E. Quantification of Rga4-GFP intensity within the cytoplasm of interphase cells in *spn1$^+$*

and *spn1Δ* conditions measured by averaging three unique regions within the cytoplasm

of each cell [**$P{\leq}0.01$, Student's T test]. **F.** Quantification of Rga4-GFP localization to

growing and non-growing cell ends of interphase cells in *spn1$^+$* and *spn1Δ* conditions.

[N=3, *n*=24-28 cells each, ****$P{\leq}0.0001$, ***$P{\leq}0.001$, **$P{\leq}0.01$, *$P{\leq}0.05$, *ns*=not

significant, one-way ANOVA, scale bar=5μm]. **G.** Montage of Rga4-GFP in *spn1$^+$* cells

during interphase, during mitosis, and in *spn1Δ* cells during interphase, taken over a 5-

minute time-lapse. Red arrowhead marks an Rga4-GFP puncta at the start of the time-

lapse. **H.** Montage of Rga4-GFP recovery after photobleaching during interphase and

mitosis in *spn1$^+$* and *spn1Δ* cells taken in 15-second intervals. Red circles indicate the

photobleached region of interest (ROI). Red lightning bolt indicates the time point of

photobleaching. **I.** FRAP recovery rates of Rga4-GFP signal intensity within the ROI

over time in *spn1$^+$* and *spn1Δ* cells during interphase and mitosis. **J.** $T_{1/2}$ fluorescence

recovery times of listed conditions. [N=3, *n*=10 cells each, **$P \le 0.01$, *$P \le 0.05$, *ns*=not

significant, one-way ANOVA, scale bar=5µm].

Figure 2. The septin cytoskeleton promotes bipolar growth. A. Representative

images of calcofluor white-stained *spn1$^+$* and *spn1Δ* cells. Blue asterisks highlight

bipolar (*spn1$^+$*) growth and red asterisks highlight monopolar (*spn1Δ*) growth. **B.**

Quantification of monopolarity index in *spn1$^+$* and *spn1Δ* cells. [N=3, *n*=65-100,

*$P \le 0.05$, **$P \le 0.01$, student's t-test, scale bars=5µm]. **C.** Quantification of the septation

index of *spn1$^+$* and *spn1Δ* cells. [N=3, *n*>200, *$P \le 0.01$, student's t-test]. **D.**

Representative images of calcofluor white-stained *spnX$^+$* and *spn2Δ, spn3Δ,* and *spn4Δ*

cells, respectively. Red asterisks highlight monopolar cells. Red arrowheads highlight

septated cells that failed to separate. **E.** Quantification of monopolarity index in *spnX$^+$*

and *spn2Δ, spn3Δ,* and *spn4Δ* cells, respectively. **F.** Quantification of the septation

index of *spnX$^+$* and *spn2Δ, spn3Δ,* and *spn4Δ* cells, respectively. [N=3, *n*=>110 cells

each, ****$P \le 0.0001$, ns=not significant, one-way ANOVA, multiple comparison test,

scale bars=5µm].

Figure 3. The septin cytoskeleton promotes bipolar Cdc42 activation. A.

Representative SUM Z-Projection images of *rga4$^+$ spn1$^+$*, *rga4$^+$ spn1Δ, rga4Δ spn1$^+$*,

and *rga4Δ spn1Δ* cells expressing CRIB-3xGFP, respectively. Red arrowheads highlight

Cdc42 activity at the ends of growing cells. **B.** Quantification of the total levels of CRIB-

3xGFP intensity via measuring integrated density at growing cell ends of the

aforementioned cells in A [N=3, *n*>50, ns=not significant, one-way ANOVA, multiple

comparison test, scale bar=5µm]. **C.** Heat map of the fraction of CRIB-GFP intensity at

each cell end over the total intensity with increasing cell size in *spn1$^+$* and *spn1Δ* cells.

D, E. Representative images of the oscillatory dynamics of Cdc42 activity via the active

Cdc42 biomarker, CRIB-3xGFP, over time in *spn1⁺* and *spn1Δ* cells. Images taken at 1-

minute intervals over a 60-minute time-lapse. **F, G.** Quantification of CRIB-3xGFP

intensity at the old and new cell ends of a representative *spn1⁺* and *spn1Δ* cell over 60-

minute time-lapse**. H.** A representative time-lapse of CRIB-3xGFP and Rlc1tdTomato in

spn1⁺ and *spn1Δ* cells. Red asterisk marks the onset of stable CRIB-3xGFP localization

at the cell ends after the completion of ring constriction marked with a purple asterisk. **I.**

Frequency plot of the timing of the onset of stable Cdc42 activity at the cell ends as

indicated by CRIB-3xGFP after the completion of ring constriction in *spn1⁺* and *spn1Δ*

cells (*n*>30). [scale bars=5µm].

Figure 4. Septins form linear filaments along the cell sides in interphase. A.

Localization patterns of Spn1-mEGFP, Spn1-tdTomato, Spn2-GFP, Spn3-GFP, and

Spn4-tdTomato. All septin proteins localize to the cell sides as linear filaments (red

arrowhead) and form a ring at the division site during mitosis. **B.** Spn2-GFP, Spn3-GFP,

and Spn4-tdTomato fail to localize to the cell sides in *spn1Δ* mutants. **C.** Spn3-GFP and

Spn1-tagRFP colocalize to the short linear filaments along the cell sides in interphase

and within a ring during division. **D.** Spn3-GFP and Spn1-tdTomato colocalize in linear

filaments along the cell cortex during interphase, and within a ring during division. Red

and white arrowheads highlight filaments. Scale bar=5µm.

Figure 5. Septins recede from the cell sides during mitosis to eventually form a

medial ring as Anaphase B ends. A. Representative time-lapse of Spn3-GFP, CRIB-

mCherry, and Sad1-mCherry expressing cells starting at the onset of mitosis. Purple

arrow highlights the time point when Cdc42 activity is totally lost from the cell ends.

Green arrow highlights the maturation of the septin ring. Black arrow highlights the end

of Anaphase B. **B.** Frequency plot of the timing of loss of active Cdc42 (CRIB-mCherry)

from the cell ends, septin ring formation, and end of Anaphase B with respect to the

onset of mitosis. (n=10). **C.** Montage of time-lapse of Spn4-tdTomato and Myo2-GFP

Sid4-GFP expressing cells during mitosis. Red asterisk marks the onset of mitosis, the

green asterisk marks the onset of septin ring formation, and the purple asterisk marks

the end of mitosis. Scale bar=2µm. **D.** Quantification over time of Spn4-tdTomato levels

along the cell side of the cell, shown in **C**. **E.** Quantification of Spn4-tdTomato along the

cell sides with increasing spindle pole body distances. [N=3, n>20, *$P \le 0.05$, one-way

ANOVA, multiple comparison test]. **F.** Montage of time-lapse of Spn3-GFP and Rlc1-

tdTomato Sad1-mCherry expressing cells during mitosis. Red asterisk marks the onset

of mitosis. Red boxes highlight decreasing Spn3-GFP filament intensity along the cell

side throughout mitosis. **G.** Quantification of Spn3-GFP intensity along the cell end and

sides compared to the cell middle throughout mitosis. [N=3, *n*=10-19, ****$P \le 0.0001$,

student's t-test]. Scale bar=5µm.

Figure 6. The septin cytoskeleton does not colocalize or interact with Rga4. A.

Rga4-GFP and Spn1-tdTomato show distinct localization patterns along the sides of

interphase cells. Red arrowheads highlight zones with Rga4-GFP but no Spn1-

tdTomato along the cell side. The middle plane inset shows partial colocalization of

Rga4-GFP with Spn1-tdTomato. The top plane inset shows distinct localization patterns

of these proteins. **B.** Rga4-GFP and Spn1-TagRFP show distinct localization patterns in

interphase cells. Rga4-GFP localizes as a corset at the cell middle (red bracket) while

Spn1-tagRFP appears closer to the cell ends (green bracket). The image is a SUM-

projection of a z-stack. **C.** Spn3-GFP mislocalizes to Tom20-GBP-RFP at the

mitochondria. **D.** Spn3-GFP and Spn1-tdTomato simultaneously mislocalize to the

mitochondria in Tom20-GBP expressing cells. **E.** Rga4-GFP mislocalizes to the

mitochondria, but not Spn1-tdTomato in Tom20-GBP expressing cells. **F.** Spn3-GFP

mislocalizes to the mitochondria, but not Rga4-mCherry, in Tom20-GBP expressing

cells. **G.** Quantification of the coefficient of variation of Rga4-mCherry heterogeneity

along cell sides in interphase cells, Spn3-GFP and Tom20-GBP expressing cells, as

shown in E. [N=3, n=20-26 cells each, *P≤0.05, student's t-test, scale bar=5µm].

Figure 7. The septin cytoskeleton organizes lipids and lipid-associated enzymes

along the cell membrane during interphase. A. Representative middle plane images

of the PI(4,5)P_2 fluorescent biomarker GFP-2x-PH$_{Plc}$ in $spn1^+$ and $spn1\Delta$ cells. Blue and

red arrowheads highlight decreased and increased GFP-2x-PH$_{Plc}$ intensity at the border

between the cell side and cell end, respectively. **B.** Quantification of GFP-2x-PH$_{Plc}$

intensity along the cell cortex, measured from the very end of the cell to the middle.

[N=3, n= 12-22 cells each, ****P≤0.0001, student's t-test]. **C.** Representative middle

plane images of the PI(4)P fluorescent biomarker GFP-P4C-SidC in $spn1^+$ and $spn1\Delta$

cells. Blue and red arrowheads highlight increased and decreased GFP-P4C-SidC

intensity at the cell ends, respectively. **D.** Quantification of GFP-P4C-SidC intensity

along the cell cortex, measured from the very end of the cell to the middle. [N=3, n=12-

20 cells each, ****P≤0.0001, student's t-test]. **E.** Representative middle plane images of

the PI(4)P kinase Its3-mNG in $spn1^+$ and $spn1\Delta$ cells. Blue and red arrowheads

highlight decreased and increased Its3-mNG intensity at the cell ends, respectively. **F.**

Quantification of Its3-mNG intensity along the cell cortex, measured from the very end

of the cell to the middle. [N=3, n=20-23 cells each, ****P≤0.0001, student's t-test, scale

bar=5µm].

Figure 8. Disrupting PI(4,5)P$_2$ levels at the cell cortex results in increased Rga4-

GFP membrane binding and puncta formation. A. Representative images of top and

middle plane images of Rga4-GFP puncta in *its3*[+] and *its3-1* cells grown at the semi-

restrictive temperature of 32°C for 2 hours. Red arrows highlight Rga4-GFP puncta at

the cell cortex. Red asterisks highlight differences in cytoplasmic Rga4-GFP intensity,

particularly towards the cell ends. **B.** Quantification of the average number per cell of

Rga4-GFP puncta at the cell cortex present in middle plane images of *its3*[+] and *its3-1*

cells. [N=3, n>20, **P≤0.01, student's t-test]. **C.** Representative images of Rga4-GFP in

both *its3*[+] or *its3-1* cells growing at the semi-restrictive temperature of 32°C for 2 hours.

Images consist of superimposed middle-plane images taken at 10 second intervals over

a 5-minute time-lapse. **D.** Quantification of the membrane-to-cytoplasmic ratio of Rga4-

GFP in *its3*[+] or *its3-1* cells grown for 2 hours at the semi-restrictive temperature of 32°C.

[N=3, n>25, *P≤0.05, student's t-test, scale bar=5µm]. **E.** Schematic model of Rga4 and

septins in *S. pombe* throughout the cell cycle. Green dots indicate Rga4-GFP, and

yellow dots indicate septins. Increased green highlighting of the cytoplasm denotes the

increasing Rga4-GFP signal in the cytoplasm of *spn1*[+] cells throughout the cell cycle

(Created with BioRender). **F.** Schematic model of the effects of the septin cytoskeleton

on membrane lipid organization and Rga4 membrane localization. **Top.** Septin filaments

help organize lipids along the plasma membrane, in turn facilitating immobile Rga4

puncta on the membrane. **Bottom.** The loss of the septin cytoskeleton results in

disorganized membrane lipids, in turn resulting in a decreased ability of Rga4 to bind

the membrane, instead promoting increased diffusion along the membrane, and

increased exchange with the cytoplasm (Created with BioRender).

Figure 1

Figure 2.

Figure 3.

A CRIB-3xGFP

spn1⁺ *spn1Δ*

rga4Δ *rga4Δ spn1Δ*

B CRIB-GFP at growing cell ends

ns

Intensity (a.u.)

spn1⁺ rga4⁺ *spn1Δrga4⁺* *spn1⁺rga4Δ* *spn1Δrga4Δ*

C *spn1⁺*

spn1Δ

Cell Length (μm)

Density

D *spn1⁺*

CRIB-3xGFP

Time (min) 0 4 8 12 16 20 24 28

E *spn1Δ*

0 4 8 12 16 20 24 28

F *spn1⁺* CRIB-3xGFP cell end intensities

Mean Intensity (A.U.)

Time (min)

Old End
New End

G *spn1Δ* CRIB-3xGFP cell end intensities

Mean Intensity (A.U.)

Time (min)

Old End
New End

H

CRIB-3xGFP

spn1⁺

Rlc1-tdT

CRIB-3xGFP

spn1Δ

Rlc1-tdT

-10 -8 -6 -4 -2 0 2 4 6 8 10 12 14 16 18 20 22 24

Time (min)

I Cdc42 reactivation at cell ends

Frequency disribution (%)

Time (min)

spn1⁺
spn1Δ

Figure 4.

Figure 5

Figure 6.

Figure 7.

A GFP-2x-PH$_{Plc}$ – PI(4,5)P$_2$ Marker

spn1$^+$ *spn1Δ*

B GFP-2xPH$_{Plc}$ Intensity Along the Cell Cortex

spn1$^+$
spn1Δ

Intensity (A.U.)
Length from Cell End (µm)

C GFP-P4C-SidC – PI(4)P

spn1$^+$ *spn1Δ*

D GFP-4C-SidC Intensity Along the Cell Cortex

spn1$^+$
spn1Δ

Intensity (A.U.)
Distance from Cell Tip (µm)

E Its3-mNG – PI(4)P Kinase

spn1$^+$ *spn1Δ*

F Its3-mNG Intensity Along the Cell Cortex

Intensity (A.U.)
Distance from Cell Tip (µm)

spn1$^+$
spn1Δ

Figure 8.

A Rga4-GFP 32°C 2-hour

Top Plane | Middle Plane | Top Plane | Middle Plane

its3⁺ | *its3-1*

B Average Number of Rga4-GFP Puncta

C Rga4-GFP 32°C 2-hour

its3⁺ | *its3-1*

D Rga4-GFP Membrane to Cytoplasmic Ratio

E Interphase → Early Mitosis → Late Mitosis

Rga4
Septin

F *spnX⁺*

Plasma membrane
Cytoplasm
Septin barrier
Rga4

spnXΔ

Disorganized plasma membrane
Cytoplasm
Membrane binding & unbinding
Rga4