# A Vulnerable Subtype of Dopaminergic Neurons Drives Early Motor Deficits in Parkinson's Disease

Akira Fushiki[1,2,9]*, David Ng[2], Zachary R. Lewis[1], Archana Yadav[3], Tatiana Saraiva[4,5], Luke A. Hammond[2,6], Christoph Wirblich[7], Bosiljka Tasic[1], Vilas Menon[3], Joaquim Alves da Silva[4,8], and Rui M. Costa[1,2,9]*

Allen Institute, Seattle, WA 98109, USA

Zuckerman Mind Brain Behavior Institute, Columbia University, New York, NY 10027, USA

Center for Translational and Computational Neuroimmunology, Department of Neurology Columbia University Irving Medical Center, New York, NY 10032, USA

Champalimaud Research, Champalimaud Foundation, Lisbon 1400-038, Portugal

Department of Neurology, University Hospital of Würzburg, Würzburg 97080, Germany

Department of Neurology, The Ohio State University, Columbus, OH 43210, USA

Department of Microbiology and Immunology, Sidney Kimmel Medical College, Thomas Jefferson University, Philadelphia, PA 19107, USA

NOVA Medical School, Universidade Nova de Lisboa, Lisbon 1169-056, Portugal

Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD, 20815

*Correspondence: akira.fushiki@alleninstitute.org (AF) and rui.costa@alleninstitute.org (RMC)

## Abstract

In Parkinson's disease, dopaminergic neurons (DANs) in the midbrain gradually degenerate, with ventral substantia nigra pars compacta (SNc) DANs exhibiting greater vulnerability. However, it remains unclear whether specific molecular subtypes of ventral SNc DANs are more susceptible to degeneration in PD, and if they contribute to the early motor symptoms associated with the disease. We identified a subtype of *Sox6+* DANs, *Anxa1+*, which are selectively lost earlier than other DANs, and with a time course that aligns with the development of motor symptoms in MitoPark mice. We generated a knock-in Cre mouse line for *Anxa1+* DANs and showed differential anatomical inputs and outputs of this population. Further, we found that the inhibition of transmitter release in *Anxa1+* neurons led to bradykinesia and tremor. This study uncovers the existence of a selectively vulnerable subtype of DANs that is sufficient to drive early motor symptoms in Parkinson's disease.

## Keywords

Parkinson's disease, dopamine, cell types, neurodegeneration, vulnerability, tremor, motor symptoms

**Introduction**

Parkinson's disease (PD) is a progressive neurological disorder that primarily affects movement. It develops gradually, sometimes starting with a barely noticeable tremor in one hand[1]. While tremors, rigidity and akinesia are well-known signs of PD, the most distinctive symptom is bradykinesia, characterized by the slowing of movements. The main pathological hallmark of PD is the loss of dopaminergic neurons (DANs) in the substantia nigra[2,3] and the progressive reduction of DANs worsens the motor symptoms over time.

DANs in the midbrain are heterogeneous and they differ in their cellular properties and functions[4,5]. This heterogeneity is evident across several dimensions, including their molecular profiles and connectivity[4,6]. Transcriptomic studies have revealed that populations of midbrain DANs can be differentiated by distinct gene expression profiles[4], a finding that extends beyond the conventional anatomical division of midbrain DANs into the substantia nigra pars compacta (SNc, A9) and the ventral tegmental area (VTA, A10). Recent findings have discovered that molecularly unique subtypes of DANs innervate different subregions of the striatum, suggesting a basis for the variety of behavioral responses linked to these distinct neural pathways[5,6]. In PD, the degeneration of DANs follows a stereotypical pattern where some populations of neurons are more vulnerable than others[7]. Within the SNc, there is a distinction between the ventral-tier and dorsal-tier DANs, where the ventral-tier neurons tend to degenerate first[7]. Other studies have focused on changes in gene expression associated with PD compared to healthy controls[8]. Transcriptomic analysis revealed significant upregulation alpha-synuclein[9] and downregulation of parkin[10], which are critical to dopamine synthesis and degradation pathways, respectively. This distinct expression pattern suggests a disruption in the proteostasis network that could be pivotal in disease progression. Additionally, genes related to mitochondrial function showed decreased expression levels, consistent with the known impact of mitochondrial dysfunction in Parkinson's pathology[11]. This differential gene expression or susceptibility aligns with the recent discovery of the distinct axonal projection patterns identified in dopaminergic neuron subtypes. These patterns determine specific neural circuits affected during disease progression, providing a mechanistic basis for the diverse behavioral changes observed. For instance, subtypes projecting to motor-related areas may influence movement deficits, while those targeting limbic regions might underlie emotional or cognitive disturbances, illustrating how circuit-specific vulnerability translates into varied clinical manifestations[4–6]. However, it is not completely understood if specific molecularly identifiable subtypes of DANs in the ventral SNc are especially vulnerable in PD, and if the loss of function of these early

vulnerable populations is just a biomarker of the disease or contributes to the early motor

symptoms associated with PD.

Here, we characterized the progression of behavioral symptoms and dopaminergic

degeneration in MitoPark mice, a well-established PD mouse model[12], which replicates key

aspects of the disease through the targeted deletion of the *Tfam* (mitochondrial transcription

factor A) gene in dopaminergic neurons using the Cre-LoxP system. We characterized their

motor behaviors in an open field with a motion sensor and identified bradykinesia and a tremor

gradually emerging as MitoPark mice showed progressive dopaminergic degeneration in the

SNc. We then performed single-nucleus RNA-sequencing (snRNA-seq) in mutant and matched

littermate controls at different ages to identify transcriptomic changes in vulnerable

dopaminergic populations, and correlated the susceptibility of different cell types to

degeneration with the emergence of behavioral phenotypes. We identified specific dopaminergic

cell types, highly ventrally localized *Sox6+* neurons, whose loss was timeline-wise correlated

with the onset of motor symptoms. Further, we found that a subpopulation of the *Sox6+* neurons

– the *Anxa1+* subset is more vulnerable than other *Sox6+* populations. We next created an

Anxa1-Cre knock-in mouse to study the circuit architecture and function of this dopaminergic

neuron subtype. We characterized the anatomical inputs and outputs and observed a specific

pattern of strong projections of *Anxa1+* neurons to the dorsal striatum, not seen in other DAN

populations, and observed that this subtype also receives different inputs than others. Finally,

we showed that specific functional impairment of *Anxa1+* neurons is sufficient to induce

slowness of movements and motor tremors, corresponding to the early PD-like symptoms.

These results suggest that a specific vulnerable subtype of DANs is responsible for a range of

early motor symptoms of PD, potentially highlighting the role of different dopaminergic neurons

subtypes in the progression of the disease.

**Results**

**Progressive development of PD-like motor deficits in MitoPark mice**

The MitoPark mouse (genotype: *Tfam-loxP/loxP*, *+/DAT-cre*) is a transgenic model of

Parkinson's disease (PD) where the mitochondrial transcription factor A ( *Tfam*) is removed

selectively in dopaminergic neurons (DANs)[12]. Loss of *Tfam* in DANs leads to a progressive

degeneration of these neurons and the manifestation of motor symptoms that are PD-like[12,13].

We characterized spontaneous movement of MitoPark mice between 8 and 24 weeks of

age in a square open field without any external cues or reward. These mice showed gradual

development of locomotor deficits post-natal with decreased average speed (bradykinesia),

increased immobility time and increased number of stops (akinesia) (**Figures 1A-1D**). Given

that typical PD has other motor symptoms, such as tremor in addition to bradykinesia and

akinesia, we used a wireless motion sensor to detect subtle motor changes that are not easily

observable by video-characterization. The sensor measures tri-axial acceleration with high

resolution and allows for the continuous monitoring of movement with high frequency (200Hz),

which could enable us to detect tremors that are usually poorly documented in genetically

engineered rodent models of PD[14]. Given that tremor is defined as an involuntary rhythmic and

oscillatory movement of a body part[15], we explored spectral analysis of accelerometer data to

quantify the potential emergence of abnormal oscillations with the progressive degeneration of

DANs. Interestingly, we observed a specific oscillation within the 12-18 Hz frequency band that

emerged in MitoPark mice at 16 weeks which gradually intensified with age (**Figures 1E-1H**

**and S1**). Notably, this oscillation was present even when the mice were immobile (**Figures 1F**

**and 1H**).

We next examined whether depletion of SNc DANs would be sufficient to lead to the

development of PD-like symptoms similar to those observed in MitoPark mice. We therefore

genetically ablated the neurons in the SNc of DAT-Cre mice by bilateral stereotactic injection of

an adeno-associated virus (AAV) to express a Cre-dependent apoptotic gene Caspase-3 (AAV-

Ef1a-FLEX-taCasp3)[16] in the SNc of DAT-Cre mice. Selective losses of the SNc DANs and their

projections were observed after 4 weeks of viral expression (**Figure 1I**). We evaluated mice

behavior in the open field, and found that the depletion of SNc DANs reproduced the PD-like

phenotype observed in the 24 weeks of MitoPark mice (**Figures 1A-1D, 1J and 1L**). We also

performed a cylinder task to observe rearing behavior, which is more effortful and therefore

more likely to be affected in PD, where there is a tendency towards less vigorous movements[17].

We indeed confirmed a significant reduction in rearing in the animals with ablation of SNc DANs

(**Figures 1K and 1M**). Furthermore, we also replicated the emergence of the tremor as

observed in MitoPark mice (**Figures 1N and 1O**). These results suggest that the death of

neurons in the SNc is sufficient to produce the bradykinesia, akinesia, and tremor observed in

the MitoPark mice.

**Progressive loss of dopamine neurons in the ventral midbrain in MitoPark mice**

To determine whether the mitochondrial *Tfam* deficit leads to changes in the number of midbrain

DANs, we surveyed Th expression in MitoPark mice and monitored changes in *Th*+ cells in the

SNc of MitoPark mice at 8, 16 and 24 weeks (**Figure 2A and 2B**). We also examined Th+

axonal projections from the SNc of MitoPark mice to striatum and performed brain

reconstruction. Given that the current Allen Common Coordinate Framework (Allen CCF) does

not have anatomical subdivisions of the striatum[18], we created an updated map based on

cortical projections[19] and applied the imaging analysis pipeline BrainJ as previously described[20]

(**Figure S2**). This updated atlas allowed us to quantitatively analyze the density of projections

neurons in subregions of the striatum and visualize the prominent loss of *Th+* axonal projections

in MitoPark mice (**Figure 2C**). We observed that the projections from the SNc DANs to the

striatum were intact at 8 weeks in MitoPark mice. However, by 16 weeks, their projections were

significantly reduced compared to age-matched controls, especially for those targeting the

dorsal striatum. By 24 weeks, the projections were largely depleted. (**Figure 2A and 2C**).

We also employed single-nucleus RNA sequencing (snRNA-seq) of the ventral midbrain

of MitoPark mice, their littermate controls (genotype: *Tfam-loxP/loxP*, +/+ or *Tfam-loxP/+*, +/+)

and C57BL/6 (B6) mice at 8, 16, and 24 weeks of age (8 mice in each group; 4 males and 4

females). We used a previously described experimental approach to isolate nuclei and profile

their transcriptomes from mice[21] (**Figure 2D**, see "Methods" section). We obtained a total of

293,435 nuclei, of which 93,703 were from MitoPark mice, 100,266 were from their littermate

controls and 99,466 were from B6 mice. We mapped cell annotations to our data with the Allen

Brain Cell (ABC) Atlas using their MapMyCells tool[22] (**Figures 2E and S3A**). Dopaminergic

nuclei were identified as the subclass annotation of 215_SNc_VTA_RAmb_Foxa1_Dopa at the

ABC Atlas, within the class annotation of 21_MB_Dopa (**Figure S3A**). The dopaminergic

population was also clearly distinguished by expression of the canonical marker gene, *Th*

(**Figure 2F**). Using *Rbfox3* to mark neuronal cells, we identified other neuronal cell types such

as excitatory (*Slc17a6* positive) and inhibitory (*Slc32a1* positive) cell types (**Figure 2F and**

**S3B**). Oligodendrocytes were characterized by the expression of *Mog*, while Oligodendrocyte

precursor cells (OPCs) expressed high levels of *Pdgfra*. Microglia were characterized by *Ptprc*

expression, while astrocytes and endothelial cells were marked by specific expression of *Gja1*

and *Nostrin*, respectively (**Figure S3B**). Consistent with our histological findings, we observed a

significant decrease in the number of dopaminergic cells in MitoPark mice between 8 weeks and

24 weeks (**Figure S4A**). To address whether cell types are enriched or depleted in MitoPark

mice, we compared the proportions of all cell-types across the ventral midbrain at each stage.

We found that the dopaminergic populations significantly decreased with age compared with

other cell types (**Figure S4B**). We also performed mixed-effects modeling (MASC)[23] that

accounts for technical and biological confounders to identify cellular populations associated with

the disease models (**Figures 2G-2I and S4C-S4E**). MitoPak mice had a significant reduction of

DANs at 16 and 24 weeks relative to their littermate controls, while the other cell-types were

stable (**Figures 2G, 2H, S4C and S4D**). A similar reduction of DANs was also observed

between 8 and 16 weeks (**Figure 2I and S4E**). Together, these data demonstrate that *Tfam*

depletion in MitoPark mice induces robust and selective dopaminergic degeneration without

altering the number of other neighboring cell types (**Figures 2G-2I and S4B-S4E**).

**Vulnerability of *Sox6+* population during the progression of PD-like phenotypes**

We next aimed to identify changes within specific subtypes of DANs during the progression of

neurodegeneration. We isolated 18,611 DANs from nuclei of the ventral midbrain, and snRNA-

seq data was processed to identify distinct groups representing unique neuronal subtypes

(**Figures 3A, S5A and S5B**). We identified 8 DAN subtypes and confirmed the expression of

well-known subtype markers *Sox6, Slc17a6, and Calb1* (**Figure 3B**)[4]. Each group was

segregated by distinct genetic markers, which included known dopaminergic neuron subtype-

specific genes (**Figures S5C, S6A-S6C**). To determine which neuronal subtypes were most

affected as neurodegeneration progressed, we analyzed the cell composition within each

identified group from MitoPark mice and their age-matched controls. Comparative analysis

showed significant changes in the composition of dopaminergic neuronal subtypes (**Figure 3C**).

Notably, the proportion of *Sox6+* neurons was markedly reduced in MitoPark mice, whereas the

proportion of other neuron cell types such as *Megf11+* or *Synpr+* remained 'stable' throughout

the degeneration process. As the overall number of dopamine neurons decreases in MitoPark

mice, the proportion of these 'stable' neurons increased, suggesting they are more resilient

compared to other dopamine subtypes that are more susceptible to degeneration (**Figure 3C**).

These neurons are mostly *Vglut2+* (Slc17a6) (**Figures 3A and 3B**), which has previously been

reported to play a potential neuroprotective role in the survival of DANs in PD model

animals[24,25]. Additionally, there are dopamine neuron subtypes that undergo uniform

degeneration over time (*Plekhg1+*, *Cck+*, *Sema5b+*, *Etv1+*, *Lypd1+*) (**Figures 3C**). Using the

MASC approach, we further analyzed the differential proportion of different cell-types or states

across different groups (**Figures 3D-3F**) while adjusting for age, sex, and batch as covariates.

The *Sox6+* population showed a significant decrease in the odds ratio in MitoPark mice

compared to their littermate controls, especially at 16 or 24 weeks (**Figures 3D and 3E**), and it

was not due to age-related effects alone (**Figures S6D-S6G**). We also conducted the test

across different ages within the same genotype to assess developmental changes in cell

population dynamics (**Figure 3F**). The odds ratio of *Sox6+* population still declined, suggesting

that this subtype exhibits greater vulnerability than other DANs in PD, consistent with findings

from human PD transcriptomic studies[8].

**A subset of *Sox6*+ DANs are more vulnerable in MitoPark mice**

*Sox6* is involved in the development and differentiation of DANs, particularly in the ventral
midbrain DANs[26]. The *Sox6*+ supertype (ID: 0882 in 215_SNc_VTA_RAmb_Foxa1_Dopa) can
be further subdivided according to their anatomical location, distinct molecular signatures, and
functional properties[4]. To explore the diversity of *Sox6*+ neurons, we conducted transcriptomic
analysis of the subtypes and performed the cluster annotation based on the ABC atlas[22]. We
found 43 clusters (ID: 3837-3879) under the supertype (ID: 0880-0887) annotation, of which, the
*Sox6*+ supertype can be divided into 7 subsets (ID: 3853-3859) (**Figure S5A**). We measured
the proportions of the *Sox6*+ dopaminergic subtypes in MitoPark and control animals and found
that *Anxa1*+ (ID: 3857) and *Arpp21*+ (ID: 3859) subsets significantly decreased compared to
controls, with *Anxa1*+ showing a decrease compared to controls at both 16 and 24 weeks,
whereas *Arpp21*+ just showed a significant decrease at 24 weeks (**Figure 4A**). This suggests
that the *Anxa1*+ subtype of *Sox6*+ DANs is especially vulnerable early in the disease. We also
performed an additional differential abundance analysis using Milo, a method that offers a
structured approach to exploring variations in cell compositions across diverse experimental
conditions[27] (**Figures 4B-4J and S7A-S7I**). Milo analysis confirmed a significant depletion of
*Sox6*+ population in MitoPark mice at 16 and 24 weeks compared to their littermate controls,
consistent with previous analyses (**Figures S7A-I**). We then applied the cluster annotation and
found that *Glra2+, Anxa1+, Hs6st3+, Grin2c+ and Csf2rb2*+ subsets exhibited substantial
 fold changes in gene expression at 16 weeks (**Figures 4B, 4E, and 4H**). We also performed
the analysis between 8 and 16 weeks in MitoPark mice and found these same subpopulations
(ie. *Glra2+, Anxa1+, Hs6st3+, Grin2c+ and Csf2rb2*+) were affected (**Figures 4D, 4G, 4J, and
S7J-S7L**). To verify the transcriptional reduction of DAN subsets in the SNc of MitoPark mice,
we conducted immunostaining with an antibody targeting *Aldh1a1*, a well-established marker for
ventral DANs[4,5] (**Figure 4K**). We found a remarkable loss of *Aldh1a1*+ DANs in MitoPark mice
at 16 weeks, particularly in the rostral-ventral region (**Figure 4K**). These results demonstrate
that *Sox6*+ DANs exhibited differential susceptibility, with *Anxa1+;Sox6*+ DANs being among
the earliest to show vulnerability compared to other *Sox6*+ (ie. *Anxa1-;Sox6*+) DANs.

**Whole-brain efferent and afferent connectivity of *Anxa1*+ DANs**

The results above indicate a subset of *Sox6*+ DANs, the *Anxa1*+ neurons*,* is more vulnerable
than others. However, it remains unclear whether loss of these neurons is an early biomarker of
disease, or are actually driving early symptoms in PD. To address this, we generated an Anxa1-

Cre knock-in mouse line (see 'Method' section), with Cre recombinase under transcriptional

control of an Anxa1-specific promoter and characterized its expression using GFP to visualize

cell bodies and axon projections to output structures. To achieve this, we injected Cre-

dependent AAV-GFP into the SNc of Anxa1-Cre mice and counted the number of *Th*+ cells with

*Calb1* antibody, which labels the dorsal dopaminergic populations in the SNc (**Figures 5A and**

**5B**). In all mice, we observed robust co-localization of GFP with *Th*+ cells but not *Calb1*+ cells,

suggesting that most of the GFP+ cells in the SNc are ventral specific DANs (**Figures 5A and**

**5B**). We then examined GFP+ axonal projections from the SNc of all DANs or *Anxa1*+ DANs

and performed whole brain reconstruction using BrainJ (**Figures 5C and 5D)**. The quantitative

analysis revealed prominent axonal projections of *Anxa1*+ DANs to the dorsal striatum (**Figure**

**5D**). We also created a DAT-Flp knock-in mouse line (**Figure S9A;** see 'Method' section) and

compared the axonal projections of SNc *Anxa1*+ DANs with other DANs subtypes such as

*Vglut2*+ or *Calb1*+ (**Figure S9B**). As shown previously[6], these subsets have distinct axonal

projections to the striatum; *Vglut2*+ DANs project to the dorsolateral and tail striatum, while

*Calb1*+ DANs project to the ventral striatum (**Figure S9B**). We also quantitatively analyzed SNc

*Vglut2*+ and *Calb1*+ DAN axonal projections throughout the whole brain, which make specific

projections to many different regions. The fact that different populations of DANs have such

distinctive projections opens the possibility that different populations modulate different

behaviors or behavioral parameters (**Figure S9C)**.

To identify the afferent inputs onto *Anxa1*+ dopaminergic neuronal subtypes and how

these inputs differ from other DAN subtypes, we used CVS-N2c rabies tracing, a strain with

enhanced transsynaptic labeling[28], with DAT-Cre+, Anxa1-Cre+ (and also DAT-Flp+;Anxa1-

Cre+), DAT-Flp+;Vglut2-Cre+ and DAT-Flp+;Calb1-Cre+ mice. We first examined the inputs to

all DANs in the SNc by injecting two Cre-dependent helper viruses (TVA and N2cG) to express

the avian receptor (TVA protein) and the rabies glycoprotein G (N2cG) unilaterally into the SNc

of DAT-Cre+ mice. Two weeks later, CVS rabies virus (CVS-tdTomato) was injected into the

same site (**Figure S10A;** see "Methods" section**)**. We observed tdTomato-labeled neurons in

more brain regions than previous studies[29,30], especially detected much more inputs for cortical

regions, likely attributed to the higher transsynaptic property of the N2C strain. These labeled

neurons, primarily restricted ipsilateral to the injection site, represent monosynaptic afferent

inputs to the SNc DANs (**Figures 5F, 5G and S10C**). The same viral injection strategy was also

used using Anxa1-Cre+ (and DAT-Flp+;Anxa1-Cre+), DAT-Flp+;Vglut2-Cre+, and DAT-

Flp+;Calb1-Cre+ mice, which also labeled an abundant number of tdTomato+ neurons in

various parts of brain regions (**Figures 5E, 5H, 5I and S9D-9G**). The dopaminergic neuron

subtypes exhibit differences in their ipsilateral inputs, particularly with respect to inputs from

cortical regions. For example, while *Anxa1+* DANs receive strong inputs from the motor cortex,

this subtype does not receive significant inputs from the somatosensory, prelimbic, or orbital

cortex. In contrast, *Vglut2+* DANs receive minimal inputs from the ventral regions of the anterior

cingulate and retrosplenial cortices compared to other dopamine subtypes. In most other cases,

however, *Anxa1+* DANs receive similar inputs to the other dopaminergic subtypes, suggesting a

large shared organization of inputs into SNc dopaminergic circuits (**Figures 5E-5I and S9D-**

**S9G**).

**Inhibition of *Anxa1+* DANs is sufficient to cause early PD-like motor deficits**

To test if the loss of *Anxa1+* neurons is merely an early biomarker of disease, or could actually

be the cause of early symptoms in PD, we selectively silenced synaptic release of *Anxa1+*

DANs with an AAV expressing tetanus toxin[31] (AAV-FLEX-TeLC-EYFP; **Figure 6**). We found

that silencing of *Anxa1+* neurons caused a reduction in locomotor speed, suggesting it is

sufficient to cause bradykinesia. However, it did not affect immobility time or number of stops,

suggesting that it was not sufficient to cause akinesia. (**Figures 6B-6D**). In a cylinder task, we

observed the significant drops in rearing, suggesting that *Anxa1+* dopaminergic neuron activity

is important for effortful movements (**Figure 6C**). We then performed spectral analysis on mice

depleted of *Anxa1+* SNc dopaminergic neurons and found a slight shift in the frequency of

tremor, resembling the phenotype of MitoPark mice at 16 weeks (**Figures 6E**). Taken together,

these data show that inhibition of *Anxa1+* DANs is sufficient to drive early PD-like motor

symptoms in movement speed (bradykinesia), movement vigor, and tremor, but not sufficient to

cause akinesia.

**Discussion**

The data presented in this study revealed that a subtype of DANs is selectively vulnerable in a

mouse model of PD and that its loss is sufficient to drive PD-like motor symptoms that are

observed early in the disease. We conducted a precise longitudinal analysis of the behavioral

changes exhibited by MitoPark mice, a Parkinson's disease (PD) model. We identified

progressive changes in bradykinesia, akinesia, and in tremor. Concomitantly, these mice

demonstrated a progressive degeneration of dopaminergic neurons (DANs) in the midbrain,

specifically within the substantia nigra pars compacta (SNc). These phenotypes observed in

late-stage MitoPark mice were replicated by ablating dopaminergic neurons in the SNc.

Additionally, we performed a comprehensive characterization of the molecular profiles of

midbrain neurons as the disease progressed, and uncovered selective vulnerability in the *Sox6+*
DANs, and in particular in the *Anxa1+* subtype. Finally, we observed that silencing this
dopamine subtype was sufficient to produce bradykinesia and tremor of the magnitude
observed early in the progression of the degeneration, but not akinesia. This reveals that the
loss of these vulnerable neurons is not only a biomarker of early disease, it also drives the early
motor symptoms observed early in the disease.

Our behavioral analysis rigorously mapped the progression of motor symptoms in
MitoPark mice revealing early symptoms that were not previously reported. Furthermore, it
revealed a previous unobserved phenotype associated with dopamine loss. By using a motion
sensor in an open field behavioral assay, we have enhanced our capability to detect more
subtle motor phenotypes that are not detected using conventional video-based characterization
methods. This approach led us to identify tremors in MitoPark mice, which developed
incrementally with the progressive degeneration of DANs in the brain. Furthermore, specific
ablation and silencing of SNc DANs and the *Anxa1+* subtype revealed their important role in this
tremor phenotype. Although these oscillations were present at rest, the frequency of the tremors
we observed are not within the typical band of 4-6 Hz seen in patients with PD. However, tremor
frequency differences between species have been well documented, such in the case of the
harmaline model of essential tremor with frequencies ranging from 8–10, 10–12 and 11–14 Hz
in monkeys, rats and mice respectively[32]. Considering that there are several types of tremors
observed in humans and the extent to which these tremors resemble those seen in human
diseases has not yet been fully established, a more comprehensive characterization is
imperative

We conducted an extensive longitudinal molecular profiling of the cell types in the ventral
midbrain as the symptoms progressed in MitoPark mice, utilizing a newly developed atlas, the
Allen Brain Cell (ABC) Atlas, for annotation[22]. This detailed examination identified a vulnerable
population of DANs, *Sox6+*, the characteristics of which align closely with those identified in
previous human studies[8]. In contrast, other subtypes, such as *Megf11+* (ID: 0880), and *Synpr+*
(ID: 0887) displayed resilience to degeneration. We identified several additional subtypes of
*Sox6+* DANs, with different vulnerability. *Anxa1+* (ID: 3857) neurons showed the earliest
significant decrease compared to WT controls, but *Arpp21+* (ID: 3859) also showed vulnerability
in late stages compared to other subtypes of *Sox6+* that did not show increased vulnerability.
This variance in vulnerability and resilience indicates that the degeneration of DANs occurs at
different rates in different types, and that the molecular composition of these subtypes may
underly the mechanisms for vulnerability and resilience. Consequently, a deeper understanding

of the molecular processes driving the degeneration and resilience of these various DAN
subtypes is critical for understanding the progression of the disease, and could be harnessed for
the development of biomarkers to monitor progression and eventually the response to disease
modifying therapies.
We also characterized with unprecedented depth the afferent and efferent connectivity of
three distinct subtypes of DANs in the SNc, roughly representing lateral, dorsal, and ventral SNc
DANs, identified by the expression of genes *Vglut2+*, *Calb1+*, and *Anxa1+* respectively.
Consistent with prior findings [6], these subtypes exhibit unique projection patterns within the
striatum, occupying distinct regions. This supports the conclusion that these subtypes display
divergent behaviors in motor control and reward processing[5]. However, our results also reveal
that, despite their distinct projection patterns, these subtypes receive surprisingly similar inputs
across the brain, with some exceptions of cortical regions. This does not support the idea that
different function or vulnerability is related to massively different inputs. One possibility is that
despite being subtle, the differences in inputs matter. Another is that positionally, the different
populations of DANs receive inputs from different regions of the same area (different neurons,
with different functions), and that results in different functions. Finally, it is also possible that
neurons in SNc have similar processing functions with subtle differences in their response, but
result in different functions because of their more specific output projections. Those projections
could target not only differential brain regions, but also different cell types within those regions
(e.g. excitatory vs. inhibitory neurons). Another layer of complexity is added by the presence of
a recurrent network where SNc DANs receive robust recurrent inhibition from medium spiny
neurons from striatum[33,34] and in turn project to the striatum. This specific projection pattern of
each dopaminergic neuron subtype could modulate activity in the striatum, which in turn,
through inhibitory feedback, could fine-tune the DAN response during different behaviors or
learning processes. Additionally, a more detailed classification of the inputs and projections of
single neurons in each DAN subtype, provide the resolution necessary to discern more specific
input and output patterns, akin to observations made in fruit flies[35]. This granular approach may
reveal intricate neurobiological dynamics obscured when broader classifications are used.
Finally, although targeted manipulation of *Anxa1+* neurons led to significant bradykinesia
and tremor reminiscent of the early stages of disease in MitoPark mice, it did not affect immobile
time or akinesia. These discrepancies might be explained by a dosage effect (i.e. akinesia
requires the loss of more dopamine neurons than bradykinesia), or by the involvement of other
dopaminergic neuron subtypes in different phenotypes. The possibility of homeostatic
compensation within DAN neural networks, with different neurons stepping in to sustain the

functionality of behaviors affected by the loss of others, should also be discussed. We observed

considerable overlap in the brain regions providing afferent connectivity to each dopamine

subtype and this observation could be critical to understand the adaptability of these circuits. A

critical approach to further unravel these complexities of adaptation as disease develops would

involve recording whole-brain activity in Parkinson's model mice. Gaining a deeper

understanding of the resilience and adaptability of neural circuits surrounding dopaminergic

neurons will enhance our grasp of Parkinson's disease and potentially foster innovative

therapeutic strategies in the future.

## Materials and Methods

### Contact for reagents and resource sharing

Further information and requests for reagents should be directed to and will be fulfilled by the lead contact, Rui M. Costa (rui.costa@alleninstitute.org).

### Mouse breeding and husbandry

All experimental protocols were performed according to National Institutes of Health (NIH) guidelines and in compliance with the regulations of the Institutional Animal Care and Use Committee at Columbia University. All experimental animals were 2 to 6 month-old mice housed on a 12 hr light/dark cycle with unrestricted access to food and water. Mice used for behavioral experiments were individually housed. The strain and lines used were: C57BL6/J (Jackson Laboratories, 000664), MitoPark mouse[12], DAT-Cre mouse (Jackson Laboratories, 006660), VGlut2-Cre mouse (Jackson Laboratories, 028863), and Calb1-IRES2-Cre mouse (Jackson Laboratories, 028532). Anxa1-Cre and DAT-Flp mouse lines were developed and utilized in our lab (details are provided below). MitoPark mice ($DAT\text{-}Cre^{+/-};Tfam^{loxP/loxP}$) were obtained by crossing $DAT\text{-}Cre^{+/-}; Tfam^{loxP/+}$ mice with $Tfam^{loxP/loxP}$ mice to selectively knockout $Tfam$ in dopaminergic neurons.

### Generation of transgenic mice (Anxa1-Cre and DAT-Flp)

Anxa1-Cre and DAT-Flp knock-in mice were generated by Cyagen (Santa Clara, CA) using CRISPR/Cas9. To generate Anxa1-Cre mice, one cell mouse embryos were microinjected with Cas9 protein, a gRNA (5' GACATCCCAACTATTCTGCA-AGG 3') targeting exon 13 in the vicinity of the stop codon, and a repair template comprised of homology arms and a 'P2A-Cre-rBG pA' cassette. Homology arms were generated by PCR from BAC clones RP23-53N8 and RP23-149B23 as template. Similarly, to generate DAT-Flp mice, a gRNA (5' CCAACAGCCAATGGCGCAGC-TGG 3') targeting exon 15, and a template with a '2A-FlpO-rBG pA' cassette were used. Homology arms were PCR amplified from BAC clones RP23-150M11 and RP23-34F24. Two silent mutations, aa 613L (CTG to TTA), and aa 614R (CGC to AGG) were introduced to prevent re-cleavage by Cas9 after homology-directed repair.

### Nuclear suspension, FACS and Single-nucleus RNAseq

We used procedures as previously described[36,37]. Briefly, MitoPark mice and their littermate controls (P56~62 for 8 weeks, P112~118 for 16 weeks and P168~174 for 24 weeks; 4 female and 4 male at each time point) were anaesthetized with isoflurane and perfused with cold artificial

cerebrospinal fluid (ACSF) containing N-methyl-d-glucamine (NMDG; 92 mM), KCl (2.5 mM),
NaH2PO4 (1.2 mM), NaHCO3 (30 mM), HEPES (20 mM), D-Glucose (25 mM), Sodium L-
ascorbate (5 mM), Sodium pyruvate (3 mM), MgSO4 (10mM), CaCl2 (0.5 mM), bubbled with
carbogen gas (95% O2 and 5% CO2) at a pH of 7.2–7.4. The brain was sectioned at 400um using
a vibratome (VT1200S, Leica Microsystems) on ice, and the ventral midbrain (SNc and VTA) were
microdissected from three consecutive sections (-2.48 to -3.88mm from Bregma; Franklin and
Paxinos, 2008). The tissues were transferred to microcentrifuge tubes, flash frozen in dry ice with
ethanol, and stored at -80C until later use. For nuclei isolation[21], frozen tissues were placed into
a homogenization buffer that consisted of 10mM Tris pH 8.0, 250mM Sucrose, 25mM KCl, 5mM
MgCl2, 0.1% Triton-X 100, 1X Protease inhibitor (Promega G6521), 0.5% RNasin Plus RNase
inhibitor (Promega N2615) and 0.1mM DTT (Promega P1171). Tissues were placed into a 2ml
dounce homogenizer (Sigma D8938) and homogenized using 10 strokes of the loose dounce
pestle followed by 10 strokes of the tight pestle to liberate nuclei. Homogenate was strained
through a 30µm cell strainer (Miltenyi Biotech 130-098-458) and centrifuged at 900xg for 10
minutes to pellet nuclei. Nuclei were then resuspended in blocking buffer containing 1X PBS
supplemented with 0.8% nuclease-free BSA (Sigma, OmniPur 2905) and 0.5% RNasin Plus
RNase inhibitor. Prior to fluorescence-activated nuclei sorting (FACS), DAPI (Sigma, D9542) was
applied to nuclei suspensions at a final concentration of 0.1µg/ml and nuclei suspensions were
filtered through a 30µm nylon mesh (Sysmex, 04-004-2326) to remove aggregates. FACS was
performed at the Zuckerman Institute Flow Cytometry platform using a MoFlo Astrios EQ
(Beckman Coulter) sorter. Single nuclei were captured by gating DAPI+ events, excluding debris
and doublets. Both cell suspension and collection chamber were kept at 4C during sorting. Sorted
nuclei were spun down immediately and barcoded using 10x Genomics Chromium Single Cell 3'
Reagent Kit v3 according to the manufacturer's protocol. Samples were processed and libraries
were prepared/sequenced by the Columbia JP Sulzberger Genome Center Single Cell Analysis
Core.
444

**RNAseq data processing and analysis**
Sequencing data were aligned and quantified using the Cell Ranger Single-Cell Software Suite
(v.6.1.2) with default parameters against the ENSEMBL GRCh38 reference genome[38]. The pre-
filtered datasets were then imported into the R Seurat package for downstream analysis (v5.0.2)[39].
We first removed low-quality cells (cell containing < 200 genes and genes expressed in <3 cells
of the data). After the filtering, contaminating ambient RNA reads were filtered by using SoupX[40].
Doublets were predicted using the DoubletFinder R package[41], and were excluded for

downstream analysis. After quality control and cell filtering (1000<nFeature_RNA<7500;

nCount_RNA<40000; percent_mt<1, percent_ribo<1, percent_hemo<1), the datasets were

processed for identifying cell populations by Seurat pipeline. Briefly, pre-processed gene

expression matrices were normalized by the NormalizedData function. A total of 2,000 most

variable features were then selected with the FindVariableFeatures and scaled by ScaleData

command (while regressing out mitochondrial genes) for the Principal Component Analysis (PCA),

which was performed by the RunPCA command for 30 PCs. After integration with Harmony[42], the

data was subsequently processed with Louvain clustering and dimensionality reduction via

Uniform Manifold Approximation and Projection (UMAP) using 30 principal components, reduction

= "harmony", and resolution of 0.8, to visualize nuclear transcriptomic profiles in two-dimensional

space. For annotation, we used the Allen Brain Cell (ABC) Atlas tool MapMyCells

(RRID:SCR_024672, the reference taxonomy: 10x Whole Mouse Brain (CCN20230722),

mapping algorithm: hierarchical mapping)[22] to assign neuronal classes or subclasses to nuclei

from a whole dataset. To subset the dopaminergic population, we first used the annotation

(21_MB_Dopa) to identify and extract the relevant population. We then re-scaled the extracted

data and applied the supertype or cluster annotation to the subset and performed further analysis.

Differentially expressed genes in each cluster were identified by Wilcoxon test (the FindAllMarkers

function in Seurat),

**Differential cell abundance**

To identify differentially abundant cell populations in each dataset, we used two methods: MASC

(version 0.10)[23] and Milo (version 1.99)[27]. MASC is a single cell association method for testing

whether a certain status influences the membership of single cells while accounting for technical

confounds and biological variation. For all datasets, we included sex as a fixed effect, and

individual as a random effect in the model. We then tested the significance of association between

the status (i.e. MitoPark mice and their littermate controls at each stage, or MitoPark mice at

different stages) with the clusters. Cell subpopulations were considered significantly associated

with a status at FDR-adjusted P<0.05 and absolute odds ratio >0. Milo is a tool for the test on k-

nearest neighbor (KNN) graph neighborhoods. Briefly, we performed PCA dimensionality

reduction and KNN graph embedding on the samples. We define a neighborhood as the group of

cells that are connected to a sampled cell by an edge in the KNN graph. For each analysis, a k-

nearest neighbors (KNN) graph was constructed using the graph slot from the adjacency matrix

of previously processed Seurat objects and cells were assigned to neighborhoods (k=30, d=30).

For each neighborhood we then perform hypothesis testing between conditions to identify

differentially abundant cell states whilst controlling the FDR across the graph neighborhoods. We
tested for differences in abundance between the cells at different stages of MitoPark mice and
their littermate controls.
489
495  **Gene ontology enrichment analysis**
491  We used the g:Profiler web tool and gprofiler2 R package (version 0.2.3)[43] to identify significantly
492  enriched gene ontology (GO) terms from upregulated or downregulated gene lists in dopaminergic
493  clusters.
494
**Stereotaxic viral injections**
Before starting the surgery mice were subcutaneously injected with Buprenorphine XR (0.5–1 mg
per kg body weight). The mouse head was shaved, cleaned with 70% alcohol and iodine, an
intradermic injection of bupivacaine (2 mg per kg body weight) was administered, and a small
incision from anterior to posterior was made on the skin to allow for aligning the head and drilling
the hole for the injection site. Surgeries were performed under sterile conditions and isoflurane
(1%–5%, plus oxygen at 1-1.5 l/min) anesthesia on a motorized stereotactic frame (David Kopf
Instruments, Model 900SD). Throughout each surgery, mouse body temperature was maintained
at 37C using an animal temperature controller (ATC2000, World Precision Instruments). For
anterograde tracing experiments, animal were unilaterally injected with 300nL of AAV5-CAG-
FLEX-GFP (titer: 4.5E12 vg/mL; UNC Vector Core) into the right hemisphere of the substantia
nigra pars compacta (SNc; AP -3.16 mm, ML 1.3 mm, DV -4.0 mm) using a Nanoject III Injector
(Drummond Scientific, USA) at a pulse rate of 5.1 nL per second, 20 nL per pulse every 5 s. After
injection, the pipette was held in place for 5 minutes before raising to ensure minimize viral efflux.
For whole-brain retrograde tracing experiments using rabies virus, 100nL of a 2:1 mixture of the
helper viruses, AAV1-CAG-FLEX-N2cG-mKate2.0 (titer: 9.83E12 vg/mL; Janelia Viral Core) or
AAVDJ-CAG-FlpX-N2cG-mKate2.0 (titer: 2.60E12 vg/mL; Zuckerman Institute Vector Core) and
AAV5-Esyn-DIO-TVA950-EYFP-WPRE (titer: 1.44E13 vg/mL; Salk Viral Vector Core), was
injected into the SNc. Two weeks later, 300nl of EnvA-N2c-deltaG-tdTomato (titer: 1.5E9 ffu/ml;
Center for Neuroanatomy with Neurotropic Viruses, CNNV) was injected within the same area at
a 10-degree angle (SNc; AP -3.16 mm, ML 2.0mm, DV -4.0 mm) to prevent labeling the injection
tract with TVA. For monosynaptic labeling from the primary motor cortex or the dorsal striatum,
100nL of a 2:1 mixture of the helper viruses, AAV1-CAG-FLEX-N2cG-mKate2.0 (titer: 9.83E12
vg/mL; Janelia Viral Core) and AAV5-Esyn-DIO-TVA950-EYFP-WPRE (titer: 1.44E13 vg/mL;
Salk Viral Vector Core), was injected into the SNc. 100nL of AAVDJ-Ef1a-fDIO-mScarlet (titer:

6.30E12 vg/mL; Zuckerman Institute Vector Core) was injected in the motor cortex or the striatum.

Two weeks later, 300nl of EnvA-N2c-deltaG-H2B-HA-FlpO (titer: 1.0E9 ffu/ml; Zuckerman

Institute Vector Core) was injected within the same area at a 10-degree angle (SNc; AP -3.16 mm,

ML 2.0mm, DV -4.0 mm). After the injection, the skull was cleaned and the skin sealed with

sutures and Vetbond tissue adhesive (3M, Maplewood, MN, USA). For ablation experiments, 150

nL of AAV5-EF1a-FLEX-taCasp3-TEVp virus (titer 4.20E12 vg/mL; UNC Vector Core) was

bilaterally injected into the SNc of DAT-Cre at two specific coordinates (SNc-medial: AP -3.16

mm, ML 1.10 mm, DV -4.1 mm; SNc-lateral: AP -3.16 mm, ML 1.50 mm, DV -4.1 mm). These

coordinates effectively targeted and ablated SNc dopaminergic neurons. For silencing

neurotransmission, 300nL of AAVDJ-hSyn-FLEX-TeLC-EYFP (titer 7.40E12 vg/mL; Zuckerman

Institute Vector Core) was bilaterally injected in the SNc of Anxa1-Cre (SNc; AP -3.16 mm, ML

1.3 mm, DV -4.0 mm).

**Behavior**

Mice were habituated with a wireless motion sensor in a square arena (40 x 40 x 20 cm, length x

width x height) housed inside a sound-attenuating chamber for 30 minutes on the first day.

Imaging was conducted using a 1280 x 1024 pixel FLIR camera (Point Grey Flea) with a CS

mount lens, recording at 30 Hz. For the open-field test, mice were imaged for 30 minutes from a

top view, and for the cylinder test (a cylinder arena: 15cm diameter x 30cm length), they were

imaged for 15 minutes from a side view. Video acquisition was performed using the commercial

software FlyCapture and the open-source visual language Bonsai[44]. Mouse centroid coordinates

(i.e., spine2 in DeepLabCut tracking, details provided below) during the open-field recording were

used to calculate basic measures of activity (e.g., trajectory or velocity). Acceleration data was

collected at a sampling rate of 200 Hz (details provided below).

**Pose estimation and tracking analysis**

DeepLabCut[45] was used to estimate the pose of the mice and track their movements throughout

the trials. In the open-field test, seven body parts (nose, right ear, left ear, spine1, spine2, spine3,

tail) were labeled to create a skeleton for pose estimation. For the cylinder test, two body parts

(motion sensor and nose) were labeled. A total of 1,100 frames were labeled for the open-field

test (training: 1.03 million iterations, training error: 1.72 pixels, test error: 2.76 pixels), and 1,040

frames were labeled for the cylinder test (training: 1.03 million iterations, training error: 2.31 pixels,

test error: 6.8 pixels). The pose data was filtered using a median filter and verified through visual

inspection. For the analysis in the open-field test, to ensure consistent data, we only used data

from the 5 to 20-minute mark (15 minutes total) within a 30-minute recording, as mice tend to move differently during the first few minutes after being placed from their home cage into the field. Immobility was defined as any tracking point with a speed of less than 0.5 cm/s for a duration of more than 0.5 seconds, and estimates with a likelihood lower than 0.95 were excluded. Average speed was calculated only during periods when the animal was mobile. In the cylinder test, the full 15 minutes of data were used for analysis, since mice often exhibit exploratory rearing behaviors during the first few minutes. Rearing behaviors were defined by the height of the motion sensor, visible from 360 degrees, and were considered valid if sustained for at least 0.5 seconds in the same position. The height thresholds were determined using DeepLabCut's output (i.e., filtered trajectory plots), and the number of rearing instances was recorded.

**Motion sensor acceleration data analysis**

Wireless motion sensor data was collected using the WEAR system developed by the Champalimaud Hardware Platform. The sensor is compact and lightweight (~1.8 g), capable of sampling nine-axis motion data from a three-axis accelerometer, gyroscope, and magnetometer at rates up to 200 Hz. This device communicates with computers through a base station designed by the platform, utilizing the Harp system. The base station is accessible via a software GUI (e.g., Harp Wear), allowing for easy adjustment of sensor parameters to best fit experimental needs. Additionally, it is compatible with Bonsai software[44], enabling synchronization of the WEAR sensor data with other data sources, such as cameras. The data was first filtered using a $5^{th}$ order, 1Hz low-pass Butterworth filter to separate the gravitational acceleration component of each axis. This component was then subtracted to the original acceleration data to obtain the fast acceleration changes induced by mice's movements (body acceleration). Total body acceleration was determined by calculating the vector norm of the three body acceleration axis. To assess the emergence of rhythmic, oscillatory movements, we performed a spectral analysis of the body's dorso-ventral acceleration axis. We used the signal.spectrogram function of the Scipy python library[46] to calculate a spectrogram with consecutive Fourier transforms, using segments of 256 samples with a 50% overlap. The spectrogram was normalized by transforming the power values of each window to a 0-1 range (maximum= 1 and minimum=0). For each mouse a final measure was obtained by calculating the median normalized power considering all the windows of the spectrogram.

**Histology**

Mice were deeply anesthetized with isoflurane and transcardially perfused with PBS, followed by
ice-cold 4% paraformaldehyde. Brains were then removed for histological analysis. Coronal
sections were cut at 30 µm for immunostaining or 75 µm for 3D reconstruction using a Leica
VT1000 vibratome. The tissue was rinsed twice in 1x PBS and then permeabilized in PBS
containing 0.4% Triton X-100 (PBST). For quantification of *Th+* cells in MitoPark mice and Anxa1-
Cre mouse line, immunohistochemistry was performed with primary antibodies by incubating the
sections with an anti-GFP antibody (Chicken, Aves labs Inc, GFP-1020) and a TH antibody
(Mouse, ImmunoStar, 22941) diluted at 1:2000 in 0.4% Triton X-100 PBS (PBST) overnight at 4C.
To visualize the dorsal-tier or ventral-tier of the SNc DANs, we also used a Calbindin D28K (Rabbit,
Synaptic Systems, 214002) or an Aldh1a1 antibody (Goat, R&D Systems, AF5869). The sections
were then incubated with secondary antibodies (Donkey anti-Chicken Alexa Fluor 488,
JacksonImmunoResearch, 703-545-155; Donkey anti-Goat Alexa Fluor 568, Invitrogen, A11057;
Donkey anti-Mouse Alexa Fluor 647, Invitrogen, A31571) diluted at 1:2000 in 0.4% PBST
overnight at 4C. DAPI (Sigma D9542; 1:1000) was used as a counterstain in all experiments.
601

**Image acquisition, processing, and analysis**
Coronal sections (30µm or 75 µm) were serially mounted on slides and sections were imaged
using an automated slide scanner (Nikon AZ100 Multizoom microscope) equipped with a 4x
0.4NA Plan Apo objective (Nikon Instruments Inc) and P200 slide loader (Prior Scientific),
controlled by NIS-Elements using custom acquisition scripts (Nikon Instruments Inc.). Image
processing and analysis using BrainJ proceeded as previously described[20]. Briefly, brain sections
were aligned and registered using two-dimensional (2D) rigid-body registration. After background
subtraction ilastik was used to detect cell bodies and neuronal processes[47]. Probability images
for class (soma and neuronal processes) were generated using the pixel classification approach
in ilastik, ensuring improved accuracy for segmentation over fluorescence intensity alone. To map
the location of these structures to an annotated brain atlas, serial tissue sections were registered
into a single volume before 3D registration to the template brain (Allen Brain Atlas Common
Coordinate Framework) using Elastix[48]. Initial measurements and data visualization were
performed in ImageJ, and subsequent analyses were performed in Python.
616

**Statistics and reproducibility**
All statistical analyses were conducted using R or Python. Data are presented as mean with 95%
confidence intervals (CI) or as the standard error of the mean (SEM), unless otherwise specified.
Statistical significance was assessed using a two-sided Mann–Whitney U test, and for multiple

comparisons, a two-way ANOVA followed by Tukey's HSD test was applied (*p < 0.05, **p < 0.01,

***p < 0.001, ****p < 0.0001).

## Data availability

625 Single-cell RNA-sequencing data will be deposited in the Gene Expression Omnibus website

626 ( https://www.ncbi.nlm.nih.gov/geo/info/seq.html ). All other data that support the findings of this

627 study will be available at Zenodo

## Code availability

630 Custom code used in this study will be available at Github (https://github.com/afushiki).

## Acknowledgments

We would like to thank Zuckerman Institute's Cellular Imaging platform for providing instrument

use and technical advice. Special thanks to Gabriela J. Martins, Mariana L. Correia and Helena

M Seuffert for mouse maintenance and Drew Baughman, Mafalda Vicente and Chrissy Weber-

Schmidt for additional support with lab and mouse management. We also appreciate Helio

Rodrigues for designing and constructing behavioral equipment, and Kim Ritola for creating

custom viral constructs. Our gratitude extends to Ira Schieren for assisting with FACS sorting.

This work was supported by grants from the Aligning Science Across Parkinson's (ASAP-020551)

through the Michael J. Fox Foundation for Parkinson's Research (MJFF), the Parkinson's Disease

Foundation (PDFPF-RCE-1948), the National Institute of Health (U19: 5U19NS104649-03) and

a NARSAD Young Investigator Grant from the Brain & Behavior Research Foundation. This

research was funded in part through the NIH/NCI Cancer Center Support Grant P30CA013696

and used the Genomics and High Throughput Screening. Rabies viruses were produced by the

Center for Neuroanatomy with Neurotropic Viruses, supported by the NIH (CNNV; P40

OD010996). For the purpose of open access, the author has applied a CC BY public copyright

license to all Author Accepted Manuscripts arising from this submission.

## Author contributions

A.F. and R.M.C. designed the study, interpreted the results, and wrote the manuscript. A.F.

conducted the experiments and analyzed the data. D.N. generated viral constructs and assisted

with the production of transgenic mice. Z.R.L., A.Y., B.T., and V.M. contributed to transcriptome

data analysis and interpretation of sequencing results. L.A.H. developed the automated pipeline

for anatomical reconstruction and analysis, and contributed to histological assessments. C.W.

produced the rabies virus. T.S. and J.A.d.S. assisted with accelerometer experiments and data

analysis. R.M.C. supervised the project. All authors contributed to the interpretation of results and

manuscript editing.

**Declaration of interests**

The authors declare no competing interests.

**References**

1. Barrett, M.J., Wylie, S.A., Harrison, M.B., and Wooten, G.F. (2011). Handedness and
motor symptom asymmetry in Parkinson's disease. J Neurol Neurosurg Psychiatry *82*,
1122–1124. https://doi.org/10.1136/jnnp.2010.209783.
2. Hirsch, E.C., Graybiel, A.M., and Agid, Y. (1988). Melanized dopaminergic neurons are
differentially affected in Parkinson's disease. Nature *334*, 345--348.
3. HORNYKIEWICZ, O. (1963). [The tropical localization and content of noradrenalin and
dopamine (3-hydroxytyramine) in the substantia nigra of normal persons and patients
with Parkinson's disease]. Wien Klin Wochenschr *75*, 309—312.
4. Poulin, J.F., Gaertner, Z., Moreno-Ramos, O.A., and Awatramani, R. (2020).
Classification of Midbrain Dopamine Neurons Using Single-Cell Gene Expression
Profiling Approaches. Preprint at Elsevier Ltd, https://doi.org/10.1016/j.tins.2020.01.004
https://doi.org/10.1016/j.tins.2020.01.004.
5. Azcorra, M., Gaertner, Z., Davidson, C., He, Q., Kim, H., Nagappan, S., Hayes, C.K.,
Ramakrishnan, C., Fenno, L., Kim, Y.S., et al. (2023). Unique functional responses
differentially map onto genetic subtypes of dopamine neurons. Nat Neurosci *26*, 1762–
1774. https://doi.org/10.1038/s41593-023-01401-9.
6. Poulin, J.F., Caronia, G., Hofer, C., Cui, Q., Helm, B., Ramakrishnan, C., Chan, C.S.,
Dombeck, D.A., Deisseroth, K., and Awatramani, R. (2018). Mapping projections of
molecularly defined dopamine neuron subtypes using intersectional genetic approaches.
Nat Neurosci *21*, 1260–1271. https://doi.org/10.1038/s41593-018-0203-4.
7. Damier, P., Hirsch, E.C., Agid, Y., and Graybiel, A.M. (1999). The substantia nigra of the
human brain: II. Patterns of loss of dopamine-containing neurons in Parkinson's disease.
Brain *122*, 1437–1448. https://doi.org/10.1093/brain/122.8.1437.
8. Kamath, T., Abdulraouf, A., Burris, S.J., Langlieb, J., Gazestani, V., Nadaf, N.M.,
Balderrama, K., Vanderburg, C., and Macosko, E.Z. (2022). Single-cell genomic profiling
of human dopamine neurons identifies a population that selectively degenerates in
Parkinson's disease (Springer US) https://doi.org/10.1038/s41593-022-01061-1.
9. Spillantini, M.G., Schmidt, M.L., Lee, V.M.-Y., Trojanowski, J.Q., Jakes, R., and Goedert,
M. (1997). α-Synuclein in Lewy bodies. Nature *388*, 839–840.
https://doi.org/10.1038/42166.
10. Kitada, T., Asakawa, S., Hattori, N., Matsumine, H., Yamamura, Y., Minoshima, S.,
Yokochi, M., Mizuno, Y., and Shimizu, N. (1998). Mutations in the parkin gene cause
autosomal recessive juvenile parkinsonism. Nature *392*, 605–608.
https://doi.org/10.1038/33416.

11.     Schapira, A.H. V, Cooper, J.M., Dexter, D., Clark, J.B., Jenner, P., and Marsden, C.D.
698             (1990). Mitochondrial Complex I Deficiency in Parkinson's Disease. J Neurochem *54*,
699             823–827. https://doi.org/https://doi.org/10.1111/j.1471-4159.1990.tb02325.x.
700     12.     Ekstrand, M.I., Terzioglu, M., Galter, D., Zhu, S., Hofstetter, C., Lindqvist, E., Thams, S.,
701             Bergstrand, A., Hansson, F.S., Trifunovic, A., et al. (2007). Progressive parkinsonism in
702             mice with respiratory-chain-deficient dopamine neurons. Proc Natl Acad Sci U S A *104*,
703             1325–1330. https://doi.org/10.1073/pnas.0605208103.
704     13.     Beckstead, M.J., and Howell, R.D. (2021). Progressive parkinsonism due to mitochondrial
705             impairment: Lessons from the MitoPark mouse model. Preprint at Academic Press Inc.,
706             https://doi.org/10.1016/j.expneurol.2021.113707
707             https://doi.org/10.1016/j.expneurol.2021.113707.
708     14.     Crabtree, D.M., and Zhang, J. (2012). Genetically engineered mouse models of
709             Parkinson's disease. Preprint, https://doi.org/10.1016/j.brainresbull.2011.07.019
710             https://doi.org/10.1016/j.brainresbull.2011.07.019.
711     15.     Bhatia, K.P., Bain, P.G., Bajaj, N.P.S., Elble, R.J., Hallett, M., Louis, E.D., Raethjen, J.,
712             Stamelou, M., Testa, C.M., and Deuschl, G. (2018). Consensus Statement on the
713             classification of tremors. from the task force on tremor of the International Parkinson and
714             Movement Disorder Society. Movement Disorders *33*.
715     16.     Yang, C.F., Chiang, M.C., Gray, D.C., Prabhakaran, M., Alvarado, M., Juntti, S.A., Unger,
716             E.K., Wells, J.A., and Shah, N.M. (2013). Sexually dimorphic neurons in the ventromedial
717             hypothalamus govern mating in both sexes and aggression in males. Cell *153*, 896–909.
718             https://doi.org/10.1016/j.cell.2013.04.017.
719     17.     Mazzoni, P., and Krakauer, J.W. (2006). An implicit plan overrides an explicit strategy
720             during visuomotor adaptation. Journal of Neuroscience *26*, 3642–3645.
721             https://doi.org/10.1523/JNEUROSCI.5317-05.2006.
722     18.     Wang, Q., Ding, S.L., Li, Y., Royall, J., Feng, D., Lesnar, P., Graddis, N., Naeemi, M.,
723             Facer, B., Ho, A., et al. (2020). The Allen Mouse Brain Common Coordinate Framework:
724             A 3D Reference Atlas. Cell *181*, 936-953.e20. https://doi.org/10.1016/j.cell.2020.04.007.
725     19.     Hintiryan, H., Foster, N.N., Bowman, I., Bay, M., Song, M.Y., Gou, L., Yamashita, S.,
726             Bienkowski, M.S., Zingg, B., Zhu, M., et al. (2016). The mouse cortico-striatal projectome.
727             Nat Neurosci *19*, 1100–1114. https://doi.org/10.1038/nn.4332.
728     20.     Botta, P., Fushiki, A., Vicente, A.M., Hammond, L.A., Mosberger, A.C., Gerfen, C.R.,
729             Peterka, D., and Costa, R.M. (2020). An Amygdala Circuit Mediates Experience-
730             Dependent Momentary Arrests during Exploration. Cell *183*, 605-619.e22.
731             https://doi.org/10.1016/j.cell.2020.09.023.
732     21.     Allen Institute for Brain Science Isolation of Nuclei from Adult Mouse Brain Tissue
733             Protocol V.2. https://doi.org/dx.doi.org/10.17504/protocols.io.bq7emzje.
734     22.     Yao, Z., van Velthoven, C.T.J., Kunst, M., Zhang, M., McMillen, D., Lee, C., Jung, W.,
735             Goldy, J., Abdelhak, A., Aitken, M., et al. (2023). A high-resolution transcriptomic and
736             spatial atlas of cell types in the whole mouse brain. Nature *624*, 317–332.
737             https://doi.org/10.1038/s41586-023-06812-z.
738     23.     Fonseka, C.Y., Rao, D.A., Teslovich, N.C., Korsunsky, I., Hannes, S.K., Slowikowski, K.,
739             Gurish, M.F., Donlin, L.T., Lederer, J.A., Weinblatt, M.E., et al. (2018). Mixed-effects

association of single cells identifies an expanded effector CD4+ T cell subset in
rheumatoid arthritis. Sci Transl Med *10*. https://doi.org/10.1126/scitranslmed.aaq0305.

24. Steinkellner, T., Conrad, W.S., Kovacs, I., Rissman, R.A., Lee, E.B., Trojanowski, J.Q.,
Freyberg, Z., Roy, S., Luk, K.C., Lee, V.M., et al. (2022). Dopamine neurons exhibit
emergent glutamatergic identity in Parkinson's disease. Brain *145*, 879–886.
https://doi.org/10.1093/brain/awab373.

25. Shen, H., Marino, R.A.M., McDevitt, R.A., Bi, G.H., Chen, K., Madeo, G., Lee, P.T.,
Liang, Y., De Biase, L.M., Su, T.P., et al. (2018). Genetic deletion of vesicular glutamate
transporter in dopamine neurons increases vulnerability to MPTP-induced neurotoxicity in
mice. Proc Natl Acad Sci U S A *115*, E11532–E11541.
https://doi.org/10.1073/pnas.1800886115.

26. Pereira Luppi, M., Azcorra, M., Caronia-Brown, G., Poulin, J.F., Gaertner, Z., Gatica, S.,
Moreno-Ramos, O.A., Nouri, N., Dubois, M., Ma, Y.C., et al. (2021). Sox6 expression
distinguishes dorsally and ventrally biased dopamine neurons in the substantia nigra with
distinctive properties and embryonic origins. Cell Rep *37*.
https://doi.org/10.1016/j.celrep.2021.109975.

27. Dann, E., Henderson, N.C., Teichmann, S.A., Morgan, M.D., and Marioni, J.C. (2022).
Differential abundance testing on single-cell data using k-nearest neighbor graphs. Nat
Biotechnol *40*, 245–253. https://doi.org/10.1038/s41587-021-01033-z.

28. Reardon, T.R., Murray, A.J., Turi, G.F., Wirblich, C., Croce, K.R., Schnell, M.J., Jessell,
T.M., and Losonczy, A. (2016). Rabies Virus CVS-N2cδG Strain Enhances Retrograde
Synaptic Transfer and Neuronal Viability. Neuron *89*, 711–724.
https://doi.org/10.1016/j.neuron.2016.01.004.

29. Watabe-Uchida, M., Zhu, L., Ogawa, S.K., Vamanrao, A., and Uchida, N. (2012). Whole-
Brain Mapping of Direct Inputs to Midbrain Dopamine Neurons. Neuron *74*, 858–873.
https://doi.org/10.1016/j.neuron.2012.03.017.

30. Menegas, W., Bergan, J.F., Ogawa, S.K., Isogai, Y., Venkataraju, K.U., Osten, P.,
Uchida, N., and Watabe-Uchida, M. (2015). Dopamine neurons projecting to the posterior
striatum form an anatomically distinct subclass. Elife *4*, 2014–2016.
https://doi.org/10.7554/eLife.10032.

31. Zhang, Y., Zhao, S., Rodriguez, E., Takatoh, J., Han, B.X., Zhou, X., and Wang, F.
(2015). Identifying local and descending inputs for primary sensory neurons. Journal of
Clinical Investigation *125*, 3782–3794. https://doi.org/10.1172/JCI81156.

32. Nicoletti, V., Cecchi, P., Pesaresi, I., Frosini, D., Cosottini, M., and Ceravolo, R. (2020).
Cerebello-thalamo-cortical network is intrinsically altered in essential tremor: evidence
from a resting state functional MRI study. Sci Rep *10*, 16661.
https://doi.org/10.1038/s41598-020-73714-9.

33. Crittenden, J.R., Tillberg, P.W., Riad, M.H., Shima, Y., Gerfen, C.R., Curry, J., Housman,
D.E., Nelson, S.B., Boyden, E.S., and Graybiel, A.M. (2016). Striosome-dendron
bouquets highlight a unique striatonigral circuit targeting dopamine-containing neurons.
Proc Natl Acad Sci U S A *113*, 11318–11323. https://doi.org/10.1073/pnas.1613337113.

34. Evans, R.C., Twedell, E.L., Zhu, M., Ascencio, J., Zhang, R., and Khaliq, Z.M. (2020).
Functional Dissection of Basal Ganglia Inhibitory Inputs onto Substantia Nigra

Dopaminergic Neurons. Cell Rep *32*, 108156.
https://doi.org/10.1016/j.celrep.2020.108156.

35.   Eschbach, C., Fushiki, A., Winding, M., Schneider-Mizell, C.M., Shao, M., Arruda, R.,
Eichler, K., Valdes-Aleman, J., Ohyama, T., Thum, A.S., et al. (2020). Recurrent
architecture for adaptive regulation of learning in the insect brain. Nat Neurosci *23*, 544–
555. https://doi.org/10.1038/s41593-020-0607-9.

36.   Tasic, B., Menon, V., Nguyen, T.N., Kim, T.K., Jarsky, T., Yao, Z., Levi, B., Gray, L.T.,
Sorensen, S.A., Dolbeare, T., et al. (2016). Adult mouse cortical cell taxonomy revealed
by single cell transcriptomics. Nat Neurosci *19*, 335–346. https://doi.org/10.1038/nn.4216.

37.   Tasic, B., Yao, Z., Graybuck, L.T., Smith, K.A., Nguyen, T.N., Bertagnolli, D., Goldy, J.,
Garren, E., Economo, M.N., Viswanathan, S., et al. (2018). Shared and distinct
transcriptomic cell types across neocortical areas. Nature *563*, 72–78.
https://doi.org/10.1038/s41586-018-0654-5.

38.   Schneider, V.A., Graves-Lindsay, T., Howe, K., Bouk, N., Chen, H.C., Kitts, P.A., Murphy,
T.D., Pruitt, K.D., Thibaud-Nissen, F., Albracht, D., et al. (2017). Evaluation of GRCh38
and de novo haploid genome assemblies demonstrates the enduring quality of the
reference assembly. Genome Res *27*, 849–864. https://doi.org/10.1101/gr.213611.116.

39.   Hao, Y., Stuart, T., Kowalski, M.H., Choudhary, S., Hoffman, P., Hartman, A., Srivastava,
A., Molla, G., Madad, S., Fernandez-Granda, C., et al. (2024). Dictionary learning for
integrative, multimodal and scalable single-cell analysis. Nat Biotechnol *42*, 293–304.
https://doi.org/10.1038/s41587-023-01767-y.

40.   Young, M.D., and Behjati, S. (2020). SoupX removes ambient RNA contamination from
droplet-based single-cell RNA sequencing data. Gigascience *9*.
https://doi.org/10.1093/gigascience/giaa151.

41.   McGinnis, C.S., Murrow, L.M., and Gartner, Z.J. (2019). DoubletFinder: Doublet
Detection in Single-Cell RNA Sequencing Data Using Artificial Nearest Neighbors. Cell
Syst *8*, 329-337.e4. https://doi.org/10.1016/j.cels.2019.03.003.

42.   Korsunsky, I., Millard, N., Fan, J., Slowikowski, K., Zhang, F., Wei, K., Baglaenko, Y.,
Brenner, M., Loh, P. ru, and Raychaudhuri, S. (2019). Fast, sensitive and accurate
integration of single-cell data with Harmony. Nat Methods *16*, 1289–1296.
https://doi.org/10.1038/s41592-019-0619-0.

43.   Reimand, J., Arak, T., Adler, P., Kolberg, L., Reisberg, S., Peterson, H., and Vilo, J.
(2016). g:Profiler-a web server for functional interpretation of gene lists (2016 update).
Nucleic Acids Res *44*, W83–W89. https://doi.org/10.1093/NAR/GKW199.

44.   Lopes, G., Bonacchi, N., Frazão, J., Neto, J.P., Atallah, B. V., Soares, S., Moreira, L.,
Matias, S., Itskov, P.M., Correia, P.A., et al. (2015). Bonsai: An event-based framework
for processing and controlling data streams. Front Neuroinform *9*.
https://doi.org/10.3389/fninf.2015.00007.

45.   Mathis, A., Mamidanna, P., Cury, K.M., Abe, T., Murthy, V.N., Mathis, M.W., and Bethge,
M. (2018). DeepLabCut: markerless pose estimation of user-defined body parts with deep
learning. Nat Neurosci *21*, 1281–1289. https://doi.org/10.1038/s41593-018-0209-y.

46.   Virtanen, P., Gommers, R., Oliphant, T.E., Haberland, M., Reddy, T., Cournapeau, D.,
Burovski, E., Peterson, P., Weckesser, W., Bright, J., et al. (2020). SciPy 1.0:

826   fundamental algorithms for scientific computing in Python. Nat Methods *17*, 261–272.
827   https://doi.org/10.1038/s41592-019-0686-2.

828 47. Berg, S., Kutra, D., Kroeger, T., Straehle, C.N., Kausler, B.X., Haubold, C., Schiegg, M.,
829   Ales, J., Beier, T., Rudy, M., et al. (2019). ilastik: interactive machine learning for
830   (bio)image analysis. Nat Methods *16*, 1226–1232. https://doi.org/10.1038/s41592-019-
831   0582-9.

832 48. Klein, S., Staring, M., Murphy, K., Viergever, M.A., and Pluim, J.P.W. (2010). elastix: A
833   Toolbox for Intensity-Based Medical Image Registration. IEEE Trans Med Imaging *29*,
834   196–205. https://doi.org/10.1109/TMI.2009.2035616.

# Figure1

**Figure1. Progressive development of PD-like motor deficits in MitoPark mice**

**A.** Cumulative trajectories of a representative MitoPark mouse and their control at different

stages (8, 16 and 24wks) in the open-field test (black: control animals, red: MitoPark mice).

**B-D**. Behavioral characterization of MitoPark mice and their littermate controls at each stage.

**(B)** Immobility time: MitoPark mice at later stages spent significantly more time immobile

compared to controls. The inset shows tracking performed using DeepLabCut. **(C)** Average

speed during mobility: the speed of MitoPark mice was significantly lowered in 16 and 24wks.

**(D)** Number of stops: the number of pauses in MitoPark mice showed significant differences

compared to the controls. Statistical significance was determined by two-sided Mann–

Whitney U-test (***$p < 0.001$; ns = not statistically significant). Error bars show 95% confidence

intervals. MitoPark 8weeks, n=6; Littermate controls 8weeks, n=6; MitoPark 16weeks, n=12;

Littermate controls 16weeks, n=8; MitoPark 24weeks, n=9; Littermate controls 24weeks, n=15.

**E-H.** Frequency analysis of total time **(E)** or immobility periods **(F)** at different ages of the

animals. Mitopark mice showed a pronounced increase in the power of acceleration oscillations

at the 12-18Hz range that becomes more prominent in the later stages. **(G, H)** The scatter plots

show the correlation between age and mean power in the 12-18Hz frequency band for Mitopark

mice and control animals during total period **(G)** or immobile period **(H)**. Pearson's correlation

coefficient reveals a strong positive correlation during both the total period (G: red, r=0.88,

$p<0.001$) and the immobile period (H: red, r=0.77, $p<0.01$) in MitoPark mice. The solid line

represents the linear regression model (G: red, $y = 6.3e\text{-}03x + 0.036$, $R^2 = 0.78$; H: red, $y =$

$5.6e\text{-}03x + 0.036$, $R^2 = 0.60$), demonstrating a statistically significant relationship between the

two variables in MitoPark mice. MitoPark 8weeks, n=3; Littermate controls 8weeks, n=3;

MitoPark 16weeks, n=4; Littermate controls 16weeks, n=3; MitoPark 24weeks, n=5; Littermate

controls 24weeks, n=6. **I-O.** Injections of Casp3 virus in the SNc of DAT-Cre mice led to the

similar phenotype of MitoPark mice 24 weeks. **(I)** The images demonstrated a specific reduction

in dopamine neurons (DANs) within the SNc, as confirmed by staining with anti-TH antibody

(green). **(J-M)** The DAT-Casp3 animals showed motor deficits characterized by diminished

movement speed, increased incidences of halting and reduced the frequency of rearing

behaviors. Statistical significance was determined by two-sided Mann–Whitney U-test (*$p <$

$0.05$; ns = not statistically significant). Error bars show 95% confidence intervals. DAT-Casp3

animals, n=5; Controls, n=5. **(N, O)** The virus injection led to the presentation of the same 12-

18Hz oscillation seen in MitoPark mice at 24 weeks.

# Figure2

**Figure 2. Progressive degeneration of dopamine neurons in MitoPark mice**

**A-C.** TH immunohistochemistry of MitoPark mice and their littermate controls at 8, 16 and 24

weeks. (A) Histological images of the mice at each stage. Scale bars: 400µm for the substantia

nigra (top) and 1mm for the striatum (bottom). (B-C) MitoPark mice exhibited a gradual and

preferential loss of DANs in the substantia nigra, and their projections (each n=1). **D.** The

workflow for single-nucleus RNA (snRNA) sequencing commenced with the collection of all

tissues from the ventral midbrain (red dashed lines). The tissue samples underwent nuclei

isolation, followed by the selection of DAPI+ nuclei. These nuclei were subsequently processed

using the 10x Genomics platform for sequencing. **E.** UMAP visualization of snRNA-seq data

from the ventral midbrain. Data are combined across all animals (MitoPark mice and their

controls) and classified by the Allen Brain Cell (ABC) Atlas class annotations. Nuclei that are

limited in number or belong to infrequent categories are designated as 'others.' **F.** Feature plot

showing expression of marker genes used to label the main class of cells: *Th* (DANs), *Slc17a6*

(excitatory neurons) and *Slc32a1* (inhibitory neurons). **G-I.** Dots and whiskers represent odds-

ratio (OR) with 95% confidence interval (CI) obtained from MASC. OR estimates of major cell

types associated with MitoPark mice (color; false discovery rate (FDR)-adjusted P<0.05). **(G)**

MitoPark 16 weeks versus their littermate controls; 21_MB_Dopa (OR=-0.95, FDR-adjusted

P<0.05). **(H)** MitoPark 24 weeks versus their littermate controls; 21_MB_Dopa (OR=-1.26, FDR-

adjusted P<0.05). **(I)** MitoPark 16 weeks versus 8 weeks; 21_MB_Dopa (OR=-0.51, FDR-

adjusted P=0.057).

# Figure3

**Figure3. Subtype-specific degeneration within the dopaminergic populations**

**A.** UMAP representation of 18,611 dopamine neuron nuclei, colored by the supertype
annotation of the Allen Brain Cell (ABC) Atlas. In supertype annotation of the ABC Atlas, DANs
are largely classified into 8 types. **B.** Representative feature plots showing gene expression
associated with dopamine used to label the main class of cells. **C.** Distribution of dopamine
subtype proportions across MitoPark and their control animals. 0882_Sox6 is mostly affected
among other dopamine neuron subtypes. Statistical significance was determined by two-way
ANOVA (p-value indicates the interaction effect between genotype and age) followed by
Tukey's HSD test for multiple comparisons (*p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001).
**D-F.** Odds-ratio estimates of each dopamine subpopulation in **(D)** MitoPark 16 weeks versus
their littermate controls; 0882_Sox6 (OR=-1.72, FDR-adjusted P<0.05), 0880_Megf11

907    (OR=1.00, FDR-adjusted P<0.05) and 0883_Cck (OR=1.13, FDR-adjusted P<0.05). **(E)**

908    MitoPark 24 weeks versus their littermate controls; 0882_Sox6 (OR=-2.49, FDR-adjusted

909    P<0.05), 0887_Synpr (OR=1.76, FDR-adjusted P<0.05) and 0880_Megf11 (OR=1.31, FDR-

910    adjusted P<0.05). **(F)** MitoPark 16 weeks versus 8 weeks; 0882_Sox6 (OR=-0.93, FDR-

911    adjusted P=0.057).

**Figure4**

**Figure4. A subset of *Sox6*+ population is more vulnerable than others in MitoPark mice**

Differential abundance analysis for the dopamine clusters of MitoPark mice and their littermate controls. The analysis reveals a subtype-specific reduction in MitoPark mice at a later stage. **A.** Distribution of *Sox6*+ dopamine subtype proportions across MitoPark and their control animals. 3857_Anxa1 is significantly affected among other dopamine neuron subtypes at 16 and 24 weeks. Statistical significance was determined by two-way ANOVA (p-value indicates the interaction effect between genotype and age) followed by Tukey's HSD test for multiple comparisons (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$). **B-D.** UMAP plots showing the dopamine subpopulations including MitoPark mice and their littermate controls (n=8 at each stage and genotype). Forty-three clusters are annotated by the Allen Brain Cell (ABC) Atlas. **E-G.** Neighborhood (Nhood) graphs with the results from Milo differential abundance testing between MitoPark mice and controls (E:16 weeks, F: 24 weeks), and between MitoPark mice at different stages (G: 8 and 16 weeks). Nodes represent neighborhoods, colored by their log fold change compared to their controls (blue: less abundant, red: more abundant, gray: non-differentially abundant). Graph edges show the number of cells shared between two neighborhoods. **H-J**. Beeswarm plots display the distribution of log-fold changes in cell abundance. Dots representing neighborhoods that overlap with the same cell populations are grouped together. Colors are represented similarly to **E-G**. **K.** Histological images representing 8 and 16 weeks of MitoPark mice reveal that significant cell depletion in the ventral region, particularly in the rostral part of the SNc (TH: green, ALDH1A1: magenta). Scale bar 400μm.

# Figure5

**Figure5. Input-output circuit architecture of *Anxa1+* DANs in the SNc**

**A.** In the Anxa1-Cre line, the majority of neurons within the SNc are positive for *Th* and negative for *Calbindin1* (TH, red; CALB1, magenta; n=2). Scale bars: 400µm. **B.** Axonal projections of *Anxa1+* DANs in the SNc. They preferentially project to dorsal striatum that integrates sensorimotor information from cortical and thalamic regions. Scale bars: 1mm. **C-D.** Brain regions to which all DANs or *Anxa1+* DANs in the SNc project, measured as the fraction

of neurites found within those brain structures defined Allen brain atlas (each n=3). Blue

horizontal line indicates dorsal striatum and the red line indicates ventral striatum. **E.**

Representative images of rabies infected cells (tdTomato). Rabies virus was injected in the SNc

of Anxa1-Cre counterstained with DAPI in blue. Scale bar 1mm. **F-I.** Statistical analysis of the

whole-brain distribution of ipsilateral (left) or contralateral (right) monosynaptic inputs to all

DANs (F, G) or *Anxa1+* dopamine neuron subtype (H, I) in the SNc. Average proportion of

tdTomato-labeled neurons in approximately 50 brain regions, each with more than 200 cells,

was greater than 0.2% of the total input to DANs in DAT-Cre mice (n=7) and Anxa1-Cre mice

(n=5; 4 animals for Anxa1-Cre+ and 1 animal for DAT-Flp+;Anxa1-Cre+). Brain areas are color-

coded by the Allen Brain Atlas.

**[Acronym]** ACAd5, Anterior cingulate area, dorsal part, layer 5; ACAv5, Anterior cingulate area,

ventral part, layer 5; ACB, Nucleus accumbens; AId5, Agranular insular area, dorsal part, layer

5; AId6a, Agranular insular area, dorsal part, layer 6a; APN, Anterior pretectal nucleus; BLAa,

Basolateral amygdalar nucleus, anterior part; CEAc, Central amygdalar nucleus, capsular part;

CEAl, Central amygdalar nucleus, lateral part; CEAm, Central amygdalar nucleus, medial part;

CPc_d, Caudal Caudoputamen, dorsal; CPc_i, Caudal Caudoputamen, intermediate; CPc_v,

Caudal Caudoputamen, ventral; CPi_dl, Intermediate Caudoputamen, dorsolateral; CPi_dm,

Intermediate Caudoputamen, dorsomedial; CPi_vl, Intermediate Caudoputamen, ventrolateral;

CPi_vm, Intermediate Caudoputamen, ventromedial; CPr_imd, Rostral Caudoputamen,

intermediate dorsal; CPr_imv, Rostral Caudoputamen, intermediate ventral; CPr_l, Rostral

Caudoputamen, lateral; CPr_m, Rostral Caudoputamen, medial; EW, EdingerWestphal nucleus;

FF, Fields of Forel; FS, Fundus of striatum; Gpe, Globus pallidus, external segment; Gpi,

Globus pallidus, internal segment; HY, Hypothalamus; IF, Interfascicular nucleus raphe; IPRL,

Interpeduncular nucleus, rostrolateral; LHA, Lateral hypothalamic area; MB, Midbrain; MOp5,

Primary motor area, Layer 5; MOp6a, Primary motor area, Layer 6a; MOs5, Secondary motor

area, layer 5; MOs6a, Secondary motor area, layer 6a; MRN, Midbrain reticular nucleus; MT,

Medial terminal nucleus of the accessory optic tract; ORBl5, Orbital area, lateral part, layer 5;

OT, Olfactory tubercle; PAG, Periaqueductal gray; PAL, Pallidum; PIL, Posterior intralaminar

thalamic nucleus; PL5, Prelimbic area, layer 5; PN, Paranigral nucleus; PP, Peripeduncular

nucleus; PST, Preparasubthalamic nucleus; PSTN, Parasubthalamic nucleus; RL, Rostral linear

nucleus raphe; RN, Red nucleus; RR, Midbrain reticular nucleus, retrorubral area; RSPd5,

Retrosplenial area, dorsal part, layer 5; RSPv5, Retrosplenial area, ventral part, layer 5; SCdg,

Superior colliculus, motor related, deep gray layer; Scig, Superior colliculus, motor related,

intermediate gray layer; Sciw, Superior colliculus, motor related, intermediate white layer; SI,

Substantia innominata; SNr, Substantia nigra, reticular part; SPFp, Subparafascicular nucleus,

parvicellular part; SSpbfd5, Primary somatosensory area, barrel field, layer 5; SSpll5, Primary

somatosensory area, lower limb, layer 5; SSpm5, Primary somatosensory area, mouth, layer 5;

SSpul5, Primary somatosensory area, upper limb, layer 5; SSs5, Supplemental somatosensory

area, layer 5; STN, Subthalamic nucleus; STR, Striatum; TEa5, Temporal association areas,

layer 5; VISC5, Visceral area, layer 5; VTA, Ventral tegmental area; ZI, Zona incerta

# Figure6

**Figure6. Inhibition of *Anxa1+* DANs leads to motor deficits resembling early onset PD-like symptoms**

**A.** In the Anxa1-Cre line, targeted expression of the tetanus toxin light chain (TeLC) specifically eliminated dopamine synaptic transmission in the dorsal striatum (white arrows). Scale bars: 1mm for the striatum (top) and 400μm for the substantia nigra (bottom). **B-D.** Behavioral characterization of Anxa1-TeLC animals. **(B)** Cumulative trajectories of a representative Anxa1-TeLC mouse and their control in the open-field test (black: a control animal, red: Anxa1-TeLC mouse). **(C)** Anxa1-TeLC animals demonstrated slowed movement; however, there were no significant differences observed in terms of immobility duration or the frequency of stops. The frequency of rearing behaviors in the Anxa1-TeLC animals was significantly reduced. Statistical significance was determined by two-sided Mann–Whitney U-test (**p < 0.01, *p < 0.05; ns = not

statistically significant). Error bars show 95% confidence intervals. Anxa1-TeLC animals, n=5;

Controls, n=4. **(D)** This figure shows the probability density function (PDF) for stop duration (left)

and move duration (right), with duration (in seconds) on the x-axis and probability density on the

y-axis. There is no significant difference in the density of stop or move durations between

Anxa1-TeLC and control animals. **E.** Anxa1-TeLC animals showed the similar 12-18Hz

oscillation seen in MitoPark mice at 16 weeks.

# Figure S1.

**A** MitoPark 24wks

**B** MitoPark 24wks

Control 24wks

Control 24wks

**C** Stop duration: MitoPark 8wks — Stop duration: MitoPark 16wks — Stop duration: MitoPark 24wks — **E** Stop duration: DAT-Casp3

**D** Move duration: MitoPark 8wks — Move duration: MitoPark 16wks — Move duration: MitoPark 24wks — **F** Move duration: DAT-Casp3

1000

**Figure S1. A specific oscillation within the 12-18Hz frequency band in the 24 weeks of**

**MitoPark mice, Related to Figure1.**

**A-B.** Examples of raw traces of dynamic acceleration in MitoPark animal 24 weeks (red) and a

control animal (black), with an enlarged view (gray square in A) shown in **B**. MitoPark mice often

show specific 12-18Hz oscillations. **C-D.** These figures show the probability density function

(PDF) for stop duration (C) and move duration (D), with duration (in seconds) on the x-axis and

probability density on the y-axis. In MitoPark mice, stop duration increased, while move duration

decreased at the later stage. **E-F.** The PDF for DAT-Casp3 animals, which also showed an

increase in stop duration and a decrease in move duration, similar to the pattern observed in the

later stage of MitoPark mice.

1011

**Figure S2.**

1012

**Figure S2. An updated atlas incorporates striatal subregions and enhances the analysis**

**of cells and projections, Related to Figure2 and Figure5.**

**A.** Top: The original Allen Common Coordinate Framework (Allen CCF; version 3 released in

2017). Middle: An updated version of the Allen CCF, incorporating subregions of the striatum

(CP: caudate putamen) based on the cortical projections shown at the bottom (CPr_imd: Rostral

CP – Intermediate dorsal, CPr_l: Rostral CP – Lateral, CPr_imv: Rostral CP – Intermediate

ventral, CPr_m: Rostral CP – Medial, CPi_dm: Intermediate CP – Dorsomedial, CPi_dl:

Intermediate CP – Dorsolateral, CPi_vl: Intermediate CP – Ventrolateral, CPi_vm: Intermeidiate

CP – Ventromedial, CPc_d: Caudal CP – Dorsal, CPc_i: Caudal CP – Intermediate, CPc_v:

Caudal CP – Ventral). **B.** Alignment of coordinates between the ABA (Allen Brain Atlas) and

Allen CCFv3, with manually segmented subregions of the striatum. **C**. Comparison of projection

density of dopamine neurons in SNc using the original atlas versus the updated atlas. The

original atlas (top) does not include subdivisions of the caudate-putamen (CP), while the

updated atlas (bottom) defines 11 subregions based on cortical projections, providing a more

detailed representation of local projections within the striatum.

# Figure S3.

1028

**Figure S3. The integrated dataset with the Allen Brain Cell (ABC) Atlas, Related to Figure2.**

**A.** UMAP plots showing all nuclei in the ventral midbrain of MitoPark mice and their controls with the class (left) or subclass (right) annotation of the ABC Atlas. Dopamine cluster is classified as Class: 21_MB_Dopa or Subclass: 215_SNc_VTA_RAmb_Foxa1_Dopa. **B.** Representative feature plots to label main class of cells. Neuronal cells, *Rbfox3*; Oligodendrocytes, *Mog*; Oligodendrocyte precursor cells (OPCs), *Pdgfra*; Microglia, *Ptprc*; Astrocytes, *Gja*; Endothelial cells, *Nostrin*.

# Figure S4.

**Figure S4. A significant reduction in DANs was observed in the later weeks of MitoPark mice, Related to Figure2.**

**A.** UMAP plots showing all nuclei in the ventral midbrain of MitoPark mice and their controls with the class annotation of the ABC Atlas. Dopamine cluster (black arrow) becomes visibly thinner than other cell types in MitoPark mice. **B.** Box and whisker plots showing the fraction of major cell types originating from the ventral midbrain of MitoPark mice, their littermate controls and B6

WT mice (each n=8), with plot center, box and whiskers corresponding to median, interquartile

range (IQR) and 1.5xIQR, respectively. MitoPark mice in the later stages exhibited a significant

reduction in the proportion of DANs (21_MB_Dopa), without affecting other cell types. Statistical

significance was determined by two-way ANOVA (p-value indicates the interaction effect

between genotype and age) followed by Tukey's HSD test for multiple comparisons (*$p < 0.05$,

**$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$). **C-E.** Odds-ratio estimated of major cell types

annotated by subclass. **(C)** MitoPark 16 weeks versus their littermate controls;

215_SNc_VTA_RAmb_Foxa1_Dopa (OR=-0.95, FDR-adjusted P<0.05). **(D)** MitoPark 24 weeks

versus their littermate controls; 215_SNc_VTA_RAmb_Foxa1_Dopa (OR=-1.26, FDR-adjusted

P<0.05). **(E)** MitoPark 16 weeks versus 8 weeks; 215_SNc_VTA_RAmb_Foxa1_Dopa (OR=-

0.51, FDR-adjusted P=0.086).

**Figure S5.**

1055

**Figure S5. The dopaminergic cluster with distinct marker genes, Related to Figure3.**

**A.** UMAP plots showing dopamine nuclei in the MitoPark mice and their controls with the

annotations (Class, Subclass, Supertype and Cluster) of the ABC Atlas. **B.** UMAP plots

illustrating the distribution of dopamine nuclei, categorized by supertype annotation based on

the species and age. **C.** Top five marker genes for each dopamine neuron subtype categorized

by supertype annotation

**Figure S6.**

1062

**Figure S6. Marker feature expression in the dopaminergic cluster, Related to Figure3.**

**A.** Feature plots depicting the expression of marker genes across dopamine clusters, classified

according to supertype annotations.**B.** Violin plot indicating the expression distribution of the top

marker gene of each cluster. **C.** Clustered dot plot profiling of marker genes for each dopamine

cluster. X-axis represents the dopamine clusters, and the Y-axis shows the top three marker

genes. Dot diameter indicates the proportion of cluster nuclei expressing a given gene. The

color scale represents the gene expression levels, with black indicating high expression and

white indicating low expression. **D.** Distribution of dopamine subtype proportions across

MitoPark, their control animals and B6 WT animals. **E-G.** There is no significant age-related

degeneration in dopamine populations. **(E)** MitoPark littermate controls versus B6 WT animals

at 8 weeks; 0882_Sox6 (OR=-0.14, FDR-adjusted P=0.80). **(E)** MitoPark littermate controls

versus B6 WT animals at 16 weeks; 0882_Sox6 (OR=-0.21, FDR-adjusted P<0.41). **(F)**

MitoPark littermate controls versus B6 WT animals at 24 weeks; 0882_Sox6 (OR=0.20, FDR-

adjusted P<0.11).

# Figure S7.

1077

**Figure S7. Selective vulnerability of *Sox6*+ dopaminergic population revealed by**

**differential abundance test with Milo, Related to Figure4.**

Differential abundance testing for the dopamine subpopulations with supertype annotations by

the ABC atlas. **A-C.** UMAP plots showing the dopamine subpopulations of MitoPark mice and

their controls. **D-F.** Neighborhood graphs with the results from Milo differential abundance

analysis between MitoPark mice and their littermate controls (D:16 weeks, E: 24 weeks), and

1084      between MitoPark mice at different stage (F: 8 weeks and 16 weeks). **G-I**. Beeswarm plots

1085      showing the distribution of log-fold changes in each supertype group. Colors are represented

1086      similarly to **D-F**. **J-L**. Beeswarm plots showing the distribution of log-fold changes in each

1087      cluster. Colors are represented similarly to **Figure4D-F**.

# Figure S8.

**A**

### GO Table
Based on dopamine subcluster

| Source | Term name | Term ID | Cluster | log10(1/padj) |
|---|---|---|---|---|
| GO:BP | nervous system development | GO:0007399 | 3854_Grin2c | 85.68 |
| GO:BP | neurogenesis | GO:0022008 | 3854_Grin2c | 76.55 |
| GO:BP | generation of neurons | GO:0048699 | 3854_Grin2c | 74.69 |
| GO:CC | cell junction | GO:0030054 | 3854_Grin2c | 111.69 |
| GO:CC | synapse | GO:0045202 | 3854_Grin2c | 97.04 |
| GO:CC | cytoplasm | GO:0005737 | 3854_Grin2c | 79.54 |
| GO:MF | protein binding | GO:0005515 | 3854_Grin2c | 85.87 |
| GO:MF | binding | GO:0005488 | 3854_Grin2c | 33.21 |
| GO:MF | ion binding | GO:0043167 | 3854_Grin2c | 32.86 |
| HP | Abnormal muscle tone | HP:0003808 | 3854_Grin2c | 5.27 |
| HP | Abnormal nervous system physiology | HP:0012638 | 3854_Grin2c | 5.27 |
| HP | Abnormality of mental function | HP:0011446 | 3854_Grin2c | 4.99 |
| KEGG | Axon guidance | KEGG:04360 | 3854_Grin2c | 9.54 |
| KEGG | Glutamatergic synapse | KEGG:04724 | 3854_Grin2c | 8.91 |
| KEGG | Morphine addiction | KEGG:05032 | 3854_Grin2c | 8.70 |
| MIRNA | mmu-miR-15a-5p | MIRNA:mmu-miR-15a-5p | 3854_Grin2c | 3.45 |
| MIRNA | mmu-miR-340-5p | MIRNA:mmu-miR-340-5p | 3854_Grin2c | 3.33 |
| MIRNA | mmu-miR-149-5p | MIRNA:mmu-miR-149-5p | 3854_Grin2c | 3.01 |
| REAC | Neuronal System | REAC:R-MMU-112316 | 3854_Grin2c | 19.47 |
| REAC | Transmission across Chemical Synapses | REAC:R-MMU-112315 | 3854_Grin2c | 10.89 |
| REAC | Protein-protein interactions at synapses | REAC:R-MMU-6794362 | 3854_Grin2c | 8.66 |
| TF | Factor: ZF5; motif: GSGCGCGR; match class: 1 | TF:M00716_1 | 3854_Grin2c | 112.68 |
| TF | Factor: ZF5; motif: GSGCGCGR | TF:M00716 | 3854_Grin2c | 98.98 |
| TF | Factor: ZF5; motif: NRNGNGCGCGCWN; match class: 1 | TF:M00333_1 | 3854_Grin2c | 95.72 |
| WP | Dravet syndrome Scn1a A1783V point mutation model | WP:WP5298 | 3854_Grin2c | 4.88 |
| WP | Splicing factor NOVA regulated synaptic proteins | WP:WP1983 | 3854_Grin2c | 3.98 |
| WP | G protein signaling pathways | WP:WP232 | 3854_Grin2c | 3.49 |

**B**

### GO Table
Based on dopamine subcluster

| Source | Term name | Term ID | Cluster | log10(1/padj) |
|---|---|---|---|---|
| GO:BP | nervous system development | GO:0007399 | 3855_Arhgap28 | 72.23 |
| GO:BP | neurogenesis | GO:0022008 | 3855_Arhgap28 | 62.71 |
| GO:BP | system development | GO:0048731 | 3855_Arhgap28 | 59.98 |
| GO:CC | cell junction | GO:0030054 | 3855_Arhgap28 | 62.84 |
| GO:CC | neuron projection | GO:0043005 | 3855_Arhgap28 | 59.86 |
| GO:CC | synapse | GO:0045202 | 3855_Arhgap28 | 56.83 |
| GO:MF | protein binding | GO:0005515 | 3855_Arhgap28 | 39.92 |
| GO:MF | cytoskeletal protein binding | GO:0008092 | 3855_Arhgap28 | 19.07 |
| GO:MF | binding | GO:0005488 | 3855_Arhgap28 | 18.82 |
| HP | Tremor | HP:0001337 | 3855_Arhgap28 | 4.95 |
| HP | Myoclonus | HP:0001336 | 3855_Arhgap28 | 3.86 |
| HP | Involuntary movements | HP:0004305 | 3855_Arhgap28 | 3.84 |
| KEGG | Morphine addiction | KEGG:05032 | 3855_Arhgap28 | 9.82 |
| KEGG | Glutamatergic synapse | KEGG:04724 | 3855_Arhgap28 | 7.65 |
| KEGG | GABAergic synapse | KEGG:04727 | 3855_Arhgap28 | 7.18 |
| MIRNA | mmu-miR-758-3p | MIRNA:mmu-miR-758-3p | 3855_Arhgap28 | 3.94 |
| MIRNA | mmu-miR-425-5p | MIRNA:mmu-miR-425-5p | 3855_Arhgap28 | 2.87 |
| REAC | Neuronal System | REAC:R-MMU-112316 | 3855_Arhgap28 | 15.38 |
| REAC | Transmission across Chemical Synapses | REAC:R-MMU-112315 | 3855_Arhgap28 | 9.58 |
| REAC | Neurotransmitter receptors and postsynaptic signal transmission | REAC:R-MMU-112314 | 3855_Arhgap28 | 2.62 |
| TF | Factor: Sp1; motif: GGNGGGGGNGGGGGMGGGGCNGGG | TF:M10375 | 3855_Arhgap28 | 33.09 |
| TF | Factor: CTCF; motif: NAGGGGCGCNNKNNNN; match class: 1 | TF:M08995_1 | 3855_Arhgap28 | 27.03 |
| TF | Factor: ZF5; motif: NRNGNGCGCGCWN; match class: 1 | TF:M00333_1 | 3855_Arhgap28 | 25.14 |
| WP | Dravet syndrome Scn1a A1783V point mutation model | WP:WP5298 | 3855_Arhgap28 | 5.10 |
| WP | Dopaminergic neurogenesis | WP:WP1498 | 3855_Arhgap28 | 3.95 |
| WP | Splicing factor NOVA regulated synaptic proteins | WP:WP1983 | 3855_Arhgap28 | 3.32 |

**C**

### GO Table
Based on dopamine subcluster

| Source | Term name | Term ID | Cluster | log10(1/padj) |
|---|---|---|---|---|
| CORUM | Sh3kbp1-Cblb-Egfr complex | CORUM:5172 | 3857_Anxa1 | 1.30 |
| GO:BP | nervous system development | GO:0007399 | 3857_Anxa1 | 91.15 |
| GO:BP | neurogenesis | GO:0022008 | 3857_Anxa1 | 83.86 |
| GO:BP | system development | GO:0048731 | 3857_Anxa1 | 82.54 |
| GO:CC | cell junction | GO:0030054 | 3857_Anxa1 | 113.87 |
| GO:CC | synapse | GO:0045202 | 3857_Anxa1 | 99.92 |
| GO:CC | neuron projection | GO:0043005 | 3857_Anxa1 | 89.89 |
| GO:MF | protein binding | GO:0005515 | 3857_Anxa1 | 74.58 |
| GO:MF | binding | GO:0005488 | 3857_Anxa1 | 32.25 |
| GO:MF | ion binding | GO:0043167 | 3857_Anxa1 | 25.68 |
| HP | Autistic behavior | HP:0000729 | 3857_Anxa1 | 7.33 |
| HP | Abnormal communication | HP:0034434 | 3857_Anxa1 | 5.91 |
| HP | Abnormality of speech or vocalization | HP:0002167 | 3857_Anxa1 | 5.58 |
| KEGG | Axon guidance | KEGG:04360 | 3857_Anxa1 | 9.78 |
| KEGG | GABAergic synapse | KEGG:04727 | 3857_Anxa1 | 9.63 |
| KEGG | Glutamatergic synapse | KEGG:04724 | 3857_Anxa1 | 9.39 |
| MIRNA | mmu-miR-758-3p | MIRNA:mmu-miR-758-3p | 3857_Anxa1 | 2.53 |
| MIRNA | mmu-miR-425-5p | MIRNA:mmu-miR-425-5p | 3857_Anxa1 | 2.44 |
| MIRNA | mmu-miR-9-5p | MIRNA:mmu-miR-9-5p | 3857_Anxa1 | 2.41 |
| REAC | Neuronal System | REAC:R-MMU-112316 | 3857_Anxa1 | 29.87 |
| REAC | Transmission across Chemical Synapses | REAC:R-MMU-112315 | 3857_Anxa1 | 19.52 |
| REAC | Neurotransmitter receptors and postsynaptic signal transmission | REAC:R-MMU-112314 | 3857_Anxa1 | 10.51 |
| TF | Factor: ZF5; motif: GSGCGCGR; match class: 1 | TF:M00716_1 | 3857_Anxa1 | 75.40 |
| TF | Factor: ZF5; motif: GSGCGCGR | TF:M00716 | 3857_Anxa1 | 65.82 |
| TF | Factor: ZF5; motif: NRNGNGCGCGCWN; match class: 1 | TF:M00333_1 | 3857_Anxa1 | 63.72 |
| WP | Dravet syndrome Scn1a A1783V point mutation model | WP:WP5298 | 3857_Anxa1 | 8.91 |
| WP | Calcium regulation in cardiac cells | WP:WP553 | 3857_Anxa1 | 6.34 |
| WP | Cholesterol biosynthesis | WP:WP103 | 3857_Anxa1 | 6.15 |

**D**

### GO Table
Based on dopamine subcluster

| Source | Term name | Term ID | Cluster | log10(1/padj) |
|---|---|---|---|---|
| CORUM | Caveolar macromolecular signaling complex | CORUM:384 | 3858_Glra2 | 1.31 |
| CORUM | Caveolar macromolecular signaling complex, using anti-Cav-3 | CORUM:385 | 3858_Glra2 | 1.31 |
| GO:BP | system development | GO:0048731 | 3858_Glra2 | 32.92 |
| GO:BP | generation of neurons | GO:0048699 | 3858_Glra2 | 32.89 |
| GO:BP | neuron differentiation | GO:0030182 | 3858_Glra2 | 32.85 |
| GO:CC | synapse | GO:0045202 | 3858_Glra2 | 57.14 |
| GO:CC | cell junction | GO:0030054 | 3858_Glra2 | 53.72 |
| GO:CC | neuron projection | GO:0043005 | 3858_Glra2 | 41.44 |
| GO:MF | protein binding | GO:0005515 | 3858_Glra2 | 19.51 |
| GO:MF | gated channel activity | GO:0022836 | 3858_Glra2 | 13.58 |
| GO:MF | monoatomic ion channel activity | GO:0005216 | 3858_Glra2 | 12.19 |
| HP | Epileptic encephalopathy | HP:0200134 | 3858_Glra2 | 1.66 |
| KEGG | Axon guidance | KEGG:04360 | 3858_Glra2 | 8.92 |
| KEGG | Oxytocin signaling pathway | KEGG:04921 | 3858_Glra2 | 6.14 |
| KEGG | Aldosterone synthesis and secretion | KEGG:04925 | 3858_Glra2 | 5.31 |
| REAC | Neuronal System | REAC:R-MMU-112316 | 3858_Glra2 | 17.64 |
| REAC | Transmission across Chemical Synapses | REAC:R-MMU-112315 | 3858_Glra2 | 7.65 |
| REAC | Axon guidance | REAC:R-MMU-422475 | 3858_Glra2 | 4.81 |
| TF | Factor: CTCF; motif: NAGGGGCGCNNKNNNN; match class: 1 | TF:M08995_1 | 3858_Glra2 | 15.79 |
| TF | Factor: ZF5; motif: NRNGNGCGCGCWN; match class: 1 | TF:M00333_1 | 3858_Glra2 | 15.69 |
| TF | Factor: FOXN4; motif: NNWANNCGWMCGCGTCNNNNMT; match class: 1 | TF:M04662_1 | 3858_Glra2 | 15.56 |
| WP | Splicing factor NOVA regulated synaptic proteins | WP:WP1983 | 3858_Glra2 | 5.19 |
| WP | Calcium regulation in cardiac cells | WP:WP553 | 3858_Glra2 | 3.40 |
| WP | Myometrial relaxation and contraction pathways | WP:WP385 | 3858_Glra2 | 2.58 |

1088

1089 **Figure S8. Gene Ontology (GO) enrichment analysis for vulnerable dopaminergic**

1090 **population, Related to Figure4.**

1091    Enrichment analysis of gene ontology (GO) terms for key genes in a representative vulnerable

1092    dopamine cluster (A, 3854_Grin2c; B, 3855_Arhgap28; C, 3857_Anxa1; D, 3858_Glra2).

# Figure S9.

**Figure S9. Whole-brain efferent and afferent connectivity of dopaminergic neuron**

**subtypes, Related to Figure5.**

**A.** Characterization of the DAT-Flp line using a reporter line (Ai65) confirmed that tdTomato+

cells are *Th+* neurons. Scale bars: 1mm for the striatum (left) and 400μm for the substantia

nigra (right). **B.** Representative images of axonal projections of *Vglut2+*, *Calb1+* and *Anxa1+*

dopamine neuron subtypes in the SNc. They preferentially project to dorsolateral striatum,

ventral medial striatum and dorsal striatum. Scale bar 1mm. **C.** Brain regions to which *Vglut2+*,

*Calb1+* and *Anxa1+* dopamine neuron project, measured as the fraction of neurites found within

those brain structures defined Allen brain atlas (n=4 *Calb1+* and n=3 for *Vglut2+* and *Anxa1+*).

Blue horizontal line indicates dorsal striatum and red line indicates ventral striatum. **D-G.**

Statistical analysis of the whole-brain distribution of ipsilateral (left) or contralateral (right)

monosynaptic inputs to *Vglut2+* neurons (D, E) or *Calb1+* dopamine neuron subtype (F, G) in

the SNc. Brain areas are color-coded by the Allen Brain Atlas.

## Figure S10.

**1108** **Figure S10. Direct afferent inputs from the motor cortex or striatum to the SNc DANs,**

**1109** **Related to Figure5.**

**A.** Schematic of the rabies injection procedure and experimental timeline for helper viruses and

rabies viruses in the DAT-Cre or Anxa1-Cre. **B.** Whole-brain 3D reconstruction using BrainJ,

color coded by brain regions (ref, left: sagittal view, right: coronal view). **C.** Representative

images of monosynaptic inputs to DANs from the whole brain (c1-c8, from the rostral to caudal).

The small white rectangle indicates the injection site, the SNc. Scale bar 1mm. **D-E.**

Monosynaptic projections from primary inputs include cells associated with motor function to the

SNc DANs (D, motor cortex; E, striatum). Scale bars: 1mm for the striatum and 400µm for the

substantia nigra.