Genetic analyses suggest that larval and adult stages of *Lucilia cuprina* employ different sensory systems

Abstract

The blowfly *Lucilia cuprina* is a destructive parasite of sheep that causes flystrike or myiasis. Larvae consume the animal's living flesh, producing large wounds that can lead to death. Growing resistance to conventional control methods has prompted the analysis of alternative strategies. RNA-Seq analysis of whole larvae at different stages and third instar head and gut tissues, suggested that odorant (OR), gustatory, ionotropic and *pickpocke*t receptors may not play a central role in the *L. cuprina* larval sensory signaling and digestive systems. Rather, ABC transporters were highly enriched in head and gut RNA, and OBPs only in the head. To confirm ORs are not essential for larval detection of rotten beef, diet-choice assays were performed including larvae and adults homozygous for a null mutation in the odorant coreceptor (*LcupOrco*) gene. While the attraction of adult females to rotten beef was fully disrupted, *LcupOrco* mutant larvae showed no change in diet preference.

Introduction

Immature stages of blowfly species (Diptera: Calliphoridae) display a wide spectrum of feeding specializations [1]. Based on their feeding habits, species within this family can be classified from obligate parasites of live animals to obligate necrophagous (dead tissue consumers) with many facultative forms in between [1–3]. Some of these species are considered livestock pests, such as *Lucilia cuprina* (Wiedemann, 1830), *Lucilia sericata* (Meigen, 1826) and *Cochliomyia hominivorax* Coquerel, 1858 [2,4]. *Lucilia cuprina* is a facultative ectoparasite and one of the main causative agents of flystrike in sheep in Oceania [5,6]. Conventional insecticides constitute the current control method for this pest [7], however, an increasing resistance to organophosphates and benzoylphenyl ureas [7], have motivated the development of alternative genetic strategies for population control such as the sterile insect technique (SIT) and female specific release of insects carrying a dominant female lethal gene (fsRIDL) [8–10].

Flystrikes on live animals are associated with the female oviposition behavior in blowfly parasitic species [2]. *Lucilia cuprina* females lay eggs in open wounds or moist tissues of host animals, clustering them in masses of ~200 eggs per ovarian cycle [11,12]. After egg hatching, the immature larval stages congregate in clusters of maggots called myiasis [12]. Within the myiasis the larvae complete the three larval (L) stages, from L1 to L3 [12]. This maggot arrangement

favors the action of the larval secretions and excretions (SE) improving larval feeding and

survival [13]. *Lucilia cuprina* larval SE includes several components such as gut microbiota,

lipases, peptidases and ammonia [14,15]. Peptidases and lipases together with the gut microbiota

facilitate host-tissue ingestion and digestion [15]. Ammonia, along with other SE, has been shown

to control deleterious fungi and bacteria that impair larval survival [16]. Maggot masses produce

heat, reaching up to 18.7°C above ambient temperature depending on the number of larvae [17].

Temperature is a key factor driving *L. cuprina* larval development and survival, and larvae are

constantly moving in and out of the maggot mass to avoid overheating [18]. The highest survival

rate for this species, considering the whole life cycle, was observed at 24±1°C [19,20].

Temperatures below and above this range can be deleterious to *L. cuprina* decreasing larval

survivorship [18].

The behavior of *L. cuprina* and other blowflies in locating hosts for oviposition has been

extensively studied [13]. The main attractants to gravid females are sulfur-rich compounds,

ammonia and other volatiles emitted from decomposing organic material, such as meat or feces

[1,13,21]. Dimethyl disulfide (DMDS), dimethyl trisulfide (DMTS), Indole and p-cresol are among the

most relevant blowfly attractants collected from decomposing organic material [22–24]. Adult gravid

females and the last larval stage of *L. cuprina* are more attracted to 5-day-old rotten beef

compared to fresh beef [25,26]. In addition, larvae preferred cold beef (~25°C) rather than warm

(~33°C) [26]. These observations could point to sensory systems conserved between the adult

and larval stages of *L. cuprina*.

RNA-Seq studies have been addressed different larval biological processes in blowfly

species, such as development [27], metabolism [28], aging [29], adaptation to temperature changes

[30,31] and immune response [32]. However, none of these studies focused on larval sensory

receptors potentially associated with physiological and behavioral processes. In *Drosophila*

melanogaster Meigen, 1830, different sensory receptors, including odorant (ORs) and gustatory

receptors (GRs), have been linked to many larval biological processes such as chemotaxis,

thermotaxis and locomotion [33,34]. For example, *DmelOR42a* and *DmelOR42b* appear to be

important for larval olfaction and locomotion [35–39]. Regarding GRs, such as *DmelGr64a*, have

been associated with taste [40], whereas others such as *DmelGR28* have been associated with

light avoidance and thermal sensing [41,42]. Ionotropic receptors (IRs) were also associated with

taste and thermal sensing in the *Drosophila* larvae [43,44].

Among other sensory receptors, the transient receptor potential channels (TRPs) and

Pickpocket receptors (PPKs) were identified in the *Drosophila* larval chordotonal organs and

showed a role in proprioception, thermotaxis and locomotion [45–48]. DmelTRPN1 (*nompC* gene)

and other TRPs belonging to the ankyrin and vanilloid subfamilies were associated with

proprioception [48] and thermotaxis [46,47,49]. Among odor-molecule carriers, odorant-binding

proteins (OBPs) and chemosensory proteins (CSPs) were identified in the larva of *D.*

melanogaster [50]. However, the role of these proteins in the larval olfaction and other

physiological aspects has not been extensively studied. Further, the role of other chemosensory

proteins previously studied in adult flies [51–53], such as the superfamily ML (MD-2 (myeloid

differentiation factor-2)-related Lipid-recognition and Niemann-Pick C2 disease proteins

(ML/NPC2), CD36-sensory neuron membrane proteins (CD36/SNMPs), and ammonium

transporters (AMTs), remains to be addressed.

The main objective of the present work was to identify sensory receptors and other

genes relevant to the physiology of *L. cuprina* larvae. We aimed to highlight genes potentially

associated with the processes mentioned above, and other physiological functions such as

feeding, digestion, excretion, diuresis and immune response. Our RNA-Seq analysis of different

larval stages and tissues suggested that the ORs, GRs, IRs and PPKs may not play a central

role in the larval sensory signaling and digestive systems of *L. cuprina*, except for *LcupGR94a*

which was biased to the larval gut. ABC transporters and OBPs were the most relevant within

the sensory-related gene families in terms of number of identified sequences and gene

expression, both in the larval head and across all larval stages. Since the odorant coreceptor

(*Orco*) is a necessary element for the normal functioning of all ORs [54], we tested both adult

females and larvae homozygous for a loss-of-function mutation created by gene editing.

Behavioral assays showed that *LcupOrco* was essential for the female response to rotten beef,

but not for larval behavior. These findings are consistent with high levels of expression of

LcupOrco in the adult female antenna [25] but very low expression in larvae (present work).

Results and Discussion

RNA-Seq overview

To identify genes that are differentially expressed at the very anterior end of the larvae, the first

segment (excluding the brain) was dissected from early feeding third instar (Supplementary Fig.

1). RNA was isolated from gut dissected from the same stage to identify genes important for

meat digestion. Lastly RNA was obtained from whole first, second and third instar (early and

late) for reference for the two tissues, and for a larval developmental series. RNA-Seq output

including reads per library, total number of sequences mapped and subtotals per type of RNA is

provided in Supplementary Table 1A-B. Considering only transcripts surpassing 5 TPMs, the

number of transcripts ranged between 5,617 and 6,642 for different larval stages and tissues

(Supplementary Data 1A-C and 2A-C). Further, a PCA analysis showed that samples belonging

to each larval stage clustered together, highlighting the differences between larval stages over

those between the samples within each group (Supplementary Fig. 2). An RNA-Seq analysis of

different larval tissues of *L. sericata* showed a similar number of transcripts expressed over the

same expression threshold, between ~ 4,400 and 6,000 [30]. Transcripts below this expression

threshold for each library are provided in Supplementary Data 1D-S and 2D-N.

Lucilia cuprina larval sensory receptors

ABC transporters and OBPs compromised about 80% of the sensory receptors/carriers showing

a TPM expression ≥ 5, and the number of receptors/carriers per gene family was as follows: 49

ABCs; 25 OBPs; eleven CD36/SNMPs; six ML/NPC2s; three CSPs and one for each of the

following gene families, AMTs, GRs and TRPs (Supplementary Data 3 and 4). Furthermore,

OR, GR, IR and PPK receptors were not observed above the expression threshold of 5 TPM,

except for *LcupGR94a* in the larval gut (Supplementary Data 4A). The expression level of ORs,

GRs and IRs suggests that the same may not have a central role in the sensory system of *L.*

cuprina larvae. For instance, *LcupOrco,* a highly expressed sensory receptors in the *L. cuprina*

antenna [25] showed a TPM expression < 0.5 in whole larvae and in head and gut (Table 1).

However, in *D. melanogaster* larvae, *Orco* expression is restricted to the olfactory dorsal organs

at the anterior tip [55] and is essential for chemotaxis [56]. Furthermore, the role of many ORs, GRs

and IRs has been confirmed by *in situ* hybridization and functional assays [33–36,41–44].

Consequently, more experiments are required to elucidate the role of the specified receptors

within the sensory system of the *L. cuprina* larvae.

Table 1. *Lucilia cuprina Orco* expression in adult and larval tissues

Tissue	TPM expression	Cite
Whole larva stage 1	0.48	This study
Whole larva stage 2	0.15	This study
Whole larva early stage 3	0.02	This study
Whole larva late stage 3	0.02	This study
First segment of larva early stage 3	0.38	This study
Gut of larva early stage 3	0.003	This study
Antenna of gravid female 10-d-old	1074	Wulff *et al*. 2024
Antenna of mated male 10-d-old	924	Wulff *et al*. 2024

The ABC transporters are classified into eight subfamilies [57]. A phylogenetic analysis

showed a possible clade expansion for the subfamily G in *L. cuprina* (Fig. 1 and Supplementary

Fig. 3). Similar expansions for the same subfamily were observed in hematophagous dipterans

[58]. In *L. sericata*, orthologs of LcupABCGs 4, 6 and 7 were highly expressed in larval Malpighian

tubules (MT) and fat body (FB) [30]. Here we found that members of this subfamily are highly

enriched in the larval anterior end (see below). In contrast to the apparent expansion of

subfamily G, subfamily A showed a possible clade contraction compared to *D. melanogaster*

(Fig. 1), with only four receptors identified for *L. cuprina* (Supplementary Data 5). In line with

these findings, clade contractions and expansions are common for subfamilies A and G along

insect species [59]. Sequences used in the analysis are provided in Supplementary Data 5A-D;

annotation of two sequences was corrected and transcript expression and differential

expression calculated using genome assembly GCA_001187945.1 (Supplementary Data 6).

Figure 1. **Phylogenetic analysis of the ABC transporters**. Three Diptera species were used for the phylogenetic

analysis as follows: *L. cuprina* (Lcup, gold), *L. sericata* (Lser, green) and *D. melanogaster* (Dmel, black). ABC

subfamilies from A to H and node bootstrap supporting values were detailed. Only *L. cuprina* receptors with a TPM

expression ≥ 5 in larval head, gut or whole body were used for the analysis, but all ABC transporters of D.

melanogaster were included. In addition, ABC transporters previously identified as larval biased in *L. sericata* 28 were

added to the analysis. If there was more than one isoform per gene, only isoforms producing a different amino acid

sequence and showing a dissimilar transcript expression were added. Potential expanded *L. cuprina* ABC

transporters belonging to subfamily G were detailed with an asterisk. For more details about sequences used in the

analysis refer to Supplementary Data 5.

Lcuppixie was the most highly expressed ABC transporter (Supplementary Data 3A and

4B) and upregulated in the first two instar compared with third (Fig. 2A). This gene belongs to

subfamily E and is essential for protein translation [60]. Receptors belonging to subfamilies F,

associated with protein translation and immune response [61], and B, associated with xenobiotic

detoxification and insecticide resistance [57], were also upregulated in the first two instar larva vs.

L3 (Fig. 2A, Supplementary Data 3A). No member of these subfamilies was head-biased, but

five ABCs of subfamily B were gut-biased (Fig. 3A-C, Supplementary Data 4B). Subfamily G

showed higher expression in the latter instars (L2 or L2/L3) compared with the first instar (Fig.

2B-D). In addition, this subfamily had the highest number of head-biased genes (Fig. 3A and

Supplementary Data 4B), with some showing a very high enrichment compared with gut or

whole-body (Supplementary Data 7A). This subfamily includes orthologs of the well-studied

Drosophila pigmentation genes *white* and *scarlet* [57], which are L2-biased in *L. cuprina* (Fig. 2B)

but do not show any enrichment in head or gut (Fig. 3 A-B).

Figure 2. **Expression and differential expression of the ABC transporters in *L. cuprina* larval stages**. The TPM expression of ABC receptors was plotted clustering the same according to differential expression between the three larval stages as follows: (A) includes receptors biased to the first two larval stages (L1 and L2) vs. L3; (B) shows receptors only biased to the L2 stage; (C) groups receptors biased to the last two stages (L2 and L3) vs. L1; and (D) the receptors only biased to the last stage (L3). Only receptors with a TPM expression ≥ 5 were added. Dots represent the gene TPM expression of each library; the average mean expression per gene plus standard deviation (SD) between libraries were also plotted. For more details about the expression level and differential expression of each receptor refer to Supplementary Data 1 and 3A.

Figure 3. **Expression and differential expression of the ABC transporters in *L. cuprina* third instar larval tissues**. The TPM expression of ABC receptors was plotted clustering the same according to differential expression between different tissues as follows: (A) includes receptors biased to the larval head (first segment of the larval body) vs. whole-body; (B) shows receptors biased to the gut vs. whole-body; and (C) groups receptors biased to head plus the gut vs. whole-body. Only receptors with a TPM expression ≥ 5 were added. Dots represent the gene TPM expression of each library; the average mean expression per gene plus standard deviation (SD) between libraries were also plotted. Symbols next to receptor's names represent larval stage biased expression (plotted in Fig. 2) as follows: # L1/L2-biased; $ L2/L3-biased; & L3-biased. For more details about the expression and differential expression of each receptor refer to Supplementary Data 2 and 4B.

Subfamily H was also well represented in the larval head, with the ortholog of

Lcuposkyddad (*osy*) showing head-biased expression (Fig. 3A) and the ortholog of

Lcupsnustorr (*snu*) enriched in head and gut (Fig. 3C). In *D. melanogaster*, *osy* and *snu* are

associated with lipid deposition on the cuticle [62,63]. In the larval gut, the best represented

subfamily was C followed by B (Fig. 3B). Both subfamilies share a role in xenobiotics

detoxification, multidrug resistance and protein translation [57]. ABC-Cs are also involved in the

binding of sulfur-rich compounds [64], and ABC-Bs in ion and lipid transportation [57] and iron/heme

metabolism in hematophagous insects [65]. Sulfur-rich compounds are common blowfly

attractants [13] and iron and heme are both present in the *L. cuprina* larval meat diet [66]. Other

genes engaged in processing these compounds, such as *heme oxygenase 1* and *ferritin* were

also highly expressed and biased to the larval gut (Supplementary Data 2B and 8A-B).

The second highest number of chemosensory proteins identified in all *L. cuprina* larval

stages and tissue transcriptomes were grouped by OBPs (Supplementary Data 3B and 4C).

This gene family is classified based on a cysteine (C) conserved motif, and modifications of the

same change the OBP-ligand binding affinity [54]. The classic subfamily includes a six C-motif and

in the Dimer subfamily, the whole motif has been duplicated [54]. Other subfamilies are Minus-C,

where the C2 and 5 from the classic motif have been lost, and Plus-C, that includes the classic

motif plus three extra conserved Cs and a conserved proline immediately after the Cs [54]. The

OBP phylogenetic analysis suggested a clade expansion for *L. cuprina* and *L. sericata* classic

OBPs 39 to 44 (Fig. 4). The *L. sericata* OBPs included in the expanded clade were highly

expressed in the FB followed by the MT [30], suggesting they may be associated with lipid

binding. OBPs grouped within the clade LcupOBPs 39-44 and 49, were the highest expressed

genes among the L3 biased (Fig. 5C-E and Supplementary Data 3B), and head-biased (Fig. 6,

Supplementary Data 4C and 8C-D). Sequences used for the analysis are provided in

Supplementary Data 9A-C.

291
292
Figure 4. **Phylogenetic analysis of the OBPs**. Three Diptera species were used for the phylogenetic analysis as
follows: *L. cuprina* (Lcup, gold), *L. sericata* (Lser, green), *D. melanogaster* (Dmel, black). OBP subfamilies classified
according to the cysteine motif (Classic, Minus-C, Plus-C and Dimer), and node bootstrap supporting values were
detailed. Only *L. cuprina* OBPs with a TPM expression ≥ 5 in larval head, gut or whole body were used for the
analysis, but all *D. melanogaster* OBPs were included. In addition, OBPs previously identified as larval biased in *L.*
sericata 28 were added to the analysis. If there was more than one isoform per gene, only isoforms producing a
different amino acid sequence and showing a dissimilar transcript expression were added. Potential expanded *L.*
cuprina OBPs 39 to 44 belonging to subfamily Classic were detailed with an asterisk. For more details about
sequences used in the analysis refer to Supplementary Data 9.

Figure 5. **Expression and differential expression of the OBPs across *L. cuprina* larval stages**. The TPM expression of OBPs was plotted clustering the same according to differential expression between the three larval stages as follows: (A) includes OBPs biased to the first two larval stages (L1 and L2) vs. L3; (B) shows OBPs only biased to the L2 stage; (C) groups OBPs biased to the last two stages (L2 and L3) vs. L1; (D) OBPs only biased to the last stage (L3); and (E) OBPs biased to the first and last stage (L1 and L3). Only receptors with a TPM expression ≥ 5 were added. Dots represent the gene TPM expression of each library; the average mean expression per gene plus standard deviation (SD) between libraries were also plotted. For more details about the expression and differential expression of each receptor refer to Supplementary Data 1 and 3B.

Figure 6. **Expression and differential expression of larval head-biased OBPs**. Only the TPM expression of head-biased OBPs was plotted. Dots represent the gene TPM expression of each library; the average mean expression per gene plus standard deviation (SD) between libraries were also plotted. Only receptors with a TPM expression ≥ 5 were added. Symbols next to receptor's names represent larval stage biased expression (plotted in Fig. 5) as follows: # L1/L2-biased; @ L2-biased; $ L2/L3-biased; & L3-biased; Ψ L1/L3-biased. For more details about the expression and differential expression of each receptor refer to Supplementary Data 2 and 4C.

In insects, OBPs are also involved in lipid deposition on the cuticle [67], and some OBPs expressed at the MT and cuticle were associated with xenobiotics detoxification [68–70]. Based on their expression in the *L. cuprina* larvae, OBPs with a lipid biding affinity may be involved in energy homeostasis, avoiding larval desiccation and assisting detoxification. Remarkably, ABC transporters with similar functions, such as receptors from subfamilies G and H, shared a common expression pattern in the *L. cuprina* larvae. Part of the Minus-C and most of the Plus-C OBPs were L1/L2 biased vs. L3 (Fig. 5A-B and Supplementary Data 3B). The Minus-C *LcupOBP50* and Plus-C *LcupOBP52* were among the most highly expressed sequences within these groups and biased to L1/L2 (Fig. 5A-B and Supplementary Data 3B). Orthologs of these receptors, namely DmelOBPs 44a and 49a respectively, were studied in *D. melanogaster*. *DmelOBP44a* is involved in lipid binding [71] and *DmelOBP49a* associated with taste in the same species [72]. Minus-C and Plus-C OBPs have been associated with binding of sesquiterpenes and derived sesquiterpenoids such as Juvenile Hormone (JH) in other insects [73–75], and they could have a similar role in *L. cuprina*. Only a few OBPs were expressed in the larval gut, and none of

them biased to the same tissue (Supplementary Data 4C). The Minus-C *LcupOBP22* and

LcupOBP24 were among the highest expressed in the gut (Supplementary Data 4C). In insect,

JH regulates the function of the adult and larval gut [76,77], and in *D. melanogaster* this hormone is

synthesized in the *corpora allata* but also in the adult gut [76]. Consequently, the above OBPs

could be involved in the JH-gut signaling system.

Lcuppainless was the only TRP above a 5 TPM expression threshold and biased to the

larval head (Supplementary Data 3C and 4D). In *D. melanogaster* this gene has been

associated with taste, thermal sensing and proprioception [78–81]. Accurate temperature detection

is indispensable to ensuring *L. cuprina* larval survival [17,19], and this receptor may be involved in

thermal sensing. Of the CD36/SNMP genes, the scavenger receptor class B1 (SR-BI,

XP_023294991.1) was upregulated in the L2/3 stages when compared to L1 (Supplementary

Data 3D), and the only member of this family showing a head bias (Supplementary Data 4E).

Interestingly this gene was highly expressed in the *L. cuprina* adult antenna [25], indicating a

possible role in larval and adult olfaction. Other members of the same family, *e.g. protein*

croquemort-like were biased to the larval gut (Supplementary Data 4E). In *D. melanogaster*

members of the CD36/SNMP gene family were associated with lipid binding and their

expression in the gut of the same species [82] suggest a role in feeding and metabolism.

ML/NPC2 transporters have been associated with lipid binding, immune response and

olfaction in different insect species [25,83–85]. Members of this family were highly expressed in the

L. cuprina L2 stage and the larval gut (Supplementary Data 3E and 4F). The *L. cuprina*

ecdysteroid-regulated 16 kDa receptor (*LcupESR16*) was the only head-biased gene within the

family (Supplementary Data 4F). The same gene was the highest expressed in the *L. cuprina*

adult antenna among this family [25]. Further, *ESR16* orthologs were found expressed in the

antenna of many insect species [25,84,85] suggesting a possible role in olfaction.

Three CSPs were expressed in *L. cuprina* larvae, and two of them were biased to the

head (Supplementary Data 3F and 4G). In *Bradysia odoriphaga* Winnertz (Diptera: Sciaridae), a

CSP protein was associated with sulfur-rich compounds detection [86], a common blowfly

attractant [13]. It is possible that CSP proteins may have a role in detection of sulfur-rich

compounds in *L. cuprina* larvae. The *ammonium transporter Rhesus type B* (*RHBG*) was biased

to the *L. cuprina* L2 stage and the larval gut (Supplementary Data 3G and 4H). This receptor is

involved in ammonia detection [52,87] and excretion at the MT [88] in *D. melanogaster*. Ammonia-rich

compounds are attractants to adult females of *L. cuprina* and *L. sericata* [13]. In adult antennae,

LcupRHBG was highly expressed in both sexes with a female-bias [25]. In the larva of *L. sericata*,

LserRHBG showed the highest expression in SG followed by MT [30]. Consequently, *LcupRHBG*

might be related to olfaction of adult flies, but involved in other processes in *L.cuprina* larvae,

such as feeding and excretion.

Odorant receptors are not essential for *L. cuprina* larval response to rotten beef

To determine the importance of ORs in larval behavior, CRISPR/Cas9 was used to obtain a null

mutation of the *LcupOrco* gene following previous protocols used in blowflies [89,90]. We identified

a deletion mutation that introduced a premature stop codon at the end of the first exon, 265

nucleotides from the start codon (Fig. 7 and Supplementary Figs. 4-12). Without the presence of

the odorant coreceptor (*Orco*), no odorant receptor can complete its olfactory function [54]. Larvae

that were a mix of wild-type (*wt*), *LcupOrco heterozygotes* and *homozygotes*, were tested for

diet preference in a choice assay. Overall, there was no major change in preference, with larvae

preferring rotten beef that was at room temperature as previously reported [26] (Fig. 8 A). No

correlation was found between diet preferences and larval sizes, either for the genotyped (Fig.

8B) or non-genotyped (Fig. 8C) larvae, ANOVA, P = 0.22 and P = 0.25 respectively. This lack of

correlation between genotype and diet preference was confirmed by a Chi-Square analysis, P =

0.66 (Fig. 8D and Supplementary Table 2A). In the blowfly *C. hominivorax, ChomOrco*

expression was minimal compared to adult flies showing the higher expression at the L1 stage

[90], as observed here for *L. cuprina* larvae (Table 1). The authors were able to rear a *ChomOrco*

silenced strain indicating that this gene may not be essential for *C. hominivorax* to complete the

larval cycle. Our results suggest that *LcupOrco* is not necessary for *L. cuprina* larvae to respond

to the complex mixture of odors emitted by decomposing beef, in contrast to observations in *D.*

melanogaster larvae, where *Orco* is essential for the response to odors [33–36].

Figure 7. CRISPR/Cas9-mediated loss-of-function mutation of the *L. cuprina* odorant coreceptor (*LcupOrco*) gene. A deletion mutation in the first exon resulted in the introduction of a premature stop codon at the end of the first exon. (A) *LcupOrco* gene detailing exons (E) and introns (I) with lengths in base pairs (bp), the guide RNA target site, and PAM sequence (underlined); (B) alignment between the *wt* and mutated *LcupOrco* DNA sequences, detailing deletions and insertions, including a 8 bp gap that introduces a premature stop codon at 265 bp from the start codon, producing a 73 amino acids (aa) truncated protein of 478 aa for the *wt* protein; and (C) LcupORCO protein sequence detailing the premature stop codon position located after the glutamic acid (E) 73, the seven transmembrane (TM) regions and cytoplasmic and external loops. The interaction between Orco protein and any odorant receptor to produce a functional dimer occurs at the Orco C-term region (-COOH in the figure), which is absent in the mutated version of LcupOrco protein. For more details about *LcupOrco* silencing refer to Supplementary Figs. 4-12 and Supplementary Notes 1 and 2.

Figure 8. Odorant receptors are not essential for *L. cuprina* larval response to rotten beef. A larval choice
assay was conducted to evaluate the larval response to fresh (F) and rotten (R) beef at two temperatures, cold
25±1°C (C) and hot 33±1°C (H). A total of 120 early L3 (4 days after oviposition) were tested, and 48 larvae randomly
collected from individuals attracted to each diet preference were genotyped. Larvae were obtained from crossing
LcupOrco heterozygous flies giving an approximate 3:1 ratio of *LcupOrco* non-silenced (wild-type plus heterozygous,
WT + HET) and silenced larvae for *LcupOrco* (homozygous, HOM). (A) the numbers on top of the bars represent total
larvae attracted to each diet preference, *i.e.* rotten cold (RC), rotten hot (RH), fresh cold (FC), fresh hot (FH) and non-
choice (NC); (B-C) show the weight of larvae in milligrams (mg) attracted to each diet preference, either for
genotyped (B) or non-genotyped (C) larvae. Dots represent single larvae, numbers on top of each preference or
genotype mean total number of larvae, and equal letters indicate no statistically significant difference (ANOVA, P =
0.22 for B and P = 0.25 for C); (D) shows how genotypes were randomly associated to each diet preference (Chi-
Square analysis, P = 0.66). Numbers on top of the bars represent total larvae per diet preference. For more details
about results obtained in this assay refer to Supplementary Figs. 6A, 7A and Supplementary Table 2A.

Disruption of *LcupOrco* affected the *L. cuprina* female oviposition behavior with egg
laying significantly delayed compared to heterozygous siblings (Supplementary Figure 8A-G).
This was not due to a feeding disorder as *LcupOrco*[-/-] homozygotes showed a fully developed

ovary (Supplementary Figure 9A-O). Further, a mating assay using transgenic males expressing

a green fluorescent protein, showed that only *LcupOrco* heterozygous females laid eggs the first

week after crossing (Supplementary Fig. 8A). After a second week, *LcupOrco* silenced females

(homozygous) started to lay non-fertilized eggs (Supplementary Fig. 8B-C), which did not

produce any offspring. *LcupOrco* silenced females needed a third week to start to lay fertilized

eggs giving healthy progeny (Supplementary Fig. 8D-G). Notably, *LcupOrco* silenced males

showed an average fertility of 88% including 30 males tested across four generations. These

results were in line with previous observations in *Harpegnathos saltator* T. C. Jerdon, 1851

(Hymenoptera: Formicidae), where silencing of *Orco* produced males with a normal mating

behavior and females with a delayed and decreased oviposition [91]. However, silencing of *Orco*

in *Helicoverpa armigera* Hübner, 1808 (Lepidoptera: Noctuidae), produced the opposite

phenotype, *i.e.* sterile males and fertile females [92], indicating that the function of *Orco* may

change among species.

We next asked if *LcupOrco* is important for detecting rotten beef in adult females using a

spatial olfactometer. A mix of *wt*, *LcupOrco* heterozygotes and homozygotes was obtained from

crossing *LcupOrco* heterozygous flies. This mix included an approximate 3:1 ratio of *LcupOrco*

non-silenced to silenced females. Also, a different group of *L. cuprina wt* females were used for

a further comparison with *LcupOrco* females. As previously reported [25], *wt* females were more

attracted to the rotten beef vs. fresh beef, or did not respond to any of the stimulus (non-choice,

NC), ANOVA, P = 0.0001 (Fig. 9A). However, for the *LcupOrco* mix, the number of non-

responding (NC) females was higher than any of other two preferences, ANOVA, P = 0.0001

(Fig. 9B). Genotyping of randomly selected females from tested groups, showed that *LcupOrco*

silenced females were overrepresented among the non-responding flies (NC), Chi-Square, P =

0.0001 (Fig. 9C and Supplementary Table 2B). These observations showed that *LcupOrco* is

essential for odor detection in adult females, and our results are consistent with previous

observations in *C. hominivorax*, where disruption of *ChomOrco* impaired the foraging and host-

seeking behaviors of adult flies [90].

Figure 9. Disruption of *LcupOrco* impairs *L. cuprina* adult female olfaction. Females were obtained from crossing *LcupOrco* heterozygous flies giving an approximate 3:1 ratio of *LcupOrco* non-silenced (wild-type plus heterozygous, WT + HET) and silenced females for *LcupOrco* (homozygous, HOM). A different group of L. cuprina *wt* females were used for a further comparison with *LcupOrco* females. All flies used for the olfaction assay were ~ 10-day-old gravid females and were separated from males the day before the assay using CO2 to anesthetize the flies. The attractants used to lure the flies were 2 grams of fresh (F) and rotten (R) beef at room temperature. A total of 80 *wt* and 174 *LcupOrco* females were tested, and 72 of the latter were genotyped. (A) the *wt* females were more attracted to the rotten beef vs. fresh beef, or did not respond to any of the stimulus (NC), ANOVA, P = 0.0001; (B) in females belonging to the *LcupOrco* mix, the number of non-responding (NC) females was higher than any of other two preferences, ANOVA, P = 0.0001; (C) genotyping of randomly selected females from tested groups, showed that *LcupOrco* silenced females were overrepresented among the non-responding flies (NC), Chi-Square, P = 0.0001. Different letters indicate statistically significant difference. For more details about results obtained in this assay refer to Supplementary Figs. 6B, 7B and Supplementary Table 2B.

Non-sensory genes expressed and biased to the larval head

Transcripts encoding the *L. cuprina* orthologs of the *D. melanogaster yellow*, *yellow-h* and

ebony genes, with roles in cuticle pigmentation and hardening (tanning) [93–95], were among the

highly expressed plus head-biased genes vs. the whole-body (Supplementary Data 7A, 8C-D).

The *L. cuprina eclosion hormone* gene (*LcupEH*) was also enriched in the larval head in

comparison to the whole-body (Supplementary Data 7A). In insects, *EH* is associated with

cuticle tanning through the control of *Bursicon* (*bur*) and *Partner of Bursicon* (*pbur*) [96]. The

presence of these transcripts suggests a role in cuticle and/or mouthparts tanning of the *L.*

cuprina larvae. In *D. melanogaster*, *yellow* controls the tanning of the larval mouth parts [97].

Remarkably, the *tan* gene which display an antagonistic role to *ebony* in cuticle tanning in *D.*

melanogaster [93,98] was downregulated in the head vs. the whole-body (Supplementary Data

8D).

Transcripts encoding for mucins such as *Lcupmucin-5AC* were highly expressed in the

head compared to whole-body, and head-biased (Supplementary Data 7A and 8C). Mucins

have been studied in the insect gut, playing a role in food processing and supporting the

immune system [99]. Enrichment of mucin transcripts in the larval head could suggest a role at the

anterior part of the digestive system. We previously found that members of this gene family

were highly expressed in the *L. cuprina* adult antenna [25], and suggested that mucins may be

important for olfaction, a role confirmed in vertebrates [100] but not yet addressed in insects. It is

possible that mucins may also be important for olfaction in *L. cuprina* larvae.

Genes potentially associated with *L. cuprina* larva feeding and digestion

Proteases and lipases such as *trypsin delta* and *lipase 3* respectively, were among the genes

highly enriched in the larval gut compared to whole-body and biased to the larval gut of *L.*

cuprina (Supplementary Data 7B and 8A-B). These findings are consistent with a larval diet

composed mainly of meat with a small fat content [66]. Neuropeptides *Allastostanine*, *CCHamide-*

2 and *long neuropeptide F* (*NPF*), that modulate food intake and digestion in *D. melanogaster*

[101,102], were also expressed and biased to the larval gut (Supplementary Data 7B and 8A-B).

Neuropeptide *Diuretic hormone class 2* (*DH31*), which play a role in diuresis, gut contractions

and elimination of virulent bacteria in *D. melanogaster* [103] showed the same expression pattern

as the above neuropeptides (Supplementary Data 7B and 8A-B). In addition, transcripts

encoding for mucins and cytochrome P450s (CYPs) were overrepresented in the larval gut

(Supplementary Data 7B and 8A-B). CYPs play a crucial role in the insect gut metabolizing

endogenous and exogenous substances [104], and mucins in the aggregation of harmful elements

[99]. These enzymes may be involved together in detoxifying harmful compounds and

metabolizing dietary components in the gut of *L. cuprina* larva.

Gene expression and differential expression across larval stages

The most highly expressed sequences in the whole-larva of any stage were genes associated

with cuticle (*e.g. larval cuticle-protein 65Ag1-like, Lcuplcp65Ag1*), muscle formation (*e.g.*

myosin-regulatory light chain 2), and RNA translation factors (Supplementary Data 1A-C).

These transcripts were also abundant in larval head, but with an overrepresentation of

sequences coding for cuticle proteins and sensory receptors (Supplementary Data 2A). The

larval cuticle is essential for larval life, acting as both skin and exoskeleton, with the muscles

linked directly to the same [105], and the interaction between cuticle and muscles allows larval

locomotion [105]. The cuticle also protects the larva from desiccation and abrasion, includes

sensory and feeding organs and constitutes the first barrier of the immune system [106].

Genes associated with cuticle formation, *e.g. Lcuplcp65Ag1*, showed the highest fold-

change (FC) increases in L2 vs. L1 (Supplementary Data 10A). Other genes potentially

associated with cuticle tanning such as the orthologs of the *D. melanogaster yellow, yellow-h*

and *EH* genes were also upregulated in L2 vs. L1 (Supplementary Data 10A). An ortholog of the

Adipokinetic hormone (*AKH*), which is involved in mobilizing lipids from storage tissues in

different insects [107], showed a high FC increase in L2 vs. L1 (Supplementary Data 10A). Other

genes involved in protein and fat digestion, such as *Lcuptrypsin-1* and *Lcuplipase 3*

respectively, were upregulated in the L2 stage (Supplementary Data 10A). Interestingly,

orthologs of the antimicrobial peptides *Lucifensin* [108] and *defense protein-l(2)34Fc* (*l(2)34Fc*) [109],

were also increased in L2 vs. L1 (Supplementary Data 10A). Larvae are reared on fresh meat

under non-sterile conditions [9] and thus could be exposed to a higher bacterial load at later

stages. On the contrary, the expression of orthologs of feeding modulators such as *short*

neuropeptide F (*sNPF*) and *SIFamide* [110,111] was downregulated in L2 vs. L1 (Supplementary

Data 10A). In *D. melanogaster* larvae, *sNPF* increases food intake but does not prolong the

feeding period or modulate food preferences, unlike *NPF* [111]. Both, *NPF* and *sNPF* are major

modulators of feeding and antimicrobial immune response [112,113].

In the late L3, arylphorin genes showed the highest FC increases compared to previous

larval stages (Supplementary Data 10B-C). In insects, these proteins have a role in storage and

cuticle formation during the prepupal period [114]. Other genes associated with cuticle formation,

such as *Lcuplcp65Ag1*, were also upregulated in L3 (Supplementary Data 10 B-C). Genes

potentially involved in larval energy homeostasis such as *LcupAKH* and *LcupCCHAmide-2*, also

showed a biased expression to L3 (Supplementary Data 10B-C). In contrast, genes potentially

involved in cuticle tanning of *L. cuprina* larva such as *Lcupyellow* and *LcupEH*, and orthologs of

feeding modulators such as *sNPF*, *NPF* and *SIFamide* were downregulated in L3 vs. L1/L2

(Supplementary Data 10B-C). In addition, orthologs of the antimicrobial peptides *Lucifensin* and

l(2)34Fc, and chymotrypsin inhibitors associated with protease inactivation [115] were also

downregulated in L3 vs. L1/L2 (Supplementary Data 10B-C).

Materials and Methods

Insect rearing conditions

All larvae and flies used in the present study were of the sub-species *Lucilia cuprina cuprina*,

called "*L. cuprina*" to ease the reading of the manuscript. The *L. cuprina* LA07 wild-type (*wt*)

colony is currently at approximately 225 generations and was established in 2010 using 300

individuals kindly provided by Dr. Aaron Tarone (Texas A&M, TX, USA). The LA07 colony was

established by Dr. Tarone from multiple collections of individuals (300 to 500) from the

University of Southern California campus and the Miracle Mile neighborhood, in Los Angeles,

CA, USA in 2007. All larvae and adult flies used for the experiments were reared using

protocols previously described [9]. Adult flies and immature stages were kept at 23.5±1°C a non-

controlled photoperiod (~13:11 light/dark).

To complete the RNA-Seq analysis, *wt* and heterozygous individuals were used, the

latter obtained from crossing *wt* and SLAM5X colonies. SLAM5X carries an X-linked

constitutively expressed DsRed fluorescent protein gene plus the tetracycline transactivator

coding sequence driven by the early embryo *Chslam* promoter [116]. For the behavioral assays,

LcupOrco larvae or adult females were obtained from crossing heterozygous *LcupOrco*

(*LcupOrco$^{+/-}$*) siblings. The resulting *LcupOrco* mixture was ~ 25% wt, 50% *LcupOrco$^{+/-}$* and 25%

LcupOrco$^{-/-}$. In addition, adult males from the *NPF$^{-/-}$* colony expressing the ZsGreen were used

for the mating assay.

RNA-Seq experiments

Experimental design and sample collection

Two RNA-Seq experiments, designated EXP-1 and EXP-2 were performed. EXP-1 addressed

gene expression pattern across the three larval development stages using RNA isolated from

whole first, second and late wandering third instar (L1, L2, L3). EXP-2 used RNA isolated from

head, gut and whole-body early feeding third instar. For EXP-1 there were five replicates for L1

and L2 and six replicates for L3. Samples one to three of L1 and L2, and all samples for L3,

were *wt* larvae. Samples four and five of L1 and L2 were collected from the *wt*-SLAM5X cross.

Each replicate was a pool including 10 to 15 larvae for L1 and L2, and three pooled individuals

631 for L3. L1 samples were collected between 20 to 30 hours (h) after egg laying, 50 to 55 h for L2

632 and between 130 to 135 h for L3 (Supplementary Fig. 1A-D). All samples were flash frozen

633 using liquid nitrogen after collection and stored at -80 °C until use.

634 Samples included in EXP-2 were the first segment of the larva designated as head (H), the gut

635 including the midgut plus the hindgut and without the crop and Malpighian tubules designated

636 as gut (G), and the whole-larva (WL) including all tissues (Supplementary Fig. 1E-G). The crop

637 was removed from the gut because the presence of meat from the larval diet withing this organ

638 could bias the analysis. All tissues were collected from *wt* L3s of ~ 75 h after egg laying. Four

639 replicates were collected for H and G and three for WL, and each replicate was a pool including

640 50 first larval segments for H, 10 guts for G and five whole larvae per pool for WL. Tissues were

641 dissected using dissection forceps and vannas scissors (WPI, Worcester, MA, USA), and

642 immediately after placed in a 1.5 mL Eppendorf tube filled with 300 µl of cold RNAlater™

643 (Thermo-Fisher, Waltham, MA, USA, Cat. # AM7020) and stored at -80°C until use.

645 **RNA isolation**

646 RNA extraction of samples associated with EXP-1 and EXP-2 was completed using two

647 different protocols as follows. Whole-larvae of EXP-1 were transferred using dissection forceps

648 to 2.0 mL tubes prefilled with zirconium beads (Benchmark Scientific, Tempe, AZ, USA, Cat. #

649 D1032-30). Subsequently, the larvae were resuspended in 250 µl of cold Trizol™ (Thermo-

650 Fisher, Cat. # 15596026) and disrupted using a benchtop homogenizer (Benchmark Scientific,

651 Cat. # Z742475). The homogenizer was set to 6.5 m/s and samples were disrupted for 1 min,

652 another 250 µl of cold Trizol™ were added to the tubes and the homogenization process was

653 repeated once more. After homogenization, total RNA was extracted using the RNeasy Kit

654 (Qiagen, Hilden, Germany, Cat. # 74104), and genomic DNA (gDNA) was removed using the

655 DNase kit (Qiagen, Cat. # 79254) following manufacturer's specifications.

656   Before RNA extraction of samples belonging to EXP-2, the RNAlater™ was removed

657 from tubes and tissues were rinsed two times with 1 mL of cold 50% ethanol. After rinsing, pools

658 of heads, guts and whole larvae were resuspended in 250, 500 and 1000 µl of cold Trizol™

659 respectively. Samples were kept on ice and tissues were disrupted using pellet pestles (Bel-Art,

660 Wayne, NJ, USA, Cat. # F65000-0002) with a homogenizer (Bel-Art, Cat. # F65100-0000) for 90

661 seconds (sec). Subsequently, samples were centrifuged at 16000 x g and 4°C for 5 min to pellet

662 part of lipids, cuticle and other debris. After centrifugation the supernatant was transferred to a

663 clean 1.5 mL tube. Total RNA was extracted using the Quick-RNA™ kit (Zymo Research, Irvine,

CA, USA, Cat. # R1050) according to the manufacturer's specifications. The exception was the

addition of a second deoxyribonuclease 1 (DNase 1) step to ensure the absence of gDNA

contamination in the samples, as follows: 1) during the RNA extraction 30 U of DNAse1 plus 75

µl of digestion buffer (DB) were added to the crude samples followed by 15 min of incubation at

RT; 2) after elution of the clean RNA in nuclease-free water (NF-water), another 5 U of DNAse1

+ 2.5 µl of DB were added followed by an incubation time same as above. After DNAse1

treatments, the RNA Clean & Concentrator™-5 kit (Zymo Research, Cat. # R1013), was used to

clean the samples following the manufacturer's specifications.

Clean RNA belonging to samples from EXP-1 and EXP-2 was eluted in 30 and 12 µl of

NF-water, respectively, and was quantified (1:10 dilution) using a Qubit-4™ fluorometer and the

HS kit (Thermo-Fisher, Cat. # Q33120). Samples were kept at -80°C until use.

RNA-Seq data analysis

Sample quality control, library construction and sequencing services were provided by

Novogene Inc. (Sacramento, CA, USA), following protocols described by the provider

(https://www.novogene.com/us-en/resources/downloads/). The sequencing platform was

Illumina NovaSeq 6000 (Illumina, San Diego, CA, USA), using paired end 150 bp reads, and the

sequencing depth coverage was ~ 50 and 400 million raw reads per library for EXP-1 and EXP-

2, respectively.

The Geneious Prime® software v2023.2.1 (https://www.geneious.com) and associated

plugin packages, were used to complete all the bioinformatic analysis and generate the RNA-

Seq data output. BBDuk plugin v38.88 was utilized to remove Illumina adaptors and trim off low-

quality bases at the 5′ and 3′ends using the following parameters: kmer length: 27; trim both

ends: minimum quality Q20; trim adapters based on paired reads overhangs: minimum overlap

22; discard short reads: minimum length 20 bp. Trimmed reads were mapped to the reference

L. cuprina genome assembly NCBI ID ASM2204524v1, using the default Geneious Prime

mapper set to detect all types of RNA sequences and low-medium sensitivity. A second *L.*

cuprina genome assembly, NCBI ID GCA_001187945.1, was used to analyze specific genes

with annotation errors in the first assembly. Gene expression was calculated only for identified

coding sequence (CDS) and the normalization method used was transcript per million (TPM),

but RPKMs (Reads Per Kilobase per Million mapped reads) FPKMs (Fragments Per Kilobase

per Million mapped fragments) were also provided with the raw data.

A Principal Components Analysis (PCA) was performed to compare EXP-1 libraries with

the R packages FactoMiner [117] and ggcorrplot [118] used for analysis and plotting, respectively.

Gene expression in TPMs was plotted using GraphPad Prism v9 software (San Diego, CA,

USA), showing the mean of each analyzed gene and the standard deviation (SD) between

libraries for the same gene. Differential gene expression between groups was calculated using

DESeq2 [119]. The EXP-1 included five libraries for groups L1 and 2, and six for L3, and EXP-2,

four libraries for groups H and G, and three for WL. A Benjamini-Hochberg (BH) adjusted value

< 0.05 and a FC > 0 was used to define differentially expressed (DE) transcripts between

groups. Sequences with a TPM expression ≥ 5 were separated from all DE transcripts and were

used for all analyses. Transcripts with low expression may not be translated into proteins in the

tissues analyzed or may be associated with pseudo- genes or isoforms [120]. Consequently, DE

ABC and OBP gene families surpassing 5 TPMs were plotted. Results for other sequences

outside this selection were provided as Supplementary Data Tables.

Phylogenetic analysis

The phylogenetic analysis was completed for the ABC and OBP sensory families. ABC protein

sequences were identified by running a NCBI BLASTp against the *L. cuprina* genome assembly,

NCBI ID ASM2204524v1, and using *D. melanogaster* ABC protein sequences as query, the

latter obtained from the FlyBase database (https://flybase.org/). *L. cuprina* OBP curated protein

sequences were obtained from a previous study performed in the same species [25]. Only *L.*

cuprina sensory receptors showing a TPM expression ≥ 5 in the present work were included in

the final phylogenetic analysis. If there was more than one isoform per gene, only isoforms with

different transcript expression were added. Further, OBPs and ABC transporters previously

identified as larval biased in *L. sericata* [30] were added to the analysis.

Prior to phylogenetic analysis, a sequence alignment was performed using the G-INS-i

method in MAFFT v7 [121]. A preliminary phylogenetic analysis including all ABC receptors

identified for *L. cuprina* with a TPM expression > 0, using the Neighbor-Joining (NJ) algorithm

[122] and the software MEGA v11 [123] was performed to rename *L. cuprina* ABCs based on direct

orthology to *D. melanogaster* sequences. Subsequently, the Maximum likelihood (ML) method

set for 1000 bootstrap replications (using the same software) was utilized to improve a second

NJ tree completed including only ABC receptors with a TPM expression ≥ 5. The best

substitution model for each ML run was determined by using the tool "Find Best DNA/Protein

Models" of the same software, prior to running the phylogenetic test. Tree figures were

produced using the interactive Tree of Life (iTOL) v6.8.2 software (https://itol.embl.de/). The

analysis for the OBP sequences was completed using the same protocol except for the

renaming process.

Odorant coreceptor (*Orco*) experiment

Gene editing using CRISPR/Cas9

LcupOrco gene sequence, guide RNA and Cas9 in vitro assay

The guide RNA (gRNA) selected for gene editing was the ortholog of a gRNA previously used to

edit the *ChomOrco* gene in *C. hominivorax* [90]. Supplementary Note 1 provides further details

about the *LcupOrco* gene sequence including exonic regions, gRNA position, primers used to

amplify the *LcupOrco* gene fragment encompassing the Cas9 cutting site, and the Polymerase

Chain Reaction (PCR) cycling parameters. The *LcupOrco* gene model showing the gene editing

site was edited with Adobe Illustrator (San Jose, CA, USA), and the LcupOrco protein

secondary structure generated with the same purpose was plotted with the Protter software

(https://wlab.ethz.ch/protter/start/). The gRNA used was CRISPR-Cas9_crRNA [Integrated DNA

Technologies (IDT), Coralville, IA, USA]. The latter was mixed with the CRISPR-Cas9_tracrRNA

(IDT, Cat. # 1072532), using an equal volume of both RNAs after diluting them to the same

concentration. Subsequently, the mix was heated at 95ºC for 5 min, cooled to ~ 15ºC for 5 min

and kept at -20ºC until use.

The gRNA cutting efficiency was tested *in vitro* using Cas9 protein [New England

Biolabs (NEB), Ipswich, MA, USA, Cat. # M0646T]. Cas9 and gRNA were mixed for a final

concentration of 66.6 and 160 nM, respectively, and incubated for 20 min at 25°C to form the

Cas9 ribonucleoprotein (RNP) complex. A PCR-generated fragment containing the targeted

sequence (Supplementary Note 1), was added to the mix for a final concentration of 3 nM and

incubated for 60 min at 25°C. The reaction (RXN) was ended by adding 1 µl of Proteinase K 10

µg/µl (Zymo, Cat. # D3001-2-20) and incubated for 10 min. The RXN mix was then loaded into a

1.5% agarose gel in parallel to uncut gDNA used before as a template (same concentration),

electrophoresed at 60 volts for 90 min and visualize using a Gel Doc™ EZ System (Bio-Rad,

Hercules, CA, USA).

Insert construction, CRISPR/Cas9 mix and embryo injections

A Synthetic DNA fragment including two sections of the *LcupOrco* gene, plus restriction cutting

sites for the enzymes XhoI (NEB, Cat. # R0146S) and NotI (NEB, Cat. # R0189S) was acquired

cloned within the plasmid pUCIDT-AMP GoldenGate (IDT). The ZsGreen fluorescent marker

later added to the same construct was obtained and purified from a donor plasmid [124]. The

pUCIDT-AMP GoldenGate plasmid and ZsGreen fragment were ligated using T4 Ligase (NEB,

Cat. # M0202S) and then used for transformation of commercial competent *Escherichia coli*

cells (NEB, Cat. # C3019H) following the manufacturer's specifications. Plasmids were obtained

from 10 single clones using the ZR Miniprep Kit (Zymo, Cat. # D4016) following the

manufacturer's specifications. Subsequently, plasmids were evaluated by restriction enzyme

digestion analyses. The whole plasmid DNA sequencing was performed with plasmids that

showed the correct banding patterns (MGH CCIB DNA core, Boston, MA). For embryo

microinjection, plasmid DNA was obtained by using a Midiprep Kit (Zymo, Cat. # D4200) and

further purified using the DNA Clean Kit (Zymo, Cat. # D4004) following the manufacturer's

specifications. Prior to injection the DNA was diluted in NF-water to 1250 ng/μl concentration.

For further details refer to Supplementary Fig. 10.

The Cas9 (IDT, Cat. # 1081058) was used for the *in vivo* embryo injections; the same

was mixed with the duplex crRNA::tracrRNA, 1M Potassium chloride (KCl) and 1 μl of 3.1 buffer

(NEB, Cat. # B6003S) and incubated at 25°C for 20 min to form the RNP complex. The mixture

was cooled to room temperature for 5 min and the plasmid carrying the *LcupOrco* insert was

added to the mix to reach a final plasmid concentration of 500 ng/μl. Final concentrations for the

other compounds were as follows: Cas9 = 3.1 μM (500 ng/μl); crRNA::tracrRNA = 7.5 μM (250

ng/μl) and KCl = 250 mM. The mix was cleaned before injections by 2 min centrifugation at 4°C

using a 0.45 μM filter column (Sigma-Aldrich, Burlington, MS, USA, Cat. # UFC30HV) and keep

on ice all the time.

The needles used for microinjections were generated by pulling filamented quartz

capillaries (Sutter Instrument, Novato, CA, USA, Cat. # QF100-70-10). The puller used was a

Sutter Instrument P-2000 set for: Heat = 690; Filament = 4; Velocity = 60; Delay = 160; Pull =

170. After pulling, needle's tips were beveled for ~ 10 seconds (sec) using a BV-10 micropipette

beveler (Sutter Instrument), a fine abrasive plate and an angle of 22.5°.

Before injections, microinjection mix was loaded into the needles using a Microloader™

(Eppendorf, Hamburg, Germany, Cat. # 930001007). The microinjector used was a XenoWorks

Digital Microinjector (Sutter Instrument), set for continuous flow, 1000 hectopascal (hPa) of

pressure and +5 hPa of transfer pressure, together with a motorized XenoWorks

Micromanipulator (Sutter Instrument) and a stereomicroscope M125 (Leica, Wetzlar, Germany).

Flies used for egg collection were LA07 8-day-old gravid females kept with males of the same

age in a plastic container (Supplementary Fig. 11). The flies were stimulated to begin ovipositing

by adding 93/7% lean-fat fresh grounded beef placed in a 35 mm Petri dish for 15 min.

Following, another 35 mm-plate with fresh meat was added to collect new eggs for 5 min. An

egg cluster was transferred to a piece of tissue paper (Kimberly-Clark, Roswell, GA, USA),

moistened with distilled (DI)-water and located in one of the depressions of a double-depression

microscope slide (VWR, Radnor, PA, USA, Cat. # 470003-480). Twenty eggs per round of

injections were separated under an Olympus stereomicroscope SZX10 (Olympus, Center

Valley, PA, USA), using a 00-paint brush and accommodated onto a Scotch® double-sided film

tape (3M, Saint Paul, MN, USA), located in the other depression of the same slide. The eggs

were then desiccated for 5 min using a desiccation chamber of 150 mm in OD (Nalgene,

Rochester, NY, USA) containing a desiccant cartridge (SP Bel-art, Wayne, NJ, USA, Cat. #

F42049-0100). Subsequently eggs were covered with oil (Sigma-Aldrich, Cat. # H8773), rest for

3 min and injected at the posterior end for ~ 1 sec. The developmental time for the injected

embryos was between 18 to 24 min, for the first and last injected embryo, respectively. After

injection, embryos were kept overnight in an incubator chamber (Billups-Rothenberg, Del Mar,

CA, USA, Cat. # MIC-101) filled with oxygen and protected from light using a carboard cover

until eggs hatched, *i.e.* between 19 to 27 h at 23.5°C. After hatching, newly emerged L1s were

transferred using a 00-paint brush to moistened tissue paper (Kimberly-Clark) and then moved

to a 3-ounces clear plastic cup filled with 50 grams of the same meat mentioned above and let

larvae develop until tested at L3 stage.

Fly screening, crosses and genotyping

Injected embryos were screened 7 days after injections (L3 stage) using a stereo microscope

M205 FA (Leica) and a green filter. Only mosaic embryos from generation (G) 0 showing

transient expression of the plasmid were retained to obtain adult flies. Emerged adult flies were

crossed following the protocol described in Supplementary Fig. 12. G1 and G2 eggs obtained

from G0 and G1 crosses, respectively, were tested for positive fluorescence using the same

microscope, and only fluorescent eggs were kept.

At G3, a middle leg was collected using dissection forceps and scissors from 2-day-old

flies for genotyping. gDNA was extracted by placing legs in a DNA extraction buffer (Biosearch

Technologies, Hoddesdon, UK, Cat. # QE09050) for 15 min at 65°C and 2 min at 98°C. Three

strategies were used to determine if the *LcupOrco* insert was landed at the correct *L. cuprina*

genome position, *i.e.* withing the *LcupOrco* coding region: 1) amplification of left side of the

insert + upstream genome; 2) amplification of right side of the insert + downstream genome;

and 3) amplification of the whole insert including upstream and downstream genome regions;

for further details refer to Supplementary Note 2. In parallel, potential indels and point mutations

at the Cas9 cutting site were investigated using same primers described in Supplementary Note

1. After PCR amplification, the gDNA was purified using the DNA Clean Kit and sequenced by

Sanger Sequencing. Amplicons >1 Kb were cloned using the pGEM®-T system (Promega,

Fitchburg, WI, USA, Cat. # A1360) together with competent cells following the manufacturer's

specifications. Plasmids were obtained, purified and sequenced following the same protocols

described in Supplementary Fig. 10.

Behavioral assays

Larval diet preference test

The larval diet preference test was previously used to evaluate *L. cuprina* larvae olfaction

behavior [26]. The larvae used for the assay were obtained using protocols described at the *Insect*

rearing conditions section. Before the assay, ~ 250 L1s were transferred to 50 g of fresh ground

beef 1 day after egg hatching. Two days after this step a total of 120 3-day-old larvae were

tested following protocols described at [26]. The options offered to the larvae were rotten (R) and

fresh (F) ground beef, either at 25±1 or 33±1°C, designated as cold (C) and hot (H) respectively,

and for further details about beef conditions refer to [26]. Those larvae that did not choose any

option after 10 min were considered as non-choice (NC). After the assay, larvae were placed in

separated 1.5 tubes filled with 300 µl of DNA preservative buffer (Zymo, Cat. # R1100-50) and

kept at -80°C until DNA extraction. Subsequently, all larvae were dried using tissue paper and

separately weighed using a LA204E analytical balance (Metter Toledo, Columbus, OH, USA).

Following, a small section of each larva was collected using dissection forceps and scissors and

placed back in the same buffer. Approximately one third of the larvae associated with each diet

preference, giving a total of 48 larvae, were genotyped using same protocols described at the

Fly screening, crosses and genotyping section and Supplementary Note 1.

Adult female olfaction assay

The adult female olfaction assay was completed using the same protocol, spatial olfactometer

and room conditions previously used to evaluate *L. cuprina* female olfaction behavior [25]. The

olfactometer consists of four separate chambers (replicates) where the flies were released.

Each of these chambers are connected with two smaller collection chambers A and B, where

meat samples were placed to attract and trap the flies. The attractants used to lure the flies

were 2 g of fresh (F) and rotten (R) beef at room temperature, placed in chambers A and B,

respectively.

The females used for the olfaction assay were from the same cohort of individuals used

for the larval diet preference test (see above). In addition, LA07 *wt* females were tested using

the same assay conditions, but on different days without mixing them with the *LcupOrco* flies.

All flies used for the olfaction assay were 10-day-old gravid females and were separated from

males the day before the assay using CO_2 to anesthetize the flies under a stereomicroscope

SMZ745 (Nikon, Melville, NY, USA). After sexes separation, females were kept in the same

room where the assay was performed for habituation, provided with tap water and cane sugar.

After the assay, females were kept in separated 1.5 mL empty tubes at -80°C until DNA

extraction. A total of 80 *wt* and 174 *LcupOrco* females were tested, and 72 of the latter,

randomly selected from all assayed *LcupOrco* females, were chosen for genotyping using the

same protocols described in the previous section. After the assay, females were separated into

three groups: 1) attracted to fresh (F) beef; 2) attracted to rotten (R) beef; and 3) flies that did

not respond to any stimulus, designated as non-choice (NC).

Mating assay

To evaluate the mating performance of the *LcupOrco* homozygous females (*LcupOrco*[-/-]), 5-day-

old females previously determined as heterozygous or homozygous females by genotyping

protocols described above, were mixed with twice as many adult *NPF*[-/-] males of approximate

same age, constitutively expressing the ZsGreen fluorescent protein marker. Fifty grams of

fresh grounded beef was offered O/N to females at 3, 10 and 17 days after mixing them with the

NPF[-/-] males. After oviposition, eggs were visualized and photographed using a stereo

microscope M205 FA (Leica), a green filter and a camera Leica K SC.

Data availability
All relevant data are in the manuscript. Raw sequences were uploaded to the NCBI database
under Bioproject PRJNA1195419.
890

891   **Code availability**
892   All software used in the present study has been previously reported and were cited in the body
893   of the present manuscript.
894
895   **References**
896

1.   Benbow, M. E., Tomberlin, J. K. & Tarone, A. M. *Carrion Ecology, Evolution, and Their Applications*. *Carrion*
ecology, evolution, and their applications (CRC Press, 2015). doi:10.1201/b18819.

2.   Wall, R., Rose, H., Ellse, L. & Morgan, E. Livestock ectoparasites: Integrated management in a changing
climate. *Vet Parasitol* **180**, 82–89 (2011).

3.   Getz, W. M. Biomass transformation webs provide a unified approach to consumer-resource modelling.
Ecol Lett **14**, 113–24 (2011).

4.   Mastrangelo, T. & Welch, J. An Overview of the Components of AW-IPM Campaigns against the New World
Screwworm. *Insects* **3**, 930–955 (2012).

5.   Heath, A. C. G. & Bishop, D. M. Flystrike in New Zealand: An overview based on a 16-year study, following
the introduction and dispersal of the Australian sheep blowfly, *Lucilia cuprina* Wiedemann (Diptera:
Calliphoridae). *Vet Parasitol* **137**, 333–344 (2006).

6.   Wardhaugh, K. G., Morton, R., Bedo, D., Horton, B. J. & Mahon, R. J. Estimating the incidence of fly myiases
in Australian sheep flocks: development of a weather-driven regression model. *Med Vet Entomol* **21**, 153–
167 (2007).

7.   Heath, A. & Levot, G. Parasiticide resistance in flies, lice and ticks in New Zealand and Australia:
mechanisms, prevalence and prevention. *N Z Vet J* **63**, 199–210 (2015).

8.   Sandeman, R. M. *et al.* Control of the sheep blowfly in Australia and New Zealand – are we there yet? *Int J*
Parasitol **44**, 879–891 (2014).

9.   Li, F., Wantuch, H. A., Linger, R. J., Belikoff, E. J. & Scott, M. J. Transgenic sexing system for genetic control
of the Australian sheep blow fly *Lucilia cuprina*. *Insect Biochem Mol Biol* **51**, 80–88 (2014).

10.   Yan, Y. & Scott, M. J. A transgenic embryonic sexing system for the Australian sheep blow fly *Lucilia*
cuprina. *Sci Rep* **5**, 16090 (2015).

11.   Byrd, J. H. & Tomberlin, J. K. *Forensic Entomology: The Utility of Arthropods in Legal Investigations*. (CRC
press, 2019).

12.   Fenton, A., Wall, R. & French, N. P. The effects of oviposition aggregation on the incidence of sheep blowfly
strike. *Vet Parasitol* **83**, 137–150 (1999).

13.   Ashworth, J. R. & Wall, R. Responses of the sheep blowflies *Lucilia sericata* and *L. cuprina* to odour and the
development of semiochemical baits. *Med Vet Entomol* **8**, 303–309 (1994).

14.   Lennox, F. G. Distribution of Ammonia in Larvæ of *Lucilia cuprina*. *Nature* **146**, 268–268 (1940).

15.   Casu, R. E. *et al.* Excretory/secretory chymotrypsin from *Lucilia cuprina*: purification, enzymatic specificity
and amino acid sequence deduced from mRNA. *Insect Mol Biol* **3**, 201–211 (1994).

16.   Arora, S., Baptista, C. & Lim, C. S. Maggot metabolites and their combinatory effects with antibiotic on
Staphylococcus aureus. *Ann Clin Microbiol Antimicrob* **10**, 6 (2011).

17.   Kotzé, Z., Villet, M. H. & Weldon, C. W. Heat accumulation and development rate of massed maggots of the
sheep blowfly, *Lucilia cuprina* (Diptera: Calliphoridae). *J Insect Physiol* **95**, 98–104 (2016).

18.   Kotzé, Z., Villet, M. H. & Weldon, C. W. Effect of temperature on development of the blowfly, *Lucilia*
cuprina (Wiedemann) (Diptera: Calliphoridae). *Int J Legal Med* **129**, 1155–1162 (2015).

19.   Bambaradeniya, Y. T. B., Karunaratne, W. A. I. P., Tomberlin, J. K., Goonerathne, I. & Kotakadeniya, R. B.
Temperature and tissue type impact development of *Lucilia cuprina* (Diptera: Calliphoridae) in Sri Lanka. *J*
Med Entomol **55**, 285–291 (2018).

20.   Kotze, A. & James, P. Control of sheep flystrike: what's been tried in the past and where to from here. *Aust*
Vet J **100**, 1–19 (2022).

21.   Forbes, S. L. & Perrault, K. A. Decomposition Odour Profiling in the Air and Soil Surrounding Vertebrate
Carrion. *PLoS One* **9**, e95107 (2014).

22.   Zhu, J. J., Chaudhury, M. F., Tangtrakulwanich, K. & Skoda, S. R. Identification of Oviposition Attractants of
the Secondary Screwworm, *Cochliomyia macellaria* (F.) Released from Rotten Chicken Liver. *J Chem Ecol*
39, 1407–1414 (2013).

23.   Chaudhury, M. F., Skoda, S. R., Sagel, A. & Welch, J. B. Volatiles Emitted From Eight Wound-Isolated
Bacteria Differentially Attract Gravid Screwworms (Diptera: Calliphoridae) to Oviposit. *J Med Entomol* **47**,
349–354 (2010).

24. Liu, W., Longnecker, M., Tarone, A. M. & Tomberlin, J. K. Responses of *Lucilia sericata* (Diptera:
Calliphoridae) to compounds from microbial decomposition of larval resources. *Anim Behav* **115**, 217–225
(2016).
25. Wulff, J. P. *et al.* Antennal transcriptome analysis reveals sensory receptors potentially associated with
host detection in the livestock pest *Lucilia cuprina*. *Parasit Vectors* **17**, 308 (2024).
26. Cunha, V. A. S., Tandonnet, S., Ferreira, D. L., Rodrigues, A. V. & Torres, T. T. Exploring Life History Choices:
Using Temperature and Substrate Type as Interacting Factors for Blowfly Larval and Female Preferences.
Journal of Visualized Experiments e65835 (2023) doi:10.3791/65835.
27. Guerrero, F. D. *et al.* A Database of Expressed Genes From *Cochliomyia hominivorax* (Diptera:
Calliphoridae). *J Med Entomol* **46**, 1109–1116 (2009).
28. Zhang, M. *et al.* Analysis of the Transcriptome of Blowfly *Chrysomya megacephala* (Fabricius) Larvae in
Responses to Different Edible Oils. *PLoS One* **8**, e63168 (2013).
29. Hjelmen, C. E. *et al.* Identification and Characterization of Small RNA Markers of Age in the Blow Fly
Cochliomyia macellaria (Fabricius) (Diptera: Calliphoridae). *Insects* **13**, 948 (2022).
30. Davis, R. J. *et al.* Genome and transcriptome sequencing of the green bottle fly, *Lucilia sericata*, reveals
underlying factors of sheep flystrike and maggot debridement therapy. *Genomics* **113**, 3978–3988 (2021).
31. Liu, Z., Han, H., Meng, F., Jiang, Y. & Cai, J. Dynamic transcriptome profiling exploring cold tolerance in
forensically important blow fly, *Aldrichina grahami* (Diptera: Calliphoridae). *BMC Genomics* **21**, 92 (2020).
32. McKenna, C. H. *et al.* Gene expression in *Lucilia sericata* (Diptera: Calliphoridae) larvae exposed to
Pseudomonas aeruginosa and *Acinetobacter baumannii* identifies shared and microbe-specific induction of
immune genes. *Insect Mol Biol* **31**, 85–100 (2022).
33. Fishilevich, E. *et al.* Chemotaxis Behavior Mediated by Single Larval Olfactory Neurons in *Drosophila*.
Current Biology **15**, 2086–2096 (2005).
34. Gerber, B. & Stocker, R. F. The *Drosophila* Larva as a Model for Studying Chemosensation and
Chemosensory Learning: A Review. *Chem Senses* **32**, 65–89 (2007).
35. Gomez-Marin, A. & Louis, M. Multilevel control of run orientation in *Drosophila* larval chemotaxis. *Front
Behav Neurosci* **8**, (2014).
36. Gomez-Marin, A., Stephens, G. J. & Louis, M. Active sampling and decision making in *Drosophila*
chemotaxis. *Nat Commun* **2**, 441 (2011).
37. Louis, M., Huber, T., Benton, R., Sakmar, T. P. & Vosshall, L. B. Bilateral olfactory sensory input enhances
chemotaxis behavior. *Nat Neurosci* **11**, 187–199 (2008).
38. Kreher, S. A., Mathew, D., Kim, J. & Carlson, J. R. Translation of Sensory Input into Behavioral Output via an
Olfactory System. *Neuron* **59**, 110–124 (2008).
39. Grillet, M., Campagner, D., Petersen, R., McCrohan, C. & Cobb, M. The peripheral olfactory code in
Drosophila larvae contains temporal information and is robust over multiple timescales. *Proceedings of the
Royal Society B: Biological Sciences* **283**, 20160665 (2016).
40. Jiao, Y., Moon, S. J. & Montell, C. A *Drosophila* gustatory receptor required for the responses to sucrose,
glucose, and maltose identified by mRNA tagging. *Proceedings of the National Academy of Sciences* **104**,
14110–14115 (2007).
41. Ni, L. *et al.* A gustatory receptor paralogue controls rapid warmth avoidance in *Drosophila*. *Nature* **500**,
580–584 (2013).
42. Sang, J., Rimal, S. & Lee, Y. *Gustatory receptor 28b* is necessary for avoiding saponin in *Drosophila
melanogaster*. *EMBO Rep* **20**, (2019).
43. Ni, L. *et al.* The Ionotropic Receptors IR21a and IR25a mediate cool sensing in *Drosophila*. *Elife* **5**, (2016).
44. Stewart, S., Koh, T.-W., Ghosh, A. C. & Carlson, J. R. Candidate ionotropic taste receptors in the *Drosophila*
larva. *Proceedings of the National Academy of Sciences* **112**, 4195–4201 (2015).
45. Ainsley, J. A. *et al.* Enhanced Locomotion Caused by Loss of the *Drosophila* DEG/ENaC Protein Pickpocket1.
Current Biology **13**, 1557–1563 (2003).
46. Klein, M. *et al.* Sensory determinants of behavioral dynamics in *Drosophila* thermotaxis. *Proceedings of the
National Academy of Sciences* **112**, E220–E229 (2015).

47. Kwon, Y., Shen, W. L., Shim, H.-S. & Montell, C. Fine Thermotactic Discrimination between the Optimal and Slightly Cooler Temperatures via a TRPV Channel in Chordotonal Neurons. *The Journal of Neuroscience* **30**, 10465–10471 (2010).

48. Walker, R. G., Willingham, A. T. & Zuker, C. S. A *Drosophila* Mechanosensory Transduction Channel. *Science (1979)* **287**, 2229–2234 (2000).

49. Pulver, S. R., Pashkovski, S. L., Hornstein, N. J., Garrity, P. A. & Griffith, L. C. Temporal Dynamics of Neuronal Activation by Channelrhodopsin-2 and TRPA1 Determine Behavioral Output in *Drosophila* Larvae. *J Neurophysiol* **101**, 3075–3088 (2009).

50. Rihani, K., Ferveur, J.-F. & Briand, L. The 40-Year Mystery of Insect Odorant-Binding Proteins. *Biomolecules* **11**, 509 (2021).

51. Leitch, O., Papanicolaou, A., Lennard, C., Kirkbride, K. P. & Anderson, A. Chemosensory genes identified in the antennal transcriptome of the blowfly *Calliphora stygia*. *BMC Genomics* **16**, 255 (2015).

52. Vulpe, A. *et al.* An ammonium transporter is a non-canonical olfactory receptor for ammonia. *Current Biology* **31**, 3382-3390.e7 (2021).

53. Rondón, J. J. *et al.* Evolution of the odorant-binding protein gene family in *Drosophila*. *Front Ecol Evol* **10**, (2022).

54. Wicher, D. & Marion-Poll, F. Editorial: Function and Regulation of Chemoreceptors. *Front Cell Neurosci* **12**, (2018).

55. Task, D. *et al.* Chemoreceptor co-expression in *Drosophila melanogaster* olfactory neurons. *Elife* **11**, (2022).

56. Utashiro, N., Williams, C. R., Parrish, J. Z. & Emoto, K. Prior activity of olfactory receptor neurons is required for proper sensory processing and behavior in *Drosophila* larvae. *Sci Rep* **8**, 8580 (2018).

57. Dermauw, W. & Van Leeuwen, T. The ABC gene family in arthropods: Comparative genomics and role in insecticide transport and resistance. *Insect Biochem Mol Biol* **45**, 89–110 (2014).

58. Kumar, V. *et al.* Delineating the Role of *Aedes aegypti* ABC Transporter Gene Family during Mosquito Development and Arboviral Infection via Transcriptome Analyses. *Pathogens* **10**, 1127 (2021).

59. Wu, C., Chakrabarty, S., Jin, M., Liu, K. & Xiao, Y. Insect ATP-Binding Cassette (ABC) Transporters: Roles in Xenobiotic Detoxification and Bt Insecticidal Activity. *Int J Mol Sci* **20**, 2829 (2019).

60. Andersen, D. S. & Leevers, S. J. The Essential *Drosophila* ATP-binding Cassette Domain Protein, Pixie, Binds the 40 S Ribosome in an ATP-dependent Manner and Is Required for Translation Initiation. *Journal of Biological Chemistry* **282**, 14752–14760 (2007).

61. Fostier, C. R. *et al.* ABC-F translation factors: from antibiotic resistance to immune response. *FEBS Lett* **595**, 675–706 (2021).

62. Wang, Y. *et al.* Dysfunction of *Oskyddad* causes Harlequin-type ichthyosis-like defects in *Drosophila melanogaster*. *PLoS Genet* **16**, e1008363 (2020).

63. Zuber, R. *et al.* The ABC transporter Snu and the extracellular protein Snsl cooperate in the formation of the lipid-based inward and outward barrier in the skin of *Drosophila*. *Eur J Cell Biol* **97**, 90–101 (2018).

64. Abo-Elghar, G. E., Fujiyoshi, P. & Matsumura, F. Significance of the sulfonylurea receptor (SUR) as the target of diflubenzuron in chitin synthesis inhibition in *Drosophila melanogaster* and *Blattella germanica*. *Insect Biochem Mol Biol* **34**, 743–752 (2004).

65. Whiten, S. R., Eggleston, H. & Adelman, Z. N. Ironing out the Details: Exploring the Role of Iron and Heme in Blood-Sucking Arthropods. *Front Physiol* **8**, (2018).

66. Selem, G. Sh., Geden, C. J., Khater, H. & Khater, K. S. Effects of larval diets on some biological parameters and morphometric and biochemical analysis of ovaries of *Lucilia cuprina* (Wiedemann) (Diptera: Calliphoridae). *Journal of Vector Ecology* **48**, (2023).

67. Lockey, K. H. Lipids of the insect cuticle: origin, composition and function. *Comparative Biochemistry and Physiology Part B: Comparative Biochemistry* **89**, 595–645 (1988).

68. Zhang, Y. *et al.* Odorant-Binding Proteins Contribute to the Defense of the Red Flour Beetle, *Tribolium castaneum*, Against Essential Oil of *Artemisia vulgaris*. *Front Physiol* **11**, (2020).

69. Abendroth, J. A., Moural, T. W., Wei, H. & Zhu, F. Roles of insect odorant binding proteins in communication and xenobiotic adaptation. *Frontiers in Insect Science* **3**, (2023).

70. Zhao, R. *et al.* Omics Analysis of Odorant-Binding Proteins and Cuticle-Enriched *SfruOBP18* Confers Multi-Insecticide Tolerance in *Spodoptera frugiperda*. *J Agric Food Chem* **72**, 22532–22544 (2024).

71. Cotten, M. L. *et al.* NMR chemical shift assignment of *Drosophila* odorant binding protein 44a in complex
with 8(Z)-eicosenoic acid. *Biomol NMR Assign* (2024) doi:10.1007/s12104-024-10178-2.

72. Jeong, Y. T. *et al.* An Odorant-Binding Protein Required for Suppression of Sweet Taste by Bitter Chemicals.
Neuron **79**, 725–737 (2013).

73. González-González, A. *et al.* Virtual Screening of Plant Volatile Compounds Reveals a High Affinity of
Hylamorpha elegans (Coleoptera: Scarabaeidae) Odorant-Binding Proteins for Sesquiterpenes From Its
Native Host. *Journal of Insect Science* **16**, 30 (2016).

74. Giordano, D., Facchiano, A., D'Auria, S. & Loreto, F. A hypothesis on the capacity of plant odorant-binding
proteins to bind volatile isoprenoids based on in silico evidences. *Elife* **10**, (2021).

75. Qiao, H. *et al.* Discrimination of alarm pheromone (E)-β-farnesene by aphid odorant-binding proteins.
Insect Biochem Mol Biol **39**, 414–419 (2009).

76. Rahman, M. M., Franch-Marro, X., Maestro, J. L., Martin, D. & Casali, A. Local Juvenile Hormone activity
regulates gut homeostasis and tumor growth in adult *Drosophila*. *Sci Rep* **7**, 11677 (2017).

77. Taracena-Agarwal, M. L. *et al.* Juvenile Hormone as a contributing factor in establishing midgut microbiota
for fecundity and fitness enhancement in adult female *Aedes aegypti*. *Commun Biol* **7**, 687 (2024).

78. Sokabe, T. & Tominaga, M. A temperature-sensitive TRP ion channel, *painless*, functions as a noxious heat
sensor in fruit flies. *Commun Integr Biol* **2**, 170–173 (2009).

79. Charlu, S., Wisotsky, Z., Medina, A. & Dahanukar, A. Acid sensing by sweet and bitter taste neurons in
Drosophila melanogaster. *Nat Commun* **4**, 2042 (2013).

80. Al-Anzi, B., Tracey, W. D. & Benzer, S. Response of *Drosophila* to Wasabi Is Mediated by painless, the Fly
Homolog of Mammalian TRPA1/ANKTM1. *Current Biology* **16**, 1034–1040 (2006).

81. Xu, J., Sornborger, A. T., Lee, J. K. & Shen, P. *Drosophila* TRPA channel modulates sugar-stimulated neural
excitation, avoidance and social response. *Nat Neurosci* **11**, 676–682 (2008).

82. Herboso, L., Talamillo, A., Perez, C. & Barrio, R. Expression of the Scavenger Receptor Class B type I (SR-BI)
family in *Drosophila melanogaster*. *Int J Dev Biol* **55**, 603–611 (2011).

83. Shi, X.-Z., Zhong, X. & Yu, X.-Q. *Drosophila melanogaster* NPC2 proteins bind bacterial cell wall components
and may function in immune signal pathways. *Insect Biochem Mol Biol* **42**, 545–556 (2012).

84. Chen, M.-M. *et al.* Characterization of an Ecdysteroid-Regulated 16 kDa Protein Gene in Chinese Oak
Silkworm, *Antheraea pernyi* (Lepidoptera: Saturniidae). *Journal of Insect Science* **20**, (2020).

85. Wulff, J. P., Traverso, L. M., Latorre-Estivalis, J. M., Segura, D. F. & Lanzavecchia, S. B. Identification of
candidate genes associated with host-seeking behavior in the parasitoid wasp *Diachasmimorpha*
longicaudata. *BMC Genomics* **25**, 147 (2024).

86. Yang, Y., Hua, D., Zhu, J., Wang, F. & Zhang, Y. Chemosensory protein 4 is required for *Bradysia odoriphaga*
to be olfactory attracted to sulfur compounds released from Chinese chives. *Front Physiol* **13**, (2022).

87. Delventhal, R. *et al.* The taste response to ammonia in *Drosophila*. *Sci Rep* **7**, 43754 (2017).

88. Browne, A. & O'Donnell, M. J. Ammonium secretion by Malpighian tubules of *Drosophila melanogaster*:
application of a novel ammonium-selective microelectrode. *Journal of Experimental Biology* **216**, 3818–
3827 (2013).

89. Paulo, D. F., Williamson, M. E. & Scott, M. J. CRISPR/Cas9 Genome Editing in the New World Screwworm
and Australian Sheep Blowfly. in 173–201 (2022). doi:10.1007/978-1-0716-2301-5_10.

90. Paulo, D. F. *et al.* Disruption of the odorant coreceptor *Orco* impairs foraging and host finding behaviors in
the New World screwworm fly. *Sci Rep* **11**, 11379 (2021).

91. Yan, H. *et al.* An Engineered *Orco* Mutation Produces Aberrant Social Behavior and Defective Neural
Development in Ants. *Cell* **170**, 736-747.e9 (2017).

92. Fan, X.-B. *et al.* Mutagenesis of the odorant receptor co-receptor (*Orco*) reveals severe olfactory defects in
the crop pest moth *Helicoverpa armigera*. *BMC Biol* **20**, 214 (2022).

93. Lamb, A. M., Wang, Z., Simmer, P., Chung, H. & Wittkopp, P. J. *ebony* Affects Pigmentation Divergence and
Cuticular Hydrocarbons in *Drosophila americana* and *D. novamexicana*. *Front Ecol Evol* **8**, (2020).

94. Drapeau, M. D., Radovic, A., Wittkopp, P. J. & Long, A. D. A gene necessary for normal male courtship,
yellow, acts downstream of *fruitless* in the *Drosophila melanogaster* larval brain. *J Neurobiol* **55**, 53–72
(2003).

95. Barek, H., Zhao, H., Heath, K., Veraksa, A. & Sugumaran, M. *Drosophila yellow-h* encodes dopaminechrome tautomerase: A new enzyme in the eumelanin biosynthetic pathway. *Pigment Cell Melanoma Res* **35**, 26–37 (2022).

96. Song, Q. & An, S. Bursicon, a Neuropeptide Hormone That Controls Cuticle Tanning and Beyond. in *Recent Advances in Entomological Research* (eds. Liu, T. & Kang, L.) 132–149 (Springer Berlin Heidelberg, Berlin, Heidelberg, 2011). doi:10.1007/978-3-642-17815-3_8.

97. Chia, W. *et al.* Molecular analysis of the *yellow* locus of *Drosophila*. *EMBO J* **5**, 3597–3605 (1986).

98. Richardt, A. *et al.* Ebony, a Novel Nonribosomal Peptide Synthetase for β-Alanine Conjugation with Biogenic Amines in *Drosophila*. *Journal of Biological Chemistry* **278**, 41160–41166 (2003).

99. Khan, S. A., Kojour, M. A. M. & Han, Y. S. Recent trends in insect gut immunity. *Front Immunol* **14**, 1272143 (2023).

100. Kennel, C. *et al.* Differential Expression of Mucins in Murine Olfactory Versus Respiratory Epithelium. *Chem Senses* **44**, 511–521 (2019).

101. Sano, H. Coupling of growth to nutritional status: The role of novel periphery-to-brain signaling by the CCHa2 peptide in *Drosophila melanogaster*. *Fly (Austin)* **9**, 183–187 (2015).

102. Ren, G. R. *et al.* CCHamide-2 Is an Orexigenic Brain-Gut Peptide in *Drosophila*. *PLoS One* **10**, e0133017 (2015).

103. Benguettat, O. *et al.* The DH31/CGRP enteroendocrine peptide triggers intestinal contractions favoring the elimination of opportunistic bacteria. *PLoS Pathog* **14**, e1007279 (2018).

104. Zhang, X. *et al.* Gut transcriptome analysis of P450 genes and cytochrome P450 reductase in three moth species feeding on gymnosperms or angiosperms. *Front Ecol Evol* **10**, (2022).

105. Lahiri, S. *et al.* Two Alternating Motor Programs Drive Navigation in *Drosophila* Larva. *PLoS One* **6**, e23180 (2011).

106. Ferveur, J.-F., Cortot, J., Rihani, K., Cobb, M. & Everaerts, C. Desiccation resistance: effect of cuticular hydrocarbons and water content in *Drosophila melanogaster* adults. *PeerJ* **6**, e4318 (2018).

107. Van der Horst, D. J., Van Marrewijk, W. J. A. & Diederen, J. H. B. Adipokinetic hormones of insect: Release, signal transduction, and responses. in *International Review of Cytology* vol. 211 179–240 (Academic Press, 2001).

108. Čeřovský, V. & Bém, R. Lucifensins, the Insect Defensins of Biomedical Importance: The Story behind Maggot Therapy. *Pharmaceuticals* **7**, 251–264 (2014).

109. De Gregorio, E. The Toll and Imd pathways are the major regulators of the immune response in *Drosophila*. *EMBO J* **21**, 2568–2579 (2002).

110. Martelli, C. *et al.* SIFamide Translates Hunger Signals into Appetitive and Feeding Behavior in *Drosophila*. *Cell Rep* **20**, 464–478 (2017).

111. Lee, K.-S., You, K.-H., Choo, J.-K., Han, Y.-M. & Yu, K. *Drosophila* Short Neuropeptide F Regulates Food Intake and Body Size. *Journal of Biological Chemistry* **279**, 50781–50789 (2004).

112. Cholewiński, M. *et al.* Short neuropeptide F in integrated insect physiology. *Journal of Zhejiang University-SCIENCE B* **25**, 389–409 (2024).

113. Deng, L. & Chiu, I. M. A neuropeptide regulates immunity across species. *Neuron* **110**, 1275–1277 (2022).

114. Willis, J. H., Iconomidou, V. A., Smith, R. F. & Hamodrakas, S. J. Cuticular Proteins. in *Comprehensive Molecular Insect Science* (ed. Gilbert, L. I.) 79–109 (Elsevier, Amsterdam, 2005). doi:10.1016/B0-44-451924-6/00050-8.

115. Kang, S. H. & Fuchs, M. S. Isolation of a chymotrypsin inhibitor from *Drosophila melanogaster*. *Insect Biochem* **4**, 1–8 (1974).

116. Williamson, M. E., Yan, Y. & Scott, M. J. Conditional knockdown of transformer in sheep blow fly suggests a role in repression of dosage compensation and potential for population suppression. *PLoS Genet* **17**, e1009792 (2021).

117. Lê, S., Josse, J. & Husson, F. **FactoMineR** : An *R* Package for Multivariate Analysis. *J Stat Softw* **25**, (2008).

118. Valero-Mora, P. M. **ggplot2:** Elegant Graphics for Data Analysis. *J Stat Softw* **35**, (2010).

119. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, (2014).

120.   Mika, K. & Benton, R. Olfactory Receptor Gene Regulation in Insects: Multiple Mechanisms for Singular
Expression. *Front Neurosci* **15:738088**, (2021).
121.   Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7: improvements in
performance and usability. *Mol Biol Evol* **30**, 772–80 (2013).
122.   Saitou, N. & Nei, M. The neighbor-joining method: a new method for reconstructing phylogenetic trees.
Mol Biol Evol (1987) doi:10.1093/oxfordjournals.molbev.a040454.
123.   Tamura, K., Stecher, G. & Kumar, S. MEGA11: molecular evolutionary genetics analysis version 11. *Mol Biol*
Evol **38**, 3022–3027 (2021).
124.   Concha, C. *et al.* Efficient germ-line transformation of the economically important pest species *Lucilia*
cuprina and *Lucilia sericata* (Diptera, Calliphoridae). *Insect Biochem Mol Biol* **41**, 70–75 (2011).
1161
1162

1163 **Funding**

1164 This work was supported by the National Science Foundation Grant No. DEB-2030345.

1165

1166 **Author information**

1167

Authors and Affiliations

Juan P. Wulff [a], Rachel K. Laminack [a], and Maxwell J. Scott [a]
[a] Entomology and Plant Pathology, NCSU, Raleigh, NC, USA.
JPW: jpwulff@ncsu.edu
RKL: rklamina@ncsu.edu
MJS: mjscott3@ncsu.edu

1174

1175 ## Author contributions

1176 MJS and JPW conceptualized the experiments. JPW and RKL collected the samples and
1177 performed the experiments, JPW completed the analyses and produced the figures. JPW and
1178 MJS wrote the manuscript and RKL reviewed and edited the manuscript. All authors read and
1179 approved the final manuscript.

1180

1181 ## Corresponding author

1182 Maxwell J. Scott
1183 MJS: mjscott3@ncsu.edu

1184

1185 **Ethics declarations**

1186

1187 ## Competing interests

1188 The authors declare that there are no competing financial interests.

1189

1190 ## Consent for publication

1191 Not applicable.

1192