Genetic and behavioral analyses suggest that larval and adult stages of *Lucilia cuprina* employ different sensory systems to detect rotten beef

Juan P. Wulff [a], Rachel K. Laminack [a], and Maxwell J. Scott [a*]

[a] Entomology and Plant Pathology, NCSU, Raleigh, NC, USA.

*Correspondence: mjscott3@ncsu.edu

E-mail addresses:
JPW: jpwulff@ncsu.edu
RKL: rklamina@ncsu.edu
MJS: mjscott3@ncsu.edu

ORCIDs:
MJS: https://orcid.org/0000-0001-6536-4735
JPW: https://orcid.org/0000-0002-5773-4684

Abstract

Background: The blowfly *Lucilia cuprina* is a destructive parasite of sheep that causes flystrike or myiasis. Larvae consume the animal's living flesh, producing large wounds that can lead to death. Growing resistance to conventional control methods has prompted the analysis of alternative strategies.

Methods: An RNA-Seq analysis was used to identify sensory receptors and other genes relevant to the physiology of *L. cuprina* larvae. Adult females and larvae of the same species carrying a loss-of-function mutation for the *L. cuprina* odorant coreceptor gene (*LcupOrco*) were obtained by gene editing. Their response to fresh and rotten meat at different temperatures was evaluated.

Results: The RNA-Seq analysis of whole larvae at different stages and third instar head and gut tissues, suggested that odorant (OR), gustatory, ionotropic and *pickpocke*t receptors may not play a central role in the *L. cuprina* larval sensory signaling and digestive systems. Rather, ATP-binding cassettes (ABCs) were highly enriched in head and gut RNA, and odorant-binding proteins (OBPs) only in the head. To confirm that ORs are not essential for larval detection of rotten beef, diet-choice assays were performed including larvae and adults homozygous for a null mutation in *LcupOrco*. While the attraction of adult females to rotten beef was fully disrupted, *LcupOrco* mutant larvae showed no change in diet preference.

Conclusions: The expression pattern of the ABC and OBP gene families suggests a central role in the sensory system of the *L. cuprina* larva for these receptors. Behavioral assays showed that ORs are essential for the adult female response to rotten beef, but not for larval behavior.

These findings are consistent with high levels of expression of *LcupOrco* in the adult female

antenna but very low expression in larvae.

Keywords: Myiasis; Blowfly livestock pest; Australian sheep blowfly; host seeking; RNA-Seq;

DESeq2; adult stage olfaction; larval stage olfaction; chemoreceptor; *Orco*.

Background

Immature stages of blowfly species (Diptera: Calliphoridae) display a wide spectrum of feeding

specializations [1]. Based on their feeding habits, species within this family can be classified

from obligate parasites of live animals to obligate necrophagous (dead tissue consumers), with

many facultative forms in between [1–3]. Some of these species are considered livestock pests,

such as *Lucilia cuprina* (Wiedemann, 1830), *Lucilia sericata* (Meigen, 1826) and *Cochliomyia*

hominivorax Coquerel, 1858 [2,4]. *Lucilia cuprina* is a facultative ectoparasite and one of the

main causative agents of flystrike in sheep in Oceania [5,6]. Conventional insecticides constitute

the current control method for this pest [7], however, an increasing resistance to

organophosphates and benzoylphenyl ureas [7], have motivated the development of alternative

genetic strategies for population control, such as the sterile insect technique (SIT) and female

specific release of insects carrying a dominant female lethal gene (fsRIDL) [8–10].

Flystrikes on live animals are associated with the female oviposition behavior in blowfly

parasitic species [2]. *Lucilia cuprina* females lay eggs in open wounds or moist tissues of host

animals, clustering them in masses of ~200 eggs per ovarian cycle [11,12]. After egg hatching,

the immature larval stages congregate in clusters of maggots called myiasis [12]. Within the

myiasis the larvae complete the three larval (L) stages, from L1 to L3 [12]. This maggot

arrangement favors the action of the larval secretions and excretions (SE) improving larval

feeding and survival [13]. *Lucilia cuprina* larval SE includes several components such as gut

microbiota, lipases, peptidases and ammonia [14,15]. Peptidases and lipases together with the

gut microbiota facilitate host-tissue ingestion and digestion [15]. Ammonia, along with other SE,

has been shown to control deleterious fungi and bacteria that impair larval survival [16].

Furthermore, maggot masses produce heat, reaching up to 18.7°C above ambient temperature

depending on the number of larvae [17]. Temperature is a key factor driving *L. cuprina* larval

development and survival, and larvae are constantly moving in and out of the maggot mass to

avoid overheating [18]. The highest survival rate for this species, considering the whole life

cycle, was observed at 24±1°C [19,20]. Temperatures below and above this range can be

deleterious to *L. cuprina* decreasing larval survivorship [18].

The behavior of *L. cuprina* and other blowflies in locating hosts for oviposition has been

extensively studied [13]. The main attractants to gravid females are sulfur-rich compounds,

ammonia and other volatiles emitted from decomposing organic material, such as meat or feces

[1,13,21]. Dimethyl disulfide (DMDS), dimethyl trisulfide (DMTS), Indole and p-cresol are among

the most relevant blowfly attractants collected from decomposing organic material [22–24]. Adult

gravid females and the last larval stage of *L. cuprina* are more attracted to 5-day-old rotten beef

compared to fresh beef [25,26]. In addition, larvae preferred cold beef (~25°C) rather than warm

(~33°C) [26]. These observations could point to sensory systems conserved between the adult

and larval stages of *L. cuprina*.

RNA-Seq studies have been addressed different larval biological processes in blowfly

species, such as development [27], metabolism [28], aging [29], adaptation to temperature

changes [30,31] and immune response [32]. However, none of these studies focused on larval

sensory receptors potentially associated with physiological and behavioral processes. In

Drosophila melanogaster Meigen, 1830, different sensory receptors, including odorant (ORs)

and gustatory receptors (GRs), have been linked to many larval biological processes such as

chemotaxis, thermotaxis and locomotion [33,34]. For example, *DmelOR42a* and *DmelOR42b*

appear to be important for larval olfaction and locomotion [35–39]. Regarding GRs, such as

DmelGr64a, have been associated with taste [40], whereas others such as *DmelGR28* have

been associated with light avoidance and thermal sensing [41,42]. Ionotropic receptors (IRs)

were also associated with taste and thermal sensing in the *Drosophila* larvae [43,44].

Among other sensory receptors, the transient receptor potential channels (TRPs) and

Pickpocket receptors (PPKs) were identified in the *Drosophila* larval chordotonal organs and

showed a role in proprioception, thermotaxis and locomotion [45–48]. DmelTRPN1 (*nompC*

gene) and other TRPs belonging to the ankyrin and vanilloid subfamilies were associated with

proprioception [48] and thermotaxis [46,47,49]. Among odor-molecule carriers, odorant-binding

proteins (OBPs) and chemosensory proteins (CSPs) were identified in the larva of *D.*

melanogaster [50]. However, the role of these proteins in the larval olfaction and other

physiological aspects has not been extensively studied. Further, the role of other chemosensory

proteins previously studied in adult flies [51–53], such as the superfamily ML (MD-2 (myeloid

differentiation factor-2)-related Lipid-recognition and Niemann-Pick C2 disease proteins

(ML/NPC2), CD36-sensory neuron membrane proteins (CD36/SNMPs), and ammonium

transporters (AMTs), remains to be addressed.

The main objective of the present work was to identify sensory receptors and other

genes relevant to the physiology of *L. cuprina* larvae. We aimed to highlight genes potentially

associated with the processes mentioned above, and other physiological functions such as

feeding, digestion, excretion, diuresis and immune response. Our RNA-Seq analysis of different

larval stages and tissues suggested that the ORs, GRs, IRs and PPKs may not play a central

role in the larval sensory signaling and digestive systems of *L. cuprina*, except for *LcupGR94a*

which was biased to the larval gut. ABC transporters and OBPs were the most relevant within

the sensory-related gene families in terms of number of identified sequences and gene

expression, both in the larval head and across all larval stages. Since the odorant coreceptor

(*Orco*) is a necessary element for the normal functioning of all ORs [54], we tested both adult

females and larvae homozygous for a loss-of-function mutation created by gene editing.

Behavioral assays showed that *LcupOrco* was essential for the female response to rotten beef,

but not for larval behavior. These findings are consistent with high levels of expression of

LcupOrco in the adult female antenna [25] but very low expression in larvae (present work).

Methods

Insect rearing conditions

All larvae and flies used in the present study were of the sub-species *Lucilia cuprina cuprina*,

called "*L. cuprina*" to ease the reading of the manuscript. The *L. cuprina* LA07 wild-type (*wt*)

colony is currently at approximately 225 generations and was established in 2010 using 300

individuals kindly provided by Dr. Aaron Tarone (Texas A&M, TX, USA). The LA07 colony was

established by Dr. Tarone from multiple collections of individuals (300 to 500) from the

University of Southern California campus and the Miracle Mile neighborhood, in Los Angeles,

CA, USA in 2007. All larvae and adult flies used for the experiments were reared using

protocols previously described [9]. Adult flies and immature stages were kept at 23.5±1°C a

non-controlled photoperiod (~13:11 light/dark).

To complete the RNA-Seq analysis, *wt* and heterozygous individuals were used, the

latter obtained from crossing *wt* and SLAM5X colonies. SLAM5X carries an X-linked

constitutively expressed DsRed fluorescent protein gene plus the tetracycline transactivator

coding sequence driven by the early embryo *Chslam* promoter [55]. For the behavioral assays,

LcupOrco larvae or adult females were obtained from crossing heterozygous *LcupOrco*

(*LcupOrco*$^{+/-}$) siblings. The resulting *LcupOrco* mixture was ~ 25% *wt*, 50% *LcupOrco*$^{+/-}$ and 25%

LcupOrco$^{-/-}$. In addition, adult males from the *NPF*$^{-/-}$ colony expressing the ZsGreen were used

for the mating assay.

RNA-Seq experiments

Experimental design and sample collection

Two RNA-Seq experiments, designated EXP-1 and EXP-2 were performed. EXP-1 addressed

gene expression pattern across the three larval development stages using RNA isolated from

whole first, second and late wandering third instar (L1, L2, L3). EXP-2 used RNA isolated from

head, gut and whole body early feeding third instar. For EXP-1 there were five replicates for L1

and L2 and six replicates for L3. Samples one to three of L1 and L2, and all samples for L3,

were *wt* larvae. Samples four and five of L1 and L2 were collected from the *wt*-SLAM5X cross.

Each replicate was a pool including 10 to 15 larvae for L1 and L2, and three pooled individuals

for L3. L1 samples were collected between 20 to 30 hours (h) after egg laying, 50 to 55 h for L2

and between 130 to 135 h for L3 (Fig. S1A-D). All samples were flash frozen using liquid

nitrogen after collection and stored at -80 °C until use.

Samples included in EXP-2 were the first segment of the larva designated as head (H), the gut

including the midgut plus the hindgut and without the crop and Malpighian tubules (MT)

designated as gut (G), and the whole larva (WL) including all tissues (Fig. S1E-G). The crop

was removed from the gut because the presence of meat from the larval diet withing this organ

could bias the analysis. All tissues were collected from *wt* L3s of ~ 75 h after egg laying. Four

replicates were collected for H and G and three for WL, and each replicate was a pool including

50 first larval segments for H, 10 guts for G and five whole larvae per pool for WL. Tissues were

dissected using dissection forceps and vannas scissors (WPI, Worcester, MA, USA), and

immediately after placed in a 1.5 mL Eppendorf tube filled with 300 µl of cold RNAlater^{TM}

(Thermo-Fisher, Waltham, MA, USA, Cat. # AM7020) and stored at -80°C until use.

RNA isolation

RNA extraction of samples associated with EXP-1 and EXP-2 was completed using two

different protocols as follows. Whole larvae of EXP-1 were transferred using dissection forceps

to 2.0 mL tubes prefilled with zirconium beads (Benchmark Scientific, Tempe, AZ, USA, Cat. #

D1032-30). Subsequently, the larvae were resuspended in 250 µl of cold Trizol^{TM} (Thermo-

Fisher, Cat. # 15596026) and disrupted using a benchtop homogenizer (Benchmark Scientific,

Cat. # Z742475). The homogenizer was set to 6.5 m/s and samples were disrupted for 1 min,

another 250 µl of cold Trizol^{TM} were added to the tubes and the homogenization process was

repeated once more. After homogenization, total RNA was extracted using the RNeasy Kit

(Qiagen, Hilden, Germany, Cat. # 74104), and genomic DNA (gDNA) was removed using the

DNase kit (Qiagen, Cat. # 79254) following manufacturer's specifications.

Before RNA extraction of samples belonging to EXP-2, the RNAlater^{TM} was removed

from tubes and tissues were rinsed two times with 1 mL of cold 50% ethanol. After rinsing, pools

of heads, guts and whole larvae were resuspended in 250, 500 and 1000 µl of cold Trizol^{TM}

respectively. Samples were kept on ice and tissues were disrupted using pellet pestles (Bel-Art,

Wayne, NJ, USA, Cat. # F65000-0002) with a homogenizer (Bel-Art, Cat. # F65100-0000) for 90

seconds (sec). Subsequently, samples were centrifuged at 16000 x g and 4°C for 5 min to pellet

part of lipids, cuticle and other debris. After centrifugation the supernatant was transferred to a

clean 1.5 mL tube. Total RNA was extracted using the Quick-RNA^{TM} kit (Zymo Research, Irvine,

CA, USA, Cat. # R1050) according to the manufacturer's specifications. The exception was the

addition of a second deoxyribonuclease 1 (DNase 1) step to ensure the absence of gDNA

contamination in the samples, as follows: 1) during the RNA extraction 30 U of DNAse1 plus 75

µl of digestion buffer (DB) were added to the crude samples followed by 15 min of incubation at

RT; 2) after elution of the clean RNA in nuclease-free (NF) water, another 5 U of DNAse1 + 2.5

µl of DB were added followed by an incubation time same as above. After DNAse1 treatments,

the RNA Clean & Concentrator^{TM}-5 kit (Zymo Research, Cat. # R1013), was used to clean the

samples following the manufacturer's specifications.

Clean RNA belonging to samples from EXP-1 and EXP-2 was eluted in 30 and 12 µl of

NF-water, respectively, and was quantified (1:10 dilution) using a Qubit-4^{TM} fluorometer and the

HS kit (Thermo-Fisher, Cat. # Q33120). Samples were kept at -80°C until use.

RNA-Seq data analysis

Sample quality control, library construction and sequencing services were provided by

Novogene Inc. (Sacramento, CA, USA), following protocols described by the provider

(https://www.novogene.com/us-en/resources/downloads/). The sequencing platform was

Illumina NovaSeq 6000 (Illumina, San Diego, CA, USA), using paired end 150 bp reads, and the

sequencing depth coverage was ~ 50 and 400 million raw reads per library for EXP-1 and EXP-

2, respectively.

The Geneious Prime® software v2023.2.1 (https://www.geneious.com) and associated

plugin packages, were used to complete all the bioinformatic analysis and generate the RNA-

Seq data output. BBDuk plugin v38.88 was utilized to remove Illumina adaptors and trim off low-

quality bases at the 5′ and 3′ends using the following parameters: kmer length: 27; trim both

ends: minimum quality Q20; trim adapters based on paired reads overhangs: minimum overlap

22; discard short reads: minimum length 20 bp. Trimmed reads were mapped to the reference

L. cuprina genome assembly NCBI ID ASM2204524v1, using the default Geneious Prime

mapper set to detect all types of RNA sequences and low-medium sensitivity. A second *L.*

cuprina genome assembly, NCBI ID GCA_001187945.1, was used to analyze specific genes

with annotation errors in the first assembly. Gene expression was calculated only for identified

coding sequence (CDS) and the normalization method used was transcript per million (TPM),

but RPKMs (Reads Per Kilobase per Million mapped reads) FPKMs (Fragments Per Kilobase

per Million mapped fragments) were also provided with the raw data.

A principal components analysis (PCA) was performed to compare EXP-1 libraries with

the R packages FactoMiner [56] and ggcorrplot [57] used for analysis and plotting, respectively.

Gene expression in TPMs was plotted using GraphPad Prism v9 software (San Diego, CA,

USA), showing the mean of each analyzed gene and the standard deviation (SD) between

libraries for the same gene. Differential gene expression between groups was calculated using

DESeq2 [58]. The EXP-1 included five libraries for groups L1 and 2, and six for L3, and EXP-2,

four libraries for groups H and G, and three for WL. A Benjamini-Hochberg (BH) adjusted value

< 0.05 and a FC > 0 were used to define differentially expressed (DE) transcripts between

groups. Sequences with a TPM expression ≥ 5 were separated from all DE transcripts and were

used for all analyses. Transcripts with low expression may not be translated into proteins in the

tissues analyzed or may be associated with pseudo- genes or isoforms [59]. Consequently, DE

ABC and OBP gene families surpassing 5 TPMs were plotted. Results for other sequences

outside this selection were provided as supplementary data tables.

Phylogenetic analysis

The phylogenetic analysis was completed for the ABC and OBP sensory families. ABC protein

sequences were identified by running a NCBI BLASTp against the *L. cuprina* genome assembly,

NCBI ID ASM2204524v1, and using *D. melanogaster* ABC protein sequences as query, the

latter obtained from the FlyBase database (https://flybase.org/). *L. cuprina* OBP curated protein

sequences were obtained from a previous study performed in the same species [25]. Only *L.*

cuprina sensory receptors showing a TPM expression ≥ 5 in the present work were included in

the final phylogenetic analysis. If there was more than one isoform per gene, only isoforms with

different transcript expression were added. Further, OBPs and ABC transporters previously

identified as larval biased in *L. sericata* [30] were added to the analysis.

Prior to phylogenetic analysis, a sequence alignment was performed using the G-INS-i

method in MAFFT v7 [60]. A preliminary phylogenetic analysis including all ABC receptors

identified for *L. cuprina* with a TPM expression > 0, using the Neighbor-Joining (NJ) algorithm

[61] and the software MEGA v11 [62] was performed to rename *L. cuprina* ABCs based on

direct orthology to *D. melanogaster* sequences. Subsequently, the Maximum likelihood (ML)

method set for 1000 bootstrap replications (using the same software) was utilized to improve a

second NJ tree completed including only ABC receptors with a TPM expression ≥ 5. The best

substitution model for each ML run was determined by using the tool "Find Best DNA/Protein

Models" of the same software, prior to running the phylogenetic test. Tree figures were

produced using the interactive Tree of Life (iTOL) v6.8.2 software (https://itol.embl.de/). The

analysis for the OBP sequences was completed using the same protocol except for the

renaming process.

Odorant coreceptor (*Orco*) experiment

Gene editing using CRISPR/Cas9

LcupOrco gene sequence, guide RNA and Cas9 in vitro assay

The guide RNA (gRNA) selected for gene editing was the ortholog of a gRNA previously used to

edit the *ChomOrco* gene in *C. hominivorax* [63]. Note S1 provides further details about the

LcupOrco gene sequence including exonic regions, gRNA position, primers used to amplify the

LcupOrco gene fragment encompassing the Cas9 cutting site, and the Polymerase Chain

Reaction (PCR) cycling parameters. The *LcupOrco* gene model showing the gene editing site

was edited with Adobe Illustrator (San Jose, CA, USA), and the LcupOrco protein secondary

structure generated with the same purpose was plotted with the Protter software

(https://wlab.ethz.ch/protter/start/). The gRNA used was CRISPR-Cas9_crRNA [Integrated DNA

Technologies (IDT), Coralville, IA, USA]. The latter was mixed with the CRISPR-Cas9_tracrRNA

(IDT, Cat. # 1072532), using an equal volume of both RNAs after diluting them to the same

concentration. Subsequently, the mix was heated at 95ºC for 5 min, cooled to ~ 15ºC for 5 min

and kept at -20ºC until use.

The gRNA cutting efficiency was tested *in vitro* using Cas9 protein [New England
Biolabs (NEB), Ipswich, MA, USA, Cat. # M0646T]. Cas9 and gRNA were mixed for a final
concentration of 66.6 and 160 nM, respectively, and incubated for 20 min at 25°C to form the
Cas9 ribonucleoprotein (RNP) complex. A PCR-generated fragment containing the targeted
sequence (Note S1), was added to the mix for a final concentration of 3 nM and incubated for
60 min at 25°C. The reaction (RXN) was ended by adding 1 µl of Proteinase K 10 µg/µl (Zymo,
Cat. # D3001-2-20) and incubated for 10 min. The RXN mix was then loaded into a 1.5%
agarose gel in parallel to uncut gDNA used before as a template (same concentration),
electrophoresed at 60 volts for 90 min and visualize using a Gel Doc™ EZ System (Bio-Rad,
Hercules, CA, USA).
278
Insert construction, CRISPR/Cas9 mix and embryo injections
A Synthetic DNA fragment including two sections of the *LcupOrco* gene, plus restriction cutting
sites for the enzymes XhoI (NEB, Cat. # R0146S) and NotI (NEB, Cat. # R0189S) was acquired
cloned within the plasmid pUCIDT-AMP GoldenGate (IDT). The ZsGreen fluorescent marker
later added to the same construct was obtained and purified from a donor plasmid [64]. The
pUCIDT-AMP GoldenGate plasmid and ZsGreen fragment were ligated using T4 Ligase (NEB,
Cat. # M0202S) and then used for transformation of commercial competent *Escherichia coli*
cells (NEB, Cat. # C3019H) following the manufacturer's specifications. Plasmids were obtained
from 10 single clones using the ZR Miniprep Kit (Zymo, Cat. # D4016) following the
manufacturer's specifications. Subsequently, plasmids were evaluated by restriction enzyme
digestion analyses. Plasmid DNA sequencing was performed for plasmids that showed the
correct banding patterns (MGH CCIB DNA core, Boston, MA). For embryo microinjection,
plasmid DNA was obtained by using a Midiprep Kit (Zymo, Cat. # D4200) and further purified
using the DNA Clean Kit (Zymo, Cat. # D4004) following the manufacturer's specifications. Prior
to injection the DNA was diluted in NF-water to 1250 ng/µl concentration. For further details
refer to Fig. S2A-F.
The Cas9 (IDT, Cat. # 1081058) was used for the *in vivo* embryo injections; the same
was mixed with the duplex crRNA::tracrRNA, 1M Potassium chloride (KCl) and 1 µl of 3.1 buffer
(NEB, Cat. # B6003S) and incubated at 25°C for 20 min to form the RNP complex. The mixture
was cooled to room temperature for 5 min and the plasmid carrying the *LcupOrco* insert was
added to the mix to reach a final plasmid concentration of 500 ng/µl. Final concentrations for the
other compounds were as follows: Cas9 = 3.1 µM (500 ng/µl); crRNA::tracrRNA = 7.5 µM (250

ng/µl) and KCl = 250 mM. The mix was cleaned before injections by 2 min centrifugation at 4°C

using a 0.45 µM filter column (Sigma-Aldrich, Burlington, MS, USA, Cat. # UFC30HV) and keep

on ice all the time.

The needles used for microinjections were generated by pulling filamented quartz

capillaries (Sutter Instrument, Novato, CA, USA, Cat. # QF100-70-10). The puller used was a

Sutter Instrument P-2000 set for: Heat = 690; Filament = 4; Velocity = 60; Delay = 160; Pull =

170. After pulling, needle's tips were beveled for ~ 10 seconds (sec) using a BV-10 micropipette

beveler (Sutter Instrument), a fine abrasive plate and an angle of 22.5°.

Before injections, microinjection mix was loaded into the needles using a Microloader™

(Eppendorf, Hamburg, Germany, Cat. # 930001007). The microinjector used was a XenoWorks

Digital Microinjector (Sutter Instrument), set for continuous flow, 1000 hectopascal (hPa) of

pressure and +5 hPa of transfer pressure, together with a motorized XenoWorks

Micromanipulator (Sutter Instrument) and a stereomicroscope M125 (Leica, Wetzlar, Germany).

Flies used for egg collection were LA07 8-day-old gravid females kept with males of the same

age in a plastic container (Fig. S3A-B). The flies were stimulated to begin ovipositing by adding

93/7% lean-fat fresh grounded beef placed in a 35 mm Petri dish for 15 min. Following, another

35 mm-plate with fresh meat was added to collect new eggs for 5 min. An egg cluster was

transferred to a piece of tissue paper (Kimberly-Clark, Roswell, GA, USA), moistened with

distilled (DI)-water and located in one of the depressions of a double-depression microscope

slide (VWR, Radnor, PA, USA, Cat. # 470003-480). Twenty eggs per round of injections were

separated under an Olympus stereomicroscope SZX10 (Olympus, Center Valley, PA, USA),

using a 00-paint brush and accommodated onto a Scotch® double-sided film tape (3M, Saint

Paul, MN, USA), located in the other depression of the same slide. The eggs were then

desiccated for 5 min using a desiccation chamber of 150 mm in outer diameter (Nalgene,

Rochester, NY, USA) containing a desiccant cartridge (SP Bel-art, Wayne, NJ, USA, Cat. #

F42049-0100). Subsequently eggs were covered with oil (Sigma-Aldrich, Cat. # H8773), rest for

3 min and injected at the posterior end for ~ 1 sec. The developmental time for the injected

embryos was between 18 to 24 min, for the first and last injected embryo, respectively. After

injection, embryos were kept overnight in an incubator chamber (Billups-Rothenberg, Del Mar,

CA, USA, Cat. # MIC-101) filled with oxygen and protected from light using a carboard cover

until eggs hatched, *i.e.* between 19 to 27 h at 23.5°C. After hatching, newly emerged L1s were

transferred using a 00-paint brush to moistened tissue paper (Kimberly-Clark) and then moved

to a 3-ounces clear plastic cup filled with 50 grams of the same meat mentioned above and let

larvae develop until tested at L3 stage.

Fly screening, crosses and genotyping

Injected embryos were screened 7 days after injections (L3 stage) using a stereo microscope

M205 FA (Leica) and a green filter. Only mosaic embryos from generation (G)0 showing

transient expression of the plasmid were retained to obtain adult flies. Emerged adult flies were

crossed following the protocol described in Fig. S4. G1 and G2 eggs obtained from G0 and G1

crosses, respectively, were tested for positive fluorescence using the same microscope, and

only fluorescent eggs were kept.

At G3, a middle leg was collected using dissection forceps and scissors from 2-day-old

flies for genotyping. gDNA was extracted by placing legs in a DNA extraction buffer (Biosearch

Technologies, Hoddesdon, UK, Cat. # QE09050) for 15 min at 65°C and 2 min at 98°C. Three

strategies were used to determine if the *LcupOrco* insert was landed at the correct *L. cuprina*

genome position, *i.e.* withing the *LcupOrco* coding region: 1) amplification of left side of the

insert + upstream genome; 2) amplification of right side of the insert + downstream genome;

and 3) amplification of the whole insert including upstream and downstream genome regions;

for further details refer to Note S2. In parallel, potential indels and point mutations at the Cas9

cutting site were investigated using the same primers described in Note S1. After PCR

amplification, the gDNA was purified using the DNA Clean Kit and sequenced by Sanger

Sequencing. Amplicons >1 Kb were cloned using the pGEM®-T system (Promega, Fitchburg,

WI, USA, Cat. # A1360) together with competent cells following the manufacturer's

specifications. Plasmids were obtained, purified and sequenced following the same protocols

described in Fig. S2.

Behavioral assays

Larval diet preference test

The larval diet preference test was previously used to evaluate *L. cuprina* larvae olfaction

behavior [26]. The larvae used for the assay were obtained using protocols described at the

Insect rearing conditions section. Before the assay, ~ 250 L1s were transferred to 50 g of fresh

ground beef 1 day after egg hatching. Two days after this step a total of 120 3-day-old larvae

were tested following protocols described at [26]. The options offered to the larvae were rotten

(Ro) and fresh (Fr) ground beef, either at 25±1 or 33±1°C, designated as cold (Co) and hot (Ho)

respectively, and for further details about beef conditions refer to [26]. Those larvae that did not

choose any option after 10 min were considered as non-choice (NC). After the assay, larvae

were placed in separated 1.5 tubes filled with 300 µl of DNA preservative buffer (Zymo, Cat. #

R1100-50) and kept at -80°C until DNA extraction. Subsequently, all larvae were dried using

tissue paper and separately weighed using a LA204E analytical balance (Metter Toledo,

Columbus, OH, USA). Following, a small section of each larva was collected using dissection

forceps and scissors and placed back in the same buffer. Approximately one third of the larvae

associated with each diet preference, giving a total of 48 larvae, were genotyped using same

protocols described at the *Fly screening, crosses and genotyping* section and Note S1.

Adult female olfaction assay

The adult female olfaction assay was completed using the same protocol, spatial olfactometer

and room conditions previously used to evaluate *L. cuprina* female olfaction behavior [25]. The

olfactometer consists of four separate chambers (replicates) where the flies were released.

Each of these chambers are connected with two smaller collection chambers A and B, where

meat samples were placed to attract and trap the flies. The attractants used to lure the flies

were 2 g of fresh (Fr) and rotten (Ro) beef at room temperature, placed in chambers A and B,

respectively.

The females used for the olfaction assay were from the same cohort of individuals used

for the larval diet preference test (see above). In addition, LA07 *wt* females were tested using

the same assay conditions, but on different days without mixing them with the *LcupOrco* flies.

All flies used for the olfaction assay were 10-day-old gravid females and were separated from

males the day before the assay using CO_2 to anesthetize the flies under a stereomicroscope

SMZ745 (Nikon, Melville, NY, USA). After sexes separation, females were kept in the same

room where the assay was performed for habituation, provided with tap water and cane sugar.

After the assay, females were kept in separated 1.5 mL empty tubes at -80°C until DNA

extraction. A total of 80 *wt* and 174 *LcupOrco* females were tested, and 72 of the latter,

randomly selected from all assayed *LcupOrco* females, were chosen for genotyping using the

same protocols described in the previous section. After the assay, females were separated into

three groups: 1) attracted to fresh (Fr) beef; 2) attracted to rotten (Ro) beef; and 3) flies that did

not respond to any stimulus, designated as non-choice (NC).

Mating assay

To evaluate the mating performance of the *LcupOrco* homozygous females (*LcupOrco*[-/-]), 5-day-

old females previously determined as heterozygous or homozygous females by genotyping

protocols described above, were mixed with twice as many adult *NPF*[-/-] males of approximate

same age, constitutively expressing the ZsGreen fluorescent protein marker. Fifty grams of

fresh grounded beef was offered O/N to females at 3, 10 and 17 days after mixing them with the

NPF[-/-] males. After oviposition, eggs were visualized and photographed using a stereo

microscope M205 FA (Leica), a green filter and a Leica K5C camera.

Results and Discussion

RNA-Seq overview

To identify genes that are differentially expressed at the very anterior end of the larvae, the first

segment (excluding the brain) was dissected from early feeding third instar (Fig. S1F). RNA was

isolated from gut dissected from the same stage to identify genes important for meat digestion.

Lastly RNA was obtained from whole first, second and third instar (early and late) for reference

for the two tissues, and for a larval developmental series. RNA-Seq output including reads per

library, total number of sequences mapped and subtotals per type of RNA is provided in Table

S1A-B. Considering only transcripts surpassing 5 TPMs, the number of transcripts ranged

between 5,617 and 6,642 for different larval stages and tissues (Data S1A-C and S2A-C).

Further, a PCA analysis showed that samples belonging to each larval stage clustered together,

highlighting the differences between larval stages over those between the samples within each

group (Fig. S5). An RNA-Seq analysis of different larval tissues of *L. sericata* showed a similar

number of transcripts expressed over the same expression threshold, between ~ 4,400 and

6,000 [30]. Transcripts below this expression threshold for each library are provided in Data

S1D-S and S2D-N.

Lucilia cuprina **larval sensory receptors**

ABC transporters and OBPs compromised about 80% of the sensory receptors/carriers showing

a TPM expression ≥ 5, and the number of receptors/carriers per gene family was as follows: 49

ABCs; 25 OBPs; eleven CD36/SNMPs; six ML/NPC2s; three CSPs and one for each of the

following gene families, AMTs, GRs and TRPs (Data S3 and S4). Furthermore, OR, GR, IR and

PPK receptors were not observed above the expression threshold of 5 TPM, except for

LcupGR94a in the larval gut (Data S4A). The expression level of ORs, GRs and IRs suggests

that the same may not have a central role in the sensory system of *L. cuprina* larvae. For

instance, *LcupOrco,* a highly expressed sensory coreceptor in the *L. cuprina* antenna [25]

showed a TPM expression < 0.5 in whole larvae and in head and gut (Table 1). However, in *D.*

melanogaster larvae, *Orco* expression is restricted to the olfactory dorsal organs at the anterior

tip [65] and is essential for chemotaxis [66]. Furthermore, the role of many ORs, GRs and IRs

has been confirmed by *in situ* hybridization and functional assays [33–36,41–44]. Consequently,

more experiments are required to elucidate the role of the specified receptors within the sensory

system of the *L. cuprina* larvae.

The ABC transporters are classified into eight subfamilies [67]. A phylogenetic analysis

showed a possible clade expansion for the subfamily G in *L. cuprina* (Fig. 1 and Fig. S6).

Similar expansions for the same subfamily were observed in hematophagous dipterans [68]. In

L. sericata, orthologs of LcupABCGs 4, 6 and 7 were highly expressed in larval Malpighian

tubules (MT) and fat body (FB) [30]. Here we found that members of this subfamily were highly

enriched in the larval anterior end (see below). In contrast to the apparent expansion of

subfamily G, subfamily A showed a possible clade contraction compared to *D. melanogaster*

(Fig. 1), with only four receptors identified for *L. cuprina* (Data S5B). In line with these findings,

clade contractions and expansions are common for subfamilies A and G along insect species

[69]. Sequences used in the analysis are provided in Data S5A-D; annotation of two sequences

was corrected and transcript expression and differential expression calculated using genome

assembly GCA_001187945.1 (Data S6).

Lcuppixie was the most highly expressed ABC transporter (Data S3A and S4B) and

upregulated in the first two instar compared with third (Fig. 2A). This gene belongs to subfamily

E and is essential for protein translation [70]. Receptors belonging to subfamily F, associated

with protein translation and immune response [71], and B, associated with xenobiotic

detoxification and insecticide resistance [67], were also upregulated in the first two instar larva

vs. L3 (Fig. 2A, Data S3A). No member of these subfamilies was head-biased, but five ABCs of

subfamily B were gut-biased (Fig. 3A-C, Data S4B). Subfamily G showed higher expression in

the latter instars (L2 or L2/L3) compared with the first instar (Fig. 2B-D). In addition, this

subfamily had the highest number of head-biased genes (Fig. 3A and Data S4B), with some

showing a very high enrichment compared with gut or whole body (Data S7A). This subfamily

includes orthologs of the well-studied Drosophila pigmentation genes *white* and *scarlet* [67],

which are L2-biased in *L. cuprina* (Fig. 2B) but do not show any enrichment in head or gut (Fig.

3A-B).

465   Subfamily H was also well represented in the larval head, with the *L. cuprina* ortholog of

466 the Drosophila *oskyddad* (*osy*) showing head-biased expression (Fig. 3A) and the ortholog of

467 *snustorr* (*snu*) enriched in head and gut (Fig. 3C). In *D. melanogaster*, *osy* and *snu* are

468 associated with lipid deposition on the cuticle [72,73]. In the larval gut, the best represented

469 subfamily was C followed by B (Fig. 3B). Both subfamilies share a role in xenobiotics

470 detoxification, multidrug resistance and protein translation [67]. ABC-Cs are also involved in the

471 binding of sulfur-rich compounds [74], and ABC-Bs in ion and lipid transportation [67] and

472 iron/heme metabolism in hematophagous insects [75]. Sulfur-rich compounds are common

473 blowfly attractants [13] and iron and heme are both present in the *L. cuprina* larval meat diet

474 [76]. Other genes engaged in processing these compounds, such as *heme oxygenase 1* and

475 *ferritin* were also highly expressed and biased to the larval gut (Data S2B and S8A-B).

476   The second highest number of chemosensory proteins identified in all *L. cuprina* larval

477 stages and tissue transcriptomes were grouped by OBPs (Data S3B and S4C). This gene family

478 is classified based on a cysteine (C) conserved motif, and modifications of the same change the

479 OBP-ligand binding affinity [54]. The classic subfamily includes a six C-motif and in the Dimer

480 subfamily, the whole motif has been duplicated [54]. Other subfamilies are Minus-C, where the

481 C2 and 5 from the classic motif have been lost, and Plus-C, that includes the classic motif plus

482 three extra conserved Cs and a conserved proline immediately after the Cs [54]. The OBP

483 phylogenetic analysis suggested a clade expansion for *L. cuprina* and *L. sericata* classic OBPs

484 39 to 44 (Fig. 4). The *L. sericata* OBPs included in the expanded clade were highly expressed in

485 the FB followed by the MT [30], suggesting they may be associated with lipid binding and

486 excretion. OBPs grouped within the clade LcupOBPs 39-44 and 49, were the highest expressed

487 genes among the L3 biased (Fig. 5C-E and Data S3B), and head-biased (Fig. 6, Data S4C and

488 S8C-D). Sequences used for the phylogenetic analysis are provided in Data S9A-C.

489   In insects, OBPs are also involved in lipid deposition on the cuticle [77], and some OBPs

490 expressed at the MT and cuticle were associated with xenobiotics detoxification [78–80]. Based

491 on their expression in the *L. cuprina* larvae, OBPs with a lipid biding affinity may be involved in

492 energy homeostasis, avoiding larval desiccation and assisting detoxification. Remarkably, ABC

493 transporters with similar functions, such as receptors from subfamilies G and H, shared a

494 common expression pattern in the *L. cuprina* larvae. Part of the Minus-C and most of the Plus-C

495 OBPs were L1/L2 biased vs. L3 (Fig. 5A-B and Data S3B). The Minus-C *LcupOBP50* and Plus-

496 C *LcupOBP52* were among the most highly expressed sequences within these groups and

497 biased to L1/L2 (Fig. 5A-B and Data S3B). Orthologs of these receptors, namely DmelOBPs

44a and 49a respectively, were studied in *D. melanogaster*. *DmelOBP44a* is involved in lipid

binding [81] and *DmelOBP49a* associated with taste in the same species [82]. Further, Minus-C

and Plus-C OBPs have been associated with binding of sesquiterpenes and derived

sesquiterpenoids such as Juvenile Hormone (JH) in other insects [83–85], and they could have

a similar role in *L. cuprina*. Only a few OBPs were expressed in the larval gut, and none of them

biased to the same tissue (Data S4C). The Minus-C *LcupOBP22* and *LcupOBP24* were among

the highest expressed in the gut (Data S4C). In insect, JH regulates the function of the adult and

larval gut [86,87], and in *D. melanogaster* this hormone is synthesized in the *corpora allata* but

also in the adult gut [86]. Consequently, the above OBPs could be involved with the JH-gut

signaling system.

Lcuppainless was the only TRP above a 5 TPM expression threshold and biased to the

larval head (Data S3C and S4D). In *D. melanogaster* this gene has been associated with taste,

thermal sensing and proprioception [88–91]. Accurate temperature detection is indispensable to

ensuring *L. cuprina* larval survival [17,19], and this receptor may be involved in thermal sensing.

Of the CD36/SNMP genes, the scavenger receptor class B1 (SR-BI, XP_023294991.1) was

upregulated in the L2/3 stages when compared to L1 (Data S3D), and the only member of this

family showing a head bias (Data S4E). Interestingly this gene was highly expressed in the *L.*

cuprina adult antenna [25], indicating a possible role in larval and adult olfaction. Other

members of the same family, *e.g. protein croquemort-like* were biased to the larval gut (Data

S4E). In *D. melanogaster* members of the CD36/SNMP gene family were associated with lipid

binding and their expression in the gut of the same species [92] suggest a role in feeding and

metabolism.

ML/NPC2 transporters have been associated with lipid binding, immune response and

olfaction in different insect species [25,93–95]. Members of this family were highly expressed in

the *L. cuprina* L2 stage and the larval gut (Data S3E and S4F). The *L. cuprina ecdysteroid-*

regulated 16 kDa receptor (*LcupESR16*) was the only head-biased gene within the family (Data

S4F). The same gene was the highest expressed in the *L. cuprina* adult antenna among this

family [25]. Further, *ESR16* orthologs were found expressed in the antenna of many insect

species [25,94,95] suggesting a possible role in olfaction.

Three CSPs were expressed in *L. cuprina* larvae, and two of them were biased to the

head (Data S3F and S4G). In *Bradysia odoriphaga* Winnertz (Diptera: Sciaridae), a CSP protein

was associated with sulfur-rich compounds detection [96], a common blowfly attractant [13]. It is

possible that CSP proteins may have a role in detection of sulfur-rich compounds in *L. cuprina*

larvae. The *ammonium transporter Rhesus type B* (*RHBG*) was biased to the *L. cuprina* L2

stage and the larval gut (Data S3G and S4H). This receptor is involved in ammonia detection

[52,97] and excretion at the MT [98] in *D. melanogaster*. Ammonia-rich compounds are

attractants to adult females of *L. cuprina* and *L. sericata* [13]. In adult antennae, *LcupRHBG* was

highly expressed in both sexes with a female-bias [25]. In the larva of *L. sericata*, *LserRHBG*

showed the highest expression in salivary glands followed by MT [30]. Consequently,

LcupRHBG might be related to olfaction of adult flies, but involved in other processes in

L.cuprina larvae, such as feeding and excretion.

Odorant receptors are not essential for L. cuprina larval response to rotten beef

To determine the importance of ORs in larval behavior, CRISPR/Cas9 was used to obtain a null

mutation of the *LcupOrco* gene following previous protocols used in blowflies [63,99]. We

identified a deletion mutation that introduced a premature stop codon at the end of the first

exon, 265 nucleotides from the start codon (Fig. 7 and Figs. S2-S4, S7-S12). Without the

presence of the odorant coreceptor (*Orco*), no odorant receptor can complete its olfactory

function [54]. Larvae that were a mix of wild-type (*wt*), *LcupOrco* heterozygotes and

homozygotes, were tested for diet preference in a choice assay. Overall, there was no major

change in preference, with larvae preferring rotten beef that was at room temperature as

previously reported [26] (Fig. 8A). No correlation was found between diet preferences and larval

sizes, either for the genotyped (Fig. 8B) or non-genotyped (Fig. 8C) larvae, ANOVA, P = 0.22

and P = 0.25 respectively. This lack of correlation between genotype and diet preference was

confirmed by a Chi-Square analysis, P = 0.66 (Fig. 8D and Table S2A). In the blowfly *C.*

hominivorax, ChomOrco expression was minimal compared to adult flies showing the higher

expression at the L1 stage [63], as observed here for *L. cuprina* larvae (Table 1). The authors

were able to rear a *ChomOrco* silenced strain indicating that this gene may not be essential for

C. hominivorax to complete the larval cycle. Our results suggest that *LcupOrco* is not necessary

for *L. cuprina* larvae to respond to the complex mixture of odors emitted by decomposing beef,

in contrast to observations in *D. melanogaster* larvae, where *Orco* is essential for the response

to odors [33–36].

Disruption of *LcupOrco* affected the *L. cuprina* female oviposition behavior with egg

laying significantly delayed compared to heterozygous siblings (Fig. S11A-G). This was not due

to a feeding disorder as *LcupOrco*$^{-/-}$ homozygotes showed a fully developed ovary (Fig. S12A-

O). Further, a mating assay using transgenic males expressing a green fluorescent protein,

showed that only *LcupOrco* heterozygous females laid eggs the first week after crossing (Fig.

S11A). After a second week, *LcupOrco* silenced females (homozygous) started to lay non-

fertilized eggs (Fig. S11B-C), which did not produce any offspring. *LcupOrco* silenced females

needed a third week to start to lay fertilized eggs giving healthy progeny (Fig. S11D-G). Notably,

LcupOrco silenced males showed an average fertility of 88% including 30 males tested across

four generations. These results were in line with previous observations in *Harpegnathos saltator*

T. C. Jerdon, 1851 (Hymenoptera: Formicidae), where silencing of *Orco* produced males with a

normal mating behavior and females with a delayed and decreased oviposition [100]. However,

silencing of *Orco* in *Helicoverpa armigera* Hübner, 1808 (Lepidoptera: Noctuidae), produced the

opposite phenotype, *i.e.* sterile males and fertile females [101], indicating that the function of

Orco may change among species.

We next asked if *LcupOrco* is important for detecting rotten beef in adult females using a

spatial olfactometer. A mix of *wt*, *LcupOrco* heterozygotes and homozygotes was obtained from

crossing *LcupOrco* heterozygous flies. This mix included an approximate 3:1 ratio of *LcupOrco*

non-silenced to silenced females. Also, a different group of *L. cuprina wt* females were used for

a further comparison with *LcupOrco* females. As previously reported [25], *wt* females were more

attracted to the rotten beef vs. fresh beef, or did not respond to any of the stimulus (non-choice,

NC), ANOVA, P = 0.0001 (Fig. 9A). However, for the *LcupOrco* mix, the number of non-

responding (NC) females was higher than any of other two preferences, ANOVA, P = 0.0001

(Fig. 9B). Genotyping of randomly selected females from tested groups, showed that *LcupOrco*

silenced females were overrepresented among the non-responding flies (NC), Chi-Square, P =

0.0001 (Fig. 9C and Table S2B). These observations showed that *LcupOrco* is essential for

odor detection in adult females, and our results are consistent with previous observations in *C.*

hominivorax, where disruption of *ChomOrco* impaired the foraging and host-seeking behaviors

of adult flies [63].

Non-sensory genes expressed and biased to the larval head

Transcripts encoding the *L. cuprina* orthologs of the *D. melanogaster yellow*, *yellow-h* and

ebony genes*,* with roles in cuticle pigmentation and hardening (tanning) [102–104], were among

the highly expressed head-biased genes vs. the whole body (Data S7A, S8C-D). The *L. cuprina*

eclosion hormone gene (*LcupEH*) was also enriched in the larval head in comparison to the

whole body (Data S7A). In insects, *EH* is associated with cuticle tanning through the control of

Bursicon (*bur*) and *Partner of Bursicon* (*pbur*) [105]. The presence of these transcripts suggests

a role in cuticle and/or mouthparts tanning of the *L. cuprina* larvae. In *D. melanogaster*, *yellow*

controls the tanning of the larval mouth parts [106]. Remarkably, the *tan* gene which display an

antagonistic role to *ebony* in cuticle tanning in *D. melanogaster* [102,107] was downregulated in

the head vs. the whole body (Data S8D).

Transcripts encoding for mucins such as *Lcupmucin-5AC* were highly expressed in the

head compared to whole body, and head-biased (Data S7A and S8C). Mucins have been

studied in the insect gut, playing a role in food processing and supporting the immune system

[108]. Enrichment of mucin transcripts in the larval head could suggest a role at the anterior part

of the digestive system. We previously found that members of this gene family were highly

expressed in the *L. cuprina* adult antenna [25], and suggested that mucins may be important for

olfaction, a role confirmed in vertebrates [109] but not yet addressed in insects. It is possible

that mucins may also be important for olfaction in *L. cuprina* larvae.

Genes potentially associated with L. cuprina larva feeding and digestion

Proteases and lipases such as *trypsin delta* and *lipase 3* respectively, were among the genes

highly enriched in the larval gut compared to whole body and biased to the larval gut of *L.*

cuprina (Data S7B and S8A-B). These findings are consistent with a larval diet composed

mainly of meat with a small fat content [76]. Neuropeptides *Allastostanine*, *CCHamide-2* and

long neuropeptide F (*NPF*), that modulate food intake and digestion in *D. melanogaster*

[110,111], were also expressed and biased to the larval gut (Data S7B and S8A-B).

Neuropeptide *Diuretic hormone class 2* (*DH31*), which play a role in diuresis, gut contractions

and elimination of virulent bacteria in *D. melanogaster* [112] showed the same expression

pattern as the above neuropeptides (Data S7B and S8A-B). In addition, transcripts encoding for

mucins and cytochrome P450s (CYPs) were overrepresented in the larval gut (Data S7B and

S8A-B). CYPs play a crucial role in the insect gut metabolizing endogenous and exogenous

substances [113], and mucins in the aggregation of harmful elements [108]. These enzymes

may be involved together in detoxifying harmful compounds and metabolizing dietary

components in the gut of *L. cuprina* larva.

Gene expression and differential expression across larval stages

The most highly expressed sequences in the whole larva of any stage were genes associated

with cuticle (*e.g. larval cuticle-protein 65Ag1-like, Lcuplcp65Ag1*), muscle formation (*e.g.*

myosin-regulatory light chain 2), and RNA translation factors (Data S1A-C). These transcripts

were also abundant in larval head, but with an overrepresentation of sequences coding for

cuticle proteins and sensory receptors (Data S2A). The larval cuticle is essential for larval life,

acting as both skin and exoskeleton, with the muscles linked directly to the same [114], and the

interaction between cuticle and muscles allows larval locomotion [114]. The cuticle also protects

the larva from desiccation and abrasion, includes sensory and feeding organs and constitutes

the first barrier of the immune system [115].

636   Genes associated with cuticle formation, *e.g. Lcuplcp65Ag1*, showed the highest fold-

change (FC) increases in L2 vs. L1 (Data S10A). Other genes potentially associated with cuticle

tanning such as the orthologs of the *D. melanogaster yellow, yellow-h* and *EH* genes were also

upregulated in L2 vs. L1 (Data S10A). An ortholog of the *Adipokinetic hormone* (*AKH*), which is

involved in mobilizing lipids from storage tissues in different insects [116], showed a high FC

increase in L2 vs. L1 (Data S10A). Other genes involved in protein and fat digestion, such as

Lcuptrypsin-1 and *Lcuplipase 3* respectively, were upregulated in the L2 stage (Data S10A).

Interestingly, orthologs of the antimicrobial peptides *Lucifensin* [117] and *defense protein-*

l(2)34Fc (*l(2)34Fc*) [118], were also increased in L2 vs. L1 (Data S10A). Larvae are reared on

fresh meat under non-sterile conditions [9] and thus could be exposed to a higher bacterial load

at later stages. On the contrary, the expression of orthologs of feeding modulators such as *short*

neuropeptide F (*sNPF*) and *SIFamide* [119,120] was downregulated in L2 vs. L1 (Data S10A).

In *D. melanogaster* larvae, *sNPF* increases food intake but does not prolong the feeding period

or modulate food preferences, unlike *NPF* [120]. Both, *NPF* and *sNPF* are major modulators of

feeding and antimicrobial immune response [121,122].

651   In the late L3, arylphorin genes showed the highest FC increases compared to previous

larval stages (Data S10B-C). In insects, these proteins have a role in storage and cuticle

formation during the prepupal period [123]. Other genes associated with cuticle formation, such

as *Lcuplcp65Ag1*, were also upregulated in L3 (Data S10B-C). Genes potentially involved in

larval energy homeostasis such as *LcupAKH* and *LcupCCHAmide-2*, also showed a biased

expression to L3 (Data S10B-C). In contrast, genes potentially involved in cuticle tanning of *L.*

cuprina larva such as *Lcupyellow* and *LcupEH*, and orthologs of feeding modulators such as

sNPF, NPF and *SIFamide* were downregulated in L3 vs. L1/L2 (Data S10B-C). In addition,

orthologs of the antimicrobial peptides *Lucifensin* and *l(2)34Fc*, and chymotrypsin inhibitors

associated with protease inactivation [124] were also downregulated in L3 vs. L1/L2 (Data

S10B-C).

Conclusions

Based on their expression patterns, ORs, GRs, IRs and PPKs may not play a central role in the *L. cuprina* larval sensory signaling and digestive systems, but further studies are needed to confirm this observation. In contrast, ABCs were highly enriched in head and gut RNA and OBPs only in the head, indicating that both gene families are essential to the physiology of the *L. cuprina* larva. Notably, ABCs were also highly enriched in the *L. cuprina* adult antenna [25], suggesting a role in olfaction across all *L. cuprina* stages for this gene family. Behavioral assays showed that ORs were essential for the adult female response to rotten beef, but not for the larval response. These findings are consistent with high levels of expression of *LcupOrco* in the adult female antenna [25] but very low expression in larvae (present work). Among other relevant genes in *L. cuprina* larvae, the TRP *Lcuppainless* may play a role in environmental sensing such as thermal, taste or proprioception, and the ammonia transporter *LcupRHBG* in food digestion and excretion. Future studies using gene editing techniques in combination with behavioral assays will be conducted to assess the function of the genes highlighted in the present study.

Abbreviations
ABC: ATP-binding cassette transporter; AKH: Adipokinetic hormone: AMT: ammonium transporter; BH: Benjamini-Hochberg; BLAST: basic local alignment search tool; C: cysteine; CDS: coding sequence; CD36/SNMP: CD36 and sensory neuron membrane proteins; CSP: chemosensory protein; Co: cold; CYP: cytochrome P450; DE: differential expressed; DGE: differential gene expression; DH3: diuretic hormone class 2; DNA: deoxyribonucleic acid; DMDS: dimethyl disulfide; DMTS: dimethyl trisulfide; E: exon; EH: eclosion hormone; FB: fat body; FC: Fold-Change; Fr: fresh; G: gut; g: gram; gDNA: genomic DNA; GR: gustatory receptor; H: head; h: hour; Ho: hot; hPa: hectopascal; I: intron; IPM: integrated pest management; IR: ionotropic receptor; iTOL: interactive Tree of Life; JH: juvenile hormone; KB: kilobase; KCl: potassium chloride; L: larva; LS: left side; ML: Maximum likelihood; ML/NPC2: Myeloid lipid-recognition (ML) and Niemann-Pick C2 disease proteins; mRNA: messenger RNA; MT: malpighian tubules; NCBI: national center for biotechnology information; NC: non-choice; NF: nuclease free; NJ: Neighbor-Joining; NPF: long neuropeptide F; OBP: odorant-binding protein; OR: odorant receptor; *Orco*: odorant receptor coreceptor; PCA: principal components analysis; PE: paired end; PPK: pickpocket receptor; RH: relative humidity; RHBG: ammonium transporter Rhesus type B; RNA: Ribonucleic Acid; RNA-Seq: RNA-sequencing; Ro: rotten; RS: right side; RT: room temperature; SD: standard deviation; SE: secretions and excretions; SIT: sterile insect technique; sNPF: short neuropeptide F; SRB1: Scavenger Receptor Class B Type I; TM: transmembrane; TPM: transcripts per million; TRP: Transient receptor potential; UTR: untranslated; WL: whole larva; *wt*: wild-type.

Availability of data and materials
All relevant data are in the manuscript. Raw sequences were uploaded to the NCBI database under Bioproject PRJNA1195419.

705
706 **References**
707
1. Benbow ME, Tomberlin JK, Tarone AM. Carrion Ecology, Evolution, and Their Applications [Internet].

Benbow ME, Tomberlin JK, Tarone AM, editors. Carrion ecology, evolution, and their applications. CRC

Press; 2015. Available from: https://www.taylorfrancis.com/books/9781466575479

2. Wall R, Rose H, Ellse L, Morgan E. Livestock ectoparasites: Integrated management in a changing

climate. Vet Parasitol [Internet]. 2011;180:82–9. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0304401711003839

3. Getz WM. Biomass transformation webs provide a unified approach to consumer-resource modelling.

Ecol Lett [Internet]. 2011;14:113–24. Available from: http://www.ncbi.nlm.nih.gov/pubmed/21199247

4. Mastrangelo T, Welch J. An Overview of the Components of AW-IPM Campaigns against the New

World Screwworm. Insects [Internet]. 2012;3:930–55. Available from: http://www.mdpi.com/2075-

4450/3/4/930

5. Heath ACG, Bishop DM. Flystrike in New Zealand: An overview based on a 16-year study, following the

introduction and dispersal of the Australian sheep blowfly, *Lucilia cuprina* Wiedemann (Diptera:

Calliphoridae). Vet Parasitol [Internet]. 2006;137:333–44. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0304401706000227

6. Wardhaugh KG, Morton R, Bedo D, Horton BJ, Mahon RJ. Estimating the incidence of fly myiases in

Australian sheep flocks: development of a weather-driven regression model. Med Vet Entomol

[Internet]. 2007;21:153–67. Available from:

https://resjournals.onlinelibrary.wiley.com/doi/10.1111/j.1365-2915.2007.00675.x

7. Heath A, Levot G. Parasiticide resistance in flies, lice and ticks in New Zealand and Australia:

mechanisms, prevalence and prevention. N Z Vet J [Internet]. 2015;63:199–210. Available from:

http://www.tandfonline.com/doi/full/10.1080/00480169.2014.960500

8. Sandeman RM, Levot GW, Heath ACG, James PJ, Greeff JC, Scott MJ, et al. Control of the sheep

blowfly in Australia and New Zealand – are we there yet? Int J Parasitol [Internet]. 2014;44:879–91.

Available from: https://linkinghub.elsevier.com/retrieve/pii/S0020751914002318

9. Li F, Wantuch HA, Linger RJ, Belikoff EJ, Scott MJ. Transgenic sexing system for genetic control of the

Australian sheep blow fly *Lucilia cuprina*. Insect Biochem Mol Biol [Internet]. 2014;51:80–8. Available

from: https://linkinghub.elsevier.com/retrieve/pii/S0965174814000952

10. Yan Y, Scott MJ. A transgenic embryonic sexing system for the Australian sheep blow fly *Lucilia*

cuprina. Sci Rep [Internet]. 2015;5:16090. Available from: https://www.nature.com/articles/srep16090

11. Byrd JH, Tomberlin JK. Forensic entomology: the utility of arthropods in legal investigations. CRC

press; 2019.

12. Fenton A, Wall R, French NP. The effects of oviposition aggregation on the incidence of sheep

blowfly strike. Vet Parasitol [Internet]. 1999;83:137–50. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0304401799000473

13. Ashworth JR, Wall R. Responses of the sheep blowflies *Lucilia sericata* and *L. cuprina* to odour and

the development of semiochemical baits. Med Vet Entomol [Internet]. 1994;8:303–9. Available from:

https://resjournals.onlinelibrary.wiley.com/doi/10.1111/j.1365-2915.1994.tb00093.x

14. Lennox FG. Distribution of Ammonia in Larvæ of *Lucilia cuprina*. Nature [Internet]. 1940;146:268–

268. Available from: https://www.nature.com/articles/146268a0

15. Casu RE, Pearson RD, Jarmey JM, Cadogan LC, Riding GA, Tellam RL. Excretory/secretory

chymotrypsin from *Lucilia cuprina*: purification, enzymatic specificity and amino acid sequence deduced

from mRNA. Insect Mol Biol [Internet]. 1994;3:201–11. Available from:

https://resjournals.onlinelibrary.wiley.com/doi/10.1111/j.1365-2583.1994.tb00168.x

16. Arora S, Baptista C, Lim CS. Maggot metabolites and their combinatory effects with antibiotic on

Staphylococcus aureus. Ann Clin Microbiol Antimicrob [Internet]. 2011;10:6. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/21299858

17. Kotzé Z, Villet MH, Weldon CW. Heat accumulation and development rate of massed maggots of the

sheep blowfly, *Lucilia cuprina* (Diptera: Calliphoridae). J Insect Physiol [Internet]. 2016;95:98–104.

Available from: https://linkinghub.elsevier.com/retrieve/pii/S002219101630227X

18. Kotzé Z, Villet MH, Weldon CW. Effect of temperature on development of the blowfly, *Lucilia cuprina*

(Wiedemann) (Diptera: Calliphoridae). Int J Legal Med. 2015;129:1155–62.

19. Bambaradeniya YTB, Karunaratne WAIP, Tomberlin JK, Goonerathne I, Kotakadeniya RB.

Temperature and tissue type impact development of *Lucilia cuprina* (Diptera: Calliphoridae) in Sri Lanka.

J Med Entomol. 2018;55:285–91.

20. Kotze A, James P. Control of sheep flystrike: what's been tried in the past and where to from here.

Aust Vet J [Internet]. 2022;100:1–19. Available from:

https://onlinelibrary.wiley.com/doi/10.1111/avj.13131

21. Forbes SL, Perrault KA. Decomposition Odour Profiling in the Air and Soil Surrounding Vertebrate

Carrion. Hansen IA, editor. PLoS One [Internet]. 2014;9:e95107. Available from:

https://dx.plos.org/10.1371/journal.pone.0095107

22. Zhu JJ, Chaudhury MF, Tangtrakulwanich K, Skoda SR. Identification of Oviposition Attractants of the

Secondary Screwworm, *Cochliomyia macellaria* (F.) Released from Rotten Chicken Liver. J Chem Ecol

[Internet]. 2013;39:1407–14. Available from: http://link.springer.com/10.1007/s10886-013-0359-z

23. Chaudhury MF, Skoda SR, Sagel A, Welch JB. Volatiles Emitted From Eight Wound-Isolated Bacteria

Differentially Attract Gravid Screwworms (Diptera: Calliphoridae) to Oviposit. J Med Entomol [Internet].

2010;47:349–54. Available from: http://openurl.ingenta.com/content/xref?genre=article&issn=0022-

2585&volume=47&issue=3&spage=349

24. Liu W, Longnecker M, Tarone AM, Tomberlin JK. Responses of *Lucilia sericata* (Diptera: Calliphoridae)

to compounds from microbial decomposition of larval resources. Anim Behav [Internet]. 2016;115:217–

25. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0003347216001056

25. Wulff JP, Hickner P V., Watson DW, Denning SS, Belikoff EJ, Scott MJ. Antennal transcriptome

analysis reveals sensory receptors potentially associated with host detection in the livestock pest *Lucilia*

cuprina. Parasit Vectors [Internet]. 2024;17:308. Available from:

https://parasitesandvectors.biomedcentral.com/articles/10.1186/s13071-024-06391-6

26. Cunha VAS, Tandonnet S, Ferreira DL, Rodrigues A V., Torres TT. Exploring Life History Choices: Using

Temperature and Substrate Type as Interacting Factors for Blowfly Larval and Female Preferences.

Journal of Visualized Experiments [Internet]. 2023;e65835. Available from:

https://app.jove.com/t/65835

27. Guerrero FD, Dowd SE, Djikeng A, Wiley G, Macmil S, Saldivar L, et al. A Database of Expressed Genes

From *Cochliomyia hominivorax* (Diptera: Calliphoridae). J Med Entomol [Internet]. 2009;46:1109–16.

Available from: https://academic.oup.com/jme/article-lookup/doi/10.1603/033.046.0518

28. Zhang M, Yu H, Yang Y, Song C, Hu X, Zhang G. Analysis of the Transcriptome of Blowfly *Chrysomya*

megacephala (Fabricius) Larvae in Responses to Different Edible Oils. Zhou D, editor. PLoS One

[Internet]. 2013;8:e63168. Available from: https://dx.plos.org/10.1371/journal.pone.0063168

29. Hjelmen CE, Yuan Y, Parrott JJ, McGuane AS, Srivastav SP, Purcell AC, et al. Identification and

Characterization of Small RNA Markers of Age in the Blow Fly *Cochliomyia macellaria* (Fabricius)

(Diptera: Calliphoridae). Insects [Internet]. 2022;13:948. Available from: https://www.mdpi.com/2075-

4450/13/10/948

30. Davis RJ, Belikoff EJ, Dickey AN, Scholl EH, Benoit JB, Scott MJ. Genome and transcriptome

sequencing of the green bottle fly, *Lucilia sericata*, reveals underlying factors of sheep flystrike and

maggot debridement therapy. Genomics [Internet]. 2021;113:3978–88. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0888754321003669

31. Liu Z, Han H, Meng F, Jiang Y, Cai J. Dynamic transcriptome profiling exploring cold tolerance in

forensically important blow fly, *Aldrichina grahami* (Diptera: Calliphoridae). BMC Genomics [Internet].

2020;21:92. Available from: http://www.ncbi.nlm.nih.gov/pubmed/31996132

32. McKenna CH, Asgari D, Crippen TL, Zheng L, Sherman RA, Tomberlin JK, et al. Gene expression in

Lucilia sericata (Diptera: Calliphoridae) larvae exposed to *Pseudomonas aeruginosa* and *Acinetobacter*

baumannii identifies shared and microbe-specific induction of immune genes. Insect Mol Biol [Internet].

2022;31:85–100. Available from: https://resjournals.onlinelibrary.wiley.com/doi/10.1111/imb.12740

33. Fishilevich E, Domingos AI, Asahina K, Naef F, Vosshall LB, Louis M. Chemotaxis Behavior Mediated

by Single Larval Olfactory Neurons in *Drosophila*. Current Biology [Internet]. 2005;15:2086–96. Available

from: https://linkinghub.elsevier.com/retrieve/pii/S0960982205013783

34. Gerber B, Stocker RF. The *Drosophila* Larva as a Model for Studying Chemosensation and

Chemosensory Learning: A Review. Chem Senses [Internet]. 2007;32:65–89. Available from:

https://academic.oup.com/chemse/article-lookup/doi/10.1093/chemse/bjl030

35. Gomez-Marin A, Louis M. Multilevel control of run orientation in *Drosophila* larval chemotaxis. Front

Behav Neurosci. 2014;8.

36. Gomez-Marin A, Stephens GJ, Louis M. Active sampling and decision making in *Drosophila*

chemotaxis. Nat Commun [Internet]. 2011;2:441. Available from:

https://www.nature.com/articles/ncomms1455

37. Louis M, Huber T, Benton R, Sakmar TP, Vosshall LB. Bilateral olfactory sensory input enhances

chemotaxis behavior. Nat Neurosci [Internet]. 2008;11:187–99. Available from:

https://www.nature.com/articles/nn2031

38. Kreher SA, Mathew D, Kim J, Carlson JR. Translation of Sensory Input into Behavioral Output via an

Olfactory System. Neuron [Internet]. 2008;59:110–24. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0896627308005035

39. Grillet M, Campagner D, Petersen R, McCrohan C, Cobb M. The peripheral olfactory code in

Drosophila larvae contains temporal information and is robust over multiple timescales. Proceedings of

the Royal Society B: Biological Sciences [Internet]. 2016;283:20160665. Available from:

https://royalsocietypublishing.org/doi/10.1098/rspb.2016.0665

40. Jiao Y, Moon SJ, Montell C. A *Drosophila* gustatory receptor required for the responses to sucrose,

glucose, and maltose identified by mRNA tagging. Proceedings of the National Academy of Sciences

[Internet]. 2007;104:14110–5. Available from: https://pnas.org/doi/full/10.1073/pnas.0702421104

41. Ni L, Bronk P, Chang EC, Lowell AM, Flam JO, Panzano VC, et al. A gustatory receptor paralogue

controls rapid warmth avoidance in *Drosophila*. Nature. 2013;500:580–4.

42. Sang J, Rimal S, Lee Y. *Gustatory receptor 28b* is necessary for avoiding saponin in *Drosophila*

melanogaster. EMBO Rep [Internet]. 2019;20. Available from:

https://www.embopress.org/doi/10.15252/embr.201847328

43. Ni L, Klein M, Svec K V, Budelli G, Chang EC, Ferrer AJ, et al. The Ionotropic Receptors IR21a and

IR25a mediate cool sensing in *Drosophila*. Elife [Internet]. 2016;5. Available from:

https://elifesciences.org/articles/13254

44. Stewart S, Koh T-W, Ghosh AC, Carlson JR. Candidate ionotropic taste receptors in the *Drosophila*

larva. Proceedings of the National Academy of Sciences [Internet]. 2015;112:4195–201. Available from:

https://pnas.org/doi/full/10.1073/pnas.1503292112

45. Ainsley JA, Pettus JM, Bosenko D, Gerstein CE, Zinkevich N, Anderson MG, et al. Enhanced

Locomotion Caused by Loss of the *Drosophila* DEG/ENaC Protein Pickpocket1. Current Biology [Internet].

2003;13:1557–63. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0960982203005967

46. Klein M, Afonso B, Vonner AJ, Hernandez-Nunez L, Berck M, Tabone CJ, et al. Sensory determinants

of behavioral dynamics in *Drosophila* thermotaxis. Proceedings of the National Academy of Sciences

[Internet]. 2015;112:E220–9. Available from: https://pnas.org/doi/full/10.1073/pnas.1416212112

47. Kwon Y, Shen WL, Shim H-S, Montell C. Fine Thermotactic Discrimination between the Optimal and

Slightly Cooler Temperatures via a TRPV Channel in Chordotonal Neurons. The Journal of Neuroscience

[Internet]. 2010;30:10465–71. Available from:

https://www.jneurosci.org/lookup/doi/10.1523/JNEUROSCI.1631-10.2010

48. Walker RG, Willingham AT, Zuker CS. A *Drosophila* Mechanosensory Transduction Channel. Science

(1979) [Internet]. 2000;287:2229–34. Available from:

https://www.science.org/doi/10.1126/science.287.5461.2229

49. Pulver SR, Pashkovski SL, Hornstein NJ, Garrity PA, Griffith LC. Temporal Dynamics of Neuronal

Activation by Channelrhodopsin-2 and TRPA1 Determine Behavioral Output in *Drosophila* Larvae. J

Neurophysiol [Internet]. 2009;101:3075–88. Available from:

https://www.physiology.org/doi/10.1152/jn.00071.2009

50. Rihani K, Ferveur J-F, Briand L. The 40-Year Mystery of Insect Odorant-Binding Proteins.

Biomolecules [Internet]. 2021;11:509. Available from: https://www.mdpi.com/2218-273X/11/4/509

51. Leitch O, Papanicolaou A, Lennard C, Kirkbride KP, Anderson A. Chemosensory genes identified in the

antennal transcriptome of the blowfly *Calliphora stygia*. BMC Genomics [Internet]. 2015;16:255.

Available from: https://bmcgenomics.biomedcentral.com/articles/10.1186/s12864-015-1466-8

52. Vulpe A, Kim HS, Ballou S, Wu S-T, Grabe V, Nava Gonzales C, et al. An ammonium transporter is a

non-canonical olfactory receptor for ammonia. Current Biology [Internet]. 2021;31:3382-3390.e7.

Available from: https://linkinghub.elsevier.com/retrieve/pii/S0960982221007144

53. Rondón JJ, Moreyra NN, Pisarenco VA, Rozas J, Hurtado J, Hasson E. Evolution of the odorant-binding

protein gene family in *Drosophila*. Front Ecol Evol [Internet]. 2022;10. Available from:

https://www.frontiersin.org/articles/10.3389/fevo.2022.957247/full

54. Wicher D, Marion-Poll F. Editorial: Function and Regulation of Chemoreceptors. Wicher D, Marion-

Poll F, editors. Front Cell Neurosci [Internet]. 2018;12. Available from:

https://www.frontiersin.org/article/10.3389/fncel.2018.00496/full

55. Williamson ME, Yan Y, Scott MJ. Conditional knockdown of transformer in sheep blow fly suggests a

role in repression of dosage compensation and potential for population suppression. PLoS Genet

[Internet]. 2021;17:e1009792. Available from: http://www.ncbi.nlm.nih.gov/pubmed/34662332

56. Lê S, Josse J, Husson F. **FactoMineR**: An *R* Package for Multivariate Analysis. J Stat Softw [Internet].

2008;25. Available from: http://www.jstatsoft.org/v25/i01/

57. Valero-Mora PM. **ggplot2:** Elegant Graphics for Data Analysis. J Stat Softw [Internet]. 2010;35.

Available from: http://www.jstatsoft.org/v35/b01/

58. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data

with DESeq2. Genome Biol. 2014;15.

59. Mika K, Benton R. Olfactory Receptor Gene Regulation in Insects: Multiple Mechanisms for Singular

Expression. Front Neurosci. 2021;15:738088.

60. Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7: improvements in

performance and usability. Mol Biol Evol [Internet]. 2013;30:772–80. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/23329690

61. Saitou N, Nei M. The neighbor-joining method: a new method for reconstructing phylogenetic trees.

Mol Biol Evol [Internet]. 1987; Available from:

https://academic.oup.com/mbe/article/4/4/406/1029664/The-

62. Tamura K, Stecher G, Kumar S. MEGA11: molecular evolutionary genetics analysis version 11. Mol

Biol Evol. 2021;38:3022–7.

63. Paulo DF, Junqueira ACM, Arp AP, Vieira AS, Ceballos J, Skoda SR, et al. Disruption of the odorant

coreceptor *Orco* impairs foraging and host finding behaviors in the New World screwworm fly. Sci Rep

[Internet]. 2021;11:11379. Available from: https://www.nature.com/articles/s41598-021-90649-x

64. Concha C, Belikoff EJ, Carey B, Li F, Schiemann AH, Scott MJ. Efficient germ-line transformation of the

economically important pest species *Lucilia cuprina* and *Lucilia sericata* (Diptera, Calliphoridae). Insect

Biochem Mol Biol [Internet]. 2011;41:70–5. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0965174810002122

65. Task D, Lin C-C, Vulpe A, Afify A, Ballou S, Brbic M, et al. Chemoreceptor co-expression in *Drosophila*

melanogaster olfactory neurons. Elife [Internet]. 2022;11. Available from:

https://elifesciences.org/articles/72599

66. Utashiro N, Williams CR, Parrish JZ, Emoto K. Prior activity of olfactory receptor neurons is required

for proper sensory processing and behavior in *Drosophila* larvae. Sci Rep [Internet]. 2018;8:8580.

Available from: https://www.nature.com/articles/s41598-018-26825-3

67. Dermauw W, Van Leeuwen T. The ABC gene family in arthropods: Comparative genomics and role

in insecticide transport and resistance. Insect Biochem Mol Biol [Internet]. 2014;45:89–110. Available

from: https://linkinghub.elsevier.com/retrieve/pii/S0965174813001951

68. Kumar V, Garg S, Gupta L, Gupta K, Diagne CT, Missé D, et al. Delineating the Role of *Aedes aegypti*

ABC Transporter Gene Family during Mosquito Development and Arboviral Infection via Transcriptome

Analyses. Pathogens [Internet]. 2021;10:1127. Available from: https://www.mdpi.com/2076-

0817/10/9/1127

69. Wu C, Chakrabarty S, Jin M, Liu K, Xiao Y. Insect ATP-Binding Cassette (ABC) Transporters: Roles in

Xenobiotic Detoxification and Bt Insecticidal Activity. Int J Mol Sci [Internet]. 2019;20:2829. Available

from: https://www.mdpi.com/1422-0067/20/11/2829

70. Andersen DS, Leevers SJ. The Essential *Drosophila* ATP-binding Cassette Domain Protein, Pixie, Binds

the 40 S Ribosome in an ATP-dependent Manner and Is Required for Translation Initiation. Journal of

Biological Chemistry [Internet]. 2007;282:14752–60. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0021925820636274

71. Fostier CR, Monlezun L, Ousalem F, Singh S, Hunt JF, Boël G. ABC-F translation factors: from

antibiotic resistance to immune response. FEBS Lett [Internet]. 2021;595:675–706. Available from:

https://febs.onlinelibrary.wiley.com/doi/10.1002/1873-3468.13984

72. Wang Y, Norum M, Oehl K, Yang Y, Zuber R, Yang J, et al. Dysfunction of *Oskyddad* causes Harlequin-

type ichthyosis-like defects in *Drosophila melanogaster*. Stern DL, editor. PLoS Genet [Internet].

2020;16:e1008363. Available from: https://dx.plos.org/10.1371/journal.pgen.1008363

73. Zuber R, Norum M, Wang Y, Oehl K, Gehring N, Accardi D, et al. The ABC transporter Snu and the

extracellular protein Snsl cooperate in the formation of the lipid-based inward and outward barrier in

the skin of *Drosophila*. Eur J Cell Biol [Internet]. 2018;97:90–101. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0171933517301851

74. Abo-Elghar GE, Fujiyoshi P, Matsumura F. Significance of the sulfonylurea receptor (SUR) as the

target of diflubenzuron in chitin synthesis inhibition in *Drosophila melanogaster* and *Blattella*

germanica. Insect Biochem Mol Biol [Internet]. 2004;34:743–52. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0965174804000499

75. Whiten SR, Eggleston H, Adelman ZN. Ironing out the Details: Exploring the Role of Iron and Heme in

Blood-Sucking Arthropods. Front Physiol [Internet]. 2018;8. Available from:

http://journal.frontiersin.org/article/10.3389/fphys.2017.01134/full

76. Selem GSh, Geden CJ, Khater H, Khater KS. Effects of larval diets on some biological parameters and

morphometric and biochemical analysis of ovaries of *Lucilia cuprina* (Wiedemann) (Diptera:

Calliphoridae). Journal of Vector Ecology [Internet]. 2023;48. Available from:

https://bioone.org/journals/journal-of-vector-ecology/volume-48/issue-2/1081-1710-48.2.72/Effects-

of-larval-diets-on-some-biological-parameters-and-morphometric/10.52707/1081-1710-48.2.72.full

77. Lockey KH. Lipids of the insect cuticle: origin, composition and function. Comparative Biochemistry

and Physiology Part B: Comparative Biochemistry [Internet]. 1988;89:595–645. Available from:

https://linkinghub.elsevier.com/retrieve/pii/0305049188903057

78. Zhang Y, Gao S, Xue S, Zhang K, Wang J, Li B. Odorant-Binding Proteins Contribute to the Defense of

the Red Flour Beetle, *Tribolium castaneum*, Against Essential Oil of *Artemisia vulgaris*. Front Physiol

[Internet]. 2020;11. Available from: https://www.frontiersin.org/article/10.3389/fphys.2020.00819/full

79. Abendroth JA, Moural TW, Wei H, Zhu F. Roles of insect odorant binding proteins in communication

and xenobiotic adaptation. Frontiers in Insect Science [Internet]. 2023;3. Available from:

https://www.frontiersin.org/articles/10.3389/finsc.2023.1274197/full

80. Zhao R, Wang H-H, Wang Z, Xiao X, Yin X-H, Hu S-Y, et al. Omics Analysis of Odorant-Binding Proteins

and Cuticle-Enriched *SfruOBP18* Confers Multi-Insecticide Tolerance in *Spodoptera frugiperda*. J Agric

Food Chem [Internet]. 2024;72:22532–44. Available from:

https://pubs.acs.org/doi/10.1021/acs.jafc.4c05737

81. Cotten ML, Starich MR, He Y, Yin J, Yuan Q, Tjandra N. NMR chemical shift assignment of *Drosophila*

odorant binding protein 44a in complex with 8(Z)-eicosenoic acid. Biomol NMR Assign [Internet]. 2024;

Available from: https://link.springer.com/10.1007/s12104-024-10178-2

82. Jeong YT, Shim J, Oh SR, Yoon HI, Kim CH, Moon SJ, et al. An Odorant-Binding Protein Required for

Suppression of Sweet Taste by Bitter Chemicals. Neuron [Internet]. 2013;79:725–37. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0896627313005412

83. González-González A, Palma-Millanao R, Yáñez O, Rojas M, Mutis A, Venthur H, et al. Virtual

Screening of Plant Volatile Compounds Reveals a High Affinity of *Hylamorpha elegans* (Coleoptera:

Scarabaeidae) Odorant-Binding Proteins for Sesquiterpenes From Its Native Host. Journal of Insect

Science [Internet]. 2016;16:30. Available from: https://academic.oup.com/jinsectscience/article-

lookup/doi/10.1093/jisesa/iew008

84. Giordano D, Facchiano A, D'Auria S, Loreto F. A hypothesis on the capacity of plant odorant-binding

proteins to bind volatile isoprenoids based on in silico evidences. Elife [Internet]. 2021;10. Available

from: https://elifesciences.org/articles/66741

85. Qiao H, Tuccori E, He X, Gazzano A, Field L, Zhou J-J, et al. Discrimination of alarm pheromone (E)-β-

farnesene by aphid odorant-binding proteins. Insect Biochem Mol Biol [Internet]. 2009;39:414–9.

Available from: https://linkinghub.elsevier.com/retrieve/pii/S0965174809000551

86. Rahman MM, Franch-Marro X, Maestro JL, Martin D, Casali A. Local Juvenile Hormone activity

regulates gut homeostasis and tumor growth in adult *Drosophila*. Sci Rep [Internet]. 2017;7:11677.

Available from: https://www.nature.com/articles/s41598-017-11199-9

87. Taracena-Agarwal ML, Walter-Nuno AB, Bottino-Rojas V, Mejia APG, Xu K, Segal S, et al. Juvenile

Hormone as a contributing factor in establishing midgut microbiota for fecundity and fitness

enhancement in adult female *Aedes aegypti*. Commun Biol [Internet]. 2024;7:687. Available from:

https://www.nature.com/articles/s42003-024-06334-y

88. Sokabe T, Tominaga M. A temperature-sensitive TRP ion channel, *painless*, functions as a noxious

heat sensor in fruit flies. Commun Integr Biol [Internet]. 2009;2:170–3. Available from:

http://www.tandfonline.com/doi/abs/10.4161/cib.7708

89. Charlu S, Wisotsky Z, Medina A, Dahanukar A. Acid sensing by sweet and bitter taste neurons in

Drosophila melanogaster. Nat Commun [Internet]. 2013;4:2042. Available from:

https://www.nature.com/articles/ncomms3042

90. Al-Anzi B, Tracey WD, Benzer S. Response of *Drosophila* to Wasabi Is Mediated by painless, the Fly

Homolog of Mammalian TRPA1/ANKTM1. Current Biology [Internet]. 2006;16:1034–40. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S096098220601428X

91. Xu J, Sornborger AT, Lee JK, Shen P. *Drosophila* TRPA channel modulates sugar-stimulated neural

excitation, avoidance and social response. Nat Neurosci [Internet]. 2008;11:676–82. Available from:

https://www.nature.com/articles/nn.2119

92. Herboso L, Talamillo A, Perez C, Barrio R. Expression of the Scavenger Receptor Class B type I (SR-BI)

family in *Drosophila melanogaster*. Int J Dev Biol [Internet]. 2011;55:603–11. Available from:

http://www.intjdevbiol.com/paper.php?doi=103254lh

93. Shi X-Z, Zhong X, Yu X-Q. *Drosophila melanogaster* NPC2 proteins bind bacterial cell wall components

and may function in immune signal pathways. Insect Biochem Mol Biol [Internet]. 2012;42:545–56.

Available from: https://linkinghub.elsevier.com/retrieve/pii/S0965174812000513

94. Chen M-M, Zhong L, Zhao C-S, Wang F-C, Ji W-J, Zhang B, et al. Characterization of an Ecdysteroid-

Regulated 16 kDa Protein Gene in Chinese Oak Silkworm, *Antheraea pernyi* (Lepidoptera: Saturniidae).

Swevers L, editor. Journal of Insect Science [Internet]. 2020;20. Available from:

https://academic.oup.com/jinsectscience/article/doi/10.1093/jisesa/ieaa033/5836191

95. Wulff JP, Traverso LM, Latorre-Estivalis JM, Segura DF, Lanzavecchia SB. Identification of candidate

genes associated with host-seeking behavior in the parasitoid wasp *Diachasmimorpha longicaudata*.

BMC Genomics [Internet]. 2024;25:147. Available from:

https://bmcgenomics.biomedcentral.com/articles/10.1186/s12864-024-10034-6

96. Yang Y, Hua D, Zhu J, Wang F, Zhang Y. Chemosensory protein 4 is required for *Bradysia odoriphaga*

to be olfactory attracted to sulfur compounds released from Chinese chives. Front Physiol [Internet].

2022;13. Available from: https://www.frontiersin.org/articles/10.3389/fphys.2022.989601/full

97. Delventhal R, Menuz K, Joseph R, Park J, Sun JS, Carlson JR. The taste response to ammonia in

Drosophila. Sci Rep [Internet]. 2017;7:43754. Available from:

https://www.nature.com/articles/srep43754

98. Browne A, O'Donnell MJ. Ammonium secretion by Malpighian tubules of *Drosophila melanogaster*:

application of a novel ammonium-selective microelectrode. Journal of Experimental Biology [Internet].

2013;216:3818–27. Available from:

https://journals.biologists.com/jeb/article/doi/10.1242/jeb.091082/257472/Ammonium-secretion-by-

Malpighian-tubules-of

99. Paulo DF, Williamson ME, Scott MJ. CRISPR/Cas9 Genome Editing in the New World Screwworm and

Australian Sheep Blowfly. 2022. p. 173–201. Available from: https://link.springer.com/10.1007/978-1-

0716-2301-5_10

100. Yan H, Opachaloemphan C, Mancini G, Yang H, Gallitto M, Mlejnek J, et al. An Engineered *Orco*

Mutation Produces Aberrant Social Behavior and Defective Neural Development in Ants. Cell [Internet].

2017;170:736-747.e9. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0092867417307705

101. Fan X-B, Mo B-T, Li G-C, Huang L-Q, Guo H, Gong X-L, et al. Mutagenesis of the odorant receptor co-

receptor (*Orco*) reveals severe olfactory defects in the crop pest moth *Helicoverpa armigera*. BMC Biol

[Internet]. 2022;20:214. Available from: https://bmcbiol.biomedcentral.com/articles/10.1186/s12915-

022-01411-2

102. Lamb AM, Wang Z, Simmer P, Chung H, Wittkopp PJ. *ebony* Affects Pigmentation Divergence and

Cuticular Hydrocarbons in *Drosophila americana* and *D. novamexicana*. Front Ecol Evol [Internet].

2020;8. Available from: https://www.frontiersin.org/article/10.3389/fevo.2020.00184/full

103. Drapeau MD, Radovic A, Wittkopp PJ, Long AD. A gene necessary for normal male courtship, *yellow*,

acts downstream of *fruitless* in the *Drosophila melanogaster* larval brain. J Neurobiol [Internet].

2003;55:53–72. Available from: https://onlinelibrary.wiley.com/doi/10.1002/neu.10196

104. Barek H, Zhao H, Heath K, Veraksa A, Sugumaran M. *Drosophila yellow-h* encodes dopaminechrome

tautomerase: A new enzyme in the eumelanin biosynthetic pathway. Pigment Cell Melanoma Res

[Internet]. 2022;35:26–37. Available from: https://onlinelibrary.wiley.com/doi/10.1111/pcmr.13008

105. Song Q, An S. Bursicon, a Neuropeptide Hormone That Controls Cuticle Tanning and Beyond. In: Liu

T, Kang L, editors. Recent Advances in Entomological Research [Internet]. Berlin, Heidelberg: Springer

Berlin Heidelberg; 2011. p. 132–49. Available from: http://link.springer.com/10.1007/978-3-642-17815-

3_8

106. Chia W, Howes G, Martin M, Meng YB, Moses K, Tsubota S. Molecular analysis of the *yellow* locus

of *Drosophila*. EMBO J [Internet]. 1986;5:3597–605. Available from:

https://onlinelibrary.wiley.com/doi/10.1002/j.1460-2075.1986.tb04688.x

107. Richardt A, Kemme T, Wagner S, Schwarzer D, Marahiel MA, Hovemann BT. Ebony, a Novel

Nonribosomal Peptide Synthetase for β-Alanine Conjugation with Biogenic Amines in *Drosophila*. Journal

of Biological Chemistry [Internet]. 2003;278:41160–6. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0021925820829159

108. Khan SA, Kojour MAM, Han YS. Recent trends in insect gut immunity. Front Immunol [Internet].

2023;14:1272143. Available from: http://www.ncbi.nlm.nih.gov/pubmed/38193088

109. Kennel C, Gould EA, Larson ED, Salcedo E, Vickery T, Restrepo D, et al. Differential Expression of

Mucins in Murine Olfactory Versus Respiratory Epithelium. Chem Senses [Internet]. 2019;44:511–21.

Available from: https://academic.oup.com/chemse/article/44/7/511/5531912

110. Sano H. Coupling of growth to nutritional status: The role of novel periphery-to-brain signaling by

the CCHa2 peptide in *Drosophila melanogaster*. Fly (Austin) [Internet]. 2015;9:183–7. Available from:

http://www.tandfonline.com/doi/full/10.1080/19336934.2016.1162361

111. Ren GR, Hauser F, Rewitz KF, Kondo S, Engelbrecht AF, Didriksen AK, et al. CCHamide-2 Is an

Orexigenic Brain-Gut Peptide in *Drosophila*. Blenau W, editor. PLoS One [Internet]. 2015;10:e0133017.

Available from: https://dx.plos.org/10.1371/journal.pone.0133017

112. Benguettat O, Jneid R, Soltys J, Loudhaief R, Brun-Barale A, Osman D, et al. The DH31/CGRP

enteroendocrine peptide triggers intestinal contractions favoring the elimination of opportunistic

bacteria. Talavera K, editor. PLoS Pathog [Internet]. 2018;14:e1007279. Available from:

https://dx.plos.org/10.1371/journal.ppat.1007279

113. Zhang X, Shi Z, Yang C, Li J, Liu J, Zhang A. Gut transcriptome analysis of P450 genes and cytochrome

P450 reductase in three moth species feeding on gymnosperms or angiosperms. Front Ecol Evol

[Internet]. 2022;10. Available from: https://www.frontiersin.org/articles/10.3389/fevo.2022.948043/full

114. Lahiri S, Shen K, Klein M, Tang A, Kane E, Gershow M, et al. Two Alternating Motor Programs Drive

Navigation in *Drosophila* Larva. Skoulakis EMC, editor. PLoS One [Internet]. 2011;6:e23180. Available

from: https://dx.plos.org/10.1371/journal.pone.0023180

115. Ferveur J-F, Cortot J, Rihani K, Cobb M, Everaerts C. Desiccation resistance: effect of cuticular

hydrocarbons and water content in *Drosophila melanogaster* adults. PeerJ [Internet]. 2018;6:e4318.

Available from: https://peerj.com/articles/4318

116. Van der Horst DJ, Van Marrewijk WJA, Diederen JHB. Adipokinetic hormones of insect: Release,

signal transduction, and responses. Int Rev Cytol [Internet]. Academic Press; 2001. p. 179–240. Available

from: https://linkinghub.elsevier.com/retrieve/pii/S0074769601110193

117. Čeřovský V, Bém R. Lucifensins, the Insect Defensins of Biomedical Importance: The Story behind

Maggot Therapy. Pharmaceuticals [Internet]. 2014;7:251–64. Available from:

https://www.mdpi.com/1424-8247/7/3/251

118. De Gregorio E. The Toll and Imd pathways are the major regulators of the immune response in

Drosophila. EMBO J [Internet]. 2002;21:2568–79. Available from:

http://emboj.embopress.org/cgi/doi/10.1093/emboj/21.11.2568

119. Martelli C, Pech U, Kobbenbring S, Pauls D, Bahl B, Sommer MV, et al. SIFamide Translates Hunger

Signals into Appetitive and Feeding Behavior in *Drosophila*. Cell Rep [Internet]. 2017;20:464–78.

Available from: https://linkinghub.elsevier.com/retrieve/pii/S2211124717308574

120. Lee K-S, You K-H, Choo J-K, Han Y-M, Yu K. *Drosophila* Short Neuropeptide F Regulates Food Intake

and Body Size. Journal of Biological Chemistry [Internet]. 2004;279:50781–9. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0021925820693620

121. Cholewiński M, Chowański S, Lubawy J, Urbański A, Walkowiak-Nowicka K, Marciniak P. Short

neuropeptide F in integrated insect physiology. Journal of Zhejiang University-SCIENCE B [Internet].

2024;25:389–409. Available from: https://link.springer.com/10.1631/jzus.B2300355

122. Deng L, Chiu IM. A neuropeptide regulates immunity across species. Neuron [Internet].

2022;110:1275–7. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0896627322003075

123. Willis JH, Iconomidou VA, Smith RF, Hamodrakas SJ. Cuticular Proteins. In: Gilbert LI, editor.

Comprehensive Molecular Insect Science [Internet]. Amsterdam: Elsevier; 2005. p. 79–109. Available

from: https://linkinghub.elsevier.com/retrieve/pii/B0444519246000508

124. Kang SH, Fuchs MS. Isolation of a chymotrypsin inhibitor from *Drosophila melanogaster*. Insect

Biochem [Internet]. 1974;4:1–8. Available from:

https://linkinghub.elsevier.com/retrieve/pii/0020179074900365

1096
1097 **Declarations**
1098
Ethics approval and consent to participate
Not applicable.
Consent for publication
Not applicable.
Competing interests
The authors declare that there are no competing financial interests.
Funding
This work was supported by the National Science Foundation Grant No. DEB-2030345.
Author contributions

MJS and JPW conceptualized the experiments. JPW and RKL collected the samples and
performed the experiments, JPW completed the analyses and produced the figures. JPW and
MJS wrote the manuscript and RKL reviewed and edited the manuscript. All authors read and
approved the final manuscript.
1112
1113 **Figure legends**
1114
Figure 1. **Phylogenetic analysis of the ABC transporters.** Three Diptera species were used for the
phylogenetic analysis as follows: *L. cuprina* (Lcup, gold), *L. sericata* (Lser, green) and *D. melanogaster*
(Dmel, black). ABC subfamilies from A to H and node bootstrap supporting values were detailed. Only *L.*
cuprina receptors with a TPM expression ≥ 5 in larval head, gut or whole body were used for the analysis,
but all ABC transporters of *D. melanogaster* were included. In addition, ABC transporters previously
identified as larval biased in *L. sericata* [30] were added to the analysis. If there was more than one
isoform per gene, only isoforms producing a different amino acid sequence and showing a dissimilar
transcript expression were added. Potential expanded *L. cuprina* ABC transporters belonging to subfamily
G were detailed with an asterisk. For more details about sequences used in the analysis refer to Data
S5A-D.
1125
Figure 2. Expression and differential expression of the ABC transporters in *L. cuprina* larval
stages. The TPM expression of ABC receptors was plotted clustering the same according to differential
expression between the three larval stages as follows: (**A**) includes receptors biased to the first two larval
stages (L1 and L2) vs. L3; (**B**) shows receptors only biased to the L2 stage; (**C**) groups receptors biased
to the last two stages (L2 and L3) vs. L1; and (**D**) the receptors only biased to the last stage (L3). Only
receptors with a TPM expression ≥ 5 were added. Dots represent the gene TPM expression of each
library; the average mean expression per gene, plus the SD between libraries were also plotted. For more
details about the expression level and differential expression of each receptor refer to Data S3A.
1134
Figure 3. Expression and differential expression of the ABC transporters in *L. cuprina* third instar
larval tissues. The TPM expression of ABC receptors was plotted clustering the same according to
differential expression between different tissues as follows: (**A**) includes receptors biased to the larval
head (first segment of the larval body) vs. whole body; (**B**) shows receptors biased to the gut vs. whole
body; and (**C**) groups receptors biased to head plus the gut vs. whole body. Only receptors with a TPM
expression ≥ 5 were added. Dots represent the gene TPM expression of each library; the average mean
expression per gene, plus the SD between libraries were also plotted. Symbols next to receptor's names
represent larval stage biased expression (plotted in Fig. 2) as follows: # L1/L2-biased; $ L2/L3-biased; &
L3-biased. For more details about the expression and differential expression of each receptor refer to
Data S4B.
1145
Figure 4. **Phylogenetic analysis of the OBPs.** Three Diptera species were used for the phylogenetic
analysis as follows: *L. cuprina* (Lcup, gold), *L. sericata* (Lser, green), *D. melanogaster* (Dmel, black). OBP
subfamilies classified according to the cysteine motif (Classic, Minus-C, Plus-C and Dimer), and node
bootstrap supporting values were detailed. Only *L. cuprina* OBPs with a TPM expression ≥ 5 in larval
head, gut or whole body were used for the analysis, but all *D. melanogaster* OBPs were included. In
addition, OBPs previously identified as larval biased in *L. sericata* [30] were added to the analysis. If there
was more than one isoform per gene, only isoforms producing a different amino acid sequence and
showing a dissimilar transcript expression were added. Potential expanded *L. cuprina* OBPs 39 to 44
belonging to subfamily Classic were detailed with an asterisk. For more details about sequences used in
the analysis refer to Data S9A-C.
1156
Figure 5. Expression and differential expression of the OBPs across *L. cuprina* larval stages. The
TPM expression of OBPs was plotted clustering the same according to differential expression between
the three larval stages as follows: (**A**) includes OBPs biased to the first two larval stages (L1 and L2) vs.
L3; (**B**) shows OBPs only biased to the L2 stage; (**C**) groups OBPs biased to the last two stages (L2 and
L3) vs. L1; (**D**) OBPs only biased to the last stage (L3); and (**E**) OBPs biased to the first and last stage

(L1 and L3). Only receptors with a TPM expression ≥ 5 were added. Dots represent the gene TPM
expression of each library; the average mean expression per gene, plus the SD between libraries were
also plotted. For more details about the expression and differential expression of each receptor refer to
Data S3B.
1166
Figure 6. Expression and differential expression of larval head-biased OBPs. Only the TPM
expression of head-biased OBPs was plotted. Dots represent the gene TPM expression of each library;
the average mean expression per gene plus, the SD between libraries were also plotted. Only receptors
with a TPM expression ≥ 5 were added. Symbols next to receptor's names represent larval stage biased
expression (plotted in Fig. 5) as follows: # L1/L2-biased; @ L2-biased; $ L2/L3-biased; & L3-biased; Ψ
L1/L3-biased. For more details about the expression and differential expression of each receptor refer to
Data S4C.
1174
Figure 7. CRISPR/Cas9-mediated loss-of-function mutation of the *L. cuprina* odorant coreceptor
(*LcupOrco*) gene. A deletion mutation in the first exon resulted in the introduction of a premature stop
codon at the end of the first exon. (**A**) *LcupOrco* gene detailing exons (E) and introns (I) with lengths in
base pairs (bp), the guide RNA target site, and PAM sequence (underlined); (**B**) alignment between the
wt and mutated *LcupOrco* DNA sequences, detailing deletions and insertions, including a 8 bp gap that
introduces a premature stop codon at 265 bp from the start codon, producing a 73 amino acids (aa)
truncated protein of 478 aa for the *wt* protein; and (**C**) LcupORCO protein sequence detailing the
premature stop codon position located after the glutamic acid (E) 73, the seven transmembrane (TM)
regions and cytoplasmic and external loops. The interaction between Orco protein and any odorant
receptor to produce a functional dimer occurs at the Orco C-term region (-COOH in the figure), which is
absent in the mutated version of LcupOrco protein. For more details about *LcupOrco* silencing refer to
Figs. S2-S4, S7-S12 and Notes S1 and S2.
1187
Figure 8. Odorant receptors are not essential for *L. cuprina* larval response to rotten beef. A larval
choice assay was conducted to evaluate the larval response to fresh (Fr) and rotten (Ro) beef at two
temperatures, cold 25±1°C (Co) and hot 33±1°C (Ho). A total of 120 early L3 (4 days after oviposition)
were tested, and 48 larvae randomly collected from individuals attracted to each diet preference were
genotyped. Larvae were obtained from crossing *LcupOrco* heterozygous flies giving an approximate 3:1
ratio of *LcupOrco* non-silenced (wild-type plus heterozygous, *wt* + HET) and silenced larvae for *LcupOrco*
(homozygous, HOM). (**A**) the numbers on top of the bars represent total larvae attracted to each diet
preference, *i.e.* rotten cold (RoCo), rotten hot (RoHo), fresh cold (FrCo), fresh hot (FrHo) and non-choice
(NC); (**B-C**) show the weight of larvae in milligrams (mg) attracted to each diet preference, either for
genotyped (**B**) or non-genotyped (**C**) larvae. Dots represent single larvae, numbers on top of each
preference or genotype mean total number of larvae, and equal letters indicate no statistically significant
difference (ANOVA, P = 0.22 for B and P = 0.25 for C); (**D**) shows how genotypes were randomly
associated to each diet preference (Chi-Square analysis, P = 0.66). Numbers on top of the bars represent
total larvae per diet preference. For more details about results obtained in this assay refer to Figs. 9A,
10A and Table S2A.
1203
Figure 9. Disruption of *LcupOrco* impairs *L. cuprina* adult female olfaction. Females were obtained
from crossing *LcupOrco* heterozygous flies giving an approximate 3:1 ratio of *LcupOrco* non-silenced
(wild-type plus heterozygous, *wt* + HET) and silenced females for *LcupOrco* (homozygous, HOM). A
different group of *L. cuprina wt* females were used for a further comparison with *LcupOrco* females. All
flies used for the olfaction assay were ~ 10-day-old gravid females and were separated from males the
day before the assay using CO_2 to anesthetize the flies. The attractants used to lure the flies were 2
grams of fresh (Fr) and rotten (Ro) beef at room temperature. A total of 80 *wt* and 174 *LcupOrco* females
were tested, and 72 of the latter were genotyped. (**A**) the *wt* females were more attracted to the rotten
beef vs. fresh beef, or did not respond to any of the stimulus (NC), ANOVA, P = 0.0001; (**B**) in females
belonging to the *LcupOrco* mix, the number of non-responding (NC) females was higher than any of other
two preferences, ANOVA, P = 0.0001; (**C**) genotyping of randomly selected females from tested groups,
showed that *LcupOrco* silenced females were overrepresented among the non-responding flies (NC),

Chi-Square, P = 0.0001. Different letters indicate statistically significant difference. For more details about
results obtained in this assay refer to Figs. S9B, S10B and Table S2B.
1218
1219 **Supplementary information**
1220
Additional file 1: Fig. S1: larval stages and tissues used for the RNA-Seq experiment. **(A)** whole larva
stage 1 (L1); **(B)** whole larva stage 2 (L2); **(C)** lateral view of a whole late larva stage 3 (L3) and **(D)**
ventral view; **(E)** whole early larval stage 3 (WL); **(F)** first segment designated as "head" (H) from an early
L3, detailed within a dashed white circle; and **(G)** gut from an early L3, where crop have been separated
from the cardia originally connected by the foregut (not present in the picture); **Note S1:** *LcupOrco* gene,
guide RNA (gRNA) sequence, and primers used to determine potential indels and point mutations at the
gRNA cutting site (genotyping); **Fig. S2:** *LcupOrco* insert construction. **(A)** *LcupOrco* gene fragment
including two sequences corresponding to the left (LHA) and right (RHA) homology arms of 1000 bp in
length each one, necessary for the CRISPR homology directed repair (HDR) protocol cloned using
plasmid pUCIDT-AMP GoldenGate (ordered from IDT). In addition, a DNA section including the restriction
cutting sites (RCS) for enzymes XhoI (NEB) and NotI (NEB) was added in between of both sequences in
the same plasmid; **(B)** donor plasmid including the ZsGreen Marker followed by the *Hsp83* promoter
located between positions 1891 to 7441 [2]; **(C)** ZsGreen Marker obtained from the donor plasmid. The
plasmid was linearized using restriction enzymes BpmI (NEB), XhoI and NotI at 37°C for 60 min and the
reaction was finished at 65°C for 20 min. After restriction digestion the mix was electrophoresed in a 1.5%
agarose gel at 60 volts for 120 min. The band with the size corresponding to the marker (~5 Kb) was
recovered from the gel and purified using the Zymoclean Gel DNA Recovery Kit (Zymo) following the
manufacturer's specifications; **(D)** to obtain the final construct, 50 ng of linearized pUCIDT-AMP
GoldenGate plasmid + 175.5 ng of the marker fragment (1:3 ratio) were ligated overnight (O/N) at 16°C
using the T4 DNA Ligase (NEB). After ligation, 1 µl of the reaction mix was used to transform 10-beta
competent *Escherichia coli* cells (NEB) following the manufacturer's specifications. Seventy-five µl of
undiluted transformed 10-beta cells were poured into an Ampicillin 20 mg/ml (+Amp) Lysogeny Broth (LB)
plate and incubated O/N at 37°C; **(E)** ten clones were recovered from the LB plate and grow in 3 mL of
liquid LB +Amp O/N at 37°C and 250 rpm. To obtain the plasmids from clones the ZR Plasmid Miniprep
Kit (Zymo) was used following the manufacturer's specifications and evaluated with the BpmI, XhoI and
NotI restriction enzymes following the same protocols described above for restriction digestion and
electrophoresis; **(F)** three positive plasmids for the selected restriction cutting sites were sequenced using
Oxford Nanopore and aligned to the *in-silico* construct to confirm the sequence identity. Abbreviations: Kb
= kilobase; MW = molecular weight; **Fig. S3:** containers used for fly rearing and crosses. **(A)** plastic bottle
used for fly rearing; **(B)** sandglass shaped arrangement made using two five ounces clear-plastic cups
joined from removed bottoms used for fly crosses after embryo injections. These containers also included:
a top holed cap and a bottom cap with an opening of ~ 1 inch in diameter; a rounded piece of white paper
towel; a small glass vial filled with tap water and a small protein cookie made with yeast, milk and egg
powder and cane sugar; **Fig. S4:** fly crosses completed after embryos microinjections. Abbreviations: G =
generation; *wt* = wild-type; **Note S2:** primers used for *LcupOrco* insert detection of landing site. Three
types of pair of primer associated with three different strategies were used to determine if the landing site
of the *LcupOrco* insert was withing the *LcupOrco* locus in the *L. cuprina* genome. First one was
associated with the left side (LS) of the insert including the upstream genome region, the *Hsp83* promoter
and part of the green marker; the second one was designed to amplify the right side (RS) of the insert
including the part of Tub3'-UTR and the downstream genome region; and third one was to amplify the
whole *LcupOrco* insert (All) including from upstream to downstream *L. cuprina* genome regions flanking
the *LcupOrco* locus; **Table S1:** RNA-Seq data overview. **(A)** each sample represents an RNA-Seq library.
Sample's names, description and number of raw, trimmed and mapped reads per library are provided; **(B)**
total sequences identified for the RNA-Seq experiment, divided by RNA type; **Fig. S5:** principal
component analysis (PCA) completed using transcripts with a TPM expression ≥ 5 for all libraries
(samples) associated with EXP-1 (larval stages). Samples WL1_1 to 3, WL2_1 to 3 and WL3_1 to 6 were
collected from the LA07 colony. Samples WL1_4 and 5 and WL2_4 and 5 were collected from
heterozygous individuals obtained from the crossing between the LA07 and the SLAM5X colonies. The
PCA analysis did not show differences between samples of the same larval stage, collected from different

colonies. Abbreviations: L1 = first larval state; L2, second larval stage; L3 = third larval stage; WL = whole
larva sample; TPM = transcript per million; **Fig. S6:** ABC transporters preliminary Neighbor Joining
phylogenetic analysis. Abbreviations: Dmel = *Drosophila melanogaster*; Lcup = *Lucilia cuprina*; Lser =
Lucilia sericata; **Fig. S7:** guide RNA (gRNA) cutting efficiency tested *in vitro* using the EnGen Spy Cas9.
The amplicon size of *LcupOrco* gene fragment without cutting was 439 bp and after Cas9 *in vitro* cutting
generated two overlapped amplicons of 224 and 215 bp, respectively. Abbreviations: C = control, uncut
genomic DNA; MW = molecular weight; T = treated, gDNA + Cas9; **Fig. S8:** *Lucilia cuprina* eggs, larva
stage 3 (L3) and adult flies expressing the ZsGreen marker before the genotyping analysis to determine
LcupOrco insert landing site. **(A)** L3 showing transient expression of the marker 7 days after *wt* eggs
injection; **(B)** eggs obtained from crossing a *LcupOrco* mosaic male with a *wt* female; **(C)** *wt* L3 on top vs.
a heterozygous ZsGreen L3 at the bottom; **(D)** heterozygous ZsGreen L3 on top vs. homozygous for the
same marker at the bottom; **(E)** dorsal view of a *wt* adult male (left) vs. G1 heterozygous ZsGreen male
(right) under bright field; **(F)** same males showed in E under a green filter; ventral view of the same males
using bright field **(G)** and a green filter **(H)**; **Fig. S9:** PCR amplification of gDNA of samples collected from
LcupOrco larval diet preference test and female olfaction assay. **(A)** samples of larval olfaction assay; **(B)**
samples of adult female olfaction assay. Primers, amplicon sequence and cycling parameters are detailed
in **Note S1** (see Supplementary Materials and Methods section). Abbreviations: MW = molecular weight;
Fig. S10: Sanger sequencing results for samples collected from *LcupOrco* larval diet preference test and
adult female olfaction assay. (**A**) genotyped samples from # 1 to 48 of larval diet preference test; and (**B**)
genotyped samples from # 1 to 72 of adult female olfaction assay. Synthego software was used to
analyze sample's chromatograms and indels % = 0 corresponds to *wt* samples, from 1 to 89% to
heterozygous samples, and ≥ 90% to homozygous samples. Results were compiled along with single
larva preferences in **Table S2 A-B**; **Table S2:** compiled data of samples associated with *LcupOrco*
behavioral assays. **(A)** larval diet preference test; **(B)** adult female olfaction assay. The numbers one
through eight after the letters in table B, refer to the main chambers of the olfactometer and numbers after
dots, to single females from the same chamber. Abbreviations: Fr = fresh beef (room temperature); FrCo
= fresh cold beef; FrHo = fresh hot beef; HET = heterozygous; HOM = homozygous; NC = non-choice;
NG = non-genotyped; Ro = rotten beef (room temperature); RoCo = rotten cold beef (25±1°C); RoHo =
rotten hot beef (33±1°C); *wt* = wild-type. Hyphen symbol means non-tested by Sanger sequencing; **Fig.**
S11: eggs obtained from *LcupOrco* mutated females. **(A)** eggs obtained from 8-day-old *LcupOrco*$^{+/-}$
females 3 days after crossing them to *NPF*$^{-/-}$ males expressing the ZsGreen marker. The same marker
was removed from *LcupOrco* females after confirming that the *LcupOrco* insert landing site was not
located withing the *LcupOrco* gene locus in the *L. cuprina* genome. The insert was removed by selecting
non-fluorescent larvae and crossing males obtained from them for two generations vs. *wt* females. In
addition, males from each generation were genotyped using protocols described in **Note S1** (see
Supplementary Materials and Methods section) to confirm the presence of indels and point mutations
within the *LcupOrco* coding region. The fluorescence in eggs confirmed the mating between the
LcupOrco$^{+/-}$ females and *LcupNPF*$^{-/-}$ males and egg fertilization. *LcupOrco*$^{-/-}$ females did not lay eggs at
this time; **(B)** part of *LcupOrco*$^{-/-}$ females laid eggs 10 days after mixing them with *LcupNPF*$^{-/-}$ males but
eggs were not fertilized as showed under green filter **(C)**, and no larvae emerged from them; part of the
LcupOrco$^{-/-}$ females laid fertilized eggs 17 days after mixing with *LcupNPF*$^{-/-}$ males, as showed under
bright field **(D-F)** and green filter **(E-G)**; these eggs produced healthy progeny; **Fig. S12:** ovaries from 10-
day-old *L. cuprina wt* and *LcupOrco* mutated females. Three females per condition were dissected
(columns). All females were provided with tap water and sugar, but the presence of protein in the diet and
mating condition changed between groups; *wt* virgin females not fed **(A-C)** and fed with protein **(D-F)**; *wt*
mated females not fed **(G-I)** and fed with protein **(J-L)**; *LcupOrco-/-* mutated females fed with protein **(M-**
O).
1317
1318   **Additional file 2: Data S1. TPM transcript expression for *L. cuprina* larval stages. A** to **C**
1319   compile all libraries from each stage, namely L1, 2 and 3, showing the average expression for
1320   each transcript, only for transcripts with a TPM expression ≥ 5. **D** to **S** show the expression of all
1321   transcripts for all libraries.
1322

1323 **Additional file 3: Data S2. TPM transcript expression for *L. cuprina* early L3 tissues. A** to
1324 **C** compile all libraries for the larval head, gut and whole larva, respectively. Same tabs also
1325 show the average expression for each transcript, only for transcripts with a TPM expression ≥ 5.
1326 **D** to **N** show the expression of all transcripts for all libraries.
1327
Additional file 4: Data S3. Sensory receptors expressed in *L. cuprina* larval stages. A)
ATP-binding cassette (ABC); **B**) odorant-binding protein (OBP); **C**) transient receptor potential
channel (TRP); **D**) CD36-sensory neuron membrane proteins (CD36/SNMPs); **E**) ML (MD-2
(myeloid differentiation factor-2)-related Lipid-recognition and Niemann-Pick C2 disease
proteins (ML/NPC2); **F**) chemosensory protein (CSP); **G**) ammonia transporter (AMT). Only
receptors with a TPM expression ≥ 5 are displayed. Differential expressions between stages for
each receptor were listed.
1335
Additional file 5: Data S4. Sensory receptors expressed in *L. cuprina* early L3 tissues. A)
gustatory receptor (GR); **B**) ATP-binding cassette (ABC); **C**) odorant-binding protein (OBP); **D**)
transient receptor potential channel (TRP); **E**) CD36-sensory neuron membrane proteins
(CD36/SNMPs); **F**) ML (MD-2 (myeloid differentiation factor-2)-related Lipid-recognition and
Niemann-Pick C2 disease proteins (ML/NPC2); **G**) chemosensory protein (CSP); **H**) ammonia
transporter (AMT). Only receptors with a TPM expression ≥ 5 are displayed. Differential
expressions between stages for each receptor were listed.
1343
1344 **Additional file 6: Data S5. ABC protein sequences included in the phylogenetic analysis.**
1345 **A**) *D. melanogaster*, **B**) *L. cuprina*; **C**) *L. cuprina* receptors with a TPM expression ≥ 5; **D**) larval
1346 biased *L. sericata* ABC transporters vs. adult flies.
1347
1348 **Additional file 7: Data S6. TPM transcript expression and differential expression of the**
1349 **ABC transporter LcupABCC13_Sur and LcupABCG6.** Annotation of both receptors was
1350 corrected using assembly GCA_001187945.1, and the transcript expression and differential
1351 expression between **A**) larval stages and **B**) tissues was calculated.
1352
1353 **Additional file 8: Data S7. Transcripts enriched in A) the larval head or B) larval gut,**
1354 **considering only TPM expression ≥ 5.**
1355
1356 **Additional file 9: Data S8. Differential expressed (DE) transcripts between different *L.***
1357 ***cuprina* early L3 tissues.** Head and gut biased transcripts with a TPM expression ≥ 5 are
1358 provided in tabs **A** and **C**. All DE expressed transcripts between the head or the gut vs. the
1359 whole larva are displayed in tabs **B** and **D**, respectively.
1360
1361 **Additional file 10: Data S9. OBP protein sequences included in the phylogenetic analysis.**
1362 **A**) *D. melanogaster*, **B**) *L. cuprina* receptors with a TPM expression ≥ 5; **C**) larval biased *L.*
1363 *sericata* OBPs vs. adult flies.
1364
1365 **Additional file 11: Data S10. Differential expressed (DE) transcripts between different *L.***
1366 ***cuprina* larval stages. A** to **C** compile those DE transcripts for L2 vs. L1, L3 vs. L2 and L3 vs.
1367 L1, respectively. Only transcripts with a TPM expression ≥ 5 are displayed. From **D** to **N** same
1368 groups comparisons are shown but including all DE transcripts.
1369
1370 **Table 1. *Lucilia cuprina* Orco expression in adult and larval tissues**

Tissue	TPM expression	Cite

Whole larva stage 1	0.48	This study
Whole larva stage 2	0.15	This study
Whole larva early stage 3	0.02	This study
Whole larva late stage 3	0.02	This study
First segment of larva early stage 3	0.38	This study
Gut of larva early stage 3	0.003	This study
Antenna of gravid female 10-d-old	1074	Wulff *et al.* 2024
Antenna of mated male 10-d-old	924	Wulff *et al.* 2024

1371

1372

1373

1374

1375

A CRISPR/Cas9 guide RNA
GACGAAGTAAATGAGTTGTC <u>CGG</u> (+)

100 bp

LcupOrco^wt

I1 382 bp I2 89 bp I3 91 bp I4 4472 bp I5 1529 bp I6 4260 bp

| E1 308 bp | E2a 136 bp | E2b 378 bp | E3 203 bp | E4 205 bp | E5 156 bp | E6 51 bp |

TM1 TM2 TM3 TM4 TM5 TM6 TM7

B
LcupOrco ^wt AAATGAGTTGTCCGGTAATACAATTACCG
LcupOrco ^-/- AAATGAA--------**TAA**AAGAAT-ACCG
           * * * * * *   * * *   *   * * *   * * * *

STOP

C

Extracellular

STOP

-COOH

TM1 TM2 TM3 TM4 TM5 TM6 TM7

Cytoplasm

H₂N-