A simple method to efficiently generate structural variation in plants

Lindsey L. Bechen[1], Naiyara Ahsan[2,4], Alefiyah Bahrainwala[2], Mary Gehring[1,3,5*], PRV

Satyaki[1,2,4*]

1 Whitehead Institute for Biomedical Research, Cambridge, MA, USA

2 Dept. of Biological Sciences, University of Toronto Scarborough, Toronto, ON,

Canada

3 Dept. of Biology, Massachusetts Institute of Technology, Cambridge, MA USA

4 Dept. of Cell and Systems Biology, University of Toronto, Toronto, ON, Canada

5 Howard Hughes Medical Institute

*co-corresponding authors

mgehring@wi.mit.edu

satyaki.rajavasireddy@utoronto.ca

Key Words: structural variation, etoposide, Arabidopsis, mutagenesis

Abstract

Phenotypic variation is essential for the selection of new traits of interest. Structural variants, consisting of deletions, duplications, inversions, and translocations, have greater potential for phenotypic consequences than single nucleotide variants. Pan-genome studies have highlighted the importance of structural variation in the evolution and selection of novel traits. Here, we describe a simple method to induce structural variation in plants. We demonstrate that a short period of growth on the topoisomerase II inhibitor etoposide induces heritable structural variation and altered phenotypes in *Arabidopsis thaliana* at high frequency. Using long-read sequencing and genetic analyses, we identified deletions and inversions underlying semi-dominant and recessive phenotypes. This method requires minimal resources, is potentially applicable to any plant species, and can replace irradiation as a source of induced large-effect structural variation.

Introduction

Genomic structural variation (SV) – insertions, deletions, duplications, translocations and inversions – are an important source of genetic and phenotypic novelty. Structural variants create genetic novelty by multiple means, including fusing the coding regions of two genes, altering cis-regulatory environments and gene expression patterns, and by altering or suppressing recombination, among other mechanisms. In addition, deletions and duplications alter gene or chromatin dosage [1–3]. Novel structural variants have larger phenotypic effects per mutation than single nucleotide variants [4] and have been selected for when they underlie traits beneficial to the organism or desirable to agriculturists [5,6] .

Recent pan-genome sequencing of multiple plant species has underscored the versatility and significance of structural variation to phenotypic variation, modern crop traits, and crop improvement. Lower seed sets in watermelons and grapes, important for the development of popular "seedless" varieties, have been linked to large inversions that cause meiotic defects [7,8]. SVs have also been linked to the domestication of broomcorn millet [9] and sorghum [10]. In pearl millets, gain of heat tolerance has been linked to SVs at multiple loci [11]. SVs are also associated with variation in immunity and plant defense responses [12–14]. In Arabidopsis, a multi-enzyme pathway synthesizes the glucosinolate family of defense compounds. SVs that either delete or fuse enzyme-encoding paralogs modify enzymatic pathways and are responsible for ecotype-specific variation in glucosinolates [15]. Studies of *Oryza sativa* provide numerous examples of how SVs influence phenotypes. Change in copy number of *VIL1* is associated with changes to flowering time and grain number variation [16]. In addition, duplications of the *KALA4* gene – a regulator of the anthocyanin biosynthesis pathway – created a novel cis-regulatory region that promoted ectopic expression of anthocyanins and caused a "black pericarp" phenotype [17]. In peaches, fruit flesh color around the stone and fruit shape are dependent on a deletion and a mega-base scale inversion [18]. Modern varieties of sweet corn were created by selecting for an inversion that creates a loss-of-function mutation in the *shrunken-2* gene, which encodes the first

enzyme of the starch biosynthesis pathway [19,20]. In tomatoes, a duplication of a

cytochrome P450 gene was linked to increased fruit weight [5].

Extant crop genetic diversity is unlikely to provide all possible SV diversity that could

impact crop improvement. Induced structural variation can therefore provide novel

germplasm for breeding programs, information that can be used for targeted CRISPR-

mediated restructuring of crop genomes, and novel genetic variants for understanding

fundamental aspects of plant biology. The most common method employed to induce

random structural variation is exposure to ionizing radiation, including X-rays, gamma

irradiation, and heavy ions [21–26]. Since the 1960's, irradiation has been used to

induce structural variation in model organisms and in numerous crops including, more

recently, rapeseed [27], wheat [28], cotton [29], rice [23,30], poplar [22,31], soybean

[32,33], and buckwheat [34]. Populations carrying such structural variation have been

used for the isolation of mutations that abolish hybrid sterility in rice [35] and the

creation of seeds differing in oil composition [21,27].

Although radiation-induced structural variation remains an important element of modern

plant breeding, there are logistical challenges. In contrast to the induction of single

nucleotide variants that researchers can conveniently pursue in their own labs using

ethyl methanesulfonate (EMS) or sodium azide [36], the induction of structural variants

requires access to radiation sources. Radiation sources are heavily regulated and

access to nuclear reactors, particle accelerators, and other sources of radiation is a

bottleneck in the creation of structural variant libraries [26]. Additionally, application of

irradiation can be tedious. For example, the creation of a mutant population in poplar

required irradiation of collected and dried pollen [22]. One alternative is the extended

TAQing approach, in which a transgene expresses a restriction enzyme to induce

conditional double-stranded DNA breaks, which then leads to shuffling of the plant

genome [37]. However, this approach can be challenging as the commercial varieties of

many crops are not easily transformed.

To surmount the current challenges of creating SVs in plants, we sought a simple

method that could be applied to any plant species of interest to induce SVs with high

frequency, drawing inspiration from the ubiquity and ease of EMS chemical

mutagenesis to induce single nucleotide variants. One possible class of mutagens for

SV induction is DNA topoisomerase II (Topo II) inhibitors. Topo II relaxes torsional

stress from DNA supercoiling generated during DNA replication or transcription by

transiently breaking both strands and then ligating them after passing a DNA segment

through the break. Between strand breakage and ligation, Topo II is covalently linked to

DNA via a tyrosine residue, forming a topoisomerase cleavage complex [38]. This

complex is stabilized by the inhibitor etoposide. A collision between covalently-linked

Topo II and DNA polymerases during DNA replication, or with RNA polymerases during

transcription, leads to removal of the Topo II enzyme, which results in the generation of

double-stranded breaks (DSBs) [39–42]. The imprecise repair of DSBs leads to

genomic rearrangements and structural variation in mouse spermatocytes, fibroblasts,

and in human cells [43–45]. Previously, it was shown that treatment with etoposide

impacts genome stability and inhibits plant growth in *Arabidopsis thaliana*, *Allium cepa*,

and *Lathyrus sativus* [46–49] and causes chromosomal fragmentation during meiosis in

Arabidopsis [48] . However, its potential as a mutagen that can induce structural

variation in plants has not been investigated. Here, we provide proof-of-principle that the

chemotherapeutic drug etoposide efficiently generates novel genomic structural

variation in *Arabidopsis thaliana*.

Results

Treatment with etoposide results in mutants with a variety of heritable

phenotypes

To test whether etoposide can induce heritable structural variation in plants, we

examined the effects of prolonged exposure to etoposide on Arabidopsis. Wild-type Col-

0 seeds were germinated and grown on MS media supplemented with sucrose and

differing concentrations of etoposide (0 μM (DMSO only), 20 μM, 40 μM, 80 μM, 160

μM, 320 μM, or 640 μM) for up to two weeks (Figure 1A). Plants grown on DMSO alone

or 20 μM etoposide showed no observable growth defects (Figure 1A). Two-week-old

seedlings grown on 40 μM etoposide exhibited gnarled leaves but no differences in size.

Figure 1: Etoposide induces novel heritable phenotypes. (A) Arabidopsis seeds germinated on increasing concentrations of etoposide show a dose-responsive growth defect (upper panel). Seedlings grown on 320 μM etoposide did not exhibit true leaves. This generation is referred to as M1. Growth defects persisted after transplantation to soil (lower panel). **(B)** M1 was self-fertilized to give rise to M2. Subsequent rounds of self-fertilization produced M3, M4, and M5 generations. A visual examination of M2 generation identified multiple phenotypes, including: **(C)** a *brassinosteroid-like dwarf* phenotype, **(D)** a fertile dwarf with short internodes, **(E)** a *virescent* phenotype with yellowish leaves, and **(F)** a *variegated* phenotype. **(G-I)** describe crosses to the wild-type and subsequent selfing to identify the nature of inheritance of the mutant phenotypes and the number of contributing loci. **(G)** The *BR-like dwarf* phenotype is incompletely dominant and displays three phenotypic classes. F2 segregation data suggests that this phenotype is caused by mutation at a single locus. **(H)** The *short-internode dwarf* phenotype is recessive and is likely caused by more than one locus. **(I)** The *virescent* phenotype is recessive and caused by a single locus.

Figure S1: Additional phenotyping of progeny of etoposide-treated plants. Visual observation of M2 plants arising from different etoposide-treated M1 progenitors identified multiple lines with mutant phenotypes, including: early flowering **(A)**, sterility **(B)**, dwarfism **(C)**, variegation **(D)**, altered leaf shape **(E-G)** . Lines with altered leaf shape or size were visually identified. These differences were confirmed by measuring leaf size (blade area, length, perimeter, width, and petiole length) and shape (cirularity) of leaves with the LeafJ plugin for ImageJ software. Leaves from multiple M2 siblings of line 35 are shown in **(G)**. 35A_1 and 35A_2 show a dramatic decrease in size parameters while 35A_3 shows an intermediate decrease. 35A_4 and 5 show no change compared to DMSO-treated lineages. **(H)** *short-internode dwarf* plants have the same number of nodes as DMSO-treated lineages and a genetic relative that lacks the phenotype, but **(I)** decreased distance between nodes, as measured on the primary inflorescence.

Seedlings grown on 80 µM etoposide showed marginally reduced growth (Figure 1A). Of the seedlings grown on 160 µM of etoposide, less than half (75/184) developed true leaves. In addition, these plants displayed stunted root development (Figure 1A). Seedlings grown on 320 µM or 640 µM etoposide showed little root or shoot growth and high seedling lethality. Seedlings grown on DMSO only, 80 µM, and 160 µM etoposide (referred to as M1 plants) were transplanted to soil and compared at maturity. Those exposed to 160 µM of etoposide exhibited significantly more abnormal phenotypes than DMSO only or 80 µM etoposide plants, including loss of apical dominance, gnarled leaves, reduced plant size, seed abortion, and lower seed number at maturity (Figure 1A). To test if these abnormal plants produced progeny with mutant phenotypes, we scored for six visible phenotypes—dwarfism, loss of apical dominance, seed size or abortion, leaf shape or size, flowering time, and leaf pigmentation—among the progeny (M2 generation) of plants treated with 160 µM etoposide. Of M2 progeny derived from 42 different M1 parents, 29 lines exhibited at least one obvious visible phenotype (Figure 1B-F, Figure S1, Table S1). These phenotypes included: variegated or albino plants, altered flowering time, altered leaf shape and size, sterility or seed abortion, and dwarfism (Figure S1, Table S1). The large proportion of plants showing visible phenotypes suggested that etoposide could be an excellent mutagen for efficiently creating large-effect mutations.

To assess the heritability of the mutant phenotypes and to establish genotype-phenotype relationships, we closely examined four M1 lines with visible phenotypes: a dwarf line reminiscent of weak brassinosteroid-insensitive mutants (Figure 1C) [50], referred to as *brassinosteroid-like (BR-like) dwarf* (line 1A); a dwarf line with short internodes (line 34C), or *short-internode dwarf* (Figure 1D); a line with delayed greening (line 13B), termed *virescent* (Figure 1E); and a line with a *variegated* phenotype (line 5A) (Figure 1F). The semi-sterile *BR-like dwarf* line (Figure 1C) exhibited short, thick stems that bore fleshy leaves. In contrast, the *short-internode dwarf* mutant, though short in stature, was fertile, produced the same number of nodes on the primary inflorescence as WT plants (Figure S1H,I), and lacked the fleshy leaf phenotype observed in the *BR-like dwarf* (Figure 1D). In the *virescent* line, juvenile plants exhibited reduced chlorophyll pigmentation, whereas adult plants exhibited normal pigmentation

Figure S2: Segregation of the *variegated* phenotype from green parents. Green plants from variegated line 5A produce either approximately 25% variegated progeny or all green progeny. Number of seedlings assesed is annotated above each bar. M2 plant 5A_58 is the parent of the M3 progeny assayed.

(Figure 1E). *Variegated* mutants displayed light- and temperature-dependent variations

in the proportions of green and white sectors on cotyledons and true leaves (Figure 1F).

The *BR-like dwarf, short-internode dwarf,* and *virescent* phenotypes were transmitted

from self-fertilized parents to offspring for at least three additional monitored

generations, indicating that the phenotype was true-breeding and the underlying mutant

genotype was stable (Figure 1B). To assess whether these phenotypes were dominant

or recessive, mutant lines were crossed to wild-type. The resultant F1 progeny from the

BR-like dwarf exhibited intermediate dwarfing, suggesting that the phenotype was

incompletely dominant (Figure 1G). The F1 progeny from the *short-internode dwarf* line

and the *virescent* line displayed a wild-type phenotype, indicating that these mutant

phenotypes were recessive (Figure 1H, I). *variegated* plants did not flower readily, so

this phenotype was maintained as a heterozygous stock for the same number of

generations. When self-fertilized, plants from this line either produced all green progeny

or one-quarter variegated progeny, indicating that this phenotype is recessive (Figure

S2).

To estimate the number of loci contributing to the *BR-like dwarf*, *short-internode dwarf*,

and *virescent* phenotypes, we examined F2 progeny obtained by self-fertilizing the F1

plants. Of F2 progeny of *BR-like dwarf* plants with an intermediate dwarfing phenotype,

23.4% had a wild-type phenotype, 49.6% exhibited an intermediate phenotype, and

27.0% had a severe mutant phenotype (Figure 1G). This result suggested that a single

incompletely dominant locus causes the *BR-like dwarf* phenotype (58:123:67

severe:intermediate:wild-type, H_0=1:2:1 ratio, χ^2=0.669; df=2, *p*>0.5). The *short-*

internode dwarf phenotype was observed in 1/8 of progeny obtained by self-fertilizing

F1 plants (28:195 *short-internode dwarf*:wild-type) (Figure 1H). This ratio is consistent

with the recessive phenotype being caused by mutation in two linked loci, among other

possibilities. For the *virescent* line, 20.9% of the F2 progeny displayed the mutant

phenotype (Figure 1I). This is consistent with the phenotype being caused by a mutation

at a single locus (29:114 *virescent*:wild-type, H_0=1:3, χ^2= 1.69; df=1, 0.25>*p*>0.1).

These observations indicate that etoposide mutagenesis can efficiently create novel

recessive and dominant phenotypes caused by large-effect alteration to one or more
loci.

RNA-Seq identifies genes that are associated with mutant phenotypes

To further characterize the four phenotypes studied in depth, we performed RNA
sequencing of rosette leaves for M3 plants of *BR-like dwarf, short-internode dwarf,*
virescent, and *variegated* lines (Table S2). As controls, we included genetic relatives
that lacked the phenotype and M3 progeny of DMSO-treated plants. To obtain a broad
overview of changes in gene expression, we performed gene set enrichment analysis
(GSEA) for biological process gene ontology (GO) on the full ranked list of genes
(ranked by $\log_2$ fold change) for each phenotypic line. We also identified differentially
expressed genes for each phenotype, defined as genes with an adjusted *p*-value < 0.01
and a |$\log_2$(fold change)| > 1 (Tables S3-S5).

For the *BR-like dwarf* phenotype, a total of 309 genes were differentially expressed in
BR-like dwarf plants compared to non-phenotypic plants (114 downregulated and 195
upregulated; Figure 2A, Table S3). The five most significant GO terms that were
overrepresented among upregulated genes were: meristem development, anatomical
structure formation involved in morphogenesis, response to auxin, plant organ
formation, and post-embryonic plant morphogenesis (Figure S3A). *AS1* was a
differentially expressed gene ($\log_2$FC = -2.62, adjusted *p*-value = 6.37 x 10^{-16}), along
with several genes it is known to interact with either directly or indirectly such as
KNAT1, *KNAT2*, *KNAT6*, *STM*, *BOP1*, *BOP2*, and *LOB* (Figure 2A, Table S3) [51–54].
The phenotypes we observed in the *BR-like dwarf* plants were strikingly similar to those
described for *as1* mutants, including a compact rosette with lobed and curled leaves
[51,55].

In contrast to *BR-like dwarf* plants, *short-internode dwarf* plants showed considerably
fewer gene expression changes in leaves. A total of 44 genes (18 down and 26 up)
were differentially expressed in *short-internode dwarf* leaves compared to control leaves
(Figure 2B, Table S4). GSEA yielded primarily stress-associated GO terms (Figure
S3B). Interestingly, both the largest subunit of RNA polymerase III, *NRPC1,* and the

Figure 2: Etoposide-induced mutants exhibit changes in gene expression. Volcano plots summarizing differential expression analysis of phenotypic vs. non-phenotypic plants for **(A)** *BR-like dwarf*, **(B)** *short-internode dwarf*, and **(C)** *virescent* mutants. Significantly differentially expressed genes (|log₂FC| > 1 and adjusted *p*-value < .01) are highlighted in (A) blue, (B) pink, and (C) green. Select genes are labeled in each volcano plot. **(D)** Heatmap of expression of genes known to be associated with variegation, in the *variegated* line. The most downregulated gene among them, *IMMUTANS,* is annotated with a *.

Figure S3: GO term enrichment for genes differentially expressed between plants exhibiting a phenotype and those without a phenotype. GSEA was used to identify enriched GO terms. Ridge plots depict significant terms for **(A)** *BR-like dwarf*, **(B)** *short-internode dwarf*, and **(C)** *virescent* phenotypes.

repressor of RNA polymerase III, *MAF1*, were upregulated. It is possible that *short-*

internode dwarf plants exhibit changes in gene expression in other plant organs besides

the leaf or perhaps exhibit a global change in gene expression that cannot be detected

with standard RNA-seq approaches.

Virescent plants had more dysregulated genes than either dwarf line, with a total of

1273 differentially expressed genes (668 up and 605 down; Figure 2C, Table S5). The

most significant GO terms overrepresented amongst genes with increased expression

include cell division, plant-type cell wall organization/biogenesis, mitotic cell cycle,

microtubule-based process, and meiotic cell cycle (Figure S3C). The most significantly

overrepresented GO terms among genes with decreased expression include response

to chitin, response to hypoxia, and response to oxygen levels (Figure S3C).

Interestingly, genes related to ethylene synthesis and signaling including *ERF022*,

ERF014, and *ACS5* were among the most differentially expressed genes (Table S5).

The mis-regulation of these genes might be associated with the *virescent* phenotype, as

ethylene is involved in greening of etiolated seedlings after light exposure, and

conversely degradation of chlorophyll during leaf senescence [56,57]. Other

differentially expressed genes of note include several subunits of photosystem I and II

and *GUN4* (Table S5).

Differential expression analysis could not reliably be conducted for the *variegated* line,

as only one variegated plant was recovered. As an alternative strategy, we examined

the expression of 18 genes known to be associated with variegation (Figure 2D). In the

one variegated plant we recovered and performed RNA-seq on, the *IMMUTANS* (*IM*;

AT4G22260) gene stood out as highly downregulated (Figure 2D). The sensitivity of the

variegation to light and temperature observed in *variegated* plants is consistent with that

displayed in *im* mutants [58,59].

Etoposide treatment induces a spectrum of structural variation types

To characterize the molecular nature of etoposide-induced mutations, we employed

Illumina short-read sequencing technology to sequence the genomes of 32 M2 or M3

progeny of 15 etoposide-treated M1 plants (Table S6). Sequenced plants included

Figure 3: Impact of etoposide on SNVs and repetitive regions. (A) Total number of indels and single nucleotide variants (SNVs), genome-wide, identified in etoposide-treated (purple) and control lines (blue). **(B)** Box-plot describes the composition of nucleotide changes in progeny of plants exposed to etoposide and progeny of plants exposed to DMSO. All SNVs passing quality filters were included here. REF indicates reference allele and ALT represents the alternate SNV. Wilcoxon test was used to determine the significance of differences in SNV numbers between the progeny of etoposide and DMSO treated plant; n.s indicates *p >0.05*. **(C)** For Illumina short-read libraries of each control and etoposide-treated plant that we sequenced, total number of reads aligning to the telomere (CCCTAAA), NOR2, and NOR4 were normalized to total numbers of aligned reads in each library. Number of thousand reads, per million sequenced reads, for all control and etoposide treated libraries are represented in this boxplot. Wilcoxon test shows no statistical difference in the median value of read-depth over repetitive regions in control and etoposide-treated plants.

those with no scored mutant phenotype and those expressing a phenotype, including

BR-like dwarf, *short-internode dwarf*, and *virescent,* among others. We also sequenced

genetic relatives of these three mutants that displayed wild-type phenotypes (see Table

S7 for relationships). To enhance the accuracy of our structural variation (SV) and other

mutation calls, we also identified mutations segregating in our laboratory Col-0

population by sequencing four M2 plants descended from four control M1 lines treated

only with DMSO.

As the inhibition of topoisomerase II activity increases double-stranded DNA breaks in

the genome [60], the short-read data was used to test if the repair of these breaks

results in increased single nucleotide variation (SNV). However, SNV analysis identified

a comparable spectrum and number of SNVs in etoposide-treated and control lines

(Figure 3A,B), suggesting that etoposide did not induce excess SNVs.

To test the potential for the induction of SVs in difficult-to-map repetitive regions, we

measured read depth at rDNA and telomeres and assessed if etoposide treatment was

linked to a higher degree of repeat copy number variation. While the number of control

plants are insufficient to make a definitive statement, we found that the variation in read

depth among etoposide-treated lines was comparable to that in control lines, suggesting

that etoposide treatment likely did not trigger additional genomic instability in repetitive

DNA (Figure 3C).

To identify etoposide-induced SVs, we first adapted a computational approach that uses

genome-wide coverage to identify large segmental deletions or duplications [61,62] .

Using this technique, we identified 10 duplications and 1 deletion that were between

100 kb and 1 Mb in length (Figure 4A, Table S8). The robustness of this analytical

approach is underscored by detecting the same duplications in M2 siblings (Figure S4,

Table S8). Next, we employed Lumpy Express [63] to identify structural variants (SVs)

in mutant and control plants. This pipeline identified 2,224 variants that included

inversions, deletions, duplications, and breakends, which were then filtered for high-

confidence events. Breakends were excluded from further analyses because of the

challenges of identifying their molecular nature. A high proportion of predicted SVs

Figure 4: Whole genome sequencing identifies structural variation in progeny of etoposide treated plants. (A) Duplications in etoposide-treated lines as assessed by short-read sequencing. Dosage across the indicated chromosome is shown. Chromosomal segments with one extra copy are indicated with +. **(B)** Nanopore long-read sequencing identifies structural variation in lines that show the *BR-like dwarf*, *short-internode dwarf*, *virescent,* and *variegated* phenotypes. SVs verified via PCR are indicated with *. Putative causal mutations were identified by sequencing two plants with a phenotype and one relative without a phenotype. Only one mutant plant was sequenced for the *variegated* phenotype. This plant was green but produced ~25% *variegated* progeny. Three PCR-verified mutations in (B) are described in (C-E). **(C)** A genome browser snapshot with reads identifying a 184 bp insertion on Chr5 in the *short-internode dwarf* line. **(D)** Reads in the genome browser snapshot indicate an 81 bp deletion on Chr5 in the *variegated* line. **(E)** Snapshot of long-read alignments to reference *IMMUTANS*, showing a 37 bp deletion in the third exon. Deleted sequence shown in grey and coverage is shown above read alignments. The deletion is heterozygous in the sequenced plant, which was green and produced ~25% *variegated* progeny. **(F)** A ribbon plot with alignments indicating a 1.52 Mb chromosome 2 inversion in the *BR-like dwarf* line (top) and a detailed view of a representative read spanning the right breakend. The right breakend is in the 5'UTR of the *AS1* gene.

Figure S4: Duplications detected in the 26A lineage. Each track of the circos plot represents genome-wide read coverage analysis for an M2 sibling generated by selfing the 26A M1 mutant. Chromosome 1 duplications are shared by three M2 siblings while a Chromosome 3 duplication is shared by two of three siblings.

mapped to Nucleolar Organizer Regions (NOR), plastid genomes, and a small subset of other genomic loci; however these SVs were also found in control plants and read depth at NORs among etoposide-treated lines was comparable to that in control lines (Figure 3C), suggesting that these SVs represent mapping artifacts or naturally occurring SVs extant in wild-type plants. Based on these observations, we excluded SVs mapping to these regions from further analysis. To further filter for etoposide-independent SVs, those that were found in four or more etoposide-derived lineages were excluded, although we cannot exclude the possibility that these represent common fragile sites. After applying these stringent filters, we identified 27 unique SVs including 16 deletions and three duplications that ranged between 35 bp and 950 bp in size (Figure S5, Figure S6, Table S9). We also identified eight inversions that span between 124 bp and 4 Mb in length (Figure S5, Figure S7, Table S9). In sum, the coverage-based approach and Lumpy express analysis suggest that etoposide treatment induces structural variation in Arabidopsis.

Long-read sequencing can identify structural variants that are not easily resolved or cannot be identified by short-read sequencing [64]. Using Nanopore sequencing technology, we sequenced the genomes of two M3 plants with the *BR-like dwarf*, *short-internode dwarf,* or *virescent* phenotypes; an M3 relative of each corresponding line that lacked the phenotype; and a single M3 plant from the *variegated* line that was green but produced a proportion of variegated progeny (Table S10). To control for structural variation already present in our wild-type lab Col-0 stock compared to the reference Col-0 genome, we also sequenced two M2 plants that were the progeny of M1 plants grown on DMSO only and an untreated Col-0 plant. Reads were aligned to the Col-CEN v1.2 reference genome [65] and SVs were called using cuteSV [66,67]. Structural variants present in the DMSO and/or Col-0 controls, present in more than one M1 line, or with a quality less than 20 were discarded. In total we identified 12 SVs: seven deletions (ranging from 37 bp to 176 bp), two insertions (71 bp and 184 bp), two inversions (1.52 Mb and 3.54 Mb), and one translocation (Figure 4B; Table S11). We also chose to confirm several of the SVs via PCR (Figure S8): a 184 bp insertion with homology to an intergenic region of chromosome 2 present in the *short-internode dwarf* line (Figure 4C); a 37 bp and an 81 bp deletion present in the *variegated* line (Figure 4D,E); and a 1.52

Figure S5: Structural variants identified by Lumpy Express in short-read data. A total of 27 SVs (16 deletions, 3 duplications, and 8 inversions) were identified. Eleven sequenced samples did not have any detected SVs.

Figure S6: Examples of deletions in an etoposide-treated line identified by Lumpy Express using short reads. JBrowse snapshots of split reads and all reads overlapping *LHP1* **(A)** and *FKBP12* **(B)** identifies short deletions in two independent lines.

Line 12_11

Figure S7: An example of an intrachromosomal inversion in an etoposide-treated line detected by Lumpy Express using short reads. Split reads and discordant reads indicate an intra-chromosomal inversion on Chr3. Black lines link split and discordant reads mapping to two different parts of the chromosome.

Figure S8: PCR verification of four strucutral variants identified via long-read sequencing. All plants sampled for PCR verification were relatives of the plants sequenced. **(A)** Diagram depicting primer design for verifcation of 1.52 Mb inversion identified in *BR-like dwarf* plants. **(B)** Image of gel electrophoresis of PCR products generated with different combinations of primers in (A); the inversion is present only in plants exhibiting the *BR-like dwarf* phenotype. **(C)** Gel image for verification of 184 bp insertion identified in *short-internode dwarf* plants. Primers were designed to flank the insertion. **(D)** Gel image for verification of 81 bp deletion identified in the plant sequenced from the *variegated* line. Primers were designed to flank the deletion; all plants tested are homozygous for the deletion, except for one plant (lane 3) for which the PCR failed. **(E)** Alignment of sequenced PCR products to *IMMUTANS* reference sequence. Primers were designed to flank the deletion. Individuals in the line exhibiting the *variegated* phenotype are homozygous for the deletion while green plants are either heterozygous or homozygous for the WT allele.

Mb inversion present on chromosome 2 in the *BR-like dwarf* line (Figure 4F). Together,

these data indicate that etoposide treatment generates a range of SV types and sizes.

Finally, we evaluated the relationships between structural variation identified by

Nanopore sequencing and gene expression in *BR-like dwarf, short-internode dwarf,* and

variegated mutants. Both sequenced *BR-like dwarf* plants contain a 1.52 Mb inversion

on chromosome 2 that is not present in non-phenotypic control plants of the same line

(Figure 4B,F, Table S11) and we investigated gene expression surrounding this locus

further. The 1.52 Mb inversion was significantly enriched for differentially expressed

genes, as determined by a permutation test (Figure S9A,B). Examining the inversion

breakpoints more closely, the right breakend separated the first 42 bp of the 5'UTR of

ASYMMETRIC LEAVES 1 (AS1; AT2G37630) from the rest of the gene (Figure 4F).

Near the left breakend of the inversion, the expression of an AP2/B3-like transcriptional

factor family protein (AT2G33720) was increased in leaves (Figure 2A, Figure 4F, Table

S3). Taken together, these data suggest that the 1.52 Mb inversion resulted in an

altered *AS1*-mediated transcript abundance, influencing leaf patterning in *BR-like dwarf*

plants.

Three SVs were common to the two *short-internode dwarf* plants that were sequenced

via Nanopore: a 176 bp deletion on chromosome 3, a 3.54 Mb inversion on

chromosome 4, and a 184 bp insertion on chromosome 5 (Figure 4B,C, Table S11).

The large inversion on chromosome 4 was not enriched for DEGs, and neither were

regions surrounding the deletion on chromosome 3 or the insertion on chromosome 5

(Figure S9C-L). However, since gene expression data was derived from only the leaves

of *short-internode dwarf* plants, we cannot exclude the possibility that these structural

variants have an impact on gene expression in other plant organs. No structural variants

were identified in *virescent* plants utilizing either short- or long-read data. Although it is

possible that no structural variation is present in this line, it is also possible that

virescent plants contain an SV that cannot be detected with our methods due to its

complexity or genomic location.

Figure S9: Permutation tests to determine if structural variants are associated with location of differentially expressed genes. (A) Log$_2$(fold change) of genes within and 1 Mb outside the 1.52 Mb inversion in *BR-like dwarf.* DEGs are in blue and SVs are shaded in light blue. Dark blue line represents locally estimated scatterplot smoothing to aid pattern visualization. **(B)** Permutation test to determine if the 1.52 Mb inversion in *BR-like dwarf* plants contains more DEGs than expected by chance. All permutation tests were performed by selecting 10,000 genomic regions of an equivalent size to the region of interest at random. Histogram plots the distribution of number of DEGs contained within the randomly selected regions. Red line marks the number of DEGs in the region of interest. P-values calculated by the probability that permuted values exceed the observed value; if p-value is < 0.05 we conclude the region of interest is enriched for DEGs. **(C)** Log$_2$(fold change) of genes within and 2 Mb outside the 3.54 Mb inversion in *short-internode dwarf.* DEGs are in pink and SVs are shaded in light pink. Dark pink line represents locally estimated scatterplot smoothing to aid pattern visualization. **(D)** Permutation test for 3.54 Mb inversion in *short-internode dwarf* plants. **(E)** Log$_2$(fold change) of genes 2 Mb outside 176 bp deletion in *short-internode dwarf.* DEGs are in pink and SVs are shaded in light pink. Dark pink line represents locally estimated scatterplot smoothing to aid pattern visualization. Permutation tests for **(F)** 5 kb, **(G)** 10 kb, and **(H)** 50 kb flanking the deletion on Chr3 in *short-internode dwarf* plants. **(I)** Log$_2$(fold change) of genes 2 Mb outside 184 bp insertion in *short-internode dwarf.* DEGs are in pink and SVs are shaded in light pink. Dark pink line represents locally estimated scatterplot smoothing to aid pattern visualization. Permutation tests for **(J)** 5 kb, **(K)** 10 kb, and **(L)** 50 kb flanking the insertion on Chr5 in *short-internode dwarf* plants.

Long-read sequencing of the *variegated* line identified a 37 bp deletion in the third exon

of *IM*; this caused a frame-shift resulting in a premature stop codon (Figure 4E).

Crossing plants of the *variegated* phenotype to known *immutans* mutants failed to

complement the variegated phenotype (32/32 variegated when crossed to *im-spotty*

CS3639 [68], 34/34 variegated when crossed to *im-spotty* CS73029 [69] and 42/42

variegated when crossed to *im* CS3157), indicating that the 37 bp deletion in *IM* is

indeed the causal locus for *variegation*.

In summary, our genomic analyses of mutants shows that etoposide treatment can

create novel plant traits via the induction of SVs including inversions, deletions,

duplications, and translocations.

Discussion

We describe a simple and convenient technique that enables researchers or breeders

with limited resources to induce structural variation. We found that early exposure of *A.*

thaliana to the Topo II inhibitor etoposide induces heritable, large-effect mutations at

high frequency. Examination of M2 plants found that 24/37 mutant lineages had at least

one out of six assessed phenotypes. By short- and long-read genomic sequencing of

mutagenized lines—including those that displayed novel phenotypes—we identified the

full spectrum of structural variations. Analysis of short-read data identified 2.53 SVs per

mutagenized M1 line whereas long-read data identified 3.25 SVs per mutagenized M1

line, with one event being detected in both short and long-read data. The lower number

of SVs identified by short-read data is consistent with the poorer performance of short-

read SV callers relative to callers using long-reads [64]. Our data suggest that mutation

rate and type induced by etoposide exposure is comparable to mutant populations

generated by irradiation. Analysis of Arabidopsis seeds irradiated with γ rays or carbon

ions showed mutations occurring at a rate of 0.8-1.6 SVs per mutant lineage [70]. Thus,

etoposide treatment out-performed irradiation by producing more than 2.5 events per

mutagenized plant. A further increase in the number of mutations per M1 parent could

perhaps be attained either by prolonging exposure to etoposide beyond two weeks or

by germinating seeds on media containing a cocktail of etoposide and other drugs that

induce genomic instability such as topoisomerase I inhibitors [71] or the DNA damage-

inducing agent cisplatin [72].

Analysis of all detected SVs suggests that etoposide disproportionately impacts genes,

although our analytical approaches cannot conclusively identify all SVs in

heterochromatin. The SVs that we identified disrupt one or more genes simultaneously

or create novel cis-regulatory regions. The effect of this genic bias is striking; even in

our relatively small mutant population, multiple loss-of-function mutations were identified

in a single line. This suggests that etoposide treatment can be used to create novel

polygenic traits.

What explains this bias for mutations in genes? Topoisomerase II has been shown to

associate with actively transcribed regions and open chromatin regions [42]. This

association with actively transcribed regions suggests a future method to enrich for

mutations in condition-specific genetic pathways by simultaneously exposing seeds to

etoposide and another condition. For example, etoposide might promote mutagenesis of

immunity genes whose transcription is triggered when seeds are germinated in the

presence of microbial pathogens. In another scenario, seeds germinated in the

presence of a hormone and etoposide might accumulate mutations in genes that

respond to said hormone.

The inversions and translocations generated by this technique might also disrupt

meiosis and distort segregation, but this possibility has not been tested. Etoposide also

created duplications and deletions that could potentially alter gene dosage over mega-

base scale regions. In allopolyploids, such segmental deletions can provide a deficiency

chromosome that can be used to assess the relative contribution of homeologs to

different traits.

How do we identify genetic changes that are causative for phenotypes of interest?

Future studies will accelerate candidate-gene discovery by employing structural-variant

callers, *de novo* genome assembly-based approaches, and RNA-seq based mapping

[64]. Although our study did not aim to use RNA-seq to identify mutations, it provides an

example of how RNA-seq data in tandem with genome sequencing can help shortlist

potential causal mutations. In cases where a potential causative variant is not obvious,

these strategies can be combined with traditional genetic mapping approaches.

However, mapping-by-sequencing approaches might not be easily applicable to some

classes of mutants—for example, inversions or translocations can suppress

recombination and reduce the efficacy of mapping-by-sequencing.

In our current study, sequencing and close analysis of the genomes and transcriptomes

of *BR-like dwarf, short-internode dwarf, virescent,* and *variegated* plants allowed us to

link two of these phenotypes to strong candidate mutations. The *BR-like dwarf*

phenotype is likely caused by an inversion that disrupts the expression of

ASYMMETRIC LEAVES 1 and the *variegated* phenotype is caused by a 37 bp deletion

in *IMMUTANS.* Strikingly, nearly 60 years ago, the *asymmetric leaves* and *immutans*

phenotypes were amongst the first X-ray induced mutations isolated in *A. thaliana* by

Redei [73,74]. These results suggest that our easy-to-use chemical mutagenesis

approach provides plant biologists and breeders with a new tool that can replace

irradiation and reduce barriers to access tools necessary for the creation of structural

variation.

Materials and Methods

Drug treatment

Etoposide (Abcam AB120227) was dissolved in DMSO to obtain a 100 mM solution that

was added to 0.5x Murashige and Skoog (MS) media (supplemented with 2% sucrose

and Phytoagar) to final concentrations of 0, 20, 40, 80, 160, 320, and 640 µM. We

noticed decreased solubility of etoposide in aqueous solutions at concentrations beyond

160 µM. Wild-type Col-0 seeds were sterilized by incubating in 2% (v/v) PPM (Plant Cell

Technologies) for three days at 4°C and plated on MS media containing etoposide or

DMSO. Seeds were germinated and grown on this media for two weeks, and seedlings

were examined for signs of stunted root or shoot growth. Two- to three-week-old

seedlings (M1 plants), including several showing stunted growth, were transferred to

soil and allowed to set seed. M2 seeds were collected from mutagenized M1 plants.

Plant Growth Conditions

Arabidopsis thaliana was grown in a Conviron growth chamber at 22°C with 16 h of light per day (120 µmol).

Phenotyping

Between 25 and 75 seeds from each M1 line were planted over three phases and M2 plants with visible phenotypes were manually identified. To identify mutants with leaf-shape differences, we measured leaf shape and size in mutant lineages where M2/M3 populations included three or more individuals displaying similar mutant phenotypes. Leaves were excised at the base of the petiole, arranged in decreasing size order on white paper, and scanned at 600 dpi on an Epson V600 scanner. Six automated leaf shape measurements were obtained (blade area, circularity, length, perimeter, width, and petiole length) using the LeafJ plugin for ImageJ software [75]. Leaf shape was manually traced in cases where leaf boundaries were not distinguished by ImageJ.

RNA sequencing and analysis

RNA sequencing was conducted on M3 generation plants. Six plants of each *BR-like dwarf, short-internode dwarf*, and *virescent* phenotypes, as well as four non-phenotypic individuals corresponding to sibling lines of each phenotype were sequenced. For the fraction variegated line, thirteen green plants (three of which produced ~25% variegated progeny) and one white plant were sampled. Nine M3 progeny of DMSO-treated plants were included as controls.

Mature leaf tissue from 32-day-old plants was flash frozen in liquid nitrogen. Frozen tissue was ground into a powder with a Tissuelyser II (QIAGEN; 30 beats/sec for 30 sec). RNA was extracted using an RNeasy Plant Mini Kit (QIAGEN). Library preparation was performed at the MIT BioMicro Center using an NEBNext Ultra II Directional RNA Library Prep Kit for Illumina (New England Biolabs). Samples were multiplexed on one lane of an Illumina NovaSeq 6000 S4 flow cell and sequenced at the MIT BioMicro Center with 150 bp paired end reads.

Trim Galore [76] was used to remove adapters and low-quality ends from reads (--

paired --retain_unpaired -q 25). Salmon [77] was used to quantify the expression of

transcripts, utilizing the Araport11 transcriptome as the reference and the Col-CEN v1.2

genome [65] as decoy sequences. Five samples were identified as outliers and

removed from the dataset, based upon the PCAgrid function in the R package rrcov

[78,79] and preliminary analyses. DESeq2 [80] was used to quantify differential

expression; for a gene to be considered it must have at least 10 normalized counts in

four or more samples. Each phenotype of interest was analyzed independently;

DESeq2 design (*~ treatment + phenotype*) incorporated both treatment (etoposide or

DMSO) and phenotype (present or absent). The R package clusterProfiler [81] was

used to conduct gene set enrichment analysis. Data was visualized utilizing R and

packages ggplot2, EnhancedVolcano [82] , and pheatmap [83].

Illumina sequencing and identification of structural variants

DNA was extracted from the rosette leaves of M2 and M3 plants using a CTAB protocol.

Up to 15 ng of genomic DNA was used by the BioMicro Center at MIT to construct DNA

sequencing libraries using the Nextera DNA Flex library preparation kit. Libraries were

multiplexed and sequenced in 150 bp paired-end format using an Illumina NovaSeq

6000 S4. The first 10 bp of each read as well as any adaptor sequence was removed

using Trimmomatic [84] and aligned to the Col-0 Centromere assembly [65] using BWA-

MEM [85] (*bwa mem -R "@RG\tID:id\tSM:sample\tLB:lib" reference.fasta*

sample.R1.fastq sample.R2.fastq \| samblaster --excludeDups --addMateTags --

maxSplitCount 2 --minNonOverlap 20). These alignments were used to identify

structural variants by two separate methods. Larger deletions and duplications were

identified using a method proposed by the Comai Lab [61,62]. Coverage was calculated

for each 100 kb window for properly paired and aligned reads using Bedtools [86]

(*bedtools coverage -a 100kb_windows.bed -b sample_paired_sorted.bam -counts*) and

read-counts/window for each line was normalized to library size as reads per million.

The median normalized read-counts for all lines for each window was calculated. As

most lines are not expected to share a structural variant, this median read depth is

considered to be the wild-type genome dosage and is set to a ploidy or chromosomal

dosage of two. The chromosomal dosage of any window for a given line is then

calculated as (*normalized read-counts / median read-counts) x2.* Windows with a

coverage score ≥3 were considered heterozygous duplications, ≥4 to be homozygous

duplications, ≤1 to be heterozygous deletion and ≤0 to be a homozygous deletion.

In addition, we also identified structural variation using a benchmarked Lumpy Express

pipeline [12,63]. Briefly, Samtools [87,88] was used to identify discordant and split

reads from the SAM file that was output by BWA [85]. Files containing discordant reads,

split reads, and all reads were used by Lumpy Express to identify putative structural

variants. Due to mapping artifacts, Lumpy Express and other variant callers often

identify false-positive structural variant candidates at repetitive regions [89]. To address

this problem, we filtered out variants from ChrC, ChrM, and the NOR on chromosome 2.

To remove SVs present in lab wild-type Col-0 stocks, we removed SVs within 2 kb of

any SV found in lines only treated with DMSO, or SVs shared by four or more unrelated

etoposide-treated mutant lineages. Finally, only those deletions and duplications that

were supported by split-reads were retained in the final list of SVs.

To assess if etoposide causes single nucleotide variants, we used BWA-MEM to align

reads from 32 M2/M3 progeny of etoposide-treated plants and four M2 progeny of

control plants treated with DMSO to the Col-0 centromere assembly [65]. Aligned reads

were sorted and converted to variant calling format (*bcftools mpileup -O b -o file.bcf -f*

ref.fasta file.sorted.bam) followed by SNP calling (*bcftools call --ploidy 2 -m -v -o file.vcf*

file.bcf) [87]. Variants that were called by bcftools were filtered to retain only calls with a

quality score higher than or equal to 40 (*bcftools view -O z -o filtered.vcf.gz -e*

'QUAL<=40' in.vcf.gz). Variants shared by all untreated lines were extracted (*bcftools*

isec -n=5 -p common_positions.vcf input_files_filtered.vcf.gz), and then excluded from

all other treated lines (*bcftools view -T common_positions.vcf filtered_files.vcf -O z -o*

untreated_filtered_files.vcf).

Nanopore sequencing and identification of structural variants

Two mature leaves from 32-day-old M3 plants were flash frozen in liquid nitrogen.

Frozen tissue was ground into a powder with a Tissuelyser II (QIAGEN; 30 beats/sec for

1 min). High molecular weight DNA was extracted using a Wizard HMW DNA Extraction

Kit (Promega). DNA was sheared by passing through a 23G needle twice.

Contaminating RNA and proteins were removed using RNase A (New England Biolabs)

and Proteinase K (New England Biolabs). DNA was further cleaned and small

fragments removed using a modified SPRI bead mixture in which the buffer of

Agencourt AMPure XP beads was exchanged with a custom buffer of 10 mM Tris-HCl,

1 mM EDTA pH 8, 1.6 M NaCl, 11% w/v PEG 8000 [90]. To perform size selection, in

brief, 0.7x volume of modified bead solution was added to each DNA sample, then

incubated on a nutator for 10 min. Samples were placed on a magnetic stand to bind

the beads to the side of the tube, then washed twice with 70% ethanol. Beads were not

allowed to dry before adding nuclease-free water to elute the DNA. To increase yield

and remove residual ethanol, samples were incubated at 50°C for 2 min and then at

room temperature for two hours with the lids open.

All library preparation and sequencing was performed at the MIT BioMicro Center. For

two samples showing the *BR-like dwarf* phenotype, high molecular weight DNA

extractions had low yield. For these samples and corresponding controls (individual

from sibling line without dwarf phenotype and individual from a line treated with DMSO),

library preparation was performed utilizing a Rapid Barcoding Kit V10 (Oxford Nanopore

Technologies; SQK-RBK110-96). These four samples were multiplexed and sequenced

on a single PromethION R9.4.1 flow cell (Oxford Nanopore Technologies; FLO-

PRO002). For all other samples, library preparation was performed with a Native

Barcoding Kit V14 (Oxford Nanopore Technologies; SQK-NBD114-24). These samples

were multiplexed into groups of three and each group was sequenced on its own

PromethION R10.4.1 flow cell (Oxford Nanopore Technologies; FLO-PRO114M). Both

runs utilized the following software versions for sequencing and base-calling: MinKNOW

23.04.6, Bream 7.5.10, Configuration 5.5.14, Guppy 6.5.7, and MinKNOW Core 5.5.5.

Nanopore reads were filtered utilizing Filtlong v0.2.1 [91] to select the best reads in

terms of length and quality, up to a max coverage of 50x (*--min_length 1000 --*

min_window_q 40 --trim --split 1500 --target_bases 6605000000). Filtered reads were

mapped to the Col-CEN v1.2 genome [65] using Vulcan dual-mode alignment pipeline

[92], which leverages both minimap2 [93] and NGMLR [94], using default settings for

ONT reads. Structural variants in each sample were detected with cuteSV *(--*

max_cluster_bias_INS 100 --diff_ratio_merging_INS 0.3 --max_cluster_bias_DEL 100 --

diff_ratio_merging_DEL 0.3 --min_size 30 --max_size -1 –genotype; [66,67]). SVs

identified in all samples were merged with SURVIVOR (*merge 1000 1 1 1 0 30;* [95]),

then samples were genotyped with cuteSV for all SVs identified (*--min_mapq 20 --*

max_cluster_bias_INS 100 --diff_ratio_merging_INS 0.3 --max_cluster_bias_DEL 100 --

diff_ratio_merging_DEL 0.3 --min_size 30 --max_size -1). Structural variants present in

the DMSO and/or Col-0 controls, or present in more than one phenotypic line, or with a

quality less than 20 were removed. Data was visualized in R using ggplot2.

Data Availability

536 Short-read DNA sequencing, long-read DNA sequencing, and mRNA-seq data are

537 available in GEO records GSE284455, GSE284456 and GSE284457.

Declaration of Interest

539 Whitehead Institute has filed a patent application related to this work.

Acknowledgements

We thank Aysha Hussain, Jaza Alam, and Lingfeng Shi for assistance with phenotyping,

Dyna Louis for assistance with verifying structural variants, and Xinlei Gao for

assistance with statistical testing. This research was funded by grants to MG from The

Abdul Latif Jameel Water and Food Systems Lab at the Massachusetts Institute of

Technology, from the Professor Amar G. Bose Research Grants, and from a gift to MG

from the Dr. Vincent J. Ryan Orphan Plant Project. MG is an Investigator of the Howard

Hughes Medical Institute.

Supplemental Information

Figure S1: Additional phenotyping of progeny of etoposide-treated plants.

Figure S2: Segregation of the *variegated* phenotype from green parents.

Figure S3: GO term enrichment for genes differentially expressed between plants

exhibiting a phenotype and a genetic relative without a phenotype.

Figure S4: Duplications detected in the 26A lineage.

Figure S5: Structural variants identified by Lumpy Express in short-read data.

Figure S6: Examples of deletions in an etoposide-treated line identified by Lumpy

Express using short reads

Figure S7: An example of an intrachromosomal inversion in an etoposide-treated line

identified by Lumpy Express using short reads.

Figure S8: PCR verification of four structural variants identified via long-read

sequencing.

Figure S9: Permutation tests to determine if structural variants are associated with

location of differentially expressed genes.

Table S1: List of phenotypes observed in M2 progeny of etoposide-treated plants.

Table S2: List of RNA-Seq libraries.

Table S3: DESeq2 output identifying mis-regulated genes in *BR-like dwarf.*

Table S4: DESeq2 output identifying mis-regulated genes in *short-internode dwarf.*

Table S5: DESeq2 output identifying mis-regulated genes in virescent.

Table S6: List of samples sequenced with Illumina.

Table S7: Relationship between the sequenced lines, treatments, and nature of

sequencing.

Table S8: Deletions and duplications identified by read coverage analysis.

Table S9: Structural variants detected by LUMPY Express using short-read sequencing.

Table S10: List of samples sequenced with Nanopore.

Table S11: Structural variants identified using Nanopore long-read data.

References

1. Brown EJ, Nguyen AH, Bachtrog D. The Drosophila Y Chromosome Affects Heterochromatin Integrity Genome-Wide. Molecular Biology and Evolution. 2020;37: 2808–2824. doi:10.1093/molbev/msaa082

2. Henikoff S. Dosage-dependent modification of position-effect variegation in Drosophla. BioEssays. 1996;18: 401–409. doi:10.1002/bies.950180510

3. Murphy PJ, Berger F. The chromatin source-sink hypothesis: a shared mode of chromatin-mediated regulations. Development. 2023;150: dev201989. doi:10.1242/dev.201989

4. Chiang C, Scott AJ, Davis JR, Tsang EK, Li X, Kim Y, et al. The impact of structural variation on human gene expression. Nat Genet. 2017;49: 692–699. doi:10.1038/ng.3834

5. Alonge M, Wang X, Benoit M, Soyk S, Pereira L, Zhang L, et al. Major Impacts of Widespread Structural Variation on Gene Expression and Crop Improvement in Tomato. Cell. 2020;182: 145-161.e23. doi:10.1016/j.cell.2020.05.021

6. Zhou Y, Minio A, Massonnet M, Solares E, Lv Y, Beridze T, et al. The population genetics of structural variants in grapevine domestication. Nat Plants. 2019;5: 965–979. doi:10.1038/s41477-019-0507-8

7. Tian S, Ge J, Ai G, Jiang J, Liu Q, Chen X, et al. A 2.09 Mb fragment translocation on chromosome 6 causes abnormalities during meiosis and leads to less seed watermelon. Horticulture Research. 2021;8: 256. doi:10.1038/s41438-021-00687-9

8. Wang X, Liu Z, Zhang F, Xiao H, Cao S, Xue H, et al. Integrative genomics reveals the polygenic basis of seedlessness in grapevine. Current Biology. 2024;34: 3763-3777.e5. doi:10.1016/j.cub.2024.07.022

9. Chen J, Liu Y, Liu M, Guo W, Wang Y, He Q, et al. Pangenome analysis reveals genomic variations associated with domestication traits in broomcorn millet. Nat Genet. 2023;55: 2243–2254. doi:10.1038/s41588-023-01571-z

10. Tao Y, Luo H, Xu J, Cruickshank A, Zhao X, Teng F, et al. Extensive variation within the pan-genome of cultivated and wild sorghum. Nat Plants. 2021;7: 766–773. doi:10.1038/s41477-021-00925-x

11. Yan H, Sun M, Zhang Z, Jin Y, Zhang A, Lin C, et al. Pangenomic analysis identifies structural variation associated with heat tolerance in pearl millet. Nat Genet. 2023;55: 507–518. doi:10.1038/s41588-023-01302-4

12. Göktay M, Fulgione A, Hancock AM. A New Catalog of Structural Variants in 1,301 A. thaliana Lines from Africa, Eurasia, and North America Reveals a Signature of

Balancing Selection at Defense Response Genes. Molecular Biology and
Evolution. 2021;38: 1498–1511. doi:10.1093/molbev/msaa309

13. Hufford MB, Seetharam AS, Woodhouse MR, Chougule KM, Ou S, Liu J, et al. De
novo assembly, annotation, and comparative analysis of 26 diverse maize
genomes. Science. 2021;373: 655–662. doi:10.1126/science.abg5289

14. Mizuno H, Katagiri S, Kanamori H, Mukai Y, Sasaki T, Matsumoto T, et al.
Evolutionary dynamics and impacts of chromosome regions carrying R-gene
clusters in rice. Sci Rep. 2020;10: 872. doi:10.1038/s41598-020-57729-w

15. Katz E, Li J-J, Jaegle B, Ashkenazy H, Abrahams SR, Bagaza C, et al. Genetic
variation, environment and demography intersect to shape Arabidopsis defense
metabolite variation across Europe. Schuman MC, Korte A, editors. eLife. 2021;10:
e67784. doi:10.7554/eLife.67784

16. Qin P, Lu H, Du H, Wang H, Chen W, Chen Z, et al. Pan-genome analysis of 33
genetically diverse rice accessions reveals hidden genomic variations. Cell.
2021;184: 3542-3558.e16. doi:10.1016/j.cell.2021.04.046

17. Oikawa T, Maeda H, Oguchi T, Yamaguchi T, Tanabe N, Ebana K, et al. The Birth
of a Black Rice Gene and Its Local Spread by Introgression. Plant Cell. 2015;27:
2401–2414. doi:10.1105/tpc.15.00310

18. Guo J, Cao K, Deng C, Li Y, Zhu G, Fang W, et al. An integrated peach genome
structural variation map uncovers genes associated with fruit traits. Genome Biol.
2020;21: 258. doi:10.1186/s13059-020-02169-y

19. Hu Y, Colantonio V, Müller BSF, Leach KA, Nanni A, Finegan C, et al. Genome
assembly and population genomic analysis provide insights into the evolution of
modern sweet corn. Nat Commun. 2021;12: 1227. doi:10.1038/s41467-021-21380-
4

20. Kramer V, Shaw JR, Senior ML, Hannah LC. The sh2-R allele of the maize
shrunken-2 locus was caused by a complex chromosomal rearrangement. Theor
Appl Genet. 2015;128: 445–452. doi:10.1007/s00122-014-2443-3

21. Du Y, Luo S, Li X, Yang J, Cui T, Li W, et al. Identification of Substitutions and
Small Insertion-Deletions Induced by Carbon-Ion Beam Irradiation in Arabidopsis
thaliana. Front Plant Sci. 2017;8. doi:10.3389/fpls.2017.01851

22. Henry IM, Zinkgraf MS, Groover AT, Comai L. A System for Dosage-Based
Functional Genomics in Poplar. The Plant Cell. 2015;27: 2370–2383.
doi:10.1105/tpc.15.00349

23. Ichida H, Morita R, Shirakawa Y, Hayashi Y, Abe T. Targeted exome sequencing of
unselected heavy-ion beam-irradiated populations reveals less-biased mutation

characteristics in the rice genome. The Plant Journal. 2019;98: 301–314.
doi:10.1111/tpj.14213

24. Ishii K, Kazama Y, Hirano T, Fawcett JA, Sato M, Hirai MY, et al. Genomic view of
heavy-ion-induced deletions associated with distribution of essential genes in
Arabidopsis thaliana. Front Plant Sci. 2024;15. doi:10.3389/fpls.2024.1352564

25. Jankowicz-Cieslak J, Hofinger BJ, Jarc L, Junttila S, Galik B, Gyenesei A, et al.
Spectrum and Density of Gamma and X-ray Induced Mutations in a Non-Model
Rice Cultivar. Plants (Basel). 2022;11: 3232. doi:10.3390/plants11233232

26. Ma L, Kong F, Sun K, Wang T, Guo T. From Classical Radiation to Modern
Radiation: Past, Present, and Future of Radiation Mutation Breeding. Front Public
Health. 2021;9: 768071. doi:10.3389/fpubh.2021.768071

27. Havlickova L, He Z, Berger M, Wang L, Sandmann G, Chew YP, et al. Genomics of
predictive radiation mutagenesis in oilseed rape: modifying seed oil composition.
Plant Biotechnol J. 2024;22: 738–750. doi:10.1111/pbi.14220

28. Xiong H, Guo H, Fu M, Xie Y, Zhao L, Gu J, et al. A large-scale whole-exome
sequencing mutant resource for functional genomics in wheat. Plant Biotechnol J.
2023;21: 2047–2056. doi:10.1111/pbi.14111

29. Zhao Z, Liu Z, Zhou Y, Wang J, Zhang Y, Yu X, et al. Creation of cotton mutant
library based on linear electron accelerator radiation mutation. Biochem Biophys
Rep. 2022;30: 101228. doi:10.1016/j.bbrep.2022.101228

30. Li F, Shimizu A, Nishio T, Tsutsumi N, Kato H. Comparison and Characterization of
Mutations Induced by Gamma-Ray and Carbon-Ion Irradiation in Rice (Oryza
sativa L.) Using Whole-Genome Resequencing. G3 Genes|Genomes|Genetics.
2019;9: 3743–3751. doi:10.1534/g3.119.400555

31. Guo W, Comai L, Henry IM. Chromoanagenesis from radiation-induced genome
damage in Populus. PLOS Genetics. 2021;17: e1009735.
doi:10.1371/journal.pgen.1009735

32. Bolon Y-T, Stec AO, Michno J-M, Roessler J, Bhaskar PB, Ries L, et al. Genome
resilience and prevalence of segmental duplications following fast neutron
irradiation of soybean. Genetics. 2014;198: 967–981.
doi:10.1534/genetics.114.170340

33. Feng Z, Du Y, Chen J, Chen X, Ren W, Wang L, et al. Comparison and
Characterization of Phenotypic and Genomic Mutations Induced by a Carbon-Ion
Beam and Gamma-ray Irradiation in Soybean (Glycine max (L.) Merr.).
International Journal of Molecular Sciences. 2023;24: 8825.
doi:10.3390/ijms24108825

34.   Yasui Y, Mori M, Aii J, Abe T, Matsumoto D, Sato S, et al. S-LOCUS EARLY
683   FLOWERING 3 Is Exclusively Present in the Genomes of Short-Styled Buckwheat
684   Plants that Exhibit Heteromorphic Self-Incompatibility. PLOS ONE. 2012;7:
685   e31264. doi:10.1371/journal.pone.0031264

686   35.   Koide Y, Ogino A, Yoshikawa T, Kitashima Y, Saito N, Kanaoka Y, et al. Lineage-
687   specific gene acquisition or loss is involved in interspecific hybrid sterility in rice.
688   Proceedings of the National Academy of Sciences. 2018;115: E1955–E1962.
689   doi:10.1073/pnas.1711656115

690   36.   Olsen O, Wang X, Von Wettstein D. Sodium azide mutagenesis: preferential
691   generation of A.T-->G.C transitions in the barley Ant18 gene. Proc Natl Acad Sci
692   USA. 1993;90: 8043–8047. doi:10.1073/pnas.90.17.8043

693   37.   Tanaka H, Muramoto N, Sugimoto H, Oda AH, Ohta K. Extended TAQing system
694   for large-scale plant genome reorganization. The Plant Journal. 2020;103: 2139–
695   2150. doi:10.1111/tpj.14888

696   38.   Wu C-C, Li T-K, Farh L, Lin L-Y, Lin T-S, Yu Y-J, et al. Structural Basis of Type II
697   Topoisomerase Inhibition by the Anticancer Drug Etoposide. Science. 2011;333:
698   459–462. doi:10.1126/science.1204117

699   39.   Ban Y, Ho C-W, Lin R-K, Lyu YL, Liu LF. Activation of a Novel Ubiquitin-
700   Independent Proteasome Pathway when RNA Polymerase II Encounters a Protein
701   Roadblock. Molecular and Cellular Biology. 2013;33: 4008–4016.
702   doi:10.1128/MCB.00403-13

703   40.   Canela A, Maman Y, Huang SN, Wutz G, Tang W, Zagnoli-Vieira G, et al.
704   Topoisomerase II-Induced Chromosome Breakage and Translocation Is
705   Determined by Chromosome Architecture and Transcriptional Activity. Molecular
706   Cell. 2019;75: 252-266.e8. doi:10.1016/j.molcel.2019.04.030

707   41.   Tammaro M, Barr P, Ricci B, Yan H. Replication-Dependent and Transcription-
708   Dependent Mechanisms of DNA Double-Strand Break Induction by the
709   Topoisomerase 2-Targeting Drug Etoposide. PLOS ONE. 2013;8: e79202.
710   doi:10.1371/journal.pone.0079202

711   42.   Uusküla-Reimand L, Wilson MD. Untangling the roles of TOP2A and TOP2B in
712   transcription and cancer. Science Advances. 2022;8: eadd4920.
713   doi:10.1126/sciadv.add4920

714   43.   Amoiridis M, Verigos J, Meaburn K, Gittens WH, Ye T, Neale MJ, et al. Inhibition of
715   topoisomerase 2 catalytic activity impacts the integrity of heterochromatin and
716   repetitive DNA and leads to interlinks between clustered repeats. Nat Commun.
717   2024;15: 5727. doi:10.1038/s41467-024-49816-7

718   44.   Marchetti F, Bishop JB, Lowe X, Generoso WM, Hozier J, Wyrobek AJ. Etoposide
719   induces heritable chromosomal aberrations and aneuploidy during male meiosis in

720  the mouse. Proceedings of the National Academy of Sciences. 2001;98: 3952–
721  3957. doi:10.1073/pnas.061404598

722 45. Quispe-Tintaya W, Gorbacheva T, Lee M, Makhortov S, Popov VN, Vijg J, et al.
723  Quantitative detection of low-abundance somatic structural variants in normal cells
724  by high throughput sequencing. Nat Methods. 2016;13: 584–586.
725  doi:10.1038/nmeth.3893

726 46. Sugimoto-Shirasu K, Roberts GR, Stacey NJ, McCann MC, Maxwell A, Roberts K.
727  RHL1 is an essential component of the plant DNA topoisomerase VI complex and
728  is required for ploidy-dependent cell growth. Proceedings of the National Academy
729  of Sciences. 2005;102: 18736–18741. doi:10.1073/pnas.0505883102

730 47. Samanta A, Maity TR, Das S, Datta AK, Datta S. Effect of etoposide on grass pea
731  DNA topoisomerase II: an in silico, in vivo, and in vitro assessments. Bull Natl Res
732  Cent. 2019;43: 170. doi:10.1186/s42269-019-0217-4

733 48. Parra-Nunez P, Cooper C, Sanchez-Moran E. The Role of DNA Topoisomerase
734  Binding Protein 1 (TopBP1) in Genome Stability in Arabidopsis. Plants. 2021;10.
735  doi:10.3390/plants10122568

736 49. Żabka A, Winnicki K, Polit JT, Maszewski J. The effects of anti-DNA topoisomerase
737  II drugs, etoposide and ellipticine, are modified in root meristem cells of *Allium*
738  *cepa* by MG132, an inhibitor of 26S proteasomes. Plant Physiology and
739  Biochemistry. 2015;96: 72–82. doi:10.1016/j.plaphy.2015.07.016

740 50. Noguchi T, Fujioka S, Choe S, Takatsuto S, Yoshida S, Yuan H, et al.
741  Brassinosteroid-Insensitive Dwarf Mutants of Arabidopsis Accumulate
742  Brassinosteroids1. Plant Physiology. 1999;121: 743–752.
743  doi:10.1104/pp.121.3.743

744 51. Byrne ME, Simorowski J, Martienssen RA. ASYMMETRIC LEAVES1 reveals knox
745  gene redundancy in Arabidopsis. Development. 2002;129: 1957–1965.
746  doi:10.1242/dev.129.8.1957

747 52. Ha CM, Jun JH, Nam HG, Fletcher JC. BLADE-ON-PETIOLE1 and 2 Control
748  Arabidopsis Lateral Organ Fate through Regulation of LOB Domain and Adaxial-
749  Abaxial Polarity Genes. The Plant Cell. 2007;19: 1809–1825.
750  doi:10.1105/tpc.107.051938

751 53. Ikezaki M, Kojima M, Sakakibara H, Kojima S, Ueno Y, Machida C, et al. Genetic
752  networks regulated by ASYMMETRIC LEAVES1 (AS1) and AS2 in leaf
753  development in Arabidopsis thaliana: KNOX genes control five morphological
754  events. The Plant Journal. 2010;61: 70–82. doi:10.1111/j.1365-313X.2009.04033.x

755 54. Jun JH, Ha CM, Fletcher JC. BLADE-ON-PETIOLE1 coordinates organ
756  determinacy and axial polarity in arabidopsis by directly activating ASYMMETRIC
757  LEAVES2. Plant Cell. 2010;22: 62–76. doi:10.1105/tpc.109.070763

55. Sun Y, Zhou Q, Zhang W, Fu Y, Huang H. ASYMMETRIC LEAVES1, an
Arabidopsis gene that is involved in the control of cell differentiation in leaves.
Planta. 2002;214: 694–702. doi:10.1007/s004250100673

56. Qiu K, Li Z, Yang Z, Chen J, Wu S, Zhu X, et al. EIN3 and ORE1 Accelerate
Degreening during Ethylene-Mediated Leaf Senescence by Directly Activating
Chlorophyll Catabolic Genes in Arabidopsis. PLOS Genetics. 2015;11: e1005399.
doi:10.1371/journal.pgen.1005399

57. Zhong S, Zhao M, Shi T, Shi H, An F, Zhao Q, et al. EIN3/EIL1 cooperate with
PIF1 to prevent photo-oxidation and to promote greening of Arabidopsis seedlings.
Proceedings of the National Academy of Sciences. 2009;106: 21431–21436.
doi:10.1073/pnas.0907670106

58. Rédei GP. Biochemical aspects of a genetically determined variegation in
Arabidopsis. Genetics. 1967;56: 431–443. doi:10.1093/genetics/56.3.431

59. Rédei GP. Somatic Instability Caused by a Cysteine-Sensitive Gene in
Arabidopsis. Science. 1963;139: 767–769.

60. Bromberg KD, Burgin AB, Osheroff N. A Two-drug Model for Etoposide Action
against Human Topoisomerase IIα *. Journal of Biological Chemistry. 2003;278:
7406–7412. doi:10.1074/jbc.M212056200

61. Tan E, Comai L, Henry I. Chromosome Dosage Analysis in Plants Using Whole
Genome Sequencing. BIO-PROTOCOL. 2016;6. doi:10.21769/BioProtoc.1854

62. Tan EH, Henry IM, Ravi M, Bradnam KR, Mandakova T, Marimuthu MP, et al.
Catastrophic chromosomal restructuring during genome elimination in plants. eLife.
2015;4: e06516. doi:10.7554/eLife.06516

63. Layer RM, Chiang C, Quinlan AR, Hall IM. LUMPY: a probabilistic framework for
structural variant discovery. Genome Biology. 2014;15: R84. doi:10.1186/gb-2014-
15-6-r84

64. Mahmoud M, Gobet N, Cruz-Dávalos DI, Mounier N, Dessimoz C, Sedlazeck FJ.
Structural variant calling: the long and the short of it. Genome Biology. 2019;20:
246. doi:10.1186/s13059-019-1828-7

65. Naish M, Alonge M, Wlodzimierz P, Tock AJ, Abramson BW, Schmücker A, et al.
The genetic and epigenetic landscape of the Arabidopsis centromeres. Science.
2021;374: eabi7489. doi:10.1126/science.abi7489

66. Jiang T, Liu Y, Jiang Y, Li J, Gao Y, Cui Z, et al. Long-read-based human genomic
structural variation detection with cuteSV. Genome Biology. 2020;21: 189.
doi:10.1186/s13059-020-02107-y

67. Jiang T, Cao S, Liu Y, Liu S, Liu B, Wang G, et al. Regenotyping structural variants
through an accurate force-calling method. 2022. doi:10.1101/2022.08.29.505534

68. Fu A, Park S, Rodermel S. Sequences Required for the Activity of PTOX
(IMMUTANS), a Plastid Terminal Oxidase: In vitro and in planta mutagenesis of
iron-binding sites and a conserved sequence that corresponds to exon 8*. Journal
of Biological Chemistry. 2005;280: 42489–42496. doi:10.1074/jbc.M508940200

69. Wetzel CM, Jiang C-Z, Meehan LJ, Voytas DF, Rodermel SR. Nuclear—organelle
interactions: the immutans variegation mutant of Arabidopsis is plastid autonomous
and impaired in carotenoid biosynthesis. The Plant Journal. 1994;6: 161–175.
doi:10.1046/j.1365-313X.1994.6020161.x

70. Hase Y, Satoh K, Kitamura S. Comparative analysis of seed and seedling
irradiation with gamma rays and carbon ions for mutation induction in Arabidopsis.
Front Plant Sci. 2023;14. doi:10.3389/fpls.2023.1149083

71. Pommier Y. Topoisomerase I inhibitors: camptothecins and beyond. Nat Rev
Cancer. 2006;6: 789–802. doi:10.1038/nrc1977

72. Weimer AK, Biedermann S, Harashima H, Roodbarkelari F, Takahashi N, Foreman
J, et al. The plant-specific CDKB1-CYCB1 complex mediates homologous
recombination repair in Arabidopsis. The EMBO Journal. 2016;35: 2068–2086.
doi:10.15252/embj.201593083

73. Rédei GP. Non-Mendelian Megagametogenesis in Arabidopsis. Genetics. 1965;51:
857–872.

74. Rodermel S. Arabidopsis Variegation Mutants. Arabidopsis Book. 2002;1: e0079.
doi:10.1199/tab.0079

75. Maloof JN, Nozue K, Mumbach MR, Palmer CM. LeafJ: An ImageJ Plugin for
Semi-automated Leaf Shape Measurement. JoVE. 2013; 50028.
doi:10.3791/50028

76. Krueger F. FelixKrueger/TrimGalore. 2024. Available:
https://github.com/FelixKrueger/TrimGalore

77. Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. Salmon provides fast and
bias-aware quantification of transcript expression. Nat Methods. 2017;14: 417–419.
doi:10.1038/nmeth.4197

78. Chen X, Zhang B, Wang T, Bonni A, Zhao G. Robust principal component analysis
for accurate outlier sample detection in RNA-Seq data. BMC Bioinformatics.
2020;21: 269. doi:10.1186/s12859-020-03608-0

79. Todorov V, Filzmoser P. An Object-Oriented Framework for Robust Multivariate
Analysis. Journal of Statistical Software. 2010;32: 1–47. doi:10.18637/jss.v032.i03

80. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. Genome Biol. 2014;15: 550. doi:10.1186/s13059-
014-0550-8

81. Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, et al. clusterProfiler 4.0: A universal
enrichment tool for interpreting omics data. Innovation (Camb). 2021;2: 100141.
doi:10.1016/j.xinn.2021.100141

82. Blighe K, Rana S, Lewis M. EnhancedVolcano: Publication-ready volcano plots
with enhanced colouring and labeling. 2018. Available:
https://github.com/kevinblighe/EnhancedVolcano

83. Kolde R. pheatmap: Pretty Heatmaps. 2019. Available: https://cran.r-
project.org/web/packages/pheatmap/index.html

84. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics. 2014;30: 2114–2120.
doi:10.1093/bioinformatics/btu170

85. Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-
MEM. arXiv; 2013. doi:10.48550/arXiv.1303.3997

86. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic
features. Bioinformatics. 2010;26: 841–842. doi:10.1093/bioinformatics/btq033

87. Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, et al. Twelve
years of SAMtools and BCFtools. GigaScience. 2021;10: giab008.
doi:10.1093/gigascience/giab008

88. Li H, Durbin R. Fast and accurate short read alignment with Burrows–Wheeler
transform. Bioinformatics. 2009;25: 1754–1760. doi:10.1093/bioinformatics/btp324

89. Zverinova S, Guryev V. Variant calling: Considerations, practices, and
developments. Hum Mutat. 2022;43: 976–985. doi:10.1002/humu.24311

90. Schalamun M, Nagar R, Kainer D, Beavan E, Eccles D, Rathjen JP, et al.
Harnessing the MinION: An example of how to establish long-read sequencing in a
laboratory using challenging plant tissue from Eucalyptus pauciflora. Molecular
Ecology Resources. 2019;19: 77–89. doi:10.1111/1755-0998.12938

91. Wick R. rrwick/Filtlong. 2024. Available: https://github.com/rrwick/Filtlong

92. Fu Y, Mahmoud M, Muraliraman VV, Sedlazeck FJ, Treangen TJ. Vulcan:
Improved long-read mapping and structural variant calling via dual-mode
alignment. GigaScience. 2021;10: giab063. doi:10.1093/gigascience/giab063

93. Li H. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics.
2018;34: 3094–3100. doi:10.1093/bioinformatics/bty191

94. Sedlazeck FJ, Rescheneder P, Smolka M, Fang H, Nattestad M, von Haeseler A,
et al. Accurate detection of complex structural variations using single-molecule
sequencing. Nat Methods. 2018;15: 461–468. doi:10.1038/s41592-018-0001-7

95. Jeffares DC, Jolly C, Hoti M, Speed D, Shaw L, Rallis C, et al. Transient structural
variations have strong effects on quantitative traits and reproductive isolation in
fission yeast. Nat Commun. 2017;8: 14061. doi:10.1038/ncomms14061