Location-Dependent Differences in Cardiac and Skeletal Muscle Dysfunction

Associated With Truncating Titin (*ttn.2*) Variants

Celine F. Santiago[1,2]ORCID Inken G Huttner[1,2]*, Ailbhe O Brien[3]*, Pauline M. Bennett[4], Jasmina Cvetkovska[1], Renee Chand[1], Mark Holt[3]ORCID, Gunjan Trivedi[1], Louis W. Wang[1,2] ORCID, Kelly A. Smith[5]ORCID, Mathias Gautel[3] **ORCID§ , Diane Fatkin[1,2,6] **ORCID§ Yaniv Hinits[3] **ORCID§

[1] Molecular Cardiology and Biophysics Division, Victor Chang Cardiac Research Institute, Darlinghurst, New South Wales, Australia.

[2] School of Clinical Medicine, Faculty of Medicine and Health, UNSW Sydney, Kensington, New South Wales, Australia

[3] Randall Centre for Cell and Molecular Biophysics, and British Heart Foundation Centre of Research Excellence, King's College London, London SE1 1UL, UK.

[4] Randall Centre for Cell and Molecular Biophysics, King's College London, London SE1 1UL, UK.

[5] Department of Anatomy & Physiology, School of Biomedical Sciences, University of Melbourne, Melbourne, Victoria, Australia.

[6] Cardiology Department, St. Vincent's Hospital, Darlinghurst, New South Wales, Australia.

Short title: *TTN*tv-associated cardiac and skeletal myopathy

* Contributed equally

** Co-senior authors

§Corresponding authors:

Mathias Gautel; e-mail: mathias.gautel@kcl.ac.uk

Diane Fatkin e-mail: d.fatkin@victorchang.edu.au

29 Yaniv Hinits; e-mail: yaniv.hinits@kcl.ac.uk

30 Randall Centre for Cell and Molecular Biophysics, New Hunt's House, 18-20 Newcomen

31 Street, King's College London BHF Centre of Research Excellence, London SE1 1UL, UK

32
33 Abstract

34 Background

35 Truncating variants (*TTN*tv) in the *TTN* gene, encoding the giant sarcomeric protein titin,

36 cause a range of human cardiac and skeletal muscle disorders of varying penetrance and

37 severity. The effects of variant location on clinical manifestations are incompletely

38 understood.

40 Methods

41 We generated six zebrafish lines carrying tv in the *ttn.2* gene at the Z-disk, I-band, A-band,

42 and M-band titin regions. Expression of titin transcripts was evaluated using qPCR.

43 Phenotype analysis was performed during embryonic development and in adult hearts.

45 Results

Using location-specific primers, we found a significant reduction of Z-disk and I-band *ttn.2*

transcripts in all homozygous embryos but levels of A-band and M-band transcripts were

reduced only in lines with truncations distal to the *cronos* promoter. Homozygous embryos

uniformly died by 7-10 days post fertilization with marked impairment of cardiac

morphology and function. Skeletal muscle motility and sarcomere organization were more

disrupted in mutants with truncations distal to the *cronos* promoter compared to those

proximal. The most C-terminal line (e232), which lacked only the titin kinase and M-band

regions, differed from other lines, with homozygous embryos showing incorporation of

truncated Ttn.2/Cronos protein and normal sarcomere assembly, but selective degradation of

fast skeletal muscle sarcomeres. Heterozygous embryos were phenotypically

indistinguishable from wild-type. High-frequency echocardiography in adult heterozygous

fish showed reduced ventricular contraction under resting conditions in A-band mutants.

Heterozygous Z-disk and I-band mutants had no significant baseline impairment but were

unable to augment ventricular contraction in response to acute adrenaline exposure,

indicating a lack of cardiac reserve.

Conclusions

Our data suggest that cardiac and skeletal muscle dysfunction associated with truncating *ttn.2*

variants is influenced by age, variant location, and the amount of functional titin protein. The

distinctive phenotype associated with distal C-terminal truncations may reflect different

requirement for C-terminal titin for maintenance of fast, slow and cardiac muscle sarcomeres.

Key words

zebrafish, titin, dilated cardiomyopathy, skeletal myopathy

71
72

Introduction

Titin is a giant protein that spans half-sarcomeres from the Z-disk to the M-band in the heart and skeletal muscle. It is a key determinant of passive muscle stiffness and active contraction in health and disease [1]. In recent years, clinical interest in titin has escalated due to discoveries of genetic variants in patients with cardiac and skeletal myopathies. Truncating variants in the *TTN* gene (*TTN*tv) are now known to be the most common genetic risk factor for dilated cardiomyopathy (DCM) [2, 3]. While *TTN*tv-associated DCM is typically a dominant adult-onset condition, albeit with variable penetrance, individuals who have two abnormal *TTN* alleles leading to truncated proteins (compound heterozygotes or homozygotes) present with childhood-onset skeletal myopathy [4-8]. *TTN*tv also cause cardiac and skeletal myopathy in compound heterozygosity with destabilising titin missense mutations [5, 7]. *TTN*tv have been reported in up to 3% of the general population [2] and frequently arise as incidental findings in genetic testing for unrelated disorders and many parental carriers of *TTN*tv remain asymptomatic beyond their sixties.

Effects due to variant location have been proposed as a useful way to stratify *TTN*tv. Most DCM-associated *TTN*tv arise in the titin A-band region, while skeletal muscle-associated variants were proposed to cluster in the M-band [2-4, 9]. This simplistic view does not account for the presence of *TTN*tv in non-A-band locations in DCM patients, nor A-band *TTN*tv in healthy subjects. Although most patients with A-band *TTN*tv have cardiac presentations, there are increasing reports of overt or subclinical skeletal muscle involvement [10]. Further, children with congenital recessive *TTN*tv-associated skeletal myopathy may progressively develop DCM [4, 8]. There are numerous developmental and tissue-specific titin isoforms that arise due to complex alternate splicing. The extent to which any single exon is represented across the range of titin isoforms is indicated by the percent spliced-in (PSI) score. DCM-associated *TTN*tv are thought to be preferentially distributed in exons that have

high PSI scores in adult heart [3]. Whether PSI scores are useful in the context of skeletal

myopathy has yet to be clarified. Identifying the subset of *TTN*tv that have the greatest risk of

pathogenicity is therefore crucial for informed medical management and genetics counselling

of patients and their families.

Zebrafish are a useful animal model to study *TTN*tv effects. There are two titin genes

in zebrafish, *ttn.2* (*ttna*) and *ttn.1* (*ttnb*). During early development, *ttn.2* is expressed in both

cardiac and skeletal muscle and is the primary titin gene in the heart [11]. Both heart and

skeletal muscle phenotypes have been reported in homozygous *ttn.2* mutants [12]. We have

previously reported adult-onset DCM in heterozygous *ttn.2* zebrafish generated to carry a

human A-band *TTN*tv [13]. *ttn.1* is mainly expressed in skeletal muscle and *ttn.1* mutants have

been reported to have predominant skeletal myopathy phenotypes [14, 15]. An alternative

promoter in *ttn.2* encodes Cronos, a short titin isoform comprising A-band and M-band

domains. First discovered in zebrafish, the Cronos isoform is conserved in mouse and human

[16, 17]. *Cronos* transcripts are abundant in skeletal muscle, but low in developing hearts.

The aim of our study was to investigate the impact of variant location on cardiac and

skeletal muscle structure and function in embryonic and adult zebrafish, using a series of

mutant lines carrying *ttn.2* variants predicted to truncate Z-disk, I-band, A-band, and M-band

regions.

Methods

Detailed methods are provided in the Supplementary Material. Data are available from the

corresponding authors on reasonable request.

Results

***ttn.2* zebrafish models**

We generated a series of six zebrafish *ttn.2* lines carrying variants predicted to result in a

truncated titin protein (Figures 1A, S1). The e5 line has a nonsense mutation that introduces a

premature stop codon in exon 5 before the Z-repeats in the proximal Z-disk region. The e25

line has a 2 bp deletion and premature stop codon in exon 25 in the proximal I-band. In the

e105 line, a 7 bp deletion in exon 105 is predicted to shift the reading frame with 12 residues

of neo-sequence followed by a premature stop codon at the amino acid corresponding to

human *TTNtv*, p.Y12304*. Exon 105 is in the distal I-band proximal to the *cronos* promoter.

The e129 line has a 5 bp deletion and stop codon in exon 129 distal to the *cronos* promoter

near the I-band/A-band junction. The e201 line, modelling the human *TTN*tv p.R26331*, has

previously been described [13] and carries an 8 bp deletion and stop codon in exon 201 in the

mid A-band. The e232 line has a 1 bp deletion and 8 bp insertion in exon 232 in the distal A-

band, upstream of the titin kinase domain and M-band [18]. Zebrafish exons targeted in these

mutants have high sequence homology to the corresponding human *TTN* exons, all of which

have high PSI scores in adult human heart (Table S5).

Wild-type (WT) and mutant *ttn.2* alleles were quantified by qPCR analysis of pooled

whole embryos (3-5 dpf). Using WT allele-specific primers, levels of WT *ttn.2* transcript

were reduced by ~50% in heterozygous (+/-) mutants and absent in homozygous (-/-) mutants

(Figure 1B). In e129-/-, e201-/- and e232 -/- embryos, titin transcripts were globally reduced

as indicated by qPCR using location-specific primers across Z-, I-, A- and M-band regions.

However, in e5-/-, e25-/- and e105-/- embryos, only Z- and I-band transcripts were reduced,

while A- and M-band transcripts were expressed at normal levels (Figure 1C). We

hypothesized that these differences might be explained by differential upregulation of the

titin Cronos isoform [16]. Indeed, quantification of *cronos* transcripts revealed significantly

increased expression in e5-/-, e25-/- and e105-/- embryos (proximal to the *cronos* promoter)

but not in e129-/-, e201-/- and e232-/- embryos (distal to the *cronos* promoter) (Figure 1C).

This may suggest up-regulation of *cronos* due to mechanism triggered by degraded mutant

mRNA [19]. There was modest up-regulation of the WT *ttn.1* transcript in all lines but this was

statistically significant only for e129-/- and e201-/- (Figure 1C).

Embryonic cardiac phenotypes

All in-crosses of heterozygous alleles gave rise to Mendelian-ratio progenies. Homozygous

embryos from e5, e25, e105, e129 and e201 lines were visually identified from 24 hpf by a

dent in the yolk caused by pericardial effusion shortly after the heart tube begins to contract

at 23 hpf [20]. By 3 dpf, there was severe pericardial edema, an indicator of impaired

ventricular contraction, as well as small eyes and craniofacial dysplasia (Figure 2A, B). The

heart chambers were small and linearly arranged, indicating poor development and impaired

looping (Figure 2B). e232-/- embryos were a notable exception, with overall wild-type

appearance and cardiac morphology. All homozygous embryos died between 4 to 10 dpf

(Figure 2C); e232-/- mutants died as larvae but survived relatively longer than other lines.

There were no differences in survival between heterozygotesand WT siblings.

Cardiac function was evaluated in embryonic fish using video-microscopy. For all

lines except e232, homozygous mutants showed a marked reduction in ventricular and atrial

contractile function (assessed by fractional area change) from 3 dpf onwards (Figure 2D, E).

Cardiac function in heterozygous *ttn.2* embryos was indistinguishable from WT siblings

(Figure S3). Heart rates were similar in WT and mutant embryos.

Myocardial architecture in embryonic fish

Embryonic hearts were evaluated using immunofluorescence staining. At 3 dpf, the intensity

of myosin heavy chain (MyHC) staining in e5-/-, e25-/-, e105-/-, e129-/-, and e201-/-

embryonic hearts was reduced, with numerous aggregates and few very disorganized fibrils

(Figures 3A, 3B, S4A,B). Z-disk titin interacts with the cytoskeletal protein, α-actinin [21]. In

e5-/-, e25-/-, e105-/-, e129-/-, and e201-/- embryos, α-actinin staining showed aggregates and

poorly organized striations, suggesting that some Z-I brushes may have formed in the

absence of titin (Figure 3A, 3B, 3D). Myosin binding protein C (MyBP-C), which is in close

contact with the C-zone of A-band titin [18], was absent from the cardiac sarcomeres of these

mutants (Figure 3C). Intriguingly, this suggests that despite the apparent absence of direct

titin-MyBP-C contacts in the assembled (relaxed) thick filaments, A-band titin is

indispensable for the correct formation of the myosin-MyBP-C complex *in vivo.* Myomesin-2

(Myom2) interacts with titin in the M-band in normal hearts (reviewed in [22]). Myom2 was

disorganized and present only at very low levels, suggestive of impaired M-band myosin-titin

arrays (Figure 3D).

Cardiac expression of Ttn.2, Ttn.1, and Cronos was evaluated using *in situ*

hybridization and immunostaining. In WT embryos, there was strong staining using the T12

antibody recognizing an epitope in I2-I3 at the start of the I-band in both Ttn.2 and Ttn.1 [23]

(Figure 3C), and a novel antibody that recognizes the Ttn.2 M6 M-band domain (Figures 3A,

S5A,B). Another antibody against the Ttn.1 M8-M9 region, but detecting both Ttn.1 and

Ttn.2 (Figures S2, S5A,C), showed normal cardiac staining but reduced M-band staining in

ttn.1 e7-/- embryos somitic muscle sarcomeres that express both Ttn.1 and Ttn.2 (Figures

S2B, S6M-X). In WT embryos, *ttn.1* transcripts express only weakly in the heart at 1 dpf

(Figure S6B, S6D, S6F and [11]) with negligible expression by 3 dpf (Figure S6H, S6J, S6L).

Like previous studies [14], we found that embryonic heart development in *ttn.1* mutant lines is

normal (Figure S2C, C'). These finding suggest Ttn.2 but not Ttn.1 is essential for normal

sarcomere assembly and formation of a functional heart.

For all homozygous embryos except e232-/-, there was no detectable Ttn.2 protein or Ttn.1 in

the heart above background levels (Figure 3A, 3C and S5). Our qPCR data in whole embryos

(Figure 1) suggested that *cronos* upregulation might be a potential compensatory mechanism

for titin deficiency. However, these changes appear to be driven primarily by *cronos*

expression in skeletal muscle, since in situ hybridization failed to detect *cronos* in the heart

(Figure S6Y, S6Z and [16]).

In keeping with preserved cardiac function, sarcomere organization was

indistinguishable in T12 stained e232-/- embryos compared to WT siblings (Figure 3B).

Given the negligible expression of Ttn.1 and Cronos in 3 dpf hearts, we speculate that e232

transcripts may be incompletely degraded and that truncated e232 Ttn.2 protein is

incorporated into cardiac sarcomeres and able to facilitate near-normal organization of

myosin and sarcomeric proteins such as Myom2 (Figure 3D).

Skeletal muscle embryonic phenotypes

Skeletal muscle motility and structure was evaluated in *ttn.2* embryos. Embryos with

truncating variants proximal to the *cronos* promoter (e5-/-, e25-/-, e105-/-) had reduced

motility compared with sibling control embryos at ~24hpf, which was limited to just a tremor

when stimulated by touch by 3 dpf (Figure S7A). Immunofluorescence staining for myosin

and actin in these 3 dpf mutants showed structured sarcomeres, but some myofibrils were

mis-aligned and break points were evident in the middle of both slow and fast muscle fibers

(Figures 4A, S7B-D). MyBP-C and α-actinin appeared normal (Figure S7E, S7F).

Immunostaining with T12, recognizing an epitope 3' of the *ttn.2* truncation site, showed

strong staining in both slow and fast muscle fibers, which most likely results from Ttn.1

(Figures S7E and S6N, S6R, S6T, S6X). Ttn.2 M6 was also detected strongly in e5-/-, e25-/-

and e105-/- embryos (Figure S7F-H) and is likely to result from Cronos protein. Other M-

band proteins localized correctly to the M-band; Myom2 in fast muscle (Figure 4B) and

Myom1 in both fast and slow muscle fibers (Figure S7I-K). Overall, these data suggest that

Ttn.1 and Cronos have a ruler function and allow many proteins to find their position in the

sarcomere in the absence of full-length Ttn.2. However, reduced motility and fiber breakage

indicate that sarcomere structure is weaker and prone to damage once contractions

commence.

Severe loss of skeletal muscle sarcomeric structure in embryos with *ttn.2* truncations

distal to the *cronos* promoter

The severity of skeletal muscle phenotypes was greater in embryos with truncating variants

distal to the *cronos* promoter (e129-/-, 201-/-) compared to those with proximal truncations

(e5-/-, e25-/-, e105-/-). The former embryos were barely motile at 24 hpf, showing only weak

twitches. By 3 dpf, embryos were paralyzed with no response to mechano-stimulation (Figure

S8A). Already at 24 hpf, sarcomeres in e129-/- embryos had immature thin myofibrils

resembling early differentiated myofibrils [24] with only small areas of low-level sarcomeric

organization (Figures 4C, S8D). By 2 and 3 dpf, immunofluorescence staining for myosin

and actin showed a loss of axial fibrillar organization in both slow and fast fibers (Figures

4A, 4C, and S8B, S8C). Rather, short lengths of ~1 μm structures were revealed by staining

for α-actinin (Figure 4E) and antibodies that detect Ttn.1, such as Z1Z2, T12, and Ttn.1 M8-

M9 (Figures 4F, 4G, S8B). The directionality of these structures was random, differing

greatly from normal organized sarcomeres (Figure 4L, 4M) and resembling I-Z-I complexes.

I-Z-I complexes are structures formed during myofibrillogenesis that contain Z-disk

components as well as α-actin and titin. These can be put in register on actin filamentous

structures and can form in the absence of MyHC [25]. Ttn.2 (M6 antibody) was not detected in

these mutants (Figures 4E, S8E) while other M-band proteins such as Myom2 were only

weakly detected (Figure 4D). This suggests that fibrils expressing only Ttn.1 have lost their

rigidity or their ability to associate side by side, so that they grow in a wavering, random way

when viewed longitudinally. Electron microscopy supports this, with a few repeats of

sarcomere-like structures running in different directions (Figure 4H-4K). Between the Z-

disks and their associated thin filaments, groups of thick filaments were seen. However, they

did not appear to be organized as A-bands and no M-band structures were seen. Furthermore,

the separation of the thick filaments in cross section was greater than seen in WT siblings.

The sarcoplasmic reticulum was present, and some triad structures remain associated with Z-

disks (Figure 4J). Overall, the results show that in skeletal muscles of homozygous embryos

with *ttn.2* truncations distal to the *cronos* promoter, structures containing I-Z-I brushes and

Ttn.1 are formed, but Ttn.1 alone cannot support proper organization of A- and M-bands.

Sarcomere structure disintegrates in fast fibers of embryos with M-band ttn.2

truncation

Similar to the cardiac findings, the skeletal muscle phenotype in e232-/- embryos was distinct

to the other mutants. The overall appearance of e232-/- embryos was indistinguishable from

WT siblings at 3 dpf (Figure 2A). However, whereas coiling movement within the chorion

was normal at 24 hpf, e232-/- embryos had an abnormal swimming pattern at 48 hpf, which

involved slower tail movement and less motion of the trunk (Supplementary Video). By 5

dpf, e232-/- embryos lacked burst-swimming upon touch, unlike WT siblings. However,

spasms of the trunk, pectoral fin movement and increased eye movement in response to touch

were detected, indicating intact sensory reflexes (Figure S9A-S9C). Burst-swimming

behaviour is mediated primarily by fast fibers, whilst the early coiling movement is mediated

by early differentiating slow muscle [26, 27]. By 5 dpf, e232-/- mutants developed curvature of

the trunk and failed to inflate their swim bladders (Figure 5A, Supplementary Video) [27].

At 24 hpf shortly after differentiation, sarcomeric structure in both slow and fast

fibers in e232-/- embryos appeared normal (Figure 5B-5D). As in cardiac sarcomeres, T12

signal was present (Figure 5G) but Ttn.2 M6 was absent (Figure 5C), suggesting that

truncated Ttn.2 and Cronos were incorporated into the sarcomere. By 3 dpf, however, while

slow muscle fibers maintained their regular sarcomeric structure, fast muscle sarcomeres

were severely disrupted as indicated by myosin, MyBP-C, α-actinin and T12 (detecting Ttn.2

upstream from the truncation domain) staining patterns (Figures 4A, 5E-5G). The M-band

protein Myom1 is normally present in both slow and fast fibers but was only detected in

e232-/- embryo slow fibers (Figure 5E). Myom2, normally present only in fast fibers, was

severely diminished in e232-/- embryos (Figure 5F). Electron microscopy revealed normal

sarcomeric structure in e232-/- slow muscle fibers, whereas no sarcomeres were visible in

fast muscle fibers and thick filaments appeared to be highly disordered (Figure 5H, 5I). A

small number of Z-discs and triads were also visible, but not associated.

Adult cardiac phenotypes

Heterozygous *ttn.2* mutant fish survived to adulthood with overall normal appearance and

swimming behaviour. *ttn.2* transcript expression in mutant lines was reduced by ~50%

relative to WT (Figure 1B). Confirming previous studies [14], neither reduced levels of titin

protein nor truncated titin proteins were detectable using Western blots with an N-terminal

specific titin antibody, or Coomassie stained polyacrylamide gels (Figure S10). Due to the

lack of embryonic cardiac phenotype in e232-/- fish, e232+/- fish were not included in the

adult analyses.

Cardiac function in 12-month-old fish was evaluated using high frequency

echocardiography (Figure 6, Table S6). Ventricular size (EDV/BSA) in the heterozygous

ttn.2 fish was unchanged compared to WT, except for e201+/- that showed modest dilatation

(Figure 6A), in line with our previous findings [13]. Significant impairment of ventricular

contractile function was seen only for e129+/- and e201+/- lines (Figure 6B through E). In

addition to systolic impairment, these lines also had prolonged isovolumic relaxation time,

indicating diastolic dysfunction (Figure 6F). Ventricular function in the A-band mutant,

e201+/- was similar in *ttn^xu071* double mutant (*ttn.2* A-band + *ttn.1* Z-disk truncations, [14]),

implicating predominant *ttn.2* effects (Figure S11).

We hypothesized that fish with *ttn.2* truncations outside the A-band might be in a

compensated state at baseline, and that exposure to additional stress might unmask latent

cardiac dysfunction. We therefore subjected 10-15 month-old e5+/- and e105+/- fish to acute

adrenaline stress, as previously described [28]. WT fish demonstrated the expected adrenaline-

induced increases in heart rate and contractility (Figure 6D-F). In contrast, these responses

were not seen in e5+/- and e105+/- fish (Figure 6D-F), indicating a lack of cardiac reserve.

Myocardial structure was further interrogated in e5+/- and e201+/- fish, hypothesizing

that these two lines are representative of compensated *vs.* penetrant disease states,

respectively. No overt differences in wall thickness or trabeculation were observed in

haematoxylin and eosin-stained tissue sections in either of these mutant lines relative to WT

siblings (Figure S12A, S12D). Sarcomere structure, content and length were also similar to

WT (Figure S12B-E).

Discussion

Zebrafish *TTN*tv models exhibit a varying extent of cardiac and skeletal muscle dysfunction

that differs with age, variant location, and the expected amount of normally functioning titin

protein. Previous zebrafish studies have focussed mainly on the developmental impact of titin

deficiency using homozygous embryos with selected Z-disk and/or A-band *ttn.2* truncations

[12]. We now extend these findings by directly comparing *ttn.2* truncations distributed

throughout the titin protein, including the first zebrafish model that truncates the M-band. We

also provide new data about cardiac phenotypes in a range of adult heterozygous fish. Our

data expand understanding of the roles of titin in heart and skeletal muscle and have potential

implications for human titinopathies.

Homozygous *ttn.2* embryos showed marked cardiac dysfunction and early death,

similar to reported *ttn.2* mutants [12]. These findings are most readily explained by a lack of

functional Ttn.2 protein in the heart and the lack of effective compensation by Ttn.1 or

Cronos. Homozygous *ttn.2* embryos also showed abnormalities of skeletal muscle structure

and function, the severity of which appeared to be determined by location with respect to the

cronos promoter. For proximal truncations, the presence of intact Cronos and Ttn.1 allows

formation of fragile sarcomeres that fail following onset of contraction. However, for distal

truncations, both Cronos and Ttn.2 are affected and although intact, Ttn.1 appears to be

insufficient to support viable sarcomere assembly.

The e232-/- embryos were a notable exception with respect to both the cardiac and

skeletal muscle phenotypes. A near-complete A-band in e232-/-, capable of interacting with

myosin and organizing MyBP-C, seems essential for incorporation and retention of truncated

Ttn.2 in sarcomeres. Remarkably, although loss of full-length Ttn.2 protein prevents

sarcomere formation, C-terminal titin appears to be dispensable for cardiomyocyte sarcomere

assembly, contraction, and normal heart morphology during early development. In contrast to

other lines, minimally truncated Ttn.2 protein is stable enough to allow integration into the

sarcomere. Factors such as the truncation position at the distal end of the giant titin protein

and the large exon size may result in reduced efficiency of nonsense mediated decay of e232

alleles [29]. While this might be sufficient to support sarcomere development, the presence of

truncated titin seems to affect protein interactions in this region. Homozygous C-terminal

titin truncation in mouse develop a late cardiac phenotype and differ greatly from other

truncations [30, 31]. Of note, sarcomeric integration of C-terminal truncated titin has been

reported in human muscle biopsies from patients with M-band *TTN*tv associated with

autosomal recessive skeletal ± cardiac myopathy [4-7].

In e232-/- embryos, sarcomere organization was preserved in cardiac and slow

muscle, but not in fast muscle fibers. Since *ttn.2* and *ttn.1* are expressed in both slow and fast

fibers, these observations suggest that there are fundamental differences in the requirement

for C-terminal titin for the assembly and maintenance in these muscle types. One difference

between slow and fast skeletal muscle fibers in the M-band region is expression of

myomesin. Whereas Myom1 is present in cardiac, slow and fast muscle in zebrafish, Myom2

is only expressed in fast skeletal muscle, similar to mammalian muscle (reviewed in [22]) . In

contrast, Myom2 is present in the murine heart from e14.5 [22], while in fish hearts, expression

of *myom2a*, one of the fish *MYOM2* homologues, is detected from the early embryonic heart

tube stage (Fig. S5AA). Collectively, our data suggest that Myom2 is not essential for

sarcomere assembly but may have a role in maintenance of sarcomere organization in fast

skeletal muscle. Other proteins that interact with M-band titin, such as obscurin, calpain-3

and myospryn, may be important for differentiating normally functioning sarcomeres in fast

versus slow and cardiac muscle fibers. Human patients with C-terminal truncations localize

Myom1 and obscurin/Obsl1 to the M-band, but with some disruptions [7, 32]. Similarly, e232-/-

mutants recruited Myom1 to slow and cardiac M-band sarcomeres, possibly via dimerization

of Myom1 and binding of Myom1 to obscurin/Obsl1 and myosin. However, absent M-

band/titin connections likely gradually result in structural instability and may underpin the

disruption seen in human patient samples [32]. The selective loss of fast skeletal muscle fibers

may reflect susceptibility to damage in response to mechanical and metabolic stress [33].

Interestingly, dystrophin-deficient mdx mice have disproportionate exercise-induced damage

in fast skeletal myofibers, and human patients with Salih myopathy and multi minicore

disease associated with C-terminal *TTN*tv show a predominance of slow (type I) fibers from a

very early age [4, 5].

The location of *TTN*tv has emerged as an important issue in human titinopathies.

Congenital skeletal titinopathy is recessive and typically associated with biallelic *TTN*tv,

except for a rare dominant *TTN*tv in exon 327 [34]. Cohort studies have demonstrated that these

disease-causing *TTN*tv can arise throughout the entire *TTN* gene [8, 35]. Cardiac involvement is

seen in approximately 50% patients and is more likely when both *TTN*tv are in exons that are

constitutively expressed in adult heart [8]. Notably, the age of disease onset is much later in

patients in whom at least one of the *TTN*tv is present in the last three M-band exons [35]. Two

autosomal dominant forms of skeletal myopathy, tibial muscular dystrophy and hereditary

myopathy with early respiratory failure, have been associated with single *TTN*tv in the final

six M-band exons [8]. These disorders typically occur in adult life and cardiac involvement is

variable. Taken together, these findings suggest that *TTN* location and titin "dose" are

important determinants of clinical manifestations and disease severity.

The C-terminal part of titin shows unexpected complexity in its integration into the myosin

filament, as revealed by recent cryo-electron tomographic (cryoET) structures. Three double

strands of titin (dubbed titin alpha and beta) with six molecules in total run along the thick

filament [18]. Both titin alpha and beta interact outside the M-band at the first levels of myosin

motor domains. However, whereas titin alpha continues into the M-band, the C-terminal

regions of titin beta from about the second Ig-domains after the kinase domain could not be

traced in the cryoET structure [18], suggesting that titin beta is either truncated or follows a

disordered path. The non-equivalence of 3 of the 6 titin filaments of each half thick filament

may account for the different impact of truncations N-and C-terminal to the kinase domain.

Structural studies of the corresponding zebrafish mutants may therefore provide valuable

insight into the different pathomechanism of these mutations.

*TTN*tv are the most common genetic cause of autosomal dominant adult-onset DCM,

with the majority of these being in the titin A-band or I/A junction [2, 3]. Interpretation of the

clinical significance of *TTN*tv outside this region has remained a clinical conundrum. Our

evaluation of adult zebrafish hearts confirms the importance of A-band *TTN*tv, with

significant changes in ventricular contraction seen under resting conditions in e129-/- and

e201+/- fish. However, e5+/- and e105+/- fish cannot be considered to have normal hearts as

their failure to augment ventricular contraction in response to an adrenaline challenge is

indicative of impaired contractile reserve. A recent study found that zebrafish from another

ttn.2 e5+/- mutant had no overt DCM but some mild abnormalities [36]. All our zebrafish lines

had *ttn.2* truncating variants in zebrafish exons that show high sequence homology to

constitutively expressed human *TTN* exons. Collectively, our data suggest that *TTN*tv in high

PSI cardiac exons confer a genetic susceptibility to DCM and that overt clinical manifestation

may depend on patient-related factors such as age, sex, and clinical risk factors, particularly

those that increase cardiac mechanical stress. Further human *TTN*tv cohort studies are needed

to investigate these points.

Whereas our data, and those from human C-terminal *TTN*tvs [4-7], show clear evidence for

integration of truncated titin protein into the sarcomere, we have no evidence to support such

phenomena for more N-terminal truncations. Recent studies have reported the persistence of

some truncated titin protein from DCM *TTN*tv cardiomyocytes, but with conflicting evidence

regarding its localization in the sarcomere or in intracellular aggregates [37-40]. Our study is not

without limitations: there remains a lack of definitive evidence for reduced titin protein

expression, and for established data for titin exon usage in fetal heart and in fetal and adult

skeletal muscle. Due to potential species differences, our findings may not be directly

translatable to humans. Nevertheless, our findings provide new insights into the variability

underpinning *TTN*tv effects and their potential fundamental impact on thick-filament

assembly and are hypothesis-generating for human clinical studies.

Acknowledgments

We thank Dan Hesselson and Xiaolei Xu for assistance with zebrafish lines; Alexander

Alexandrovich and Atsushi Fukuzawa for assistance with antigen expression and

purification; Wolfgang Linke for expert assistance with titin protein analysis; the Centre for

Ultrastructural Imaging, King's College London, for electron microscopy services; and the

Victor Chang Cardiac Research Institute aquarium staff, in particular Aaron Hay and Cecelia

Jenkin, for assistance with zebrafish husbandry. The authors acknowledge the Victor Chang

Cardiac Research Institute Innovation Centre, funded by the NSW Government, including the

Preclinical Imaging Facility.

Sources of Funding

AOB was supported by the PhD programme of the King's College London British Heart

Foundation (BHF) Centre of Research Excellence. MG holds the BHF Chair of Molecular

Cardiology. MG and YH are supported by the European Research Council Synergy Grant

856118 StuDySARCOMERE. CFS was supported by the Simon Lee Foundation and the

Australian Department of Education & Training Research Training Program Scholarship. DF

is supported by NSW Health, and the Medical Research Futures Fund.

Disclosures

None.

References

1.   Loescher CM, Hobbach AJ, Linke WA. Titin (ttn): From molecule to modifications, mechanics,
and medical significance. *Cardiovasc Res*. 2022;118:2903-2918

2.   Herman DS, Lam L, Taylor MR, Wang L, Teekakirikul P, Christodoulou D, Conner L,
DePalma SR, McDonough B, Sparks E, Teodorescu DL, Cirino AL, Banner NR, Pennell DJ,
Graw S, Merlo M, Di Lenarda A, Sinagra G, Bos JM, Ackerman MJ, Mitchell RN, Murry CE,
Lakdawala NK, Ho CY, Barton PJ, Cook SA, Mestroni L, Seidman JG, Seidman CE.
Truncations of titin causing dilated cardiomyopathy. *N Engl J Med*. 2012;366:619-628

3.   Roberts AM, Ware JS, Herman DS, Schafer S, Baksi J, Bick AG, Buchan RJ, Walsh R, John
S, Wilkinson S, Mazzarotto F, Felkin LE, Gong S, MacArthur JA, Cunningham F, Flannick J,
Gabriel SB, Altshuler DM, Macdonald PS, Heinig M, Keogh AM, Hayward CS, Banner NR,
Pennell DJ, O'Regan DP, San TR, de Marvao A, Dawes TJ, Gulati A, Birks EJ, Yacoub MH,
Radke M, Gotthardt M, Wilson JG, O'Donnell CJ, Prasad SK, Barton PJ, Fatkin D, Hubner N,
Seidman JG, Seidman CE, Cook SA. Integrated allelic, transcriptional, and phenomic
dissection of the cardiac effects of titin truncations in health and disease. *Sci Transl Med*.
2015;7:270ra276

4.   Carmignac V, Salih MAM, Quijano-Roy S, Marchand S, Al Rayess MM, Mukhtar MM,
Urtizberea JA, Labeit S, Guicheney P, Leturcq F, Gautel M, Fardeau M, Campbell KP, Richard
I, Estournet B, Ferreiro A. C-terminal titin deletions cause a novel early-onset myopathy with
fatal cardiomyopathy. *Ann. Neurol.* 2007;61:340-351

5.   Chauveau C, Bonnemann CG, Julien C, Kho AL, Marks H, Talim B, Maury P, Arne-Bes MC,
Uro-Coste E, Alexandrovich A, Vihola A, Schafer S, Kaufmann B, Medne L, Hubner N, Foley
AR, Santi M, Udd B, Topaloglu H, Moore SA, Gotthardt M, Samuels ME, Gautel M, Ferreiro
A. Recessive ttn truncating mutations define novel forms of core myopathy with heart disease.
Hum Mol Genet. 2014;23:980-991

6.   Ceyhan-Birsoy O, Agrawal PB, Hidalgo C, Schmitz-Abe K, DeChene ET, Swanson LC,
Soemedi R, Vasli N, Iannaccone ST, Shieh PB, Shur N, Dennison JM, Lawlor MW, Laporte J,
Markianos K, Fairbrother WG, Granzier H, Beggs AH. Recessive truncating titin gene, ttn,
mutations presenting as centronuclear myopathy. *Neurology*. 2013;81:1205-1214

7.   Rees M, Nikoopour R, Fukuzawa A, Kho AL, Fernandez-Garcia MA, Wraige E, Bodi I,
Deshpande C, Ozdemir O, Daimaguler HS, Pfuhl M, Holt M, Brandmeier B, Grover S, Fluss
J, Longman C, Farrugia ME, Matthews E, Hanna M, Muntoni F, Sarkozy A, Phadke R,
Quinlivan R, Oates EC, Schroder R, Thiel C, Reimann J, Voermans N, Erasmus C, Kamsteeg
EJ, Konersman C, Grosmann C, McKee S, Tirupathi S, Moore SA, Wilichowski E,
Hobbiebrunken E, Dekomien G, Richard I, Van den Bergh P, Dominguez-Gonzalez C, Cirak
S, Ferreiro A, Jungbluth H, Gautel M. Making sense of missense variants in ttn-related
congenital myopathies. *Acta Neuropathol*. 2021;141:431-453

8.   Oates EC, Jones KJ, Donkervoort S, Charlton A, Brammah S, Smith JE, 3rd, Ware JS, Yau KS,
Swanson LC, Whiffin N, Peduto AJ, Bournazos A, Waddell LB, Farrar MA, Sampaio HA,
Teoh HL, Lamont PJ, Mowat D, Fitzsimons RB, Corbett AJ, Ryan MM, O'Grady GL,
Sandaradura SA, Ghaoui R, Joshi H, Marshall JL, Nolan MA, Kaur S, Punetha J, Topf A, Harris
E, Bakshi M, Genetti CA, Marttila M, Werlauff U, Streichenberger N, Pestronk A, Mazanti I,
Pinner JR, Vuillerot C, Grosmann C, Camacho A, Mohassel P, Leach ME, Foley AR,
Bharucha-Goebel D, Collins J, Connolly AM, Gilbreath HR, Iannaccone ST, Castro D,
Cummings BB, Webster RI, Lazaro L, Vissing J, Coppens S, Deconinck N, Luk HM, Thomas
NH, Foulds NC, Illingworth MA, Ellard S, McLean CA, Phadke R, Ravenscroft G, Witting N,
Hackman P, Richard I, Cooper ST, Kamsteeg EJ, Hoffman EP, Bushby K, Straub V, Udd B,
Ferreiro A, North KN, Clarke NF, Lek M, Beggs AH, Bonnemann CG, MacArthur DG,
Granzier H, Davis MR, Laing NG. Congenital titinopathy: Comprehensive characterization and
pathogenic insights. *Ann Neurol*. 2018;83:1105-1124

9. Savarese M, Sarparanta J, Vihola A, Udd B, Hackman P. Increasing role of titin mutations in
neuromuscular disorders. *J Neuromuscul Dis*. 2016;3:293-308

10. Skriver SV, Krett B, Poulsen NS, Krag T, Walas HR, Christensen AH, Bundgaard H, Vissing
J, Vissing CR. Skeletal muscle involvement in patients with truncations of titin and familial
dilated cardiomyopathy. *JACC Heart Fail*. 2024;12:740-753

11. Seeley M, Huang W, Chen Z, Wolff WO, Lin X, Xu X. Depletion of zebrafish titin reduces
cardiac contractility by disrupting the assembly of z-discs and a-bands. *Circ Res*.
2007;100:238-245

12. Santiago CF, Huttner IG, Fatkin D. Mechanisms of ttntv-related dilated cardiomyopathy:
Insights from zebrafish models. *J Cardiovasc Dev Dis*. 2021;8

13. Huttner IG, Wang LW, Santiago CF, Horvat C, Johnson R, Cheng D, Frieling-Salewsky Mv,
Hillcoat K, Bemand TJ, Trivedi G, Braet F, Hesselson D, Alford K, Hayward CS, Seidman JG,
Seidman CE, Feneley MP, Linke WA, Fatkin D. A-band titin truncation in zebrafish causes
dilated cardiomyopathy and hemodynamic stress intolerance. *Circulation: Genomic and
Precision Medicine*. 2018;11:e002135

14. Shih YH, Dvornikov AV, Zhu P, Ma X, Kim M, Ding Y, Xu X. Exon- and contraction-
dependent functions of titin in sarcomere assembly. *Development*. 2016;143:4713-4722

15. Steffen LS, Guyon JR, Vogel ED, Howell MH, Zhou Y, Weber GJ, Zon LI, Kunkel LM. The
zebrafish runzel muscular dystrophy is linked to the titin gene. *Developmental Biology*.
2007;309:180-192

16. Zou J, Tran D, Baalbaki M, Tang LF, Poon A, Pelonero A, Titus EW, Yuan C, Shi C, Patchava
S, Halper E, Garg J, Movsesyan I, Yin C, Wu R, Wilsbacher LD, Liu J, Hager RL, Coughlin
SR, Jinek M, Pullinger CR, Kane JP, Hart DO, Kwok P-Y, Deo RC. An internal promoter
underlies the difference in disease severity between n- and c-terminal truncation mutations of
titin in zebrafish. *eLife*. 2015;4:e09406

17. Zaunbrecher RJ, Abel AN, Beussman K, Leonard A, von Frieling-Salewsky M, Fields PA,
Pabon L, Reinecke H, Yang X, Macadangdang J, Kim DH, Linke WA, Sniadecki NJ, Regnier
M, Murry CE. Cronos titin is expressed in human cardiomyocytes and necessary for normal
sarcomere function. *Circulation*. 2019;140:1647-1660

18. Tamborrini D, Wang Z, Wagner T, Tacke S, Stabrin M, Grange M, Kho AL, Rees M, Bennett
P, Gautel M, Raunser S. Structure of the native myosin filament in the relaxed cardiac
sarcomere. *Nature*. 2023

19. El-Brolosy MA, Kontarakis Z, Rossi A, Kuenne C, Gunther S, Fukuda N, Kikhi K, Boezio
GLM, Takacs CM, Lai SL, Fukuda R, Gerri C, Giraldez AJ, Stainier DYR. Genetic
compensation triggered by mutant mrna degradation. *Nature*. 2019;568:193-197

20. Bakkers J. Zebrafish as a model to study cardiac development and human cardiac disease.
Cardiovasc Res. 2011;91:279-288

21. Gautel M, Djinovic-Carugo K. The sarcomeric cytoskeleton: From molecules to motion. *J Exp
Biol*. 2016;219:135-145

22. Lange S, Pinotsis N, Agarkova I, Ehler E. The m-band: The underestimated part of the
sarcomere. *Biochim Biophys Acta Mol Cell Res*. 2020;1867:118440

23. Sebestyen MG, Wolff JA, Greaser ML. Characterization of a 5.4 kb cdna fragment from the z-
line region of rabbit cardiac titin reveals phosphorylation sites for proline-directed kinases. *J
Cell Sci*. 1995;108 (Pt 9):3029-3037

24. Hinits Y, Hughes SM. Mef2s are required for thick filament formation in nascent muscle fibres.
Development. 2007;134:2511-2519

25.   Schultheiss T, Lin ZX, Lu MH, Murray J, Fischman DA, Weber K, Masaki T, Imamura M,
541   Holtzer H. Differential distribution of subsets of myofibrillar proteins in cardiac nonstriated
542   and striated myofibrils. *J Cell Biol*. 1990;110:1159-1172

543   26.   Saint-Amant L, Drapeau P. Time course of the development of motor behaviors in the zebrafish
544   embryo. *J Neurobiol*. 1998;37:622-632

545   27.   Naganawa Y, Hirata H. Developmental transition of touch response from slow muscle-
546   mediated coilings to fast muscle-mediated burst swimming in zebrafish. *Dev Biol*.
547   2011;355:194-204

548   28.   Wang LW, Huttner IG, Santiago CF, Fatkin D. Bradycardia in zebrafish heart failure: A true
549   physiological response or anesthetic-induced red herring? *Zebrafish*. 2016;13:475-476

550   29.   Lindeboom RGH, Vermeulen M, Lehner B, Supek F. The impact of nonsense-mediated mrna
551   decay on genetic disease, gene editing and cancer immunotherapy. *Nat Genet*. 2019;51:1645-
552   1651

553   30.   Radke MH, Polack C, Methawasin M, Fink C, Granzier HL, Gotthardt M. Deleting full length
554   titin versus the titin m-band region leads to differential mechanosignaling and cardiac
555   phenotypes. *Circulation*. 2019;139:1813-1827

556   31.   Weinert S, Bergmann N, Luo X, Erdmann B, Gotthardt M. M line-deficient titin causes cardiac
557   lethality through impaired maturation of the sarcomere. *J Cell Biol*. 2006;173:559-570

558   32.   Fukuzawa A, Lange S, Holt M, Vihola A, Carmignac V, Ferreiro A, Udd B, Gautel M.
559   Interactions with titin and myomesin target obscurin and obscurin-like 1 to the m-band –
560   implications for hereditary myopathies. *Journal of Cell Science*. 2008;121:1841-1851

561   33.   Lloyd EM, Pinniger GJ, Murphy RM, Grounds MD. Slow or fast: Implications of myofibre
562   type and associated differences for manifestation of neuromuscular disorders. *Acta Physiol
563   (Oxf)*. 2023;238:e14012

564   34.   Claeys KG, Savarese M, Jonson PH, Goosens V, Topf A, Vihola A, Straub V, Udd B. A titin
565   truncating variant causing a dominant myopathy with cardiac involvement in a large family:
566   The exception that proves the rule. *Neurol Genet*. 2024;10:e200185

567   35.   Savarese M, Vihola A, Oates EC, Barresi R, Fiorillo C, Tasca G, Jokela M, Sarkozy A, Luo S,
568   Diaz-Manera J, Ehrstedt C, Rojas-Garcia R, Saenz A, Muelas N, Lonardo F, Fodstad H,
569   Qureshi T, Johari M, Valipakka S, Luque H, Petiot P, de Munain AL, Pane M, Mercuri E,
570   Torella A, Nigro V, Astrea G, Santorelli FM, Bruno C, Kuntzer T, Illa I, Vilchez JJ, Julien C,
571   Ferreiro A, Malandrini A, Zhao CB, Casar-Borota O, Davis M, Muntoni F, Hackman P, Udd
572   B. Genotype-phenotype correlations in recessive titinopathies. *Genet Med*. 2020;22:2029-2040

573   36.   Zhu P, Li J, Yan F, Islam S, Lin X, Xu X. Allelic heterogeneity of ttntv dilated cardiomyopathy
574   can be modeled in adult zebrafish. *JCI Insight*. 2024;9

575   37.   Fomin A, Gartner A, Cyganek L, Tiburcy M, Tuleta I, Wellers L, Folsche L, Hobbach AJ, von
576   Frieling-Salewsky M, Unger A, Hucke A, Koser F, Kassner A, Sielemann K, Streckfuss-
577   Bomeke K, Hasenfuss G, Goedel A, Laugwitz KL, Moretti A, Gummert JF, Dos Remedios CG,
578   Reinecke H, Knoll R, van Heesch S, Hubner N, Zimmermann WH, Milting H, Linke WA.
579   Truncated titin proteins and titin haploinsufficiency are targets for functional recovery in
580   human cardiomyopathy due to ttn mutations. *Sci Transl Med*. 2021;13:eabd3079

581   38.   McAfee Q, Caporizzo MA, Uchida K, Bedi KC, Jr., Margulies KB, Arany Z, Prosser BL.
582   Truncated titin protein in dilated cardiomyopathy incorporates into the sarcomere and transmits
583   force. *J Clin Invest*. 2024;134

584   39.   McAfee Q, Chen CY, Yang Y, Caporizzo MA, Morley M, Babu A, Jeong S, Brandimarto J,
585   Bedi KC, Jr., Flam E, Cesare J, Cappola TP, Margulies K, Prosser B, Arany Z. Truncated titin
586   proteins in dilated cardiomyopathy. *Sci Transl Med*. 2021;13:eabd7287

587   40.   Kellermayer D, Tordai H, Kiss B, Torok G, Peter DM, Sayour AA, Polos M, Hartyanszky I,
588   Szilveszter B, Labeit S, Gango A, Bedics G, Bodor C, Radovits T, Merkely B, Kellermayer
589   MS. Truncated titin is structurally integrated into the human dilated cardiomyopathic
590   sarcomere. *J Clin Invest*. 2024;134

592
593

Figure legends

Figure 1. Zebrafish *ttn.2* truncating variants. **A,** Schematic of the Ttn.2 protein showing

major domains and the *cronos* translation start site (TSS). Locations of *ttn.2* truncations (red

arrows) and the *ttn*[x071] mutant [14] (asterisk) are indicated. **B,** qPCR analysis of homozygous (-

/-), heterozygous (+/-) and wild-type (WT; +/+) whole embryos (left panel) and adult hearts

(right panel) using primers specific for the wild-type allele. **C,** qPCR analysis of homozygous

ttn.2 whole embryos and WT controls using primer pairs specific for various *ttn.2* and *ttn.1*

regions. qPCR data (**B,C**) were normalized to house-keeping genes and expressed as absolute

levels of abundance. Two-way ANOVA, mean ± SD.

Figure 2. Cardiac morphology and function in homozygous *ttn.2* embryos. A, Brightfield

images of 3 dpf homozygous and sibling control embryos shown in lateral view, anterior to

left, with (**B**) higher magnification of the heart. With the exception of e232-/-, mutant

embryos show pericardial edema (magenta arrowhead), as well as small eyes and head (cyan

arrowhead). **C,** Survival in homozygous and wild-type (WT) embryos between 0 and 20 dpf;

Mantel-Cox test. Ventricular (**D**) and atrial (**E**) contractile function were assessed using

video-microscopy. Data are expressed as fractional area change (FAC; %). Unpaired t-tests,

mean ± SD. A denotes atrium, OED, edema, OFT, outflow tract, and V, ventricle.

Figure 3. Cardiac sarcomere assembly in *ttn.2* embryos. A and **B,** Single confocal scans

through ventricular sarcomeres of 3 dpf homozygous and sibling (sib) control embryos,

immunostained for myosin heavy chain (MyHC), α-actinin, and Ttn.2 M6. Mutants show

punctate aggregates (arrowheads) except e232-/- embryos in which sarcomeres appear

normal.   No titin protein is detected by Ttn.2 M6 antibodies in e129-/- hearts. **C,** Confocal

stacks of 3 dpf hearts of homozygous *ttn.2* and sibling control embryos, immunostained for

Ttn T12 and myosin binding protein-C (MyBP-C), shown in ventral view, anterior to left.

Heart chambers show very low levels of Ttn T12 and MyBP-C. **D**, Single confocal scans

through ventricular sarcomeres of e25-/-, e232-/-, and sibling control embryos,

immunostained for α-actinin and myomesin-2 (Myom2). e25-/- embryos show punctate

aggregates of α-actinin and only low levels of Myom2. In (**C**) and (**D**), cardiac morphology

and sarcomeres in e232-/- embryos are similar to control. A denotes atrium, SH,

sternohyoideus muscle, and V, ventricle.

Figure 4. Skeletal muscle phenotypes in *ttn.2* embryos. A through **G,** Immunofluorescence

staining of 3 dpf homozygous and sibling (sib) control embryos for myosin heavy chain

(MyHC), fast myosin light chain (Fast MyLC), α-actinin, myomesin-2 (Myom2), Ttn.2 M6,

Ttn T12, and Ttn.1 M8-M9, shown in lateral view, anterior to left. MyHC is strongly

expressed in superficial slow fibers, while MyLC labels fast fibers. **A**, e25-/- embryos show

some good sarcomeric structure as well as regions of damaged sarcomeres in the middle of

the fiber (white arrowheads). e129-/- embryos have disorganized sarcomeres in both slow

(yellow arrowheads) and fast fibers, while e232-/- embryos have striated slow fibers but no

organized thick filaments in fast fibers. **B**, In e25-/- embryos, α-actinin (slow and fast fibers)

and Myom2 (fast only) seem striated. **C** through **G**, In e129-/- embryos, abnormal striation is

detected from 24 hpf to 3 dpf; MyHC and α-actinin and show no normal striation but some

radial organization is seen around random points in the fiber (**C,E**), Myom2 is abolished (**D**),

and Ttn.2 M6 is completely missing (**E**). This abnormal striation is also seen with Ttn T12

(**F**) and Ttn.1 M8-M9 (**G**). **H** through **M,** Electron micrographs of longitudinal sections of 3

dpf e129-/- and sibling control embryos. Thin filaments in mutant embryos are arranged in I-

Z-I brushes that are organized in radiating patterns (**H**, red arrows), highlighting structural

features (**I**), including triads (blue arrowheads), Z-disks (pink arrowheads) and thin filaments

(cyan arrowheads). **J**, Schematic illustrating arrangement of thick and thin filaments. **K,**

Electron micrographs of transverse sections of fibers in 3 dpf embryos showing lattice pattern

of thick and thin filaments in sibling controls and increased interfilament space in e129-/-. **L**

and **M,** Quantification of directionality of fibres. Rose diagram of representative somitic

muscle (**L**) and graph comparing the variance of the orientations (**M**). The possible range is 0

to 1, where 0 is completely ordered and 1 is completely random. Unpaired t-tests, mean ±

SEM.

Figure 5. Skeletal muscle phenotypes in e232-/- embryos. A, Brightfield images of 5 dpf

e232-/- and sibling (sib) control embryos, lateral view, anterior to left. Mutants show a

distinctive trunk curvature (red arrowhead) and fail to inflate swim bladder (black

arrowheads). **B** through **G,** Immunofluorescence staining of e232-/- and sibling (sib) control

embryos for myosin heavy chain (MyHC), fast myosin light chain (Fast MyLC), α-actinin,

Ttn.2 M6, Ttn.1 M8-M9, myomesin-1 (Myom1), myosin binding protein-C (MyBP-C),

myomesin-2 (Myom2) and Ttn T12, shown in lateral view, anterior to left. At 24 hpf (**B-D**),

MyHC, fast MyLC, α-actinin and Ttn.1 M8-M9 (slow) show regular sarcomeres although

Ttn.2 M6 is not detected. By 3 dpf (**E-H**), Myom1 (strong in fast fibers) and MyBP-C

(strong in slow fibers) are present in both slow (**E**, arrows) and fast fibers (**E**, arrowheads) in

controls but are detected only in slow fibers in e232-/- embryos. Myom2 is expressed only in

fast fibers (**F**, arrowheads), whereas α-actinin is in both slow (**F**, arrows) and fast (**F**,

arrowheads) fibers. In e232-/- embryos, α-actinin is normal in slow fibers (arrows) but is

disorganized and aggregated in the deeper fast fibers. Myom2 shows diminished signal above

background. **G**, Fibers are scanned at different levels to highlight superficial slow muscle

(left panels) and deeper medial fast fibers (right panels). Ttn T12 striation patterns are similar

in e232-/- and control embryos in slow fibers but mutants have a punctate disorganised

staining fast fibers. **H** through **I,** Electron micrographs of longitudinal muscle sections of 3

dpf e232-/- and sibling control embryos. Control embryos have organized sarcomeric

structure with clear Z-disk (Z) and M-band (M) visible in both slow and fast muscle fibers.

e232-/- embryos have organized sarcomeres only in slow muscle fibers. Thick filaments (red

arrowheads) are present in mutant fast fibers but are not arranged into sarcomeres.

Sarcoplasmic reticulum (orange arrowheads) is disordered in mutant fast fibers. Blue kites

indicate EM artefacts. Triads (green arrowheads) are located at the Z-disk in control fast

fibers. In mutant fast fibers, there are a small number of triads and Z-disk but these are not

associated together.

Figure 6. Ventricular size and function in adult heterozygous *ttn.2* zebrafish. Cardiac

function was assessed in 12 month-old wild-type (WT) and heterozygous *ttn.2* fish using high

frequency echocardiography. **A** through **F** show ventricular function under resting conditions:

A, indexed ventricular end-diastolic volume (EDV/BSA), **B**, ejection fraction (EF); **C**, global

longitudinal strain (GLS), **D**, peak ventricular outflow tract velocity (VOT), **E**, peak systolic

global longitudinal strain rate (GLSRs), **F**, isovolumic relaxation time (IVRT). **G** through **I**

show responses (at 2 hr) to acute adrenaline administration in 10-15 month-old fish. Two-

way ANOVA, mean $\pm$ SD, or Δmean $\pm$ SD.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6