Hydraulic modelling reveals untreated sewage, not pharmaceutical waste, drives antimicrobial resistance in a small river running through a big city

Vikas Sonkar[a,*], Arun Kashyap[a], Rebeca Pallares-Vega[b], Sai Sugitha Sasidharan[a], Ankit Modi[c], Cansu Uluseker[d], Sangeetha Chandrakalabai Jambu[a], Pranab Kumar Mohapatra[c], Joshua Larsen[e], David W Graham[b], Shashidhar Thatikonda[a,*], Jan-Ulrich Kreft[d,*], on behalf of the AMRflows consortium

[a]One Health Research Lab, Department of Civil Engineering (Environmental Engineering), Indian Institute of Technology Hyderabad, Kandi, Sangareddy, 502284, Telangana, India

[b]School of Engineering, Newcastle University, Newcastle upon Tyne, NE1 7RU, UK

[c]Department of Civil Engineering, Indian Institute of Technology Gandhinagar, Gandhinagar, Gujarat, 382355, India

[d]Institute of Microbiology and Infection & School of Biosciences, University of Birmingham, Edgbaston, Birmingham, B15 2TT, UK

[e]School of Geography, Earth and Environmental Sciences, University of Birmingham, Edgbaston, Birmingham, B15 2TT, UK

*Corresponding authors

E-mail addresses:

vikassonkar629@gmail.com (V. Sonkar)

shashidhar@ce.iith.ac.in (S. Thatikonda)

j.kreft@bham.ac.uk (J.-U. Kreft)

Abstract

Quantifying sources of antimicrobial resistance (AMR) in rivers receiving various waste streams is essential for targeting mitigation strategies yet rarely performed. This study combined field monitoring with hydraulic modelling and mass balance calculations to attribute sources of AMR in the Musi River running through Hyderabad, India, a city renowned for pharmaceutical manufacturing. We quantified antibiotic resistance genes (ARGs), resistant bacteria (ARBs), and physicochemical parameters in water and sediment samples in the dry and wet season. Absolute ARG and ARB abundances spiked in the city, declining again downstream. Changes were more gradual in the wet season. Pollution levels were significantly different between upstream, city and downstream stretches and seasons. Hydraulic modelling revealed that 60-80% or 20-40% of the river water in the city derived from untreated sewage during the dry or wet seasons, respectively. This established municipal waste, not pharmaceutical sources, as the dominant driver of AMR in the Musi. Linear discriminant analysis identified dissolved oxygen and total nitrogen as reliable proxies for distinguishing AMR-polluted from less-polluted sites. The Musi, with insufficient wastewater treatment and limited dilution of point-source loadings, is typical for many urban rivers in resource-limited countries highlighting the urgent need for improved wastewater management to reduce AMR exposures.

Keywords: Antibiotic resistance; Sewage; Pollution; Source apportionment; Mass balances; Water quality proxies

1. Introduction

Antimicrobial resistance (AMR) is among the greatest health threats of the 21[st] century, with an estimated 1.14 (95% UI 1.00–1.28) million deaths attributable to bacterial AMR in 2021, projected to rise to 1.91 (1.56–2.26) million deaths by 2050 (UNEP 2023; Naghavi et al. 2024). Although the primary human health risk stems from increasing resistance in human pathogens, their resistance is primarily conferred by antimicrobial resistance genes (ARGs) that can be acquired through horizontal gene transfer (HGT) from any antimicrobial resistant bacteria (ARBs) *via* mobile genetic elements (MGEs) (Martínez 2008). Consequently, ARGs, ARBs and MGEs are often used as surrogates for monitoring AMR spread across One Health sectors.

AMR prevalence is increasing disproportionately in Low- and Middle-Income Countries (LMICs) due to inadequate waste management and less regulated antimicrobial use, resulting in AMR-laden wastes released into the environment, fuelling AMR transmission and spread *via*

polluted water across One Health sectors (Klein et al. 2018; Karkman et al. 2019; Kashyap et al. 2023; Sonkar et al. 2024). In parallel, many LMICs are experiencing accelerated urbanisation that strains civil infrastructure. For example, 34% of India's population lived in urban areas in 2021, which is increasing at 2.4% per year (The World Bank 2022), exacerbating pressures on already overloaded infrastructure. This is especially acute for wastewater treatment plants (WWTPs). Although functional WWTPs reduce AMR release into receiving waters (Pallares-Vega et al. 2019; Beltrán de Heredia et al. 2023; Yang et al. 2024; Sonkar et al. 2025a), insufficient treatment capacity results in untreated discharges that transform receiving water bodies into conduits for AMR spread, including transmission of pan-resistant strains, and emergence through mutation and HGT (Graham et al. 2014; Wang et al. 2021). Despite the importance of quantitative source apportionment, it has remained rare in LMIC settings, limiting our ability to prioritize intervention strategies based on evidence.

The consequences of inadequate waste management and AMR spread are exemplified by conditions in the Musi River (**Fig. 1**). The Musi has significant economic and societal importance (Diwan et al. 2018), but has also become a major waste disposal site for municipal and industrial wastewater from the growing city of Hyderabad, Telangana (southern India). Importantly, Hyderabad is a global pharmaceutical manufacturing hub, infamous for high antibiotic pollution (Larsson et al. 2007; Gothwal and Shashidhar 2017; Konda et al. 2022, 2024). The construction of common effluent treatment plants (CETP) for industrial effluents and a zero liquid discharge policy for industries offers an opportunity to assess drivers of AMR spread in the Musi River, which is key to prioritising remedial actions to reduce AMR exposures across the watershed.

The Musi serves as a good model for assessing AMR fate and transport in urban river catchments in regions with warm climates (enabling rapid microbial turnover), insufficient WWTP capacity, and low dilution ratio where river discharge is small relative to waste loading (Caucci et al. 2016; Herrig et al. 2020; Kumar et al. 2020; Beltrán de Heredia et al. 2023). These conditions are representative of many rivers including several studied rivers such as the Code, Indonesia (Muurinen et al. 2022), Bagmati, Nepal (Amarasiri et al. 2022), Brisbane, Australia (Ahmed et al. 2021), Virilla, Costa Rica (Barrantes-Jiménez et al. 2025) and the Indian rivers Sabarmati (Ram and Kumar 2020), Mutha (Marathe et al. 2017) and Kshipra (Hanna et al. 2020). Nevertheless, past studies have rarely explained spatiotemporal variation of AMR levels in terms of point-source discharges and mass loadings using hydraulic modelling approaches.

Such quantitative frameworks are crucial for distinguishing levels of contribution from
industrial and various municipal inputs.

Here, we integrated field monitoring and hydraulic modelling to quantify drivers of AMR in
the Musi River running through Hyderabad city. Water and sediment samples were collected at
10 locations along different parts of the catchment spanning 153 km and during the wet and dry
season to assess how spatial and seasonal factors impact *in situ* AMR. The objectives were to:
(1) characterise the spatial and seasonal distribution of AMR determinants and physicochemical
conditions required for mathematical modelling and risk analysis, (2) assess correlations
between the different AMR determinants and with environmental factors, (3) quantify
contributions from municipal versus industrial point-sources through hydraulic modelling to
explain observed patterns, and (4) identify possible water quality proxies for cost-effective
AMR hotspot detection to guide interventions and prioritise sites for more detailed AMR
monitoring and analyses.

2. Materials and methods

100 This study adhered to EMBRACE-WATERS reporting guidelines (Hassoun-Kheir et al. 2021)
101 wherever possible, any deviations are indicated.

2.1. Study area and sample collection

The Musi is ~260 km long and a major tributary of the Krishna River. The average annual
rainfall in its catchment is ~800 mm, most of it falling in the monsoon or wet season (June to
September). The Musi riverbed is generally rocky, consisting primarily of pink granite
(Narshimha et al. 2020). Two reservoirs, Osman Sagar and Himayat Sagar, exist where the Musi
and Musa Rivers (a tributary of the Musi 60 km upstream of Hyderabad city) are blocked by
dams, respectively. These two dams are closed most of the year except for spillway overflows
during the wetter months. Therefore, flows in the Musi are often very low upstream of the city
(dependent on leaky dams), and most of the flow within the city is from wastewater discharges,
about 1.2 million m^3/d of partially treated municipal wastewater, mixed with industrial wastes
from across the city.

The wastewater discharged into the Musi is either untreated or released from municipal WWTPs
(**Supplementary File 1**). Two Common Effluent Treatment Plants (CETPs) for pharmaceutical
and industrial waste also discharge into the Musi catchment: the Patancheru Enviro Tech Ltd.
(PETL) and Jeedimetla Effluent Treatment Plant (JETL) release 1,600–2,000 and 50,000
m^3/day, respectively. Effluents from the PETL and JETL flow through a 26 km pipeline into the

Amberpet WWTP (339,000 m^3/day), where the industrial discharges mix with untreated

wastewater, before treatment and discharge into the Musi itself **(Fig. 1)** (CPCB 2005, 2010;

Lübbert et al. 2017). In terms of land use, the Musi catchment is both rural and urban, including

many residential communities, hospitals and factories. Downstream of the city, land use

includes agriculture and aquaculture, with less human impact than within the city. Sampling

sites were chosen to contrast upstream water conditions with sites in the city and downstream.

Since the Musi flows from a reservoir upstream of the city, only one upstream sampling point

was possible. Within the city, many locations were sampled, reflecting different inputs along

the river.

Water and sediment grab samples were collected at 10 locations along the Musi River flowing

through Hyderabad city, Telangana, Southern India during the wet (18-19 August 2022) and dry

(11-12 March 2023) seasons **(Fig. 1;** site details in **Supplementary File 1)**. Sampling occurred

on days without rainfall for the preceding five days to minimize storm-related variability. At

most sites, duplicate water samples were collected mid-stream from bridges using a bucket

suspended on a rope and transferred to pre-sterilised 2-L HDPE bottles. Sub-samples were

preserved immediately with H_2SO_4 for chemical oxygen demand (COD) analysis, whereas

water temperature, pH, dissolved oxygen (DO), total dissolved solids (TDS) and conductivity

were measured on-site with an Orion Star A329 portable multimeter with smart probes

(ThermoFisher Scientific, USA). Duplicate sediment samples (0-15 cm depth) were collected

3 metres from the riverbank using a grab dredger, sieved through a 2 mm sterile mesh, and

transferred into pre-sterilised polypropylene bags (approx. 1 kg). All samples were stored in the

dark, on ice in a cooler and transported back to the lab for further processing and analysis.

2.2. Water and sediment quality analysis

Water quality and sediment physicochemical parameters were measured using standard

protocols (**Table S1** and **S2**). When required for analysis (*i.e.*, organic carbon, total nitrogen

(TN), ammonia, nitrate, and nitrite), water samples were filtered through 0.45 μm

polyvinylidene difluoride (PVDF) membrane filters (Merck Life Sciences, Merck, India) before

analysis. To establish a common basis for concentrations of bacteria and genes across sediment

samples with different mineral content, dry matter (DM) and organic matter (OM) fractions

were quantified using standard methods (**Table S2**).

Fig. 1. The basin of the Musi River in Telangana, Southern India with the locations of sampling sites (numbered 1-10) and WWTP effluent discharges. A total of 25 point-sources of untreated sewage discharged through drains or treated sewage and other sources were identified within Hyderabad, between sampling points 1 and 8, see **Supplementary File 1** for details.

2.3. Enumeration of *E. coli*, heterotrophic bacteria and ARBs

Seven bacterial targets were enumerated in water and sediment samples across seasons following methods described in Sonkar et al. (2025a). Briefly, *E. coli* was quantified on Tryptone Bile X-β-D-glucuronide agar (TBX) (HiMedia, India) as blue-green colonies (TBXB), incubated at 44°C (ISO 16649-3:2015 2015), as an indicator of faecal pollution and sentinel for AMR prevalence (Anjum et al. 2021). 'Total' Gram-negative (GN) heterotrophic bacteria were enumerated on SDS-supplemented OECD synthetic sewage medium (OSS) at both 30°C for environmental and 44°C for gut-associated bacteria. Extended-spectrum β-lactamase (ESBL)-producing and carbapenem-resistant (CARB) GN heterotrophic subpopulations were enumerated on OSS supplemented with appropriate selective antibiotics (Sonkar et al. 2025a). These clinically relevant resistance phenotypes were selected as GN bacteria account for the majority of global AMR-attributable mortality (Naghavi et al. 2024). All samples were plated in triplicate within 24 hours of collection, and colony counts recorded after 24 hours of incubation.

2.4. Measurement of ARGs by qPCR

DNA was extracted as previously described Sonkar et al. (2025a). The 16S rDNA gene was quantified as a proxy for total prokaryotes and to compare with culturing data. ARG targets were chosen according to Berendonk et al. (2015) and to reflect resistance to antibiotics used and produced in Hyderabad (**Fig. S1**). Genes included: fluoroquinolones (*qnrS*), aminoglycosides (*aph(3″)-Ib*), sulfonamides (*sul2*), carbapenems and third-generation cephalosporins (*bla$_{NDM}$* and *bla$_{CTX-M}$*), macrolides (*ermF*), and tetracyclines (*tetW*). Two more genes were included: *uidA* to quantify *E. coli* numbers (Frahm and Obst 2003; Silkie et al. 2008) and *intI1* to quantify class I integrons as a proxy for MGEs (Gillings et al. 2015). Based on preliminary analysis, DNA extracts were diluted 10-fold to reduce inhibition. The protocol was adapted from Pallares-Vega et al. (2019) as previously described Sonkar et al. (2025a). Concentrations of forward and reverse primers, amplicon sizes of target genes and annealing temperatures are summarised in **Table S3**.

2.5. Quantifying pollutant point-sources along the Musi River

To quantify the drivers of AMR in the Musi River, we developed a hydraulic mass balance model to estimate river discharge and pollutant loadings from point-sources. We identified 25 point-source inputs throughout the catchment, including untreated municipal wastewater, treated WWTP effluents, agricultural runoff, and mixed wastes, and estimated volumetric flow and mass loadings from each source (**Fig. S2, Supplementary File 1**).

For each location, wastewater production rates were estimated based on local population sizes and the average municipal wastewater production rate (**Fig. S3**) specifically for Hyderabad (detailed methods given in **SI Section 1.1, data in Supplementary File 2**). Further, flow records of WWTP and reservoir discharges were obtained. Catchment sizes and locations were based on official records and digital elevation maps. For corroboration, an alternative approach to estimate population sizes was followed, based on census data combined with satellite images to identify dwellings and allocate people to dwellings (High Resolution Population Density Maps + Demographic Estimates).

Results from the two approaches were reasonably consistent, comparing estimates using a hypothetical tracer that reflected wastewater inputs from point-sources. Our "tracer" was meant to reflect the typical composition of wastewater and was set to measured values of COD in untreated (normalised to 100%) and treated wastewater (16%) (Sonkar et al. 2025a). Values for VS (12%), 16S (19%) and total ARGs (22%) in treated wastewater were similar (see **Supplementary File 1**). Mass balance and loading calculations for the tracer were performed using upstream river discharge and concentration data and estimated point-source discharges to determine the relative impact of each point-source on river water quality (**Supplementary File 1**). For simplicity, the tracer was assumed to be inert.

2.6. Data processing and statistical analysis

Statistical analyses were performed with R 4.3.0 (R Core Team 2018), mainly packages ade4, vegan and MASS using the RStudio 2023.03.0 desktop (https://www.rstudio.com/) and fully documented in an R notebook (**Supplementary File 3**). The methods and approach, including Hierarchical clustering, Principal Component Analysis (PCA), Linear Discriminant Analysis (LDA) and PERMANOVA, were guided by Borcard et al. (2018), but we are responsible for the choices made. Two clustering methods and two dissimilarity measures were used to ensure that results were robust and not dependent on specific choices. We only report results supported by all method combinations. Ricotta and Podani (2017) was used for guidance on the choice of dissimilarity measures. Whether data should be scaled or standardised was decided after exploring and analysing the data. It was not possible to perform sample size (statistical power) calculations *a priori* due to a lack of preliminary data.

3. Results and discussion

3.1. Spatio-temporal patterns of environmental conditions and AMR determinants along the river

3.1.1. Environmental condition gradients

All analysed water quality (WQ) parameters followed a distinct spatial gradient along the Musi River (**Fig. S4**). Water from city sites was severely contaminated across seasons, with high concentrations (mean±SD, mg/L) of COD (168±97), TOC (14±5), NH_3-N (62±55), TN (26±13) and TP (26±17), whereas DO remained near-anoxic (<1 mg/L). Several WQ parameters (pH, EC, COD, etc.) exceeded permissible limits set by India's Central Pollution Control Board (CPCB) within the city, classifying the water as 'very poor' (CPCB, 2005). Downstream sites exhibited natural attenuation of pollution (COD 74±29; DO 5±1), though levels remained elevated relative to the upstream site (COD 37 ±4; DO 7±2) (**Fig. S4**). In contrast, only some of the corresponding sediment physicochemical conditions showed distinct gradients with conductivity, ammonia and organic matter elevated in the city (**Fig. S5**).

3.1.2. Antibiotic resistance and other genes

Seven ARGs and the *intI1, uidA* and 16S rDNA genes were quantified in river water (**Fig. 2**) and sediment samples (**Fig. S6**) from the wet and dry seasons. The most abundant genes ($\log_{10}$ copies/mL) in city water samples for both seasons were the 16S rDNA (7.81±0.81), followed by *intI1* (6.53±0.96), *sul2* (6.17±0.82), *aph(3")-Ib* (6.11±1.09), *tetW* (5.56±1.06), *ermF* (5.55±1.53) and *qnrS* (5.03±1.09). Conversely, *bla*$_{CTX-M}$ (3.97±0.84), *bla*$_{NDM}$ (3.72±0.84), and *uidA* (3.35±1.24) concentrations were lowest in both seasons. The abundances of all ten genes followed consistent spatial gradients similar to WQ, with the lowest levels found at the upstream site, highest within the city, dropping to more or less lower levels downstream (**Fig. 2**). During the wet season, water column gene levels increased more gradually as water passed through the city and also declined more gradually downstream. The ARG abundances in the city sites were substantially higher than those in anthropogenically-impacted European rivers such as the Suze River in Switzerland (Lee et al. 2021) and the Danube River in the most polluted stretches in Central Serbia and the tributary Arges river in Romania (Kirschner et al. 2024; Schachner-Groehs et al. 2024), but comparable to concentrations in raw municipal wastewater in Hyderabad (Javvadi and Mohan 2023, 2024; Sonkar et al. 2025a).

Unlike water samples, no consistent seasonal patterns in gene levels were evident in the sediment samples, although concentrations were generally higher, albeit more variable, within the city (**Fig. S6**). The high concentrations in the sediments suggest that sediment acts as an

important ARG reservoir, accumulating ARGs from the point-sources under low-flow
conditions during the dry season. Overall, the analysed genes in water and sediment columns
indicated that the Musi in Hyderabad was highly polluted with ARGs in both seasons, although
spatial patterns of detected genes differed between sediment and water column samples. On a
positive note, ARGs declined downstream of the city, suggesting natural recovery was rapid
once pollution dropped.

Fig. 2. Absolute abundance of genes along the Musi River in dry (red) and wet season (blue) water samples. All panels have the same scale

showing log transformed absolute abundances of 16S rDNA, *uidA*, *intI1* and seven ARGs. The river can be divided into three stretches: (a) upstream

(light blue), (b) within the city (dark blue) and (c) downstream of the city (medium blue). Point-sources of raw wastewater, treated wastewater, a

mixture of both, reservoir flows, and agricultural runoff are indicated by symbols. See **Fig. S6** for equivalent sediment sample data.

3.1.3. Bacterial abundances and their correlations with genes

'Total' Gram-negative heterotrophic bacteria (OSS), *E. coli* (TBXB), and ESBL- and CARB-resistant bacteria were quantified in water and sediment samples. In general, incubation at 30°C or 44°C did not alter spatio-temporal distribution patterns of total GN-heterotrophic bacteria or *E. coli* in the water samples (**Fig. S7**; see **SI Section 2.1.** for details). That is, viable bacterial concentrations ($\log_{10}$ CFU/mL) were lower upstream (0.46), much higher in the city (4.88±0.49), and lower again downstream (2.15±1.17) in the dry season. These spatial patterns were similar to those observed for ARGs (**Section 3.1.1**), indicating consistent pollution sources driving both bacterial and resistance gene abundances. In contrast to the dry season, bacterial abundances were elevated everywhere in the wet season, similar to levels in dry season city samples (**Fig. S7A**). This pattern aligns with studies of the Ganga river, India (Prasad et al. 2024) and the Danube river, Europe (Schachner-Groehs et al. 2024), where municipal wastewater discharges drive distinct spatial patterns of bacterial contamination.

The pattern for ESBL-resistant bacteria was also similar with abundance below detection limit (BDL) upstream, higher in the city ($\log_{10}$ 4.54±0.63 CFU/mL), and returning to BDL downstream in the dry season (**Fig. S8A**). In the wet season, levels in the city were similar to those found within the city during the dry season. Conversely, CARB-resistant bacteria were BDL in the city (and elsewhere) in the dry season, while the wet season pattern reflected ESBL. The consistently higher ESBL and CARB counts in the wet season (**Fig. S8A**) suggest elevated flows reduce sedimentation of water column bacteria and/or increase scouring of organisms previously deposited in sediments, which can occur in rivers in monsoonal climates (Knapp et al. 2012).

Sediment bacterial abundances exhibited seasonal variation similar to water samples, generally higher in the wet season than in the dry season across all media (**Fig. S7B**) (for values see **Supplementary File 4**). In contrast to the distinct spatial gradients observed in water samples, sediments showed no discernible upstream to downstream patterns during the dry season. During the wet season, abundances were slightly higher in the city and downstream sediments compared with upstream, though these differences were less dramatic than those in the water samples. ESBL- and CARB-resistant GN heterotrophs were with one exception all BDL in sediments in the dry season, while they were all at similarly higher levels in the wet season (**Fig. S8B**). This seasonal dichotomy of bacterial abundance suggests that increased wet season flows cause sediment resuspension, mobilising sediment-associated resistant bacteria into water

column, while concurrent urban runoff contributes additional microbial inputs to the river (Heß et al. 2018).

Five hypotheses were tested regarding relationships between viable counts at different conditions and viable counts versus genes quantified by qPCR: (i) counts at 44°C would correlate with but be less than counts at 30°C on the same medium; (ii) counts on CARB media would correlate with but be less than counts on ESBL media; (iii) counts on ESBL media would correlate with *E. coli* counts because both relate to levels of municipal wastewater pollution; (iv) counts on OSS media would best correlate with 16S rDNA gene data; and (v) counts of *E. coli* would correlate with *uidA* gene concentrations. Data were consistent with all five hypotheses, but no firm conclusions could be made because of the highly scattered viable count data (**Figs. S9, S10, S11** and **Table S4**). The gene data were clearly more consistent across samples and therefore used in most subsequent numerical analyses.

3.2. Multivariate analysis of patterns of AMR determinants and environmental conditions
Visualizing gene abundances, bacterial counts, and water quality parameters indicated that the Musi catchment can be divided into three stretches: upstream, city, and downstream (**Figs. 2, S4, S7**). To objectively validate these patterns and test for statistical significance, hierarchical cluster analysis (HCA), principal component analysis (PCA), and permutational multivariate analysis of variance (PERMANOVA) were employed.

3.2.1. Water column patterns
Hierarchical clustering of water samples revealed season-dependent spatial patterns. In the dry season, city sites formed a distinct cluster, whereas upstream and downstream sites clustered together (**Fig. 3A**). Conversely, in the wet season, city sites split with late-city sites remained a distinct cluster, but early-city sites now clustered with downstream samples—the upstream site remained distinct (**Fig. 3B**). This in-city split suggests the city stretch contains two sub-stretches in the wet season, that merge into a single pollution zone during the dry season. These clustering patterns were based on UPGMA clustering with Euclidean distance, but similar clusters were obtained for all four combinations (UPGMA or Ward's clustering combined with Euclidean distance or Bray-Curtis dissimilarity), confirming the robustness of the results (see **Supplementary File 3**). PCA corroborated these patterns, showing a tight cluster of all city sites in the dry season and a more spread and mixed pattern in the wet season (**Fig. 3C-F**). All seven ARG vectors pointed in similar directions in PCA ordination space, indicating similar AMR pollution sources in the river.

Since the sampling design was not balanced (one upstream, seven city, and two downstream sites), and variables were not independent and normally distributed, PERMANOVA (adonis2 from the vegan package) was used to test for significant spatial and seasonal differences. Both stretch and season were significantly different ($p < 0.05$) for most water column variables (**Table S5**). Stretch was significant for all 10 genes, six of seven WQ parameters (except NO_2^-), and all bacterial counts, *i.e.*, 19 of 20 parameters (ARB data were insufficient for analysis). Season was significant for nine of 10 genes (except *tetW*), five of seven WQ parameters (except DO and NO_2^-), and all bacterial counts, *i.e.*, 17 of 20 variables. These results confirm that both spatial location and seasonal hydrology significantly structure AMR determinant distributions in the Musi water.

3.2.2. Sediment column patterns

In contrast to water sites, sediment sites showed weaker spatial and temporal differentiation. Stretch was significant for only five of 10 genes (*intI1, aph(3")-Ib, ermF, qnrS, tetW*), the sum of ARGs, two of seven SQ parameters (total nitrogen and conductivity), and two of three bacterial counts, *i.e.*, nine of 20 parameters in total. Season was significant for only two genes (*uidA, bla_{CTX-M}*), but six of seven SQ parameters, and all bacterial counts (**Table S5**). The attenuated spatial differentiation in sediments likely reflect long-term accumulation coupled with transport of ARGs, which obscures the longitudinal gradients observed in water samples. Thus, sediments function as integrative archives of historical contamination rather than reflecting instantaneous pollution dynamics (Heß et al. 2018).

Fig. 3. Hierarchical clustering and ordination of water samples comparing dry and wet seasons based on Log₁₀ transformed absolute abundances of seven ARGs. Sampling locations are numbered along the river: upstream (1), city (2-8) and downstream (9-10). **(A-B)** Hierarchical clustering (UPGMA) of Euclidean distances shows the separation of the city from up- and downstream samples in the dry season, whereas only late city sites cluster in the wet season. **(C-F)** PCA biplots for dry **(C-D)** versus wet **(E-F)** seasons. Scaling 1 correctly represents sites with their ARG profiles while Scaling 2 correctly represents genes (vectors).

3.2.3. Relationships between water and sediment conditions

Before making comparisons, the water and sediment ARG concentrations were scaled to account for different types of concentrations, *i.e.*, gene copies per mL for water samples and gene copies per mg of organic matter (OM) for sediments (see R notebook for scaling details). Concentrations were expressed relative to organic matter rather than dry mass because inorganic particles such as sand contributes to dry mass but are inaccessible to bacteria and their genes, and inorganic content varied substantially across sites.

These scaled data revealed season-dependent coupling between water and sediment ARG profiles (**Fig. 4**). In the wet season, sediment and water samples clustered separately, suggesting higher river flows decoupled their ARG content. This decoupling likely reflects increased lateral dispersion in the water column and increased sediment resuspension combined with reduced exchange between consolidated sediments and overlying water. Conversely, dry season clustering showed selective coupling: city water samples remained distinct, while upstream and downstream water samples clustered to some extent with their corresponding sediments. Inner-city sediment samples formed separate sub-clusters in both seasons, indicating persistent, high-level AMR contamination year-round in central Hyderabad. This seasonal shift suggests spatially homogeneous AMR pollution during low-flow conditions, with spatial differentiation emerging during elevated flows as it takes more point-sources to pollute the river after entering the city and then spreading further downstream. Further interpretations of these multivariate patterns and their connections to linear discriminant analysis are provided in **Section 3.4 (SI Section 2.2)**.

Fig. 4. Joint clustering of sediment and water samples along the Musi in the wet and dry seasons. Sampling locations are numbered along the river: upstream (1), city (2-8) and downstream (9-10), while W represents water and S represents sediment samples. This joint analysis of water and sediment samples is based on Log$_{10}$-transformed absolute abundances of ARGs. Since water and sediment ARG concentrations had a different basis, data were scaled to make them more comparable. (**A-B**) Joint hierarchical clustering of the Euclidean distances between ARGs, either in the dry (**A**) or wet season (**B**). (**C-D**) PCA biplots, either in the dry (**C**) or wet (**D**) season. These biplots used Scaling 2, which correctly depicts genes (vectors).

3.3. Source apportionment of AMR determinants in the Musi

3.3.1. Hydraulic modelling and point-source loading quantification

To explain observed spatial patterns of ARGs and ARBs in the Musi, we identified and quantified 25 point-sources of wastewater and other inputs within the area of investigation and calculated mass loadings using discharge estimates and a conservative tracer approach

(**Supplementary File 1**). Point-source discharges were quantified using operational records for WWTP discharges and population-based estimates for raw wastewater production using population density and per capita wastewater production rates from local government and other sources (150±25 L/d; see **SI Section 1.1**). The hypothetical inert tracer concentration was set to 1.0 for untreated sewage and 0.16 for treated WWTP effluent, based on measured COD remaining in WWTP effluents and consistent with what remains of volatile solids (12%), 16S rDNA (19%), and total ARGs (22%) in effluents from the four main WWTPs along the Musi (Sonkar et al. 2025a). Details of point-sources, sampling sites and discharge, tracer, and loading calculations are provided in **Supplementary File 1**.

Mass balance calculations for the tracer suggested that 60-80% of river water in the city stretch was derived from raw wastewater inputs in the dry season, with total bacterial abundance (16S rDNA) ranging from $2\text{-}6\times10^8$ copies/mL (**Figs. 5A, S12**). Sharp increases in bacterial abundance coincided spatially with tracer spikes at point-source discharge locations, validating the mass balance approach. In contrast, during the wet season, monsoon runoff and increased discharges released from Osman Sagar and Himayat Sagar reservoirs provided higher dilution, reducing raw wastewater contributions to 20-40% of river flow and decreasing bacterial abundance by an average of 1.52 $\log_{10}$ ($0.19\text{-}8\times10^7$ copies/mL) compared to the dry season (**Figs. 5C, S12**). Importantly, WWTP discharge points (tracer = 0.16) led to localized decreases in bacterial abundance and ARG levels, demonstrating that WWTPs improved local water quality (**Fig. 5A**). This is in stark contrast to countries where the bulk of the wastewater is treated before entering rivers, so WWTPs pollute the rivers rather than making them cleaner as in the case of the Musi (Brown et al. 2019; Lee et al. 2021; Kirschner et al. 2024).

Predicted spikes of the tracer and untransformed 16S rDNA gene abundances (**Fig. 5A, C**) both indicated that bacterial levels (which mirror ARGs and other measured variables) increased sharply at point-source locations. However, 16S abundances declined progressively downstream of these inputs, in contrast with the tracer that was assumed to be inert. The downstream decline of the 16S abundance was likely due to microbial die-off, which was not quantified and may vary with environmental conditions. Nevertheless, the tracer calculations allow the quantification of untreated municipal wastewater inputs into the river. Interestingly, the observed data imply that the Musi River itself acts as a "pseudo-WWTP" for the city's wastewater, reducing carbon levels by about 59% (see **Supplementary File 1**). A similar phenomenon was seen in community sewer lines in an European study on the fate of AMR at urban catchment scales (Li et al. 2021). However, the ecological consequences of this

wastewater dominance are severe as city stretch waters caused 100% mortality in zebrafish (*Danio rerio*) embryos within 96 hours post fertilization, compared to 0% mortality when exposed to upstream water (Sonkar et al. 2025b), indicating that the Musi River has been transformed from a viable aquatic ecosystem into a de facto wastewater conveyance channel.

These mass balance calculations show the importance of considering point-source mass loadings (*i.e.*, mass transferred per time), rather than just concentrations, when developing practical solutions to problems like AMR spread in rivers. The current "WHO Guidance on waste and wastewater management in pharmaceutical manufacturing with emphasis on antibiotic production" is based on point-source concentrations, assuming a fixed dilution factor of 10 upon entry into a river. This concentration-focused approach creates a regulatory loophole as facilities can achieve compliance through pre-discharge dilution without reducing actual pollutant mass inputs. Such a loophole can be addressed by also stipulating load limits in environmental standards, especially when considering AMR genes and bacteria released into the environment.

Fig. 5. Direct comparison of the absolute abundance of the 16S gene in the river with the estimated wastewater fraction in both dry and wet seasons. The tracer concentration was set to 1 in raw wastewater and to 0.16 in treated wastewater to track wastewater loads. The tracer was assumed to be inert in contrast to 16S rDNA levels (•), which declined along the river. Normalized discharge (blue stairs), normalized loadings (+) and the fraction of wastewater in the river based on tracer mass balance calculations (brown stairs).

3.3.2. Pharmaceutical manufacturing contributed far less than municipal wastewater

India, the world's third-largest pharmaceutical producer with over 400 rivers and approximately 2.4 million other freshwater bodies (Rathinavelu et al. 2024), faces dual AMR pollution risks. Two questions guided this analysis: 1) how typical is the Musi of Indian rivers and 2) do manufacturing waste inputs alter AMR in the river?

Regarding the first question, rapid urbanisation in India has greatly outpaced the development of new WWTPs and supporting sewerage infrastructure. The "National Inventory of Sewage Treatment Plants" of India (CPCB 2021) indicates that only 28% of municipal wastewater is treated nationwide and only 17% in WWTPs compliant with discharge standards. Hyderabad treats about 41% of its wastewater (see **Supplementary File 1**), above national average, yet almost 60% of municipal wastewater remains untreated and is discharged directly into the environment. This infrastructure deficit makes the Musi representative of many Indian rivers where treatment capacity lags behind population growth, a similar pattern also observed globally in rapidly urbanising countries (Jones et al. 2021).

Regarding the second question, a distinctive aspect of the Hyderabad sewershed is that it has extensive antimicrobial manufacturing and processing units, which discharged wastewater into the local sewer drains until the construction of two treatment plants (see below). Hyderabad has been infamous for pollution from manufacturing since Larsson et al. (2007) found ciprofloxacin concentrations in surface waters near the city up to 31 mg/L. Subsequently, ciprofloxacin concentrations as high as 5.5 mg/L were found in the Musi during the dry season (Gothwal and Shashidhar 2017). These concentrations were measured after the construction of two Common Effluent Treatment Plants (PETL in 1995 and JETL in 1989) designed specifically for the antimicrobial manufacturing cluster, which discharge treated effluents into the Amberpet municipal WWTP for additional treatment and dilution before reaching the Musi River, indicating ongoing illegal dumping. Recent studies found much lower concentrations of antimicrobials in surface waters in the region, suggesting stricter enforcement has reduced illegal discharges into the Musi (Konda et al. (2024, 2022); **Table S6**). For example, ciprofloxacin levels were much lower than previous observations, *i.e.*, only 19.3 µg/L in local sewer drains and 7.28 µg/L in local industrial effluents and much lower in the Musi water. Based on our mass-balance calculations, the PETL/JETL discharge becomes diluted upon entering Amberpet WWTP and again upon entering the Musi to the extent that only 4% of the river water after Amberpet WWTP derives from PETL/JETL discharge (dilution factor of 27), and these calculations did not consider treatment and other losses along the sewer flow path (see

Supplementary File 1). Therefore, the manufacturing waste inputs and impacts on the Musi

River appear to be small relative to municipal inputs at the catchment scale. Previous studies

detected several antibiotics and ARGs (*aph(3″)-Ib*, *ermF*, *qnrS*, *sul2*, *bla$_{CTX-M}$*, *bla$_{NDM}$*) in drains

of Hyderabad (Javvadi and Mohan 2023, 2024), the same genes were abundant in the Musi as

expected from local antibiotic sales patterns in Hyderabad (**Fig. S1**). However, the spatial

concordance between ARG levels and municipal drain inputs (**Figs. 2, 5**), combined with

volumetric dominance demonstrates that nowadays untreated municipal wastewater drives

AMR pollution in the Musi River at the catchment scale.

3.4. Water quality parameters as proxies to predict AMR hotspots

3.4.1.   Linear discriminant analysis identifies DO and TN as suitable proxies

Having established that municipal wastewater drives AMR pollution in the Musi (**Section 3.3**),

we investigated whether simple water quality parameters could serve as cost-effective proxies

for identifying AMR hotspots, enabling rapid screening in resource-limited settings. Water

quality indicators (DO, COD, TOC, TN, etc.) exhibited spatial patterns mirroring ARG and

bacterial distributions: low pollution upstream, very high in the city, and moderate downstream

(**Fig. S4;** detailed explanation in **SI Section 2.3**). This concordance suggests wastewater

pollution as the common cause for water quality degradation and AMR prevalence, indicating

that traditional water quality indicators may serve as surrogates for AMR contamination.

To quantitatively test whether water quality parameters could discriminate river stretches across

seasons and predict AMR prevalence, Linear Discriminant Analysis (LDA) was performed on

z-score standardised water quality (WQ) data (**Fig. S13**). Our full LDA model using all seven

non-collinear WQ parameters (WQ7: pH, DO, COD, $NO_2^-, NO_3^-, NH_4^+, TN$) clearly

discriminated between stretches and seasons with 100% classification accuracy (**Fig. 6A**). To

develop the simplest field deployable model, parameters were removed from the full model in

stages (since there were 127 possible combinations of the seven parameters). By removing

variables individually, two were found to have little effect, resulting in a model with four

variables (WQ4: pH, DO, COD, TN) with 95% accuracy that maintained clear stretch

separation and was logical in terms of river water chemistry and microbiology (**Fig. 6B**).

Further simplification demonstrated that the two-parameter model combining DO and TN

achieved 90% discrimination accuracy, superior to all other two-parameter combinations,

representing a good balance between analytical simplicity and predictive power (**Fig. 6C**).

Although the DO+TN model was less accurate than WQ4, it could provide a practical, cost-

effective tool for rapid AMR hotspot identification in resource-limited settings.

Ott et al. (2021) had previously proposed the use of DO and ammonia as proxies for AMR pollution based on the Skudai River, Malaysia. Applying their DO+ammonia model to the Musi data provided similar results to our DO+TN model, while the latter proved better for the Musi data (**Fig. S14**). This difference reflects distinct nitrogen cycling dynamics between rivers, as explained below.

3.4.2. Nitrogen cycling dynamics explains why TN is a good proxy in rivers like the Musi

The superiority of TN over individual nitrogen species (ammonia, nitrite, nitrate) warrants a mechanistic explanation especially as this has practical implications for monitoring diverse aquatic environment. Wastewater inputs introduce organic carbon and nitrogen while stimulating microbial respiration, which depletes DO. In the Musi, DO declined to near-anoxic levels (<1 mg/L) in the city stretch during the dry season, creating conditions for denitrification that consumed residual nitrite and nitrate (**Fig. S4**). Downstream, reduced wastewater loading and re-oxygenation increased DO, halting denitrification and enabling nitrification, which oxidised ammonia to nitrite and nitrate. This caused transient spikes in oxidized nitrogen species at the downstream site 9 (**Fig. S4**) despite declining wastewater influence. Consequently, ammonia, nitrite, and nitrate concentrations fluctuated along the river dependent on the presence of oxygen, making each individual species an unreliable indicator of total wastewater loading. In contrast, TN integrates all nitrogen forms and is conserved during these transformations, providing a more stable proxy for cumulative wastewater inputs regardless of local redox conditions. PERMANOVA confirmed TN significantly distinguished river stretches ($p < 0.001$) (**Table S5**). This explains why Ott et al. (2021) found ammonia to be a reliable proxy as the Skudai River remains oxygenated with low N-cycling, so ammonia concentrations directly reflect wastewater inputs. However, when DO varies like in the Musi, representative of many LMIC urban rivers receiving high organic loads, nitrogen speciation varies dynamically, suggesting DO+TN could be a more robust proxy for wastewater pollution and associated AMR.

A LDA of 6 groups explained by water quality parameters: accuracy = 1, terms used: pH, DO, COD, Nitrate, Ammonium, Nitrite, TN Text represents actual groups, colour represents classification

B LDA of 6 groups explained by water quality parameters: accuracy = 0.95, terms used: pH, DO, COD, TN Text represents actual groups, colour represents classification

C LDA of 6 groups explained by water quality parameters: accuracy = 0.9, terms used: DO, TN Text represents actual groups, colour represents classification

Fig. 6. Linear discriminant analysis (LDA) showing that the three river stretches identified by clustering and ordination of ARGs can be discriminated by environmental characteristics in both seasons (resulting in 6 groups). Text labels combine actual stretch (Up, City, Down) and season (Dry, Wet) while colours indicate the LDA classifications of the sites, which are mostly consistent (accuracy of classification 90%-100%). Ellipses show 95% confidence intervals. **(A)** Discrimination using the full set of seven independent water quality parameters (WQ7). **(B)** Discrimination without the N-cycling ammonium, nitrite and nitrate (WQ4). **(C)** Discrimination with only DO and TN as water quality parameters (DO plus TN).

3.5. Study limitations

Although EMBRACE-WATERS guidelines for studying AMR in wastewater and aquatic environments were followed (Hassoun-Kheir et al. 2021), our study had sampling limitations and analytical challenges. Downstream sampling sites were too far away to be sampled on the same day. Resource limitations restricted sampling to once per season, precluding a more extensive temporal analysis. The study faced three analytical challenges. First, variability in sediment compositions, ranging from mineral-rich (sandy) to organic-rich samples, complicated comparisons between samples. We addressed this by normalising concentrations to the organic fraction (organic matter) of the sediments. This approach, while not ideal, was deemed the best available option. Further research across diverse sediment types is necessary to refine this method for future updates of the EMBRACE-WATERS guidelines. Second, because of using organic matter as denominator for concentrations in sediment samples, comparisons to water samples with volume as denominator were not straightforward. To facilitate this comparison, ARG concentrations in water samples were scaled to align with average sediment concentrations, but this is unlikely to have made water and sediment concentrations fully comparable. Third, correlations between viable bacterial counts and gene copy numbers from qPCR were poor due to large scatter in count data, likely due to differences in the sample preparation methods: larger water volumes (~200 mL) for qPCR should be less affected by small-scale environmental heterogeneity than the diluted 100 µL samples for viable counts. Moreover, viable counts will be affected by variable detachment of cells from particle surfaces during sample homogenisation while filtration for DNA extraction would not be affected. This highlights the need for methodological improvements and standardisation to enable reliable data integration across methods.

4. Conclusions and future directions

This study provides the first comprehensive apportionment of AMR sources in an Indian urban river by integrating field monitoring and hydraulic modelling to quantify pollution sources identified practical AMR surveillance tools for rapidly urbanising regions. Mass balances

suggested that, in the city stretch, 60-80% and 20-40% of river flow derived from untreated municipal wastewater during dry and wet seasons, respectively. Strong correlations between predicted wastewater fractions and measured AMR determinants validated this attribution. Despite Hyderabad's concentration of pharmaceutical industries, pharmaceutical manufacturing and processing, waste discharges into the Musi likely had minimal impact on *in situ* AMR as only ~4% of the river water following the Amberpet WWTP effluent point-source derived from industrial treatment plant effluent. Given that Hyderabad has greater than average WWTP coverage than other cities in India, it is probable that sources and conditions observed in the Musi are common in India and other countries with insufficient sewerage and wastewater treatment infrastructure. LDA showed that simple water quality markers such as DO and TN can serve as proxies for local stakeholders to identify AMR hotspots in freshwaters and to target interventions. DO and TN mirror sewage inputs, which often dominate AMR determinant loadings, particularly in rapidly urbanising countries where population growth has outpaced treatment infrastructure. These proxies are especially valuable because DO and TN are inexpensive to measure and DO is already included in current river health monitoring programmes. However, these proxies need validation in other rivers, especially when coupled with field-based methods such as DO test strips, to empower citizens to monitor and identify local AMR hotspots in their own catchments. Finally, this work highlights the value of simple hydrological and hydraulic modelling in predicting pollution loads, dilution factors, and concentrations in receiving rivers based on publicly available data. This approach identifies areas of greatest AMR exposure risk and transmission potential using mass loading data rather than concentration data in field assessments. Our methodology provides a framework for identifying and quantifying exposure pathways in future risk assessments, which are critical to better understand, prevent, and prioritise mitigation actions against AMR in places with insufficient WWTP capacities like the Musi River.

CRediT authorship contribution statement

Vikas Sonkar: Conceptualization, Methodology, Data curation, Investigation, Formal analysis, Validation, Visualization, Writing – original draft, review & editing. **Arun Kashyap:** Conceptualization, Methodology, Data curation, Investigation. **Rebeca Pallares-Vega:** Conceptualization, Methodology, Investigation, Formal analysis, Validation, Writing – review & editing. **Sai Sugitha Sasidharan:** Methodology, Data curation, Investigation. **Ankit Modi:** Methodology, Investigation, Data curation, Formal analysis. **Cansu Uluseker:** Methodology, Investigation, Data curation. **Sangeetha Chandrakalabai Jambu:** Conceptualization, Writing – review & editing, Project administration. **Pranab Kumar Mohapatra:** Conceptualization, Formal analysis, Writing – review & editing, Supervision, Funding acquisition, Project administration. **Joshua Larsen:** Conceptualization, Formal analysis, Writing – review & editing, Supervision, Funding acquisition. **David W Graham:** Conceptualization, Formal analysis, Writing – original draft, review & editing, Funding acquisition. **Shashidhar Thatikonda:** Conceptualization, Investigation, Methodology, Writing – review & editing, Supervision, Funding acquisition, Project administration. **Jan-Ulrich Kreft:** Conceptualization, Formal Analysis, Methodology, Writing – original draft, review & editing, Supervision, Funding acquisition, Project administration.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

All data and their statistical analysis are made available as part of supplementary materials and files.

Acknowledgements

This work was part of the project "AMRflows: antimicrobials and resistance from manufacturing flows to people: joined up experiments, mathematical modelling and risk analysis" supported by the Department of Biotechnology (DBT) in India (Computer No. 8981 || BT/IN/Indo-UK/AMR-Env/03/ST/2020-21 || AMRFlows) and the Natural Environment Research Council (NERC) in the UK (NE/T013222/1). Vikas Sonkar gratefully acknowledges support through the University Grants Commission (UGC) Junior Research Fellow (JRF) PhD Scholarship (grant number F. 82-1/2018 (SA-III)). The work of our AMRflows team has benefitted from numerous discussions with colleagues from the other four projects funded by the "India-UK Tackling AMR in the Environment from Antimicrobial Manufacturing Waste"

626 call through the NERC-funded Programme Coordination Team (PCT). We would also like to

627 thank colleagues from IIT Hyderabad for assisting with the field sampling (Jilla Adithya,

628 Vineeth Jupaka, Sreshtha Baidya and Nikash Naorem) and Mohammed Azharuddin for his

629 assistance with plotting the graphs in R.

References

Ahmed W, Gyawali P, Hamilton KA, et al (2021) Antibiotic Resistance and Sewage-Associated Marker Genes in Untreated Sewage and a River Characterized During Baseflow and Stormflow. 12:1–13. https://doi.org/10.3389/fmicb.2021.632850

Amarasiri M, Takezawa T, Malla B, et al (2022) Prevalence of antibiotic resistance genes in drinking and environmental water sources of the Kathmandu Valley, Nepal. Front Microbiol Volume 13-2022:

Anjum MF, Schmitt H, Börjesson S, et al (2021) The potential of using *E. coli* as an indicator for the surveillance of antimicrobial resistance (AMR) in the environment. Curr Opin Microbiol 64:152–158. https://doi.org/10.1016/j.mib.2021.09.011

Barrantes-Jiménez K, Mendoza-Guido B, Morales-Mora E, et al (2025) Influence of Seasonality and Pollution on the Presence of Antibiotic Resistance Genes and Potentially Pathogenic Bacteria in a Tropical Urban River. Antibiotics 14:798

Beltrán de Heredia I, Garbisu C, Alkorta I, et al (2023) Spatio-seasonal patterns of the impact of wastewater treatment plant effluents on antibiotic resistance in river sediments. Environ Pollut 319:120883. https://doi.org/https://doi.org/10.1016/j.envpol.2022.120883

Berendonk TU, Manaia CM, Merlin C, et al (2015) Tackling antibiotic resistance: The environmental framework. Nat Rev Microbiol 13:310–317. https://doi.org/10.1038/nrmicro3439

Borcard D, Gillet F, Legendre P (2018) Numerical Ecology with R, Second Edi. Springer, New York

Brown PC, Borowska E, Schwartz T, Horn H (2019) Impact of the particulate matter from wastewater discharge on the abundance of antibiotic resistance genes and facultative pathogenic bacteria in downstream river sediments. Sci Total Environ 649:1171–1178. https://doi.org/10.1016/j.scitotenv.2018.08.394

Caucci S, Karkman A, Cacace D, et al (2016) Seasonality of antibiotic prescriptions for outpatients and resistance genes in sewers and wastewater treatment plant outflow. FEMS Microbiol Ecol 92:fiw060. https://doi.org/10.1093/femsec/fiw060

CPCB (2005) Performance Status of Common Effluent Treatment Plants in India. East Arjun Nagar, Delhi

CPCB (2010) Final Action Plan for Improvement of Environmental Parameters in "Patancheru-Bollaram Cluster" Andhra Pradesh. Santhnagar, Hyderabad

CPCB (2021) National Inventory of Sewage Treatment Plants (2021). East Arjun Nagar, Delhi

Diwan V, Hanna N, Purohit M, et al (2018) Seasonal variations in water-quality, antibiotic residues, resistant bacteria and antibiotic resistance genes of *Escherichia coli* isolates from water and sediments of the Kshipra River in Central India. Int J Environ Res Public Health 15:1–16. https://doi.org/10.3390/ijerph15061281

Frahm E, Obst U (2003) Application of the fluorogenic probe technique (TaqMan PCR) to the detection of *Enterococcus spp.* and *Escherichia coli* in water samples. J Microbiol Methods 52:123–131. https://doi.org/10.1016/S0167-7012(02)00150-1

Gillings MR, Gaze WH, Pruden A, et al (2015) Using the class 1 integron-integrase gene as a proxy for anthropogenic pollution. ISME J 9:1269–1279. https://doi.org/10.1038/ismej.2014.226

Gothwal R, Shashidhar (2017) Occurrence of High Levels of Fluoroquinolones in Aquatic Environment due to Effluent Discharges from Bulk Drug Manufacturers. J Hazardous, Toxic, Radioact Waste 21:05016003. https://doi.org/https://doi.org/10.1061/(ASCE)HZ.2153-5515.0000346

Graham DW, Collignon P, Davies J, et al (2014) Underappreciated role of regionally poor water quality on globally increasing antibiotic resistance. Environ Sci Technol 48:11746–11747. https://doi.org/https://doi.org/10.1021/es504206x

Hanna N, Purohit M, Diwan V, et al (2020) Monitoring of Water Quality, Antibiotic Residues, and Antibiotic-Resistant *Escherichia coli* in the Kshipra River in India over a 3-Year Period. Int. J. Environ. Res. Public Health 17:7706

Hassoun-Kheir N, Stabholz Y, Kreft J-U, et al (2021) EMBRACE-WATERS statement: Recommendations for reporting of studies on antimicrobial resistance in wastewater and related aquatic environments. One Heal 13:100339

Herrig I, Fleischmann S, Regnery J, et al (2020) Prevalence and seasonal dynamics of blaCTX-M antibiotic resistance genes and fecal indicator organisms in the lower Lahn River, Germany. PLoS One 15:e0232289. https://doi.org/10.1371/journal.pone.0232289

Heß S, Berendonk TU, Kneis D (2018) Antibiotic resistant bacteria and resistance genes in the bottom sediment of a small stream and the potential impact of remobilization. FEMS Microbiol Ecol 94:fiy128. https://doi.org/10.1093/femsec/fiy128

ISO 16649-3:2015 (2015) Microbiology of the food chain — Horizontal method for the enumeration of *Escherichia coli* Part 3: Detection and most probable number technique using 5-bromo-4-chloro-3-indolyl- ß-D-glucuronide. Geneva, Switzerland

Javvadi Y, Mohan SV (2023) Understanding the distribution of antibiotic resistance genes in an urban community using wastewater-based epidemiological approach. Sci Total Environ 868:161419. https://doi.org/10.1016/j.scitotenv.2023.161419

Javvadi Y, Mohan SV (2024) Temporal dynamics and persistence of resistance genes to broad spectrum antibiotics in an urban community. npj Clean Water 7:1–12. https://doi.org/10.1038/s41545-024-00349-y

Jones ER, Van Vliet MTH, Qadir M, Bierkens MFP (2021) Country-level and gridded estimates of wastewater production, collection, treatment and reuse. Earth Syst Sci Data 13:237–254. https://doi.org/10.5194/essd-13-237-2021

Karkman A, Pärnänen K, Larsson DGJ (2019) Fecal pollution can explain antibiotic resistance gene abundances in anthropogenically impacted environments. Nat Commun 10:80. https://doi.org/10.1038/s41467-018-07992-3

Kashyap A, Nishil B, Thatikonda S (2023) Experimental and numerical elucidation of the fate and transport of antibiotics in aquatic environment: A review. Environ Monit Assess 195:942. https://doi.org/10.1007/s10661-023-11482-5

Kirschner AKT, Schachner-Groehs I, Kavka G, et al (2024) Long-term impact of basin-wide wastewater management on faecal pollution levels along the entire Danube River. Environ Sci Pollut Res 31:45697–45710. https://doi.org/10.1007/s11356-024-34190-0

Klein EY, Van Boeckel TP, Martinez EM, et al (2018) Global increase and geographic convergence in antibiotic consumption between 2000 and 2015. Proc Natl Acad Sci U S A 115:E3463–E3470. https://doi.org/10.1073/pnas.1717295115

Knapp CW, Lima L, Olivares-Rieumont S, et al (2012) Seasonal variations in antibiotic resistance gene transport in the Almendares River, Havana, Cuba. Front Microbiol 3:1–11. https://doi.org/10.3389/fmicb.2012.00396

Konda S, Asati A, Williams M, Mudiam MKR (2022) Development and evaluation of a multi-class analytical method based on solid-phase extraction combined with liquid chromatography-tandem mass spectrometry for the analysis of pharmaceuticals and personal care products in urban wastewater samples. Sep Sci Plus 5:105–119. https://doi.org/10.1002/sscp.202100041

Konda S, Nagendla NK, Williams M, Mudiam MKR (2024) Investigating pharmaceuticals and personal care products in Musi River: A exploratory study in Hyderabad, India. J Hazard Mater Adv 14:100420. https://doi.org/10.1016/j.hazadv.2024.100420

Kumar M, Sulfikar, Chaminda T, et al (2020) Prevalence of antibiotic resistance in the tropical rivers of Sri Lanka and India. Environ Res 188:109765. https://doi.org/10.1016/j.envres.2020.109765

Larsson DGJ, de Pedro C, Paxeus N (2007) Effluent from drug manufactures contains extremely high levels of pharmaceuticals. J Hazard Mater 148:751–755. https://doi.org/10.1016/j.jhazmat.2007.07.008

Lee J, Ju F, Maile-Moskowitz A, et al (2021) Unraveling the riverine antibiotic resistome: The downstream fate of anthropogenic inputs. Water Res 197:117050. https://doi.org/10.1016/j.watres.2021.117050

Li L, Nesme J, Quintela-Baluja M, et al (2021) Extended-Spectrum β-Lactamase and Carbapenemase Genes are Substantially and Sequentially Reduced during Conveyance and Treatment of Urban Sewage. Environ Sci Technol 55:5939–5949. https://doi.org/10.1021/acs.est.0c08548

Lübbert C, Baars C, Dayakar A, et al (2017) Environmental pollution with antimicrobial agents from bulk drug manufacturing industries in Hyderabad, South India, is associated with dissemination of extended-spectrum beta-lactamase and carbapenemase-producing pathogens. Infection 45:479–491. https://doi.org/10.1007/s15010-017-1007-2

Marathe NP, Pal C, Gaikwad SS, et al (2017) Untreated urban waste contaminates Indian river sediments with resistance genes to last resort antibiotics. Water Res 124:388–397. https://doi.org/10.1016/j.watres.2017.07.060

Martínez JL (2008) Antibiotics and Antibiotic Resistance Genes in Natural Environments. Science (80-) 321:365–367. https://doi.org/DOI: 10.1126/science.1159483

Muurinen J, Muziasari WI, Hultman J, et al (2022) Antibiotic Resistomes and Microbiomes in the Surface Water along the Code River in Indonesia Reflect Drainage Basin Anthropogenic Activities. Environ Sci Technol 56:14994–15006. https://doi.org/10.1021/acs.est.2c01570

Naghavi M, Vollset SE, Ikuta KS, et al (2024) Global burden of bacterial antimicrobial resistance 1990-2021: a systematic analysis with forecasts to 2050. Lancet 404:1199–1226. https://doi.org/10.1016/S0140-6736(24)01867-1

Narshimha C, Reddy UVB, Praveen K, Ramesh S (2020) Geochemistry data of the Nadergul granodiorite of the late Archean Hyderabad granite batholith, part of Eastern Dharwar craton, India; implications for composition of the lower continental crust. Data Br 31:105768. https://doi.org/https://doi.org/10.1016/j.dib.2020.105768

Ott A, O'Donnell G, Tran NH, et al (2021) Developing Surrogate Markers for Predicting Antibiotic Resistance "Hot Spots" in Rivers Where Limited Data Are Available. Environ Sci Technol 55:7466–7478. https://doi.org/10.1021/acs.est.1c00939

Pallares-Vega R, Blaak H, van der Plaats R, et al (2019) Determinants of presence and removal of antibiotic resistance genes during WWTP treatment: A cross-sectional study. Water Res 161:319–328. https://doi.org/10.1016/j.watres.2019.05.100

Prasad DK, Shukla R, Ahammad SZ (2024) Antibiotic resistance in the Ganga River: Investigation of antibiotic resistant bacteria and antibiotic resistance genes, and public health risk assessment. J Environ Chem Eng 12:114931. https://doi.org/https://doi.org/10.1016/j.jece.2024.114931

Ram B, Kumar M (2020) Correlation appraisal of antibiotic resistance with fecal, metal and microplastic contamination in a tropical Indian river, lakes and sewage. npj Clean Water 3:1–12. https://doi.org/10.1038/s41545-020-0050-1

Rathinavelu S, Uluseker C, Sonkar V, et al (2024) Mapping the scarcity of data on antibiotics in natural and engineered water environments across India. Front Antibiot 3:1–17. https://doi.org/10.3389/frabi.2024.1337261

Ricotta C, Podani J (2017) On some properties of the Bray-Curtis dissimilarity and their ecological meaning. Ecol Complex 31:201–205. https://doi.org/10.1016/j.ecocom.2017.07.003

Schachner-Groehs I, Koller M, Leopold M, et al (2024) Linking antibiotic resistance gene patterns with advanced faecal pollution assessment and environmental key parameters along 2300 km of the Danube River. Water Res 252:121244. https://doi.org/10.1016/j.watres.2024.121244

Silkie SS, Tolcher MP, Nelson KL (2008) Reagent decontamination to eliminate false-positives in *Escherichia coli* qPCR. J Microbiol Methods 72:275–282. https://doi.org/10.1016/j.mimet.2007.12.011

Sonkar V, Kashyap A, Pallares-Vega R, et al (2025a) Removal of antimicrobial resistance determinants from wastewater: Role of capacity overloading and treatment technology in wastewater treatment plants. J Environ Manage 395:127897. https://doi.org/https://doi.org/10.1016/j.jenvman.2025.127897

Sonkar V, Sasidharan SS, Jambu SC, et al (2025b) A multi-endpoint approach to ecotoxicological assessment of wastewater polluted rivers using zebrafish. Environ Res 282:121996. https://doi.org/10.1016/j.envres.2025.121996

Sonkar V, Venu V, Nishil B, Thatikonda S (2024) Review on antibiotic pollution dynamics: insights to occurrence, environmental behaviour, ecotoxicity, and management strategies. Environ Sci Pollut Res 31:51164–51196. https://doi.org/10.1007/s11356-024-34567-1

The World Bank (2022) Urban population growth (annual %) - India. https://data.worldbank.org/indicator/SP.URB.GROW?locations=IN. Accessed 24 May

2023

UNEP (2023) Bracing for Superbugs: Strengthening environmental action in the One Health response to antimicrobial resistance. Geneva

Wang H, Hou L, Liu Y, et al (2021) Horizontal and vertical gene transfer drive sediment antibiotic resistome in an urban lagoon system. J Environ Sci 102:11–23. https://doi.org/https://doi.org/10.1016/j.jes.2020.09.004

Yang H, Li X, Li X, et al (2024) The fate of antibiotic resistance genes and their correlation with microbial communities and wastewater quality/parameters in a wastewater treatment plant under different seasons. J Water Process Eng 60:105156. https://doi.org/10.1016/j.jwpe.2024.105156

AMRflows consortium

[1]Panagiota Adamou, [2]Shubham Anurag, [2]Sounak Banerjee, [1]Ewelina Bien, [3]Sangeetha Chandrakalabai Jambu, [1]David W Graham, [1]Colin Harwood, [4]Siu Fung Stanley Ho, [5]Rupert Hough, [2]Chaitanya Janivara Chandregowda, [1]Kelly Jobling, [3]Arun Kashyap, [4,6]Jan-Ulrich Kreft, [2]Soumendra Nath Kuiry, [7]Joshua Larsen, [2]Thara Methale Velakkath Paramba, [8]Ankit Modi, [8]Pranab Kumar Mohapatra, [2]Anantha Barathi Muthukrishnan, [2]Indumathi Manivannan Nambi, [1]Julián Osvaldo Ovis Sánchez, [1]Rebeca Pallarés Vega, [2]Ravikrishna Raghunathan, [2,4,6]Sasikaladevi Rathinavelu, [3]Aravind Kumar Rengan, [8]Wahidullah Hakim Safi, [2]Shubhangi Sanjeev, [2,3]Sai Sugitha Sasidharan, [2]Vandit Kumar Shah, [3]Vikas Sonkar, [3]Rupali Srivastava, [3]Shashidhar Thatikonda, [5]Mads Troldborg, [4,6]Cansu Uluşeker, [9]Rama Vaidyanathan, [4]Willem van Schaik, [4,6]Anjali Vasudevan, [2]Kiruthika Eswari Velmaiel, [2]Arathy Viswanathan.

[1]School of Engineering, Newcastle University, Newcastle upon Tyne, NE1 7RU, UK

[2]Environmental & Water Resources Engineering Division, Department of Civil Engineering, Indian Institute of Technology Madras, Chennai, 600036, Tamil Nadu, India

[3]Environmental Microbiology Laboratory, Department of Civil Engineering (Environmental Engineering), Indian Institute of Technology Hyderabad, Kandi, Sangareddy, 502284, Telangana, India

[4]Institute of Microbiology and Infection, University of Birmingham, Edgbaston, Birmingham, B15 2TT, UK

[5]The James Hutton Institute, Craigiebuckler, AB15 8QH, Aberdeen, UK

[6]School of Biosciences, University of Birmingham, Edgbaston, Birmingham, B15 2TT, UK

[7]School of Geography, Earth and Environmental Sciences, University of Birmingham,

Edgbaston, Birmingham, B15 2TT, UK

[8]Department of Civil Engineering, Indian Institute of Technology Gandhinagar, Gandhinagar,

382355, Gujarat, India

[9]Dr. M.G.R. Educational and Research Institute, E.V.R. Periyar Salai, Adayalampattu, Chennai,

600 095, Tamil Nadu, India