Protocol for assessing DNA damage levels *in vivo* in rodent testicular germ cells in the Alkaline Comet Assay

Ann-Karin Olsen[1,2,6,7*]**, Xiaoxiong Ma**[2,3]**, Congying Zheng**[4]**, Hildegunn Dahl**[1]**, Yvette Dirven**[1]**, Anne-Mette Zenner Boisen**[5]**, Anoop Sharma**[5]**, Dag. M. Eide**[1]** and Gunnar Brunborg**[1,4]

[1]Department of Chemical Toxicology, Division of Climate and Environmental Health, Norwegian Institute of Public Health, NO-0213 Oslo, Norway
[2]Section of Health Effects, Department of Environmental and Health, The Climate and Environmental Research Institute NILU, NO-2027 Kjeller, Norway
[3]SpringBiolabs, Oslo Cancer Cluster, Radiumhospitalet, Oslo, 0379, Norway
[4]NorGenoTech, Oslo Cancer Cluster, Radiumhospitalet, Oslo, 0379, Norway
[5]Division for Risk Assessment and Nutrition, Group for Chemical Risk Assessment and GMO, National Food Institute, Technical University of Denmark, 2800 Lyngby, Denmark
[6]Technical contact
[7]Lead contact
*Correspondence: anka@nilu.no

Summary

Few protocols are available for retrieving male germ cell-specific information on DNA damage dynamics or assessing the potential harmful effects of environmental contaminants, drugs, or lifestyle factors related to genotoxicity. Here, we present a protocol for evaluating testicular germ cell genotoxicity using a modified version of the alkaline comet assay in rodent testicular germ cells. The protocol includes experimental design, preparation of testicular cell suspensions, comet analysis, germ cell-specific scoring, and data curation methods to collect information on testicular male germ cells, specifically spermatids and spermatocytes.

For complete details on the use of this protocol please refer to Olsen et al., in preparation

Highlights

- DNA damage levels can be specifically measured in testicular germ cells using the developed revised version of the alkaline comet assay

- 1C spermatids as well as 4C primary spermatocytes can be assessed

- Both manual- and modeling-based approaches were developed that facilitate user-friendly protocols to select 1C spermatid comets.

- This protocol expands the limited methodologies available to study germ cell DNA damage dynamics

42 Graphical abstract
43

44
45
46
47

Before you begin

This protocol describes how to measure germ cell-specific DNA damage levels in *in vivo* rodent studies. The protocol may be used for both mice and rats. Rats are used herein as an example in the description of the protocol.

Institutional permissions

All experiments and procedures described herein were performed according to a protocol approved by the Danish Animal Testing Authority, application number: 2021-15-0201-01102. It is required that you acquire the necessary permissions from the relevant institutions for your studies.

Preparation 1 (optional, only if relevant): Treatment of rodents.

If not relevant, continue to Preparation step 2.

Timing: The timing depends on the design of the experiment. Typically, exposure is for 3-28 days plus one day for euthanasia and tissue harvest.

1) Follow recommended protocols and guidelines for animal experiments for treatment or exposure of animals.
2) Arbitrarily designate animals to each exposure group. Tag animals individually to enable blinding of research personnel during all procedures. Then use a randomization tool like Excel to assign animals to groups.
3) Prepare stock solutions of your test compound/agent.
4) Prepare serial dilutions of your compound stock of the concentrations to be tested.
5) Weigh and expose the animals using the relevant route, in a standardized manner concerning exposure technique and timing between animals. The timing should be followed accordingly for euthanasia and tissue harvest

Preparation 2: Preparation of comet stock solutions

Timing: 1-2 days

Use your comet protocol of preference and prepare the corresponding stock solutions. Alternatively, use the comet protocol by Gutzkow et al., 2013 (1) or other standard protocols (2-4), recently summarized by Collins and co-authors (5).

1) Prepare stock comet lysis solution. For fresh biomaterials Dimethyl sulfoxide (DMSO) can be omitted from the lysis solution.
2) Prepare the comet electrophoresis stock solution, if such a stock solution is part of your comet protocol

Note: The comet platform used in this project (1) provides a medium-high throughput variant of alkaline comet assay with testicular germ cells. Efficient workflow is highly encouraged to minimize endogenous DNA damage and repair. We further recommend preparing all required equipment and solutions before starting. All solutions should be freshly made and kept cool, except the agarose solution that is maintained at 37 °C in a closed container to prevent evaporation. We suggest using chilled metal plates (custom-made by NorGenoTech) placed on ice. It allows easy cold handling during tissue and cell processing and efficient and rapid solidification of comet gels during casting of cells.

Preparation 3: Preparation of assay solutions

Timing: At the day of rodent termination; 2h

1) Prepare working solutions:
   a) Lysis solution with DMSO. Cool in refrigerator.
   b) Electrophoresis solution. Place on ice.

c) Merchants/Fetal bovine serum/DMSO (M/F/D)-solution. Place on ice.
Note: Alternative solutions to Merchant solutions can be used as long as they are isotonic and
contain 10-20% EDTA.
2) Prepare containers with ice-cold PBS to immediately chill the testes after euthanasia and tissue
excision.
3) Set a heating block to 37 °C.
4) Melt 0.75 % low melting point agarose solution carefully in the microwave oven. Swirl gently to
see that all agarose is melted and homogenously distributed in the tube, i.e. that it is completely
clear. Prepare low-melt agarose for each experiment in one batch. Keep it in a container with a
securely attached lid to avoid evaporation that could lead to changes in agarose concentrations. If
needed, aliquot the agarose solution into suitable smaller containers
a) Place the melted agarose in the preheated heating block to adjust to 37 °C.
5) Prepare polystyrene box with ice to work in, alternatively chill metal metal plates to 4 °C
6) Mark tubes for cell suspensions, dilutions and Gelbond®-films/slides
7) Cut/prepare gauze and nylon mesh filter pieces to filter testicular cell suspensions
8) Prepare the genomic tips or cut 1 mL tips to obtain a wide bore to avoid spurious induction of DNA
damage due to mechanical stress.
9) Place Petri dishes on ice or cold metal plate
10) Prepare Bürker chambers and microscope for counting cells before casting gels.
122
123

124 # Key resources table
125
126

REAGENT or RESOURCE	SOURCE	IDENTIFIER, Cat #
Chemicals, Peptides, and Recombinant Proteins		
Fetal bovine serum, heat inactivated	Thermo Fisher Scientific	16140071
Dimethyl sulfoxide (DMSO)	Sigma Aldrich	5895691000
Experimental Models: Organisms/Strains		
Standard laboratory strains of healthy young sexually mature male rodents, Here: Male Wistar Han IGS rats. 9-11 weeks of age at arrival.	Scanbur (Karlslunde, Denmark)	
Software and Algorithms		
Comet assay IV software	Perceptive Instruments Ltd., Suffolk, UK	N/A
R-studio	https://posit.co/products/open-source/rstudio/	N/A
BioRender	https://www.biorender.com	N/A
Other		
Cotton gauze	VWR	BRAU9030409W
Nylon Membrane Filter, 100.0 µm Pore Size (Alternatives exist: Can be purchased by the meter)	Millipore	NY1H04700
Bacteria spreader, triangle shaped	VWR	612-2688
Petri dish, 100 x 15 mm	Merck	P5731
Surgical scissors	VWR	140981
Scalpel blade, #10	VWR	76549-476
Fine-point forceps, curved	VWR	470005-440
Fine-point forceps, straight	VWR	470007-560
Dissecting scissors, curved	VWR	76457-366
Microcentrifuge tubes, 1.5 mL	Avantor by VWR	211-2610
Wide bore pipette tips, 1000 uL	Merck	Z709999-960EA
VWR® VisiScope® 254, Upright Microscopes	VWR	630-1929
Bürker counting chambers, disposable, C-Chip	NanoEnTek, Avantor	DHC-B01

128 # Materials and equipment (optional)

131 **Merchant's solution**

Reagent	Final concentration	Amount
NaCl	0.14 M	16.4 g
KH_2PO_4	1.47 mM	400.2 mg
KCl	2.7 mM	402.6 mg
Na_2HPO_4	8.1 mM	2.3 g
Ethylenediaminetetraacetic acid disodium salt dihydrate; EDTA-Na_2	10 mM	7.4 g
ddH_2O	n/a	1.5 L first to dissolve the salts (add NaOH dropwise to dissolve the salts, see footnote). Adjust to 2 L
Total	**n/a**	**2 L**

Adjust with NaOH dropwise until the components dissolve. Take care that the pH does not exceed 7.4.
Adjust to 2L with distilled water. Continue to add NaOH dropwise until pH=7.4. Sterilize the solution by
autoclaving or sterile filtering.
[Note on storage conditions: temperature: 4°C, maximum storage time: 2 years]
Note: The EDTA-concentration can alternatively be doubled, to 20 mM, so that the working solution has a final
EDTA-concentration of 10 mM.

Merchants/Fetal bovine serum/DMSO (M/F/D)-solution
(for freezing fresh cell suspensions, or generating single-cell suspensions from frozen tissue)

Reagent	Final concentration	Amount
Merchant's solution	50%	5 mL
Fetal bovine serum, inactivated	40%	4 mL
DMSO	10%	1 mL
Total	**n/a**	**10 mL**

141 Prepared minutes before cell isolation.
142 [Note on storage conditions: temperature: 4°C, maximum storage time: 3 h]

144 **Solutions**
145 • **Tris-EDTA (TE)-buffer**: (1 mM Na_2EDTA, 10 mM Tris-HCl), pH 7.5

Step-by-step method details

1. Overview of work-flow and options

Processing of testis tissue to generate single-cell suspensions (scs) includes two alternative approaches: a) processing of fresh testis tissue or b) snap-freeze testis tissue pieces for later generation of scs. The scs are subject to comet analyses, following recommended parameters, and scored as described below. Selection of the testicular germ cells containing 1C DNA (1C spermatids) can be done by two approaches based on comet total DNA content (Total Intensity; Tot_Int); visually, i.e. by inspection of graphs of comet data produced by a spreadsheet (The provided excel template or the R template) or by modeling using forced adaptation to a 3-population distribution (use the provided R template). Additionally, comets of primary spermatocytes (PS) can be enriched by selective scoring of PS-comets according to parameters described.

Figure 1: Overview of workflow and options.
(Created in BioRender. BioRender.com/p97s689)

CRITICAL: It is imperative that all steps are performed rapidly and tissues and samples are kept cold
(on ice) to control spurious background levels or prevent removal of DNA lesions by repair that occurs
at ambient.

Note: The protocol can be used with either fresh or frozen testis tissue.

CRITICAL: Frozen testis tissue must be processed instantly after thawing. A frozen testis piece is
often smaller than a fresh testis piece due to the requirement of rapid snap-freeze to preserve the
integrity of both cells and their DNA. Consequently, the dilution with M/D/F solution should be
accordingly adjusted.
176

177 2. Animal exposure (optional)

178
179 **Timing: [Typically 3 to 28 days]**
180

181 The aim is to expose animals to understand potential genotoxic effects.

182
1. Acclimatize the animals for at least one week, see FELASA-AALAS recommendations.
2. Randomly allocate the animals to experimental groups, tag animals and enable blinding of
research personnel during all procedures.
3. House animals according to accepted guidelines.
a. Observe the animals every day according to accepted guidelines.
4. Expose animals by suitable route.

189
190 **Note**: The conditions for exposure, euthanasia and tissue harvest should be standardized and conducted
191 following a carefully planned time schedule with a set interval between animals.

192
193 **Note**: According to OECD Test Guideline TG 489 (6) exposure is recommended on three consecutive
194 days, 24 h apart, and euthanasia should be conducted 3 h after the last exposure depending on the
195 exposure agent, pilot studies and expert judgement.

196
197 3. Animal termination and tissue harvest

198
199 **Timing: [5 min]**
200

201 The aims of this step are to terminate animals according to standard recommendations for animal
202 experiments, and harvest testis tissue by rapidly dissecting and chilling the tissues to prevent potential
203 elimination of DNA lesions by DNA repair or serendipitous induction of non-relevant DNA damage.

204
1) If Step 2 Animal exposure is non-relevant, allow the animals to acclimatize.
2) Terminate the animals using a relevant method according to approved guidelines.
a) Weigh the whole animal and terminate, one animal at a time in a timely fashion (a set # min
apart between each animal, in concert with potential exposure regime) to allow time to process
tissues and preparing single-cell suspensions as described below. Perform in a randomized and
blinded sequence.
3) Harvest testicular tissue immediately after euthanasia.

a) Open the rodent by cutting the abdominal wall along the midline using scissors. Pull the testes
from the scrotum to the abdomen. Collect the intact testes rapidly.
b) Chill the intact testes rapidly by submerging them in ice-cold Merchant's solution or
phosphate-buffered saline (PBS)
c) **Recommended**: weigh the testes rapidly (see below)
i) Return the tissue to the Merchant's solution/PBS to keep cold.
d) Immediately proceed to the next step

Note: Animal termination and testis tissue harvest should be conducted according to a timely schedule
with a set interval between animals to standardize the conditions for each animal, in a randomized and
blinded manner (exposure groups must be concealed for the research personnel performing the
procedures).

Note: According to OECD Test Guideline TG 489 (6) it is recommended to euthanize the animals 3
hours after the last exposure pending on the exposure agent, pilot studies and expert judgment.

Optional: Weigh the intact testes by blotting the chilled tissue on paper to remove excess liquid before
weighing the tissue.

CRITICAL: Work rapidly and keep everything cold.

4. Testis dissection

Timing: [1-2 min]

The aims are to dissect the testis and cut the testis tissue into small pieces for immediate preparation of
testicular single-cell suspensions (**Video 1**). Alternatively, small testis pieces are snap-frozen for later
processing (**Video 1**).

1) Place the testis in a cold Petri dish (on ice, or a metal plate)
2) Remove the capsule and the main blood vessel
3) Cut the testis tissue into small pieces
a) Use forceps to hold the testis and sharp small scissors to cut the tissue into pieces
b) Cut testis tissue into pieces of approximately 3-4 mm^3
c) Transfer the tissue pieces to a new cold Petri dish containing 1 mL ice-cold M/F/D-solution

Optional: To preserve testis tissue for later analyses, cut smaller pieces (2-3 mm^3) of testis and transfer
the pieces rapidly to a prelabelled tube and snap-freeze in liquid N2.
Optional: If the testis tissue has been snap-frozen, transfer and thaw the testis piece in a Petri dish
containing 1 mL M/F/D-solution (**Video 2**).

CRITICAL: Work rapidly and keep everything cold.

CRITICAL: Watch the instructional video before starting (**Video 1**).

5. Preparation of single-cell suspensions

259 6. **Timing: [1-3 min]**

260 The aim is to prepare testicular single-cell suspensions (scs) by mechanical release of tubular testicular
261 cells followed by filtering (**Video 3**).

1. Use forceps to fix/hold the testis tissue piece immersed in the ice-cold M/F/D-solution
2. Use a triangular bacterial spreader and gently squeeze and stroke the testicular tubules 2-4 times to release testicular cells. You may observe that the solution becoming opaque due to the released tubular cells
3. Homogenize the cell suspension by pipetting 2 times using a 1 mL pipette fitted with a wide-bore 1 mL tip. Keep the tip opening towards the bottom of the Petri dish to enhance the dislodging of testicular cells.
4. Filter the scs rapidly through a cotton gauze placed on top of a 100 µm nylon membrane filter into a pre-labeled tube (1,5 mL or 0,5 mL tube) to remove larger tissue pieces and reduce the presence of spermatozoa
5. Dilute 50 µl cells in 950 µl M/F/D-solution and rapidly count the testicular cells using a manual Bürker chamber and a phase-contrast microscope.
   a. Dilute the single cell suspension (scs) with M/F/D-solution to a concentration suitable for scoring of comets.
   b. Place the scs on ice and proceed to the next step as soon as possible.

Note: If wide-bore tips are not available conventional 1 mL tips can be cut at the end using scissors.

Note: Keep air bubbles generated during pipetting to a minimum to prevent inducing spurious DNA
lesions.

Note: Counting can be performed with cells without staining and thus rapidly processed. Testicular cell
suspensions can be challenging to count due to their small and highly variable sizes. Count cells of all
sizes. The cells with curved nuclei and tails are testicular spermatozoa. We use a concentration of 0,6 –
1,2 x 10^6 cells/mL that is further diluted 10 x during casting of gels. Do not count the cells using an
automated cell counter.

Note: Counting can be omitted by casting more than one dilution of each sample and selecting the
dilution most suitable for scoring by rapidly assessing the gels before starting to score.

CRITICAL: Watch the instructional video before processing the testis tissue (**Video 3**).

Optional: When testicular single-cell suspensions are prepared from frozen tissue, the testis tissue
piece is rapidly thawed in 1 mL cold M/F/D-solution in a Petri dish on ice before proceeding with the
same protocol as for fresh tissue (**Video 2**).

6. Comet assay

Timing: [1 day]

The aim is to cast the single-cell suspensions in gels as rapid as possible for subsequent execution of
the comet assay according to your comet protocol, following recent guidelines (5, 7, 8) and
recommendations listed below. Details and instructions regarding the comet method can be found in
published articles (1, 3, 4, 9) (5).

We recommend using the comet protocol established in your laboratory, or your comet protocol of
preference (10). We do however recommend following the instructions below noted as Critical.

We used a medium-throughput protocol with Gelbond® support films for agarose (1). Cells were
casted in low-melt agarose directly onto the Gelbond® film as 4 ul gels and 8 gels per sample) before
submerging in lysis solution. Specimens can be left in lysis solution for several days without changes in
the % tail intensity (TI) (11).

309
Some critical parameters:

CRITICAL: The time the cell suspensions are kept on ice before casting in agarose and lysis must be
kept as short as possible since single-strand breaks are generally rapidly repaired.

Note: Avoid unnecessary pipetting and the generation of air bubbles during pipetting.

Note: We recommend including somatic (liver) in addition to the testicular cell samples from each
animal on the same slide or Gelbond®-film. This may facilitate comparisons of Tot_Int mean levels in
each cell type and can be used to verify the identification of the 1C-population of germ cells in later
stages of the protocol, assuming that the induced DNA damage levels of both cell types are somewhat
similar. This is especially relevant for reprotoxic stressors that may lead to 1C-spermatid toxicity.

CRITICAL: **Use the same final concentration of low-melt agarose in all experiments**, and the
same batch of agarose within each experiment. We propose a final concentration of 0.75% LMP.
Complying with these instructions as a standard will reduce the variability within experiments and
between experiments and laboratories.

CRITICAL: The time in lysis solution should be at least 16 h and the lysis solution should contain
DMSO.

CRITICAL: **Keep equal and constant voltage in all experiments.** This will reduce the variability
between experiments and laboratories and facilitate comparisons. First measure the slide stage width
using a simple ruler. Second, fill the electrophoresis tank with the volume of the electrophoresis buffer
of your protocol. Turn on the power supply. Measure the voltage across the stage with a voltmeter by
submerging the electrodes on each side of the stage. Register the voltage. Calculate the V/cm by
dividing the V by the stage width (in cm). Change the voltage on the power supply to obtain the
voltage/cm wanted (preferably 0.86 V/cm; range 0.84-0.88 V/cm) and refine by adding or removing
electrophoresis buffer until the preferred V/cm is obtained. Take care that there is sufficient
electrophoresis buffer above the slides/gelbond®-films (~5mm). Make sure to use a power supply that
can maintain the voltage throughout electrophoresis. **When all parameters are set, run the**
electrophoresis for exactly 25 min at 8 °C and 0.84-0.88 V/cm across the stage.

Pause point: At the stage of lysis there is an opportunity to stop since the specimens can be left for
longer times (11).

Note + pause point: Slides and Gelbond® films can be air dried and stored until staining the DNA (we
use SYBR^TM Gold) and scoring of comets.

Note + pause point: Alternatively, if scoring has to be finalized within a limited period (depending on
DNA stain used, relevant for SYBR^TM Gold staining), all samples can be stained before drying
followed by re-wetting the Slides/Gelbond® films on the day of scoring.

7. Comet visualization and analysis (scoring)

344
345 **Timing, per sample: Depending on testicular cell type assessed/sample:**

Scoring: scoring of > 200 round nucleoid comets ~1h
Optional scoring step: scoring of >= 150 primary spermatocytes (4C) ~40 min

The main aim of this protocol is to obtain testicular germ cell comet data from specific stages of spermatogenesis. The scoring of testicular cells follows simple principles described below and is hence different from the conventional scoring of somatic cells. The testicular cell suspensions contain a mix of many cell types of both germ and somatic origin with variable nucleoid shape and DNA content (**Figure 2**).

For each animal, depending on whether data is warranted from both the 1C spermatids only, or also the 4C primary spermatocytes (PS), an optional second scoring for the 4C PS can be done. During the first scoring the 1C spermatids will be selected (**Videos 4-7**). During the optional second scoring 4C PS are enriched. To select 1C spermatids all round-shaped nucleoids are scored, whereas to enrich for 4C PS the particularly large round nucleoids with high total intensities are scored. (The large nucleoids, **Videos 4-7**)

Note: Comet images for instructing the scoring and a set of training comet images are provided as **supplementary files in folder "Scoring"**.

Note: Ensure blinding and random sequence of scoring samples.

Note: Your scoring system may use a different term for the total DNA content of each comet, please identify the denotation and use the appropriate term in steps below replacing Total intensity (Tot_Int).

    1.   Scoring to select 1C spermatids
        a. On each gel of each sample, score "all round-shaped nucleoids" (**Figure 2, Videos 4-7**)
            i. Do not score crescent-like comets (elongating/elongated spermatids) or spermatozoa (**Figure 2, see comets NOT scored in videos 4-7**).
           ii. Record the number of hedgehogs in each sample
        b. Score systematically across the gels and make sure to avoid the gel edges
        c. Perform the identical procedure on each gel
        d. Note the Tot_Int of the large nucleoids to instruct for the optional 4C PS scoring
        e. In total, the sum of comets from all gels/sample, score as many comets as considered appropriate to obtain a clear Tot_Int distribution of comets to select the 1C spermatids (see next steps)
    2. **Optional**: Scoring to enrich 4C Primary Spermatocytes (PS)
        a. On each gel of each sample, score the comets with particularly large nucleoids (**Figure 3, See large comets in Videos 4-7**) and verify the Tot_Int to be in the range observed for large comets during scoring of all the 1C nucleoids in the previous step
        b. Score systematically across the gel. Make sure to avoid the gel edges
        c. Perform the identical procedure on each gel
        d. The sum of comets from all gels/sample: score as many nucleoids as considered appropriate, normally >= 150 per sample is considered sufficient

391
392
393

Untreated   Treated, low damage (~20 % TI)   Treated, high damage (~60 % TI)

394
Figure 2. Scoring of "all round shaped nucleoids". The images represent untreated samples (2 to the
left), treated samples with low damage (~20 % TI; middle images) and treated samples with high
damage (~60 %TI; images to the right), with two images for each condition. The arrows indicate the
round-shaped nucleoid to be scored, whereas the nucleoids not scored are spermatozoa,
elongating/elongating spermatids, multiplets, hedgehogs or nucleoids at the edge of the image, not to
be scored. (For training of selection of which nucleoids to score see Supplemetary files in folder
"scoring").
402
403
404 **Note:** Quality assessment of the staining protocol of your laboratory can be verified by evaluating the
405 total intensities of the samples.

Figure 3. Scoring of 4C primary spermatocytes. The images represent untreated samples (top image) and samples with moderate levels of DNA damage (middle and bottom images). One typical comet score is visualized in each image, along with the high Tot_Int used to confirm the identity as a 4C cell and the % TI level recorded shown in the right box.

CRITICAL: The number of "all round nucleoid" comets from all gels/sample to score: score as many comets as considered appropriate to obtain a clear Tot_Int distribution of comets to select the 1C round spermatids (see next steps). The # of comets to score depends on each experiment. Relevant parameters include testicular germ cell toxicity and genotoxicity potency. Due to the selection of germ cell specific comets sufficient numbers of comets have to be scored for the number of selected germ cell specific comets to be acceptable.

Note: Visual instructions to determine which nucleoids to score are provided as annexes (**Annex 1, Annex 2**) containing images for training.

Note: If you plan to use the R-script for selection of 1C round spermatids, please follow the instructions in the R-script on naming of files and organization of folders.

8. Selection of testicular germ cell comets

Comets of 1C round spermatids can be selected using two different approaches, both based on distribution plots of the Tot_Int of the comets of each animal. The separating parameter is the Tot_Int, i.e. the amount of DNA in each comet. 1C comets are specific for germ cells since they do not exist among somatic cells. The 1C spermatid population is identified by either 1) Visual approach or 2) Modeling approach. The Visual approach can be done two ways, either by using the excel template or the R-script inspecting the animals one-by-one, whereas the Modeling approach processes all animals simultaneously.

Note: The **optional** selection of the 4C PS comets is based on the enrichment (increasing the relative numbers of PS in the selection of comets scored) of this cell type based on two defining nucleoid characteristics; their particularly large nucleoid size plus high Tot_Int (i.e. high DNA content). Protocol details for PS comets are described below the instructions for selection of the 1C spermatids.

Note: The Tot_Int-based identification of the 1C-population of comets can be verified by plotting the distributions of the Tot-Int of the comets of other somatic tissues, such as liver or blood, from the same animal/experiment, where the Tot_Int of the somatic 2C-population is easily identified.

CRITICAL: Your scoring system may use a different term for the total DNA content of each comet, please identify the denotation and replace that term with "Total intensity (Tot_Int)" for steps below. This is important when using the excel-template and R-script.

Optional: The Tot_Int-based identification of the 4C PS can be verified by plotting the distributions of the Tot_Int of the comets of other somatic tissues, such as liver or blood, from the same animal/experiment, where the Tot_Int of the somatic 4C-population is easily identified.

Identification of 1C spermatids

457 **Note**: use the provided excel-template and R-script.
458
1) Excel template
a) After importing the scored comets according to the instructions in the excel template the
distributions of the Tot_Int of all the scored comets of each animal are plotted as distribution
histograms with total intensities on the x-axis, %TI on the left y-axis and frequencies of comets
on the right y-axis (presented as a line).
b) Distribution histograms of the Tot_Int of all the round cell comets (with different bin numbers,
to facilitate easier identification of Tot-Int threshold) are automatically plotted (**Figure 4**).
c) Distribution histograms of the Tot_Int of all the scored comets of the other cell types (liver, 4C
PS) are automatically plotted.
d) Identify the threshold between the 1C comet population and the >1C other comets of the testis
using the frequency line distinguishing the 1C and the 2C populations (**Figure 4**, see also **Figure**
1)
i)   **Optional**: The Tot_Int threshold to identify the 1C population can be verified by comparing
472   with the somatic Tot_Int distribution populations (2C and 4C; **Figure 5**)
e) All comets with Tot_Int below the threshold are automatically selected as the 1C-population
(**Figure 6**)
f) Retrieve the median % TI values for the comets for each cell type to your software for statistical
analyses of preference
g) Repeat steps a) - f) for each animal
h) Calculate the mean +/- SD % TI of each experimental group
i) Perform statistical analyses as deemed appropriate

485
486 **Figure 4** Distribution and frequency of all round comets

DNA content (Total intensity), 75 bins

488
489 **Figure 5.** Distributions and frequencies of all round comets (grey dots and grey line), liver comets (red
490 line) and PS comets (green line)
491

DNA content (Total intensity)

492
493 **Figure 6.** Dot plot showing all comets as dots, a red stippled line showing the threshold line between 1C
494 comets (pink) and >1 C comets (grey)
495
Note: For R, first download the folders on your local desktop. Next open R-script_template.Rmd located
in the folder "R-template" where you will find example data for 5 animals and the folder structures
necessary for the R-script. Follow the description in the R-script to generate the data for these 5 animals,
with predefined cut-off and Tot_Int threshold values. Next, train by setting your cut-off and Tot_Int
threshold values on the 2 animals in the R_training data folder. To process your own data, create a folder
structure identical to the R_template.
502
503 2) R_template
504 There are two approaches to select the 1C population, manual or modelling
505 Manual: follow steps a-n, r
506 Modelling: follow steps a-h, o-r
507
508 **Note**: As an alternative to using the excel template steps a-n and r can be followed using the R-
509 script_template.Rmd.
510
a) Open the R-script in RStudio and follow the instructions given for file naming and folder
structure
b) Set up your working directory (play button #1)

c) Run R-packages (play button #2)
d) Import raw data .xls-files for all cell types of each animal in each experiment (play button #3).
The R-script collapses the comets from all gels of each cell type for each animal.
e) Create distribution plots for each animal (play button #4)
f) Export the raw data (File name:"All_comets_raw"; median %TI, # comets) for all cell types of
each animal to a txt-file (play button #5)
g) Inspect the order of your folder structure for each animal. Follow this order for the subsequent
operations (play button #6)
h) Removal of non-relevant datapoints
i) Inspect the testis (all round comets) distribution plots generated in e) to remove the non-
524   relevant datapoints (debris below 1C and >4C population) based on Tot_Int for each animal
525   ii) Insert (or replace) the debris below 1C and >4C Tot_Int cut-off values of your animals in the
526   correct order to remove the non-relevant datapoints >4C-population (play button #7)
Manual selection of 1C population
i) Generate distribution plots all round comets, liver and PS (play button #8; **Figure 8**)
j) Visually inspect the distribution plots to determine the 1C threshold level to discern the 1C
population from the >1C populations for each animal
k) Inspect the order of your folder structure for each animal. Follow this order for the subsequent
operations (play button #9)
l) Selection of 1C population
i) Manually insert the Tot_Int threshold values for the 1C population for each animal in the
535   correct order
ii) Execute play button #10 to separate the 1C population from the >1C populations
m) Export the 1C population data (File name: "1C_manual"; median %TI, # comets) of each animal
to a txt-file (play button #11)
n) Export the >1C population data (File name: "More_than_1C_manual"; median %TI, # comets)
of each animal to a txt-file (play button #12)
Automated selection of 1C population
o) Create distribution plots for each animal using Gaussian mixture model with three forced
distributions (play button #13; Package: mclust, Function: densityMclust). The distribution plots
include median %TI + 1 and 2 standard deviations of the first population (**Figure 9**). Evaluate,
based on the plots, which # +SD is appropriate for your dataset, and make sure to avoid setting
the threshold so that >1C comets are included
i) Delimit the 1C population by examining the distribution plots and the best fitting # standard
deviations of the median % Tot_Int as threshold
Optional: The modelled testis distribution plots can be compared with somatic/4C-plots
generated in j) to verify the correct identification of the first population as the 1C population.
p) Export the 1C population data (File name: "1C_modelling"; median %TI, # comets) of each
animal to a txt-file (play button #14)
q) Export the >1C population data (File name: ">1C_modelling"; median %TI, # comets) of each
animal to a txt file (play button #14)
r) Calculate the mean +/- SD % TI of each experimental group. Perform statistical analyses as
deemed appropriate
557
558   **Note**: When using the modeling inspect the population settings to safeguard that the modelling of the
559   three populations is correct.
560

562
563 **Figure 7.** Tot_Int distribution plots of "all round comets" (grey), PS (green) and liver (red) from one
564 animal derived using the R-script, step I, with Tot_Int on x-axis and frequency of comets on the y-axis.
565

566
567 **Figure 8.** Tot_Int distribution plots of "all round comets" from one animal derived using modelling (R-
568 script, step o), with Tot_Int on x-axis and frequency of comets on the y-axis and blue line is the forced
569 3-population modelling.
570
571

572 **Optional: Enrichment of 4C primary spermatocyte comets**
573 1) Import the separately scored PS comets for each animal into a suitable software, such as Excel by
574   collapsing all comets per animal and calculate the median % TI (or use the Excel-template)
575
576 2) Export the % TI values for the PS comets to your software for statistical analyses of preference
577   a) Calculate the mean +/- SD %TI of the 4C comets from each experimental group and perform
578   statistical analyses as deemed appropriate
579
580 **Optional**: Verification of the Tot_Int level of the identified 4C population can be done by comparing
581 with the Tot_Int distributions (2C and 4C) of somatic origin generated above (or use the Excel-template).
582
583
584
585

586
587

Expected outcomes

Example of derived data

Please see examples below of Tot_Int distribution diagrams for "all round comets", PS and liver processed using the R-template Visual approach with a red line indicating where the cut-off for selecting 1C spermatids should be set.

Figure 9. Tot_Int distribution plots for "all round comets", PS and liver processed using the R-template Visual approach with a dashed red line showing the Tot_Int threshold level discriminating 1C from > 1C comets.

Figure 10. DNA damage levels (% Tail DNA) in (a)1C spermatid comets and (b) enriched 4C PS from
a rat exposed to a genotoxic agent at two different dose levels. The data are obtained using the R-
template "Visual" method.

Quantification and statistical analysis (optional)

1. All comets of each cell type from all gels for each biological replicate (animal) are compiled and the median % TI is calculated.

2. For each cell type and experimental group the % TI mean +/- SD is calculated.

3. Depending on normality and homogeneity of variances within each group, parametric tests or non-parametric tests can be used to test the likelihood of significant differences in %TI between experimental groups. Transformation of non-normal data can be explored.

Considerations and Limitations

Time from animal euthanasia to generation of testicular single-cell suspension: It is imperative that these processes are conducted rapidly (in accord with the indicated timelines in the protocol) and that tissues and cells are continuously kept on ice to reduce the chance of repair of the potentially induced DNA lesions.

Standardization: The protocol benefits greatly from standardization of many parameters during the animal experiment concerning housing, treatment, timing between animals, technical personnel as well as during the cell isolation and the comet assay (agarose, duration of gel electrophoresis, staining, scoring and post-scoring data curation) to reduce variability and obtain repeatable results. Recommendations for the comet assay can be found in (5).

Scoring of comets: scoring of testicular comets is skill/time-demanding since specific nucleoids are selected (the round nucleoids and not the elongating/elongated spermatids or sperm).

Troubleshooting

Potential solution:

- [Describe the details of the solution to resolve this problem.
- You can use a bulleted list if needed.]

Problem 1: Challenging to distinguish elongating/elongated spermatids from the round nucleoids during comet scoring

Potential solution:

Use the microscope image to determine if each small nuclei has a crescent like nuclei, and if the comet scored may lie in a plane where the crescent shape may be located horizontally so that it appears similar as a round nucleoid.

Problem 2: High cell density and overlapping comets
Potential solution:

Make sure to check the cell concentration using a Bürker chamber during single cell preparation, prior to gel casting. Alternatively, add gels of more than one dilution of cells.

Problem 3: High % TI background levels
Potential solution:

- Keep time from harvest to snap-freeze of tissue pieces as short as possible.
- Keep the time from thawing of frozen testis tissue until casting of gels as short as possible.
- Make sure to avoid harsh treatment of cell suspensions.
- Minimize the time on ice for the cell suspensions.
- Make sure that the volt/cm over stage is appropriate.
- Make sure that the agarose concentration is appropriate.
- Ensure that the scoring conditions are appropriate.

Problem 4: Low % TI levels
Potential solution:

- Keep tissues/cells on ice at all times.
- Keep the time from tissue harvest/thawing to final cell solution on ice as short as possible.
- Prevent evaporation of the agarose solution that could lead to low % TI levels
- Make sure that the volt/cm over stage is appropriate

Problem 5: Poor separation between populations by forced 3-population modelling

Potential solution:

Transform raw comet data using e.g. square root, cube root or logarithmic transformations, and re-analyse.

Resource availability

Lead contact
Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Ann-Karin Hardie Olsen (anka@nilu.no).

Materials availability
Not relevant.

Data and code availability
The codes for the R-scripts are provided

Acknowledgments

Funding was received from the Nordic Working Group for Chemicals, Environment, and Health, under the Nordic Council of Ministers (2021-022, 2021-101, 2022-12, 2023-04, 2024-04), however, the content does not necessarily reflect the Nordic Council of Ministers' views, opinions, attitudes, or recommendations. Funding was also received from the Research Council of Norway through its Centers of Excellence funding scheme, project number 223268/F50 CERAD.

Sincere thanks to Linda Ly (animal experiment), Dina Behmen (comet assay) and Jill Andersen (comet scoring) for excellent technical assistance.

Author contributions

A.K.O. and G.B. conceptualized the concept. A.K.O., Y.D and A.S. designed the experiments. A.K.O., Y.D., X.M, H.D., A-M.Z.B and A.S. conducted the experiments. D.M.E. and X.M. developed the Excel-templates and R-scripts. A.K.O., X.M. and C.Z performed statistical analyses and generated figures. H.D. contributed to the Biorender figure. A.K.O., X.M and C. Z. wrote the manuscript. A.K.O., X.M, C. Z., H.D., Y.D., A-M.Z.B, A.S., D.M.E and G.B revised the manuscript. A.K.O. and A.S. supervised experiments and analyses and provided grant support.

Declaration of interests

The authors declare no competing interests.

References

1.	Gutzkow KB, Langleite TM, Meier S, Graupner A, Collins AR, Brunborg G. High-throughput comet assay using 96 minigels. Mutagenesis. 2013;28(3):333-40.

2.	Singh NP, McCoy MT, Tice RR, Schneider EL. A simple technique for quantitation of low levels of DNA damage in individual cells. Exp Cell Res. 1988;175(1):184-91.

3.	Olsen AK, Bjortuft H, Wiger R, Holme J, Seeberg E, Bjoras M, et al. Highly efficient base excision repair (BER) in human and rat male germ cells. Nucleic Acids Res. 2001;29(8):1781-90.

4.	Olsen AK, Duale N, Bjoras M, Larsen CT, Wiger R, Holme JA, et al. Limited repair of 8-hydroxy-7,8-dihydroguanine residues in human testicular cells. Nucleic Acids Res. 2003;31(4):1351-63.

5.	Collins A, Moller P, Gajski G, Vodenkova S, Abdulwahed A, Anderson D, et al. Measuring DNA modifications with the comet assay: a compendium of protocols. Nat Protoc. 2023;18(3):929-89.

6.	OECD Test No. 489: In Vivo Mammalian Alkaline Comet Assay. OECD Guidelines for the Testing of Chemicals, Section 4 Health Effects. 2016.

7.	Brunborg G, Collins A. Guidance for publishing comet assay results. Mutat Res Genet Toxicol Environ Mutagen. 2020;854-855:503146.

8.	Moller P, Azqueta A, Boutet-Robinet E, Koppen G, Bonassi S, Milic M, et al. Minimum Information for Reporting on the Comet Assay (MIRCA): recommendations for describing comet assay procedures and results. Nat Protoc. 2020;15(12):3817-26.

9.	Jansen J, Olsen AK, Wiger R, Naegeli H, de Boer P, van Der Hoeven F, et al. Nucleotide excision repair in rat male germ cells: low level of repair in intact cells contrasts with high dual incision activity in vitro. Nucleic Acids Res. 2001;29(8):1791-800.

10.	Brunborg G, Eide DM, Graupner A, Gutzkow K, Shaposhnikov S, Kruszewski M, et al. Calibration of the comet assay using ionising radiation. Mutat Res Genet Toxicol Environ Mutagen. 2023;885:503560.

11.	Enciso JM, Gutzkow KB, Brunborg G, Olsen AK, Lopez de Cerain A, Azqueta A. Standardisation of the in vitro comet assay: influence of lysis time and lysis solution composition on the detection of DNA damage induced by X-rays. Mutagenesis. 2018;33(1):25-30.