Transcriptomic analysis of adult mouse cardiac stromal cells using single-cell qRT-PCR

Rita Alonaizan[1,2,3]✉, Patricia Chaves-Guerrero[4], Sara Samari[4], Michela Noseda[4], Nicola Smart[1,2], Carolyn Carr[1]

1 Department of Physiology, Anatomy & Genetics, University of Oxford, Oxford, UK.
2 Institute of Developmental and Regenerative Medicine, Old Road Campus, University of Oxford, Oxford, UK.
3 King Faisal Specialist Hospital & Research Centre, Riyadh, Saudi Arabia.
4 National Heart and Lung Institute, Imperial College London, London, UK.
✉ Correspondence: rita.alonaizan@dpag.ox.ac.uk

Keywords: heart regeneration, cell therapy, cardiosphere-derived cells, cardiac progenitor cells, myocardial infarction, single-cell qRT-PCR.

Abstract

Fate mapping studies have challenged the longstanding view of the adult mammalian heart
as a post-mitotic organ, suggesting limited cardiomyocyte renewal. This has spurred efforts to
identify cardiac progenitor cell (CPC) populations, but their contribution to cardiac
regeneration has been found to be minimal compared to cardiomyocyte proliferation. Despite
this, CPC transplantation has shown therapeutic potential through paracrine signalling.
20
The identity of CPCs remains unclear due to overlapping characteristics with other cardiac
stromal cell populations such as fibroblasts, mesenchymal cells, and pericytes. This study
sought to optimise the isolation of CPCs by developing a cardiac collagenase-trypsin (CT)
protocol, which was compared to the established method of isolating cardiosphere-derived
cells (CDCs). The CT protocol resulted in a higher cell yield and reduced expansion time, with
both CTs and CDCs showing superior survival potential under serum starvation compared to
commercially acquired cardiac fibroblasts (CFs).
28
Single-cell qRT-PCR analysis revealed that CTs and CDCs share a similar gene expression
profile, distinct from CFs, characterised by the enrichment of cardiogenic transcription factors.
Notably, CTs exhibited higher expression of *Tcf21* and lower *Tbx5*, suggesting an epicardial-
derived fibroblast phenotype, whereas *Tbx5* was enriched in CDCs and CFs. Additionally, CTs
showed an enrichment of macrophage-associated genes *Mrc1* and *Csf1r*, possibly due to the
transdifferentiation of macrophages to or from a fibroblast phenotype in a subset of CTs. The
study concludes that CTs represent a robust and efficient source of CPCs with therapeutic
potential and offers insights into the complex identity of cardiac stromal cells.

Introduction

Evidence stemming from fate mapping studies of cardiomyocyte renewal in the adult mammalian heart challenged the dogma considering the heart as a post-mitotic organ[1-3]. Therefore, multiple research groups attempted to characterise endogenous cardiac populations that may have a role in the regenerative process. Identifiers of the presumed cardiac progenitor cell (CPC) populations include SCA1 and the SP dye-efflux phenotype, KIT, ISL1, cardiosphere- and colony-forming assays, or re-activation of the embryonic epicardial programme and WT1 upregulation[4]. More recent findings indicate that the mechanism behind cardiomyocyte renewal is cardiomyocyte proliferation, and that CPC contribution is minimal[1-6]. Despite their low cardiomyogenic potential after transplantation into the infarcted heart, CPCs have demonstrated therapeutic benefits in pre-clinical studies, primarily due to paracrine signalling[7-11].

The identity of CPCs remains unclear, with numerous studies questioning the distinctions between CPCs and various cardiac stromal cell populations, including mesenchymal cells, fibroblasts, and pericytes[12-16]. This uncertainty stems from the overlap in their morphology and gene expression profiles. For instance, cardiac fibroblasts, like CPCs, express cardiogenic transcription factors that contribute to cardiac development and repair[17]. Both cardiac and tail fibroblasts share a molecular signature similar to that of mesenchymal stem cells (MSCs)[18]. Significant efforts have been made to distinguish these cardiac stromal cells. To complicate matters further, single-cell transcriptional profiling of mouse ventricular non-myocytes has revealed subpopulation heterogeneity within the cardiac fibroblast population[19]. Additionally, markers commonly used to identify pericytes are relatively non-specific, albeit some genes are enriched in pericytes compared to other cell types[19].

Certain research groups have sorted CPCs based on surface marker expression followed by clonogenic selection, which include the derivation of SCA1+ PDGFRα+/SP cells[9,20]. These methods produced homogeneous, self-renewing cardiogenic populations that showed promise. However, these cell types are found at very low rates in the adult mouse heart, with a reported cloning efficiency of around 1-2%. Ideally, an optimal cell isolation protocol would yield a large, robust, and reproducible population of cells in the shortest possible duration. The isolation of cardiosphere-derived cells (CDCs) involves culturing cardiac explants to harvest explant-derived cells (EDCs), which are then induced to form cardiospheres. EDCs vary in morphology and time required to achieve confluent culture, with CDC expansion times ranging from 5 to 84 days[21].

To optimise CPC isolation, we developed a cardiac collagenase-trypsin (CT) protocol adapted from studies by Gharaibeh *et al*.[22] and Okada *et al*.[23] for isolating slowly adhering cells (SACs) from skeletal muscle. These SACs demonstrated superior differentiation, survival, and therapeutic potential compared to rapidly adhering cells (RACs). Comparably, we showed that these CPCs, which we term CTs, can be induced to differentiate into the cardiomyocyte lineage *in vitro*[24] using TGFβ1 stimulation[25,26]. We also showed that fatty acid supplementation triggered a metabolic switch via the PPARα pathway, evidenced by upregulated oxidative metabolism and increased expression of cardiomyocyte markers in CTs[24]. This aligns with the metabolic switch from glycolysis to fatty acid oxidation observed during cardiomyocyte maturation in development[27]. In this study, we characterise the presumed CPC populations, CTs and CDCs, in comparison to commercially acquired primary cardiac fibroblasts (CFs) (**Figure 1**) using single-cell qRT-PCR and cell survival assays. Our aim is to profile CPCs obtained without clonogenic and marker selection and compare them to the cardiac fibroblast compartment.

Figure 1. Isolation method of the three cardiac stromal populations, CTs, CDCs and
CFs. *Schematic showing the protocol used to isolate CTs, CDCs, and CFs. The isolation*
protocol of CTs and CDCs is detailed in the Methods. As CFs were purchased and not isolated
in our lab, some information is lacking e.g. age of mice, duration of cell attachment, and
composition of the accompanying media. Illustration created with BioRender.com.

Results

The CT protocol produces higher cell yield compared to CDCs, but a similar survival
potential.
CTs and CDCs were isolated from adult mouse atrial tissue and expanded for further analysis.
The average time to reach 90% confluency at passage 0 and passage 1 was considerably
shorter for CTs, taking approximately 10 days and 20 days, respectively. In contrast, CDCs
required around 40 days to passage 0 and 60 days to passage 1, due to the additional steps
involved in explant culture and cardiosphere formation (**Figure 2.A**).

After expansion to passage 3, we assessed the survival potential of CTs compared to CDCs
and commercially acquired CFs to determine their robustness for transplantation. Serum
starvation was used to challenge the cells in culture for 3 days and 10 days. CTs, CDCs, and
CFs continued to grow following serum starvation with no significant difference in cell numbers
detected at day 3. At day 10, CT cell numbers were significantly higher than CFs but not CDCs,
suggesting that CTs have a superior survival potential compared to CFs (**Figure 2.B**).

Single-cell profiling of cardiac stromal cells.
CTs, CDCs and CFs isolated from adult mouse atrial tissue were expanded to passage 3 for
single-cell qRT-PCR analysis and compared with the HL-1 cardiomyocyte line which was used
as a control (**Figure 3**). Genes selected for this experiment (detailed in supplementary table
1) included those encoding cardiogenic transcription factors, stem cell-related markers,
fibroblasts, cardiomyocytes and other genes that were differentially expressed in the SCA1
subpopulations according to the Noseda *et al*. study[9]. Moreover, we attempted to detect
different cardiac subpopulations including pericytes, endothelial cells and macrophages within
the cardiac stromal cell populations. The heatmap (**Figure 3.A**) shows that the HL-1
cardiomyocytes were distinct from the cardiac stromal cells with expression of each gene
across the different populations shown in **Figure 3.B**. The cardiac stromal populations
expressed the fibroblast-associated genes *Cd44*, *Ddr2*, *Pdgfrb*, *Acta2* (encodes αSMA), and
Vim, albeit CFs had less consistent expression of *Pdgfrb*, and *Acta2*. Interestingly, *Ddr2, Acta2*
and *Vim* were also expressed by HL-1 cardiomyocytes. Additionally, on rare occasions, CFs
(n=2/29) and CTs (n=3/55) co-expressing *Wt1* and *Tcf21* were detected, perhaps inferring an
epicardial-derived fibroblast identity. *Ly6a* (encodes SCA1) and *Pdgfra* were expressed in all
three populations. The three cardiac stromal populations expressed little or no *Prg4* and *Wif1*
but displayed expression of *Medag* and *Col1a1*. *Ms4a4d* expression was detected in CTs and
some CFs only. Regarding pericyte-associated markers, *Kcnj8* showed little or no expression
in all populations, *Colec11* showed sporadic expression, whereas *Ng2* showed high
expression in all three cardiac stromal populations.

In CTs and CDCs, the most prevalent cardiogenic transcription factors were *Gata4*, *Tbx20*,
Hand2, and *Mef2a* with little or no expression of *Hand1*, *Isl1*, and *Nkx2-5*. In contrast, CFs had
less uniform expression of *Gata4*, *Tbx20*, and *Hand2*. Unlike CDCs, both CTs and CFs
expressed *Mef2c* heterogeneously. The expression of *Tbx5* in CFs followed a similar pattern
to CDCs but was more sporadic in CTs. Interestingly, only CFs expressed *Isl1,* which has a
very limited expression profile in the adult heart[28,29]. The three cardiac stromal populations
had very little or no expression of markers of cardiomyocytes (*Nppa*, *Myh6*, and *Myl2*),
haematopoietic (*Cd45*) and endothelial cells (*Cdh5* and *Vwf*), except for *Kdr* expression, which
was detected in the CTs, CFs and HL-1 cardiomyocytes. Expression of stem cell-related
markers (*Pou5f1*, *Nanog*, *Klf4*, and *Tert*) was found in the three cardiac stromal populations.
Pou5f1, *Nanog*, and *Klf4*, but not *Tert*, were also expressed in HL-1 cardiomyocytes. As
expected, HL-1 cardiomyocytes had a distinctly different expression profile and displayed
expression of most cardiogenic transcription factors (*Gata4*, *Hand2*, *Mef2a/c*, *Nkx2-5* and
Tbx5/20), and cardiomyocyte markers (*Nppa*, *Myh6*, and *Myl2*).

Expression of Schwann cell-related markers (*Plp1* and *Kcna1*) was found sporadically in all
populations, except for HL-1 cardiomyocytes where only *Plp1*, but not *Kcna1*, was expressed.
We also assessed the expression of the macrophage-associated genes *Mrc1*, *Csf1r* and
Cx3cr1. *Cx3cr1* showed no expression in any population, and *Mrc1* was only expressed in
CTs. In contrast, *Csf1r* was expressed in all three stromal populations but showed a more
uniform pattern in CTs.

A comparison of the molecular profiles of CPCs and CFs.
By principal component analysis (PCA), cardiac stromal cells and HL-1 cardiomyocytes were
resolved as discrete groups, which is consistent with their distinct phenotypes. Gene loadings
contributing to each dimension (Dim) suggest that a small subset of genes explain the cross-
group variability captured by Dim.1 and Dim.2. The separation of HL-1 cardiomyocytes was
attributable to *Wif1*, *Nkx2-5*, *Mef2c*, *Nppa*, *Myl2*, and *Myh6* (**Figure 4.A**). Some separation of
CFs from CTs was also revealed, with the CDCs clustering between the other two populations.
This can be seen more clearly in the PCA of the stromal populations alone (**Figure 4.B**). The
separation of CTs was attributable to *Tcf21*, clearly seen in **Figure 4.B**, while the separation
of CDCs and CFs was mainly attributable to *Tbx5*. This *Tcf21* versus *Tbx5* polarity may reflect
functional or origin heterogeneity. CFs appeared more dispersed, suggesting higher
heterogeneity, in comparison to CTs and CDCs, which again is more apparent in the PCA of
stromal populations only.

Statistical analysis comparing the three cardiac stromal population revealed that CTs were
more similar to CDCs than CFs with CTs showing significantly higher expression of *Vim* ($P <$
0.05), *Kdr* ($P < 0.05$), *Mrc1* ($P < 0.001$), *Tcf21* ($P < 0.0001$), *Ms4a4d* ($P < 0.0001$), *Tbx20* (P
< 0.0001), *Csf1r* ($P < 0.0001$), *Mef2c* ($P < 0.0001$), and *Medag* ($P < 0.05$), but lower expression
of *Tbx5* ($P < 0.0001$) and *Acta2* ($P < 0.01$), compared to CDCs. In comparison to the CFs,
CTs showed significantly higher expression of *Col1a1* ($P < 0.0001$), *Pdgfrb* ($P < 0.0001$),
Hand2 ($P < 0.0001$), *Pou5f1* ($P < 0.05$), *Mrc1* ($P < 0.001$), *Tcf21* ($P < 0.0001$), *Tbx20* ($P <$
0.0001), *Ng2* ($P < 0.01$), *Mef2c* ($P < 0.05$), *Medag* ($P < 0.001$), and *Gata4* ($P < 0.0001$), but
lower expression of *Isl1* ($P < 0.0001$) and *Tbx5* ($P < 0.01$) (**Figure 5.A**).

Finally, we performed co-expression correlation to further support the inferred biological
functions of the stromal and cardiomyocyte populations (**Figure 5.B**). Correlation plot analysis
showed a polarisation between the expression of some fibroblast-associated markers and
cardiogenic transcription factors, and a positive correlation between the expression of
cardiogenic transcription factors and cardiomyocyte markers. *Tcf21* expression showed a
positive correlation with the expression of *Tbx20*, *Ms4a4d*, and macrophage-associated genes
Csf1r and *Mrc1*. This is attributed to the high co-expression of these genes in CTs. In addition,
expression of pericyte-associated gene *Kcnj8* showed a positive correlation with the
expression of *Isl1* due to their co-expression pattern in a small number of CFs. Collectively,
these data confirm the co-expression of cardiogenic markers, associated predominantly with
CTs and CDCs, as distinct from the gene profile that characterises CFs. Additionally, the
enrichment of *Tcf21* expression in CTs further delineates CTs from CFs.

	Average time to 90% confluency at P0	Average time to 90% confluency at P1
CDCs	40 days	60 days
CTs	10 days	20 days

178  *Figure 2.* **The CT protocol produces higher cell yield compared to CDCs but a**
179  **similar survival potential.** *(A) Average time to reach 90% at passage 0 and*
180  *passage 1, as assessed by cell counting, in CTs and CDCs. (B) Cell Survival*
181  *following serum starvation for 3 or 10 days, as assessed by cell counting, in CTs,*
182  *CDCs and CFs. Analysed using a two-way ANOVA with Tukey's multiple*
183  *comparisons test to determine statistical significance (n=4), * p < 0.05, ** p < 0.01.*

Figure 3. Single-cell analysis reveals an enrichment of Tcf21 and low expression of Tbx5 in CTs, compared to CDCs and CFs. (A) *Expression of 43 genes was analysed at the single-cell level in the four cell populations shown: CTs (n=55), CDCs (n=41) and CFs (n=29) at passage 3, in addition to HL-1 cardiomyocytes (n=13). Both CTs and CDCs represent 3 biological replicates. The heatmap illustrates expression as -ΔCT values (blue is low or absent; red is high). Samples were ordered based on cell type, and the genes grouped by a hierarchical clustering algorithm based on the underlying co-expression pattern. **(B)** Single-cell expression of the 43 genes tested, represented individually in the four cell populations shown with the mean represented as a black diamond in each set.*

B

Figure 4. PCA of the three cardiac stromal populations and HL-1 cardiomyocytes. (A) *PCA of the four cell populations shown: CTs (n=55), CDCs (n=41) and CFs (n=29) at passage 3, in addition to HL-1 cardiomyocytes (n=13). Both CTs and CDCs represent 3 biological replicates. Dim.1 separates the cardiac stromal populations, whereas Dim.2 shows a distinct separation between cardiac stromal and myocyte populations. Gene loadings contributing to each Dim suggest that a small subset of genes explain the cross-group variability captured by Dim.1 and Dim.2. Tcf21 is associated with CTs (emphasised in the blue circle), whereas Tbx5 is associated with CDCs and CFs (emphasised in the brown circle). The separation between cardiac stromal and myocyte populations is reflected by clustering of cardiogenic and cardiomyocyte genes (Nkx2.5, Wif1, Nppa, Myl2 and Myh6), which are emphasised in the pink circle. **(B)** PCA of the three cardiac stromal populations alone: CTs (n=28), CDCs (n=41), CFs (n=27). Both CTs and CDCs represent 3 biological replicates. Gene loadings contributing to each Dim show that Tcf21 is associated with CTs, whereas Tbx5 is associated with CDCs and CFs.*

Figure 5. Statistical analysis of the single-cell qRT-PCR and gene correlations.
(A) Statistical analysis of the four cell populations shown: CTs (n=55), CDCs (n=41) and CFs (n=29) at passage 3, in addition to HL-1 cardiomyocytes (n=13). Both CTs and CDCs represent 3 biological replicates. The heatmap illustrates P values from 0 to 0.1 (orange to blue is low to high). The genes were grouped by a hierarchical clustering algorithm based on the underlying co-expression pattern. (B) Gene correlation plot, derived from the single-cell qRT-PCR data, is represented by the colour of the dots (blue is negative; red is positive). Both the diameter and colour intensity of each dot are proportional to the level of correlation.

B

Discussion

CTs have reduced expansion time and a superior survival potential following serum starvation.

Low survival and retention of transplanted cells following ischaemic events is mainly due to the harsh conditions within the infarct region that include hypoxia, inflammation, and nutrient restriction. Serum in cell culture media provides amino and fatty acids and its withdrawal is therefore, considered a form of nutrient restriction and serum starvation is an effective way of inducing cell apoptosis[30]. Here, we found that all three cardiac stromal populations continued to grow following serum starvation with no significant difference in cell numbers detected at day 3. However, CF cell numbers at day 10 were similar to the cell number seeded initially suggesting substantial cell death. At day 10, CT cell numbers were significantly higher than CFs but not CDCs, suggesting that CTs have a superior survival potential to that of CFs. This is consistent with the single-cell qRT-PCR results showing that CTs and CDCs share a more similar gene expression profile in comparison to the CFs. Tolerance to serum starvation has also been observed in MSCs under both normoxic and hypoxic culture conditions[31,32].

Reduced robustness was noticed during routine culture of CFs in comparison to CTs and CDCs as shown by increased time to reach confluency and higher numbers of floating dead cells in culture (data not shown). The CFs displayed reduced recovery following single-cell sorting for the single-cell qRT-PCR experiment, in comparison to CTs and CDCs. It is difficult to pinpoint what is responsible for this fragility due to difference in gene expression, cell culture media and coating substrate. Differential effects of various culture coating substrate types on cell proliferation have been shown[33]. However, standardising the culture conditions of all cell types would have been counterproductive as our aim was to compare the populations as they would be cultured in other labs based on their recommended and published culture conditions. Ideally, the survival potential of these different stromal population should be compared following transplantation into infarcted hearts or in *in vitro* conditions that more closely mimic the *in vivo* environment following MI. Overall, our data suggested that CTs and CDCs were more robust than CFs, both in long-term cell expansion and survival under serum starvation. The survival potential of cardiac stromal populations may differ in response to stress *in vitro*, which can be an indication of their *in vivo* behaviour following transplantation. Additionally, the reduced expansion time needed to generate CTs, compared to CDCs, indicated that a CDC-like population for therapeutic use could be achieved more efficiently.

CPCs show a fibroblast phenotype with no detection of endothelial cells or pericytes.

A study by Skelly *et al*. 2018 examined the single-cell transcriptional profile of adult mouse ventricular nonmyocytes to identify distinct cell populations[19]. Subpopulation heterogeneity within the cardiac fibroblast population was described where most fibroblasts "fibroblasts 1" were characterised by high expression of *Medag, Ms4a4d, Pdgfra* and *Tcf21*, while a smaller distinct cluster "fibroblasts 2" expressed higher levels of *Tbx20*, *Wif1* and *Prg4*. Although we found that *Tbx20* was highly expressed in both CTs and CDCs, *Wif1* and *Prg4* showed little to no expression in all stromal populations. Moreover, *Medag* and *Pdgfra* expression was detected in all three populations, whereas *Ms4a4d* and *Tcf21* was only found in CTs and CFs. However, CFs showed less consistent expression of *Ms4a4d* compared to CTs. This suggests that none of the three stromal populations fit the expression pattern of "fibroblasts 2" and that CTs were the best match to the expression pattern of "fibroblasts 1". However, an important consideration to make is that the study by Skelly *et al*. examined ventricular rather than atrial nonmyocytes and used freshly isolated cells rather than passaged cells which may undergo phenotypic drift *in vitro*.

Additional mesenchymal- and fibroblast-associated genes were assessed including *Vim*, *Ddr2*, *Acta2*, *Cd44*, and *Col1a1*. αSMA, encoded by *Acta2,* is a protein commonly used to identify smooth muscle cells, pericytes and myofibroblasts, which acquire a contractile phenotype similar to that of smooth muscle cells upon differentiation[34]. *Vim* was the only gene

expressed in all the screened cells of all populations. Vimentin is an intermediate filament
protein that has been widely used as a reliable marker of mesenchymal cells, highly migratory
cells derived by EMT, and has a protective role against nuclear rupture and DNA damage
during migration[35]. None of these markers showed specificity to a certain population. *Vim*,
Ddr2 and *Acta2* were also expressed in HL-1 cardiomyocytes, an adult immortalised atrial cell
line that has been shown to be similar to a mitotic embryonic cardiomyocyte rather than a
mature adult cell[36], as reflected by their immature energy metabolism[37]. The expression of
mesenchymal- and fibroblast-associated genes in HL-1 cardiomyocytes may reflect the lack
of specificity of these markers in *in vitro* cultures, the immature phenotype of HL-1
cardiomyocytes and/or their dedifferentiation in culture. Genomic instability is a common
setback associated with the use of cell lines in research[38]. Nonetheless, the above data
suggest a fibroblast-like phenotype of all stromal populations. A combination of *Pdgfra, Col1a1*
and Medag, which were expressed in the stromal cells but not the HL-1 cells, might be more
reliable markers for identifying *in vitro* fibroblast cultures.

Commonly used markers to identify pericytes (e.g. *Pdgfrb* and *Ng2*) are relatively non-specific.
However, *Kcnj8* and *Colec11* have been identified as genes that show higher expression in
pericytes relative to other cell types expressing them[19]. We found little or no expression of
Kcnj8 in any population. *Colec11* showed sporadic expression, whereas *Ng2* and *Pdgfrb*
showed high expression in all three cardiac stromal populations. Therefore, there is no
conclusive evidence to suggest that these stromal populations are pericytes in their
phenotype. Our data also suggest that the specificity of *Kcnj8* and *Colec11* expression to
pericytes might be superior to that of commonly used markers. Moreover, the expression of
Kcnj8 was positively correlated with *Isl1*, attributable to co-expression in a small number of
CFs. ISL1-mediated differentiation of coronary pericytes to coronary artery smooth muscle
cells has been described in the embryonic heart[39]. Although *Isl1* expression is detected in the
developing heart and multipotent ISL1[+] progenitors play an important role in giving rise to
cardiac lineages during development[40], ISL1[+] cells are very rare in adult human and murine
hearts[41,42].

The three cardiac stromal populations had very little or no expression of markers of
cardiomyocytes (*Nppa*, *Myh6*, and *Myl2*), haematopoietic (*Cd45*), or endothelial cells (*Cdh5*
and *Vwf*). Expression of *Nppa* and *Myh6* was found in few CDCs (2 *Nppa*[+] cells, n=41) and
CTs (4 *Myh6*[+] cells, n=55). Given that these cells are at passage 3, it is highly unlikely that the
cells expressing *Nppa* or *Myh6* cells are cardiomyocytes that survived the isolation protocol.
This expression may be a result of spontaneous differentiation in culture or may reflect some
progenitor status. In addition, *Kdr* expression, a marker of endothelial cells, was detected in
the CTs, CFs and HL-1 cardiomyocytes. During development, KDR[+] progenitors, deriving from
the MESP1[+] mesodermal lineage, have a cardiovascular tri-lineage potential[43,44]. It has been
shown by lineage tracing that precursors expressing *Gata5* and *Wt1* give rise to the adult
SCA1[+]SP[+] population (PDGFRα[+]/CD31[-]), whereas the adult SCA1[+]SP[-] population (PDGFRα[-]
/CD31[+]) is mainly derived from precursors expressing *Kdr*[9]. However, CTs lack the expression
of other endothelial markers as observed in the SCA1[+]SP[-] population. *Kdr* expression has
also been described in coronary adventitial progenitors with cardiogenic potential[45] and in
endothelial progenitors[46]. Therefore, *Kdr* expression may reflect some progenitor status as
opposed to the presence of endothelial cells. CDCs have been examined at the single cell
transcriptomics level[47] and characterised as mesenchymal/stromal/fibroblast-like, with a small
subset of endothelial-like cells. Notably, transplantation of SCA1+ CDCs but not SCA1− CDCs
enhanced cardiac function after MI, highlighting the previously unappreciated functional
differences between CDC subpopulations.

A few cells co-expressing *Wt1* and *Tcf21* were detected in both the CF and CT populations.
In the developing heart, cells of the epicardium are identified through the expression of *Wt1*,
Tbx18, *Tcf21*, and *Raldh2*[48]. It has been demonstrated that the majority of TCF21[+] epicardium-
derived cells (EPDCs) are committed to the cardiac fibroblast lineage and that *Tcf21*

expression persists in cardiac fibroblasts of the adult heart[49-51]. Although WT1 is
downregulated following gestation[52,53], its expression is still detected in the epicardium and a
subset of coronary endothelial cells in the healthy adult heart[50,54]. Enrichment of the epicardial
markers *Wt1* and *Tcf21* has been shown in adult cardiac fibroblasts[17]. More recent studies
showed that *Wt1* is expressed in more cell types than previously appreciated with a study
describing a Wt1b[+] macrophage subpopulation that plays an essential role in zebrafish heart
and fin regeneration[55]. In our data, *Wt1* expression did not show a correlation with *Mrc1* or *Kdr*
but all cells expressing *Wt1* were also positive for *Tcf21* suggesting a fibroblast phenotype.

Expression of stem cell-related markers (*Pou5f1*, *Nanog*, *Klf4*, and *Tert*) was found in all three
cardiac stromal populations. In addition, *Pou5f1*, *Nanog*, and *Klf4*, but not *Tert*, also showed
expression in HL-1 cardiomyocytes. The expression of these stem cell-associated genes
(*Pou5f1*, *Klf4*, *Nanog*) has been observed in mouse cardiac stromal populations and rat
CDCs[56]. A sporadic expression of these markers has also been noted in primary neonatal
cardiomyocytes[9]. Other studies have also reported the expression of *Pou5f1*[57]*, Klf4*[58]*,
Nanog*[59,60] in adult differentiated cells. Furthermore, *Tert* is activated in human and mouse
primary fibroblasts to protect against malignant transformation[61]. As stem cell-associated
genes have been implicated in tumorigenesis and proliferation[62,63], it is not surprising to
observe their expression in the HL-1 cell line or the stromal populations.

Tcf21 versus Tbx5 polarity may represent functional heterogeneity in fibroblast
populations and reveal a macrophage phenotype in a CT subpopulation.
Noseda *et al*.[9] revealed an enrichment of *Tcf21* and *Pdgfra*, the cardiogenic genes *Gata4,*
Mef2a/c, Tbx5/20 and Hand2, and a lack of *Cdh5* and *Kdr* in the SCA1[+] SP[+] population in
comparison to total SCA1[+] and SCA1[+] SP[-] populations. In our study, a lack of *Cdh5* expression
was observed in all cardiac stromal populations. Both CTs and CDCs showed high expression
of the cardiogenic transcription factors *Gata4*, *Tbx20*, *Hand2*, and *Mef2a* with CTs expressing
Mef2c and CDCs expressing *Tbx5*. Interestingly, among these transcription factors, *Tbx5*
showed the lowest expression in the SCA1[+] SP[+] population. Moreover, these results are
consistent with the observations of Furtado *et al*. showing an enrichment of cardiogenic genes
in cardiac fibroblasts[17]. In contrast, CFs had less uniform expression of *Gata4*, *Tbx20*, and
Hand2. The PCA analysis showed an interesting polarisation between *Tcf21* and *Tbx5*, which
was also reflected on the correlation plot showing a strong negative correlation between the
two genes. Indeed, the separation in clustering of CTs and CFs can be attributed to the
expression of these two genes. Although it is unclear what this polarisation means, it is an
interesting observation that warrants further investigations in future studies as it may reflect
potential functional heterogeneity.

In addition to the enrichment of *Tcf21* in CTs as opposed to other stromal populations, the
macrophage-associated gene *Mrc1* was detected in a subset of CTs. This was also reflected
by the positive correlation between *Mrc1* and *Tcf21* versus the negative correlation between
Mrc1 and *Tbx5*. *Mrc1* has been reported to show higher expression in macrophages, relative
to other cardiac cell types[19]. *Csf1r*, enriched in M2-polarised macrophages[64], was expressed
in all stromal populations, albeit at significantly higher levels in CTs. This suggests the
possibility of a macrophage-like subpopulation in CT populations, and potentially in CDC and
CF populations as well. *Csf1r* expression has also been used as an indicator of embryonic
myeloid lineage commitment[65]. However, cells expressing *Mrc1* or *Csf1r* showed co-
expression of fibroblast markers and did not show clear separation in clustering. Cells
expressing *Mrc1* or *Csf1r* may therefore represent the transdifferentiation of macrophages in
the stromal isolations into fibroblasts or *vice versa*. Indeed, the transdifferentiation of
macrophages to fibroblast cells has been described *in vivo*[66]. Moreover, hybrid mesenchymal
phenotypes have been previously described where fibroblast-associated genes are expressed
in minor subpopulations of macrophage or endothelial cells[19]. However, due to the lack of
Cd45 expression in CTs, it is more likely that macrophage-associated genes were upregulated

in a subset of fibroblasts rather than *vice versa*. This phenomenon has been previously
reported in *in vitro* fibroblast cultures[67].

Our results suggest that (1) CTs were similar to CDCs and differed from CFs in their
expression profile, enrichment of cardiogenic factors and robustness; (2) CTs, CDCs, and CFs
resembled "fibroblasts 1", which are the predominant fibroblast subtype in the adult murine
heart; (3) CDCs clustered between CTs and CFs which separated based on their expression
of *Tcf21* and *Tbx5*, respectively; (4) CTs were enriched for the macrophage-associated genes
Mrc1 and *Csf1r*, which may be attributed to transdifferentiation of macrophages to or from a
fibroblast phenotype in a subset of CTs; (5) PCA analysis revealed a transcriptional gradient
suggesting that heterogeneity exists within fibroblast subtypes which may reflect functional
heterogeneity. Overall, our findings highlight the heterogeneity of the stromal compartment of
the heart and how the isolation protocol may enrich for specific cell subpopulations, which may
ultimately affect the therapeutic potential of the cells. Furthermore, the decreased culture time
required to generate CTs, as compared to that for the CDCs, suggests that a CPC population
for therapeutic use could be achieved more efficiently.

Methods

1. Mice

C57BL/6 mice (Harlan, Oxon, UK) were kept under in a 12-hour light-dark cycle, and controlled conditions of temperature, humidity, with free access to water and chow. All animal procedures were reviewed and approved by the University of Oxford Animal Welfare and Ethical Review Board and conforms to the Animals (Scientific Procedures) Act 1986 incorporating Directive 2010/63/EU of the European Parliament.

2. Isolation and expansion of mouse CPCs

Mice were terminally anesthetised with isoflurane and hearts were isolated and washed with Dulbecco's phosphate buffered saline (DPBS) containing 50mg of primocin (antimicrobial agent; InvivoGen). Mouse atrial appendages were dissected and minced mechanically into 1 - 2 mm^3 pieces. For CT isolation, the tissue pieces were then transferred into a digestion mix (0.1% trypsin and 0.1% Collagenase II (Calbiochem, 286U/mg) in DPBS) and incubated in a water bath at 37°C for a total of 40 minutes. Every 10 minutes the digestion mix was mechanically triturated by pipetting, left on ice for 1 minute to settle and the supernatant was collected. Fresh digestion mix was added again to the tissue pieces for a total of 4 digestions. At the final digestion step, sterile 19G needles and 1ml syringes were used to triturate the tissue. After each digestion, the supernatant was neutralised, and the cell suspension was resuspended in fresh complete explant medium (CEM; Iscove's modified Dulbecco's medium supplemented with 20% FBS, 100U/ml penicillin, 100µg/ml streptomycin and 2mM L-glutamine, ThermoFisher)[11] to be plated in a 12-well plate through a 40µm cell strainer. The wells were pre-coated with 50µl per cm^2 of fibronectin (2µg/ml in DPBS; Sigma-Aldrich) and cells were allowed to attach for 3 days without media changes. For CDC isolation[11], the tissue pieces were digested in 0.05% Trypsin-EDTA (ThermoFisher) for 4 minutes at 37°C then neutralised with CEM. Processed tissue pieces were plated on fibronectin-coated (3µg/ml in DPBS) 6-well plates containing CEM and explants were cultured for 25-30 days at 37°C in 5% CO_2 with media changes every 2-3 days. When explant-derived cells (EDCs) reached 70-80% confluency, they were harvested by washing with 0.53mM Versene (ThermoFisher) before a trypsin digestion. EDCs were resuspended in Cardiosphere Growth Medium (CGM), comprising 65% Dulbecco's modified eagle medium (DMEM/F12), 35% IMDM, 7% foetal bovine serum (FBS; ThermoFisher), 2% B27 (ThermoFisher), 25ng/ml cardiotrophin (Peprotech EC), 10ng/ml epidermal growth factor (EGF; Peprotech EC), 20ng/ml basic fibroblast growth factor (FGF; Promega) and 5 units thrombin (Sigma-Aldrich), at a concentration of $2x10^4$ cells/ml and seeded as 25µl droplets on an un-coated lid of a Petri dish. DPBS was added to the Petri dishes to keep the hanging drops humid throughout 3 days of culture allowing the spherical multicellular clusters, known as cardiospheres, to form. Cardiospheres were collected by elution and gentle pipetting with DPBS and were plated in fibronectin-coated 12-well plates with CEM allowing them to spontaneously release CDCs.

3. Cell culture

Primary CPCs were cultured in CEM and plated on fibronectin-coated flasks. HL-1 cardiomyocytes[36] were maintained in Claycomb medium (Sigma-Aldrich), supplemented with 100U/ml penicillin, 100µg/ml streptomycin and 2mM L-glutamine (ThermoFisher), 100µM norepinephrine (Sigma-Aldrich) in 30mM L-ascorbic acid (Sigma-Aldrich), and 10% FBS and plated on flaks pre-coated with 0.02% (wt/vol) gelatin (Sigma-Aldrich) containing 5µg/ml fibronectin. C57BL/6 adult mouse atrial primary cardiac fibroblasts, obtained from Cell Biologics (USA), were maintained in complete fibroblast medium (Cell Biologics – USA) supplemented with the provided supplement kit: FGF, hydrocortisone, L-glutamine, Antibiotic-Antimycotic Solution, and 10% FBS and plated on flasks pre-coated with 0.02% (wt/vol) gelatin. For serum starvation experiments, cells were switched to serum-free media for 72 hours.

4. Single-cell qRT-PCR

Single cells were sorted directly (FACSAria II) into 96-well plates containing the reaction mixture for pre-amplification, using CellsDirect One-Step qRT–PCR Kits (ThermoFisher). Pre-

amplification was performed in a Veriti Thermal Cycler (Applied Biosystems) for 22 cycles. As negative controls, at least 3–5 non-template samples were included in each run at the pre-amplification stage. Quantitative amplification was performed using Dynamic Array chips for 48 assays x 48 samples, the BioMark HD system (Fluidigm) and TaqMan probes (Appendix D) according to manufacturer's instructions. Each sample was normalised to ΔCt using the expression of *Ubc*. As described in Noseda *et al*. 2015[9], data were plotted in colour-coded heatmaps of inverted ΔCt values (blue is low or absent; red is high). Samples were ordered based on cell type, and the genes grouped by a hierarchical clustering algorithm based on the underlying co-expression pattern. Differences between samples were investigated using PCA. PCA applies multiple linear transformations (singular value decomposition) to the expression profiles (standardised ΔCt values) of individual samples, and identifies a series of PCs that elucidate the most distinguishing features between the samples. The linear projections (PC scores) attempt to maximise the variation between the samples, whereas the coefficients of those projections (PC loadings) measure the importance of genes in defining the underlying variability associated with each component.

Taqman probes used for single-cell qRT-PCR

Mesp1	Mm00801883_g1	*Nkx2-5*	Mm00657783_m1
Mef2c	Mm01340842_m1	*Myh11*	Mm00443013_m1
Mef2a	Mm01318990_m1	*Cnn1*	Mm00487032_m1
Hand1	Mm00433931_m1	*αSMA (Acta2)*	Mm01546133_m1
Islet1 (Isl1)	Mm00627860_m1	*Vim*	Mm01333430_m1
Tbx20	Mm00451515_m1	*Cd44*	Mm01277163_m1
Gata4	Mm00484689_m1	*Pdgfrb*	Mm00435546_m1
Hand2	Mm00439247_m1	*Ng2*	Mm01283063_m1
Tbx5	Mm00803518_m1	*Ddr2*	Mm00445615_m1
Ly6a (Sca1)	Mm00726565_s1	*Medag*	Mm00551008_m1
Wt1	Mm00460570_m1	*Ms4a4d*	Mm00656404_m1
Klf4	Mm00516104_m1	*Col1a1*	Mm00801666_g1
Pou5f1 (Oct4)	Mm03053917_g1	*Prg4*	Mm01284582_m1
Nanog	Mm02019550_s1	*Wif1*	Mm00442355_m1
Tert	Mm0352136_m1	*Colec11*	Mm01289834_m1
Kdr	Mm00440111_m1	*Kcnj8*	Mm00434620_m1
Cdh5	Mm00486938_m1	*Mrc1*	Mm01329362_m1
Vwf	Mm00550376_m1	*Cx3cr1*	Mm00438354_m1
Pdgfra	Mm01211685_m1	*Csf1 r*	Mm01266652_m1
Tcf21	Mm00448961_m1	*Kcna1*	Mm00439977_s1
Myl2	Mm00440384_m1	*Plp1*	Mm01297210_m1
Myh6	Mm00440359_m1	*Ptprc (Cd45)*	Mm01293577_m1
Nppa	Mm01255748_g1	*Ubc*	Mm01201237_m1
Hmbs	Mm01143545_m1		

459 **Supplementary Table 1.** Description of the markers used in the single-cell qRT-PCR
460 characterisation.
461

Gene	Full name of encoded protein	Type	Relevant expression pattern	Relevant function	References
Cd44	Cluster of differentiation 44	Transmembrane glycoprotein	Fibroblasts, macrophages, endothelial cells	Cellular adhesion; angiogenesis; release of cytokines; regulates inflammation and fibrosis	68-71
Cdh5	Cadherin 5 or VE-cadherin	Member of the cadherin family of calcium-dependent glycoproteins	Endothelial cells	Calcium-dependent cell adhesion; cell proliferation and migration	68,72
Col1a1	Collagen, type I, alpha 1	Member of group I collagen	Fibroblasts, smooth muscle cells	Collagen type I synthesis	68,73
Colec11	Collectin subfamily member 11	Member of the collectin family of C-type lectins	Pericytes, endothelial cells	Collectins are initiators of the lectin complement pathway, which is integral to the innate immune response; embryogenesis	19,68,74,75
Csf1r	Colony-stimulating factor 1 receptor	Receptor tyrosine kinase	Macrophages (used as an M2 marker)	Growth and differentiation of macrophages; maintenance of the intestinal stem cell niche	64,76
Cx3cr1	CX3C chemokine receptor 1	Member of the 7-transmembrane G protein-coupled receptor family	Macrophages	Macrophage survival and function	68,77,78
Ddr2	Discoidin domain receptor tyrosine kinase 2 or CD167b	Receptor tyrosine kinase	Fibroblasts, smooth muscle cells, pericytes	Activated by collagen binding; cell proliferation; adhesion and migration; ECM degradation	68,79,80
Gata4	GATA binding factor 4	Transcription factor of the zinc-finger GATA family	Cardiomyocytes, cardiac fibroblasts	Cardiogenesis	17,18,81-83
Hand1	Heart- and neural crest derivatives-expressed protein 1	Transcription factor of the basic helix-loop-helix family	FHF, reported to be undetectable in adult mouse hearts	Cardiogenesis; epicardial development and coronary vessel formation	84-87
Hand2	Heart- and neural crest derivatives-expressed protein 2	Transcription factor of the basic helix-loop-helix family	SHF, cardiomyocytes, endothelial cells, fibroblasts	Cardiogenesis; epicardial development and coronary vessel formation	9,18,84,85,87
Isl1	Insulin gene enhancer protein 1	Transcription factor of the LIM/homeodomain family	CPCs, cardiac neural crest, endothelial cells	Cardiogenesis	28,40-42,88
Kcna1	Potassium voltage-gated channel subfamily A member 1	Member of the 6-TM family	Schwann cells	Regulator of membrane excitability	19,68,89
Kcnj8	Potassium inwardly-rectifying channel, subfamily J, member 8	The inward-rectifier potassium channel family (also known as 2-TM channels)	Pericytes	Regulator of vascular tone and cardiac adaptive response to systemic metabolic stressors	19,68,90-92
Kdr	Kinase insert domain receptor or foetal liver kinase 1 (FLK1); VEGFR2; CD309	Receptor tyrosine kinase	Endothelial cells, Kdr[+] developmental cardiovascular progenitor	Transduction of VEGF-A effects	43,44,68,93
Klf4	Krüppel like factor 4	Member of the Krüppel-like family of transcription factors	Stem cells, endothelial cells, fibroblasts	Pluripotency; angiogenesis; embryonic vasculogenesis; myofibroblast differentiation	58,68,94-96
Medag	Mesenteric oestrogen-dependent	Member of the sex-steroid regulated mesenteric	Cardiac fibroblasts "1"	Adipogenesis; glucose uptake; interacts with the fatty acid	19,97,98

	adipogenesis protein	oestrogen- dependent adipogenesis proteins		transporter CD36 that is involved in the uptake of apoptotic material	
Mef2a	Myocyte-specific enhancer factor 2A	Transcription factor of the MEF2 family	Cardiomyocytes, cardiac fibroblasts, endothelial cells, smooth muscle cells, macrophages and pericytes	Cardiogenesis; vasculogenesis	[9,68,99,100]
Mef2c	Myocyte-specific enhancer factor 2C or MADS box transcription enhancer factor 2	Transcription factor of the MEF2 family	Cardiomyocytes, cardiac fibroblasts, endothelial cells, smooth muscle cells, macrophages and pericytes	Cardiogenesis; vasculogenesis	[9,17,18,68,83,101,102]
Mesp1	Mesoderm Posterior Transcription Factor 1	Transcription factor of the basic helix-loop-helix family	Mesoderm	Development of cardiac mesoderm	[103]
Mrc1	Mannose receptor C-type 1 or CD206	Transmembrane glycoprotein	Macrophages (used as an M2 marker)	Mediates the endocytic functions by macrophages and the DNA replication checkpoint	[19,68,104-106]
Ms4a4d	Membrane-spanning 4-domains subfamily A member 4D	Member of the membrane-spanning 4A (MS4A) family	Cardiac fibroblasts "1", reported to be expressed in M2 macrophages	May have a role as an adaptor protein to facilitate intracellular protein–protein interactions. Its functionality is still not fully understood.	[19,107,108]
Myh6	Myosin heavy chain 6 or αMHC	Hexameric ATPase cellular motor protein	Predominantly atrial cardiomyocytes	Cardiogenesis; force generation and muscle contraction	[109]
Myl2	Myosin light chain 2	Hexameric ATPase cellular motor protein	Predominantly ventricular cardiomyocytes	Cardiogenesis; force generation and muscle contraction	[109]
Nanog	Nanog Homeobox	Transcription factor of the Nanog homeobox family	Stem cells	Pluripotency; tumourigenicity	[59,60,110]
Ng2	Neuron-glial antigen 2 or chondroitin sulphate proteoglycan 4	Transmembrane proteoglycan	Pericytes, smooth muscle cells	Co-receptor in conjunction with PDGFRα; angiogenesis; cell survival and migration	[68,111-113]
Nkx2-5	NK2 homeobox 5	Transcription factor of the NK-2 homeobox family	Cardiomyocytes	Cardiogenesis	[68,114]
Nppa	Natriuretic Peptide A	The natriuretic peptide family	Predominantly atrial cardiomyocytes	Cardiac hormone involved in regulating blood volume and pressure and cardiovascular homeostasis	[115]
Pou5f1	Octamer-binding transcription factor 4 (Oct4) or POU class 5 homeobox 1	Transcription factor of the homeodomain POU (Pit-Oct-Unc) family.	Stem cells	Pluripotency; tumourigenicity	[116,117]
Pdgfra	Platelet-derived growth factor receptor α	Receptor tyrosine kinase	Fibroblasts, defines the SCA1$^+$SP$^+$ population (CPCs)	Development of mesenchymal cell types; cardiogenesis	[9,68,118-120]
Pdgfrb	Platelet-derived growth factor receptor β	Receptor tyrosine kinase	Pericytes, fibroblasts, endothelial cells, smooth muscle cells	Development of mesenchymal cell types; regulation of autophagic processes; cell proliferation and survival; adult mammalian cardiomyocyte proliferation; cardiogenesis	[68,120-126]
Plp1	Proteolipid protein 1	Myelin proteolipid protein	Schwann cells, endothelial cells, smooth muscle cells,	Myelination	[127]

			pericytes and macrophages		
Prg4	Proteoglycan 4	Mucinous glycoprotein	Cardiac fibroblasts "2"	Ligand for CD44 receptor, ECM constituent conferring compression resistance	[19,128,129]
Ptprc (Cd45)	Protein tyrosine phosphatase, receptor type, C	Transmembrane glycoprotein	Haematopoietic cells (except erythrocytes and plasma cells)	Haematopoietic cell activation and differentiation and immune Function	[130-132]
Ly6a	Stem cell antigen 1 (SCA1) or lymphocyte antigen 6 complex	Glycosyl phosphatidylinositol-anchored cell surface protein of the LY6 gene family	HSCs, fibroblasts, endothelial cells, CPCs	Tumourigenicity; cell migration; HSC self-renewal and fate determination	[9,68,133-136]
Tbx20	T-box 20	Transcription factor of the T-box family	Cardiomyocytes, cardiac pericytes and fibroblasts	Cardiogenesis	[9,68]
Tbx5	T-box 5	Transcription factor of the T-box family	Cardiomyocytes, cardiac fibroblasts	Cardiogenesis	[17,18,83]
Tcf21	Transcription factor 21	Transcription factor of the basic helix-loop-helix family	Epicardium, Cardiac fibroblasts "1"	Initiation of epicardial EMT for fibroblast development; proepicardial cell specification	[19,49,137]
Tert	Telomerase reverse transcriptase	Member of the reverse transcriptase family	Stem and progenitor cells	Cellular immortalisation tumourigenicity; cell proliferation	[138,139]
Vim	Vimentin or fibroblast intermediate filament	Type III intermediate filament protein	Fibroblasts, endothelial cell, smooth muscle cells, Schwann cells	Maintenance of cellular integrity; EMT; protective role against nuclear rupture and DNA damage during cell migration.	[35,68]
Vwf	von Willebrand factor	Adhesive plasma glycoprotein	Endothelial cells	Platelet adhesion and aggregation	[140]
Wif1	WNT inhibitory factor 1	Lipid-binding protein	Cardiac fibroblasts "2"	Mesoderm segmentation; cardiogenesis	[19,141]
Wt1	Wilms tumour 1	Transcription factor with proline / glutamine- rich DNA- binding domain	Epicardium, fibroblasts, endothelial cells, macrophages	Epicardial EMT; vasculogenesis	[17,54,55,142,143]
Acta2	Alpha smooth muscle actin (αSMA) or Actin alpha 2	Member of the globular Actin proteins family	Smooth muscle cells, myofibroblasts, pericytes	Cell motility and contractility	[34,144]

463 **Author contributions**
464 Conceptualisation: RA, NS, CC. Investigation: RA, PCG, SS, MN. Funding acquisition: RA.
465 Writing: RA, CC.
466
467 **Declaration of interests**
468 None
469
470 **Acknowledgements**

471 This work was supported by the King Faisal Specialist Hospital & Research Centre (PhD scholarship
472 to RA).

References

Ali, S. R. *et al.* Existing cardiomyocytes generate cardiomyocytes at a low rate after birth in mice. *Proc Natl Acad Sci U S A* **111**, 8850-8855 (2014). https://doi.org/10.1073/pnas.1408233111

Kimura, W. *et al.* Corrigendum: Hypoxia fate mapping identifies cycling cardiomyocytes in the adult heart. *Nature* **532**, 268 (2016). https://doi.org/10.1038/nature16177

Senyo, S. E. *et al.* Mammalian heart renewal by pre-existing cardiomyocytes. *Nature* **493**, 433-436 (2013). https://doi.org/10.1038/nature11682

Alonaizan, R. & Carr, C. Cardiac regeneration following myocardial infarction: the need for regeneration and a review of cardiac stromal cell populations used for transplantation. *Biochem Soc Trans* **50**, 269-281 (2022). https://doi.org/10.1042/BST20210231

Eschenhagen, T. *et al.* Cardiomyocyte Regeneration: A Consensus Statement. *Circulation* **136**, 680-686 (2017). https://doi.org/10.1161/CIRCULATIONAHA.117.029343

Li, Y. *et al.* Genetic Tracing Identifies Early Segregation of the Cardiomyocyte and Nonmyocyte Lineages. *Circ Res* **125**, 343-355 (2019). https://doi.org/10.1161/CIRCRESAHA.119.315280

Carr, C. A. *et al.* Cardiosphere-derived cells improve function in the infarcted rat heart for at least 16 weeks--an MRI study. *PLoS One* **6**, e25669 (2011). https://doi.org/10.1371/journal.pone.0025669

Lee, S. T. *et al.* Intramyocardial injection of autologous cardiospheres or cardiosphere-derived cells preserves function and minimizes adverse ventricular remodeling in pigs with heart failure post-myocardial infarction. *J Am Coll Cardiol* **57**, 455-465 (2011). https://doi.org/10.1016/j.jacc.2010.07.049

Noseda, M. *et al.* PDGFRalpha demarcates the cardiogenic clonogenic Sca1+ stem/progenitor cell in adult murine myocardium. *Nat Commun* **6**, 6930 (2015). https://doi.org/10.1038/ncomms7930

Smart, N. *et al.* De novo cardiomyocytes from within the activated adult heart after injury. *Nature* **474**, 640-644 (2011). https://doi.org/10.1038/nature10188

Smith, R. R. *et al.* Regenerative potential of cardiosphere-derived cells expanded from percutaneous endomyocardial biopsy specimens. *Circulation* **115**, 896-908 (2007). https://doi.org/10.1161/CIRCULATIONAHA.106.655209

Caplan, A. I. All MSCs are pericytes? *Cell Stem Cell* **3**, 229-230 (2008). https://doi.org/10.1016/j.stem.2008.08.008

13   Denu, R. A. *et al.* Fibroblasts and Mesenchymal Stromal/Stem Cells Are Phenotypically
Indistinguishable.   *Acta   Haematol*   **136**,   85-97   (2016).
https://doi.org/10.1159/000445096

14   Hematti, P. Mesenchymal stromal cells and fibroblasts: a case of mistaken identity?
Cytotherapy **14**, 516-521 (2012). https://doi.org/10.3109/14653249.2012.677822

15   Murray, I. R. & Peault, B. Q&A: Mesenchymal stem cells - where do they come from
and is it important? *BMC Biol* **13**, 99 (2015). https://doi.org/10.1186/s12915-015-
0212-7

16   Soundararajan, M. & Kannan, S. Fibroblasts and mesenchymal stem cells: Two sides of
the   same   coin?   *J   Cell   Physiol*   **233**,   9099-9109   (2018).
https://doi.org/10.1002/jcp.26860

17   Furtado, M. B. *et al.* Cardiogenic genes expressed in cardiac fibroblasts contribute to
heart   development   and   repair.   *Circ   Res*   **114**,   1422-1434   (2014).
https://doi.org/10.1161/CIRCRESAHA.114.302530

18   Furtado, M. B. *et al.* Microarray profiling to analyse adult cardiac fibroblast identity.
Genom Data **2**, 345-350 (2014). https://doi.org/10.1016/j.gdata.2014.10.006

19   Skelly, D. A. *et al.* Single-Cell Transcriptional Profiling Reveals Cellular Diversity and
Intercommunication   in   the   Mouse   Heart.   *Cell Rep* **22**,   600-610   (2018).
https://doi.org/10.1016/j.celrep.2017.12.072

20   Vicinanza, C. *et al.* Adult cardiac stem cells are multipotent and robustly myogenic: c-
kit expression is necessary but not sufficient for their identification. *Cell Death Differ*
24, 2101-2116 (2017). https://doi.org/10.1038/cdd.2017.130

21   Chan, H. H. *et al.* Human cardiosphere-derived cells from patients with chronic
ischaemic heart disease can be routinely expanded from atrial but not epicardial
ventricular   biopsies.   *J Cardiovasc Transl Res* **5**,   678-687   (2012).
https://doi.org/10.1007/s12265-012-9389-0

22   Gharaibeh, B. *et al.* Isolation of a slowly adhering cell fraction containing stem cells
from murine skeletal muscle by the preplate technique. *Nat Protoc* **3**, 1501-1509
(2008). https://doi.org/10.1038/nprot.2008.142

23   Okada, M. *et al.* Human skeletal muscle cells with a slow adhesion rate after isolation
and an enhanced stress resistance improve function of ischemic hearts. *Mol Ther* **20**,
138-145 (2012). https://doi.org/10.1038/mt.2011.229

24   Malandraki-Miller, S. *et al.* Metabolic flux analyses to assess the differentiation of
adult cardiac progenitors after fatty acid supplementation. *Stem Cell Res* **38**, 101458
(2019). https://doi.org/10.1016/j.scr.2019.101458

25   Goumans, M. J. *et al.* TGF-beta1 induces efficient differentiation of human
cardiomyocyte progenitor cells into functional cardiomyocytes in vitro. *Stem Cell Res*
1, 138-149 (2007). https://doi.org/10.1016/j.scr.2008.02.003

26   Smits, A. M. *et al.* Human cardiomyocyte progenitor cells differentiate into functional
mature cardiomyocytes: an in vitro model for studying human cardiac physiology and
pathophysiology. *Nat Protoc* **4**, 232-243 (2009).
https://doi.org/10.1038/nprot.2008.229

27   Lopaschuk, G. D. & Jaswal, J. S. Energy metabolic phenotype of the cardiomyocyte
during development, differentiation, and postnatal maturation. *J Cardiovasc
Pharmacol* **56**, 130-140 (2010). https://doi.org/10.1097/FJC.0b013e3181e74a14

28   Laugwitz, K. L. *et al.* Postnatal isl1+ cardioblasts enter fully differentiated
cardiomyocyte lineages. *Nature* **433**, 647-653 (2005).
https://doi.org/10.1038/nature03215

29   Ye, J. *et al.* Sca-1+ cardiosphere-derived cells are enriched for Isl1-expressing cardiac
precursors and improve cardiac function after myocardial injury. *PLoS One* **7**, e30329
(2012). https://doi.org/10.1371/journal.pone.0030329

30   Lu, C. *et al.* Serum starvation induces H2AX phosphorylation to regulate apoptosis via
p38 MAPK pathway. *FEBS Lett* **582**, 2703-2708 (2008).
https://doi.org/10.1016/j.febslet.2008.06.051

31   Huang, Y. C. *et al.* Isolation of mesenchymal stem cells from human placental decidua
basalis and resistance to hypoxia and serum deprivation. *Stem Cell Rev Rep* **5**, 247-255
(2009). https://doi.org/10.1007/s12015-009-9069-x

32   Peng, L. *et al.* Comparative analysis of mesenchymal stem cells from bone marrow,
cartilage, and adipose tissue. *Stem Cells Dev* **17**, 761-773 (2008).
https://doi.org/10.1089/scd.2007.0217

33   Liberio, M. S., Sadowski, M. C., Soekmadji, C., Davis, R. A. & Nelson, C. C. Differential
effects of tissue culture coating substrates on prostate cancer cell adherence,
morphology and behavior. *PLoS One* **9**, e112122 (2014).
https://doi.org/10.1371/journal.pone.0112122

34   Rockey, D. C., Weymouth, N. & Shi, Z. Smooth muscle alpha actin (Acta2) and
myofibroblast function during hepatic wound healing. *PLoS One* **8**, e77166 (2013).
https://doi.org/10.1371/journal.pone.0077166

35   Patteson, A. E. *et al.* Vimentin protects cells against nuclear rupture and DNA damage
during migration. *J Cell Biol* **218**, 4079-4092 (2019).
https://doi.org/10.1083/jcb.201902046

36 Claycomb, W. C. *et al.* HL-1 cells: a cardiac muscle cell line that contracts and retains
phenotypic characteristics of the adult cardiomyocyte. *Proc Natl Acad Sci U S A* **95**,
2979-2984 (1998). https://doi.org/10.1073/pnas.95.6.2979

37 Kuznetsov, A. V., Javadov, S., Sickinger, S., Frotschnig, S. & Grimm, M. H9c2 and HL-1
cells demonstrate distinct features of energy metabolism, mitochondrial function and
sensitivity to hypoxia-reoxygenation. *Biochim Biophys Acta* **1853**, 276-284 (2015).
https://doi.org/10.1016/j.bbamcr.2014.11.015

38 Geraghty, R. J. *et al.* Guidelines for the use of cell lines in biomedical research. *Br J
Cancer* **111**, 1021-1046 (2014). https://doi.org/10.1038/bjc.2014.166

39 Volz, K. S. *et al.* Pericytes are progenitors for coronary artery smooth muscle. *Elife* **4**
(2015). https://doi.org/10.7554/eLife.10036

40 Moretti, A. *et al.* Multipotent embryonic isl1+ progenitor cells lead to cardiac, smooth
muscle, and endothelial cell diversification. *Cell* **127**, 1151-1165 (2006).
https://doi.org/10.1016/j.cell.2006.10.029

41 Lam, J. T., Moretti, A. & Laugwitz, K. L. Multipotent progenitor cells in regenerative
cardiovascular medicine. *Pediatr Cardiol* **30**, 690-698 (2009).
https://doi.org/10.1007/s00246-009-9450-1

42 Weinberger, F. *et al.* Localization of Islet-1-positive cells in the healthy and infarcted
adult murine heart. *Circ Res* **110**, 1303-1310 (2012).
https://doi.org/10.1161/CIRCRESAHA.111.259630

43 Kattman, S. J., Huber, T. L. & Keller, G. M. Multipotent flk-1+ cardiovascular progenitor
cells give rise to the cardiomyocyte, endothelial, and vascular smooth muscle lineages.
Dev Cell **11**, 723-732 (2006). https://doi.org/10.1016/j.devcel.2006.10.002

44 Motoike, T., Markham, D. W., Rossant, J. & Sato, T. N. Evidence for novel fate of Flk1+
progenitor: contribution to muscle lineage. *Genesis* **35**, 153-159 (2003).
https://doi.org/10.1002/gene.10175

45 Mekala, S. R. *et al.* Generation of Cardiomyocytes From Vascular Adventitia-Resident
Stem Cells. *Circ Res* **123**, 686-699 (2018).
https://doi.org/10.1161/CIRCRESAHA.117.312526

46 Zampetaki, A., Kirton, J. P. & Xu, Q. Vascular repair by endothelial progenitor cells.
Cardiovasc Res **78**, 413-421 (2008). https://doi.org/10.1093/cvr/cvn081

47 Gao, L. *et al.* Single-cell transcriptomics of cardiac progenitors reveals functional
subpopulations and their cooperative crosstalk in cardiac repair. *Protein Cell* **12**, 152-
157 (2021). https://doi.org/10.1007/s13238-020-00788-6

48 Masters, M. & Riley, P. R. The epicardium signals the way towards heart regeneration.
Stem Cell Res **13**, 683-692 (2014). https://doi.org/10.1016/j.scr.2014.04.007

49 Acharya, A. *et al.* The bHLH transcription factor Tcf21 is required for lineage-specific
EMT of cardiac fibroblast progenitors. *Development* **139**, 2139-2149 (2012).
https://doi.org/10.1242/dev.079970

50 Braitsch, C. M., Kanisicak, O., van Berlo, J. H., Molkentin, J. D. & Yutzey, K. E.
Differential expression of embryonic epicardial progenitor markers and localization of
cardiac fibrosis in adult ischemic injury and hypertensive heart disease. *J Mol Cell*
Cardiol **65**, 108-119 (2013). https://doi.org/10.1016/j.yjmcc.2013.10.005

51 Kanisicak, O. *et al.* Genetic lineage tracing defines myofibroblast origin and function in
the   injured   heart.   *Nat   Commun*   **7**,   12260   (2016).
https://doi.org/10.1038/ncomms12260

52 Armstrong, J. F., Pritchard-Jones, K., Bickmore, W. A., Hastie, N. D. & Bard, J. B. The
expression of the Wilms' tumour gene, WT1, in the developing mammalian embryo.
Mech Dev **40**, 85-97 (1993). https://doi.org/10.1016/0925-4773(93)90090-k

53 Carmona, R., Gonzalez-Iriarte, M., Perez-Pomares, J. M. & Munoz-Chapuli, R.
Localization of the Wilm's tumour protein WT1 in avian embryos. *Cell Tissue Res* **303**,
173-186 (2001). https://doi.org/10.1007/s004410000307

54 Duim, S. N., Kurakula, K., Goumans, M. J. & Kruithof, B. P. Cardiac endothelial cells
express Wilms' tumor-1: Wt1 expression in the developing, adult and infarcted heart.
J Mol Cell Cardiol **81**, 127-135 (2015). https://doi.org/10.1016/j.yjmcc.2015.02.007

55 Sanz-Morejon, A. *et al.* Wilms Tumor 1b Expression Defines a Pro-regenerative
Macrophage Subtype and Is Required for Organ Regeneration in the Zebrafish. *Cell*
Rep **28**, 1296-1306 e1296 (2019). https://doi.org/10.1016/j.celrep.2019.06.091

56 Tan, S. C. *et al.* Preconditioning of Cardiosphere-Derived Cells With Hypoxia or Prolyl-
4-Hydroxylase Inhibitors Increases Stemness and Decreases Reliance on Oxidative
Metabolism.   *Cell   Transplant*   **25**,   35-53   (2016).
https://doi.org/10.3727/096368915X687697

57 Zangrossi, S. *et al.* Oct-4 expression in adult human differentiated cells challenges its
role   as   a   pure   stem   cell   marker.   *Stem Cells* **25**, 1675-1680 (2007).
https://doi.org/10.1634/stemcells.2006-0611

58 Ghaleb, A. M. & Yang, V. W. Kruppel-like factor 4 (KLF4): What we currently know.
Gene **611**, 27-37 (2017). https://doi.org/10.1016/j.gene.2017.02.025

59 Ambady, S. *et al.* Expression of NANOG and NANOGP8 in a variety of undifferentiated
and   differentiated   human   cells.   *Int J Dev Biol* **54**, 1743-1754 (2010).
https://doi.org/10.1387/ijdb.103192sa

60 Piazzolla, D. *et al.* Lineage-restricted function of the pluripotency factor NANOG in
stratified   epithelia.   *Nat   Commun*   **5**,   4226   (2014).
https://doi.org/10.1038/ncomms5226

61 Sun, L. *et al.* Transient induction of telomerase expression mediates senescence and
reduces tumorigenesis in primary fibroblasts. *Proc Natl Acad Sci U S A* **116**, 18983-
18993 (2019). https://doi.org/10.1073/pnas.1907199116

62 Wang, G., Zhou, H., Gu, Z., Gao, Q. & Shen, G. Oct4 promotes cancer cell proliferation
and migration and leads to poor prognosis associated with the survivin/STAT3
pathway in hepatocellular carcinoma. *Oncol Rep* **40**, 979-987 (2018).
https://doi.org/10.3892/or.2018.6491

63 White, M. G. *et al.* Pluripotency-associated stem cell marker expression in proliferative
cell cultures derived from adult human pancreas. *J Endocrinol* **211**, 169-176 (2011).
https://doi.org/10.1530/JOE-11-0123

64 Cannarile, M. A. *et al.* Colony-stimulating factor 1 receptor (CSF1R) inhibitors in cancer
therapy. *J Immunother Cancer* **5**, 53 (2017). https://doi.org/10.1186/s40425-017-
0257-y

65 Plein, A., Fantin, A., Denti, L., Pollard, J. W. & Ruhrberg, C. Erythro-myeloid progenitors
contribute endothelial cells to blood vessels. *Nature* **562**, 223-228 (2018).
https://doi.org/10.1038/s41586-018-0552-x

66 Sinha, M. *et al.* Direct conversion of injury-site myeloid cells to fibroblast-like cells of
granulation tissue. *Nat Commun* **9**, 936 (2018). https://doi.org/10.1038/s41467-018-
03208-w

67 Feng, R. *et al.* PU.1 and C/EBPalpha/beta convert fibroblasts into macrophage-like
cells. *Proc Natl Acad Sci U S A* **105**, 6057-6062 (2008).
https://doi.org/10.1073/pnas.0711961105

68 Franzen, O., Gan, L. M. & Bjorkegren, J. L. M. PanglaoDB: a web server for exploration
of mouse and human single-cell RNA sequencing data. *Database (Oxford)* **2019** (2019).
https://doi.org/10.1093/database/baz046

69 Huebener, P. *et al.* CD44 is critically involved in infarct healing by regulating the
inflammatory and fibrotic response. *J Immunol* **180**, 2625-2633 (2008).
https://doi.org/10.4049/jimmunol.180.4.2625

70 Sneath, R. J. & Mangham, D. C. The normal structure and function of CD44 and its role
in neoplasia. *Mol Pathol* **51**, 191-200 (1998). https://doi.org/10.1136/mp.51.4.191

71 Suleiman, M., Abdulrahman, N., Yalcin, H. & Mraiche, F. The role of CD44, hyaluronan
and NHE1 in cardiac remodeling. *Life Sci* **209**, 197-201 (2018).
https://doi.org/10.1016/j.lfs.2018.08.009

72 Harris, E. S. & Nelson, W. J. VE-cadherin: at the front, center, and sides of endothelial
cell organization and function. *Curr Opin Cell Biol* **22**, 651-658 (2010).
https://doi.org/10.1016/j.ceb.2010.07.006

73 Pan, X., Chen, Z., Huang, R., Yao, Y. & Ma, G. Transforming growth factor beta1 induces
the expression of collagen type I by DNA methylation in cardiac fibroblasts. *PLoS One*
8, e60335 (2013). https://doi.org/10.1371/journal.pone.0060335

74 Sandri, T. L. *et al.* Human collectin-11 (COLEC11) and its synergic genetic interaction
with MASP2 are associated with the pathophysiology of Chagas Disease. *PLoS Negl*
Trop Dis **13**, e0007324 (2019). https://doi.org/10.1371/journal.pntd.0007324

75 Selman, L. & Hansen, S. Structure and function of collectin liver 1 (CL-L1) and collectin
11   (CL-11,   CL-K1).   *Immunobiology*   **217**,   851-863   (2012).
https://doi.org/10.1016/j.imbio.2011.12.008

76 Sehgal, A. *et al.* The role of CSF1R-dependent macrophages in control of the intestinal
stem-cell niche. *Nat Commun* **9**, 1272 (2018). https://doi.org/10.1038/s41467-018-
03638-6

77 Lee, M., Lee, Y., Song, J., Lee, J. & Chang, S. Y. Tissue-specific Role of CX(3)CR1
Expressing Immune Cells and Their Relationships with Human Disease. *Immune Netw*
18, e5 (2018). https://doi.org/10.4110/in.2018.18.e5

78 Zheng, J. *et al.* Chemokine receptor CX3CR1 contributes to macrophage survival in
tumor metastasis. *Mol Cancer* **12**, 141 (2013). https://doi.org/10.1186/1476-4598-12-
141

79 Leitinger, B. Discoidin domain receptor functions in physiological and pathological
conditions. *Int Rev Cell Mol Biol* **310**, 39-87 (2014). https://doi.org/10.1016/B978-0-
12-800180-6.00002-5

80 Valiathan, R. R., Marco, M., Leitinger, B., Kleer, C. G. & Fridman, R. Discoidin domain
receptor tyrosine kinases: new players in cancer progression. *Cancer Metastasis Rev*
31, 295-321 (2012). https://doi.org/10.1007/s10555-012-9346-z

81 Molkentin, J. D., Lin, Q., Duncan, S. A. & Olson, E. N. Requirement of the transcription
factor GATA4 for heart tube formation and ventral morphogenesis. *Genes Dev* **11**,
1061-1072 (1997). https://doi.org/10.1101/gad.11.8.1061

82 Watt, A. J., Battle, M. A., Li, J. & Duncan, S. A. GATA4 is essential for formation of the
proepicardium and regulates cardiogenesis. *Proc Natl Acad Sci U S A* **101**, 12573-12578
(2004). https://doi.org/10.1073/pnas.0400752101

83 Xin, M., Olson, E. N. & Bassel-Duby, R. Mending broken hearts: cardiac development
as a basis for adult heart regeneration and repair. *Nat Rev Mol Cell Biol* **14**, 529-541
(2013). https://doi.org/10.1038/nrm3619

84 Barnes, R. M. *et al.* Hand2 loss-of-function in Hand1-expressing cells reveals distinct
roles in epicardial and coronary vessel development. *Circ Res* **108**, 940-949 (2011).
https://doi.org/10.1161/CIRCRESAHA.110.233171

85   McFadden, D. G. *et al.* The Hand1 and Hand2 transcription factors regulate expansion
of the embryonic cardiac ventricles in a gene dosage-dependent manner.
Development **132**, 189-201 (2005). https://doi.org/10.1242/dev.01562

86   Thattaliyath, B. D., Livi, C. B., Steinhelper, M. E., Toney, G. M. & Firulli, A. B. HAND1
and HAND2 are expressed in the adult-rodent heart and are modulated during cardiac
hypertrophy. *Biochem Biophys Res Commun* **297**, 870-875 (2002).
https://doi.org/10.1016/s0006-291x(02)02297-0

87   Tsuchihashi, T. *et al.* Hand2 function in second heart field progenitors is essential for
cardiogenesis. *Dev Biol* **351**, 62-69 (2011).
https://doi.org/10.1016/j.ydbio.2010.12.023

88   Cai, C. L. *et al.* Isl1 identifies a cardiac progenitor population that proliferates prior to
differentiation and contributes a majority of cells to the heart. *Dev Cell* **5**, 877-889
(2003). https://doi.org/10.1016/s1534-5807(03)00363-0

89   Glasscock, E. Kv1.1 channel subunits in the control of neurocardiac function. *Channels*
(Austin) **13**, 299-307 (2019). https://doi.org/10.1080/19336950.2019.1635864

90   Miki, T. *et al.* Mouse model of Prinzmetal angina by disruption of the inward rectifier
Kir6.1. *Nat Med* **8**, 466-472 (2002). https://doi.org/10.1038/nm0502-466

91   Tester, D. J. *et al.* Loss-of-function mutations in the KCNJ8-encoded Kir6.1 K(ATP)
channel and sudden infant death syndrome. *Circ Cardiovasc Genet* **4**, 510-515 (2011).
https://doi.org/10.1161/CIRCGENETICS.111.960195

92   Zingman, L. V., Alekseev, A. E., Hodgson-Zingman, D. M. & Terzic, A. ATP-sensitive
potassium channels: metabolic sensing and cardioprotection. *J Appl Physiol (1985)*
103, 1888-1893 (2007). https://doi.org/10.1152/japplphysiol.00747.2007

93   Simons, M., Gordon, E. & Claesson-Welsh, L. Mechanisms and regulation of
endothelial VEGF receptor signalling. *Nat Rev Mol Cell Biol* **17**, 611-625 (2016).
https://doi.org/10.1038/nrm.2016.87

94   Chiplunkar, A. R. *et al.* The Kruppel-like factor 2 and Kruppel-like factor 4 genes
interact to maintain endothelial integrity in mouse embryonic vasculogenesis. *BMC*
Dev Biol **13**, 40 (2013). https://doi.org/10.1186/1471-213X-13-40

95   Zhang, P., Andrianakos, R., Yang, Y., Liu, C. & Lu, W. Kruppel-like factor 4 (Klf4)
prevents embryonic stem (ES) cell differentiation by regulating Nanog gene
expression. *J Biol Chem* **285**, 9180-9189 (2010).
https://doi.org/10.1074/jbc.M109.077958

96   Zhang, Y. *et al.* Kruppel-like factor 4 transcriptionally regulates TGF-beta1 and
contributes to cardiac myofibroblast differentiation. *PLoS One* **8**, e63424 (2013).
https://doi.org/10.1371/journal.pone.0063424

97   Stolzing, A. & Grune, T. Neuronal apoptotic bodies: phagocytosis and degradation by
primary microglial cells. *FASEB J* **18**, 743-745 (2004). https://doi.org/10.1096/fj.03-
0374fje

98   Zhang, H., Chen, X. & Sairam, M. R. Novel genes of visceral adiposity: identification of
mouse and human mesenteric estrogen-dependent adipose (MEDA)-4 gene and its
adipogenic function. *Endocrinology* **153**, 2665-2676 (2012).
https://doi.org/10.1210/en.2011-2008

99   Subramanian, S. V. & Nadal-Ginard, B. Early expression of the different isoforms of the
myocyte enhancer factor-2 (MEF2) protein in myogenic as well as non-myogenic cell
lineages during mouse embryogenesis. *Mech Dev* **57**, 103-112 (1996).
https://doi.org/10.1016/0925-4773(96)00542-4

100 Wang, L., Fan, C., Topol, S. E., Topol, E. J. & Wang, Q. Mutation of MEF2A in an
inherited disorder with features of coronary artery disease. *Science* **302**, 1578-1581
(2003). https://doi.org/10.1126/science.1088477

101 Bi, W., Drake, C. J. & Schwarz, J. J. The transcription factor MEF2C-null mouse exhibits
complex vascular malformations and reduced cardiac expression of angiopoietin 1 and
VEGF. *Dev Biol* **211**, 255-267 (1999). https://doi.org/10.1006/dbio.1999.9307

102 Lin, Q., Schwarz, J., Bucana, C. & Olson, E. N. Control of mouse cardiac morphogenesis
and myogenesis by transcription factor MEF2C. *Science* **276**, 1404-1407 (1997).
https://doi.org/10.1126/science.276.5317.1404

103 Kitajima, S., Takagi, A., Inoue, T. & Saga, Y. MesP1 and MesP2 are essential for the
development of cardiac mesoderm. *Development* **127**, 3215-3226 (2000).
https://doi.org/10.1242/dev.127.15.3215

104 Komata, M., Bando, M., Araki, H. & Shirahige, K. The direct binding of Mrc1, a
checkpoint mediator, to Mcm6, a replication helicase, is essential for the replication
checkpoint against methyl methanesulfonate-induced stress. *Mol Cell Biol* **29**, 5008-
5019 (2009). https://doi.org/10.1128/MCB.01934-08

105 Staines, K., Hunt, L. G., Young, J. R. & Butter, C. Evolution of an expanded mannose
receptor gene family. *PLoS One* **9**, e110330 (2014).
https://doi.org/10.1371/journal.pone.0110330

106 Roszer, T. Understanding the Mysterious M2 Macrophage through Activation Markers
and Effector Mechanisms. *Mediators Inflamm* **2015**, 816460 (2015).
https://doi.org/10.1155/2015/816460

107 Eon Kuek, L., Leffler, M., Mackay, G. A. & Hulett, M. D. The MS4A family: counting past
1, 2 and 3. *Immunol Cell Biol* **94**, 11-23 (2016). https://doi.org/10.1038/icb.2015.48

108 Martinez, F. O., Gordon, S., Locati, M. & Mantovani, A. Transcriptional profiling of the
human monocyte-to-macrophage differentiation and polarization: new molecules and

patterns of gene expression. *J Immunol* **177**, 7303-7311 (2006).
https://doi.org/10.4049/jimmunol.177.10.7303

109 England, J. & Loughna, S. Heavy and light roles: myosin in the morphogenesis of the
heart. *Cell Mol Life Sci* **70**, 1221-1239 (2013). https://doi.org/10.1007/s00018-012-
1131-1

110 Gawlik-Rzemieniewska, N. & Bednarek, I. The role of NANOG transcriptional factor in
the development of malignant phenotype of cancer cells. *Cancer Biol Ther* **17**, 1-10
(2016). https://doi.org/10.1080/15384047.2015.1121348

111 Ampofo, E., Schmitt, B. M., Menger, M. D. & Laschke, M. W. The regulatory
mechanisms of NG2/CSPG4 expression. *Cell Mol Biol Lett* **22**, 4 (2017).
https://doi.org/10.1186/s11658-017-0035-3

112 Fukushi, J., Makagiansar, I. T. & Stallcup, W. B. NG2 proteoglycan promotes
endothelial cell motility and angiogenesis via engagement of galectin-3 and
alpha3beta1 integrin. *Mol Biol Cell* **15**, 3580-3590 (2004).
https://doi.org/10.1091/mbc.e04-03-0236

113 Goretzki, L., Burg, M. A., Grako, K. A. & Stallcup, W. B. High-affinity binding of basic
fibroblast growth factor and platelet-derived growth factor-AA to the core protein of
the NG2 proteoglycan. *J Biol Chem* **274**, 16831-16837 (1999).
https://doi.org/10.1074/jbc.274.24.16831

114 Tanaka, M., Chen, Z., Bartunkova, S., Yamasaki, N. & Izumo, S. The cardiac homeobox
gene Csx/Nkx2.5 lies genetically upstream of multiple genes essential for heart
development. *Development* **126**, 1269-1280 (1999).
https://doi.org/10.1242/dev.126.6.1269

115 Song, W., Wang, H. & Wu, Q. Atrial natriuretic peptide in cardiovascular biology and
disease (NPPA). *Gene* **569**, 1-6 (2015). https://doi.org/10.1016/j.gene.2015.06.029

116 Pan, G. J., Chang, Z. Y., Scholer, H. R. & Pei, D. Stem cell pluripotency and transcription
factor Oct4. *Cell Res* **12**, 321-329 (2002). https://doi.org/10.1038/sj.cr.7290134

117 Zeineddine, D., Hammoud, A. A., Mortada, M. & Boeuf, H. The Oct4 protein: more than
a magic stemness marker. *Am J Stem Cells* **3**, 74-82 (2014).

118 El-Rass, S. *et al.* Disruption of pdgfra alters endocardial and myocardial fusion during
zebrafish cardiac assembly. *Biol Open* **6**, 348-357 (2017).
https://doi.org/10.1242/bio.021212

119 Horikawa, S. *et al.* PDGFRalpha plays a crucial role in connective tissue remodeling. *Sci*
Rep **5**, 17948 (2015). https://doi.org/10.1038/srep17948

120 Kim, J. *et al.* PDGF signaling is required for epicardial function and blood vessel
formation in regenerating zebrafish hearts. *Proc Natl Acad Sci U S A* **107**, 17206-17210
(2010). https://doi.org/10.1073/pnas.0915016107

121 Betsholtz, C. Insight into the physiological functions of PDGF through genetic studies
in mice. *Cytokine Growth Factor Rev* **15**, 215-228 (2004).
https://doi.org/10.1016/j.cytogfr.2004.03.005

122 Hsieh, P. C., Davis, M. E., Gannon, J., MacGillivray, C. & Lee, R. T. Controlled delivery
of PDGF-BB for myocardial protection using injectable self-assembling peptide
nanofibers. *J Clin Invest* **116**, 237-248 (2006). https://doi.org/10.1172/JCI25878

123 Peng, Y., Yan, S., Chen, D., Cui, X. & Jiao, K. Pdgfrb is a direct regulatory target of
TGFbeta signaling in atrioventricular cushion mesenchymal cells. *PLoS One* **12**,
e0175791 (2017). https://doi.org/10.1371/journal.pone.0175791

124 Van den Akker, N. M. *et al.* PDGF-B signaling is important for murine cardiac
development: its role in developing atrioventricular valves, coronaries, and cardiac
innervation. *Dev Dyn* **237**, 494-503 (2008). https://doi.org/10.1002/dvdy.21436

125 Wilkinson, S., O'Prey, J., Fricker, M. & Ryan, K. M. Hypoxia-selective macroautophagy
and cell survival signaled by autocrine PDGFR activity. *Genes Dev* **23**, 1283-1288
(2009). https://doi.org/10.1101/gad.521709

126 Yue, Z. *et al.* PDGFR-beta Signaling Regulates Cardiomyocyte Proliferation and
Myocardial Regeneration. *Cell Rep* **28**, 966-978 e964 (2019).
https://doi.org/10.1016/j.celrep.2019.06.065

127 Grossi, S. *et al.* Molecular genetic analysis of the PLP1 gene in 38 families with PLP1-
related disorders: identification and functional characterization of 11 novel PLP1
mutations. *Orphanet J Rare Dis* **6**, 40 (2011). https://doi.org/10.1186/1750-1172-6-40

128 Alquraini, A. *et al.* The autocrine role of proteoglycan-4 (PRG4) in modulating
osteoarthritic synoviocyte proliferation and expression of matrix degrading enzymes.
Arthritis Res Ther **19**, 89 (2017). https://doi.org/10.1186/s13075-017-1301-5

129 Han, L., Grodzinsky, A. J. & Ortiz, C. Nanomechanics of the Cartilage Extracellular
Matrix. *Annu Rev Mater Res* **41**, 133-168 (2011). https://doi.org/10.1146/annurev-
matsci-062910-100431

130 Dawes, R. *et al.* Combinations of CD45 isoforms are crucial for immune function and
disease. *J Immunol* **176**, 3417-3425 (2006).
https://doi.org/10.4049/jimmunol.176.6.3417

131 Hermiston, M. L., Xu, Z. & Weiss, A. CD45: a critical regulator of signaling thresholds in
immune cells. *Annu Rev Immunol* **21**, 107-137 (2003).
https://doi.org/10.1146/annurev.immunol.21.120601.140946

132   Nakano, A., Harada, T., Morikawa, S. & Kato, Y. Expression of leukocyte common
868   antigen (CD45) on various human leukemia/lymphoma cell lines. *Acta Pathol Jpn* **40**,
869   107-115 (1990). https://doi.org/10.1111/j.1440-1827.1990.tb01549.x

870 133   Batts, T. D. *et al.* Stem cell antigen-1 (sca-1) regulates mammary tumor development
871   and cell migration. *PLoS One* **6**, e27841 (2011).
872   https://doi.org/10.1371/journal.pone.0027841

873 134   Bradfute, S. B., Graubert, T. A. & Goodell, M. A. Roles of Sca-1 in hematopoietic
874   stem/progenitor cell function. *Exp Hematol* **33**, 836-843 (2005).
875   https://doi.org/10.1016/j.exphem.2005.04.001

876 135   Ito, C. Y., Li, C. Y., Bernstein, A., Dick, J. E. & Stanford, W. L. Hematopoietic stem cell
877   and progenitor defects in Sca-1/Ly-6A-null mice. *Blood* **101**, 517-523 (2003).
878   https://doi.org/10.1182/blood-2002-06-1918

879 136   Upadhyay, G. *et al.* Stem cell antigen-1 enhances tumorigenicity by disruption of
880   growth differentiation factor-10 (GDF10)-dependent TGF-beta signaling. *Proc Natl*
881   *Acad Sci U S A* **108**, 7820-7825 (2011). https://doi.org/10.1073/pnas.1103441108

882 137   Tandon, P., Miteva, Y. V., Kuchenbrod, L. M., Cristea, I. M. & Conlon, F. L. Tcf21
883   regulates the specification and maturation of proepicardial cells. *Development* **140**,
884   2409-2421 (2013). https://doi.org/10.1242/dev.093385

885 138   Mattson, M. P., Fu, W. & Zhang, P. Emerging roles for telomerase in regulating cell
886   differentiation and survival: a neuroscientist's perspective. *Mech Ageing Dev* **122**, 659-
887   671 (2001). https://doi.org/10.1016/s0047-6374(01)00221-4

888 139   Yuan, X., Larsson, C. & Xu, D. Mechanisms underlying the activation of TERT
889   transcription and telomerase activity in human cancer: old actors and new players.
890   *Oncogene* **38**, 6172-6183 (2019). https://doi.org/10.1038/s41388-019-0872-9

891 140   Peyvandi, F., Garagiola, I. & Baronciani, L. Role of von Willebrand factor in the
892   haemostasis. *Blood Transfus* **9 Suppl 2**, s3-8 (2011).
893   https://doi.org/10.2450/2011.002S

894 141   Lu, D. *et al.* WIF1 causes dysfunction of heart in transgenic mice. *Transgenic Res* **22**,
895   1179-1189 (2013). https://doi.org/10.1007/s11248-013-9738-z

896 142   Smits, A. M., Dronkers, E. & Goumans, M. J. The epicardium as a source of multipotent
897   adult cardiac progenitor cells: Their origin, role and fate. *Pharmacol Res* **127**, 129-140
898   (2018). https://doi.org/10.1016/j.phrs.2017.07.020

899 143   Zhao, L. *et al.* Notch signaling regulates cardiomyocyte proliferation during zebrafish
900   heart regeneration. *Proc Natl Acad Sci U S A* **111**, 1403-1408 (2014).
901   https://doi.org/10.1073/pnas.1311705111

902   144   Schildmeyer, L. A. *et al.* Impaired vascular contractility and blood pressure
903   homeostasis in the smooth muscle alpha-actin null mouse. *FASEB J* **14**, 2213-2220
904   (2000). https://doi.org/10.1096/fj.99-0927com