Title:

**Differential expression of sex regulatory genes in gonads of *Astyanax mexicanus* surface fish**

**and cavefish**

**Authors:**

Kaitlyn A. Webster[1], Bethany Ponte[2], Hans Vasquez-Gross[3], Juli Petereit[3], John Hutchinson[4],

Misty R. Riddle[2*]

**Affiliations:**

1. Harvard Medical School, Boston, MA, USA

2. University of Nevada, Reno, Reno, Nevada, USA

3. Nevada Bioinformatics Center (RRID:SCR_017802), University of Nevada, Reno, Reno,

Nevada, USA

4. Harvard T.H. Chan School of Public Health, Boston, MA, USA

* Corresponding author: mistyriddle@unr.edu

**Abstract**

**Background:** *Astyanax mexicanus* is a single species of fish that consists of river-dwelling

(surface) and cave-dwelling morphotypes. Little is known about how sexual determination,

differentiation or reproduction have evolved in the surface morphs or cavefish, though

divergence in reproductive strategy is expected as the latter have adapted to the novel cave

environment. Evolution of the gonad transcriptome may underlie the differences in gamete

morphology, fertility, and fecundity previously reported between morphotypes.

**Results:** We compared the ovary and testis transcriptome of surface fish and cavefish at juvenile

and adult stages. We found that samples clustered by developmental stage, sex, and

morphotype identity. Several key genes that are typically associated with the female gonad in

other vertebrates showed a reversal in sexual dimorphism or were not differentially expressed

between sexes in *A. mexicanus*. In contrast, while gene expression typically associated with

male gonads was largely conserved and consistent with vertebrate testicular expression profiles.

Transcriptional and physiological differences between surface fish and cavefish morphotypes

were observed in gonads from both sexes. Cavefish ovaries exhibited unique upregulation of

neuron development and differentiation genes, and extensive innervation of the ovarian

epithelium, while cavefish testes showed increased expression of angiogenesis regulating genes,

and greater vasculature density compared to surface fish testes.

**Conclusions:** These results reveal significant gene expression differences between *A. mexicanus*

surface fish and cavefish morphotypes that may have functional consequences in gonad

morphogenesis and fertility. Our findings provide a foundation for investigating the evolution of

sex regulatory pathways and reproductive strategies in animals adapting to new and challenging

environments in which nutrient availability, temperature, and mate selection are suboptimal.

**Keywords:** *Astyanax mexicanus*, gonad, sex differentiation, transcriptome, ovary, testis, *foxl2*,

*cyp19a1a*, teleost

**Background**

Studying gonadal sex differentiation in fishes can help identify novel molecular pathways

that mediate sex determination and development. In contrast to mammals, which universally

utilize chromosomal XY sex determination directed by the *SOX3*-derived *SRY* gene, fishes display

a diversity of sex determining mechanisms (SDM) and sex regulatory genes. Across teleosts,

SDM are known to be plastic, rapidly evolving, and inclusive of all known modes of regulation:

chromosomal, polygenic, hormonal, social, and environmental (Devlin and Nagahama 2002;

Schartl 2004).  Different SDM and master sex regulatory genes can evolve even within the same

genus, as reported in closely related species of Medaka rice fish (Matsuda 2002, Myosho 2012,

Takehana 2014). Like other non-mammalian vertebrates, teleost fish sex determination can be

influenced and overridden by social dynamics and environmental conditions, even when sex

chromosomes or sex regulatory genes are employed. Notably, environmental factors such as

photoperiod, temperature, salinity, pH and hypoxia have been shown to skew sex ratios in

teleosts (Brown 2014, Baroiller 2009, Abucay 1999, Shang 2006). Despite the diversity of SDM,

most vertebrate sex regulatory genes have orthologues in teleosts, which have been well

studied in zebrafish, Medaka, and tilapia. For example, the mammalian ovary-promoting genes

*Foxl2*, *Wnt4*, and *Cyp19* (aromatase) show conserved female-specific function in teleosts, as do

the testis-promoting genes *dmrt1*, *amh*, and *sox9* (Yang 2017, Kossack 2017, Lau 2016, Guigen

2010, Webster 2017, Yan 2019, Nakamura 2012). How sex differentiation and gonad

development evolve as fishes adapt to new environments is not well understood.

The Mexican tetra, *Astyanax mexicanus,* is a species of fish that consists of river-adapted

and cave-adapted morphotypes. There are as many as 32 geographically separate cave-dwelling

"cavefish" populations that evolved from river-dwelling "surface fish" in the underground

limestone caves of the El Abra Region of Tamaulipas and San Luis Potosí, Mexico (Figure 1D)

(Espinasa et al., 2020). Based on whole genome sequencing of field-collected individuals, the *A.*

*mexicanus* phylogeny defines two lineages; cavefish from the Tinaja and Pachón caves form a

monophyletic clade separate from the Molino cavefish and Río Choy surface fish (Herman et al.,

2018). This evolutionary history indicates that cavefish traits have evolved through repeated

evolution providing researchers with natural replicates to investigate the genetic and

developmental basis of cavefish traits.

Little is known about sex and gonad development in *A. mexicanus* despite its growing

importance as a model system in understanding the genetic basis of evolution (Perera 2023).

For example, only a few studies have compared the expression or function of sex-associated

genes between surface fish and cavefish morphs. In Pachón cavefish, sex-associated genes such

as *foxl2a*, *wnt4b*, *cyp19a1a*, *dmrt1*, and *amh* were reported to be expressed in the gonads but

surprisingly *foxl2a*, *wnt4b*, and *cyp19a1a* were not found to be significantly sexually dimorphic

(Imarazene 2021a). Pachón cavefish male sex is determined by supernumerary "B-sex"

chromosomes that exhibit non-mendelian inheritance (Imarazene et al., 2021b). The gene

conferring male sex on the Pachón cavefish B chromosome is *growth differentiation factor 6b*

(*gdf6b*) which if knocked out results in male-to-female sex reversals (Imarazene et al., 2021b).

The mechanisms of sex determination in surface fish and other cavefish populations have yet to

be described. The atypical expression of conserved vertebrate sex-specific genes and

emergence of a novel sex determination mechanism in Pachón cavefish highlights the promise

of this model for the study of evolution of sex regulatory pathways and adaptive reproductive

strategies. In addition, since traits like body condition and relative organ size are sexually

dimorphic and are controlled by loci that have sex-dependent effects (Riddle et al., 2021),

revealing how sex determination and gonad development differs between *A. mexicanus*

morphs is crucial to understanding cavefish evolution.

Here, we describe the first transcriptomic analysis of ovaries and testis in *A. mexicanus,*

comparing surface fish and Pachón cavefish at juvenile and adult stages. We found that samples

clustered by developmental stage, sex, and morphotype identity. We found that in surface fish

and Pachón cavefish, typical female-associated genes are not differentially expressed between

males and females. However, typical male-associated gene expression is conserved. We

observed an upregulation of genes involved in neuron development in Pachón cavefish ovaries

compared to surface fish ovaries and found that the germinal epithelium surrounding germ

cells is extensively innervated in both morphotypes. Comparing the testis, we observed

increased expression of angiogenesis genes in Pachón cavefish compared to surface fish and

evidence of more vasculature within the Pachón cavefish testis.  Our results provide insight into

the molecular, functional, and structural differences between surface fish and cavefish gonads

which is essential for future studies examining the evolution of sex differentiation, sexual

dimorphism, and reproduction in this model system.

**Methods**

*Animal husbandry*

Río Choy surface fish and Pachón cavefish were derived from parents bred for greater

than five generations in the laboratory of Dr. Clifford Tabin at Harvard Medical School. All

animals used in this study were maintained, housed, and fed under the same conditions on a

recirculating water system (Temperature: 23-24 °C, pH: 7–7.5, Conductivity: 1200-1400µS) in a

facility with a 10:14-h light/dark cycle. Adult fish were fed with a combination of New Life

Spectrum TheraA+ small and/or medium sinking pellets (dependent on fish size)

and *Artemia* and housed at a density equal to two adult fish per liter of water. Hatched larval

fish were fed rotifers up to 14 days post fertilization (dpf) in 1-liter cups of fish-ready water,

then transferred to the recirculating system where they were fed *Artemia* through 60 dpf. At

1.5 years post fertilization, fish were euthanized in MS-222 prior to gonad dissection. We found

that adult Pachón cavefish males and females were on average larger than surface fish

counterparts with larger gonads, but did not observe significant differences in size between

morphotypes (Pachón cavefish versus surface fish average fish weight: male = 1.34g vs 0.95g, p

= 0.09, female = 1.9g vs 1.2g, p = 0.17, t-test, Pachón cavefish versus surface fish average gonad

weight: testes = 0.034g vs 0.024g, p = 0.40, ovaries = 0.165g vs .097g, p = 0.46, Additional file 1).

*Sample collection, RNA extraction and cDNA synthesis*

Juvenile (i.e., not containing ovulatory eggs or caudal duct spermatozoa) to adult fish aged

between 1-1.5 years were euthanized in 400ppm MS-222. The gonads were dissected into cold

1 X PBS, and thoroughly cleared of associated fat or viscera using a dissecting microscope.

Cleaned gonads were then collected into 0.5mL (adult gonads) or 0.25mL (juvenile gonads)

Trizol and homogenized using a hand-held pellet pestle, then stored at –80°C. Total RNA was

extracted using Zymo Research Direct-zol RNA MicroPrep with DNAse treatment according to

the manufacturers protocol. LunaScript RT Supermix kit with 1μg of RNA was used to synthesize

cDNA. Diluted cDNA samples (50ng/μL) were used for sequencing.

*RNA sequencing and gene ontology analysis*

Illumina sequencing was performed by the Nevada Genomics Center using NextSeq 2000 P3 100

cycle sequencing kit. Samples were sequenced with 50bp paired-end reads with a mean of 52.2

million reads (minimum depth of 29.6 to 58.8 million reads per sample). FastQC v0.11.9 was ran

on each sequenced sample both pre and post read trimming to assess improvement after

trimming. Trimmomatic v0.36 with parameters ILLUMINACLIP:2:30:10 SLIDINGWINDOW:4:15

MINLEN:36 was used in order to remove Illumina sequencing adapters and poor base quality

regions.

Reads were aligned to Ensembl build 2 of the *A. mexicanus* genome, augmented with transcript

information from Ensembl release 2.0.97 using STAR v2.7.5c (Dobin et al., 2013). STAR output

alignment BAM files were imported into featureCounts v2.0 from the subread package in order

to quantify reads per sample for genes. Quantified counts generated via featureCounts were

imported into the statistical software package R (R-core-team, 2019) for downstream analysis. A

low count was defined by less than 10 counts per gene. If a gene demonstrated low counts

across all samples and experimental groups, it was removed from further downstream analysis.

Differential gene analysis was performed with the DESeq2 package (Love 2014) and statistically

significant differences were defined by a Benjamini-Hochberg FDR adjusted *p*-value <0.05.

Variance stabilizing transformation (VST) was used to visualize count data in a Principal

Component Analysis plot.

To gain more insight into the differentially expressed genes, gene names were annotated using

the biomaRt package (Durinck 2005), and we determined enriched Biological Processes Gene

Ontologies (BP GO) using clusterProfiler (Yu 2012). Gene ontology maps were generated from

enriched Biological Processes GO terms for differentially expressed genes (padj < 0.05) using

clusterProfiler cnetplot function (Yu 2012). Annotation hub AH107459 was used for *Astyanax*

*mexicanus* GO term annotation.

*Ovary immunostaining*

Ovaries from adult Pachón cavefish and surface fish were dissected, fixed in 4% PFA, and

dehydrated in ethanol. After rehydration, the tissue was incubated with α-acetylated tubulin

(1:500, Sigma-Alrich, T6793) in blocking solution (BSA, sodium azide, goat serum, DMSO, PBS,

and Triton X-100) for 48 hours at room temperature, rinsed and incubated with a secondary

antibody (goat anti-mouse 488 IgG H&L, ABCam, ab150117) for another 48 hours at room

temperature. The tissue was imaged on a Leica THUNDER model organism microscope.

*Testes Vascular Staining and Quantification*

Testes were dissected from adult Pachón cavefish and surface fish and collected into 1 x PBS at

room temperature, briefly. Whole testes were then immediately incubated for 5 minutes in the

dark in a solution of o-Dianisidine (3,3▢-dimethoxybenzidine, MilliporeSigma D9143-5G)

formulated as described in O'Brien 2009. Stained testes were dehydrated, cleared in xylene, and

embedded in paraffin. Sections of 10 uM were then counterstained with modified Harris

hematoxylin solution (Sigma-Aldrich HHS32). Tissue area was calculated using ImageJ. Each

bundle of vascular staining was traced, and the area was recorded. The vascular area was

summed and divided by the total tissue area in the image to calculate percentage of testicular

vascularization.

**Results**

We compared the ovary and testes transcriptome of *A. mexicanus* surface fish and

Pachón cavefish that were 14-18 months post fertilization (Figure 1 A-E). Interestingly,

regardless of chronological age, two distinct morphological stages of gonads were observed for

each sex, referred to in this study as juvenile (not containing mature gametes) or adult

(containing mature gametes). Gonads categorized as juvenile appeared small and translucent

upon dissection, and by histology were confirmed to lack ovulatory oocytes (females) or

spermatozoa (males). Adult gonads were large and opaque, and stored mature gametes in the

urogenital ducts. Overall, the gonad morphology was similar comparing surface fish and Pachón

cavefish, with the most striking difference observed in coloration. Surface testes were sheathed

in scattered black melanocytes (Figure 1B), whereas Pachón testes lacked any melanin

pigmentation consistent with albinism (Protas 2006) (Figure 1C). Conversely, Pachón cavefish

ovaries were intensely yellow-orange compared to surface fish ovaries, consistent with

increased carotenoid accumulation in the eggs (Riddle 2020) (Figure 1E).

We first visualized transcriptome differences between sample types (n=3 per

developmental stage, sex, and morphotype) using principal component analysis and found that

samples clustered according to sex and morphotype identity (Figure 1J). We found that PC1

(57% variance) separates testes from ovaries, while PC2 (27% variance) separates surface fish

from Pachón cavefish. Juvenile and adult gonads formed overlapping clusters except for surface

fish ovaries that showed no overlap between developmental stages.  These findings indicate

differential regulation of gene expression between *A. mexicanus* morphotypes within the

gonads of males and females.

We examined the gene expression profiles that were specific to each sex to investigate

if the function of known regulators of sex differentiation in other species are likely conserved in

*A. mexicanus.* We found 17,955 differentially expressed genes (DEG) between the testes and

ovaries of *A. mexicanus* (Figure 2A). We observed expected sexually dimorphic expression of

some highly conserved and commonly studied sex regulatory genes, but also found that several

genes which are customarily restricted to the ovary or upregulated in the ovary, as best

characterized in the zebrafish model, are not sexually dimorphic (Figure 2B). For example, the

germ-plasm aggregator bucky ball (*buc*) is more highly expressed in the ovary but is also

unexpectedly abundant in the testes which do not express *buc* in other teleosts (Figure 2J).

Other ovary-associated genes such as the estrogen synthesis enzyme aromatase (*cyp19a1a*),

the gonadal somatic cell transcription factor Forkhead box protein L2 (*foxl2a*), and the secreted

growth factor wingless-type MMTV integration site family, member 4 (*wnt4a*) were either not

sexually dimorphic in the gonads (*cyp19a1a*, Figure 2C), or exhibit a sex-reversed expression

profile, i.e. were unexpectedly more highly expressed in the testes (*foxl2a* and *wnt4a*, Figure

2D-E). These results suggest possible neofunctionalization of these sex regulators in *A.*

*mexicanus.* In contrast to female-promoting genes, male-promoting genes displayed expected

patterns of expression comparing ovaries and testes. For example, the TGF-β superfamily

member anti-Müllerian hormone (*amh* Figure 2F), the androgen synthesis enzyme Cytochrome

P450, family 11 (*cyp11c1*, Figure 2G), the doublesex and mab-3 related transcription factor 1

(*dmrt1*, Figure 2H), and the teleost-specific TGF-β superfamily member gonadal soma derived

factor (*gsdf*, Figure 2I) were upregulated in the testes of both Pachón cavefish and surface fish.

We next compared the ovary transcriptome between surface fish and cavefish

morphotypes (Figure 3). Comparing juveniles, we observed 7,312 DEG between morphotypes

(Figure 3A). Interestingly, the gene that is most statistically significantly upregulated comparing

juvenile Pachón cavefish versus surface fish ovaries was *spock2* (Testican-2), which is a calcium-

binding proteoglycan that mediates oocyte maturation in mice. Comparing adults, we found

3,698 DEG (Figure 3B). We found that the top 3 gene ontology (GO) term categories for genes

upregulated in Pachón juvenile ovaries are protein phosphorylation, neuron development, and

plasma membrane bounded cell projection organization (Figure 3C).  Similarly, in Pachón adult

ovaries the top categories are nervous system development, neuron differentiation, and

generation of neurons (Figure 3D). *Semaphorin 3aa* (*sema3aa*) is one of the most upregulated

genes in Pachón ovaries of both stages and is implicated in the migration of specialized neurons

that secrete gonadotropin-releasing hormone (GnRH) – a critical mediator of puberty and

subsequent reproductive cycling (Giacobini 2015). Overall, these findings indicate an

upregulation of neuron-associated genes in the ovaries of Pachón cavefish compared to surface

fish.

In line with a role for neurons in the regulation of ovary function, we observed

substantial innervation throughout the sheath that surrounds the ovary, the geminal

epithelium, in both morphotypes (Figure 3F-H). A large neuron trunk that branches to reach

throughout the ovary germinal epithelium was visible on medial side of the ovary while the

lateral side of the ovary shows little to no innervation. A similar innervation pattern is observed

in zebrafish and is thought to be important to regulate oocyte quiescence through

norepinephrine signaling (Kim 2021). Our findings suggest that altered neuronal signaling may

be important for the evolution of cavefish ovaries.

We next compared the testes transcriptome between surface fish and cavefish

morphotypes (Figure 5). Interestingly, the testes transcriptome was more similar between

morphotypes than the ovary transcriptome. We found 1,505 DEG comparing juvenile testes and

4,038 DEG comparing adult testes of Pachón cavefish versus surface fish (Figure 4A and 4B,

respectively). Many of the most highly expressed DEG in either morph are unannotated, novel

genes. However, the most statistically significantly upregulated gene in adult Pachón testes was

Syntaxin 5a-like (*stx5al*), which is expressed in mammalian Sertoli cells (somatic nurse cells of

the testicular germ cells), but has also been reported as a primordial germ cell marker in

zebrafish single-cell RNAseq atlas Daniocell (Sur 2023). Gene set enrichment analysis of the

genes upregulated in Pachón testes revealed Biological Process GO term categories related to

angiogenesis in juveniles, and cell-cell signaling in adult testes (Figure 4C-D), likely representing

morphogenesis/patterning and sperm surface interactants, respectively. Consistent with these

gene expression differences, we found that in surface fish testes, blood vessels or vascular

bundles were scant, while in Pachón testes most testicular tubules were associated peripherally

with vascular tissue (Figure 4D). Our results at the transcriptional and morphological levels,

suggest increased vascularization of the cavefish testes which could have an impact on the

regulation of spermatogenesis.

**Discussion**

In this study, we present the first transcriptomic analysis of ovaries and testes in

*Astyanax mexicanus,* comparing gene expression between lab-raised surface fish and Pachón

cavefish at both juvenile and adult stages. Our results reveal clustering of samples based on

developmental stage, sex, and morphotype identity, demonstrating that these factors play

significant roles in shaping gonadal gene expression. This clustering supports the idea that

despite the shared ancestry of surface and cave morphotypes, their reproductive systems

exhibit distinct developmental trajectories resulting from evolution under different

environmental pressures. This transcriptome will enable the identification and characterization

of novel and conserved sex-specific genes that drive gonadal sex differentiation in juveniles and

maintain sex fate determinations in adults, which in teleost fish requires both male pathway

promotion and sustained female pathway suppression (Nishimura and Tanaka 2014). These

findings also highlight the potential for *A. mexicanus* to serve as a model for studying the rapid

evolution of highly conserved vertebrate sex regulating genes, especially in the context of

extreme environmental and nutritional deficits.

***Conservation and deviation in expression profiles of conserved sex genes***

Broadly compared to other male vertebrates, our *A. mexicanus* gonadal transcriptome

establishes a high degree of conserved gene expression associated with testicular

differentiation and spermatogenesis. All male gonad-promoting candidates investigated were

expressed uniquely in, or significantly upregulated in testes of both morphotypes, including

transcription factors (*dmrt1, sox9a*), signaling molecules (*amh*), and steroidogenic enzymes

(*cyp11c1*). These results are consistent with expression profiles reported in other fish testis

transcriptomes, and furthermore indicate homology of testicular cell subtypes relative to other

vertebrates, including humans (Nynca 2022, Qian 2022, Liu 2023). However, despite typical

expression of testis-enriched transcripts, our data show a striking deviation from other fishes or

vertebrates with respect to the expression of canonically ovary-promoting genes. This suggests

possible evolution, differential selection, or neofunctionalization of ovarian genes in *A.*

*mexicanus*.

Most strikingly, we identified a reversal of sexually dimorphic expression in *foxl2a*, an

evolutionarily conserved transcription factor produced by somatic cells of the differentiating

ovary in all metazoans. Across species, *foxl2a*/Foxl2a is typically an ovary-enriched transcript

and protein. Transcript levels of *foxl2a* will not necessarily correlate to protein products active

in the testis, however, highlighting the importance of further experiments. Foxl2 has a

complicated history of repeated gains, losses, and role changing, but generally is present in fish

genomes as *foxl2a* or *foxl2b*, with some basal teleosts retaining both *foxl2a* and *foxl2b* (Bertho

2016). We identified *foxl2a*, *foxl2b*, and the paralog *foxl3* in the *A. mexicanus* ovary and testis

transcriptomes. Surprisingly, our data show that *foxl2b* is more highly expressed in ovaries than

testes, despite its sequence being highly divergent from other teleosts exhibiting male-enriched

or non-dimorphic expression of this gene. While *A. mexicanus foxl2a* shows 75-80% identity to

its teleost orthologs, there are no known studies reporting higher testicular expression of *foxl2a*

in any other fishes, with most reporting low to no expression in testes (summarized in Bertho

2016). High expression of *FOXL2* protein has, however, been reported in American alligator

testes prior to temperature-dependent sex determination (Yatsu 2016), which may point to

latent sexual plasticity of the *A. mexicanus* testis. Taken together, it is possible that

neofunctionalization – or impairment of functionalization – of *foxl2a* in *A. mexicanus* has

conferred a novel role upon this gene in the testis, while *foxl2b* has retained an ovary-

promoting function.

Downstream of ovary-promoting transcription factors such as *foxl2a*, expression of the

steroidogenic enzyme aromatase (*cyp19a1a*) typically signals ovarian differentiation and

commitment in vertebrates via production of feminizing estradiol (E2). We found both

morphotypes to lack sexually dimorphic expression of this gene, suggesting deviation from

typical sex hormone signaling in *A. mexicanus* and possible elevation of E2 in males. *cyp19a1a* is

detected in somatic gonadal cells of both sexes, though in fish is broadly ovary-enriched and in

zebrafish is required to antagonize testis fate (Devlin and Nagahama 2002, Dranow 2016).

*cyp19a1a* can synthesize E2 through two biochemical pathways, a direct pathway that produces

E2 using testosterone produced by the steroidogenic enzyme *hsd17b3*, and an indirect pathway

that produces an estrone (E1) intermediate from androstenedione, requiring the action of

*hsd17b1* to ultimately generate E2 (Miller and Auchus 2011). Our analysis revealed sexually

dimorphic expression of both *hsd17b1* and *hsd17b3* (upregulated in ovaries and testes,

respectively), indicating that the deployment of either E2 synthesis pathway 1) differs between

sexes and 2) may ultimately result in expected rates or physiological concentrations of sex

hormones. The *hsd17b1* E2 synthesis pathway is also favored in zebrafish ovaries, though has

not been explored in testes, in which *cyp19a1a* expression is very low and dispensable (Liu

2022, Dranow 2016).  Future studies of circulating serum hormone levels will be necessary to

demonstrate whether *A. mexicanus* males exhibit physiologic aberrantly high estrogen

signaling. It is also possible that male *A. mexicanus* produce an inactive *cyp19a1a* protein due

to alternative splicing, as reported in the protandrous hermaphrodite barramundi (Domingos

2018).

***Upregulation of innervation in cavefish ovaries***

Enrichment analysis of the DEGs upregulated in the ovaries of Pachón cavefish versus

surface fish suggest possible differences in ovarian physiology. Most notably, we identified

enrichment of genes involved in neuron and axon differentiation and development. This is

consistent with our observation that the external ovarian epithelium is extensively innervated

in both morphotypes (Figure 4). In zebrafish, the ovarian epithelium demarcates a germinal

zone that is associated with nests of presumptive germline stem cells (Liu 2022). In the

mammalian ovary, nerve cells are connected to the follicles as well as gonadal blood vessels

(Malamed 2012), and neurotrophic growth factors and receptors are expressed at varying levels

dependent on oocyte stage (reviewed in Linher-Melville and Li 2013). Given this evidence that

innervation and neurotrophins influence folliculogenesis and oocyte maturation, enhanced

neurogenesis in Pachón ovaries may represent an adaptive reproductive strategy that regulates

renewal versus differentiation of ovarian germ cells in the cave environment, which is

reproductively challenging both in terms of nutrient deprivation and mate selection.

Additionally, increased innervation may facilitate reproductive success in caves in the

absence of visual stimuli, by integrating other sensory inputs from the environment that serve

as cues for spawning (e.g. temperature, water currents, precipitation). In zebrafish, production

of norepinephrine within neurons of the ovary is important for regulating oocyte maturation in

response to nutrient cues (Kim 2021). Increased sensitivity to such cues may provide cavefish

females an energy-saving advantage as nutrient availability fluctuates seasonally (Espinasa

2017). The functional significance of ovarian innervation in cavefish remains speculative, but

further studies comparing oocyte development between morphotypes may elucidate how cave

environments influence reproductive physiology.

***Upregulation of vascularization in cavefish testes***

Enrichment analysis uncovered a striking upregulation in the expression of angiogenesis-

related genes in Pachón cavefish compared to surface fish. This molecular evidence is

supported by our histological findings of increased vasculature in Pachón cavefish testes (Figure

6). Vertebrate seminiferous tubules are typically avascular and form a blood-testis barrier that

occludes the germinal epithelium (spermatogonia, spermatocytes, sperm, and Sertoli support

cells), though specialized non-fenestrated capillaries form in the interstitial space outside of the

seminiferous tubules where the steroidogenic Leydig cells reside (Ergün 2016). In mammals,

upregulation of vascularization is necessary for initiating organogenesis of the testis, in

particular ensuring development of fetal testis cords and limiting differentiation of the Leydig

cell population (reviewed in Gu 2021). Thus, cavefish testicular vascularization may indicate a

delayed or repressed state, with respect to sexual maturation. Across species, testes

vascularization allows for thermoregulation, establishing optimal timing and temperature for

spermiogenesis (Setchell 1998). In many fishes, temperature fluctuation in either direction

lowers testosterone production and inhibits germ cell differentiation (Manning and Kime 1985,

Konkal and Ganesh 2021). Testicular thermoregulation may be particularly challenging in

chronically low-temperature cave environments, and the temperature preferences exhibited by

cavefish could represent a reproductive strategy that preserves male fertility (Tabin 2018).

Cave environments are also hypoxic, a condition known to induce angiogenesis and

vascular branching via upregulated Notch signaling (Krock 2011). Pachón cavefish have adapted

to hypoxia by developing more erythrocytes and enlarged hematopoietic domains, driven by an

increase in Sonic hedgehog (*shh*) (van der Weele 2024). These studies align with our findings,

which show upregulation of *notch1b*, *notch3*, and *shh* specifically in the Pachón testes

(Additional file 2). Further comparative analysis of surface and cavefish testes will be needed to

characterize differences in rate of spermatogenesis or time to sexual maturity. Our study

provides molecular evidence for gonadal adaptation that allows for morphogenesis and

spermiogenesis to proceed in the extremes of cave environments.

**Conclusions**

Our study provides new insights into the molecular, functional, and structural

differences between surface fish and cavefish gonads, gametes, and fertility, contributing to the

broader understanding of reproductive biology in *A. mexicanus*. Reversal or loss of sex-specific

gene expression (*foxl2a, cyp19a1a*) suggests novel functional changes or plasticity emerging in

conserved sex regulating pathways in this species. The upregulation of neuron development in

ovaries and increased angiogenesis in testes highlight the potential for adaptive changes in

gonadal tissue architecture and function in response to environmental pressures in the cave.

These findings open several avenues for future research, including functional studies to

determine how these molecular changes impact reproductive success and how environmental

factors shape the evolution of sexual dimorphism and reproduction in cave-adapted

populations. Overall, our results establish a foundation for further investigations into the

evolution of sex-regulatory pathways and sexual dimorphism in organisms in extreme habitats.

**Declarations**

*Ethics approval and consent to participate*

Not applicable

*Consent for publication*

Not applicable

*Availability of data and materials*

The RNA sequencing datasets generated and analyzed during the current study are available

in the NCBI SRA repository [BioProject Reviewer Link].

*Competing interests*

The authors declare that they have no competing interests.

*Funding*

This project was supported by grants from the National Institute of General Medical Sciences

(GM103440 and 5 U54 GM104944).

*Authors' contributions*

KW conceived the study, collected samples, analyzed data, and wrote and edited manuscript

draft.  BP collected samples, analyzed data, and edited the manuscript draft. HVG, JP, and JH

processed and analyzed RNAseq data, and edited the manuscript draft. MR conceived the

study, analyzed data, and wrote and edited manuscript draft.

*Acknowledgements*

Brian Martineau, Emma Ferrante, and Taylor Rebbe provided fish husbandry in the aquatic

facility of Clifford J. Tabin at Harvard Medical School.

**References**

1. Abucay, J., Mair, J., Skibinski, D. & Beardmore, J. Environmental sex determination: the effect

of temperature and salinity on sex ratio in *Oreochromis niloticus*. *Aquaculture* 173, 219–234

(1999).

2. Baroiller, J. F., D'Cotta, H. & Saillant, E. Environmental Effects on Fish Sex Determination and

Differentiation. *Sexual Development* 3, 118–135 (2009).

3. Bertho, S. *et al.* Foxl2 and Its Relatives Are Evolutionary Conserved Players in Gonadal Sex

Differentiation. *Sexual Development* 10, 111–129 (2016).

4. Brown, E. E., Baumann, H. & Conover, D. O. Temperature and photoperiod effects on sex

determination in a fish. *Journal of Experimental Marine Biology and Ecology* 461, 39–43 (2014).

5. Devlin, R. H. & Nagahama, Y. Sex determination and sex differentiation in fish: an overview of

genetic, physiological, and environmental influences. *Aquaculture* 208, 191–364 (2002).

6. Domingos, J. A. *et al.* Sex-specific *dmrt1* and *cyp19a1* methylation and alternative splicing in

gonads of the protandrous hermaphrodite barramundi. *PLOS ONE* 13, e0204182 (2018).

7. Dranow, D. B. *et al.* Bmp15 Is an Oocyte-Produced Signal Required for Maintenance of the

Adult Female Sexual Phenotype in Zebrafish. *PLOS Genetics* 12, e1006323 (2016).

8. Durinck, S. *et al.* BioMart and Bioconductor: a powerful link between biological databases

and microarray data analysis. *Bioinformatics* 21, 3439–3440 (2005).

9. Elliott, W. R. Cave Exploration and Mapping in the Sierra de El Abra Region. in *Biology and*

*Evolution of the Mexican Cavefish* 9–40 (Elsevier, 2016). doi:10.1016/B978-0-12-802148-

4.00001-3.

10. Ergün, S., Davidoff, M. & Holstein, A. F. Capillaries in the lamina propria of human

seminiferous tubules are partly fenestrated. *Cell and Tissue Research* 286, 93–102 (1996).

11. Espinasa, L. *et al.* Discovery of Two New *Astyanax* Cavefish Localities Leads to Further

Understanding of the Species Biogeography. *Diversity* 12, 368 (2020).

12. Espinasa, L. *et al.* Contrasting feeding habits of post-larval and adult *Astyanax* cavefish.

*Subterranean Biology* 21, 1–17 (2017).

13. Giacobini, P. Shaping the Reproductive System: Role of Semaphorins in Gonadotropin-

Releasing Hormone Development and Function. *Neuroendocrinology* 102, 200–215 (2015).

14. Gu, X., Li, S. & DeFalco, T. Immune and vascular contributions to organogenesis of the testis

and ovary. *The FEBS Journal* 289, 2386–2408 (2022).

15. Guiguen, Y., Fostier, A., Piferrer, F. & Chang, C. F. Ovarian aromatase and estrogens: A pivotal

role for gonadal sex differentiation and sex change in fish. *General and Comparative*

*Endocrinology* 165, 352–366 (2010).

16. Herman, A. *et al.* The role of gene flow in rapid and repeated evolution of cave-related traits

in Mexican tetra, Astyanax mexicanus. *Molecular Ecology* 27, 4397–4416 (2018).

17. Imarazene, B. *et al.* Primordial Germ Cell Migration and Histological and Molecular

Characterization of Gonadal Differentiation in Pachón Cavefish *Astyanax mexicanus*; *Sexual*

*Development* **14**, 80–98 (2020).

18. Imarazene, B. *et al. A supernumerary "B-sex" chromosome drives male sex 1 determination*

*in the Pachón cavefish, Astyanax mexicanus 2 3.* https://ssrn.com/abstract=3875774.

19. Kim, J., Hyun, M., Hibi, M. & You, Y. J. Maintenance of quiescent oocytes by noradrenergic

signals. *Nature Communications* 12, (2021).

**20.** Konkal, P. & Ganesh, C. B. The effect of high or low temperature on testicular activity in the

cichlid fish *Oreochromis mossambicus. Fisheries Science* 87, 837–844 (2021).

21. Kossack, M. E. *et al.* Female Sex Development and Reproductive Duct Formation Depend on

Wnt4a in Zebrafish. *Genetics* 211, 219–233 (2019).

22. Krock, B. L., Skuli, N. & Simon, M. C. Hypoxia-Induced Angiogenesis: Good and Evil. *Genes &*

*Cancer* 2, 1117–1133 (2011).

23. Lau, E. S.-W., Zhang, Z., Qin, M. & Ge, W. Knockout of Zebrafish Ovarian Aromatase Gene

(cyp19a1a) by TALEN and CRISPR/Cas9 Leads to All-male Offspring Due to Failed Ovarian

Differentiation. *Scientific Reports* 6, 37357 (2016).

24. Li, F. *et al.* Gonadal Transcriptome Sequencing Analysis Reveals the Candidate Sex-Related

Genes and Signaling Pathways in the East Asian Common Octopus, *Octopus sinensis. Genes* 15,

(2024).

25. Linher-Melville, K. & Li, J. The roles of glial cell line-derived neurotrophic factor, brain-

derived neurotrophic factor and nerve growth factor during the final stage of folliculogenesis: a

focus on oocyte maturation. *REPRODUCTION* 145, R43–R54 (2013).

26. Liu, Y. *et al.* Single-cell transcriptome reveals insights into the development and function of

the zebrafish ovary. *eLife* **11**, (2022).

27. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biology* 15, 550 (2014).

28. Malamed, S., Gibney, J. A. & Ojeda, S. R. Ovarian innervation develops before initiation of

folliculogenesis in the rat. *Cell & Tissue Research* 270, 87–93 (1992).

29. Manning, N. J. & Kime, D. E. The effect of temperature on testicular steroid production in

the rainbow trout, *Salmo gairdneri, in vivo* and *in vitro*. *General and Comparative Endocrinology*

57, 377–382 (1985).

30. Matsuda, M. *et al.* DMY is a Y-specific DM-domain gene required for male development in

the medaka fish. *Nature* 417, 559–563 (2002).

31. Miller, W. L. & Auchus, R. J. The Molecular Biology, Biochemistry, and Physiology of Human

Steroidogenesis and Its Disorders. *Endocrine Reviews* 32, 81–151 (2011).

32. Myosho, T. *et al.* Tracing the emergence of a novel sex-determining gene in medaka, *Oryzias*

*luzonensis*. *Genetics* 191, 163–170 (2012).

33. Nakamura, Y. *et al.* Sox9 Is Upstream of MicroRNA-140 in Cartilage. *Applied Biochemistry*

*and Biotechnology* 166, 64–71 (2012).

34. Nishimura, T. & Tanaka, M. Gonadal development in fish. *Sexual Development* 8, 252–261

(2014).

35. Nynca, J. *et al.* Ovarian transcriptome analysis of diploid and triploid rainbow trout revealed

new pathways related to gonadal development and fertility. *Animal* 16, 100594 (2022).

36. O'Brien, B. Identification of Haemoglobin by its Catalase Reaction with Peroxide and *o* -

Dianisidine. *Stain Technology* **36**, 57–61 (1961).

37. Perera, P., Perez Guerra, D. & Riddle, M. R. The Mexican Tetra, *Astyanax mexicanus*, as a

Model System in Cell and Developmental Biology. *Annual Review of Cell and Developmental*

*Biology* **39**, 23–44 (2023).

38. Protas, M. E. *et al.* Genetic analysis of cavefish reveals molecular convergence in the

evolution of albinism. *Nature Genetics* 38, 107–111 (2006).

39. Qian, P., Kang, J., Liu, D. & Xie, G. Single Cell Transcriptome Sequencing of Zebrafish Testis

Revealed Novel Spermatogenesis Marker Genes and Stronger Leydig-Germ Cell Paracrine

Interactions. *Frontiers in Genetics* 13, (2022).

40. Riddle, M. R. *et al.* Genetic architecture underlying changes in carotenoid accumulation

during the evolution of the blind Mexican cavefish, *Astyanax mexicanus*. *Journal of Experimental*

*Zoology Part B: Molecular and Developmental Evolution* 334, 405–422 (2020).

41. Riddle, M. R. *et al.* Genetic mapping of metabolic traits in the blind Mexican cavefish reveals

sex-dependent quantitative trait loci associated with cave adaptation. *BMC Ecology and*

*Evolution* 21, 94 (2021).

42. Schartl, M. Sex chromosome evolution in non-mammalian vertebrates. *Current Opinion in*

*Genetics and Development* vol. 14 634–641 at https://doi.org/10.1016/j.gde.2004.09.005

(2004).

43. Setchell, B. P. The Parkes Lecture Heat and the testis. *Reproduction* 114, 179–194 (1998).

44. Shang, E., Yu, R. M. K. & Wu, R. S. S. Hypoxia Affects Sex Differentiation and Development,

Leading to a Male-Dominated Population in Zebrafish (*Danio rerio*). *Environmental Science &*

*Technology* 40, 3118–3122 (2006).

45. Sur A, Wang Y, Capar P, Margolin G, Prochaska MK, Farrell JA. Single-cell analysis of shared

signatures and transcriptional diversity during zebrafish development. Dev Cell. 2023 Dec

18;58(24):3028-3047.e12. https://daniocell.nichd.nih.gov/gene/S/stx5al/stx5al_pgc.html

46. Tabin, J. A. *et al.* Temperature preference of cave and surface populations of Astyanax

mexicanus. *Developmental Biology* 441, 338–344 (2018).

47. Takehana, Y. *et al.* Co-option of *Sox3* as the male-determining factor on the Y chromosome

in the fish *Oryzias dancena*. *Nature Communications* 5, (2014).

48. van der Weele, C. M., Hospes, K. C., Rowe, K. E. & Jeffery, W. R. Hypoxia-sonic hedgehog axis

as a driver of primitive hematopoiesis development and evolution in cavefish. *Developmental*

*Biology* 516, 138–147 (2024).

49. Webster, K. A. *et al.* Dmrt1 is necessary for male sexual development in zebrafish.

*Developmental Biology* 422, 33–46 (2017).

50. Wilson, E. J., Tobler, M., Riesch, R., Martínez-García, L. & García-De León, F. J. Natural history

and trophic ecology of three populations of the Mexican cavefish, Astyanax mexicanus.

*Environmental Biology of Fishes* 104, 1461–1474 (2021).

51. Wu, Y. *et al.* Syntaxin binding protein 2 in Sertoli cells regulates spermatogonial stem cell

maintenance through directly interacting with connexin 43 in the testes of neonatal mice.

*Molecular Biology Reports* 49, 7557–7566 (2022).

52. Yan, Y.L. *et al.* A Hormone That Lost Its Receptor: Anti-Müllerian Hormone (AMH) in

Zebrafish Gonad Development and Sex Determination. *Genetics* 213, 529–553 (2019).

53. Yang, Y. J., Wang, Y., Li, Z., Zhou, L. & Gui, J. F. Sequential, divergent, and cooperative

requirements of *foxl2a* and *foxl2b* in ovary development and maintenance of zebrafish. *Genetics*

205, 1551–1572 (2017).

54. Yatsu, R. *et al.* RNA-seq analysis of the gonadal transcriptome during *Alligator*

*mississippiensis* temperature-dependent sex determination and differentiation. *BMC Genomics*

17, 77 (2016).

55. Yu, G., Wang, L. G., Han, Y. & He, Q. Y. ClusterProfiler: An R package for comparing biological

themes among gene clusters. *OMICS A Journal of Integrative Biology* 16, 284–287 (2012).

**Figure Legends**

**Figure 1. Transcriptome analysis of *Astyanax mexicanus* gonads.** Appearance of *A. mexicanus*

surface fish (**A**) and Pachón cavefish (**B**) males and females. Appearance of sexually mature

testis and ovaries of surface fish (**C, E**) and Pachón cavefish (**D, F**). Testes (**C, D**) consist of two

lobes of tubules containing male germ cells and sperm and terminating at the efferent

urogenital duct (white arrowheads). Ovaries (**E-F**) consist of two lobes containing clusters of

female germ cells and eggs surrounded by the germinal epithelium and terminating at the

oviduct (black arrowhead). (**G**) Principal component analysis (PCA) plot of normalized, variance

stabilized RNA sequencing expression profiles of *A. mexicanus* gonads. Each point represents an

individual sample, colored by sample type. The first principal component (PC1) explains 57% of

the total variance and separates samples by sex. The second principal component (PC2) explains

27% of the total variance and separates samples by morphotype (surface fish vs Pachón

cavefish). PCA plots based on differentially expressed genes (DEG) only (adjusted p-value less

than 0.05).

**Figure 2. Sex-specific gene expression profiles in *Astyanax mexicanus* surface fish and Pachón**

**cavefish gonads. (A)** Volcano plot showing genes that are differentially expressed between *A.*

*mexicanus* testes and ovaries (includes data from juvenile, adult, surface fish, and Pachón

cavefish gonads). The y-axis shows -log10 adjusted p-value and the x-axis shows log2 fold

change for all genes after low-count filtering (23,809 transcripts). Genes with adjusted p-value

< 0.05 and absolute log2 fold change > 1 are highlighted in red. Negative fold change indicates

testis-enriched expression, while positive fold change indicates ovary-enriched expression. **(B)**

Summary of genes with sex-specific expression profiles in the gonads of zebrafish (*Danio rerio*),

compared to the expression profiles we observed in *A. mexicanus*. "Ovary" indicates a gene

with upregulation in female gonads, whereas "testis" indicates a gene with upregulation in male

gonads. **(C-J)** Plots showing *A. mexicanus* gonad log2 transcript abundance of genes with sex-

specific expression profiles and functions in the gonads of other vertebrates. Each point

represents the transcript abundance in an individual sample, colored and organized on the x-

axis by the sample type. From left to right: cavefish (CF) juvenile ovary, CF adult ovary, surface

fish (SF) juvenile ovary, SF adult ovary.

**Figure 3. Ovary transcriptome differences between surface fish and Pachón cavefish suggest**

**enhanced neurogenesis in cavefish. (A, B)** Volcano plots showing genes that are differentially

expressed between the ovaries of *A. mexicanus* surface fish and Pachón cavefish at juvenile (**A**)

and adult (**B**) stages. The y-axis shows -log10 adjusted p-value and the x-axis shows log2 fold

change for all genes after low-count filtering.  Genes with adjusted p-value < 0.05 and absolute

log2 fold change > 1 are highlighted in green. Negative fold change indicates greater expression

in surface fish, while positive fold change indicates greater expression in cavefish. (**C**, **D**) Dot

plots displaying top 15 enriched biological process gene ontology (GO) terms identified from

gene set enrichment analysis of genes upregulated in Pachón cavefish compared to surface fish

at juvenile (**C**) and adult (**D**) stages. Each dot represents a biological process GO category, with

the size of the dot corresponding to the number of genes in the category, and the color scale

indicating the adjusted p-value. The x-axis represents the gene ratio (proportion of input genes

involved in the pathway), while the y-axis lists the enriched pathways.

**Figure 4. Innervation in the *Astyanax mexicanus* ovarian epithelium revealed by wholemount**

**immunostaining**. Brightfield images of the ovaries of adult surface fish (**A**) and Pachón cavefish

(**B**). Lobes on the lefthand side of the image are placed with the medial side upward, while

lobes on the righthand side are placed with the lateral side upward. Visualization of acetylated

tubulin in wholemount fluorescence images of the ovaries shown in A, B at low magnification

(**B, F**) and higher magnification (**C, D, G, H**) highlighting the differences in innervation in the

medial side (**C, G**) versus lateral side (**D, H**) of the ovary. A large neuron trunk (solid arrows) can

be seen on the medial side of the ovary with branches (dashed arrows) that reach throughout

the ovarian epithelium.

**Figure 5. Testis transcriptome differences between surface fish and Pachón cavefish suggests**

**enhanced angiogenesis in cavefish. (A,B)** Volcano plots showing genes that are differentially

expressed between the testes of *A. mexicanus* surface fish and Pachón cavefish at juvenile (**A**)

and adult (**B**) stages. The y-axis shows -log10 adjusted p-value and the x-axis shows log2 fold

change for all genes after low-count filtering. Genes with adjusted p-value < 0.05 and absolute

log2 fold change > 1 are highlighted in green. Negative fold change indicates greater expression

in surface fish, while positive fold change indicates greater expression in cavefish. (**C**, **D**) Dot

plots displaying top 15 enriched biological process gene ontology (GO) terms identified from

gene set enrichment analysis of genes upregulated in Pachón cavefish compared to surface fish

at juvenile (**C**) and adult (**D**) stages. Each dot represents a biological process GO category, with

the size of the dot corresponding to the number of genes in the category, and the color scale

indicating the adjusted p-value. The x-axis represents the gene ratio (proportion of input genes

involved in the pathway), while the y-axis lists the enriched pathways.

**Figure 6. Histological comparison of testis vasculature between *Astyanax mexicanus* surface**

**fish and cavefish.** Histological sections of adult testes stained to visualize blood vessels in

surface fish (A) and Pachón cavefish (B). Vasculature is stained using o-dianisidine (brown) with

hematoxylin (blue) as a counterstain. Surface fish testis section (B) shows many seminiferous

tubules and only a single vascular bundle (black arrow). In contrast, cavefish testis section

shows numerous vascular bundles around the seminiferous tubules (black arrows). se =

seminiferous epithelium; sz = spermatozoa.

**Additional material**

**Additional file 1.pptx** Boxplots displaying weight of fish (A-B) and gonad (C-D) for the samples

used for RNAseq analysis. Juvenile gonads were too small to accurately measure weight. In

boxplots central line represents median, upper and lower edges indicate first and third

quartiles, and whiskers extend to 1.5 times the interquartile range. No significant differences

were observed between sample types (t-test p > 0.05).

**Additional file 2.pptx** Plots showing *A. mexicanus* gonad log2 transcript abundance of genes

potentially involved in steroidogenesis (*hsd17b1* and *hsdb17b3*) or vascularization and

angiogenesis (*notch1b, notch3, shh*). Each point represents the transcript abundance in an

individual sample, colored and organized on the x-axis by the sample type. From left to right:

cavefish (CF) juvenile ovary, CF adult ovary, surface fish (SF) juvenile ovary, SF adult ovary.

Figure 1

**Figure 2**

Figure 3

Figure 4

Figure 5

Figure 6