Molecular analyses of individual variation in honey bee sociability

Ian M. Traniello[1†*], Arian Avalos[2], Michael J. M. Gachomba[3], Tim Gernat[1], Zhenqing Chen[1], Amy C. Cash-
Ahmed[1], Adam R. Hamilton[1], Jennifer L. Cook[3], Gene E. Robinson[1,4,5]

[1]Carl R Woese Institute for Genomic Biology, University of Illinois at Urbana-Champaign, Urbana, IL (UIUC).

[2]United States Department of Agriculture – Agricultural Research Services, Honey Bee Breeding, Genetics and
Physiology Research Laboratory, Baton Rouge, LA.

[3]School of Psychology, University of Birmingham, Birmingham, UK

[4]Neuroscience Program, University of Illinois at Urbana-Champaign, UIUC.

[5]Department of Entomology, UIUC.

[†]Current Address: Lewis-Sigler Institute for Integrative Genomics, Princeton University, Princeton, NJ.

*Corresponding authors. IMT: it4770@princeton.edu; GER: generobi@illinois.edu

Abstract

Individual variation in sociability is a central feature of every society. This includes honey bees, with
some individuals well-connected and sociable, and others at the periphery of their colony's social
network. However, the genetic and molecular bases of sociability are poorly understood. Trophallaxis –
a behavior involving sharing liquid with nutritional and signaling properties - comprises a social
interaction and a proxy for sociability in honey bee colonies: more sociable bees engage in more
trophallaxis. Here we identify genetic and molecular mechanisms of trophallaxis-based sociability by
combining genome sequencing, brain transcriptomics, and automated behavioral tracking. A genome-
wide association study (GWAS) identified 18 single-nucleotide polymorphisms (SNPs) associated with
variation in sociability. Several SNPs were localized to genes previously associated with sociability in
other species, including in the context of human autism, suggesting shared molecular mechanisms of
sociability. Variation in sociability also was linked to differential brain gene expression, particularly genes
associated with neural signaling and development. Using comparative genomic and transcriptomic
approaches, we also detected evidence for divergent mechanisms underpinning sociability across
species, including those related to reward sensitivity and encounter probability. These results highlight
both potential evolutionary conservation of the molecular roots of sociability and points of divergence.

Introduction

Consistent variation among individuals in the tendency to engage one another is a central feature of all
animal societies. "Sociability," which broadly characterizes associative interactions outside of the
context of aggression or courtship [1], falls on a spectrum, with hypo- and hyper-sociable individuals
flanking a "normal" intermediate range [2]. There are many causes of this variation, stemming from a
combination of labile drivers like motivational state [3], social status [4], and previous experience [5,6],
as well as heritable factors that drive lifelong positive or negative inclinations toward social interactions
[7].

As a trait, sociability arises from the integration of socially responsive neural circuitry, physiology, and
experience, all of which contribute to how an animal interprets and responds to social cues [8–10].
Detailed study of vertebrate brain evolution has outlined a "social behavior network" that regulates

multiple forms of social behavior [11], acting in concert with the mesolimbic reward system to influence
how an animal makes decisions in the context of various social situations [2,12,13].

Social insects, including the western honey bee (*Apis mellifera*), have tiny and very differently organized
nervous systems relative to vertebrates, but they also form large and intricate societies whose
ecological and evolutionary success are tightly linked to individual- and group-level sociability [14,15].
Honey bees form large colonies of tens of thousands of individuals and are a powerful model for further
understanding the relationship between genes and sociability. The bulk of the colony is composed of
sterile workers for whom task performance is influenced by a combination of individual age, genetics,
and the needs of the colony [16–18]. Honey bee workers lack morphologically distinct castes, and each
individual maintains the ability to perform all tasks in the colony other than reproduction required for
colony growth and development. The molecular mechanisms underlying specific behavioral states like
brood care ("nursing"), guarding the hive entrance, or foraging outside for floral resources are sensitive
to environmental demands, with clear individual differences in the likelihood of performing certain tasks
as well as predisposition toward social contact [5]. We used honey bees to study genetic and molecular
components of sociability, including comparative genomic analyses.

Comparative genomics offers a toolkit for exploring the extent to which analogous behaviors, like the
tendency to engage socially with others, have similar molecular representations in the brain across
disparate species [19,20]. While a behavioral trait may be similarly expressed across species, whether
that trait also has a common molecular basis requires observation of similarities across deeper layers of
biological organization, including at the level of the genome and/or transcriptome. This approach allows
cognitive scientists to better understand the development of the "social brain," bypassing the need to
compare distinct brain anatomies, which is often experimentally challenging, in favor of studying their
more quantifiable molecular building blocks. In addition, brain-wide spatial localization of homologous
genes or proteins supporting analogous behaviors across species can clarify how vastly different
neuroanatomies can support analogous forms of social life [13,21,22]. Drawing from studies in social
insects, variation in gene sequence and expression profile have suggested some degree of conservation
between vertebrates and invertebrates in the neurogenomic architecture of social behavior, despite
~600 million years divergence from humans [20,23,24].

Sociability is thought to have a heritable component. In humans, for example, structural variation in the
7q11.23 genomic locus can dramatically affect the developmental trajectory of the social brain, causing
either lifelong hypo- or hyper-sociability [7]. Such structural variants affect many aspects of an
individual's neurobiology and physiology, making it challenging to pinpoint direct genetic effects on the
propensity to engage in social interactions within a typical range of individual variation. Nevertheless,
such findings indicate the existence of heritable molecular substrates that mediate the intensity of
sociability.

One limitation in the study of vertebrate sociability is related to difficulties in tracking many individuals
over long periods of time – a major challenge in relating genotype and social phenotype in humans [25].
It is similarly difficult to disentangle sociability from related traits, like locomotion, that contribute to
overall activity rate: an individual may interact with others more because they are motivated to engage
such interactions, because they are simply more physically active and therefore more likely to engage
others by chance, or both. But, in some species, it is not possible to characterize individuals across
multiple social phenotypic vectors. This issue has been largely overcome for social insects, as recent

advances in automated tracking now allow for behavioral monitoring of each individual in a colony over
long portions of their lives [26–29]. These advances permit finely resolved comparisons of social
behavior, brain, and genome in a naturalistic setting.

In addition to task-related behaviors, honey bees also frequently engage in mouth-to-mouth sharing of
liquid containing nutritive and signaling components in a stereotyped process called trophallaxis.
Automated behavioral tracking of barcoded bees has shown consistent variation in trophallaxis
frequency to be a robust proxy for several aspects of an individual's life history, from task specialization
to disease status [5,30–32]. More generally, trophallaxis acts as a "social glue" mechanism central to
group cohesion in socially advanced insect societies [33], reflecting individual- and group-level
tendencies toward reciprocal social interaction. In addition, variation in the duration of trophallaxis
interaction in bees and face-to-face interactions in humans fall along similar heavy-tailed distributions,
suggesting that universal principles may underlie the biology of social interaction networks [34]. Taken
together, trophallaxis is a quantifiable analog to other measures of sociability across diverse taxa.

Though automated tracking of individuals engaged in trophallaxis has yielded new insights into the
social structure of honey bee society, there are important gaps in the analysis of trophallaxis sociability.
These include genomics, transcriptomics, and movement dynamics associated with trophallaxis within
the colony. Addressing these gaps promises to reveal deeply conserved neurobiological mechanisms
that shape the social brain [28] as well as behavioral mechanisms associated with social bond formation,
as in humans [35,36].

In honey bees, transcriptomic analyses have revealed provocative associations between variation in
brain gene expression and variation in sociality. Shpigler et al. [20] found significant enrichment for
Autism-related genes in the brains of individual honey bees that were consistently unresponsive to task-
related social stimuli. In a second study, these "socially unresponsive" bees also were found to have
fewer and briefer trophallaxis interactions compared to age-matched individuals that were more
responsive to task-related social stimuli [5]. These results established a link between variation in task
performance and variation in sociability.

Here we report on behavioral, genomic, transcriptomic, and comparative genomic analyses of
trophallaxis sociability. First, we employed automated tracking of individually barcoded bees in colonies
to reveal individual variation in this behavior. Second, we quantified movement kinematics to
disentangle trophallaxis engagement frequency from overall activity rate and test hypotheses regarding
how individual bees influence one another in terms of movement within the hive. Third, we followed
these analyses with lab-based behavioral assays that robustly indicate task specialization. We then
performed whole-genome resequencing and brain transcriptomic profiling on these extensively profiled
individuals to identify genetic variants and brain gene expression profiles associated with variation in
behavior. We also localized the expression in the brain of one particularly interesting gene, *neuroligin-2*
(*nlg2),* identified in both genetic and transcriptomic analyses. Finally, using comparative genomics, we
explored similarities and differences in variation in trophallaxis sociability, reward processing, and
aggression in honey bees and across species as well.

Methods

Animals

All molecular and bioinformatic analyses were performed on bees collected as part of a previous study
that addressed different questions [5]. The field- and lab-based behavioral experiments took place at the
University of Illinois Bee Research Facility, Urbana, IL, USA, from August to September 2017. Briefly, we
used adult worker bees collected from three colonies, each headed by a queen instrumentally
inseminated with semen from a single male (drone, "SDI queen"); individuals derived from these
matings have an average coefficient of relatedness of 0.75 due to haplodiploidy. To obtain 0- to 24-h-old
(one-day-old) bees for age-matching, honeycomb frames containing late-stage pupae were removed
from colonies, maintained in a dark incubator at 34°C and 50% relative humidity (RH), and ~500
individuals were gently swept off the frames the morning of each behavioral experiment. All colonies
were filmed when the bees were 3 to 10 days old. On Day 10 of the recording, bees were removed and
prepared for laboratory-based behavioral assays, as described below.

Automated behavioral tracking, image processing, and trophallaxis detection

Single-sided glass-walled observation hives, each containing a colony with barcoded bees, were set up
as previously described [5,27,30,31]. Individual bees were cold-anesthetized and a barcode was applied
to the dorsal thorax using a small amount of cyanoacrylate glue. Barcoded bees were then gently
transferred to an observation hive containing one honeycomb frame provisioned with honey (top 18
rows, ~200 mg per cell, ~330 g in total) and pollen paste (next six rows, ~100 mg per cell, ~45 g total;
pollen paste was made from 45% honey, 45% pollen and 10% water). A naturally mated queen,
unrelated to the workers, was lightly anesthetized with carbon dioxide, barcoded, and allowed to
recover surrounded by workers in the observation hive.

Observation hives were then transferred to a dark room connected to the outside by a foraging port
that was kept closed for the first two days of the experiment to prevent young bees with undeveloped
flight muscles from exiting the colony. For recording, we used LED lights (Smart Vision Lights, Muskegon,
MI, USA) to illuminate the honeycomb with infrared light, which the bees cannot see. Whole-colony
images were captured at a rate of 1 image/s using a Prosilic GT6600 machine vision camera (Allied
Vision, Exton, PA). To ensure sharp images, the observation hive glass window was gently replaced daily
with a clean one without disturbance to the colony.

Hive images were processed as previously described [5]. Briefly, images were resized and sharpened to
improve barcode detection rate, and each bee's location and orientation were inferred using custom
software [27], which also filtered any barcode that was unreadable or resulting from a spurious
detection. We then used a convolutional neural network to identify all trophallaxis events by recording
when two bees were close (1.7 − 7.4 mm apart), facing each other, and one bee had inserted her
extended proboscis ("tongue") between the open mouthparts of another bee [5,37]. These "raw"
trophallaxis events were merged if they were consecutive and <1 min apart, and interactions lasting <3 s
or >3 min were removed as spurious detections.

Bee kinematics quantification

We quantified bee movement as the instantaneous linear (locomotion) and angular (turning) speeds of
individuals. Linear speed was defined as the Euclidean distance between a bee's barcode center at time
points t and t+1, divided by the image rate r. Angular speed a was defined as the unsigned angle
between a bee's barcode orientation vector at t and t+1, divided by r. Mean speeds were calculated by
averaging these quantities across time. If a bee was not detected at time point t, we recorded no speed

measurements for the image captured at time point t+1. We also recorded no speed measurement if a
bee moved less than the width of a honeycomb cell (4.9 mm) and turned less than from one cell corner
to the next (60°). These thresholds were applied to differentiate actual locomotion from the small
movements that a bee may perform while she is stationary (e.g., during allogrooming and other
behaviors).

Social influence analysis

To measure the extent to which the movement of one bee influenced the movement of another bee, we
computed the information partial directed coherence (iPDC) [38], employing the AsympPDC package
https://www.lcs.poli.usp.br/~baccala/pdc/CRCBrainConnectivity/ [39]. iPDC measures the directed
coherence between two or more time-series, enabling estimation of the bidirectional transfer of
movement influence-related information. As previously done [5], we focused on the last two days of the
recording and limited analysis to periods characterized by proximal interactions (i.e., when two bees
were physically close to each other for a minimum amount of time), since it is unlikely that two bees
located far apart in the hive would influence each other's movement directly.

To identify periods of proximal interaction, we computed for each image the Euclidean distance R
between a focal bee and all other bees, using the X and Y coordinates of their barcodes. Given a focal
bee, we defined *neighbor* as another bee whose distance R to the focal bee was equal or lower than a
radius R_{max} for a duration L equal or higher than a minimum duration L_{min} ($R<=R_{max}$, $L>=L_{min}$). We chose
R_{max} = 2cm for being approximately twice the length of a bee's body, and because previous studies used
this value as a measure of proximal interaction in honey bees (Wild et al., 2021). We chose L_{min} = 30
images as a minimum duration of the proximity interaction in order to have sufficient time-series
samples for computing iPDC. In addition, since our aim was to measure social influence during
movement, we excluded images where either one of the two bees' speed was very low as well as images
where the two bees were engaging in trophallaxis, during which animals are relatively still. If a focal bee
interacted with the same neighboring bee multiple times satisfying the above criteria, we kept data
from the first interaction only. In this manner, the identity of the neighbor varied with each proximity
interaction. The number of proximity interactions varied across focal bees, depending on the number of
unique neighbors they encountered, which could be from the same or different source colony.

We retrieved bees' X and Y coordinates during each proximity interaction and built an input dataset to
which iPDC was applied. In the dataset, each interaction is represented as a bivariate time-series of at
least 30 time points (30 sec), where the first time-series is the focal bee's X or Y coordinate and the
second time-series is its neighbor's X or Y coordinate, respectively. Next, we fit a vector autoregressive
model (VAR) to the time-series in each interaction and computed from it the iPDC spectra, from the
focal bee to the neighbor and vice versa. iPDC spectra were then averaged across interactions to obtain
a single iPDC spectrum for each focal bee, in both directions. Finally, the averaged iPDC spectra were
integrated across all frequencies into information flow (I_{flow}), as adapted from equation 8 in (Takahashi
et al., 2010):

$$I_{flow} = -\frac{1}{fs} \cdot \int_0^{fs/2} log2\left(1 - iPDC(f)\right) df$$

199 where *fs* is the sampling rate.

200　We thus obtained a scalar value representing the magnitude of the influence between the focal bee and
201　its neighbors in units of information transfer (bits).

Behavioral state identification via laboratory-based assays and foraging detection

Behavioral states related to natural colony division of labor (guards, nurses, foragers, generalists, and
non-responders) were studied under controlled laboratory conditions as previously described [5].
Briefly, at the conclusion of the recording portion of the experiment, the glass observation window
covering the hive was replaced with a Plexiglas window containing resealable portholes, and groups of 9
bees were gently removed, paint-marked with a unique color applied to the dorsal thorax (Testors Paint,
Rockford, IL), and transferred to a vertically oriented 100 x 200 mm Petri dish (Thermo Fisher Scientific,
Waltham, MA). Dishes contained a tube of honey (~1.2 ml), 50% sucrose solution (2 ml), and a pollen
ball (70% pollen, 30% sucrose solution described above). Bees were given at least 60 min to acclimate to
normal fluorescent lighting prior to the start of the behavioral assays.

Established laboratory assays were used to identify aggression, which reflects guarding behavior
[40,41] and affiliative caregiving, which reflects nursing behavior [42,43]. Aggression was measured by
subjecting groups of bees to a 5-min interaction with a foreign bee ("intruder") and quantitating
aggressive biting and stinging interactions. Affiliative caregiving was measured by exposing groups of
bees to a larva for 5 min and quantitating observations of caregiving interactions like food provisioning
and inspection. Each group was exposed to both stimuli, and "guards" and "nurses" were identified
based on consistent (≥20–30 s) observations of biting and stinging or larval feeding, respectively. We
also noted instances of fanning, wax-building, and vibration-signaling behaviors (following descriptions
in [44]) to give a more comprehensive depiction of each individual's behavioral state.

Affiliative care assays were performed in a controlled environment that mimicked the interior of the
hive (34°C and 50% relative humidity) under ambient lighting. Aggression assays were performed in a
cooler room (28°C) to approximate outdoor temperatures at the hive entrance, where defensive
aggression by guards against intruders is most likely to occur. Each dish of 9 bees was tested in both
assays in a random order with the second assay performed 60 min after the first. Immediately following
the second assay, all bees were flash-frozen in liquid nitrogen, their barcodes removed for identification
purposes, and stored at -80°C prior to DNA and RNA analysis.

Because the barcoded colonies were initially composed only of one-day-old bees, precocious foraging
occurred during the observation period from days 3-10 [45]. Foragers were identified via automated
monitoring of flight in and out of the hive with an entrance monitor attached to the outer portion of the
entrance tube connected to the observation hive [30]. Though foraging typically occurs toward the end
of a honey bee's lifetime, in colonies such as these that lack older bees, a subset will exhibit
"precocious" foraging to serve the colony's nutritional needs [46,47]. The entrance monitor contained a
small enclosure with a simple maze for slowing bees down to facilitate image capture. A Raspberry Pi
Camera Module v1.3 was mounted above the enclosure and separated from the maze by a removable
glass window. The camera recorded .mjpg videos at a temporal resolution of 3 images/s from the hours
of 07:00 to 19:00 h CST, automatically adjusting for changes in light conditions throughout the day.

Videos of bee activity were first converted to still images using ffmpeg (https://ffmpeg.org/) and then
processed to detect barcoded bees, as previously described [31]. Raw incoming and outgoing "passes"
were merged by vectoring the bee's displacement toward or away from the hive entrance. To be labeled

a forager, we required bees to be detected taking at least six trips, with more than three trips per day,
for any two days of the experiment, and at least 25% of these trips had to be made during peak foraging
hours (10:00-15:00 h CST). This is consistent with honey bee foraging behavior under natural conditions
[48].

Bees that did not respond to behavioral stimuli during the lab assays and were not observed to forage
when in their observation hive were labeled "non-responders" [20]. Bees that showed mild
responsiveness (i.e. < 20 s interaction) and displayed some other behaviors (fanning, wax-building, and
vibration-signaling behaviors) were referred to as "baseline" bees for which the behavioral state could
not be robustly identified but were clearly responsive to stimuli. Bees that exhibited two or more
behaviors (i.e., brood care, aggression, and/or foraging) were "generalists."

DNA- and RNA-Sequencing and analysis

All DNA- and RNA-Sequencing data were newly generated for this study from the individuals analyzed
behaviorally as described above.

DNA-Sequencing

DNA was extracted from the thorax of 391 individual bees using the Gentra Puregene Tissue Kit
(QIAGEN, Germantown, MD, USA) as previously described (Avalos et al., 2020). Shotgun genomic
libraries were prepared using the Illumina DNA (Nextera Flex) sample preparation kit, pooled,
quantitated via qPCR, and sequenced on one S4 lane for 151 cycles from both ends of the fragments on
an Illumina NovaSeq 6000. Raw fastq files were generated and demultiplexed using the bcl2fastq v2.2
conversion software (Illumina).

Mushroom body RNA-Sequencing

We sequenced RNA from 176 individuals, a subset of the larger set used for DNA sequencing. These
individuals were selected because they each had a definable behavioral state other than "baseline,"
following our above-described assignment strategy. We did not sequence "baseline" bees, as these
individuals did not forage or robustly engage social stimuli in the dish assays, but showed higher levels
of engagement than non-responders, in addition to other well-known natural behaviors like wax-
building or fanning. As we could not reliably characterize these individuals in behavioral terms,
molecular profiling would not be as meaningful as for other groups.

Mushroom body (MB) dissection and RNA extraction were performed as previously described [5,49]. We
chipped a small window on the anterior surface (frons) of flash-frozen bee heads immersed in a dry/ice
ethanol bath before submerging the entire head in RNAlater-ICE Frozen Tissue Transition Solution
(Thermo Fisher Scientific) overnight at 20°C. The brain was then fully dissected on wet ice and the MB
was removed and stored at −80°C until RNA extraction.

We used the PicoPure RNA Isolation Kit (Thermo Fisher Scientific, Waltham, MA) with DNase treatment
(Invitrogen, Carlsbad, CA) as has been previously described [5,20] and quantified via QuBit fluorometer
(Thermo Fisher Scientific). RNA-Seq libraries were prepared with the Kapa Hyper Stranded mRNASeq
Sample Prep Kit (Illumina, San Diego, CA); libraries were pooled, quantitated via qPCR, and sequenced
on an S4 lane for 151 cycles from both ends of the fragment on an Illumina NovaSeq 6000.

Bioinformatic analyses

 DNA alignment, variant calling and Genome-Wide Association Study (GWAS)

Whole-genome resequencing yielded at least 40,000,000 paired-end reads per sample (mean [μ] $\pm$
standard deviation: μ = 64,900,620 $\pm$ 9,035,972). After adaptor trimming, raw reads were aligned to the
honey bee genome, Amel_HAv3.1 (Wallberg et al., 2019), using the Burrows-Wheeler algorithm, and
resulting alignment files were sorted and deduplicated. The Samtools "flagstat" report indicated > 99%
mapping rates for samples to the honey bee genome, and deduplication of sequenced reads suggested a
duplication rate of < 20% (μ = 16.8 $\pm$ 0.012%). Thus, each sample was sequenced at an average depth of
30x, indicating a high-quality dataset suitable for GWAS.

Variant calling was performed using the Sentieon DNAseq workflow
(https://support.sentieon.com/manual/DNAseq_usage/dnaseq/). Following indel realignment and
variant calling, which resulted in a VCF file for each of our samples, genotyping was conducted across all
samples to generate a single multi-sample VCF file. We then removed indels, multiallelic variants, and
variants localized to unplaced scaffolds or mitochondrial genes. The remaining biallelic SNPs were
filtered using joint quality measures and overall representation in the dataset, utilizing missing and low-
coverage calls in the multi-sample VCF file. No sample showed excessive missing reads. Sequenced
genomes were then phased and imputed with Beagle (v5.1) [50,51]. Linkage disequilibrium (LD) pruning
was performed using the snpgdsLDpruning command in the R package SNPRelate [52] using the D'
metric with an LD threshold set to 0.3 and mean allele frequency set to 0.2. The resulting ~2.5 million
SNPs were used for downstream analyses.

We derived a genetic relatedness matrix (GRM) using the pcrelate function in GENESIS [53] to account
for variation based on kinship, as has been previously done in honey bees [54]. Bees that did not cluster
with their colony of origin, and/or bees that were later found to have unreadable barcodes (and
therefore unreliable identifying information), were excluded from analysis, leaving us with a final sample
size of 357 individuals for DNA analysis.

For individual-level phenotypic association, we utilized the quasi-likelihood approximation in GENESIS,
which scores and assigns a *P*-value to each SNP, thus predicting the relevance of the association
between a given SNP and phenotype for each genotype. We used GENESIS to identify SNPs significantly
associated with trophallaxis sociability or kinematic metrics like locomotion and distance traveled, all
averaged over the final two days of the recording experiment.

Brain mushroom body (MB) RNA-Sequencing alignments and detection of differentially expressed genes

Each sample yielded at least 13,000 paired-end reads (μ = 17,836,571 $\pm$ 2,286,238). After
demultiplexing, raw reads for each sample were aligned to the honey bee genome, Amel_HAv3.1 [55]
using STAR v2.7.3a [56] and reads were counted using the *featureCounts* function in the Subread v2.0
package [57]. We identified the presence of Deformed Wing Virus (DWV), a virus that is naturally
widespread in honey bee apiaries and commonly found in the brain [41,58] by also aligning raw reads to
the DWV genome. All linear models for identifying differentially expressed genes (DEGs) included
percentage of total reads aligned to the DWV genome as a covariate, as well as colony of origin, in order
to control for unwanted variation due to natural viral presence or latent genetic variation across
colonies. We observed a unique mapping rate of 84.5 $\pm$ 0.0845% against the honey bee genome, and an
average of 65.7 $\pm$ 0.066% of these reads were uniquely mapped to a gene model.

We first identified DEGs associated with sociability by regressing MB gene expression using trophallaxis
sociability, averaged over the last two days of the experiment [5], as a continuous predictor. Next, we
calculated DEGs associated with each behavioral state (generalist, forager, guard, nurse, or non-
responder) by comparing each state to expression levels averaged across the remaining four groups.
Each DEG list was generated via the Wald test and corrected for multiple testing using the Benjamini-
Hochberg method, FDR < 0.05. Gene Ontology enrichment analysis was performed by converting honey
bee gene identifiers to one-to-one orthologs in *Drosophila melanogaster* via a previously published
reciprocal best-hit BLAST [58], detecting term enrichment in GOrilla [59], and visualizing results in GO-
Figure [60]. Gene list overlap analyses were performed using the GeneOverlap package
(https://bioconductor.org/packages/GeneOverlap) with Bonferroni corrections for multiple testing.

Spatial analysis of neuroligin-2 *gene expression*

In situ hybridization was performed in January 2023 to provide additional information on one of the
genes identified in DNA and RNA analyses that emerged as a particularly strong sociability-associated
candidate gene, *neuroligin 2* (*nlg2*). One-day-old individuals were collected from a honeycomb frame
containing brood from a colony kept indoors at the Bee Research Facility. They were kept in a small
Plexiglas cage at 35°C and 50% until whole-brain dissections were performed at 7 or 14 days of age.

We followed an established protocol (https://www.protocols.io/view/hcr-rna-fish-protocol-for-the-whole-
mount-brains-o-bzh5p386) with minor modifications. Brains were dissected in phosphate-buffered saline
(PBS) and fixed in 4% paraformaldehyde (in 1X PBS with Triton-X100; PBS-T) overnight at 4°C. Fixed
brains were washed three times in PBS with Tween (Fisher Scientific), dehydrated in serial
methanol/PBST washes (25%, 50%, 75% and 100%), and stored at -20°C until use.

Brains were rehydrated in the reverse methanol/PBST serial washes (75%, 50%, 25%) and washed in
PBS-T. Brains were pre-hybridized in hybridization buffer (Molecular Instruments, Los Angeles, CA)
without probes for 1 h at 37°C. Next, we combined 4 µL of 1 µM HCR detection probes targeting the
honey bee *nlg2* gene (Molecular Instruments) with 250 µL hybridization buffer, and incubated each
brain in this solution overnight at 37°C. The following day, brains were washed in preheated probe wash
buffer (Molecular Instruments) 5 times, 10 min per wash, at 37°C, and then in 5X saline-sodium citrate
twice for 5 min per wash. Washed brains were pre-amplified in amplification buffer (Molecular
Instruments). Signal amplification hairpin oligos (B1-H1-488 and B1-H2488, from Molecular Instruments)
were first denatured at 95°C for 90 s and annealed by cooling on wet ice for 30 min. Amplification
hairpins were used at a concentration of 6 µL per 100 µL amplification buffer. Brains were incubated in
the amplification probe solutions overnight, followed by two washes in probe wash buffer containing
DAPI (Fisher Scientific, R37606). The samples were washed in PBS and cleared with RapiClear® 1.49
(SunJin Lab Co, Hsinchu City, Taiwan) using an established protocol (Bekkouche et al., 2020).

Confocal imaging was performed with a Zeiss LSM 900 inverted confocal microscope (Carl Zeiss,
Oberkochen, Germany) with 405 and 488 nm excitation wavelengths. The 20X /0.8, 25X/0.8 objective
was used for scanning the whole brain in the tile-scan mode. The Plan-Apochromat 25X/0.8 and 63X/1.4
oil objectives were used for scanning MB sub-regions. Images were viewed and processed in Zeiss ZEN

Results

Genome-wide association study (GWAS) on trophallaxis sociability and movement kinematics

We first asked if there are genomic correlates of trophallaxis sociability, utilizing automated tracking of
trophallaxis interactions performed across two replicate trials with experimental colonies composed of
bees from three source colonies: "R2," "R8," and "R41." Colony 1 was composed of bees from R2 and
R41 and Colony 2 was composed of bees from R2 and R8. We generated a ~2.5 million SNP-set from
whole-genome sequencing of 357 bees that we then correlated with trophallaxis sociability. Trophallaxis
sociability was quantified as the average number of trophallaxis interactions per bee across the last two
days of recording, as in [5].

This analysis identified 18 single nucleotide polymorphisms (SNPs) localized across nine chromosomes
(Bonferroni-corrected $P < 0.05$; Fig. 1). Eleven of these SNPs localized to introns of honey bee gene
models [61], one localized to a predicted noncoding RNA (LOC113219153), and the remaining six were
found outside of genes (Fig. 1; Supp. Table S1). Among the 11 intronic SNPs, two were found in the gene
neuroligin-2, two in *glutamate receptor ionotropic, NMDA 2B,* which had robust one-to-one orthology
with the *Drosophila melanogaster* genes *neuroligin-2* and *NMDA receptor 2*, respectively. A single SNP
was found in the following genes, with robust one-to-one orthologs in *D. melanogaster* given in
parentheses: *Angiogenic factor with G patch and FHA domains 1* (CG8079), *CCR4-NOT transcription*
complex subunit 6-like (no hit in *D. melanogaster*), *nephrin* (*sidestep VIII*), *discoidin domain-containing*
receptor 2 (no hit in *D. melanogaster*)*, nuclear factor 1 X-type* (*nuclear factor 1*) and uncharacterized
genes LOC102655706 (*tartan*) and LOC100577268 (no hit in *D. melanogaster*).

In a comparative bioinformatic analysis, n*euroligin-2* (*nlg2*) and *NMDA receptor 2* (*nmdar2*) were found
to be Category 1 ("high confidence") genes in the Simons Foundation Autism Research Initiative (SFARI)
gene database (https://gene.sfari.org/), meaning they have been clearly implicated in human Autism
Spectrum Disorder. Similarly, *nuclear factor 1 X-type* (*Nf1*) is characterized as a "Syndromic" autism
gene, meaning its dysfunction is associated with the development of autistic traits that are not typically
associated with a conventional autism diagnosis.

We performed the same GWAS analysis for two other behaviors: locomotion and turning. Like
trophallaxis, these behaviors require fine motor control and therefore contribute to trophallaxis
interactions, but without obvious social components. (Fig. 1). In contrast to our findings for trophallaxis
sociability, GWAS for these two behaviors did not result in the detection of significantly associated SNPs.

Neuroligin-2 *(*nlg2*) and colony sociability*

Of the three source colonies assayed in this study, individuals from source colony R41 were found to all
be homozygous for the alternative allele (i.e., containing alleles that were different from the reference
locus in the honey bee genome) for the majority of identified SNPs (Fig. 2a; Supp. Fig. S1). In contrast,
individuals from source colonies R2 and R8 were predominantly heterozygous for the more prevalent
reference allele. This result prompted us to investigate the genetics of colony-level variation in behavior.

Trophallaxis sociability significantly varied with source colony (ANOVA: $F_{(2,354)} = 3.165$, $P = 4.24e-16$), and
bees from source colony R41 displayed significantly lower trophallaxis rates than those from source
colony R8 (Tukey HSD post-hoc test, $P = 0.034$) but not R2 ($P = 0.23$; Fig. 2b). In addition, R2 individuals
were significantly less social when co-housed with R41 individuals in Colony 1 than with R8 individuals in
Colony 2 (Wilcoxon rank-sum test: W = 6833.5, $P = 4.17e-11$), further implicating R41 as significantly less
sociable than the other genotypes.

While individuals from source colony R41 demonstrated the lowest levels of trophallaxis sociability, they
were not the slowest moving relative to bees from the other two source colonies. Locomotion was
significantly associated with source colony (ANOVA: $F_{(2,349)}$ = 12.52, P < 4.24e-16), and each colony
significantly differed from one another in this measurement (Tukey HSD post-hoc test, P < 0.005 for all
comparisons). However, movement dynamics could not explain the observed variation in trophallaxis
rate (Fig. 2c).

Social influences on movement in the hive

To explore bee social interactions further, we measured the extent to which a given bee's movement
influenced, and was influenced by, the movement of a nestmate. We implemented information Partial
Directed Coherence (iPDC), which allows for the measuring of bidirectional influences between signals
[38]. To this end, we identified periods of proximal interaction (i.e., when two bees were physically close
to each other), retrieved the X and Y coordinates of each bee relative to the upper-left corner of the
hive, and quantified iPDC from the coordinate time-series of one bee to the coordinate time-series of its
neighbor and vice versa. From iPDC, we then computed information flow (I_{flow}) (Fig. 2d-f; see also
Methods). Higher I_{flow} values indicate higher information transfer from the past location of a bee to the
current location of its neighbor, and we used this metric as measure of social influence. I_{flow} was
approximately normally distributed across bees (Fig. 2g; $I_{flow\ bee->neighbour}$ μ =0.0261 bits, σ=0.0033; I_{flow}
$_{neighbour->bee}$ μ =0.0261 bits, σ=0.0029) and the values observed are consistent with those reported for
other animal species when assessing social influence in movement dynamics [62]. We then asked
whether bees of different colonies exhibited different levels of social influence. Information flow
significantly varied with source colony (Fig. 2h; Supp. Fig. S2; one-way ANOVA $I_{flow\ bee->neighbour}$: $F_{(2,352)}$ =
12.26, P = 7.11e-06; $I_{flow\ neighbour->bee}$: $F_{(2,352)}$ = 7.1, P = 9.51e-4). Post-hoc comparisons revealed no
difference between colonies R2 and R8 (Tukey HSD $I_{flow\ bee->neighbour}$: P = 0.58; $I_{flow\ neighbour->bee}$: P = 0.17). In
contrast, consistent with the trophallaxis results, bidirectional I_{flow} was lowest for individuals from colony
R41 compared to R2 (Tukey HSD $I_{flow\ bee->neighbour}$: P = 1.37e-4; $I_{flow\ neighbour->bee}$: P = 0.045) and R8 (Tukey HSD
$I_{flow\ bee->neighbour}$: P = 7.6e-6; $I_{flow\ neighbour->bee}$: P = 5.69e-4), suggesting that R41 bees show reduced influence
on, and from, their neighbor's movement during time of social proximity.

Mushroom body (MB) transcriptomic profiles and trophallaxis sociability

We next asked how trophallaxis sociability is related to gene expression, focusing on the mushroom
bodies (MB), a higher-order center of sensory integration, decision-making [14], and social regulation
[42]. Unlike GWAS, transcriptome profiling generates arrays of genes that associate with a particular
continuous or discrete phenotypic variable. Deriving lists of genes with expression values correlated
with trophallaxis rate, for example, allows us to test for statistically significant overlap between our
findings and previous studies of other behaviors, such as reward processing. We used RNA-Sequencing
(RNA-Seq) to profile the MB transcriptome of a subset of bees included in the GWAS, selecting
individuals that were reliably identified as generalists, foragers, guards, nurses, or non-responders,
following criteria from previous studies [5,20,40].

Non-responders were transcriptionally distinct from generalists, foragers, guards, and nurses, as in [20] ;
Supp. Tables S2-S6). A GO enrichment performed on genes that were upregulated in all responders
compared to non-responders identified only "chemosensory behavior" and "learning or memory" as
enriched in responders. For genes upregulated in non-responders, only a number of less specific
metabolic terms were identified as enriched (Fig. 3a).

We identified 920 and 785 genes with expression values that positively or negatively correlated with
individual differences in sociability, respectively (Supp. Table S7). Gene Ontology (GO) enrichment
analysis identified distinct terms for each set of genes, with positively associated genes enriched for
several regulatory processes in the context of cellular communication and signaling and negatively
associated genes enriched for processes associated with translation and macromolecule biosynthesis
(Fig. 3b). In contrast to our GWAS results, we also identified a significant overlap of 1108 genes with
expression values that correlated both with trophallaxis sociability and locomotion (hypergeometric
test, fold enrichment = 3.3, P = 2.75e-239; Fig. 3c; Supp. Table S8).

Comparing our data with previously published results, we did not identify a significant overlap of genes
positively or negatively associated with trophallaxis sociability and those associated with food reward or
anticipation of food reward (hypergeometric test, P > 0.10) [63,64].

Neuroligin-2 *RNA-Seq expression and fluorescent* in situ *hybridization*

We compared *nlg2* expression in the MB in non-responders to the average expression in generalists,
foragers, guards, and nurses. Expression levels were significantly associated with trophallaxis sociability
(Spearman's rho = 0.21, P = 0.0063, Fig. 3d), source colony (Tukey HSD post-hoc test on significant
ANOVA [$F_{(2, 169)}$ = 69.762, P < 2e-16], P < 0.10 for each comparison, Fig. 3e), and overall social
responsiveness, when ("responders," Wilcoxon rank sum test, W = 3682, P = 0.0014, Fig. 3f).

Motivated by the association of *nlg2* and trophallaxis sociability in both GWAS (Fig. 1) and RNA-Seq
results (Fig 3d-f), we asked where *nlg2* is expressed in the brain, and if its regional localization overlaps
with regions known to be associated with social decision-making. We collected bees of different ages
and used HCR-FISH to visualize *nlg2* expression. We found *nlg2* localization primarily in neuronal somata
and largely absent from synaptically dense neuropil when compared to control samples where no gene-
specific probes were used (Fig. 4a-b), as expected. The Class I Kenyon cells of the mushroom bodies
(MB) appeared to have higher expression, particularly in the densely packed neurons at the center of
the MB calyces (Fig. 4c), and expression was weaker in the surrounding, more loosely packed cells (Fig.
4d). We also observed expression in MB extrinsic neurons and dorsal midline populations of cells that
compose the pars intercerebralis (Fig. 4e), a small population of large neurosecretory cells vital for
nutritional signaling [65]. Some optic lobe cell populations, especially those most proximal to the central
brain, also showed discernable fluorescence indicating transcription of *nlg2*.

As our study considered 10-day-old bees, we sought to qualitatively assess how spatiotemporal
expression of *nlg2* varied before and after this age. We performed FISH on one-, seven-, and 14-day-old
bees, and found that *nlg2* expression was present in one-day-olds but weak in seven-day-old bees
before returning to discernable levels in 14-day-old bees (Fig. 4f-k).

Discussion

Social proclivities vary within animal societies, driving a continuous distribution of individual tendencies
to associate with others. This is true for many species, from humans to honey bees: some individuals
pursue regular social contact and maintain large networks while others engage less. Social interactions
are multifaceted, as are their drivers, which integrate across physiology, developmental and
motivational state, genetics, and past experience. It thus remains challenging to quantify social
interactions and their molecular underpinnings. We used a combination of high-throughput behavioral

tracking [27,37], laboratory-based behavioral assays [5,42], genome resequencing, brain
transcriptomics, and comparative genomics to explore honey bee sociability and its molecular
underpinnings. We focused on trophallaxis interactions and inter-individual motor coordination. Our
results indicate deep homology in the molecular roots of social behavior shared between honey bees
and vertebrates while also revealing divergent mechanisms.

Our genome-wide association study (GWAS) identified 18 single nucleotide polymorphisms (SNPs)
associated with variation in trophallaxis sociability. In contrast, no SNPs were identified when a GWAS
was performed on locomotion or turning rate, kinematic metrics that contribute to trophallaxis. Like any
social interaction, trophallaxis is more than the sum of its parts: rapid movement may support more
social interactions but our results indicate it does not alone drive variation in tendency to interact in this
way. By standards of human GWAS, our sample size (N = 357 individuals) was small, but similar numbers
of sociability-associated SNPs have been identified from GWAS with much larger human cohorts [25],
reflecting the complexity of factors underpinning sociability.

Over half of the identified SNPs were localized to the intronic regions of genes. Using the Simons
Foundation Autism Research Initiative (SFARI) gene database (https://gene.sfari.org/), we identified
neuroligin-2 (*nlg2*) and *NMDA receptor 2* (*nmdar2*) as Category 1 ("high confidence") genes, meaning
they have been strongly implicated in human autism. Another gene with an intronic SNP, *nuclear factor*
1 X-type (*Nf1*) is characterized as a "Syndromic" autism gene, meaning its dysfunction is associated with
the development of autistic traits that are not typically associated with a conventional autism diagnosis.
In humans, impaired glutamatergic signaling by way of NMDAR disruption has clear associations with
the development of autism [66], while glutamate is best known in insects for its role at the
neuromuscular junction [67]. However, NMDAR activity is causally associated with memory formation in
honey bees [68], and its function may therefore contribute to the reinforcement of trophallaxis
relationships, which ultimately rely on reciprocity between nestmates.

The identification of two *nlg2* SNPs associated with variation in trophallaxis sociability contributes to a
comparative framework for understanding the molecular building blocks of social behavior across highly
diverged species. Neuroligins are cell surface proteins involved in the formation and remodeling of
synapses in the central nervous system in an isoform-specific manner, and *nlg2* acts exclusively at
inhibitory synapses [69]. In vertebrates, genomic variation in, or disrupted expression of, *nlg2* or its
interaction partners is associated with forms of neurodivergence like autism, schizophrenia, and anxiety,
each of which is linked to differences in how individuals associate with their social environment
[2,70,71]. In *Drosophila melanogaster*, social behavior also is affected by mutations in *nlg2* [72].

Several correlative findings suggest a role for *nlg2* in honey bee trophallaxis sociability. First, the two
intronic variants (SNPs) in the honey bee *nlg2* gene that we identified were predominantly restricted to
individuals from one source colony, and those individuals displayed low levels of trophallaxis sociability
compared to those from other source colonies. We probed other kinematic aspects of these bees and
found that this lower level of sociability was not due to slower movement. Second, *nlg2* expression
levels in the mushroom bodies (MB), a higher-order sensory integration center in the bee brain [14,73],
were also significantly correlated with trophallaxis rate and social responsiveness across all three tested
colonies. Third, a strong signal of *nlg2* expression was localized to the MB and appeared to overlap the
pars intercerebralis, a set of neurosecretory cells that mediate the relationship between diet, nutrition,
and division of labor [65]. Trophallaxis sociability in honey bees illustrates the importance of food-

sharing as a foundation for animal society. We speculate that *nlg2* may have played a role in the shaping
of ancestral feeding circuitry to drive stable social network formation in this social bee lineage, as may
be the case for other social animals. Consuming but not fully digesting food requires a neurobiological
change to bypass the rewarding nature of feeding events, yet such a mechanism provides a foundation
for the formation of long-term social bonds [74].

We also explored the synchronization of body movement between pairs of individual bees, some not
necessarily engaged in trophallaxis. This model of social influence is similar to trophallaxis in that it
requires coordination of one individual's body with another among honey bees located close to each
other, but also directly quantifies a measure of social influence identified in other species [62]. We
found that the *ngl2* allele associated with reduced trophallaxis also was associated with reduced motor
synchrony between bees. This means that the movement of bees in the colony with high frequencies of
this allele had less influence on, and were less influenced by, the movement of other bees. Interpersonal
synchrony characterizes temporally or physically aligned behaviors in humans that naturally emerge
between individuals in a social group, and such bidirectional motor influence tends to be diminished
among autistic individuals and their peers [75,76]. It is interesting to note that casual observations
revealed no motor deficits in the colony with high frequencies of this allele and it is therefore unclear if
negative fitness consequences exist. It may be that such variation is well-tolerated within the honey bee
population in the context of colony performance.

Social interactions help establish and reinforce group membership and, in the context of food-sharing,
facilitate mutually rewarding altruistic bonds between individuals [77,78]. In this regard, we predicted
trophallaxis to be an intrinsically rewarding behavior, as is honey bee dance communication [63].
However, this prediction was not supported by comparative transcriptomic analysis. We did not identify
gene expression similarities between the present study and previous MB transcriptomic characterization
of reward processing [63,64]. Perhaps this finding is related to the fact that dopaminergic neurons,
which contribute to reward seeking, expectation and acquisition in honey bees [79,80] are sparsely
distributed in small clusters throughout the bee brain [81,82], earlier findings corroborated by more
recent single-cell brain analyses [83,84]. Perhaps the bulk sequencing of the MB performed in our study
both diluted and excluded dopaminergic neurons. It may also be that food donation versus reception,
which we did not distinguish in this study, supports different brain gene expression signatures.

While we identified significant variation in social proclivities across and within honey bee colonies, and
have started to elucidate its molecular roots, the effects of this variation on colony health and well-
being are unknown. Basic evolutionary theory supports the speculation that socially divergent
phenotypes are either tolerated or preserved across diverse animal lineages due to supportive, or
perhaps even vital, roles they contribute to their society. It could be useful to explore this issue in honey
bees, given serious concerns about honey bee health and worldwide colony population declines over
the past two decades.

The expression of individual differences in social behavior, from humans to honeybees, is derived from a
constellation of interacting neurobiological and physiological factors. While no single gene or genetic
variant alone drives sociability, the identification of shared genomic underpinnings of social interactivity
across diverse taxa hints at conserved molecular building blocks of social life. We anticipate new tools
for automated tracking will continue to improve the granularity of such behavioral comparisons across

social animals [28,37], further promoting a role for social insects in elucidating how the brain and
genome have evolved to support the intricate social lives observed across diverse animal societies.

Acknowledgments

The authors thank A. Hernandez and C. Wright at the Roy J. Carver Biotechnology Center (UIUC) for
sequencing services and members of the Robinson and Cook labs for valuable comments and feedback
that improved the manuscript. This work was supported by the European Union's Horizon 2020
Research and Innovation Program under ERC-2017-StG Grant Agreement 757583 (Brain2Bee; to JLC and
GER). IMT is presently supported by the Lewis-Sigler Institute for Integrative Genomics as a Lewis-Sigler
Scholar.

Figures

Figure 1. Genomic correlates of honey bee trophallaxis sociability. (Top) Genome-wide association
study (GWAS) identified 17 single nucleotide polymorphisms (SNPs) significantly associated with
variation in trophallaxis (food-sharing) interactions, 12 of which were localized to the introns of gene
models in the honey bee genome (HAv3.1). In vertebrates, variation in the expression or structure of
three of these genes, *neuroligin-2*, *NMDA receptor 2*, and *Nuclear factor I* is associated with autism per
the Simons Foundation Autism Research Initiative (SFARI) database. (Middle and bottom) In contrast, we
did not identify any SNPs associated with variation in locomotion or turning kinematics, movements that
contribute to, but do not fully encompass, trophallaxis interactions. Dashed red line indicates
significance threshold (Bonferroni-corrected *P*-value < 0.05).

Figure 2. Genotype-specific variation in motor synchrony. (a) A single source colony, "R41," was found to be homozygous for *nlg2* as well as the majority of trophallaxis-associated SNPs, while other colonies in this experiment contained the more prevalent reference allele. (b) Colony "R41" also demonstrated the lowest levels of trophallaxis sociability, yet (c) was not the slowest-moving colony. (d) Images of proximal interaction were selected where two bees were physically close to each other ($R \leq 2$ cm, where R is the Euclidean distance between the two bees' barcodes). The X and Y coordinates of the barcode

represent a given bee's position in Euclidian space. (e) Drawing representing the dataset to which iPDC
was applied. Each proximal interaction (interaction 1, … interaction n) is represented as a bivariate time-
series of at least 30 time points (30 sec), where the first time-series is the focal bee's X or Y coordinate
and the second time-series is its neighbor's X *or* Y coordinate, respectively. For each focal bee, the
identity of the neighbor varied across interactions. iPDC was calculated within each paired time-series
and averaged across interactions to subsequently compute information flow in both directions. (f)
Example of two bees with different levels of social influence. Arrow thickness is proportional to
information transfer. On average, Bee #1697 influenced the movement of its neighbor with $I_{flow\ bee\text{-}>neighbor}$
= 0.035 bits, and was influenced by the movement of its neighbor with $I_{flow\ neighbor\text{-}>bee}$ = 0.019
bits. (g) Distributions of information flow across bees, representing the magnitude of movement
influence-related information. (h) As for trophallaxis sociability, "R41" bees were also least likely to
influence, or be influenced by, other bees in their experimental cohort in terms of motor synchrony.
Violin plots are constructed as follows: points represent raw data, solid black lines represent the mean,
pale white shading above and below the mean represent a 95% confidence interval, and plot shape
represents a smoothed density curve outlining the distribution of raw data. Letters above violin plots
represent significance (P-value < 0.05) from a between-group Tukey Post-Hoc analysis following a one-
way ANOVA.

Figure 3. Neuromolecular correlates of trophallaxis sociability and social responsiveness in the honey bee mushroom bodies (MB). (a) Gene Ontology (GO) analysis identified highly specific terms associated with chemosensation and learning enriched in genes upregulated in Responders, while generic metabolic process terms were enriched in genes upregulated in NonResponders. (b) GO enrichment analysis performed on genes positively correlated with trophallaxis frequency revealed the enrichment

of several terms related to cell signaling and stimulus response, while genes negatively associated with
trophallaxis were more strongly related to biosynthesis and metabolism, and (c) this gene set was
significantly similar to genes with expression levels correlated with locomotion (hypergeometric overlap
test). (d) *Nlg2* levels were weakly but significantly correlated with trophallaxis frequency, (e) significantly
lowest in colony "R41" (one-way ANOVA with Tukey post-hoc test), and (f) differentially expressed in
Responders compared to NonResponders in a lab-based assay examining behavioral responsiveness to
social stimuli (Wilcoxon rank-sum test). For GO plots, circle diameter is inversely correlated to term
specificity in the GO hierarchy, and more similar terms are more closely clustered in semantic space.
Violin plots are constructed as previously described.

Figure 4. Whole-mount fluorescent *in situ* hybridization (smFISH) reveals *neuroligin-2* (*nlg2*) expression patterns in the developing adult honey bee brain. (a) *Nlg2* is widely expressed in the adult worker brain as detected via HCR probes, while (b) no staining was observed with antisense probes. (c) Expression levels were highest in the mushroom body (MB) Kenyon cells (KC), specifically the inner-compact KC (dashed white line), yet (d) were observed in the outer-compact KC as well. (e) We also identified strong signaling along the dorsal midline of the brain, in or around the neurosecretory cells that compose the pars intercerebralis. (f-k) *Nlg2* expression was apparent throughout adulthood in the worker honey bee brain, though levels appeared highest in callows and two-week-old bees (left and right columns are color and greyscale depictions of the same image, respectively). For each panel, *nlg2* localization is shown in green and nuclei counterstained with DAPI are shown in blue.

Supplementary Figures

Supplementary Figure S1. Zygosity for trophallaxis-associated single-nucleotide polymorphisms (SNPs). All SNPs are described in Supplementary Table S1; 17 out of 18 SNPs are shown, with the remaining SNP shown in Fig. 2a.

Supplementary Figure S2. Genotype-specific variation in motor synchrony. iPDC was applied to pairs of bees' X and Y coordinate time-series, obtaining information flow (Iflow) between the movement of a bee and its neighbor along the X and Y axis, respectively. Iflow was then averaged across the two coordinates obtaining a single Iflow for each bee (Fig. 2h). Violin plots are constructed as follows: points represent raw data, solid black lines represent the mean, pale white shading above and below the mean represent a 95% confidence interval, and plot shape represents a smoothed density curve outlining the distribution of raw data. Letters above violin plots represent significance (P-value < 0.05) from a between-group Tukey Post-Hoc analysis following a one-way ANOVA (for X coordinate $I_{flow\ bee->neighbour}$: $F_{(2,352)} = 8.03$, $P = 3.87$e-04, $I_{flow\ neighbour->bee}$: $F_{(2,352)} = 7.02$, $P = 0.001$); for Y coordinate $I_{flow\ bee->neighbour}$: $F_{(2,352)}$ = 6.92, $P = 0.001$, $I_{flow\ neighbour->bee}$: $F_{(2,352)} = 1.4$, $P = 0.248$). Tukey HSD for X coordinate $I_{flow\ bee->neighbour}$: $P = 0.58$; $I_{flow\ neighbour->bee}$: $P = 0.17$).

Supplementary Tables

Supplementary Table S1: Annotations and test statistics for 18 single-nucleotide polymorphisms (SNPs) associated with variation in trophallaxis sociability via genome-wide association study (GWAS).

Supplementary Table S2: Differentially expressed genes detected in non-responders relative to remaining behavioral groups.

Supplementary Table S3: Differentially expressed genes detected in generalists relative to remaining behavioral groups.

Supplementary Table S4: Differentially expressed genes detected in foragers relative to remaining behavioral groups.

Supplementary Table S5: Differentially expressed genes detected in guards relative to remaining behavioral groups.

Supplementary Table S6: Differentially expressed genes detected in nurses relative to remaining behavioral groups.

Supplementary Table S7: Genes with expression levels that correlate with trophallaxis sociability.

Supplementary Table S8: Genes with expression levels that correlate with locomotion.

References

1. Gartland LA, Firth JA, Laskowski KL, Jeanson R, Ioannou CC. Sociability as a personality trait in animals:
methods, causes and consequences. Biol Rev. 2022;97: 802–816. doi:10.1111/brv.12823

2. Barak B, Feng G. Neurobiology of social behavior abnormalities in autism and Williams syndrome. Nat
Neurosci. 2016;19: 647–655. doi:10.1038/nn.4276

3. Bonnell TR, Vilette C, Henzi SP, Barrett L. Network reaction norms: taking account of network position
and plasticity in response to environmental change. Behav Ecol Sociobiol. 2023;77: 35.
doi:10.1007/s00265-023-03300-2

4. Friedman DA, Johnson BR, Linksvayer TA. Distributed physiology and the molecular basis of social life
in eusocial insects. Horm Behav. 2020;122: 104757. doi:10.1016/j.yhbeh.2020.104757

5. Traniello IM, Hamilton AR, Gernat T, Cash-Ahmed AC, Harwood G, Ray AM, et al. Context-dependent
influence of threat on honey bee social network dynamics and brain gene expression. J Exp Biol.
2022;225. doi:10.1242/jeb.243738

6. Taylor SE, Klein LC, Lewis BP, Gruenewald TL, Gurung RAR, Updegraff JA. Biobehavioral responses to
stress in females: Tend-and-Befriend, not Fight-or-Flight. Psychol Rev. 2000;107: 411–429.
doi:10.1037/0033-295x.107.3.411

7. López-Tobón A, Trattaro S, Testa G. The sociability spectrum: evidence from reciprocal genetic copy
number variations. Mol Autism. 2020;11: 50. doi:10.1186/s13229-020-00347-0

8. Fischer EK, O'Connell LA. Hormonal and neural correlates of care in active versus observing poison
frog parents. Horm Behav. 2020;120: 104696. doi:10.1016/j.yhbeh.2020.104696

9. Kabelik D, Weitekamp CA, Choudhury SC, Hartline JT, Smith AN, Hofmann HA. Neural activity in the
social decision-making network of the brown anole during reproductive and agonistic encounters. Horm
Behav. 2018;106: 178–188. doi:10.1016/j.yhbeh.2018.06.013

10. Johnson ZV, Hegarty BE, Gruenhagen GW, Lancaster TJ, McGrath PT, Streelman JT. Cellular profiling
of a recently-evolved social behavior in cichlid fishes. Nat Commun. 2023;14: 4891. doi:10.1038/s41467-
023-40331-9

11. Newman SW. The medial extended amygdala in male reproductive behavior: A node in the
mammalian social behavior network. Ann N York Acad Sci. 1999;877: 242–257. doi:10.1111/j.1749-
6632.1999.tb09271.x

12. O'Connell LA, Hofmann HA. The Vertebrate mesolimbic reward system and social behavior network:
A comparative synthesis. J Comp Neurol. 2011;519: 3599–3639. doi:10.1002/cne.22735

13. O'Connell LA, Hofmann HA. Evolution of a vertebrate social decision-making network. Science.
2012;336: 1154–1157. doi:10.1126/science.1218889

14. Strausfeld NJ. Arthropod Brains: Evolution, Functional Elegance, and Historical Significance. Harvard
University Press; 2012.

15. Wilson EO. The insect societies. Cambridge, MA: Harvard University Press; 1971.

16. Page RE, Robinson GE. The genetics of division of labour in honey bee colonies. Adv Insect Physiol.
1991;23: 117–169. doi:10.1016/s0065-2806(08)60093-4

17. Whitfield CW, Cziko A-M, Robinson GE. Gene expression profiles in the brain predict behavior in
individual honey bees. Science. 2003;302: 296–299. doi:10.1126/science.1086807

18. Zayed A, Robinson GE. Understanding the relationship between brain gene expression and social
behavior: Lessons from the honey bee. Genetics. 2012;46: 591–615. doi:10.1146/annurev-genet-
110711-155517

19. Cook JL, Robinson GE. Comparative genomics and the roots of human behavior. Trends Cogn Sci.
2023;27: 230–232. doi:10.1016/j.tics.2022.12.012

20. Shpigler HY, Saul MC, Corona F, Block L, Ahmed AC, Zhao SD, et al. Deep evolutionary conservation
of autism-related genes. Proc Natl Acad Sci. 2017;114: 9653–9658. doi:10.1073/pnas.1708127114

21. Rittschof CC, Bukhari SA, Sloofman LG, Troy JM, Caetano-Anollés D, Cash-Ahmed A, et al.
Neuromolecular responses to social challenge: Common mechanisms across mouse, stickleback fish, and
honey bee. Proc Natl Acad Sci. 2014;111: 17929–17934. doi:10.1073/pnas.1420369111

22. Traniello IM, Chen Z, Bagchi VA, Robinson GE. Valence of social information is encoded in different
subpopulations of mushroom body Kenyon cells in the honeybee brain. Proc R Soc B. 2019;286:
20190901. doi:10.1098/rspb.2019.0901

23. Arechavaleta-Velasco ME, Alcala-Escamilla K, Robles-Rios C, Tsuruda JM, Hunt GJ. Fine-Scale linkage
mapping reveals a small set of candidate genes influencing honey bee grooming behavior in response to
varroa mites. PLoS ONE. 2012;7: e47269. doi:10.1371/journal.pone.0047269

24. Kocher SD, Mallarino R, Rubin BER, Yu DW, Hoekstra HE, Pierce NE. The genetic basis of a social
polymorphism in halictid bees. Nat Commun. 2018;9: 4338. doi:10.1038/s41467-018-06824-8

25. Bralten J, Mota NR, Klemann CJHM, Witte WD, Laing E, Collier DA, et al. Genetic underpinnings of
sociability in the general population. Neuropsychopharmacology. 2021;46: 1627–1634.
doi:10.1038/s41386-021-01044-z

26. Gal A, Saragosti J, Kronauer DJ. anTraX, a software package for high-throughput video tracking of
color-tagged insects. eLife. 2020;9: e58145. doi:10.7554/elife.58145

27. Gernat T, Rao VD, Middendorf M, Dankowicz H, Goldenfeld N, Robinson GE. Automated monitoring
of behavior reveals bursty interaction patterns and rapid spreading dynamics in honeybee social
networks. Proc Natl Acad Sci. 2018;115: 1433–1438. doi:10.1073/pnas.1713568115

28. Traniello IM, Kocher SD. Integrating computer vision and molecular neurobiology to bridge the gap
between behavior and the brain. Curr Opin Insect Sci. 2024;66: 101259. doi:10.1016/j.cois.2024.101259

29. Wolf SW, Ruttenberg DM, Knapp DY, Webb AE, Traniello IM, McKenzie-Smith GC, et al. NAPS:
Integrating pose estimation and tag-based tracking. Methods Ecol Evol. 2023. doi:10.1111/2041-
210x.14201

30. Geffre AC, Gernat T, Harwood GP, Jones BM, Gysi DM, Hamilton AR, et al. Honey bee virus causes
context-dependent changes in host social behavior. Proc Natl Acad Sci. 2020;117: 10406–10413.
doi:10.1073/pnas.2002268117

31. Jones BM, Rao VD, Gernat T, Jagla T, Cash-Ahmed AC, Rubin BE, et al. Individual differences in honey
bee behavior enabled by plasticity in brain gene regulatory networks. eLife. 2020;9: e62850.
doi:10.7554/elife.62850

32. Wild B, Dormagen DM, Zachariae A, Smith ML, Traynor KS, Brockmann D, et al. Social networks
predict the life and death of honey bees. Nat Commun. 2021;12: 1110. doi:10.1038/s41467-021-21212-5

33. Wcislo WT. Trophallaxis in weakly social bees (Apoidea). Ecol Èntomol. 2016;41: 37–39.
doi:10.1111/een.12289

34. Choi SH, Rao VD, Gernat T, Hamilton AR, Robinson GE, Goldenfeld N. Individual variations lead to
universal and cross-species patterns of social behavior. Proc Natl Acad Sci. 2020;117: 31754–31759.
doi:10.1073/pnas.2002013117

35. Glass D, Yuill N. Social motor synchrony in autism spectrum conditions: A systematic review. Autism.
2024;28: 1638–1653. doi:10.1177/13623613231213295

36. Glass D, Yuill N. Evidence of mutual non-verbal synchrony in learners with severe learning disability
and autism, and their support workers: a motion energy analysis study. Front Integr Neurosci. 2024;18:
1353966. doi:10.3389/fnint.2024.1353966

37. Gernat T, Jagla T, Jones BM, Middendorf M, Robinson GE. Automated monitoring of honey bees with
barcodes and artificial intelligence reveals two distinct social networks from a single affiliative behavior.
Sci Rep. 2023;13: 1541. doi:10.1038/s41598-022-26825-4

38. Takahashi DY, Baccalá LA, Sameshima K. Information theoretic interpretation of frequency domain
connectivity measures. Biol Cybern. 2010;103: 463–469. doi:10.1007/s00422-010-0410-x

39. Sameshima K, Baccalá LA. Methods in brain connectivity inference through multivariate time series
analysis. CRC Press; 2014.

40. Hamilton AR, Traniello IM, Ray AM, Caldwell AS, Wickline SA, Robinson GE. Division of labor in honey
bees is associated with transcriptional regulatory plasticity in the brain. J Exp Biol. 2019;222: jeb200196.
doi:10.1242/jeb.200196

41. Shpigler HY, Saul MC, Murdoch EE, Cash-Ahmed AC, Seward CH, Sloofman L, et al. Behavioral,
transcriptomic and epigenetic responses to social challenge in honey bees. Genes, Brain Behav. 2017;16:
579–591. doi:10.1111/gbb.12379

42. Shpigler HY, Saul MC, Murdoch EE, Corona F, Cash-Ahmed AC, Seward CH, et al. Honey bee
neurogenomic responses to affiliative and agonistic social interactions. Genes, Brain Behav. 2019;18:
e12509. doi:10.1111/gbb.12509

43. Shpigler HY, Robinson GE. Laboratory assay of brood care for quantitative analyses of individual
differences in honey bee (Apis mellifera) affiliative behavior. PLoS ONE. 2015;10: e0143183.
doi:10.1371/journal.pone.0143183

44. Wilson ML. The biology of the honey bee. Harvard University Press; 1991.

45. Robinson GE, Jr. REP, Strambi C, Strambi A. Hormonal and genetic control of behavioral integration
in honey bee colonies. Science. 1989;246: 109–112. doi:10.1126/science.246.4926.109

46. Giray T, Robinson GE. Effects of intracolony variability in behavioral development on plasticity of
division of labor in honey bee colonies. Behav Ecol Sociobiol. 1994;35: 13–20. doi:10.1007/bf00167054

47. Huang Z-Y, Robinson GE. Regulation of honey bee division of labor by colony age demography. Behav
Ecol Sociobiol. 1996;39: 147–158. doi:10.1007/s002650050276

48. Johnson BR. Honey Bee Biology. Princeton University Press; 2023.

49. Lutz CC, Robinson GE. Activity-dependent gene expression in honey bee mushroom bodies in
response to orientation flight. J Exp Biol. 2013;216: 2031–2038. doi:10.1242/jeb.084905

50. Browning SR, Browning BL. Rapid and accurate haplotype phasing and missing-data inference for
whole-genome association studies by use of localized haplotype clustering. Am J Hum Genet. 2007;81:
1084–1097. doi:10.1086/521987

51. Browning BL, Zhou Y, Browning SR. A one-penny imputed genome from next-generation reference
panels. Am J Hum Genet. 2018;103: 338–348. doi:10.1016/j.ajhg.2018.07.015

52. Zheng X, Levine D, Shen J, Gogarten SM, Laurie C, Weir BS. A high-performance computing toolset
for relatedness and principal component analysis of SNP data. Bioinformatics. 2012;28: 3326–3328.
doi:10.1093/bioinformatics/bts606

53. Gogarten SM, Sofer T, Chen H, Yu C, Brody JA, Thornton TA, et al. Genetic association testing using
the GENESIS R/Bioconductor package. Bioinformatics. 2019;35: 5346–5348.
doi:10.1093/bioinformatics/btz567

54. Avalos A, Fang M, Pan H, Lluch AR, Lipka AE, Zhao SD, et al. Genomic regions influencing aggressive
behavior in honey bees are defined by colony allele frequencies. Proc Natl Acad Sci. 2020;117: 17135–
17141. doi:10.1073/pnas.1922927117

55. Wallberg A, Bunikis I, Pettersson OV, Mosbech M-B, Childers AK, Evans JD, et al. A hybrid de novo
genome assembly of the honeybee, *Apis mellifera*, with chromosome-length scaffolds. BMC Genom.
2019;20: 275. doi:10.1186/s12864-019-5642-0

56. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal RNA-seq
aligner. Bioinformatics. 2013;29: 15–21. doi:10.1093/bioinformatics/bts635

57. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for assigning sequence
reads to genomic features. Bioinformatics. 2014;30: 923–930. doi:10.1093/bioinformatics/btt656

58. Traniello IM, Bukhari SA, Kevill J, Ahmed AC, Hamilton AR, Naeger NL, et al. Meta-analysis of honey
bee neurogenomic response links Deformed wing virus type A to precocious behavioral maturation. Sci
Rep. 2020;10: 3101. doi:10.1038/s41598-020-59808-4

59. Eden E, Navon R, Steinfeld I, Lipson D, Yakhini Z. GOrilla: a tool for discovery and visualization of
enriched GO terms in ranked gene lists. BMC Bioinform. 2009;10: 48. doi:10.1186/1471-2105-10-48

60. Reijnders MJMF, Waterhouse RM. Summary visualizations of Gene Ontology terms with GO-Figure!
Front Bioinform. 2021;1: 638255. doi:10.3389/fbinf.2021.638255

61. Elsik CG, Worley KC, Bennett AK, Beye M, Camara F, Childers CP, et al. Finding the missing honey bee
genes: lessons learned from a genome upgrade. BMC Genom. 2014;15: 86. doi:10.1186/1471-2164-15-
86

62. Gachomba MJM, Esteve-Agraz J, Caref K, Maroto AS, Bortolozzo-Gleich MH, Laplagne DA, et al.
Multimodal cues displayed by submissive rats promote prosocial choices by dominants. Curr Biol.
2022;32: 3288-3301.e8. doi:10.1016/j.cub.2022.06.026

63. McNeill MS, Kapheim KM, Brockmann A, McGill TAW, Robinson GE. Brain regions and molecular
pathways responding to food reward type and value in honey bees. Genes, Brain Behav. 2016;15: 305–
317. doi:10.1111/gbb.12275

64. Naeger NL, Robinson GE. Transcriptomic analysis of instinctive and learned reward-related behaviors
in honey bees. J Exp Biol. 2016;219: 3554–3561. doi:10.1242/jeb.144311

65. Wheeler MM, Ament SA, Rodriguez-Zas SL, Southey B, Robinson GE. Diet and endocrine effects on
behavioral maturation-related gene expression in the pars intercerebralis of the honey bee brain. J Exp
Biol. 2015;218: 4005–4014. doi:10.1242/jeb.119420

66. Lee E-J, Choi SY, Kim E. NMDA receptor dysfunction in autism spectrum disorders. Curr Opin
Pharmacol. 2015;20: 8–13. doi:10.1016/j.coph.2014.10.007

67. Jan LY, Jan YN. L-glutamate as an excitatory transmitter at the *Drosophila* larval neuromuscular
junction. J Physiol. 1976;262: 215–236. doi:10.1113/jphysiol.1976.sp011593

68. Si A, Helliwell P, Maleszka R. Effects of NMDA receptor antagonists on olfactory learning and
memory in the honeybee (Apis mellifera). Pharmacol Biochem Behav. 2004;77: 191–197.
doi:10.1016/j.pbb.2003.09.023

69. Varoqueaux F, Jamain S, Brose N. Neuroligin 2 is exclusively localized to inhibitory synapses. Eur J
Cell Biol. 2004;83: 449–456. doi:10.1078/0171-9335-00410

70. Ali H, Marth L, Krueger-Burg D. Neuroligin-2 as a central organizer of inhibitory synapses in health
and disease. Sci Signal. 2020;13: eabd8379. doi:10.1126/scisignal.abd8379

71. Parente DJ, Garriga C, Baskin B, Douglas G, Cho MT, Araujo GC, et al. Neuroligin 2 nonsense variant
associated with anxiety, autism, intellectual disability, hyperphagia, and obesity. Am J Méd Genet Part A.
2017;173: 213–216. doi:10.1002/ajmg.a.37977

72. Corthals K, Heukamp AS, Kossen R, Großhennig I, Hahn N, Gras H, et al. Neuroligins Nlg2 and Nlg4
affect social behavior in *Drosophila melanogaster*. Front Psychiatry. 2017;8: 113.
doi:10.3389/fpsyt.2017.00113

73. Paffhausen BH, Fuchs I, Duer A, Hillmer I, Dimitriou IM, Menzel R. Neural correlates of social
behavior in mushroom body extrinsic neurons of the honeybee *Apis mellifera*. Front Behav Neurosci.
2020;14: 62. doi:10.3389/fnbeh.2020.00062

74. Fischer EK, O'Connell LA, Levine JD, Kronauer DJC, Dickinson MH. Modification of feeding circuits in
the evolution of social behavior. J Exp Biol. 2017;220: 92–102. doi:10.1242/jeb.143859

75. Fitzpatrick P, Romero V, Amaral JL, Duncan A, Barnard H, Richardson MJ, et al. Evaluating the
importance of social motor synchronization and motor skill for understanding autism. Autism Res.
2017;10: 1687–1699. doi:10.1002/aur.1808

76. McNaughton KA, Redcay E. Interpersonal synchrony in autism. Curr Psychiatry Rep. 2020;22: 12.
doi:10.1007/s11920-020-1135-8

77. Dridi S, Akçay E. Learning to cooperate: The evolution of social rewards in repeated interactions. Am
Nat. 2018;191: 58–73. doi:10.1086/694822

78. Walter H, Abler B, Ciaramidaro A, Erk S. Motivating forces of human actions: Neuroimaging reward
and social interaction. Brain Res Bull. 2005;67: 368–381. doi:10.1016/j.brainresbull.2005.06.016

79. Huang J, Zhang Z, Feng W, Zhao Y, Aldanondo A, Sanchez MG de B, et al. Food wanting is mediated
by transient activation of dopaminergic signaling in the honey bee brain. Science. 2022;376: 508–512.
doi:10.1126/science.abn9920

80. Liang ZS, Nguyen T, Mattila HR, Rodriguez-Zas SL, Seeley TD, Robinson GE. Molecular determinants
of scouting behavior in honey bees. Science. 2012;335: 1225–1228. doi:10.1126/science.1213962

81. Lehman HK, Schulz DJ, Barron AB, Wraight L, Hardison C, Whitney S, et al. Division of labor in the
honey bee (*Apis mellifera*): the role of tyramine β-hydroxylase. J Exp Biol. 2006;209: 2774–2784.
doi:10.1242/jeb.02296

82. Tedjakumala SR, Rouquette J, Boizeau M-L, Mesce KA, Hotier L, Massou I, et al. A tyrosine-
hydroxylase characterization of dopaminergic neurons in the honey bee brain. Front Syst Neurosci.
2017;11: 47. doi:10.3389/fnsys.2017.00047

83. Li Q, Wang M, Zhang P, Liu Y, Guo Q, Zhu Y, et al. A single-cell transcriptomic atlas tracking the neural
basis of division of labour in an ant superorganism. Nat Ecol Evol. 2022;6: 1191–1204.
doi:10.1038/s41559-022-01784-1

84. Sheng L, Shields EJ, Gospocic J, Glastad KM, Ratchasanmuang P, Berger SL, et al. Social
reprogramming in ants induces longevity-associated glia remodeling. Sci Adv. 2020;6: eaba9869.
doi:10.1126/sciadv.aba9869

889