Endocytic Patch Dynamics are Differentially Regulated at Distinct Cell Sites in Fission

Yeast

Bethany F. Campbell[1], Uma J. Patel[1], Ashlei R. Williams[2], and Maitreyi E. Das[1]*

[1]Boston College, Chestnut Hill, MA

[2]The University of Tennessee, Knoxville, TN

Correspondence: *dasma@bc.edu

Running Title: Cdc42 regulates branched actin

Keywords: Polarization, Branched Actin Networks, Clathrin-mediated Endocytosis, Cdc42

GTPase, p21-Activated Kinase, Type I Myosin

ABSTRACT

Endocytosis promotes polarity and growth in eukaryotes. In *Schizosaccharomyces pombe*

fission yeast*,* endocytosis occurs at the polarized cell ends and division site and at the non-

polarized cell sides. Our characterization of endocytic actin patches show that they are

differentially regulated. The patches at the cell ends and division site internalize successfully

while those at the sides are weak and erratic. The major regulator of cell polarity, Cdc42, and its

target Pak1 kinase only localize to the cell ends and division site. We find that these proteins

regulate assembly and internalization of patches at these sites but not at the cell sides.

Moreover, Cdc42 specifically activated by the GEF Gef1 promotes proper patch dynamics.

Endocytosis requires phosphorylation of the Type I Myosin Myo1 by the Pak1 kinase. Myo1

localizes to the cell ends, division site, and the cell sides. We find that unlike Cdc42 and Pak1,

Myo1 also promotes patch assembly at the cell sides. Our data indicate that while Myo1 can

globally promote branched actin assembly, successful endocytic patch dynamics and

internalization at polarized sites require Cdc42 and Pak1 kinase.

SUMMARY STATEMENT

Endocytic patch dynamics are differentially regulated at distinct sites such as the cell ends,

division site and the cell sides by Cdc42 and its downstream targets Pak1 kinase and the Type

1 myosin.

INTRODUCTION

Endocytosis is an essential process by which cells uptake nutrients and recycle proteins from the plasma membrane. In cell-walled organisms such as the fission yeast *Schizosaccharomyces pombe*, internal turgor pressure is high at ~10 atm (Basu et al., 2014; Lacy et al., 2018), roughly equivalent to a racing bike tire. To overcome such high turgor pressure, successful clathrin-mediated endocytosis in yeasts requires synthesis of branched actin networks to produce enough counteractive force to invaginate the plasma membrane and subsequently internalize endocytic vesicles (Aghamohammadzadeh and Ayscough, 2009; Basu et al., 2014; Nickaeen et al., 2022). In *S. pombe*, branched actin is formed by the Arp2/3 complex with the help of nucleation promoting factors such as the Type I myosin Myo1 and the WASP Wsp1 (Sirotkin et al., 2005). Actin binding proteins such as fimbrin Fim1 also localize to endocytic patches to crosslink the branched actin and stimulate force production within the patch (Skau et al., 2011). Furthermore, if endocytosis is impaired, cells cannot grow or divide. Once treated with the Arp2/3 complex inhibitor, CK-666, *S. pombe* cells halt their growth during interphase and fail to form a septum during cytokinesis (Onwubiko et al., 2019). Thus, regulation of branched actin formation is an essential cellular process for *S. pombe* cell viability.

In rod-shaped *S. pombe* cells, clathrin-mediated endocytic events primarily occur at sites of polarization, namely the cell ends during interphase and the division site during cytokinesis (Gachet and Hyams, 2005; Sirotkin et al., 2010). However, some endocytic actin patches also form at the non-growing cell sides. Previous work has shown that sites of endocytosis coincide with regions of cell growth and cytokinesis (Gachet and Hyams, 2005). The GTPase Cdc42 is the major regulator of polarity in *S. pombe* and is active only at polarized cell ends and the division site and not at the cell sides (Das et al., 2012). Additionally, Cdc42 is activated at the division site during actomyosin ring formation and persists throughout cytokinesis (Hercyk and

Das, 2019a; Wei et al., 2016). Further work has connected Cdc42 activation to Arp2/3-mediated

endocytosis since both are required for septum formation and onset of actomyosin ring

constriction in fission yeast (Campbell et al., 2022; Onwubiko et al., 2019). Studies in

mammalian MDCK cells also indicate that Cdc42 regulates both endocytic trafficking and

exocytic secretion (Kroschewski et al., 1999). Specifically, expression of dominant negative

Cdc42^{T17N} mislocalizes basolateral membrane proteins to the apical membrane, while

expression of constitutively active Cdc42Q61L depolarizes actin from the apical membrane,

causing actin to instead appear throughout the cell periphery (Kroschewski et al., 1999).

In *S. pombe*, Cdc42 is independently activated by two GEFs, a transient and mostly cytoplasmic

GEF, Gef1, which localizes to cortical puncta at polarized regions (Coll et al., 2003; Das et al.,

2015; Hercyk and Das, 2019b), as well as a strong GEF, Scd1, which localizes throughout

polarized regions and promotes sustained Cdc42 activation via a positive feedback loop (Das et

al., 2012; Das and Verde, 2013; Lamas et al., 2020). Like Cdc42, both of these GEFs are

present at polarized cell ends, but it remains unclear which one arrives first at this site to initiate

Cdc42 activation, although Scd1 has been shown to prevent ectopic localization of Gef1

(Hercyk et al., 2019). Previous research has shown that Gef1 is recruited to cortical patches via

the F-BAR protein Cdc15 (Hercyk and Das, 2019a; Hercyk and Das, 2019b), which promotes

endocytosis and is itself recruited to endocytic actin patches via the endocytic Type I myosin,

Myo1 (Arasada and Pollard, 2011; Carnahan and Gould, 2003). Conversely, Cdc15 does not

recruit Scd1 to cortical patches (Hercyk and Das, 2019b). Additionally, loss of *gef1* leads to

excessive accumulation and prolonged lifetimes of Cdc15 within endocytic patches, which can

be rescued by expression of a constitutively active *cdc42G12V* allele (Adams et al., 1990).

Together, these reports suggest that Cdc15 recruits Gef1 to endocytic patches to promote

Cdc42 activation and that this recruitment is important to regulate endocytic patch dynamics.

Prior research has additionally linked Cdc42 activity to the regulation of endocytosis via its direct

downstream effector, the p21-activated kinase (PAK) (Murray and Johnson, 2001; Ottilie et al.,

1995; Wu et al., 1997), which is activated by binding GTP-bound Cdc42 to relieve its

autoinhibition (Bokoch, 2003). Like Cdc42, PAK is conserved across eukaryotic life from yeast

to humans, functioning as a serine/threonine kinase (Manser et al., 1994; Marcus et al., 1995).

In unicellular eukaryotes, PAK is known to phosphorylate the endocytic Type I myosin in

organisms such as slime molds, amoebas, and yeasts (Brzeska et al., 1997; Lechler et al.,

2001; Lechler et al., 2000; Lee et al., 2000; Murray and Johnson, 2001; Wu et al., 1997),

including *S. pombe* (Attanapola et al., 2009). In these organisms, PAK promotes the ATPase

activity and motor function of the Type I myosin by phosphorylating a conserved serine or

threonine residue located in the motor domain known as the TEDS-site (Bement and Mooseker,

1995; Fujita-Becker et al., 2005; Pedersen et al., 2023). Specifically, TEDS-site phosphorylation

facilitates association of the myosin with actin, stabilizing the complex (Fujita-Becker et al.,

2005). In *S. pombe*, phospho-null mutation of the TEDS-site abrogates proper localization of

Myo1 to endocytic actin patches, instead causing the myosin to localize uniformly along the

plasma membrane (Attanapola et al., 2009). Additionally, we have recently shown that Pak1

promotes successful internalization of endocytic patches and even its own removal via

endocytosis at growing cell ends of *S. pombe* (Harrell et al., 2024). These findings suggest that

regulation of Cdc42 activation may be necessary to facilitate proper endocytic patch dynamics.

While we see distinct sites of endocytosis at the cells ends, division site, and the cell sides, it is

not known whether these sites are functionally similar and if they show different regulatory

patterns. Here, we use genetic approaches and quantitative live cell imaging to better define the

intricate connections between the different sites of polarization and endocytosis. We show that

endocytic actin patches display disparate dynamics at distinct cell sites. Endocytosis at the

polarized sites requires Cdc42, its target Pak1, and the downstream substrate Myo1. At the cell

sides, Myo1 but not Cdc42 and Pak1 regulates patch assembly and dynamics. While our recent

work shows that endocytosis ensures proper spatiotemporal regulation of Cdc42 activation

(Harrell et al., 2024), here we demonstrate that Cdc42 activation likewise spatiotemporally

directs endocytosis.

RESULTS

Endocytic events at distinct regions in the cell show differential dynamics

To our knowledge, the dynamic behaviors of individual endocytic patches within each distinct

region have not been characterized and reported. We were thus curious to determine how

endocytic patches at polarized regions behave compared to those at non-growing regions. To

do this, we performed 1-sec interval timelapse imaging of asynchronous cultures of wild-type

cells expressing mEGFP-tagged fimbrin Fim1, a branched actin crosslinker (Nakano et al.,

2001), which promotes actin bundling necessary for endocytic patch internalization (Skau et al.,

2011). This approach allowed us to simultaneously capture the dynamics of individual endocytic

patches at the cell ends, the cell sides, and the division site (Fig. 1 A). We then used the FIJI

plugin TrackMate (Tinevez et al., 2017) to track the trajectories of individual Fim1-mEGFP

patches. From this data, we found that Fim1-mEGFP-labeled endocytic patches have disparate

lifetimes at each region, with patches at the cell sides exhibiting the shortest lifetimes, while

patches at the division site show the longest lifetimes (Fig. 1 B). While the lifetimes at polarized

cell ends and the division site are statistically different in duration, the mean lifetimes for both

regions (~16 sec at the cell ends and ~18 sec at the division site) fall within the reported

average range for Fim1-mEGFP lifetimes (Berro and Pollard, 2014; Sirotkin et al., 2010). In

contrast, the patch lifetime at the cell sides is ~12 sec (Fig. 1 B), well below than that at the

polarized sites.

Mean patch intensity was also measured frame-by-frame throughout each patch's lifetime. This

analysis revealed that endocytic patches at the cell ends and the division site recruit Fim1-

mEGFP at the same rate and to the same peak intensity (Fig. 1 C). Conversely, endocytic

patches at the cell sides recruited ~50% less Fim1-mEGFP at peak intensity compared to the

polarized sites (Fig. 1 C). Using the R package TrackMateR (Sittewelle and Royle, 2024) , we

calculated the diffusive properties of tracked Fim1-mEGFP patches to characterize their

respective dynamics. Calculations show that endocytic patches at the division site are the

slowest moving (Fig. 1 D) and most stable with the least diffusion of all sites (Fig. 1 F). In

contrast, endocytic patches at the cell sides move the fastest (Fig. 1 D) and exhibit the highest

diffusion rate (Fig. 1 F). Patches at the cell ends display a comparatively intermediate speed

and diffusion rate (Fig. 1 D and F). However, mean squared displacement (MSD) analysis

reveals that endocytic patches at the cell ends and the division site exhibit similar degrees of

directed motion, while patches at the cell sides demonstrate more diffusive behavior overall

(Fig. 1 E). Together, these data indicate that endocytic patches at polarized regions recruit twice

as much Fim1-mEGFP, are longer-lived, and exhibit more stable and directed dynamics

compared to patches at the non-growing cell sides.

Cdc42 activity promotes timely formation of endocytic branched actin networks at the division site

Our findings that endocytic events behave differently at polarized regions compared to non-growing cell sides suggest that polarity regulators may influence endocytic dynamics. Indeed, Cdc42 has been shown to promote uptake of Lucifer Yellow via endocytosis in *S. pombe* (Murray and Johnson, 2001). Connecting previous observations that sites of endocytosis mirror sites of Cdc42 activation, we asked if Cdc42 activation is necessary for endocytosis. To test this idea, we asked if a delay in Cdc42 activation at the division site would consequently delay onset of endocytic activity within the division plane. We selected the division site since the temporal pattern of Cdc42 activation is best understood at this site. The GEF Gef1 is the first GEF to appear and activate Cdc42 at the division site, and loss of Gef1 consequently leads to a delay in Cdc42 activity at this site (Wei et al., 2016) since the other Cdc42 GEF, Scd1, depends on Gef1 for timely localization to the division site (Hercyk et al., 2019). While delayed Cdc42 activation in *gef1Δ* mutant delays actomyosin ring constriction due to delayed recruitment of Bgs1, the timing of mitotic events proceeds normally (Wei et al., 2016). Thus, we measured the timing of Fim1-mEGFP appearance at the division site relative to onset of mitosis marked by spindle pole body (SPB) separation in *gef1Δ* cells compared to *gef1$^+$* controls. These experiments show that Fim1-mEGFP first appears at the division site ~12 mins after SPB separation in *gef1$^+$* cells, while its appearance in *gef1Δ* cells is delayed to ~20 mins post-SPB separation (green dotted box, Fig. 2A and Supplementary Fig.S1A). Intriguingly, Cdc42 is also first activated at the division site ~10-12 min post-SPB separation in wild-type cells (Hercyk and Das, 2019b; Wei et al., 2016). Fim1-mEGFP patches appear uniformly throughout the division site ~20 mins post-SPB separation in *gef1$^+$* controls compared to ~28 mins post-SPB separation in *gef1Δ* mutants (solid green box, Fig. 2 A and B). These findings suggest that Cdc42 activation is indeed required for endocytosis.

We were then curious to determine how long after Cdc42 activation does Fim1 appear at the

division site in *gef1⁺* as well as *gef1Δ* cells. In *gef1⁺* cells, Fim1-mCherry first appears at the

division site (purple dotted box) ~5-6 mins after Cdc42 is activated (green box and arrowheads,

Fig. 2 D and Supplementary Fig.S1B), as visualized by the CRIB-3xGFP bio-probe which binds

active GTP-bound Cdc42 (Tatebe et al., 2008). Comparatively, in *gef1Δ* cells, Fim1-mEGFP first

appears at the division site (purple dotted box) about the same time as Cdc42 activation (green

box and arrowheads, Fig. 2 D and Supplementary Fig.S1B). Similarly, Fim1-mCherry appears

throughout the division site after Cdc42 activation at ~7-8 mins in *gef1⁺* and at ~4 mins in *gef1Δ*

cells (Fig. 2 C and D). This suggests that endocytosis starts soon after Cdc42 activation in

gef1Δ mutants since these cells have already progressed through mitosis and thus many

cellular events are already complete. Our findings suggest that *gef1Δ* cells simply await Cdc42

activation to initiate endocytosis. Together, these data suggest that Cdc42 activation is required

for the onset of endocytosis at the division site.

The GEFs Gef1 and Scd1 differentially regulate Cdc42 to direct branched actin synthesis and dynamics at sites of polarization

While the Cdc42 GEF Scd1 is required for polarized growth at the cell ends, Gef1 is required for bipolar growth (Hercyk et al., 2019). Findings that Gef1 and Scd1 differentially activate Cdc42 at polarized sites prompted us to define the manner of Cdc42 activation required to direct endocytosis. We tracked the dynamics of Fim1-mEGFP at the ends in *gef1Δ* and *scd1Δ* cells compared to *gef1⁺scd1⁺* controls, as previously described (Fig. 1). In these experiments, cells also expressed Cdc15-tdTomato to corroborate and expand upon our previous findings. As shown before (Onwubiko et al., 2019), *gef1Δ* mutants exhibit excessive accumulation and prolonged lifetimes of Cdc15-tdTomato at the cell ends compared to *gef1⁺* control cells (Fig. 3 A; Supplementary Fig. S2 A and B). We additionally find that Fim1-mEGFP accumulation within endocytic patches is likewise increased in *gef1Δ* mutants at both the cell ends and the division site compared to controls. (Fig. 3 A, B, and D). In in *gef1Δ* mutants, Fim1-mEGFP lifetimes are also prolonged at the cell ends; however, we saw no change in patch lifetime at the division site in *gef1Δ* cells (Fig. 3 H). Notably, the recruitment patterns and lifetimes of Fim1-mEGFP patches are the same at the cell sides of *gef1⁺* and *gef1Δ* cells (Fig. 3 F and H), which suggests that the dynamics of these patches are not influenced by Cdc42 or its GEFs.

In contrast, in *scd1Δ* mutants, we did not observe any change in the recruitment dynamics of either Cdc15-tdTomato or Fim1-mEGFP compared to *scd1⁺* cells (Supplementary Fig. S2 C and D; Fig. 3 A, C, E, and G). We did, however, find that Fim1-mEGFP patches are depolarized rather than concentrated at the cell ends in *scd1Δ* mutants (Supplementary Fig. S2 E). Fim1-mEGFP patches in *scd1Δ* mutants also display similar lifetimes at all regions of the cell (Fig. 3 H), which is consistent with the random, depolarized Cdc42 activation patterns that occur in these cells due to loss of Scd1-mediated restriction of ectopic Gef1 localization (Hercyk et al.,

213   2019). Together, these results suggest that Gef1 is the primary GEF that regulates Cdc42

214   activation to direct proper endocytic patch dynamics at polarized regions.

Pak1/Orb2 kinase activity promotes timely formation of endocytic branched actin networks at the division site

Next, we asked if Cdc42 regulates endocytosis via its target the Pak1 kinase. In addition to the cell ends, Pak1 is also present at the division site during cytokinesis where it promotes assembly of the contractile actomyosin ring (Magliozzi et al., 2020). Similar to previously described experiments (Fig. 2), we measured the time intervals between SPB separation and appearance of Fim1-mEGFP at the division site in $orb2^+$ cells compared to $orb2$-34, a kinase-dead, temperature-sensitive allele of *pak1* (Das et al., 2012; Verde et al., 1995). Even at permissive temperature, $orb2$-34 mutants show minimal kinase activity and display polarity defects compared to wild-type *pak1* (Das et al., 2012; Verde et al., 1995). Thus, experiments were performed under permissive 25°C conditions to minimize pleiotropic effects that may arise under temperature restriction, which causes $orb2$-34 cells to become rounded and orb-like (Verde et al., 1995). Under permissive conditions, the $orb2$-34 mutants grow as well as $orb2^+$ cells and only display cell morphology defects. Nevertheless, we do see a ~2 min delay in the appearance of Fim1-mEGFP appearance (dotted and solid green boxes) at the division site in $orb2$-34 mutants compared to controls (Fig. 4 A-C). While this is not a major delay, it is consistent and significant. It is possible that under complete abrogation of Pak1 kinase activity the difference will be enhanced. However, our experimental conditions do not allow us to expose the cells to such extreme conditions for extended timelapse imaging while avoiding pleiotropic effects. Thus, these data indicate that timely initiation of endocytosis at the division site depends both on Cdc42 and Pak1 activation.

Pak1 facilitates proper formation and dynamics of endocytic actin patches at sites of polarization

Given our recent findings that loss of Pak1 kinase function impairs internalization of endocytic patches at polarized cell ends (Harrell et al., 2024), we asked whether similar endocytic defects also occur at the division site and cell sides. To examine this, we tracked Fim1-mEGFP patches in temperature-sensitive *orb2-34* mutants and compared their dynamics to patches tracked in *orb2*+ control cells. As at the cell ends (Harrell et al., 2024), Fim1-mEGP patch internalization at the division site is impaired in *orb2-34* mutants compared to controls, especially at the 35°C restrictive temperature (Fig. 5 A). Overall, it appears that Fim1 patches largely move within the plane of the plasma membrane in the absence of Pak1 kinase function.

We also observe impaired recruitment of Fim1-mEGFP to endocytic patches in *orb2-34* mutants under permissive conditions. At 25°C, *orb2-34* cells show decreased peak Fim1-mEGFP intensity within endocytic patches at the cell ends and the division site compared to controls (Fig. 5 B and D). In contrast, at the 35°C restrictive temperature, peak Fim1-mEGFP intensity at the cell ends in *orb2-34* mutants remains the same as observed at 25°C, while *orb2*+ controls show decreased Fim1-mEGFP patch intensity at the cell ends at this temperature (Fig. 5 C). However, we observe the opposite effect at the division site. Here at 35°C, Fim1-mEGFP patch intensity dramatically decreases at the division site in *orb2-34* mutants while patch intensity is unchanged in *orb2*+ controls (Fig. 5 E). These observations suggest that Fim1 recruitment to the cell ends of *orb2-34* mutants is insensitive to temperature restriction, while Fim1 recruitment to the division site of *orb2-34* mutants is impaired under these conditions. Given that patch internalization is impaired at the cell ends in *orb2-34* mutants (Harrell et al., 2024), our data suggest the level of actin assembly and dynamics alone do not determine the ability of endocytic vesicles to successfully internalize. Consistent with previous experiments showing

that endocytic dynamics at the cell sides are not influenced by changes in polarity regulation, we

did not observe changes in Fim1-mEGFP recruitment at these sites other than some decrease

in peak patch intensity in $orb2^+$ controls at 35°C (Fig. 5 F and G).

Across the board at all cell regions, Fim1-mEGFP patch lifetimes decrease at 35°C compared to

25°C (Fig. 5 H and I), likely due to increased Brownian motion. As previously reported (Harrell et

al., 2024), Fim1-mEGFP patch lifetimes do not change at the cell ends of $orb2-34$ mutants

compared to controls (Fig. 5 H and I). Conversely, at 25°C, Fim1-mEGFP patch lifetimes are ~5

seconds shorter at the division site in $orb2-34$ cells compared to $orb2^+$ cells (Fig. 5 H), but this

difference is abrogated at 35°C where $orb2^+$ patch lifetimes decrease (Fig. 5 I). While loss of

Pak1 kinase function impairs endocytic internalization at both the cell ends and the division site,

our data suggests that Pak1 may perform additional functions at the division site since we

observe changes in Fim1 accumulation and lifetime at this site, but not at the cell ends.

Together, these results indicate that Pak1 kinase plays important and differential roles at

polarized regions to regulate endocytic patch behavior.

The Type I myosin Myo1 promotes timely internalization of endocytic actin patches via Pak1-mediated phosphorylation of its motor domain

Our data show that endocytosis is differentially regulated at distinct sites within polarized cells. While Cdc42 and Pak1 kinase are required for proper actin patch dynamics at the cell ends and the division site, they are not required for the patches at the cell sides. Given that previous reports indicate that the Pak1 kinase regulates endocytosis via Myo1 phosphorylation, we asked if Myo1 itself is differentially required for endocytosis at these distinct sites. We first analyzed Myo1-GFP localization in *S. pombe*. As expected Myo1-GFP localizes to the growing cell ends and the division site (Attanapola et al., 2009; Carnahan and Gould, 2003; Lee et al., 2000). Myo1-GFP puncta also appear along the cell sides, co-localizing with branched actin patches as labeled with Fim1-mCherry (arrowheads, Supplementary Fig. S3A) (Arasada and Pollard, 2011; Sirotkin et al., 2010). Next, we examined Fim1 dynamics in *myo1Δ* cells compared to *myo1$^+$* controls at the cell ends, division site, and the cell sides. In the absence of *myo1*, endocytic patches marked with Fim1-mCherry display weak internalization and largely slide within the plane of the plasma membrane at both the cell ends and the division site (Fig. 6 A). Additionally, fewer endocytic patches form in *myo1Δ* mutants, although Fim1-mCherry lifetimes are much longer compared to controls at all regions (Fig. 6 B). Notably, while Fim1-mCherry recruitment to endocytic patches reaches higher peak levels in *myo1Δ* cells compared to *myo1$^+$* controls, Fim1 takes longer to accumulate at patches in these mutants (Fig. 6 D). Indeed, peak Fim1-mCherry patch intensity is reached ~8 seconds later at the cell ends (Fig. 6 E) and ~11 seconds later at the division site in *myo1Δ* mutants compared to *myo1$^+$* controls due to this slower recruitment of Fim1 (Fig. 6 F). At the cell sides, peak Fim1-mCherry patch intensity is slightly increased in *myo1Δ* cells and patches also take longer to reach peak intensity (Fig. 6 G). Since Myo1 can localize to cell sides in addition to sites of polarization, this may explain the changes in Fim1 behavior at the cell sides in *myo1Δ* mutants.

While Pak1 phosphorylates Myo1 at the S361 TEDS site located within its N-terminal motor

domain (Attanapola et al., 2009), Myo1 also binds and weakly activates the branched actin

nucleating Arp2/3 complex via its C-terminal tail (Lee et al., 2000; Sirotkin et al., 2005). Thus,

Myo1 may promote endocytic patch internalization via its N-terminal motor function and/or its C-

terminal association with Arp2/3. As *myo1Δ* completely abolishes all Myo1 function,

experiments in *myo1Δ* mutants alone cannot distinguish between these mechanistic

possibilities. To determine if Pak1-mediated Myo1 phosphorylation is necessary to promote

endocytic patch internalization, we examined endocytic patch behavior in the single point TEDS

site phospho-null mutant, *myo1-S361A*, which cannot be phosphorylated by Pak1 but retains its

C-terminal Arp2/3 binding domain (Attanapola et al., 2009). In the absence of Pak1-mediated

phosphorylation of Myo1, we hypothesized that Fim1-mCherry patches would show similar

weak internalization as observed in *myo1Δ* mutants.

In the TEDS site *myo1-S361A* mutant, we find that Fim1-mCherry is adequately recruited to

endocytic patches, yet the patches do not properly internalize (Fig. 6 A and D, Fig. S2 A and B).

Additionally, we observe that Myo1-S361A-GFP intensity remains mostly stable at the plasma

membrane of polarized cell ends, while in *myo1$^+$* cells, Myo1-GFP intensity increases as Fim1-

mCherry is recruited to endocytic patches and is lost as Fim1-mCherry patches internalize at

the cell ends (Fig. S2), as reported previously (Sirotkin et al., 2005). We also observe increased

cytoplasmic intensity of Myo1-S361A-GFP compared to controls (Fig. S2 A and B). In *myo1-*

S361A mutants, Fim1-mCherry lifetimes are indistinguishable from *myo1$^+$* controls at the cell

ends and sides, but patch lifetimes are prolonged at the division site (Fig. 6 C). Similar to

myo1Δ mutants, the timing of Fim1 recruitment to endocytic patches is prolonged in *myo1-*

S361A mutants (Fig. 6 D and H; Fig. S2). At the division site, Fim1 recruitment is enhanced in

myo1-S361A mutants and peak Fim1-mCherry intensity is reached slightly later compared to

myo1⁺ controls (Fig. 6 I), mimicking *myo1Δ* mutants (Fig. 6 F). Similar to *myo1Δ* mutants, peak

Fim1-mCherry intensity at the cell ends is reached later in *myo1-S361A* mutants compared to

controls (Fig. 6 J). These observations suggest that branched actin formation can occur in the

absence of Myo1 function, even though Myo1 is a nucleation promoting factor, albeit weak

compared to Wsp1 (Sirotkin et al., 2005). We also find that while Myo1 regulates patch lifetimes

at the cell sides, the TEDS site phosphorylation is not required for the patches at these sites.

This aligns with the fact that the Pak1 kinase that phosphorylates the TEDS is also not required

for patch dynamics at the cell sides.

Cdc42 and Pak1 activity regulate the recruitment and dynamics of Myo1 at polarized sites

Our data show that Cdc42 and Pak1 are required for proper endocytic dynamics at polarized

sites. We asked if Cdc42 and Pak1 regulate endocytosis via the regulation of Myo1 recruitment

and dynamics at these sites. To test this, we first observed whether loss or delay of Cdc42 or

Pak1 activation at the division site delays Myo1 recruitment to this site. Using *gef1Δ* mutants to

delay Cdc42 activation at the division site as previously described (Fig. 2), we observe that

Myo1-GFP recruitment to the division site is consequently delayed by ~10 mins compared to

gef1$^+$ control cells (Fig. 7 A and B). Similarly, in *orb2-34* mutants lacking Pak1 kinase activity,

Myo1-GFP recruitment to the division plane is delayed by ~4 mins compared to *orb2*$^+$ controls,

even under permissive conditions (Fig. 7 C and D). These findings indicate that both Cdc42 and

Pak1 activation promote timely recruitment of Myo1 to the division site.

Following our findings at the division site, we then asked if Cdc42 and Pak1 activation similarly

regulate Myo1 recruitment to polarized cell ends. To test this, we first measured the mean

intensity of Myo1-GFP recruited to the cell ends in genotypes with varying degrees of Cdc42

activity. While loss of *gef1* decreases Cdc42 activity (Wei et al., 2016), loss of the GAPs *rga4*

and *rga6* increases Cdc42 activation levels (Campbell et al., 2022). Thus, we measured mean

Myo1-GFP intensity at the cell ends of *gef1Δ* and *rga4Δrga6Δ* mutants compared to control

cells. Intriguingly, Myo1-GFP recruitment is decreased in *gef1Δ* mutants and conversely

enhanced in *rga4Δrga6Δ* mutants compared to controls (Fig. 8 A and B). To investigate the

dynamics of Myo1 at the cell ends, we also performed 1 sec-interval time-lapse imaging and

measured the dwell time of Myo1-GFP at the plasma membrane. These analyses reveal that

Myo1-GFP dwell time is increased in both *gef1Δ* and *rga4Δrga6Δ* mutants compared to

controls, suggesting that the myosin is less dynamic in both cases (Fig. 8 C and D).

We then performed similar experiments in *orb2-34* cells to see if decreased Pak1 activation

impairs Myo1-GFP recruitment and dynamics at cell ends. Under permissive conditions at 25C°,

mean Myo1-GFP intensity is indistinguishable between *orb2-34* and *orb2*$^+$ cell ends (Fig. 8 E

and F). In contrast, temperature restriction at 35C° reduces mean Myo1-GFP intensity at the cell

ends in both *orb2*$^+$ and *orb2-34* cells, although Myo1-GFP recruitment is more severely

impacted in *orb2-34* mutants (Fig. 8 E and F). Finally, loss of Pak1 kinase function likewise

impairs Myo1-GFP dynamics at cell ends, as Myo1-GFP dwell times at the plasma membrane

are prolonged in *orb2-34* compared to *orb2*$^+$ cells under permissive conditions (Fig. 8 G and H).

Together, these data indicate that Cdc42 and Pak1 activation promote the recruitment of Myo1

to the plasma membrane in a dose-dependent manner. Additionally, these results suggest that

an intermediate range of Cdc42 activation facilitates proper Myo1 dynamics. In summary, our

findings suggest that sufficient recruitment of Myo1 to polarized sites depends on both Cdc42

and Pak1 activation.

DISCUSSION

In many cell-walled organisms, endocytosis is an actin-dependent process since cells must overcome high internal turgor pressure to properly bend and invaginate the plasma membrane (Basu et al., 2014; Lacy et al., 2018). In yeast, branched actin polymerization and crosslinking provide the force needed to promote successful patch internalization against high turgor pressure (Basu et al., 2014; Kaksonen and Roux, 2018; Skau et al., 2011). However, it is not known whether the force requirements are the same at distinct sites of endocytosis. For instance, at the division site, patches form in close proximity to the actomyosin ring, the synthesizing septum, as well as other membrane-associated cytoskeletal structures such as the septin ring (Wu et al., 2003), which may impact the force required for successful internalization. Here, we demonstrate that endocytic patch behavior is intricately linked with cell polarization. We find that endocytic patches at non-growing cell sides weakly recruit Fim1, display decreased lifetimes, and exhibit diffusive behavior. In contrast, endocytic patches within polarized regions recruit twice as much Fim1 with longer lifetimes, and exhibit directed motion during patch internalization. While it is unclear if the endocytic patches at the cell sides successfully undergo vesicle internalization. our data show that only the patches at the polarized sites exhibit the controlled and directed motion required for orderly membrane invagination and vesicle scission.

Cdc42 and Pak1 activity promote initiation of endocytosis at sites of polarization

We show that Cdc42 activation is required for the initiation of branched actin-mediated endocytosis at the division site. Similar to Cdc42 activation patterns, Fim1-mEGFP first appears at the division site of *gef1*⁺ controls during ring maturation well before the onset of ring constriction (Wang et al., 2016). However, in *gef1Δ* mutants, Fim1-mEGFP appearance is delayed to just before the onset of ring constriction, as reported for Cdc42 activation (Wei et al., 2016). Yet, once Cdc42 is finally activated in *gef1Δ* mutants, Fim1-mEGFP rapidly appears at

the division site. These results suggest that the timing of Cdc42 activation and onset of

endocytosis are tightly linked. In a similar manner, we also demonstrate that the Cdc42

downstream effector Pak1 promotes timely onset of endocytosis at the division site. Given the

previously described challenges, we have not directly shown that Cdc42 activity is essential for

endocytosis at the cell ends. However, we demonstrate that decreased Cdc42 activation via

loss of its GEF *gef1* results in excessive Fim1 recruitment yet delayed patch internalization at

the cell ends. Thus, Gef1-mediated Cdc42 activation facilitates proper patch formation and

internalization at the cell ends.

The two Cdc42 GEFs differentially regulate endocytosis

As we probed how Cdc42 activation promotes endocytosis, we found that various polarity

mutants exhibit distinct phenotypes of endocytic patch behavior (summarized in Supplementary

Table S1, S2, and S3). Individual loss of each Cdc42 GEF results in disparate phenotypes. Loss

of *gef1* —where *scd1* is the sole Cdc42 GEF— severely impacts endocytic patch behavior at

polarized sites. On the other hand, loss of *scd1* —where *gef1* is the sole GEF— depolarizes the

location of endocytic sites across the cell cortex but the patch dynamics remain mostly normal.

These findings suggest that rather than strong, sustained activation via Scd1, transient, Gef1-

mediated Cdc42 activation is required for proper endocytic patch formation and internalization.

While dispensable for proper patch behavior, Scd1 is still required to prevent ectopic Cdc42

activation and thus regulates where endocytosis occurs. We observe that loss of *gef1* results in

some differences between patch dynamics at the cell ends and the division site. In *gef1$^+$* cells,

peak Fim1-mEGFP patch intensity is reached ~2 seconds faster at the division site compared to

the cell ends. In contrast, *gef1Δ* mutants appear to recruit Fim1 to endocytic patches at both

sites with a delay. Additionally, Fim1-mEGFP patch lifetimes are increased at the cell ends of

gef1Δ mutants. Comparatively, patch lifetimes are not enhanced at the division site of *gef1Δ*

mutants, even though *gef1Δ* mutants recruit more Fim1 to endocytic patches. These

observations indicate that the rate of branched actin synthesis is likely faster in *gef1Δ* mutants,

since more branched actin is formed at the patch within the same span of time. As Cdc42 is

under strong Scd1-mediated activation in *gef1Δ* mutants, it is possible that sustained Cdc42

activity in these mutants leads to enhanced branched actin synthesis. Our results further show

that this excessive branched actin formation is actually deleterious to timely patch internalization

in *gef1Δ* mutants. Together, these findings indicate that Cdc42 activation must be tightly

regulated at the endocytic patch to form proper amounts of branched actin.

Pak1 kinase differentially regulates endocytic patch dynamics at distinct sites

Although loss of *pak1* kinase function impairs endocytic patch internalization at the cell ends

(Harrell et al., 2024), it seems that actin patch formation is most greatly impaired at the division

site of *pak1* mutants. At restrictive temperature, Fim1 recruitment to endocytic patches at the

cell ends is not impaired in *pak1* mutants, while it is greatly decreased at the division site. This

indicates that patch assembly at the cell ends is more robust to loss of *pak1* function than the

division site. However, as reported earlier, these patches do not internalize efficiently and often

fail endocytosis (Harrell et al., 2024). This suggests that Pak1 could play additional roles in

endocytosis apart from patch assembly. Alternatively, while the patch synthesizes sufficient

actin filaments, it may not be properly organized for successful internalization. We also see

enhanced patch assembly defects at the division site in *pak1* mutants. This suggests that Pak1

may play a more important role in at the division site that affects endocytosis. Already some

such roles are known since Pak1 phosphorylates several cytokinetic ring proteins (Magliozzi et

al., 2020) .

Cdc42 and Pak1 kinase regulate Type I myosin recruitment and dynamics at sites of

endocytosis

We show that Cdc42 promotes Myo1 recruitment to polarized sites in a dose-dependent

manner. Our data indicate that both insufficient and excessive Cdc42 activation leads to greater

stability of Myo1 at the plasma membrane, as demonstrated by its increased dwell time in

mutants of Cdc42 GEFs and GAPs and as well as the *pak1* kinase. This increased stability is

reminiscent of the stable localization displayed by Myo1-S361A-GFP, which cannot be

phosphorylated by Pak1 kinase at its motor domain TEDS site. Since endocytic patch

internalization defects seem to coincide with increased Myo1 stability, these results suggest that

Myo1 requires a certain level of dynamic behavior to properly promote endocytic patch

internalization. Since Myo1 is a direct target of Pak1 (Attanapola et al., 2009), we initially

expected *myo1-S361A* mutants to phenocopy the endocytic defects observed in kinase-dead

pak1 mutants. Our results, however, show distinct phenotypes of altered endocytic patch

behavior for these mutants, which indicates that Pak1 has additional roles that are yet to be

explored.

In line with previous reports, our findings suggest that the Arp2/3 complex can generate

branched actin without Pak1 phosphorylation of Myo1's TEDS site (Attanapola et al., 2009) and

even without Myo1 altogether. Thus, the internalization defect observed in *myo1-S361A*

mutants does not stem from an inability to create branched actin. Given that TEDS site

phosphorylation promotes motor function of the Type I myosin in other organisms (Bement and

Mooseker, 1995; Fujita-Becker et al., 2005; Pedersen et al., 2023), we suspect that the same

likely occurs in *S. pombe*. Indeed, the perturbed localization and increased stability we observe

for Myo1-S361A-GFP suggests that its motor function may be impaired. Such findings have

been reported in budding yeast where Type I myosin motor activity is independent of its ability

to nucleate actin (Manenschijn et al., 2019). As the other actin nucleation promoting factor, the

WASP homolog Wsp1, activates Arp2/3 independently and more strongly than Myo1 in *S.*

pombe (Sirotkin et al., 2005), we suspect that Wsp1 primarily promotes Arp2/3-dependent

branched actin formation in *myo1-S361A* mutants.

Independent of Cdc42 and Pak1, Myo1 promotes patch assembly and dynamics at the cell

sides. In agreement, Pak1-dependent TEDS site phosphorylation of Myo1 does not appear to

be required at the cell sides. The patches at the cell sides contain less actin, have decreased

lifetimes, and do not rapidly internalization into the cell interior. It is possible that while Myo1

enables actin patch assembly at the cell sides, it may not generate sufficient branched actin

required for patch internalization. Sufficient branched actin assembly likely requires Pak1-

dependent phosphorylation of Myo1. This would explain why the patches at the polarized sites

show stronger internalization.

Endocytic patch assembly and dynamics are differentially regulated at distinct cell sites

While both the growing cell ends and the division site are sites of polarization, we observe

subtle differences in endocytic patch behavior between these sites. As endocytosis is a force-

sensitive and actin-dependent process, the differences in patch dynamics at these distinct sites

suggest that the forces experienced by patches at the division site are higher than that of the

cell ends. If such is the case, patches at the division site would require stronger actin

architecture and/or additional assistance from endocytic proteins such as Myo1 to overcome

this increased force. This disparate force requirement could arise in several ways. Perhaps the

flow of polarized secretion produces some counteractive force that must be overcome for

patches to internalize. At the division site, polarized secretion occurs throughout the division

plane, while endocytosis is largely confined to the outer rim of the membrane furrow (Wang et

al., 2016). Furthermore, our observations that endocytic patches at the cell sides are weakly

formed suggest that the forces they experience are also different than those at polarized

regions. It is unclear if the patches at the cell sides internalize, but do not move as far from the

plasma membrane since the force produced within the patch is less. It is possible that these

patches do not require access to active Cdc42 to internalize. Alternatively, it is possible that

they are random assemblies of branched actin that do not represent true sites of endocytosis.

Finally, it seems from our observations in *gef1Δ* mutants that formation of branched actin

patches is not sufficient in itself to stimulate timely patch internalization, since patches

excessively recruit both Cdc15 and Fim1 yet do not timely internalize. Perhaps the actin

architecture within these mutants is improperly organized to internalize the patch. This

hypothesis fits with our observations in *myo1Δ* and *myo1-S361A* mutants, which also assemble

branched actin patches with excessive Fim1 yet exhibit severe patch internalization defects.

Together, our findings indicate a direct role for Cdc42 in regulating endocytic patch formation via

the Pak1 kinase and endocytic patch internalization via the Pak1 target, the Type I myosin, in a

site-specific manner. A recent report characterized endocytic patch dynamics in different fungal

species including *S. pombe* (Picco et al., 2024). Compared to the other species, endocytic

patches in *S. pombe* appear to rapidly internalize at very high speeds deep into the cell interior

(Picco et al., 2024; Sun et al., 2019), which suggests the patches may experience higher

amounts of force which propel them inward at high velocity. Further investigation will determine

if Cdc42, Pak1 and Myo1 differentially regulate endocytosis in different fungal species resulting

in distinct dynamics.

MATERIALS AND METHODS

Strains and cell culture

The *S. pombe* strains used in this study are listed in Table 1. All strains are isogenic to the

original strain PN567. Cells were cultured in yeast extract (YE) medium and grown exponentially

at 25°C for 3 rounds of 8 generations before imaging. Standard techniques were used for

genetic manipulation and analysis (Moreno et al., 1991).

Microscopy

Imaging was performed at room temperature (23–25°C) for all experiments except for restrictive

temperature experiments, which were performed at 35°C (Fig. 5). Images were acquired on a

spinning disk confocal system equipped with a Nikon Eclipse inverted microscope with a

100×/1.49 NA lens, a CSU-22 spinning disk system (Yokogawa Electric Corporation), and a

Photometrics EM-CCD camera (Photometrics Technology Evolve with excelon Serial No:

A13B107000). On this system, images were acquired with MetaMorph (Molecular Devices,

Sunnyvale, CA). Imaging was also performed on a 3i spinning disk confocal system using a

Zeiss AxioObserver microscope equipped with a 100×/1.49 NA objective, an integrated

Yokogawa spinning disk (Yokogawa CSU-X1 A1 spinning disk scanner), and a Teledyne

Photometrics Prime 95b back-illuminated sCMOS camera (Serial No: A20D203014). On this

system, images were acquired using SlideBook (3i Intelligent Imaging innovations). All images

were analyzed with FIJI (National Institutes of Health, Bethesda, MD (Schneider et al., 2012)).

For still imaging, the cells were mounted directly on glass slides with a #1.5 coverslip (Thermo

Fisher Scientific, Waltham, MA) and imaged immediately, and with fresh slides prepared every

10 min. *Z*-series images were acquired with a depth interval of 0.4 μm. For time-lapse images,

cells were placed in 5 mm glass-bottom culture dishes (MatTek, Ashland, MA) and overlaid with

YE medium containing 0.6% agarose and 100 µM ascorbic acid as an antioxidant to minimize

toxicity to the cell, as reported previously (Frigault et al., 2009; Wei et al., 2017).

Analysis of vesicle tracking

Wild-type and mutant cells expressing Fim1-mEGFP or Fim1-mCherry were grown to OD 0.2-

0.5 and placed in glass-bottom culture dishes as previously described for time-lapse imaging.

Cells were imaged in a single medial plane with a frame rate of 1 sec for ~3-5 minutes using the

same laser power and exposure settings for all experiments. To measure patch lifetimes and

mean fluorescence intensity of patches throughout their lifetimes, Fim1-mEGFP/Fim1-mCherry

patches were tracked using the FIJI plugin TrackMate (Tinevez et al., 2017). Background

fluorescence was subtracted from the imaging field away from cells to sharpen the signal of

patches for ease of tracking. Lifetime was measured as the time from which a patch first

displayed distinguishable fluorescence at the cell cortex to when the fluorescence was no longer

detectable. Mean speed (μm/sec), MSD (mean squared displacement), and diffusion

coefficients (μm^2/sec) were all calculated using TrackMateR, an R package that can analyze

data captured and saved by TrackMate as XML files ((Sittewelle and Royle, 2024);

https://quantixed.org/2022/09/05/tracking-announcing-new-r-package-trackmater/). To ensure

that patches were tracked from the beginning of patch formation through patch disassembly,

patches selected for tracking 1. gradually increased in intensity until a peak intensity was

reached and 2. decreased in intensity following peak intensity (Berro and Pollard, 2014; Wang

et al., 2016). Patches tracked with TrackMate were also manually reviewed to ensure tracking

accuracy.

Disruption of Pak1 kinase function via temperature-sensitive allele *orb2-34*

Pak1 kinase was inactivated via the *orb2-34* (*pak1-ts*) temperature-sensitive mutation. Cells

were grown for 2 days in YE medium. On the day of the experiment, cells were cultured to an

OD of 0.2. A subset of these cells was incubated at the permissive temperature of 25°C, while

the other set was incubated at the restrictive temperature of 35°C for 4 h. Cells were imaged

after 4 h under temperature-controlled conditions.

Analysis of Myo1-GFP mean fluorescence intensity at cell ends

Wild-type and mutant cells expressing Myo1-GFP were grown to OD 0.2-0.5 and imaged on

slides. Still Z-images with 21-24 z-planes were collected at a z-interval of 0.4 μm for the 488 nm

channel. The same number of z-slices were collected for wild-type and mutant cells for each

experiment using the same imaging settings. FIJI was used to generate sum projections from

the z-series and the polygon selection tool was used to encircle the "cap" of Myo1-GFP signal

present at individual cell ends. Mean fluorescence intensity at individual cell ends was then

measured within these manually selected Myo1-GFP "caps". The background fluorescence in a

cell-free region of the image was subtracted to normalize intensity measurements. Only cells

found within the central 80% of the imaging field were used for analysis. This was done to

account for uneven illumination of the imaging field.

Analysis of Myo1-GFP dwell time at cell ends

Wild-type and mutant cells expressing Myo1-GFP were grown to OD 0.2-0.5 and placed in

glass-bottom dishes as previously described for time-lapse imaging. Cells were imaged in a

single medial plane with a frame rate of 1 sec for 3 minutes using the same imaging settings for

all experiments. To visualize Myo1-GFP dwell time throughout the 3-minute time-lapse,

kymographs were generated from 1 pixel-wide selection lines drawn across the midline of the

cell end, parallel to the cell's long axis. These kymographs were then manually analyzed to

measure the dwell time of Myo1-GFP at individual cell ends in the following manner: Using

FIJI's rectangle selection tool, individual Myo1-GFP dwell time events were marked on each

kymograph and the length of each event was measured in pixels. Since our imaging settings

generated kymographs were 1 pixel = 1 sec, the number of pixels measured for an individual

event equals Myo1-GFP dwell time in seconds for that event.

Statistical tests

GraphPad Prism was used to determine significance. One-way ANOVA, followed by a Tukey's

multiple comparisons post-hoc test, was used to determine individual p-values when comparing

three or more samples. When comparing two samples, an unpaired Student's t-test (two-tailed,

unequal variance) was used to determine significance.

ACKNOWLEDGMENTS

We thank Drs. Vladimir Sirotkin and Daniel Mulvihill for strains; and Bret Judson for the Boston

College imaging core facility.

COMPETING INTERESTS

No competing interests declared.

FUNDING

This research was supported by the National Science Foundation CAREER award #2309328 to

MD.

AUTHOR CONTRIBUTIONS

M.E.D. and B.F.C. conceived the project. M.E.D. obtained funding for the project. B.F.C.,

A.R.W., and U.P. performed experiments and analyzed data. B.F.C. and M.E.D. wrote the

manuscript.

DATA AND RESOURCE AVAILABILITY

Data and resource availability: All relevant data and resource can be found within the article and

its supplementary information.

REFERENCES

Adams, A. E., Johnson, D. I., Longnecker, R. M., Sloat, B. F. and Pringle, J. R.

(1990). CDC42 and CDC43, two additional genes involved in budding and the establishment of

cell polarity in the yeast Saccharomyces cerevisiae. *J Cell Biol* **111**, 131-42.

Aghamohammadzadeh, S. and Ayscough, K. R. (2009). Differential requirements for

actin during yeast and mammalian endocytosis. *Nat Cell Biol* **11**, 1039-42.

Arasada, R. and Pollard, T. D. (2011). Distinct roles for F-BAR proteins Cdc15p and

Bzz1p in actin polymerization at sites of endocytosis in fission yeast. *Curr Biol* **21**, 1450-9.

Attanapola, S. L., Alexander, C. J. and Mulvihill, D. P. (2009). Ste20-kinase-

dependent TEDS-site phosphorylation modulates the dynamic localisation and endocytic

function of the fission yeast class I myosin, Myo1. *J Cell Sci* **122**, 3856-61.

Basu, R., Munteanu, E. L. and Chang, F. (2014). Role of turgor pressure in

endocytosis in fission yeast. *Mol Biol Cell* **25**, 679-87.

Bement, W. M. and Mooseker, M. S. (1995). TEDS rule: a molecular rationale for

differential regulation of myosins by phosphorylation of the heavy chain head. *Cell Motil*

Cytoskeleton **31**, 87-92.

Berro, J. and Pollard, T. D. (2014). Local and global analysis of endocytic patch

dynamics in fission yeast using a new "temporal superresolution" realignment method. *Mol Biol*

Cell **25**, 3501-14.

Bokoch, G. M. (2003). Biology of the p21-activated kinases. *Annu Rev Biochem* **72**,

743-81.

Brzeska, H., Knaus, U. G., Wang, Z. Y., Bokoch, G. M. and Korn, E. D. (1997). p21-

activated kinase has substrate specificity similar to Acanthamoeba myosin I heavy chain kinase

and activates Acanthamoeba myosin I. *Proc Natl Acad Sci U S A* **94**, 1092-5.

Campbell, B. F., Hercyk, B. S., Williams, A. R., San Miguel, E., Young, H. G. and

Das, M. E. (2022). Cdc42 GTPase activating proteins Rga4 and Rga6 coordinate septum

synthesis and membrane trafficking at the division plane during cytokinesis. *Traffic* **23**, 478-495.

Carnahan, R. H. and Gould, K. L. (2003). The PCH family protein, Cdc15p, recruits two

F-actin nucleation pathways to coordinate cytokinetic actin ring formation in

Schizosaccharomyces pombe. *J Cell Biol* **162**, 851-62.

Coll, P. M., Trillo, Y., Ametzazurra, A. and Perez, P. (2003). Gef1p, a new guanine

nucleotide exchange factor for Cdc42p, regulates polarity in Schizosaccharomyces pombe. *Mol*

Biol Cell **14**, 313-23.

Das, M., Drake, T., Wiley, D. J., Buchwald, P., Vavylonis, D. and Verde, F. (2012).

Oscillatory dynamics of Cdc42 GTPase in the control of polarized growth. *Science* **337**, 239-43.

Das, M., Nuñez, I., Rodriguez, M., Wiley, D. J., Rodriguez, J., Sarkeshik, A., Yates,

J. R., 3rd, Buchwald, P. and Verde, F. (2015). Phosphorylation-dependent inhibition of Cdc42

GEF Gef1 by 14-3-3 protein Rad24 spatially regulates Cdc42 GTPase activity and oscillatory

dynamics during cell morphogenesis. *Mol Biol Cell* **26**, 3520-34.

Das, M. and Verde, F. (2013). Role of Cdc42 dynamics in the control of fission yeast

cell polarization. *Biochem Soc Trans* **41**, 1745-9.

Frigault, M. M., Lacoste, J., Swift, J. L. and Brown, C. M. (2009). Live-cell microscopy

- tips and tools. *J Cell Sci* **122**, 753-67.

Fujita-Becker, S., Dürrwang, U., Erent, M., Clark, R. J., Geeves, M. A. and Manstein,

D. J. (2005). Changes in Mg2+ ion concentration and heavy chain phosphorylation regulate the

motor activity of a class I myosin. *J Biol Chem* **280**, 6064-71.

Gachet, Y. and Hyams, J. S. (2005). Endocytosis in fission yeast is spatially associated

with the actin cytoskeleton during polarised cell growth and cytokinesis. *J Cell Sci* **118**, 4231-42.

Harrell, M. A., Liu, Z., Campbell, B. F., Chinsen, O., Hong, T. and Das, M. (2024).

Arp2/3-dependent endocytosis ensures Cdc42 oscillations by removing Pak1-mediated negative

feedback. *J Cell Biol* **223**.

Hercyk, B. and Das, M. (2019a). Rho Family GTPases in Fission Yeast Cytokinesis.

Commun Integr Biol **12**, 171-180.

Hercyk, B. S. and Das, M. E. (2019b). F-BAR Cdc15 Promotes Cdc42 Activation During

Cytokinesis and Cell Polarization in Schizosaccharomyces pombe. *Genetics* **213**, 1341-1356.

Hercyk, B. S., Rich-Robinson, J., Mitoubsi, A. S., Harrell, M. A. and Das, M. E.

(2019). A novel interplay between GEFs orchestrates Cdc42 activity during cell polarity and

cytokinesis in fission yeast. *J Cell Sci* **132**.

Kaksonen, M. and Roux, A. (2018). Mechanisms of clathrin-mediated endocytosis. *Nat*

Rev Mol Cell Biol **19**, 313-326.

Kroschewski, R., Hall, A. and Mellman, I. (1999). Cdc42 controls secretory and

endocytic transport to the basolateral plasma membrane of MDCK cells. *Nat Cell Biol* **1**, 8-13.

Lacy, M. M., Ma, R., Ravindra, N. G. and Berro, J. (2018). Molecular mechanisms of

force production in clathrin-mediated endocytosis. *FEBS Lett* **592**, 3586-3605.

Lamas, I., Merlini, L., Vještica, A., Vincenzetti, V. and Martin, S. G. (2020).

Optogenetics reveals Cdc42 local activation by scaffold-mediated positive feedback and Ras

GTPase. *PLoS Biol* **18**, e3000600.

Lechler, T., Jonsdottir, G. A., Klee, S. K., Pellman, D. and Li, R. (2001). A two-tiered

mechanism by which Cdc42 controls the localization and activation of an Arp2/3-activating

motor complex in yeast. *J Cell Biol* **155**, 261-70.

Lechler, T., Shevchenko, A. and Li, R. (2000). Direct involvement of yeast type I

myosins in Cdc42-dependent actin polymerization. *J Cell Biol* **148**, 363-73.

Lee, W. L., Bezanilla, M. and Pollard, T. D. (2000). Fission yeast myosin-I, Myo1p,

stimulates actin assembly by Arp2/3 complex and shares functions with WASp. *J Cell Biol* **151**,

789-800.

Magliozzi, J. O., Sears, J., Cressey, L., Brady, M., Opalko, H. E., Kettenbach, A. N.

and Moseley, J. B. (2020). Fission yeast Pak1 phosphorylates anillin-like Mid1 for spatial

control of cytokinesis. *J Cell Biol* **219**.

Manenschijn, H. E., Picco, A., Mund, M., Rivier-Cordey, A. S., Ries, J. and

Kaksonen, M. (2019). Type-I myosins promote actin polymerization to drive membrane bending

in endocytosis. *Elife* **8**.

Manser, E., Leung, T., Salihuddin, H., Zhao, Z. S. and Lim, L. (1994). A brain

serine/threonine protein kinase activated by Cdc42 and Rac1. *Nature* **367**, 40-6.

Marcus, S., Polverino, A., Chang, E., Robbins, D., Cobb, M. H. and Wigler, M. H.

(1995). Shk1, a homolog of the Saccharomyces cerevisiae Ste20 and mammalian p65PAK

protein kinases, is a component of a Ras/Cdc42 signaling module in the fission yeast

Schizosaccharomyces pombe. *Proc Natl Acad Sci U S A* **92**, 6180-4.

Moreno, S., Klar, A. and Nurse, P. (1991). Molecular genetic analysis of fission yeast

Schizosaccharomyces pombe. *Methods Enzymol* **194**, 795-823.

Murray, J. M. and Johnson, D. I. (2001). The Cdc42p GTPase and its regulators Nrf1p

and Scd1p are involved in endocytic trafficking in the fission yeast Schizosaccharomyces

pombe. *J Biol Chem* **276**, 3004-9.

Nakano, K., Satoh, K., Morimatsu, A., Ohnuma, M. and Mabuchi, I. (2001).

Interactions among a fimbrin, a capping protein, and an actin-depolymerizing factor in

organization of the fission yeast actin cytoskeleton. *Mol Biol Cell* **12**, 3515-26.

Nickaeen, M., Berro, J., Pollard, T. D. and Slepchenko, B. M. (2022). A model of

actin-driven endocytosis explains differences of endocytic motility in budding and fission yeast.

Mol Biol Cell **33**, ar16.

718  **Onwubiko, U. N., Mlynarczyk, P. J., Wei, B., Habiyaremye, J., Clack, A., Abel, S. M.**

719 **and Das, M. E.** (2019). A Cdc42 GEF, Gef1, through endocytosis organizes F-BAR Cdc15

720 along the actomyosin ring and promotes concentric furrowing. *J Cell Sci* **132**.

721  **Ottilie, S., Miller, P. J., Johnson, D. I., Creasy, C. L., Sells, M. A., Bagrodia, S.,**

722 **Forsburg, S. L. and Chernoff, J.** (1995). Fission yeast pak1+ encodes a protein kinase that

723 interacts with Cdc42p and is involved in the control of cell polarity and mating. *Embo j* **14**, 5908-

724 19.

725  **Pedersen, R. T. A., Snoberger, A., Pyrpassopoulos, S., Safer, D., Drubin, D. G. and**

726 **Ostap, E. M.** (2023). Endocytic myosin-1 is a force-insensitive, power-generating motor. *J Cell*

727 *Biol* **222**.

728  **Picco, A., Toret, C. P., Rivier-Cordey, A.-S. and Kaksonen, M.** (2024). An

729 evolutionary cell biology perspective into the diverging mechanisms of clathrin-mediated

730 endocytosis in dikarya fungi. *bioRxiv*, 2024.03.28.587219.

731  **Schneider, C. A., Rasband, W. S. and Eliceiri, K. W.** (2012). NIH Image to ImageJ: 25

732 years of image analysis. *Nat Methods* **9**, 671-5.

733  **Sirotkin, V., Beltzner, C. C., Marchand, J. B. and Pollard, T. D.** (2005). Interactions of

734 WASp, myosin-I, and verprolin with Arp2/3 complex during actin patch assembly in fission

735 yeast. *J Cell Biol* **170**, 637-48.

736  **Sirotkin, V., Berro, J., Macmillan, K., Zhao, L. and Pollard, T. D.** (2010). Quantitative

737 analysis of the mechanism of endocytic actin patch assembly and disassembly in fission yeast.

738 *Mol Biol Cell* **21**, 2894-904.

739  **Sittewelle, M. and Royle, S. J.** (2024). Passive diffusion accounts for the majority of

740 intracellular nanovesicle transport. *Life Sci Alliance* **7**.

741  **Skau, C. T., Courson, D. S., Bestul, A. J., Winkelman, J. D., Rock, R. S., Sirotkin, V.**

742 **and Kovar, D. R.** (2011). Actin filament bundling by fimbrin is important for endocytosis,

743 cytokinesis, and polarization in fission yeast. *J Biol Chem* **286**, 26964-77.

Sun, Y., Schoneberg, J., Chen, X., Jiang, T., Kaplan, C., Xu, K., Pollard, T. D. and

Drubin, D. G. (2019). Direct comparison of clathrin-mediated endocytosis in budding and fission

yeast reveals conserved and evolvable features. *Elife* **8**.

Tatebe, H., Nakano, K., Maximo, R. and Shiozaki, K. (2008). Pom1 DYRK regulates

localization of the Rga4 GAP to ensure bipolar activation of Cdc42 in fission yeast. *Curr Biol* **18**,

322-30.

Tinevez, J. Y., Perry, N., Schindelin, J., Hoopes, G. M., Reynolds, G. D., Laplantine,

E., Bednarek, S. Y., Shorte, S. L. and Eliceiri, K. W. (2017). TrackMate: An open and

extensible platform for single-particle tracking. *Methods* **115**, 80-90.

Verde, F., Mata, J. and Nurse, P. (1995). Fission yeast cell morphogenesis:

identification of new genes and analysis of their role during the cell cycle. *J Cell Biol* **131**, 1529-

38.

Wang, N., Lee, I. J., Rask, G. and Wu, J. Q. (2016). Roles of the TRAPP-II Complex

and the Exocyst in Membrane Deposition during Fission Yeast Cytokinesis. *PLoS Biol* **14**,

e1002437.

Wei, B., Hercyk, B. S., Habiyaremye, J. and Das, M. (2017). Spatiotemporal Analysis

of Cytokinetic Events in Fission Yeast. *J Vis Exp*.

Wei, B., Hercyk, B. S., Mattson, N., Mohammadi, A., Rich, J., DeBruyne, E., Clark,

M. M. and Das, M. (2016). Unique spatiotemporal activation pattern of Cdc42 by Gef1 and Scd1

promotes different events during cytokinesis. *Mol Biol Cell* **27**, 1235-45.

Wu, C., Lytvyn, V., Thomas, D. Y. and Leberer, E. (1997). The phosphorylation site for

Ste20p-like protein kinases is essential for the function of myosin-I in yeast. *J Biol Chem* **272**,

30623-6.

Wu, J. Q., Kuhn, J. R., Kovar, D. R. and Pollard, T. D. (2003). Spatial and temporal

pathway for assembly and constriction of the contractile ring in fission yeast cytokinesis. *Dev*

Cell **5**, 723-34.

 FIGURE LEGENDS

Figure. 1. **Endocytic events at distinct sites in the cell show differential dynamics.**

A. Endocytic events occur at different regions of polarized *S. pombe* cells. Colored tracks

indicate trajectories of individual Fim1-mEGFP patches. **B.** Lifetimes of Fim1-mEGFP

labeled endocytic patches at each region. **C.** Mean Fim1-mEGFP peak intensity in

endocytic patches at each region. **D.** Mean speed of endocytic patches at each region

averaged from individual cells. **E.** MSD (mean square displacement) of Fim1-mEGFP

patches calculated for each region. **F.** Diffusion coefficient of Fim1-mEGFP patches at

each region. (n ≥ 28 patches per region, N=3; n ≥ 6 cells analyzed per region). Ordinary

one-way ANOVA with Tukey's multiple comparisons. ****, $p < 0.0001$; **, $p < 0.01$

Figure 2. Endocytosis follows Cdc42 activation at the division site.

A. Representative cells showing initial (green dotted box) and uniform (solid green box)

Fim1-mEGFP appearance at the division site relative to Sad1-mCherry labeled spindle

pole body (SPB) separation (purple arrowhead). **B.** Quantification of A, where time 0 =

SPB separation. **C.** Representative cells showing initial (purple dotted box) and uniform

(solid purple box) Fim1-mCherry appearance relative to CRIB-3xGFP appearance (green

box) at the division site. **D.** Quantification of C, where time 0 marks CRIB-3xGFP

appearance (green arrowheads) at the division site. (n ≥ 8 cells per genotype per

experiment). Unpaired Student's *t*-test. ****, $p < 0.0001$; ***, $p < 0.001$; **, $p < 0.01$, scale

bar=5μm

Figure 3. The Cdc42 GEFs Gef1 and Scd1 differentially regulate endocytosis.

A. Representative montages of Cdc15-tdTom and Fim1-mEGFP at growing cell ends in the indicated genotypes. **B-G.** Mean patch intensities of Fim1-mEGFP at growing cell ends, the division site, and cell sides. (n ≥ 20 endocytic patches per triplicate experiment). **H.** and **I.** Patch lifetimes of Fim1-mEGFP at each region in the indicated genotypes. (n ≥ 20 endocytic patches, N=3). Ordinary one-way ANOVA with Tukey's multiple comparisons. ****, $p<0.0001$; ***, $p<0.001$; **, $p<0.01$; *, $p< 0.05$; ns, not significant

Figure 4. Pak1/Orb2 kinase is required for timely initiation of endocytosis at the division site. A. Representative *orb2+* and *orb2-34* cells showing initial (green dotted box) and uniform (solid green box) Fim1-mEGFP appearance at the division site relative to Sad1-mCherry to mark spindle pole body separation (purple arrowhead. **B-C.** Quantification of A, where time 0 = SPB separation. (n ≥ 8 cells per genotype per experiment). Unpaired Student's *t*-test. **, $p<0.01$; *, $p< 0.05$, scale bar=5µm

Figure 5. Pak1/Orb2 kinase is required for normal endocytic patch dynamics at the cell ends.

A. Representative cells showing Fim1-mEGFP endocytic events tracked via TrackMate. **B-G.** Mean patch intensities of Fim1-mEGFP at growing cell ends, the division site, and cell sides in the indicated genotypes and conditions. (n ≥ 20 endocytic patches per experiment). **H-I.** Patch lifetimes of Fim1-mEGFP at each region in the indicated genotypes and conditions. (n ≥ 20 endocytic patches, N=3). Ordinary one-way ANOVA with Tukey's multiple comparisons. ****, $p<0.0001$; ***, $p<0.001$; **, $p<0.01$; *, $p< 0.05$; ns, not significant

Figure 6. The type 1 Myosin Myo1 regulates endocytic patch dynamics at the cell ends, division site and the cell sides.

A. Representative cells showing Fim1-mCherry endocytic events tracked via TrackMate in the indicated strains. **B.** Patch lifetimes of Fim1-mCherry at each region in *myo1*[+] and *myo1Δ*. **C**. Patch lifetimes of Fim1-mCherry at each region in *myo1*[+] and *myo1S361A*. **D.** Montages of Fim1-mCherry overtime at the cell ends of the indicated strains. **E-J.** Mean patch intensities of Fim1-mCherry at each region of the indicated strains. (n ≥ 17 endocytic patches, N=3). Ordinary one-way ANOVA with Tukey's multiple comparisons. ****, $p<0.0001$; ***, $p<0.001$; **, $p<0.01$; *, $p< 0.05$; ns, not significant

Figure 7. Myo1 localization to the division site requires Cdc42 and Pak1/Orb2 kinase.

A and **D.** Representative cells showing Myo1-GFP appearance (green box) at the division site relative to Sad1-mCherry to mark spindle pole body (SPB) separation (purple arrowhead) in the indicated strains. **B** and **C.** Quantification of A and D, respectively, where time 0 = SPB separation. (n ≥ 10 cells per genotype per experiment). Unpaired Student's *t*-test. ***, $p<0.001$; **, $p<0.01$, scale bar=5µm

Figure 8. Cdc42 and Pak1/Orb2 kinase promotes Myo1 localization to the cell ends and division site.

A and **E.** Sum Z-projections of Myo1-GFP in the indicated strains and conditions. **B** and **F.** Mean Myo1-GFP intensity measured at cell ends in the indicated strains (n=15 cells per genotype per triplicate experiment). **C** and **G.** Quantification of dwell times of Myo1-

839　GFP in indicated strains. (n≥15 dwell time events per genotype. **D** and **H.** Montages

840　showing Myo1-GFP dwell time dynamics at individual cell ends. One-way ANOVA with

841　Tukey's multiple comparisons post hoc test. ns, not significant; *, $p<0.05$; **, p.<0.01; ***,

842　$p<0.001$; ****, p<0.0001, scale bar=5μm

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8