Collecting, detecting and handling non-wear intervals in longitudinal light exposure data

Carolina Guidolin[1, 2, [0009-0007-4959-2667]], Johannes Zauner[1, 2, [0000-0003-2171-4566]], Steffen Lutz Hartmeyer[3, [0000-0002-2813-2668]], Manuel Spitschan[1, 2, 4, 5, [0000-0002-8572-9268]] *

1 TUM School of Medicine and Health, Department Health and Sports Sciences, Chronobiology & Health, Technical University of Munich, Munich, Germany

2 Max Planck Institute for Biological Cybernetics, Max Planck Research Group Translational Sensory & Circadian Neuroscience, Tübingen, Germany

3 Laboratory of Integrated Performance in Design (LIPID), School of Architecture, Civil and Environmental Engineering (ENAC), École Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland

4 TUM Institute for Advanced Study (TUM-IAS), Technical University of Munich, Garching, Germany

5 TUMCREATE Ltd., Singapore, Singapore

* Corresponding author: Manuel Spitschan

Email address: manuel.spitschan@tum.de

Phone number: +49 7071 601 1670

Address: Max Planck Institute for Biological Cybernetics, Max-Planck-Ring 8-14, 72076 Tübingen, Germany

Keywords: light exposure, behaviour, wearables, non-wear behaviour

Abstract

In field studies using wearable light loggers, participants often need to remove the devices, resulting in non-wear intervals of varying and unknown duration. Accurate detection of these intervals is an essential step during data pre-processing. Here, we deployed a multi-modal approach to collect non-wear time during a longitudinal light exposure collection campaign and systematically compare non-wear detection strategies. Healthy participants (n=26; mean age 28±5 years, 14F) wore a near-corneal plane light logger for one week and reported non-wear events in three ways: pressing an "event marker" button on the light logger, placing it in a black bag, and using an app-based Wear log. Wear log entries, checked twice daily, served as ground truth for non-wear detection, showing that non-wear time constituted 5.4±3.8% (mean±SD) of total participation time. Button presses at the start and end of non-wear intervals were identified in >85.4% of cases when considering time windows beyond one minute for detection. To detect non-wear intervals based on black bag use and lack of motion, we employed an algorithm detecting clusters of low illuminance and clusters of low activity. Performance was higher for illuminance (F1=0.78) than activity (F1=0.52). Light exposure metrics derived from the full dataset, a dataset filtered for non-wear based on self-reports, and a dataset filtered for non-wear using the low illuminance clusters detection algorithm showed minimal differences. Our results highlight that while non-wear detection may be less critical in high-compliance cohorts, systematically collecting and detecting non-wear intervals is feasible and important for ensuring robust data pre-processing.

Introduction

The use of light loggers to measure personal light exposure

Ocular light exposure in everyday life is increasingly being recognised for its potential to influence human physical and mental wellbeing, including cognitive functions, sleep, circadian health, and eye health (Brown et al., 2022; Muralidharan et al., 2021). In this framework, measuring light exposure in the real world is essential to understand what light exposure individuals receive in their daily lives and the cultural, geographical, architectural, and behavioural contexts underlying their exposure patterns (Biller et al., 2024; Guidolin et al., 2024). The measurement of light exposure in free-living conditions is largely accessible in research contexts due to the availability of light loggers, that is, wearable devices recording light exposure over extended periods of time (Hartmeyer et al., 2022; van Duijnhoven et al., 2024). Depending on the intended measurement location, light loggers come in various form factors, including wrist-watches (wrist level measurements), pendants or brooches (chest level measurements), and removable attachments to non-prescription glasses frames (near-corneal plane level measurement) (Hartmeyer et al., 2022). The collected light exposure data can either be directly analysed to classify exposure patterns or also aggregated and summarised in light exposure metrics, which can in turn be associated with health-related outcomes of interest (Spitschan et al., 2022). The versatility of wearable light loggers has led to their use for different research applications (for a detailed review of studies using light loggers in the context of circadian rhythms, alertness, cognitive performance and mood, see Hartmeyer et al., 2022). Most commonly, light loggers have been used in observational field studies aiming to capture the effects of light exposure on sleep (Mead et al., 2023), alertness (Didikoglu et al., 2023), and metabolic function (Reid et al., 2014). Furthermore, light loggers integrated into actigraphy devices have been used in large-scale epidemiological studies to assess light exposure as a potential risk factor for disease prevalence observed at follow-up (Burns et al., 2023; Windred et al., 2024; Windred et al., 2024). Finally, a few studies have employed light loggers in clinical settings to monitor compliance with prescribed light regimes (Sit et al., 2018) or in occupational health to assess whether employees meet target light levels during working hours (Peeters et al., 2020).

Data quality challenges in pre-processing light exposure data: non-wear intervals

Light logging also poses challenges to researchers interested in extracting light metrics from measured light levels, including data quality assessment. The raw wearable data is usually exported as a singular data file containing a time series of light levels and other logged variables, such as activity and external temperature, depending on the individual device (van Duijnhoven et al., 2024). These light data can contain artefacts and data recorded while the device was not worn (Hartmeyer et al., 2022). In fact, in field studies using light loggers, participants may need to remove the devices, such as when taking a shower, or during certain physical activities and sports. Moreover, participants may forget to wear the device or choose to remove it for miscellaneous reasons (Stefani et al., 2024). These periods of non-wear result in invalid data included in the collected datasets. To date, best practices for data cleaning pipelines have not been systematically investigated yet, and implemented strategies are seldom reported in the methods section of research papers (Hartmeyer et al., 2022). Hence, it is currently unknown whether non-wear intervals impact the derived light exposure metrics. It is important to note that the challenge of detecting non-wear time is not exclusive to wearable light loggers, but common across other research wearable devices, such as wrist-worn actimeters used for measuring rest and activity rhythms in field settings. In actimetry research, a variety of algorithms leveraging skin temperature, skin conductance, and (lack of) wearable movement have been developed to detect non-wear intervals (Choi et al., 2012; Pilz et al., 2022; Van Der Donckt et al., 2024). However, skin temperature and conductance might not be available in light loggers, or it might not be a useful measure to detect non-wear for light loggers that are not placed directly on the participants' skin. Furthermore, contrary to movement in actimetry, non-wear time of light loggers is not necessarily characterised by a lack of light signal, since leaving the device on a window sill could lead to light levels similar to those of someone sitting in the same room and wearing the device. Some studies have addressed this issue by relying on intervals of device inactivity to identify invalid periods, using cut-off lengths ranging from 5 to 120 minutes of inactivity (Hartmeyer et al., 2022). Other approaches to detect non-wear intervals include relying on self-reported non-wear times, combining minimal variability in light signal and activity to detect non-wear

time, and detecting stability in device tilt angles (Daugaard et al., 2017; Price et al.,
2019; Stefani et al., 2024; Van der Maren et al., 2018). However, whether these
methods were data-driven or involved manual inspection of participants' files often
remains unclear. Furthermore, even when data-driven techniques are employed, no
study thus far has systematically analysed the performance of algorithmic
approaches using a ground truth for comparison.

The current study

The current study aims to describe and compare strategies for collecting, detecting
and handling non-wear time of light exposure data using a high-quality ground truth
for non-wear time intervals. To achieve this, during a week-long ambulatory light
exposure study, we first collected non-wear information from three independent
sources: participants' self-reports, button presses on the light logger, and placement
of the light logger in an opaque black cloth bag during non-wear time. Self-reported
entries were checked twice daily to ensure high data quality and used as ground
truth for non-wear interval detection. Second, we examine how non-wear detection
varies based on non-wear sources. As part of this analysis, we employ an algorithm
to detect clusters of low illuminance resulting from black bag use during non-wear,
and clusters of low activity resulting from lack of motion during non-wear. We discuss
its performance and how it could be leveraged to develop robust non-wear detection
strategies further. Lastly, we assessed the impact of non-wear data on the accuracy
of daily light exposure metrics.

Methods and materials

Procedure

This study aimed to identify individual and contextual factors contributing to light
exposure patterns in free-living conditions, and a detailed protocol of the study was
previously described (Guidolin et al., 2024). The analyses related to this study
objective will be reported in a separate publication. In the current paper, we use the
data collected during this data collection campaign to investigate collection, detection
and handling of non-wear behaviour. Healthy participants aged 18 to 55 years were
recruited to participate in a study on light exposure in everyday life lasting for a week

(Monday to Monday) in Tübingen, Germany (48.5216° N, 9.0576° E) between

August and November 2023. At time of recruitment, we aimed for a convenience

sample of N=24 due to resource and time constraints. Later systematic analyses of

sample size and statistical power requirements for light exposure studies in the field

have validated that a sample size of N=24 provides a 80% statistical power for most

light exposure metrics (Zauner et al., 2024).

Near-corneal plane light logging

Participants wore a small commercial, multispectral 10-channel light logger

(ActLumus, Condor Instruments, São Paolo, Brazil) able to measure photopic

illuminance and melanopic equivalent daylight illuminance (mEDI). The device

further contains an accelerometer, temperature sensor, and an 'event' button

accessible to the participants. The light logger was inserted into a custom-made 3D-

printed holder, which was in turn attached to the frame of non-prescription glasses

(**Figure 1**). The glasses were available in sizes S, M and L, and provided by the

brand Firmoo (**Figure 1** and **Supplementary Materials**). This set-up was chosen to

ensure light exposure is measured at the near-corneal plane, which is less

susceptible to clothing occlusion than wrist and chest measurements and a more

accurate measurement point for measuring the physiological role of ocular light

exposure. The sampling interval of the light logger was set to 10 seconds to achieve

highly temporally resolved light exposure data. To ensure clarity in our explanations

to participants and in this document, we referred to the combination of the non-

prescription glasses and the mounted ActLumus light logger together as "light

glasses". Participants were instructed to wear the light glasses as much as possible

throughout their participation in the study and only remove them in case of sleep,

water sports, contact sports and during showering. Participants were compensated

€10 for each day of wearing the light glasses for at least 80% of their waking hours.

Logging and coding of non-wear events

Throughout the week, participants reported (non-)wear events, which included taking

the light glasses off, placing the light glasses back on, and taking the light glasses off

before sleep. These events were logged in three ways: (1) Pressing the event button

on the light logger, (2) placing the light glasses in an opaque black-cloth bag (UVEX

microfibre bag for spectacles) for the entire duration of non-wear, and (3) manually

entering the time of this event in a Wear log on the mobile app MyCap (Harris et al.,

2022). Participants were instructed as follows. When taking the light glasses off, they

were to press the event button on the light glasses and then place them in the black

bag provided. Participants were then to log this action in the app-based Wear log as

"Taking the light glasses off". Likewise, when putting the light glasses back on, they

were instructed to take them out of the black bag, press the event button and log this

action in the Wear log as "Putting the light glasses on". In the Wear log, participants

also had to indicate whether the light glasses were actually placed in the black bag.

If they forgot the black bag, they had to describe where they placed the light glasses

instead. When taking the light glasses off before sleep, participants were to place

them facing upwards on a bedside table or flat surface near their bed and to log this

action in the Wear log as "Taking the light glasses off before sleep and placing them

on a nightstand". Each Wear log entry included an automatic timestamp of the

moment this action was performed. If they forgot to log an event, participants were

allowed to retrospectively log any of the possible Wear log events as a "past event".

In this case, they were instructed to try their best to remember and retrospectively

report the times of these actions.

This strategy for collecting non-wear intervals in three ways led to three independent

non-wear sources: (1) A binary outcome for each button press (0/1) automatically

added in the raw ActLumus data file, (2) clusters of near zero-lux illuminance levels

while the device was placed in the black bag, and (3) a timestamp for each Wear log

entry within the MyCap record (**Figure 2**). Among these three sources of non-wear

information, the Wear log entries were deemed the cleanest based on continuous

data quality assurance checks (see *Continuous non-wear data quality assurance* in

Data pre-processing), and thus considered as the ground truth for our analyses (see

Analysis of button presses and *Analysis of low illuminance clusters and low activity*

clusters).

Data analysis

202   All processing steps as well as all statistical analyses were performed in R statistical

203   software (version 4.4.5) (R Core Team, 2025), using the packages "tidyverse"

204   (version 2.0.0) (Wickham et al., 2019),"LightLogR" (version 0.9.2) (Zauner et al.,

205   2025), "stats" (base package, version 4.5.0; R Core Team, 2025) and "rstatix"
206   (version 0.7.2; Alboukadel, 2024).

Data pre-processing

Pre-processing of multi-modal data sets, such as the one used in the current study,
is a time-consuming step. It demands a principled, systematic and reproducible
approach. While scientific publications often report "data cleaning" and/or "data
transformation" steps in the Methods section, the entire pre-processing pipeline is
seldom reported in a detailed, systematic way (Hartmeyer et al., 2022) that allows
other investigators to reproduce these analyses. This leads to multiple similar yet
likely not equivalent ways of approaching the same type of data. In the following
paragraphs, we summarise the pre-processing approach taken in the current study
for handling each of the three non-wear sources to inspire other researchers working
with wearable light loggers and facing similar challenges. Additional detailed
information on our pre-processing strategy can be found in the **Supplementary**
Materials.
Our pre-processing pipeline can be divided into two steps: (1) continuous non-wear
data quality assurance, taking place during the study, and (2) data cleaning and
transformation, taking place after the end of data collection.

Continuous non-wear data quality assurance

Throughout the study, we performed data quality assurance as participants were
enrolled in the study. This step of our pre-processing pipeline was only applied to the
Wear log entries, as information about button presses and the use of the black bag,
as indicated by low illuminance values, was only available upon study completion. A
researcher checked participants' incoming Wear log entries at least twice daily, once
in the morning and once in the late afternoon. This quality assurance was performed
using the online platform REDCap (Research Electronic Data Capture; Harris et al.,
2009), which was used for survey data collection and provides the backend with
MyCap (Harris et al., 2022). The goal of this continuous check was to ensure that the
collected data could be trusted and increase participants' adherence to the protocol,
thus ensuring high data quality throughout the study. For each participant, a
spreadsheet was used to log the manual quality assurance and colour-coding valid
Wear log entries as green and invalid (unclear and/or missing) entries as red. **Figure**

3 illustrates how invalid entries were identified and adjusted, and further examples of

invalid entries are provided in the **Supplementary Materials**.

Data cleaning of Wear log entries

A detailed, step-by-step description of the data cleaning and transformation

approach undertaken for the Wear log entries is provided as a flowchart in the

Supplementary Materials (**Figure S1**). The goal of the Wear log pre-processing

pipeline was to identify intervals that participants spent wearing the light glasses

(Wake 🗆on🗆), not wearing the light glasses while awake (Wake 🗆off🗆), and not

wearing the light glasses during sleep (Sleep 🗆off🗆). To further examine the validity

of the Wear log entries beyond the manual quality assurance procedures performed

during the study (see above under *Continuous non-wear data quality assurance*),

two additional types of quality checks were performed:

1) Quality check #1: Identifying identical timestamps for two consecutive Wear log states,
i.e. identical and consecutive Datetime values.
2) Quality check #2: Identifying consecutive identical Wear log states, i.e. identical and
consecutive Wake 🗆on🗆, Wake 🗆off🗆, or Sleep 🗆off🗆 values

These quality checks were performed on a total of 797 observations (of which only

n=51 retrospectively logged). Eight entries (1%) failed Quality check #1, and 14

entries (1.76%) failed Quality check #2. These invalid entries were adjusted

manually by referring to the raw files. We describe how these adjustments were

applied in the **Supplementary Materials** (Data cleaning of Wear log entries: details

on manually adjusted entries). After these adjustments, a rerun of the quality checks

showed no erroneous entries after applying these corrections. The next step was to

only include all entries up until and including the "Putting the light glasses on" entry

after sleep on the last study day (Monday), before the participants returned to the

laboratory to return the materials. Lastly, the Wear log entries were cleaned,

arranged chronologically for each participant and integrated into the light logger

dataset. This led to a final data frame where each light logger measurement,

including a measurement of illuminance and activity, was labelled based on the

Wear log *State* (Wake 🗆on🗆, Wake 🗆off🗆, Sleep 🗆off🗆).

Data cleaning of button presses and low illuminance clusters

For the other two sources of non-wear information, namely the button presses and
clusters of low illuminance levels, pre-processing was less extensive than for Wear
log entries. Button presses within the same 60-second interval were considered
invalid, i.e. caused by the participant accidentally pressing the button more than
once to indicate the same (non-)wear action. These invalid consecutive button
presses were removed, so only the first was kept. Regarding low illuminance
clusters, the mEDI values corresponding to the Wear log Sleep □off□ state were
excluded before implementing our custom-made algorithm, as the scope was to
detect either wear or non-wear time during waking hours. Further transformations on
illuminance levels throughout data analysis are detailed in the Results sections.
When visualising illuminance levels on a logarithmic scale (**Figure 7A**), 0 values
were handled by adding a value of 1 to all mEDI values. However, all calculations
related to mEDI were performed without conversion to logarithmic scale and on the
raw data, which included values of mEDI=0 lux.

Description of wear status based on ground truth

The first part of our analysis consists of a description of wear status based on the
ground truth, i.e. the (non-)wear information obtained from the Wear log. Specifically,
we examined the time spent in each Wear log state across the week, the distribution
of Wake □off□ status during weekdays and weekend days, and the timing of Wake
□off□ status across the day.

Analysis of button presses

To compare non-wear information obtained by the button presses to the ground
truth, we examined whether the start and end of each Wake □off□ interval aligned
with a button press. More specifically, we classified each Wake □off□ interval based
on the presence of a button press within a given time window from the Wear log
timestamp. Non-wear intervals were categorised as follows: intervals not bounded by
a button press at the start and/or the end were labelled as open, while intervals
bounded by two button presses, one at the start and one at the end, were labelled as
closed. To investigate how the percentage of open and closed intervals varied with
increasing time window lengths, we performed this analysis using eight windows
ranging from one to eight minutes, increasing in one-minute steps.

Analysis of low illuminance clusters and low activity clusters

To compare non-wear information obtained by the use of the black bag to that of our

ground truth, we used an algorithm (Hartmeyer et al., 2024) to detect clusters of low

illuminance (melanopic equivalent daylight illuminance, mEDI) values in the light

logger dataset. This algorithm takes three input parameters: (1) A maximum

threshold for the boolean variable of interest, for which the clusters should be found

(in our case, mEDI), (2) a minimum length of the cluster, and (3) a maximum

interruption length of the cluster by values above threshold (in our case, mEDI

values above threshold). With these three input parameters, the algorithm labels

every illuminance observation in a participant's light exposure dataset as belonging

to a cluster of low illuminance levels or not. To assess the performance of this cluster

detection algorithm in identifying the same non-wear intervals identified by our

ground truth, we constructed a precision-recall (PR) curve. PR curves provide a

graphical illustration of a classifier's performance across several threshold values (in

our case, mEDI values), with positive predictive value (PPV, or precision) on the Y-

axis and true positive rate (TRP, or recall) on the X-axis. In the context of our

classifier, precision indicates the number of instances that were predicted as non-

wear by the algorithm and were non-wear according to the Wear log. On the other

hand, recall indicates the number of instances correctly identified as non-wear by the

algorithm out of all actual non-wear instances (i.e. Wear log Wake ☐off☐ instances).

Importantly, PR curves are preferred to commonly used receiver operating

characteristic (ROC) curves in scenarios of class imbalance, as in our dataset. In this

case, the small percentage of non-wear relative to wear intervals could risk

overstating the algorithm's performance, when evaluating it with a ROC curve (Cook

& Ramadas, 2020).

To define the optimal combination of the minimum cluster length and maximum

interruption length to detect clusters of low illuminance in our dataset, PR curves

were first built with varying thresholds of these two variables, while keeping a

threshold of <1 mEDI lux (Supplementary materials, **Figure S2 and Figure S4**). A

minimum length of clusters of 21 minutes and a maximum interruption of 0 minutes

led to the highest F1 score (F1=0.78). F1 is a metric representing the harmonic

mean of precision and recall, with values ranging from 0 to 1 (Cook & Ramadas,

2020). Thus, these input parameters were used in our algorithm to build the PR

curve with varying mEDI thresholds. Because clusters of low activity are often used

to detect non-wear in field studies investigating sleep and light exposure with

wearables similar to our light logger (Hartmeyer et al., 2022), we implemented our

algorithm to also detect clusters of low activity in our dataset. Activity was quantified

from the ActLumus built-in accelerometer as proportional integrating measure (PIM),

a quantity directly calculated by the device throughout the study. As for illuminance,

we first built PR curves for minimum cluster length and maximum cluster interruption

length, while keeping an activity threshold of <5 (Supplementary materials, **Figure**

S3 and Figure S4). A minimum cluster duration of 12 minutes, and a maximum

cluster interruption of 0 minutes led to the highest F1 score (F1=0.52), and this

combination was then used to build a PR curve with varying PIM thresholds.

Comparison of light exposure metrics

To determine whether handling non-wear intervals identified by either the Wear log

or by the algorithm would result in different values of light exposure metrics

compared to not handling non-wear intervals, we calculated 14 commonly used light

exposure metrics (Hartmeyer & Andersen, 2023; Zauner, Hartmeyer, et al., 2025)

and compared them across three variants:

1.   Raw dataset: All mEDI values included, i.e., ignoring non-wear information;

2.   Wear log-corrected dataset: All mEDI values corresponding to a *Wake off* interval

replaced with "NA", R's standard placeholder for missing values;

3.   Algorithm-corrected dataset: All mEDI values corresponding to algorithm-detected

clusters of low illuminance replaced with "NA".

Specifically, we calculated the following 14 metrics:

-    Metrics representing temporal variability of light exposure: across days (interdaily

stability, IS) and within days (intradaily variability, IV) (n=2);

-    Metrics quantifying timing-related light exposure characteristics: Mean timing of light

above 250 lux ($MLiT_{250}$), mean timing of light above 1000 lux ($MLiT_{1000}$), first timing of

light above 10 lux ($FLiT_{10}$), first timing of light above 250 lux ($FLiT_{250}$), first timing of light

above 1000 lux ($FLiT_{1000}$), last timing of light above 10 lux ($LLiT_{10}$), last timing of light

above 250 lux ($LLiT_{250}$), last timing of light above 1000 lux ($LLiT_{1000}$) (n=8);

-    Metrics quantifying duration-related light exposure characteristics: Time above 250 lux

(TAT_{250}), and time above 1000 lux (TAT_{1000}) (n=2).

-    Metrics quantifying intensity of light levels: average light exposure during daytime and

mean across the brightest 10 hours (M10m).

All metrics were calculated for every day of the study where full light exposure data
were available (n=6, since the first Monday was incomplete for all participants). IS
and IV were calculated directly on the entire dataset, the outcome being a single
participant value for each metric. Average light exposure during daytime was also
calculated on the entire dataset, however, only for daytime periods, defined by the
time from civil dawn until civil dusk, using photoperiod information at our specific
latitude   and   longitude   coordinates   (see
https://tscnlab.github.io/LightLogR/reference/photoperiod.html for more information).
For the remaining metrics, values were first calculated for each of the six study days,
and then mean and standard deviation values were calculated across these six
values. This led to a single mean and standard deviation participant value for each
metric. For average daytime light exposure and M10m, mEDI values were first log10
transformed, and a small value was added to 0 values prior to transformation. Values
were then back-transformed (Zauner et al., 2025). Differences in the selected 14
metrics across the three datasets were assessed using Friedman test due non-
normality of the data. Post-hoc pairwise comparisons were then performed using
Wilcoxon test for paired data, and corrected for multiple comparisons with the
Benjamini-Hochberg method (Benjamini & Hochberg, 1995). Results are presented
here using a significance threshold of 0.05. Finally, to further explore differences in
metrics across the three datasets beyond significance testing, we conducted
equivalence testing on each post-hoc pairwise comparison to probe equivalence in
light exposure metrics. Given the non-normal distribution of the data and the
relatively small sample size, we employed bootstrapping to estimate the sampling
distribution of the effect size for each pairwise comparison. To define the smallest
effect size of interest (SESOI) for equivalence testing, we drew on prior literature
examining weekday-to-weekend variation in personal light exposure (lux*hours), a
common covariate in light exposure analysis. Based on large datasets from
Switzerland (n=20 participants, 554 participant-days) and Malaysia (n=19
participants, 509 participant-days), where participants wore a light-measuring device
continuously for one month (Biller, Zauner, et al., 2024), the observed effect sizes
were 0.27 and 0.33, respectively. We therefore set the SESOI at d=0.3 (Cohen's d,
i.e., mean difference in metric divided by the standard deviation; Cohen, 2013).
Equivalence was concluded when the SESOI (±0.3) completely covered 90% of the
bootstrapped effect sizes (90% CI).

Results

Final dataset

A total of 30 participants enrolled in the study and began data collection. Of these,

one participant withdrew from the study, and participation was terminated early for

the other two participants. This was due to a lack of compliance with and

understanding of the study procedure, whereby participants did not fill in the daily

questionnaires, despite receiving various prompts from the researchers. This led to a

sample of 27 participants who completed the study. One participant was excluded

from the analyses because of technical problems with the light logger leading to

unrecoverable data loss. Thus, our analyses were performed on a final sample of

n=26 aged between 22 and 46 years (mean±1SD: 28±5 years, 14 female).

Description of wear status based on ground truth

To explore how participants engaged in non-wear behaviours across the week, we

first visually inspected light exposure, button presses and self-reported (Wear log)

wear status for each participant across all study days (**Figure 4**). This allowed us to

gain a qualitative understanding of how the three sources of non-wear (Wear log,

button presses, and use of black bag) compared in our study. To quantify wear

status according to our ground truth, we then calculated the amount of time

participants spent in each Wear log state throughout the week: Wake 'on▢, Wake

▢off▢ and Sleep ▢off▢ (**Figure 5A**). Based on Wear log entries, participants spent

57.7% of their total participation time wearing the light logger (Wake ▢on▢ mean,

SD=6.3%, range= 42.2%–72.7%), 5.4% of time not wearing the light logger (Wake

▢off▢ mean, SD=3.8%, range=0.3%–13.0%), and 37.0% of time with the light logger

off during sleep (Sleep ▢off▢ mean, SD=4.6%, range=25.9%–46.1%). When

comparing non-wear time between weekdays and weekend days, a similar trend was

observed, with participants spending a small amount of time not wearing the light

glasses on weekdays (mean±SD= 4.7±3.8%, range=0%–16.0%) as well as during

weekend days (mean±SD=7.0±8.0%, range=0%–29.9%) (**Figure 5B**). Moreover, we

investigated non-wear time during wake periods only, as the participants were

instructed to remove the light glasses while sleeping. Non-wear time across

participants was 8.6±6.2% (mean±SD). To gain a better understanding of how the

self-reported sleeping periods in the Wear log (Sleep 'off') related to participants'
actual sleep time, we compared these timestamps with participants' self-reported
sleep time measured with a daily sleep diary. The results highlighted that, for most
participants, the two measures differed by less than 30 minutes (**Figure S5)**.

To further examine patterns in participants' non-wear behaviour, we inspected
duration and timing of reported non-wear episodes across the day. The median
duration of non-wear was 00:37, with the 25^{th} and 75^{th} percentiles at 00:19 and
01:27, respectively. The minimum and maximum durations were 00:02 and 10:24 (all
values reported as hh:mm). As a proxy of non-wear timing, we calculated the centre
point of each Wake ☐off☐ episode (**Figure 5C**). Non-wear episodes occurred
continuously across 24 hours, with the exception of two time windows during night
hours (between 02:00 and 03:00 and between 04:00 and 05:00). Density estimates
for non-wear episodes showed an increase throughout the day and reached a peak
at 20:00 with 20 non-wear episodes falling into this time bin. The number of non-
wear time episodes decreased in the evening hours.

Analysis of button presses

We next compared non-wear information obtained by the button presses to non-wear
status information from our ground truth. Visual inspection suggested that only a few
button presses (n=15) were not performed in proximity of a self-reported change in
wear status. We examined the presence of button presses at the start and end of
each Wake ☐off☐ interval using different time windows. When considering button
presses within a 1-minute window from a start and/or end of a Wake ☐off☐ interval,
only 64.1% of n=198 intervals were bounded by a button press on both ends and
thus classified as closed. Expanding window sizes to more liberal cut-offs of 2, 3, 4,
5 and 6 minutes increased this number to 85.4%, 87.4%, 88.4%, 88.9%, and 89.9%,
respectively. Adopting longer window sizes of 7 and 8 minutes did not increase the
proportion of Wake ☐off☐ intervals classified as closed, with the percentage of these
intervals seemingly asymptotic at 88.9% for these cut-offs (**Figure 6**).

Algorithmic approach for non-wear detection

To explore the possibility of using clusters of low illuminance and low activity in our
dataset as proxy of non-wear, we first examined illuminance and activity levels

across Wear log states (**Figure 7A**). Participants' median illuminance and median

activity levels across the study showed a similar pattern, whereby similarly low mEDI

and activity values were observed during Wake □off□ and Sleep □off□ intervals

(mEDI mean±SD in Wake □off□ and Sleep □off□: 0.4±1.2 lux and 0.01±0.07 lux,

respectively; activity mean±SD in Wake □off□ and Sleep □off□: 4.5±22.8 and

0.0±0.0, respectively). On the other hand, higher median illuminance and activity

values were observed during Wake □on□ intervals (mean±SD: 62.4±66.6 lux and

61.5±46.1, respectively). Based on this observation, we used a previously developed

algorithm (Hartmeyer et al., 2024) for detecting low illuminance clusters and low

activity clusters in our dataset. The performance of this algorithm was examined

using a precision-recall (PR) curve with varying maximum mEDI thresholds (from 1

lux to 10 lux, at 1-step increments), and activity thresholds (from 5 to 50, at

increments of 5) (Figure 7B). Overall, detecting clusters of low illuminance performed

better than detecting clusters of low activity in predicting non-wear. This difference

was reflected in the F1 score, with the highest score for mEDI F1=0.78 (with a mEDI

threshold of 1 lux) and the higher score for activity F1=0.52 (with an activity

threshold of 5).

For illuminance, setting the threshold to smaller mEDI values for cluster detection led

to a higher PPV (or precision) compared to higher mEDI thresholds (PPV=0.71 for

<1 mEDI lux and PPV=0.33 for <10 mEDI lux). The gain in precision with decreasing

mEDI threshold values came at a minimal cost for the TRP (or recall), with

TRP=0.89 and TRP=0.81 for thresholds of <10 mEDI lux and <1 mEDI lux,

respectively. Furthermore, we tested the algorithm performance separately for

daytime periods (starting at dawn and ending at dusk) and nighttime periods (starting

from dusk and ending at dawn). The PR curve revealed a better performance during

daytime periods (highest F1=0.80, with mEDI threshold of 2 lux), compared to

nighttime periods (highest F1=0.76, with mEDI threshold of 1 lux) (**Figure S6**). Of

note, daytime performance consistently exceeded nighttime performance, with the

lowest daytime performance (F1=0.68 for mEDI threshold <10 lux) still higher than

nighttime performance for all mEDI thresholds except <1 lux and <2 lux. As the PR

curve in **Figure 7B** shows the aggregate classification accuracy, we next probed

whether participants had idiosyncratic non-wear behaviours and analysed if

classification accuracy depended on each participant. This was indeed the case

(**Figure 8**). Furthermore, we did not identify a clear pattern for classification accuracy

of low illuminance clusters and participants' self-reported compliance with using the

black bag during non-wear time. For example, for a participant who reportedly used

the black bag for all non-wear (Wake □off□ events), both precision and recall of the

classifier were low (**Figure 8, PID 214**). Likewise, for a participant who reportedly

never placed the light glasses in the black bag when taking them off, the recall of the

classifier was very high (**Figure 8, PID 218**).

Examination of algorithm misclassification

Since detecting clusters of low activity would be an easy-to-implement strategy to

detect non-wear, we examined reasons behind the poor performance of the

algorithm for detection of non-wear based on clusters of low activity. We first

analysed whether using PIM as activity parameter was a justified choice, and

whether changing activity parameter would yield a different algorithm performance.

ActLumus files contain three parameters that quantify activity: proportional

integration mode (PIM, a measure of activity levels), time above threshold (TAT, a

measure of time spent in motion), and zero crossing mode (ZCM, a measure of

frequency of movement) (Pilz et al., 2022). Our results indicate that the performance

of our algorithm was similarly poor for all three activity-quantifying parameters

(Supplementary materials, **Figure S7**). Furthermore, we investigated whether pre-

processing activity (PIM) values would lead to a better classification performance.

We applied several pre-processing methods including log-transformation, smoothing,

and normalisation of PIM values, and showed that none of these strategies

benefitted classification performance (Supplementary materials, **Figure S8**). Given

the superior performance in detecting clusters of low illuminance using a mEDI

threshold of <1 lux (F1=0.78), we then applied our algorithm to the dataset using this

threshold, and focused subsequent analyses on illuminance rather than activity. In

order to understand in which circumstances our algorithm failed at predicting non-

wear (resulting in false positive and false negative instances), we next analysed

characteristics of these instances that might have contributed to misclassification.

Specifically, we calculated the duration of each misclassified instance, the mean

melanopic illuminance during this interval, and whether the participants reported

using the bag during this time. Using these three variables, we identified four

potential causes of misclassification and labelled each false negative and false

positive interval in our dataset accordingly:

1. Algorithm limitation or bag not actually used: False negative intervals (>3 minutes)
where participants reported using the bag and with mean mEDI >1 lux. These false
negatives could be caused by participants not using the black bag despite reporting
doing so, or by the algorithm input parameters preventing correct classification (e.g.
cluster duration <21 minutes).

2. Bag not used: False negative intervals of any duration where participants reported
not using the bag and with mean mEDI >1 lux. In this case, since the bag was
reportedly not used and the mEDI values suggest similarly, no clusters of low
illuminance could be detected by the algorithm.

3. Low illuminance during wear: False positive intervals (<3 minutes) with mean mEDI
≤1 lux. In this case, individuals received low illuminance during wear time, leading to
the algorithm detecting these intervals as non-wear clusters.

4. Transition state: False negative intervals (≤3 minutes) where participants reported
using the bag and with mean mEDI >1 lux, and false positive intervals (≤3 minutes)
with mean mEDI ≤1 lux. These brief intervals likely represent transitions from one
Wear log state to the next one, where a short time lag between logging wear status
on the Wear log and placement or removal of the light glasses from the black bag
can lead to misclassification.

A summary of misclassified and correctly classified instances can be found in **Table
1**. Furthermore, examples of the algorithm misclassification instances in **Table 1** are
shown in an exemplary participant (**Figure 9**). Due to the high number of transition
states identified as potential causes of our algorithm's misclassification (n=210, or
21.5% of 978 total instances), we implemented a modification to the algorithm-
detected clusters of low illuminance. Specifically, for each cluster of low illuminance
values (i.e. a non-wear cluster), we checked for a transition state label, identified as
described above, immediately before or immediately after the cluster. If a transition
state was detected, the non-wear cluster was expanded as follows. In case of a
transition state detected before the cluster, all observations from the start of the
transition state to the start of the non-wear cluster were re-labelled as non-wear. In
the case of a transition state after the cluster, all observations from the end of the
non-wear cluster to the end of the transition state were re-labelled as non-wear. This
adjustment ensured that each low illuminance cluster adjacent to a transition state
was "padded" with non-wear observations at both ends.

Comparison of light exposure metrics

To assess whether removal of non-wear time would affect light exposure metrics in our data, we calculated 14 light exposure metrics across three datasets. The choice of light metrics was based on commonly used metrics quantifying timing-related characteristics ($MLiT_{250}$, $MLiT_{1000}$, $FLiT_{10}$, $FLiT_{250}$, $FLiT_{1000}$, $LLiT_{10}$, $LLiT_{250}$, $LLiT_{1000}$), duration-related characteristics (TAT_{250} and TAT_{1000}), temporal dynamics (IS and IV) of light exposure patterns, as well as light levels (average light level during daytime hours and mean across the brightest 10 hours, M10m) (Hartmeyer & Andersen, 2023; Zauner et al., 2025). The relationship between each metric calculated in the raw dataset and the corrected, cleaned datasets (Wear log-corrected and algorithm-corrected), as well as a histogram of the delta of the metrics, can be found in **Figure 10**. To test for differences in metrics across the three derived datasets, we performed a Friedman test. The results indicated statistical differences in metrics between the three datasets from all four metric domains: $LLiT_{10}$, $LLiT_{1000}$, TAT_{250}, TAT_{1000}, average light exposure during daytime hours, M10m, and IV. No statistical difference was found for the following metrics quantifying timing-related patterns of light exposure and its temporal variability: $MLiT_{250}$, $MLiT_{1000}$, $LLiT_{250}$, $FLiT_{10}$, $FLiT_{250}$, $FLiT_{1000}$, and IS (**Table S1**). When examining post-hoc comparisons (n=21) using Wilcoxon signed-rank tests with Benjamini-Hochberg correction for multiple comparisons, n=9 pairwise comparisons reached statistical significance, corresponding to light exposure metrics related to duration-related patterns of light exposure as well as its intensity (TAT_{250}, TAT_{1000}, average daytime mEDI, and M10m). For metrics quantifying timing-related patterns of light exposure and its temporal variability that resulted in significantly different outcomes in the three-way comparison ($LLiT_{10}$, $LLiT_{100}$, and IV), no statistical difference across the pairwise comparisons was found. A full list of pairwise comparisons and adjusted p-values is provided in **Table 2**. Furthermore, the means and standard deviation values for all 14 metrics are reported in **Table S2**.

Among the 21 pairwise comparisons investigated, none were conclusively equivalent. For n=9 comparisons, a directed effect was found; i.e. the bootstrapped 90% confidence intervals of the effect size between metrics lay entirely above or below the SESOI region. Specifically, for metric quantifying duration-related patterns (TAT_{250} and TAT_{1000}), raw dataset values were higher than Wear log-corrected

values (TAT_{1000}) as well as algorithm-corrected values (TAT_{250}). The opposite trend
was observed for metrics quantifying light intensity (average daytime mEDI and
M10m): raw dataset values resulted lower than Wear log- and algorithm-corrected
values. The remaining n=12 comparisons were inconclusive. Equivalence test
results for each pairwise comparison are reported in **Table 2**, and corresponding
bootstrapped effect size distributions are shown in **Figure S9**. A summary of
differences in metrics from their uncorrected values (raw dataset), as well as for
differences between correction methods (Wear log-corrected and algorithm-
corrected), can be found in **Table 3.** Furthermore, **while** our choice of SESOI for
equivalence testing was based on literature for weekday-to-weekend effects, other
SESOI thresholds can be relevant in other contexts. We thus further explored how
different SESOI thresholds (0 to 1, at 0.1 steps) might influence the outcome of the
equivalence tests. **Figure S10** illustrates the proportion of bootstrapped effect sizes
falling outside of the SESOI region as a function of SESOI threshold, while **Figure
S11** summarises how the proportion of equivalence conclusions changes depending
on SESOI selection.

Discussion

To the best of our knowledge, this is the first study to systematically assess the
impact of non-wear intervals on light exposure metrics in ambulatory light logging
studies. Our protocol, characterised by continuous data quality checks, was
designed to yield a high-quality ground truth for non-wear intervals, i.e. the self-
reported Wear log entries, essential to evaluate and compare different non-wear
detection strategies. Our results indicated that button presses did not consistently
mark the start and end of non-wear intervals, and that algorithmically detecting
clusters of low illuminance resulting from black bag use had a reasonably strong
performance against ground truth labels (F1=0.78). Lastly, we found that addressing
non-wear intervals in our small, high-compliance dataset only had a marginal impact
on some of the commonly used light exposure metrics examined.

Description of wear status based on ground truth

As part of data pre-processing, we first visually inspected all three sources of non-wear. Relying on visual inspection alone to detect non-wear intervals is time consuming and requires training (Pilz et al., 2022), but it can be used as an initial step for exploring raw data. Visual inspection allowed us to gain a better understanding of participant compliance and concordance between non-wear sources, and provided a qualitative insight into which automated detections (i.e., non-wear algorithms) could be applied to our data. Visualisation of raw light exposure data is possible using open-access packages in programming environments such as the *LightLogR* package (Zauner et al., 2025) or the *pyLight* submodule of pyActigraphy (Hammad et al., 2024), the former also offering options to visualise multiple non-wear sources in the same graph, as shown in Figure 4 of this paper.

When examining self-reported Wear log entries, the relatively low non-wear time (mean±SD=5.4±3.8% of total participation time and 8.6±6.2% of wake-only time) across the sample indicates that participants were compliant with study instructions, spending a substantial amount of time wearing the light glasses during their daily activities. Compliance rates were high for both weekdays and weekend days. Our results align closely with the findings of a recent study where participants wore a near-corneal plane light logger for five days, a duration similar to that of our study. The authors report that, despite participants documenting discomfort and unwanted attention in public settings when wearing the device, adherence to the study instructions was high (89% during waking hours, excluding sleep time) (Stefani et al., 2024). As to the timing of non-wear, the highest number of reported non-wear episodes in our dataset coincided with early evening hours, peaking at 20:00. Stefani and colleagues (2024) also reported slightly higher wear compliance during daytime compared to evening hours during weekdays (92% of recording time during daytime, 71% of recording time the evening). While we lack additional data to understand which specific situation prompted the removal of the light glasses at this time, we speculate that reasons for removal might be similar to those reported in other studies that used near-corneal level light loggers: being in the shower, being in public spaces where wearing the light glasses is thought to cause undesirable reactions

655 from others, performing sports, or experiencing discomfort (Balajadia et al., 2023;
656 Stefani et al., 2024).

Analysis of button presses

When turning to button presses marking the start or end of a non-wear interval, our
results indicated that button press events mostly happened within 1 minute to 6
minutes from the corresponding Wear log entry. Extending the time window beyond
this range did not result in more non-wear intervals labelled as closed, suggesting
that participants occasionally forgot to press the button to mark a non-wear event. To
the best of our knowledge, no previous study has examined the concordance of
button presses with self-reported non-wear time, although some studies mention
instructing participants to use this when removing the light logger (Van der Maren et
al., 2018), as well as using it in visual inspection of the data (Stefani et al., 2024).
Our results suggest that button presses alone are unreliable for collecting accurate
information on non-wear intervals. However, since several available light loggers
have a built-in event marker (van Duijnhoven et al., 2024), collecting non-wear
information using button presses does not add a significant burden to participants
and can help validate non-wear self-reports. This strategy could especially benefit
research studies in which non-wear self-reports are not as clean as in the current
dataset, and no other source of non-wear is collected. Based on our findings, we
recommend using cut-off criteria larger than 1 minute for automatic detection of
button presses corresponding to a self-reported non-wear event, as stricter criteria
could compromise non-wear validation.

Data-driven non-wear detection

As part of our non-wear detection analysis, we applied an algorithm detecting
clusters of low illuminance values caused by the use of the black bag during non-
wear time, as well as clusters of low activity, which might also be indicative of non-
wear time. Evaluating the performance of our algorithm using PR curves, our results
indicate that collecting non-wear information by asking participants to place the light
logger in a black bag might be an effective strategy to facilitate automated detection
of non-wear time in ambulatory light exposure studies. When varying the input
parameters (maximum mEDI levels, minimum cluster length, and maximum cluster

interruption), we found that both the maximum mEDI threshold and the minimum
cluster length had a clear influence on performance, whereas the maximum cluster
interruption length had only a minor effect. Based on these findings, we recommend
adopting a conservative threshold for maximum light levels (<1 mEDI lux). For the
minimum cluster length, durations between 10 and 30 minutes yielded comparable
F1 scores (0.76–0.78), likely reflecting the variable duration of non-wear episodes
among participants. Therefore, in field studies where non-wear duration is
uncontrolled, we suggest prioritising medium-length minimum cluster lengths
(between 10 and 30 minutes) over very short ones (<10 minutes) to improve
detection. More broadly, our findings highlight that maximum light level thresholds
and minimum cluster length should be explicitly considered when developing more
complex algorithms for non-wear detection to ensure both accuracy and
generalizability across study designs.
Our results also highlight that photoperiod information (day and night) must be
considered when deciding on an illuminance threshold for automated non-wear
detection. When only looking at daytime periods (dawn to dusk), the strong contrast
between environmental light (e.g. daylight or electric light) illuminance levels and the
low black bag illuminance levels might lead to better algorithm performance. On the
other hand, during the night period (dusk to dawn), the low intensity of the
environmental light might impair the ability of the algorithm to accurately detect
clusters of low illuminance due to black bag use. While for both daytime and
nighttime lower mEDI thresholds led to better algorithm performance, higher mEDI
thresholds during daytime (<9 and <10 lux) led to a similar performance to low mEDI
thresholds during nighttime (<2 lux). Overall, this result suggests that when using an
automated low-illuminance detection algorithm in periods of naturally lower light
levels (i.e. nighttime), a more conservative illuminance threshold should be adopted.
Considering the poorer performance of detecting low activity clusters compared to
low illuminance clusters, it is important to consider that this was not always observed
when looking at individual PR curves (**Figure 8**). Specifically, for some individuals,
using low illuminance clusters to detect non-wear led to perfect recall, but using
clusters of low activity led to greater precision. Furthermore, the observation that
compliance with bag use did not seem to underlie better algorithm performance for
illuminance might be explained by individuals placing the device in a dark space,
such as a drawer, even when they did not place it in the black bag. We suggest that

relying on both illuminance and activity (i.e., multi-modal input to our algorithm) could
improve automated non-wear detection, particularly in accounting for non-
compliance with black bag use. This approach would increase classification
performance for non-wear intervals where the signal from the two variables is
opposite, such as when the device sits on an illuminated surface (i.e., high light
levels, low activity levels).

Upon examining the reasons behind our algorithm's misclassifications, transition
states were a major contributor to false positives and false negatives. We attribute
this to a time mismatch in non-wear logging: While participants were instructed to
first place the light glasses in the black bag and then log non-wear in the Wear log,
they might have performed these actions in reverse order. As a result, the Wear log
would indicate a non-wear period at time X, while the algorithm would detect a
cluster of low illuminance only at time X+1 minute, when the light glasses were
placed in the black bag. Consequently, the algorithm would label the instances
between X and X+1 minute as false negatives. A similar issue occurs when putting
the glasses back on after non-wear. If participants log this action before taking the
light glasses out of the bag, the algorithm detection will result in false positive
instances until the device is actually outside the bag and being worn. We addressed
this transition state problem post-hoc by padding each non-wear cluster where a
misclassification related to a transition state was identified. A further improvement to
our algorithm would be to identify transition states dynamically as part of the
classification process. It is important to note that our algorithmic approach might not
be the only one suitable for identifying non-wear time in ambulatory light exposure
studies. For example, since our algorithm employs a minimum length during which
light or activity must be below a threshold value, short intervals are likely
misclassified. Algorithms that do not rely on an interval but rather explore signal
characteristics immediately preceding or following a non-wear interval might lead to
better non-wear classification (Syed et al., 2021).

Comparison of light exposure metrics

Our analysis on commonly used light exposure metrics revealed that accounting for
non-wear time intervals only affected n=7 out of n=14 light metrics investigated. Of
note, post-hoc tests revealed that for three ($LLiT_{10}$, $LLiT_{100}$, and IV) of the n=7

affected metrics, no statistical difference across pairwise comparisons was found.
Furthermore, equivalence testing was inconclusive for a small effect size
(SESOI=±0.3), indicating insufficient evidence that the differences in metrics were
small enough to be considered practically equivalent. Thus, while the metrics are not
significantly different between the methods of non-wear handling, they might be in
more sensitive cases (i.e. with higher statistical power).
For n=9 out of n=21 post-hoc pairwise comparisons, a significant difference was
found and equivalence testing indicated a directed effect. Importantly, with the
exception of one metric type (TAT), equivalence testing indicated that differences
were found only between the raw dataset and a corrected dataset, not between the
Wear log-corrected set and the algorithm-corrected set. Interestingly, raw dataset
values were higher than Wear log-corrected values for duration-related metrics, and
the opposite trend was observed for metrics quantifying light intensity. These
diverging trends might be due to duration metrics being sensitive to peaks of high
light exposure (i.e. non-wear periods during which participants are not using the
black bag, since these metrics calculate any value above threshold), and less
sensitive to periods of non-wear in a dim or dark environment (i.e. non-wear periods
during which the black bag is used, since these contribute to zero time above
threshold). On the other hand, intensity-related metrics are more sensitive than
duration metrics to whether or not low light observations in the data are filtered (i.e.
low illuminance values resulting from black bag use). However, it is important to note
that the differences in daily mean TAT_{250} and TAT_{1000} values between the raw and
Wear log-corrected dataset are very small (~2 minutes and ~1 minute, respectively).
This means that, despite the difference being significant, the magnitude of this
difference is very small and likely driven by low variability in participant-level
differences rather than a meaningful change in the metric. The lack of differences
between light metrics could be explained by the timing of non-wear observed in our
dataset. Our participants mostly reported removing the light glasses in the evening
hours, when light levels tend to be low. Thus, the missing data during this period do
not impact light metrics quantifying exposure to bright light of 250 mEDI lux and 1000
mEDI lux. Of note, we calculated differences in ten out of the 14 metrics analysed as
differences in mean values of the metrics across the study. Thus, it could be that our
results would differ when investigating differences in metrics for each day of the
week or for specific times of the day. For example, in studies that examine the

association between evening light exposure levels and health outcomes, filtering
non-wear time using clusters of low illuminance might yield different metrics
compared to calculating metrics on the raw data. In other words, longer recording
periods might make the data more robust to artefacts and missing data, a finding
which is supported by a recent preprint by Biller, Zauner and colleagues (2024),
where multiple hours of missing daily data did not significantly influence average light
exposure metrics for a dataset that spans a whole month. However, as more studies
will need to confirm the generalisability of these findings in different, larger cohorts
and in studies of varying duration, we recommend including non-wear detection as
part of pre-processing pipelines and hereby present an easy-to-implement algorithm
that can serve this purpose.

Towards proposed best practices for collecting and reporting non-wear data

In the current study, participants had to perform three actions to report non-wear:
press a button (event marker) on the device, place it in a black bag, and log this
action in an app-based Wear log. We recognised that this could significantly burden
participants and potentially result in lower data quality and implemented various
countermeasures to minimise this risk. In this paragraph, we outline the strategies
which we found to be successful.
Since the light glasses were very visible on participants' faces, thus potentially
contributing to more instances of non-wear, transparency about the device form
factor was essential in exchanges with participants before screening. During the
screening, emphasising the importance of wearing the device as much as possible
and logging non-wear with honesty was successful in ensuring compliance. Our
Wear log questionnaire provided options for retrospective logging of non-wear
events as well as indicating where the device was located in case no black bag was
used. Although these are scenarios impacting data quality, they are inevitable in
ambulatory studies, and it is recommended to account for them and systematically
label these instances rather than disregarding their occurrence (Van Der Donckt et
al., 2024). This proactive and informed approach allows researchers to make better
choices of how to handle this data in the analyses. Furthermore, providing
participants with the rationale behind performing specific non-wear behaviour was
important for compliance. Specifically, we explained that the use of the black bag

would lead to low light levels, and that this would help our analyses. Maintaining
contact with the participants during the week, monitoring their compliance, and
reinstructing them when needed were essential to promote engagement throughout
the study. This included continuous inspection of wear behaviour and interference
when a Wear log entry appeared inconsistent with the previous entry. We
acknowledge that this continuous monitoring can increase experimenter burden and
does not scale well to studies with many participants. We recommend providing
study participants with non-wear instruction sheets that can be directly consulted by
participants in case of doubts on how to log a wear behaviour and even unexpected
behaviour of the light logger. While a comprehensive review of challenges and
solutions to promote accuracy of non-wear data collection in ambulatory light logging
field studies is out of the scope of this paper, we refer the reader to related work in
the field of actimetry, where relevant strategies to mitigate data entry challenges and
ensure participant compliance have been examined (Balbim et al., 2021; Sriram et
al., 2009; Van Der Donckt et al., 2024).

Limitations

We acknowledge several limitations of the current study. First, we consider the self-
reported information from the Wear log as ground truth. We acknowledge that self-
reports may not be reliable, and they do not represent an optimal ground truth of
when the light glasses were worn or not. While we tried our best to limit
inconsistencies in the self-reported data by performing continuous data monitoring of
Wear log entries and systematic pre-processing of this data, we ultimately cannot
determine whether these entries represent participants' actual non-wear behaviour.
Second, our collection approach and our pre-processing pipeline have to be
interpreted in light of our study sample, which consisted of healthy, compliant, and
engaged participants. The high compliance of our participants also shows in the low
number of Wear log entries that had to be cleaned (2.76% of all Wear log entries,
due to for example open intervals, incorrect timestamps and duplicate entries). In
large scale studies or clinical populations, continuously monitoring data and asking
participants to report non-wear time as performed in the current study is not possible,
and number of incorrect entries and open Wear log intervals would be higher than
the one encountered in this study. Therefore, more extensive pre-processing

pipelines might be required to ensure data quality and accurate detection of non-
wear in larger studies and in studies with a low-compliance sample. As a
consequence, we can consider the results of this study as the best possible
outcome. Third, feasibility and accuracy of non-wear collection, detection, and
handling strategies depend on specific characteristics of the light logger used in
ambulatory studies. For example, for wrist-worn light loggers, skin temperature and
conductance can be used as inputs to automated non-wear detection, and existing
algorithms might be more useful than the approach taken here (Van Der Donckt et
al., 2024). On the other hand, for chest and eye-level light loggers, different inputs
are required for automated non-wear detection, and the strategies outlined in the
current paper might be useful to further develop suitable algorithms for each light
logger form factor. Fourth, our analysis on the comparison of light exposure metrics
across the raw and non-wear corrected datasets is limited by our relatively small
sample size, and in that differences are examined considering only one metric for
each participant across the week, rather than looking at differences in daily light
exposure metrics values. Lastly, the current study had a high temporal resolution
with a measurement frequency of 10 seconds. Smoothing or re-sampling of light
data might affect the performance of automated non-wear intervals detection,
particularly for those of short duration. While more research is required on how to
best pre-process light data and whether smoothing techniques affect non-wear
detection, researchers should provide detailed information on how they apply
smoothing transformations prior to non-wear detection on their datasets.

Conclusion

The emergence of wearable light loggers in recent years has advanced the study of
light exposure in free-living conditions. However, there remains a need of
standardisation in pre-processing the large time series data produced by these
devices, including the detection of non-wear intervals. In this study, we
systematically collected non-wear data using multiple independent sources and
continuously monitored participant's self-reported non-wear behaviour. This
approach provided a highly accurate non-wear ground truth, enabling a robust
assessment of data-driven detection methods. Our findings demonstrate that, in a

high compliance cohort, using a black bag for non-wear collection combined with an
algorithm to detect resulting clusters of low illuminance is a promising strategy to
manage non-wear time. Additionally, individual-level analysis of algorithm
performance for low activity and low illuminance clusters suggests that combining
activity and illuminance data could further improve non-wear detection. We believe
that the methods and insights presented in the current study will inform the
development of more advanced automated non-wear detection, including machine
learning approaches. To support this effort, we are making our dataset available and
hope to promote continued discussion to establish best practices for data pre-
processing in ambulatory light exposure research.

Declarations

Funding

This research was supported by funds of the Max Planck Research Group Translational Sensory and Circadian Neuroscience.

Conflicts of interest

The authors have no relevant financial or non-financial interests to disclose.

Ethics approval

This study was performed in line with the principles of the Declaration of Helsinki. Approval was granted by the Ethics Committee of the Technical University of Munich (TUM Ethics Committee 2023-115-S-KK).

Consent to participate

Participants signed an informed consent prior to participation in this study.

Consent for publication

Participants consented to publication of the data from this study by signing an informed consent.

Availability of code, data and materials

All code and data analysed in this study are publicly available on GitHub (https://github.com/tscnlab/GuidolinEtAl_BehavResMethods_2025) under the MIT Licence (code) and CC-BY Licence (data).

Authors' contributions

Conceptualisation: CG, MS

Data curation: CG

Formal analysis: CG

Funding acquisition: MS

Methodology: CG, JZ, SLH

Project administration: CG

Software: CG, SLH

Resources: -

Supervision: MS

Validation: -

Visualisation: CG

Writing – original draft: CG

Writing – review & editing: CG, JZ, SLH, MS

925 Open practice statement

926 Data and analysis code are available on GitHub at:
927 https://github.com/tscnlab/GuidolinEtAl_BehavResMethods_2025. This study was
928 not pre-registered.

929 Declaration of AI-assisted technologies in the writing process

930 During the preparation of this work, the authors used ChatGPT to assist in
931 reformulating sentences and improving the clarity of the manuscript. All content was
932 subsequently reviewed and revised by the authors, who take full responsibility for the
933 final version of the publication.

934 Acknowledgments

935 We thank an anonymous reviewer for their suggestion to include equivalence testing
936 as part of our analytic strategy.

References

Alboukadel, K. (2024). *Rstatix: Pipe-Friendly Framework for Basic Statistical Tests. R package version 0.7.2, https://rpkgs.datanovia.com/rstatix/.* https://rpkgs.datanovia.com/rstatix/

Balajadia, E., Garcia, S., Stampfli, J., Schrader, B., Guidolin, C., & Spitschan, M. (2023). Usability and Acceptability of a Corneal-Plane α-Opic Light Logger in a 24-h Field Trial. *Digital Biomarkers*, *7*(1), 139–149. https://doi.org/10.1159/000531404

Balbim, G. M., Marques, I. G., Marquez, D. X., Patel, D., Sharp, L. K., Kitsiou, S., & Nyenhuis, S. M. (2021). Using Fitbit as an mHealth Intervention Tool to Promote Physical Activity: Potential Challenges and Solutions. *JMIR mHealth and uHealth*, *9*(3), e25289. https://doi.org/10.2196/25289

Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B (Methodological)*, *57*(1), 289–300. https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

Biller, A. M., Balakrishnan, P., & Spitschan, M. (2024). *Behavioural determinants of physiologically-relevant light exposure.* *2*(1), 114. https://doi.org/10.1038/s44271-024-00159-5

Biller, A. M., Zauner, J., Cajochen, C., Gerle, M. A., Kalavally, V., Mohamed, A., Rottländer, L., Ming-Yi, S., Stefani, O., & Spitschan, M. (2024). *Physiologically-relevant light exposure and light behaviour in Switzerland and Malaysia.* https://doi.org/10.1101/2025.01.07.631760

Brown, T. M., Brainard, G. C., Cajochen, C., Czeisler, C. A., Hanifin, J. P., Lockley, S. W., Lucas, R. J., Münch, M., O'Hagan, J. B., Peirson, S. N., Price, L. L. A.,

Roenneberg, T., Schlangen, L. J. M., Skene, D. J., Spitschan, M., Vetter, C.,
Zee, P. C., & Jr, K. P. W. (2022). Recommendations for daytime, evening,
and nighttime indoor light exposure to best support physiology, sleep, and
wakefulness in healthy adults. *PLOS Biology*, *20*(3), e3001571.
https://doi.org/10.1371/journal.pbio.3001571

Burns, A. C., Windred, D. P., Rutter, M. K., Olivier, P., Vetter, C., Saxena, R., Lane,
J. M., Phillips, A. J. K., & Cain, S. W. (2023). Day and night light exposure are
associated with psychiatric disorders: An objective light study in >85,000
people. *Nature Mental Health*, *1*(11), Article 11.
https://doi.org/10.1038/s44220-023-00135-8

Choi, L., Ward, S. C., Schnelle, J. F., & Buchowski, M. S. (2012). Assessment of
Wear/Nonwear Time Classification Algorithms for Triaxial Accelerometer.
Medicine & Science in Sports & Exercise, *44*(10), 2009.
https://doi.org/10.1249/MSS.0b013e318258cb36

Cohen, J. (2013). *Statistical Power Analysis for the Behavioral Sciences* (2nd ed.).
Routledge. https://doi.org/10.4324/9780203771587

Cook, J., & Ramadas, V. (2020). When to consult precision-recall curves. *The Stata
Journal*, *20*(1), 131–148. https://doi.org/10.1177/1536867X20909693

Daugaard, S., Garde, A. H., Bonde, J. P. E., Christoffersen, J., Hansen, Ä. M.,
Markvart, J., Schlünssen, V., Skene, D. J., Vistisen, H. T., & Kolstad, H. A.
(2017). Night work, light exposure and melatonin on work days and days off.
Chronobiology International, *34*(7), 942–955.
https://doi.org/10.1080/07420528.2017.1327867

Didikoglu, A., Mohammadian, N., Johnson, S., van Tongeren, M., Wright, P.,
Casson, A. J., Brown, T. M., & Lucas, R. J. (2023). Associations between light

exposure and sleep timing and sleepiness while awake in a sample of UK

adults in everyday life. *Proceedings of the National Academy of Sciences*,

120(42), e2301608120. https://doi.org/10.1073/pnas.2301608120

Guidolin, C., Aerts, S., Agbeshie, G. K., Akuffo, K. O., Aydin, S. N., Baeza-Moyano,

D., Bolte, J., Broszio, K., Cantarero-García, G., Didikoglu, A., González-

Lezcano, R. A., Joosten-Ma, H., Melero-Tur, S., Tengelin, M. N., Pérez

Gutiérrez, M. C., Stefani, O., Svensson, I., Udovičić, L., Zauner, J., &

Spitschan, M. (2024). Protocol for a prospective, multicentre, cross-sectional

cohort study to assess personal light exposure. *BMC Public Health*, *24*(1),

3285. https://doi.org/10.1186/s12889-024-20206-4

Hammad, G., Wulff, K., Skene, D. J., Münch, M., & Spitschan, M. (2024). Open-

Source Python Module for the Analysis of Personalized Light Exposure Data

from Wearable Light Loggers and Dosimeters. *LEUKOS*, *20*(4), 380–389.

https://doi.org/10.1080/15502724.2023.2296863

Harris, P. A., Swafford, J., Serdoz, E. S., Eidenmuller, J., Delacqua, G., Jagtap, V.,

Taylor, R. J., Gelbard, A., Cheng, A. C., & Duda, S. N. (2022). MyCap: A

flexible and configurable platform for mobilizing the participant voice. *JAMIA*

Open, *5*(2), ooac047. https://doi.org/10.1093/jamiaopen/ooac047

Harris, P. A., Taylor, R., Thielke, R., Payne, J., Gonzalez, N., & Conde, J. G. (2009).

Research electronic data capture (REDCap)—A metadata-driven

methodology and workflow process for providing translational research

informatics support. *Journal of Biomedical Informatics*, *42*(2), 377–381.

https://doi.org/10.1016/j.jbi.2008.08.010

Hartmeyer, S., & Andersen, M. (2023). Towards a framework for light-dosimetry

studies: Quantification metrics. *Lighting Research & Technology*,

14771535231170500. https://doi.org/10.1177/14771535231170500

Hartmeyer, S., Baldursdottir, B., Valdimarsdottir, H., Agustsson, G., Gudjonsson, I.,

& Andersen, M. (2024). Identification of spectral distributions in light-dosimetry

data: Methodology and application to an intervention field study. *Lighting*

Research & Technology, 14771535241265905.

https://doi.org/10.1177/14771535241265905

Hartmeyer, S., Webler, F., & Andersen, M. (2022). Towards a framework for light-

dosimetry studies: Methodological considerations. *Lighting Research &*

Technology, 14771535221103258.

https://doi.org/10.1177/14771535221103258

Mead, M. P., Reid, K. J., & Knutson, K. L. (2023). Night-to-night associations

between light exposure and sleep health. *Journal of Sleep Research*, *32*(2),

e13620. https://doi.org/10.1111/jsr.13620

Muralidharan, A. R., Lança, C., Biswas, S., Barathi, V. A., Wan Yu Shermaine, L.,

Seang-Mei, S., Milea, D., & Najjar, R. P. (2021). Light and myopia: From

epidemiological studies to neurobiological mechanisms. *Therapeutic*

Advances in Ophthalmology, *13*, 25158414211059246.

https://doi.org/10.1177/25158414211059246

Peeters, S. T., Smolders, K. C. H. J., & de Kort, Y. A. W. (2020). What you set is

(not) what you get: How a light intervention in the field translates to personal

light exposure. *Building and Environment*, *185*, 107288.

https://doi.org/10.1016/j.buildenv.2020.107288

Pilz, L. K., de Oliveira, M. A. B., Steibel, E. G., Policarpo, L. M., Carissimi, A.,

Carvalho, F. G., Constantino, D. B., Tonon, A. C., Xavier, N. B., da Rosa

Righi, R., & Hidalgo, M. P. (2022). Development and testing of methods for

detecting off-wrist in actimetry recordings. *Sleep*, *45*(8), zsac118.

https://doi.org/10.1093/sleep/zsac118

Price, L. L. A., Udovičić, L., & Khazova, M. (2019). Circadian light exposure of shift

working nurses. *Proceedings of the 29th Quadrennial Session of the CIE,*

Washington, DC, USA, 838–845.

R Core Team. (2025). *R Core Team (2025). _R: A Language and Environment for*

Statistical Computing_. R Foundation for Statistical Computing,   Vienna,

Austria. [Computer software]. <https://www.R-project.org/>.

Reid, K. J., Santostasi, G., Baron, K. G., Wilson, J., Kang, J., & Zee, P. C. (2014).

Timing and intensity of light correlate with body weight in adults. *PloS One*,

9(4), e92251. https://doi.org/10.1371/journal.pone.0092251

Sit, D. K., McGowan, J., Wiltrout, C., Diler, R. S., Dills, J. (Jesse), Luther, J., Yang,

A., Ciolino, J. D., Seltman, H., Wisniewski, S. R., Terman, M., & Wisner, K. L.

(2018). Adjunctive Bright Light Therapy for Bipolar Depression: A

Randomized Double-Blind Placebo-Controlled Trial. *American Journal of*

Psychiatry, *175*(2), 131–139. https://doi.org/10.1176/appi.ajp.2017.16101200

Spitschan, M., Smolders, K., Vandendriessche, B., Bent, B., Bakker, J. P.,

Rodriguez-Chavez, I. R., & Vetter, C. (2022). Verification, analytical validation

and clinical validation (V3) of wearable dosimeters and light loggers. *DIGITAL*

HEALTH,   *8*,   20552076221144858.

https://doi.org/10.1177/20552076221144858

Sriram, J., Shin, M., Kotz, D., Rajan, A., Sastry, M., & Yarvis, M. (2009). Challenges in Data Quality Assurance in Pervasive Health Monitoring Systems. In D. Gawrock, H. Reimer, A.-R. Sadeghi, & C. Vishik (Eds.), *Future of Trust in Computing: Proceedings of the First International Conference Future of Trust in Computing 2008* (pp. 129–142). Vieweg+Teubner. https://doi.org/10.1007/978-3-8348-9324-6_14

Stefani, O., Marek, R., Schwarz, J., Plate, S., Zauner, J., & Schrader, B. (2024). Wearable Light Loggers in Field Conditions: Corneal Light Characteristics, User Compliance, and Acceptance. *Clocks & Sleep*, *6*(4), Article 4. https://doi.org/10.3390/clockssleep6040042

Syed, S., Morseth, B., Hopstock, L. A., & Horsch, A. (2021). A novel algorithm to detect non-wear time from raw accelerometer data using deep convolutional neural networks. *Scientific Reports*, *11*(1), 8832. https://doi.org/10.1038/s41598-021-87757-z

Van Der Donckt, J., Vandenbussche, N., Van Der Donckt, J., Chen, S., Stojchevska, M., De Brouwer, M., Steenwinckel, B., Paemeleire, K., Ongenae, F., & Van Hoecke, S. (2024). Mitigating data quality challenges in ambulatory wrist-worn wearable monitoring through analytical and practical approaches. *Scientific Reports*, *14*(1), 17545. https://doi.org/10.1038/s41598-024-67767-3

Van der Maren, S., Moderie, C., Duclos, C., Paquet, J., Daneault, V., & Dumont, M. (2018). Daily Profiles of Light Exposure and Evening Use of Light-emitting Devices in Young Adults Complaining of a Delayed Sleep Schedule. *Journal of Biological Rhythms*, *33*(2), 192–202. https://doi.org/10.1177/0748730418757007

van Duijnhoven, J., Hartmeyer, S., Didikoğlu, A., Stefani, O., Houser, K., Kalavally, V., & Spitschan, M. (2024). *Overview of wearable light loggers: State-of-the-art and outlook*. https://www.authorea.com/users/785068/articles/958417-overview-of-wearable-light-loggers-state-of-the-art-and-outlook?commit=06505e6a52132e1fa8098b3ca400acac7c53b83c

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L. D., François, R., Grolemund, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T. L., Miller, E., Bache, S. M., Müller, K., Ooms, J., Robinson, D., Seidel, D. P., Spinu, V., … Yutani, H. (2019). Welcome to the Tidyverse. *Journal of Open Source Software*, *4*(43), 1686. https://doi.org/10.21105/joss.01686

Windred, D. P., Burns, A. C., Lane, J. M., Olivier, P., Rutter, M. K., Saxena, R., Phillips, A. J. K., & Cain, S. W. (2024). Brighter nights and darker days predict higher mortality risk: A prospective analysis of personal light exposure in >88,000 individuals. *Proceedings of the National Academy of Sciences*, *121*(43), e2405924121. https://doi.org/10.1073/pnas.2405924121

Windred, D. P., Burns, A. C., Rutter, M. K., Yeung, C. H. C., Lane, J. M., Xiao, Q., Saxena, R., Cain, S. W., & Phillips, A. J. K. (2024). Personal light exposure patterns and incidence of type 2 diabetes: Analysis of 13 million hours of light sensor data and 670,000 person-years of prospective observation. *The Lancet Regional Health – Europe*, *42*. https://doi.org/10.1016/j.lanepe.2024.100943

Zauner, J., Guidolin, C., & Spitschan, M. (2025). How to Deal With Darkness: Modeling and Visualization of Zero-Inflated Personal Light Exposure Data on a Logarithmic Scale. *Journal of Biological Rhythms*, 07487304251336624. https://doi.org/10.1177/07487304251336624

Zauner, J., Hartmeyer, S., & Spitschan, M. (2025). LightLogR: Reproducible analysis

of personal light exposure data. *Journal of Open Source Software*, *10*(107),

7601. https://doi.org/10.21105/joss.07601

Zauner, J., Udovicic, L., & Spitschan, M. (2024). Power analysis for personal light

exposure measurements and interventions. *PLOS ONE*, *19*(12), e0308768.

https://doi.org/10.1371/journal.pone.0308768

1113

1114 **Figures**

1115 **Figure 1**

1116 Light glasses worn by participants throughout the study

1117

1118 *Note.* The light glasses comprised non-prescription glasses and the ActLumus light logger mounted

1119 centrally on the bridge of the glasses frame. (1) Button (event marker) on the light logger.

1120 **Figure 2**

1121 Schematic diagram of the non-wear intervals collection strategy

1122

1123 *Note.* Three independent methods of non-wear intervals were collected, leading to three independent

1124 source of non-wear information to base our analytic strategy on. This image was partially created

1125 using *BioRender.com.*

1126 **Figure 3**

1127 Flowchart for continuous quality assurance of Wear log entries performed throughout the study

1128

1129 *Note.* Rectangles indicate processes, rhombuses are used for yes/no branching, and rectangles with

1130 smoothed corners are used for descriptions.

1131 **Figure 4**

1132 Exemplary light exposure and wear status for a participant (n=1) across the study.

1133
1134 *Note.* The continuous black line shows mEDI as measured by the light glasses. Red points represent

1135 button presses performed by participants in case of any (non-)wear behaviour. Shaded areas

1136 represent self-reported wear status (Wear log).

1137 **Figure 5**

1138 Distribution of self-reported wear status throughout the week and the day

1139

Note. (A) Percentage of time spent in each Wear log state relative to individual study duration, and

equivalent number of hours. Study duration slightly differed for each participant, as they all started the

study at different times on the first Monday. (B) Percentage of time spent not wearing the light glasses

(Wake 'off' state) relative to total weekday and weekend day time. (C) Frequency of non-wear

episodes (Y-axis) over timing of non-wear episodes (X-axis). Timing of non-wear episodes (X-axis)

was calculated as centre point of each Wake 'off' interval.

Figure 6

Classification of non-wear intervals based on presence of button press at the start and end of the

interval

1149

1150   *Note.* Each Wear log Wake ⬚off⬚ interval (n=198) was classified as open or closed based on the

1151   presence of a button press within a specified time window. Note that button presses at both the start

1152   and end of an interval were needed for an interval to qualify as closed.

1155

Note. (A) Median values of log activity and illuminance throughout the week based on Wear log state. Every data point indicates the weekly median value for a participant (n=26 points for each Wear log state). Note that for this visualisation, a value of one was added to all mEDI and PIM raw values prior to transformation to logarithmic scale. (B) Precision recall curve for detection of non-wear intervals based on clusters low illuminance (red-orange colour scale) and low activity (blue-light blue colour scale), as detected via our algorithm. A minimum cluster length of 21 minutes and a maximum interruption length of 0 minutes were used as input parameters in our algorithm for illuminance. For activity, a minimum cluster length of 12 minutes and a maximum interruption length of 0 minutes were used. Each data point on the curve represents a maximum threshold (mEDI: 1 lux to 10 lux, at 1-step increments; PIM: 5 to 50, at 5-step increments).

1166 **Figure 8**

1167 Individual precision-recall curves for detection of non-wear intervals based on clusters of low

1168 illuminance and low activity, as calculated by the algorithm

1169

Note. A minimum cluster length of 21 minutes and a maximum interruption length of 0 minutes were

used as input parameters in our algorithm for illuminance (red-orange colour scale). For activity, a

minimum cluster length of 12 minutes and a maximum interruption length of 0 minutes were used

(blue-light blue colour scale). Each panel represents a participant, and each data point within a panel

represents a maximum threshold (mEDI: 1 lux to 10 lux, at 1-step increments; PIM: 5 to 50, at 5-step

increments). The percentage value displayed on the top left of each panel shows the frequency of bag

use relative to the participant's self-reported non-wear episodes.

1179

Note. Blue and green shaded areas represent correctly classified intervals (true negative and true positive, respectively). Yellow and red areas represent misclassified intervals (false negative and false positive, respectively). Grey shaded areas represent sleep episodes (excluded from the algorithm classification). The colour of the line representing light levels (mEDI) mirrors the wear status according to the ground truth (Wear log). The black symbols indicate examples of misclassified

1185   intervals as labelled in Table 1. Arrow: example of "Misclassified: Transition state"; Open circle:

1186   example of "Misclassified: Algorithm limitation or bag not actually used" instance; Asterisk:

1187   "Misclassified: Bag not used" instance; Star: "Misclassified: Low illuminance during wear" instance.

1188

Figure 10

Comparison of 14 light exposure metrics (A-M) between the raw dataset and the clean (Wear log)

dataset and between the raw dataset and the clean (algorithm) dataset.

Note. For each metric, the first scatter plot shows the relationship between a given metric in the raw
dataset and in the clean (Wear log) dataset (A1-O1), and a second scatterplot shows the relationship
between a given metric in the raw dataset and in the clean (algorithm) dataset (A2-O2). Each data
point in the scatterplot represents the value of the metric for a given participant (n=26). Distribution of
the delta values for the clean (Wear log) dataset and clean (algorithm) dataset relative of the raw
dataset are shown in A3-M3. Note that for timing metrics (C3-J3; $MLiT^{250}$, $MLiT^{1000}$, $FLiT^{10}$, $FLiT^{250}$,
$FLIT^{1000}$, $LLiT^{10}$, $LLiT^{250}$, $LLiT^{1000}$), the label HH:MM on the X-axis of the histogram of the deltas
represents local time, whereas for duration metrics (K3-M3; TAT^{250} and TAT^{1000}), the HH:MM
represents a duration. For the intensity metrics (average mEDI and M10m), the values shown in the
graphs represent mEDI lux (N1-O1 and N2-O2).

Tables

Table 1

Summary of correctly classified and misclassified (non-)wear instances by the algorithm detecting low

illuminance clusters

Algorithm performance		
Instance	N	N (%)
Misclassified: Algorithm limitation or bag not actually used	57	5.8282
Misclassified: Bag not used	59	6.0327
Misclassified: Low illuminance during wear	105	10.7362
Misclassified: Transition state	210	21.4724
Correctly classified	547	55.9305
Total	978	100.0000

1208

1209 *Note.* Correctly classified instances represent true negative and true positive instances, and

1210 misclassified instances represent false positive and false negative instances. Potential reasons for

1211 misclassified instances were deducted from the duration of the misclassified instance, mean mEDI

1212 during this interval, and self-reported use of the black bag during this instance.

1213

1214 **Table 2**

1215 Post-hoc Wilcoxon signed rank test comparisons results and corresponding equivalence test results

Post-hoc Wilcoxon pairwise comparisons and equivalence test results

Smallest effect size of interest for equivalence testing = 0.3

Metric	Group 1	Group 2	N1	N2	Statistic	p	Adjusted p	Significance	Cohen's d	Lower CI	Upper CI	Equivalence test result	Directionality of effect
$LLiT_{10}$	Raw data	Wear log-corrected	26	26	21	0.0360	0.0890	ns	0.2285	0.2012	0.4393	Inconclusive	/
$LLiT_{10}$	Raw data	Algorithm-corrected	26	26	15	0.0590	0.0890	ns	0.2142	0.1977	0.4193	Inconclusive	/
$LLiT_{10}$	Wear log-corrected	Algorithm-corrected	26	26	0	1.0000	1.0000	ns	-0.1961	-0.3541	-0.1961	Inconclusive	/
$LLiT_{1000}$	Raw data	Wear log-corrected	26	26	15	0.0590	0.1500	ns	0.2812	0.2022	0.4564	Inconclusive	/
$LLiT_{1000}$	Raw data	Algorithm-corrected	26	26	1	1.0000	1.0000	ns	0.1961	0.1961	0.3541	Inconclusive	/
$LLiT_{1000}$	Wear log-corrected	Algorithm-corrected	26	26	0	0.1000	0.1500	ns	-0.2804	-0.4564	-0.1961	Inconclusive	/
TAT_{1000}	Raw data	Wear log-corrected	26	26	120	0.0007	0.0020	**	0.5294	0.3945	0.7089	Positive effect	Raw data > Wear log-corrected
TAT_{1000}	Raw data	Algorithm-corrected	26	26	15	0.0570	0.0570	ns	0.4297	0.2511	0.6210	Inconclusive	/
TAT_{1000}	Wear log-corrected	Algorithm-corrected	26	26	0	0.0030	0.0040	**	-0.5248	-0.7049	-0.3896	Negative effect	Wear log-corrected < Algorithm-corrected
TAT_{250}	Raw data	Wear log-corrected	26	26	171	0.0002	0.0006	***	0.5014	0.3797	0.6791	Positive effect	Raw data > Wear log-corrected
TAT_{250}	Raw data	Algorithm-corrected	26	26	66	0.0040	0.0040	**	0.7205	0.5278	0.9786	Positive effect	Raw data > Algorithm-corrected
TAT_{250}	Wear log-corrected	Algorithm-corrected	26	26	0	0.0010	0.0020	**	-0.4943	-0.6728	-0.3667	Negative effect	Wear log-corrected < Algorithm-corrected
Average daytime mEDI	Raw data	Wear log-corrected	26	26	1	0.0000	0.0000	****	-0.6871	-0.9096	-0.5603	Negative effect	Raw data < Wear log-corrected
Average daytime mEDI	Raw data	Algorithm-corrected	26	26	0	0.0000	0.0000	****	-0.7178	-0.9495	-0.5847	Negative effect	Raw data < Algorithm-corrected
Average daytime mEDI	Wear log-corrected	Algorithm-corrected	26	26	156	0.8720	0.8720	ns	0.1276	-0.2885	0.3565	Inconclusive	/
M10m	Raw data	Wear log-corrected	26	26	0	0.0000	0.0000	****	-0.8381	-1.1174	-0.6883	Negative effect	Raw data < Wear log-corrected
M10m	Raw data	Algorithm-corrected	26	26	0	0.0000	0.0000	****	-0.8472	-1.0649	-0.7029	Negative effect	Raw data < Algorithm-corrected
M10m	Wear log-corrected	Algorithm-corrected	26	26	197	0.1840	0.1840	ns	0.2388	-0.0633	0.4838	Inconclusive	/
IV	Raw data	Wear log-corrected	26	26	137	0.3400	0.6530	ns	-0.1749	-0.3600	0.2596	Inconclusive	/
IV	Raw data	Algorithm-corrected	26	26	144	0.4370	0.6530	ns	-0.1518	-0.3370	0.3964	Inconclusive	/
IV	Wear log-corrected	Algorithm-corrected	26	26	194	0.6530	0.6530	ns	0.0907	-0.2448	0.4072	Inconclusive	/

1216
1217

Note. Post-hoc Wilcoxon signed rank tests were performed for all metrics where the Friedman test
returned a significant difference. P-value adjustment was performed using the Benjamini-Hochberg
method for multiple comparison correction. Significant values are reported at threshold p<0.05.
Equivalence tests were also run on all post-hoc pairwise comparisons using a smallest effect size of
interest (SESOI) of *d* = 0.3. For comparisons yielding to positive or negative effects in equivalence
testing (i.e. both 90% confidence intervals entirely above or entirely below the SESOI), the
directionality indicates which dataset had higher or lower values.

1225

1226   **Table 3**

1227   Summary of positive, negative, and inconclusive effects identified via equivalence testing

		Difference from no correction	
		Inconclusive	**Yes**
Difference between correction methods	**Inconclusive**	$LLiT_{10}$ $LLiT_{1000}$ IV IS	Average daytime mEDI M10m
	Yes	/	TAT_{1000} TAT_{250}

Uncorrected metric value is lower
Uncorrected metric value is higher

1228   *Note.* Summary of light exposure metrics that yielded statistically meaningful positive, negative or
1229   inconclusive effects in pairwise comparisons performed in equivalence testing (SESOI = ±0.3). The
1230   metrics are categorised by differences from the raw dataset (i.e. no correction; table columns) and
1231   differences between correction methods (i.e. Wear log-corrected and algorithm-corrected; table rows).