Title: Auditory brainstem responses to speech-in-noise reflect selective attention,
comprehension, and subjective listening effort

Abbreviated title: Attention shapes brainstem responses to speech

Authors: Mankel, K.[1,2,3], Comstock, D. C.[3], Bormann, B. M.[3,4], Das, S.[3], Sagiv, D.[5], Brodie, H.[5],
Miller, L. M.[3,5,6]

Affiliations:
[1]*School of Communication Sciences & Disorders, University of Memphis, Memphis, TN, USA*
[2]*Institute for Intelligent Systems, University of Memphis, Memphis, TN, USA*
[3]*Center for Mind and Brain, University of California, Davis, Davis, CA, USA*
[4]*Neuroscience Graduate Group, University of California, Davis, Davis, CA, USA*
[5]*Department of Otolaryngology | Head and Neck Surgery, University of California, Davis, Davis,*
CA, USA
[6]*Department of Neurobiology, Physiology & Behavior, University of California, Davis, Davis, CA,*
USA

Corresponding author: Kelsey Mankel, kmankel@memphis.edu

Acknowledgements

This work was supported by the Office of the Assistant Secretary of Defense for Health Affairs
through the Congressionally Directed Medical Research Program (CDMRP) Hearing
Restoration Research Program (HRRP) under Award No. W81XWH-20-1-0485 (to LMM).
Opinions, interpretations, conclusions, and recommendations are those of the author and are
not necessarily endorsed by the Department of Defense. This work was also supported by the
Child Family Fund for the Center for Mind and Brain at the University of California, Davis (to
LMM). The authors would like to thank members of the University of California, Davis Health
audiology and clinical research team for performing initial hearing evaluations of our
participants: Dr. Robert Ivory, AuD; Dr. Mackenzie Quinn, AuD; Dr. Rachel Krager, AuD; Dr.
Steven Zurawski, AuD; Dr. Austin Childers, AuD; Dr. Kimberly Smith, AuD; Angela Beliveau, and
Randev Sandhu. We are also grateful for our team of undergraduate lab volunteers—
particularly Jillian McKie and Cathleen Chan—who assisted in countless hours of participant
recruitment, data collection, and data pre-processing. Thank you to Tiana Smith and Elyse
Ehlert for their efforts in participant recruitment.
AI was used in a limited capacity for this work, primarily assisting with minor text editing
and basic phrasing. All significant drafting, analysis, and editing were carried out by the authors.

Dedications

34 This research is dedicated in loving memory of our research team member, fellow lab mate and
35 friend, Karim Abou Najm.

Conflicts of interest

37 Lee M. Miller is an inventor on intellectual property related to chirped-speech (Cheech) owned
38 by the Regents of University of California, not presently licensed.

Abstract

The auditory brainstem plays a crucial role in speech-in-noise listening, refining numerous acoustic features such as pitch and spatial location under continual descending influence from cortex. However, the difficulty of characterizing brainstem activity during continuous speech listening has obscured its functional role in ecologically valid contexts—not only the effects of selective attention on neural responses, but also their impact on comprehension and listening effort. Here, we evaluate the role of the brainstem on speech-in-noise perception and selective attention using a continuous, speech-based stimulus with embedded chirps (Cheech) that rapidly and effectively evokes auditory brainstem responses (ABRs) while participants listen to short story narratives. The Cheech-modified stories were presented alone or in the presence of another spatially separated talker, and neural responses were measured throughout by EEG. Both word-level detection performance and narrative-level comprehension were evaluated, as well as subjective reports of listening effort. ABR wave V was modulated by the presence of a competing talker, selective attention for the target versus masker, and the talker gender. Additionally, faster wave V peak latencies and larger amplitudes were associated with identification of the target words embedded within the stories, while faster latencies additionally related to better comprehension question accuracy and lower subjective listening effort. Collectively, our results provide clear evidence for the influence of brainstem encoding processes on individual speech-listening behaviors, including the ability to selectively attend to and comprehend target speech in the presence of a competing talker.

Keywords: auditory brainstem response (ABR), chirped-speech (Cheech), speech-in-noise, EEG, auditory event-related potentials (ERPs)

Significance statement

This study highlights the crucial role of the brainstem in speech-in-noise perception and higher-level language abilities like comprehension and subjective listening effort. Through the use of a novel speech stimulus blended with embedded chirps (Cheech) to evoke brainstem responses while listening to short story narratives, we show how selective attention influences subcortical processes. Faster and more robust subcortical processes, in turn, were associated with more successful speech-listening performance across multiple behavioral metrics. These findings advance our understanding of how the brainstem supports selective attention and speech perception in challenging listening environments.

Introduction

Neural processing of speech has primarily been studied at the level of the cortex (e.g., Hickok and Poeppel, 2004). Yet, brainstem nuclei along the auditory pathway perform crucial functions relevant for processing speech such as encoding pitch and timing of spectrotemporal fluctuations (e.g., Krishnan and Gandour, 2009; Krishnan and Plack, 2011), preliminary phoneme categorization (Carter and Bidelman, 2023; Rizzi and Bidelman, 2023), and even early differentiation of competing auditory streams, likely through corticofugal modulations (Price and Bidelman, 2021). Additionally, poorer encoding of auditory brainstem responses (ABRs) has been associated with decreased speech-in-noise (SiN) performance (Papakonstantinou et al., 2011; Bramhall et al., 2015). Thus, rather than serving as a simple relay for acoustic information, the brainstem plays an active, perceptually consequential role in speech perception.

Yet, whether and how subcortical processes influence the ability to selectively attend to target sounds while filtering out background noise, as required for SiN perception, remains uncertain. In early studies, ABRs were considered immune from effects of attention and arousal states (e.g., Jewett and Williston, 1971; Connolly et al., 1989; Gregory et al., 1989). However, midbrain responses (ABR wave V) may be modulated by selective attention of audition over another modality (Lukas, 1980; Bauer and Bayles, 1990; Kumar et al., 2023), increased cognitive load (Sörqvist et al., 2012), or cognitive interference (Brännström et al., 2020). Mixed results on the influence of attention have also been demonstrated for other subcortical responses like the frequency-following response (FFR) (e.g., Hoormann et al., 2000; Galbraith et al., 2003; Varghese et al., 2015), possibly due to cortical contributions (Coffey et al., 2016; cf. Bidelman, 2018). Meanwhile, regression- or correlation-based brainstem responses derived from continuous speech are more robust for attended rather than unattended talkers (Forte et al., 2017; Etard et al., 2019), and attentional modulation of these responses relates to independent tests of SiN perception (Saiz-Alía and Reichenbach, 2020).

Historically conflicting results regarding attentional modulation of brainstem activity likely reflect differences among experimental approaches, chief among them a varying degree of ecological validity. While subcortical EEG responses are best evoked by transient artificial stimuli such as isolated clicks or tone bursts, these are not always good predictors of speech perception in real-world listening environments (Skoe and Kraus, 2010), nor do simple sounds adhere to naturalistic statistics known to influence midbrain encoding (Escabí et al., 2003). As a result, despite numerous studies of auditory brainstem function, its role in attentive listening and its relevance to real-world speech performance remain unclear. In order to shed light on the neural processes underlying SiN perception, a challenging, ecologically valid task using naturalistic speech with relevant outcome metrics should be used (Hamilton and Huth, 2020).

However, fundamental limitations characterize ABR recordings such as low signal-to-noise ratios (SNR), relatively small amplitudes, and the need for multiple stimulus repetitions. To address this, we have developed a technique for the rapid measurement of brainstem through cortical responses via "chirped-speech" (Cheech), continuous speech stimuli blended energetically and perceptually with frequency sweeps (Backer et al., 2019; Miller et al., 2020). Our technique parallels recent efforts to derive brainstem activity during continuous speech (Forte et al., 2017; Maddox and Lee, 2018; e.g., Etard et al., 2019; Polonenko and Maddox, 2021; Kulasingham et al., 2024). Compared to other methods, Cheech offers more acoustic control over the specific parameters used to elicit multiple, simultaneously recorded brain responses from naturalistic speech, all while listeners perform a challenging, ecologically valid task complete with spatial cues and attentional switching.

Here, we use Cheech-modified short stories presented both in quiet and in the presence
of a competing talker to elucidate attentional effects within the brainstem through ABRs (i.e.,
wave V). Our results suggest wave V is affected not only by the presence of another talker, but
also by selective attention to one talker over another. We also demonstrate the relationship
between ABR wave V and several higher-level speech listening behaviors, including
comprehension and subjective listening effort.
125
126
Methods

Participants

Twenty-nine participants were recruited for this study. To be eligible, prospective
participants must be between 20 and 40 years old, speak English as their first language, and
report no neurologic, psychiatric, or scalp conditions that directly affected their ability to
understand speech or interfere with electrophysiologic measurements (e.g., epilepsy, certain
strokes, ADHD, scalp problems, etc.). Participants were screened for visual acuity (i.e., $\leq$ 20/40
per Snellen chart), normal hearing (i.e., $\leq$25 dB HL air conduction thresholds between 250-8000
Hz in both ears, no air-bone gaps >10 dB within 500-2000 Hz, no asymmetries between the
ears >20 dB at 500, 1000, or 2000 Hz), and cognitive abilities (>24 Montreal Cognitive
Assessment [MoCA] score—consistent with an ongoing comparison study group of veterans;
Nasreddine et al., 2005). One participant withdrew from the study due to personal reasons; one
participant was excluded due to technical difficulties during the EEG recordings, and two
participants were excluded because they did not pass the hearing screener at the time of
testing. This resulted in a final cohort of N=25 total participants with the following demographics:
average age=22.88 ± 4.98 SD years old (range=18-38 years old); 15 females and 10 males;
N=21 self-identified handedness right, N=2 left, and N=2 ambidextrous. Participants had on
average 16.86 ± 3.19 SD years of education and 4.20 ± 3.74 SD years of formal music training.
Of those who reported some level of formal music training (N=22), the average age they began
music training was 10.05 ± 2.75 SD years old. Average MoCA score was 28.16 ± 1.46 SD
(range=25-30). Additionally, participants completed a version of the MacArthur Scale of
Subjective Socioeconomic Status where they were asked to rate their perceived socioeconomic
status (SES) relative to individuals across the US society using a ladder-rung rating from 1
(lowest SES)-10 (highest SES) (Adler et al., 2000). Mean subjective SES scores were 4.70 ±
1.30 SD, range 3-8. Average air conduction pure tone average (PTA) thresholds were 6.80 ±
3.72 SD and 7.04 ± 4.47 SD for the right and left ears, respectively, indicative of normal hearing.
All participants completed an informed consent as approved by the University of California,
Davis Institutional Review Board and were compensated financially for their participation.
155
Stimuli

The stimuli were short fairytale stories available in the public domain (e.g., on
www.librivox.com). The stories were selected for content unlikely to be widely known, to avoid
comprehension question recall based on modern renditions (i.e., popularized by children's
books or movies), and with story arc progress within a 7.5-minute block (i.e., rising action,
climax, or falling action) to ensure sufficient and generally consistent content for probing
comprehension. Thirteen stories were piloted on four raters who scored them according to
factors such as interest and engagement levels, ease of understanding, grammar or writing
style complexity, familiarity (i.e., low scores preferred), and suitability for an adult audience.

Seven stories that scored highest across these factors were ultimately retained for the
experiment.
The short stories were modified to add monosyllable color words as targets for the
spatial attention experiment (e.g., red, green, blue, white, etc.). The color words were
embedded as part of the story itself rather than unassociated, random target word interjections
(e.g., "…with a narrow **_blue_** ribbon over her shoulders"). We recorded a male talker (native
English speaker with a neutral American accent) reading the modified short stories aloud using
a Shure KSM244 vocal microphone (cardioid polar pattern, high-pass filter of 18 dB per octave
cutoff at 80 Hz) and Adobe Audition (sample rate = 48,000 Hz, 32 bit depth [float]) in a quiet,
soundproof room. Silent gaps in the recordings greater than 500 ms were shortened to 500 ms,
similar to (Broderick et al., 2018; Teoh and Lalor, 2019; Teoh et al., 2022), and then the trimmed
recordings were then cropped to 7.5 minutes per story audio.
177
<u>Chirped speech (Cheech)</u>

The short story auditory stimuli were then modified via a patented process called
"Cheech" ("chirped-speech") to evoke the auditory evoked potentials (Backer et al., 2019; Miller
et al., 2020; Figure 1). In Cheech, some of the glottal pulse energy within predefined frequency
bands are replaced with narrowband synthetic chirps, consistent with previously published
protocols from our lab (Backer et al., 2019). Aligning the chirps with glottal pulse energy inherent
in speech creates an acoustically-fused auditory perceptual object with sufficient transient chirp
activation to measure auditory brainstem through cortical responses while preserving
naturalistic speech qualities and linguistic content. First, periods of sufficient voiced power were
identified; specifically, the audio was filtered from 20 to 1000 Hz, and voiced periods at least 50
ms long were identified when the speech envelope between 20-40 Hz surpassed a threshold
~28% of overall speech root-mean-square (RMS) amplitudes. The timing of the glottal pulses
during voiced periods was determined by a speech resynthesis process using custom Matlab
code and the TANDEM-STRAIGHT toolbox that retains original speech frequency
characteristics (Kawahara et al., 1999, 2008; Kawahara and Morise, 2011). The continuous
speech was re-filtered into alternating, octave-wide frequency bands of 0-250, 500-1000, 2000-
4000, 11,000-∞ Hz, and chirp energy was constrained to frequency bands of 250-500, 1000-
2000, 4000-11,000 Hz. The chirp and speech bands in the alternating, interleaved octave-wide
bands were then added together. Each mono track was then duplicated to form stereo audio.
Audio files were normalized prior to the Cheech synthesis to ensure equivalent chirp amplitudes
relative to the speech levels across each story, and the root-mean-square (RMS) amplitudes
were re-checked across all Cheech-ed stories for equivalence after the Cheech process was
completed.
Timing of the chirps varied with the natural fluctuations in the running speech, with
special modifications to optimize measurement of the auditory ERPs. Specifically, when voicing
energy was detected, chirps were inserted at minimum 18.2 ms apart (55 Hz), skipping glottal
pulses that occur within the 18.2 ms window. The first chirp within a sequence is always
followed by a minimum of 48 ms before the next chirp is presented, skipping glottal pulses in
between, to produce a longer ISI that allows for improved evolution of middle latency responses
(MLR).
Within the dual-talker conditions, one story from each pair was pitch modulated using
MATLAB STRAIGHT prior to the Cheech process. This manipulation allowed us to compare
neural and behavioral responses for male versus female voices (between-story comparison) in
addition to attended versus unattended conditions (within-story comparison; see Experimental
Conditions section). Specifically, female-intended stories were modulated in pitch to an average

F0 of 180 Hz, while the original (male voice) audio had an average F0 of approximately 128 Hz,
and modulated in frequency to model a shorter vocal tract length. Modifying voice parameters in
this manner meant speaker characteristics such as phrasing, intonation, and pacing were
otherwise preserved because all stories were originally spoken by a single talker. Modifying one
talker's F0 also provided the added benefit of a combined voice-gender + spatial release from
masking which improved pilot test performance (Oh et al., 2021, 2022). The Cheech processing
details are summarized in Appendix B.

Figure 1: A brief segment of the chirped-speech (Cheech) stimuli. (Top) Participants listened to the Cheech-modified short stories and pressed a button when they heard the target talker speak a color word (e.g., "blue"). (Middle) Stimulus waveform and (bottom) spectrogram for the Cheech audio. Cheech interleaves speech and chirp frequency bands in a manner that preserves naturalistic speech qualities while evoking robust ABRs. See Methods for details.

Spatial filtering and attention switching task

The Cheech stimuli were spatially filtered via HRTFs to simulate audio at ±15° (SADIE II
database; Armstrong et al., 2018). Spatial audio was mirrored by two symbols on two computer
screens approximately 15° to the left and right of the midline in front of the participant. A "+"
symbol on one of the screens corresponded to a given target location ."<" and ">" symbols were
shown on the computer screen opposite from the target speech location (e.g., a ">" shown on
the left screen and a "+"on the right screen indicated a the target voice located in the right
auditory spatial field; see Figure 2). Consequently, "<" and ">" symbols indicated either the
absence of another speaker for a given spatial field, as in the mono-talker condition, or the
location of the masker story in the two-talker condition. The symbols were identical in size, color,
and line lengths to maintain equivalent luminance for the eye tracking measurements (data not
included here). The spatially filtered audio and visual symbols were then edited with custom
Matlab code to swap left and right spatial locations 75 times approximately every six seconds
(7.5 minutes ÷ 75 switches = 6 seconds; switches assigned per random distribution of 6 ± 1
seconds; see Figure 1). The visual icons switch screens instantaneously at each switch time
while the crossover fade time for the audio was 35 ms, beginning at the onset of a switch cue
(Figure 2). During piloting, this crossover time was an appropriate balance between minimizing
"pops" and other artifacts from cutting the audio across left and right audio tracks while also
minimizing the perception of a gradually moving sound between spatial locations (i.e., a sudden
jump from one side to the other without introducing robust distortions). In this manner, our
spatial attention switching task mimicked more ecologically valid listening scenarios, as if
perceptually following a conversation between two spatially separated talkers
Color words were randomly assigned to one of five possible onset categories between
0.125 and 2 seconds from the switch-to-color-word-onset (i.e., 0.125, 0.25, 0.5, 1, and 2
seconds, 5 color words per condition; see Figure 2). These categories were established after

preliminary piloting to assess whether "attentional blinks" from switching left to right spatial
attention following the cue (and vice versa) impaired color word detection across participants;
these data will be presented in a subsequent paper. Onset timings of the color words were
determined by Gentle, an audio forced aligner (https://github.com/lowerquality/gentle). The
closest preceding switch time was replaced by the assigned switch-to-color-word-onset,
resulting in 50 random switches and 25 switch-to-color-words (i.e., 33% probability of a target
color word onset within 2 seconds of a switch). The 25 words were balanced across left-to-right
and right-to-left switches within a story (minus one remainder).
254

Figure 2: Experimental paradigm. Participants are instructed to attend to the target speech (red line) and press a button when a color word is spoken ('X'). They must ignore color words spoken by the masker talker when present (i.e., dual talker condition; 'O'). The target visual cue and voice alternate between approximately +/- 15 º in the visual and auditory spatial fields on average around every 6 seconds (see Methods).

255
Experimental conditions

Following audio cropping to 7.5 minute length, the seven stories were then divided into
separate experimental conditions: two stories presented alone ("mono-talker"), two sets of
paired stories ("dual-talker"), and one for initial practice which was not included in subsequent
analyses. Within the dual-talker condition, each story was heard twice—once as the "target
story" and once as the "masker story" within the pair. This experimental manipulation thus
allowed us to measure effects of directed spatial attention because the same audio was
presented and participants were instructed to either "attend" or "ignore" the story, respectively.
Mono-talker stories were only presented once each. The target story always started on the left
side, whether presented alone (mono-talker) or in the presence of a masker (dual-talker). The

order of stories and conditions was counterbalanced with a Latin square design across
participants to reduce potential sequence-dependent learning effects. No story pair was heard
back-to-back to reduce short-term learning effects as well.
269   Story pairs in the dual-talker condition were subject to additional considerations. As
mentioned in the Cheech section, one voice within the story pair was pitch modulated to
simulate a female speaker. Consistent with combined voice-gender and spatial release of
masking outcomes observed in previous work (Oh et al., 2021, 2022), two distinct talker F0s
helped listeners better distinguish between the spatially-separated voices and more successfully
complete the task during piloting. The mono-talker stories were presented in the original,
unmodified male voice only. Given the constraint of naturalistic color word embeddings within
each story, color words occasionally overlapped within the dual-talker pairings. The story pairs
were determined according to those which had the fewest color word overlaps, and any
overlapping color words were dropped from the behavioral task analyses so behavioral
responses to target (hits) and masker color words (false alarms) would not be conflated. A few
target story switch times were also manually adjusted to ensure that no masker color word
occurred during a switch (i.e., masker color word onset was not between -0.5-0.1 ms from a
[target story-driven] switch time).
283   Additionally, half of the stories—one mono-talker story and a pair of dual-talker stories—
were subjected to additional modifications during the Cheech process. These modifications
were specifically intended to disrupt subcortical encoding mechanisms, so we do not include
analysis of those stories here. These data will be analyzed in separate reports. Rather, this
study focuses on the remaining 3 experimental stories (i.e., one mono-talker story and a pair of
dual-talker stories). For additional details on the short stories used in the current study, see
Appendix A.
290
291   <u>Stimulus presentation equipment</u>

292   Participants were seated in a soundproof, electromagnetically shielded booth (ETS
Lindgren), approximately 176 cm from two computer monitors. The Dell UltraSharp monitors
had a refresh rate of 60 Hz, and screen settings were left on standard factory defaults. Stimulus
presentation, including both audio and visual, was controlled by custom MATLAB code using
Psychtoolbox (Brainard, 1997; Pelli, 1997). Auditory stimuli were routed from a Hammerfall
audio card to an audio interface (RME Fireface UFX II), then to a headphone driver (RME ADI-
2DAC FS) which drove ER-2 insert earphones (Etymotic Research/Interacoustics). The ER-2
inserts were electromagnetically shielded to reduce EEG artifacts (Campbell et al., 2012).
Cheech stimuli were presented at 75 dB SPL; clicks for the separate ABR recordings were
presented at 80 dB SPL (Larson-Davis PRM902 preamp & 2221 power supply, B&K 4157
coupler, C-weighted average).
303
304   <u>Procedure</u>

305   In total, each participant completed three sessions as part of a larger experiment: one 1-
hour audiologic exam, one 2-hour behavioral session, and one 2.5-hour laboratory EEG
session. Several participants completed more than one session on the same day (i.e., either
audiologic + behavioral or behavioral + EEG sessions), but a long break was offered in between
sessions to reduce fatigue. The data presented here focuses on the EEG experimental session
only.

The order of events within each block is as follows: 1 minute of brief rest (for resting
state EEG analysis), 7.5 minutes of the spatial attention listening task, 5 comprehension
questions, 6 subjective task load questions (Hart and Staveland, 1988), a flash reaction time
task (5 short trials), and then a break. During the one-minute rest period, participants were
instructed to remain still with their eyes open. During the spatial attention listening task,
participants listened to the short stories, following the target story and corresponding visual cue
as they switched between left and right relative spatial locations (Figure 2). They were
instructed to press the spacebar when they heard the embedded color words within the story. In
dual-talker condition blocks, participants were told to ignore color words from the masker story
and only press the spacebar for target story color words. The audio was immediately followed
by a series of five multiple choice comprehension questions with four possible answer choice
options (chance = 25% accuracy). Comprehension questions were balanced for difficulty (not
too hard nor too easy; piloted for accuracy on 3-5 individuals per story), story time relevant for
the question (i.e., early vs. late in the 7.5-minute story segment), and memorization vs. inductive
reasoning probed by the answers. Next, participants completed the NASA Task Load Index, a
measure of subjective task-related workload or effort, modified for computerized data collection
using a sliding scale selection tool (Hart and Staveland, 1988). To measure general mental
fatigue across the blocks, we used a psychomotor vigilance-based visual reaction time task
(e.g., Basner & Dinges, 2011; Dinges & Powell, 1985; Hornsby, 2013; Wilkinson & Houghton,
1982). Participants were instructed to watch the crosshairs shown at the center of the computer
screen and press the spacebar as quickly as possible when a white circle appears on screen.
The circle flashed on screen for 40 ms across five trials. Finally, participants were offered a
break before starting the next block. The entire experiment presentation was controlled by
custom MATLAB and Psychtoolbox code.
The data presented here were collected as part of a larger study. As mentioned in the
Experimental Conditions section, participants completed three additional short story blocks with
Cheech audio designed to impair auditory brainstem encoding, so those blocks are not included
in the present analyses. Additionally, participants obtained a full audiologic evaluation and
completed a battery of cognitive tests in separate experimental sessions. These data will be
analyzed in separate reports. Many participants completed more than one session in a given
day but not all three on a single day, and they were given longer breaks in between sessions to
reduce fatigue.
343
<u>Behavioral outcome measures and analysis</u>

Selective attention listening was assessed by correct identification of color words
embedded in the target story (hits). A "hit" was defined as pressing a designated keyboard
button within 2 seconds after the onset of a target word. Any pairs of color words within +/-2
second onset across the target and masker stories in the dual talker condition were excluded
from analyses, resulting in four overlapped color words dropped from each story in the
behavioral task analyses for the pair of stories used here. Specifically, the relevant measures
were the proportion and corresponding reaction times of correct hits out of the total possible,
non-overlapping target words (n=25 in the mono talker condition, n=21 per story in the dual
talker condition). Behavioral outcomes measured after each story concluded were
comprehension question accuracy (i.e., total % correct out of five questions), task load effort
scores (i.e., average NASA task load index scores across six questions, where higher scores
means greater perceived task load; Hart and Staveland, 1988), and average reaction times on
the vigilance-based flash task in a given block.

358
<u>EEG recordings</u>

EEG recordings were obtained using a BioSemi ActiveTwo system (BioSemi B.V.,
Netherlands, www.biosemi.com) in an electrically shielded booth with a sample rate of 8192 Hz
recording data from 72 electrodes: 64 channels following the International 10-20 standard with 4
additional channel pairs around the ears (earlobe, mastoid, lower posterior: posterior to the
earlobe and inferior to the mastoid, and upper anterior: superior-anterior to the tragus near the
zygomatic arch). Electrode offsets from all channels were measured to be below 20 µV relative
to the Common Mode Sense electrode using ActiveView2 software. Data were recorded using
open-source Lab Streaming Layer (LSL) software (https://github.com/sccn/labstreaminglayer)
as XDF files on a desktop computer running Windows 10.
A pair of channel outputs containing trigger events was routed from the audio interface to
a pair of Brain Products StimTraks for monitoring triggers associated with the target and masker
stories, respectively, which then convert the audio trigger signals to trigger pulses sent to the
TriggerBox (Brain Products GmbH). This setup permitted audio playback synchronized with
chirp triggers embedded during the Cheech process with minimal latency delays (see Cheech
section); such near-simultaneous synchronicity is crucial for measuring rapid, lower-level
auditory brainstem responses during the listening task. Triggers controlled by Psychtoolbox (i.e.,
participant responses, block sequence triggers, spatial location switches, etc.) were also routed
to the TriggerBox via the computer parallel port. The TriggerBox then passed both triggers from
the parallel port and the StimTraks to the Biosemi signal receiver box.
379

<u>EEG Preprocessing</u>

EEG were preprocessed using MATLAB 2021b (MathWorks Inc.) using a combination of
EEGLAB functions (Delorme and Makeig, 2004), ERPLAB functions (Lopez-Calderon and Luck,
2014), and custom MATLAB code. Data were first imported into EEGLAB with events extracted
from the trigger channel in the XDF file using custom Matlab code. Given the large number of
tightly packed event codes for our experimental design (>130,000/participant at an average rate
of 1 event every 20ms), event codes from the 3 separate streams (parallel port, and 2
StimTraks, described above) occasionally arrived at the TriggerBox simultaneously, resulting in
correctable event code errors. Custom code was used to check the recorded events against the
expected events for a given condition to add missing events, remove extra events, and adjust
event latencies for any overlapped event. A second set of corrections were then applied to
adjust for latency differences between the acoustic signal and the event times. These
corrections took into account a 1ms delay due to audio travel time from the insert earphone
tubes, a 1.973ms delay to the audio signal due to the HRTF process, a 0.541ms delay to the
chirp triggers to account for a difference between the expected chirp timing from the Cheech
process and the chirp onsets, and a 1ms delay in one of the StimTrak channels caused by the
TriggerBox hardware.
Data were then pruned to retain only rest periods and attentive listening task periods
along with a 10 second buffer on both sides of each selected period. Following pruning, each
section of the data was DC corrected and a 2^{nd} order 0.1 Hz high-pass Butterworth filter was
applied to remove slow drifts. We then referenced the data to electrode Cz. Even though our
data were collected in an electromagnetically shielded booth, we occasionally found 60 Hz line
noise in some channels. To minimize the effects of cleaning line noise, only channels with

detected line noise were cleaned (mean = 4.4 channels ± 7.56 SD, range for a given participant
= 0-39 channels). This was carried out by first checking for line noise using a frequency tagging
approach. Spectra were calculated across all frequencies for each channel. For each frequency,
the average spectra of 4 neighboring frequencies except those immediately adjacent (x-2, x-3,
x+2, x+3) was subtracted from the spectra for each frequency. If the neighbor corrected power
at 60 Hz or any of its first four harmonics (120, 180, 240, 300 Hz) was greater than 8 standard
deviations calculated from the corrected power from remaining frequencies, that channel was
cleaned of line noise using the Cleanline plugin (https://www.nitrc.org/projects/cleanline). This
process was repeated until no further line noise was detected up to a maximum of 5 repetitions.
To clean the data of prominent artifacts such as eye blinks, eye movements, and
heartbeat signals, we used Independent Component Analysis (ICA) following the methodology
outlined by Luck (2022) which improves ICA performance, and greatly reduces the required
processing time, compared to more traditional approaches. This process involved making a
copy of the processed data (the ICA dataset), down-sampling and pre-cleaning it of prominent
(non-eye and heart) artifacts, applying ICA to select and remove eye and heart artifacts, then
transferring the calculated component weights back to the original dataset. Specifically, we first
down-sampled the data to 256 Hz to speed ICA processing, then we inspected and rejected bad
channels by eye (mean=3.72 ± 2.13 SD channels, range=0-9 channels). Following channel
rejection, we applied a non-causal 8th order Butterworth band pass filter with cutoffs at 1 and 50
Hz. Further cleaning was then undertaken using ERPLABs continuous artifact rejection function
to remove sections of data with significant movement and muscle amplitude artifacts, the
presence of which can hinder ICA's ability to isolate eye and heart components (mean=129.19 ±
86 s SD or 3.6% ± 2.35%, range=4.98-298.79 s or 0.14-7.96%). The rejection criterion operated
on a 1-s sliding window with a 0.25 s step size and an initial threshold of 200 µV. As we wanted
to retain eye activity for ICA, frontal channels that contain the largest eye related activations
(FP1, FPz, FP2, AF7, AF3, AFz, AF4, AF8) were excluded from the rejection threshold. We then
inspected and adjusted the threshold level for each dataset (mean=220 ± 28.87 SD µV,
range=200-300 µV) until only those sections of data with large muscle or movement artifacts
were rejected while any eye blink or eye movement related artifacts remained. To further speed
the ICA processing time, the number of input channels was reduced from an average of 67
channels to the first 32 principle components using Principle Component Analysis (PCA), then
ICA was performed using the infomax algorithm. Eye blink, eye movement, and (when present)
heart artifact components were manually marked for rejection (mean=2.8 ± 0.65 SD
components, range=2-4 components). Following ICA processing, we prepared the original
dataset for transferring component weights by first removing the same bad channels marked for
removal in the ICA dataset. We then transferred the calculated ICA weights from the ICA dataset
to the original dataset and rejected the marked artifact components in the original dataset. Thus,
any eye and heart artifact components were removed from the dataset with the original sample
rate (and without the aggressive pre-cleaning done to prepare the ICA dataset).
Once the data was cleaned with ICA, we applied a non-causal 8th order Butterworth
bandpass filter with cutoffs at 100 and 1500Hz and then re-referenced to the average of the left
and right earlobes. If one of the earlobe channels had been rejected as a bad channel, the
average of the left and right mastoid was used instead. Activity at channel Cz (the previous
reference) was then recalculated and added back to the dataset. We then interpolated any
rejected bad channels using spherical interpolation.
Epochs were extracted between –2ms and +15ms of chirp onset for all conditions with
baseline correction from –2 to 0ms. The number of epochs was balanced so that each condition

had the same number of chirps both in total, and from each spatial location (left, right), resulting
in each condition (mono talker male, dual target male, dual target female, dual masker male,
dual masker female) having 7,886 epochs per condition equally distributed across left and right
presentation (3,943 epochs from both left- and right-side presentations per condition). The
smallest number of chirps on any one side for all conditions was calculated and epochs were
randomly sampled so all conditions were balanced. Balancing of the number of epochs for each
condition was done to avoid any systematic condition differences due to a difference in signal to
noise (SNR) caused by some conditions having more chirps than others. This difference was
primarily due to the female-voiced stories, where more frequent glottal pulses (driven by a
higher F0) resulted in approximately 9% more chirps in the female-voiced stories compared to
the male-voiced stories. Additional differences in total chirp events were due to an approximate
3% variation of chirps across stories (i.e., subtle differences in voicing and pausing). Given
evidence for left-right asymmetries in speech processing and neural encoding (e.g., "right-ear
advantage"; Ahonniska et al., 1993; Jerger and Martin, 2004; Hornickel et al., 2008; Keefe et al.,
2008; Bidelman and Bhagat, 2015), we also balanced the number of chirps across stimulus
presentation side for each condition to ensure results were not driven by unequal ear-based
SNR in the EEG measures (approximately 5% variation in the of number of chirps across sides
within each story). Epochs with artifacts were then removed with a 2-step process: first, any
epochs with activity outside the threshold of +/-100µV were rejected, and then any remaining
epochs whose activity was outside of 6 SD for a single channel or 2 SD across all channels
were eliminated (mean=9.85 ± 6.24% SD of rejected epochs, range=3.65-29.96% rejected
epochs across individuals and conditions).
ABR Wave V peak amplitudes and latencies were extracted at channel Cz from the
average amplitude over time across the remaining epochs for each condition for each
participant. Wave V latency measures were extracted by fitting a local positive peak in a window
between 5-8 ms post chirp-onset. Wave V amplitude was computed as the peak-to-peak
amplitude between local positive peak within 5-8 ms and the subsequent local negative peak
between 7-11 ms. The measurement windows were determined based on visual inspection of all
individual Wave Vs for each condition. In cases in which more than one local peak was seen in
the wave V waveform inside the measurement window, the values from the peak with the
greatest amplitude were taken.
481
Statistical analyses

Unless otherwise specified, mixed effects linear regression models and repeated-
measures ANOVAs were used to analyze the data in SAS Studio (v.3.81). All regression models
included a random intercept of subject to account for inherent variability present in each
individual and improve generalizability of the results. Outcome measures were ABR wave V
amplitudes and latencies for the neural models, and the behavioral metrics (i.e., target word
identification accuracy, target word identification reaction time, comprehension question
accuracy, subjective task effort ratings) were included as dependent variables for the behavior-
only models as well as the brain-behavior comparisons. Independent variables were
characterized by the condition of interest (i.e., one versus two talkers; target versus masker;
also main effects and interaction terms with male versus female voices where appropriate).
When interactions were not significant in the full model, they were removed using a backwards
selection procedure, and the remaining simplified model variables are reported in the results
below. One-way repeated-measures ANOVAs were used to deconstruct within-subject condition
effects for both the brain and behavior metrics individually. All post hoc comparisons were
corrected using Tukey adjustment. Outliers were identified by residuals beyond ±2. Non-

normality of the dependent measures was evaluated using the Shaprio-Wilk test. A Box-Cox
transformation procedure was performed on non-normal variables; specifically, the first
convenient power parameter within the lambda confidence interval was used to select a
transformation method (cf. optimal power parameter; specified by the "convenient" t-option).
However, data transformations did not alter results for any non-normally distributed data, so the
non-transformed results are reported below.
504
Results

<u>ABR condition effects</u>

Masking and speech in noise effects: single- vs. dual-talker

To evaluate speech-in-noise (SiN) effects on the ABR wave V, we compared Cheech
responses for the male voice mono-talker and the dual-talker conditions (i.e., main effect
variable dummy coded for either mono- or dual-talker). Overall, the dual-talker condition was
associated with both reduced wave V amplitudes ($F_{1,99}$=25.84, p<0.0001) and delayed latencies
($F_{1,99}$=10.82, p=0.0014) compared to the mono-talker condition. A one-way, repeated-measures
ANOVA was then used to elucidate underlying relationships between all conditions (i.e., mono-
talker male target, dual-talker male target, dual-talker male masker, dual-talker female target,
and dual-talker female masker). Tukey-adjusted post hoc comparisons revealed larger
amplitudes for the mono-talker (male) condition compared to all dual-talker conditions
individually (p's<0.05). Additionally, faster latencies were observed for the mono-talker condition
compared to the dual-talker female target (Tukey-adjusted p=0.0276) and dual-talker male
masker (Tukey-adjusted p=0.0043). As expected, these results suggest poorer brainstem
encoding for stimuli in the presence of a competing masker (i.e., smaller amplitudes and
delayed latencies; Figure 3).

Attention effects

Attention effects within the brainstem were assessed as the difference between target
and masker responses across the listening blocks. Considering the effects of target versus
masker alone, no differences were observed in either wave V amplitudes ($F_{1,99}$=2.96, p=0.0885)
or latencies ($F_{1,99}$=1.75, p=0.1886). However, as suggested by the grand average waveforms
(Figure 3A), the attention effect could be modulated by the talker's voice. We therefore tested
for the interaction of the talker sex and directed attention (target versus masker) as well as
respective main effects on wave V measures. The interaction between selective attention and
voice sex was significant for both wave V amplitudes ($F_{1,97}$=7.19, p=0.0086) and wave V
latencies ($F_{1,97}$=8.17, p=0.0052). Tukey-adjusted simple effect comparisons suggested larger
amplitudes for the male target (mono- and dual-talker combined) compared to the male masker
(p=0.0032) and female target (p=0.0447). Additionally, faster latencies were observed for the
male target compared to the male masker (p=0.0055) and the female target voice (p=0.0341) in
post hoc contrasts. As described above, these effects are at least partially driven by larger
amplitudes and faster latencies for the mono-talker male condition compared to the dual-talker
conditions.

Figure 3: (A) ABR waveforms evoked by the Cheech chirps across target and masker conditions show wave V differences with the presence of another talker (speech-in-noise effects) as well as directed selective attention (target vs. masker). (B) Wave V amplitudes were larger for the mono-talker condition compared to all dual-talker conditions. The male target was also (combined mono- & dual-talker) was larger in amplitude compared to the male masker and female target. (C) Wave V latencies were faster for the mono-talker condition compared to the dual-talker conditions. Additionally, faster latencies were observed for the male target compared to the male masker and the female target voice. Violin plots in B & C map the kernel density estimates of the data capped at the range bounds. Inlaid box plots indicate the first and third quartiles (25th and 75th percentiles) with the white rectangle, median shown by the black horizontal line, and the range depicted by the whiskers up to 1.5x the inter-quartile range (IQR).

539   Behavioral performance

540   Selective attention performance was measured behaviorally as identification accuracy
and corresponding reaction times for color words embedded in the target story (Figure 4). Color
word identification accuracy significantly differed across all three experimental conditions
($F_{2,48}$=18.15, $p<0.0001$). Specifically, both the mono-talker male and dual-talker male conditions
were associated with higher identification accuracy than the dual-talker female (both Tukey-
adjusted $p<0.0001$; Figure 4A). The two male talker conditions were not significantly different

(p=0.8931), suggesting a slight behavioral identification accuracy decrement specific to the
female-voiced story, not necessarily with the general presence of a competing talker.
Meanwhile, target color word reaction times also differed across conditions ($F_{2,48}$=10.02,
p=0.0002). Specifically, faster reaction times were observed for the mono-talker block compared
to the dual-talker female (p=0.0001; Figure 4B). Although the other Tukey-adjusted post hoc
comparisons were not significant, reactions times trended towards faster reaction times for the
mono-talker male compared to the dual-talker male (p=0.0739) and for the dual-talker male
compared to the dual-talker female condition (p=0.0760).
Following the listening task, participants were asked to complete several comprehension
questions to probe their understanding of the short story segments. Response accuracy for
comprehension questions was higher for the mono-talker condition compared to either dual-
talker condition (overall $F_{2,48}$=5.97, p=0.0048; Tukey-adjusted mono-talker vs. dual-talker male
p=0.0501, mono-talker vs. dual-talker female p=0.0045; Figure 4C). No difference was observed
between the male- and female-voiced dual-talker conditions (p=0.6243). Participants then
completed the NASA Task Load Index questionnaire (Hart and Staveland, 1988) as a measure
of their subjective task load or effort. The mono-talker story was associated with lower scores
(i.e., less perceived task load) compared to either dual-talker condition (dual-talker male:
p=0.0182; dual-talker female: p=0.0009), but the dual-talker blocks did not significantly differ
from each other (p=0.5410; Figure 4D). Collectively, the results suggest that the dual-talker
scenario was associated with both greater perceived effort (task load) and impaired
comprehension abilities.

Figure 4: Behavioral results for color word identification accuracy (A), identification reaction times (B), narrative comprehension (C), and ratings of subjective task load or effort (D). Violin plots map the kernel density estimates of the data capped at the range bounds. Inlaid box plots indicate the first and third quartiles (25th and 75th percentiles) with the white rectangle, median shown by the black horizontal line, and the range depicted by the whiskers up to 1.5x IQR.

567   Relationships between behavioral measures were also evaluated using linear mixed-
568   effects regression models. All four behavioral outcomes were associated with each other (all p's
569   < 0.05). As might be expected, better behavioral performance was characterized by higher
570   identification accuracy, faster identification reaction times, higher comprehension question
571   accuracy, and lower effort ratings.
572

<u>Brainstem-behavior relationships</u>

A compelling strength of our Cheech technique is the capacity to simultaneously
measure ABRs while participants complete a listening task with continuous speech (i.e.,
identification of color words embedded in short story narratives). We first analyzed mixed effects
models with main effects and interactions of directed attention (i.e., target versus masker), talker
sex, and wave V measures to predict behavioral outcomes (i.e., separate models for latencies
and amplitudes). No 3- or 2-way interactions were observed between either wave V measure
and directed attention or talker sex variables. The only 2-way interactions between talker sex
and directed attention observed in the full model simply replicated findings previously reported
in the Behavioral Performance section.
The models were then re-analyzed to assess only main effects of ABR wave V
amplitudes and latencies separately on behavioral outcomes (Table 1). In these simplified
analyses, faster wave V latencies were associated with better target word identification
accuracy ($F_{1,99}$=8.59, p=0.0042), better comprehension question accuracy ($F_{1,99}$=4.73,
p=0.0320), and lower subjective task load ($F_{1,99}$=19.28, p<0.0001). Latencies alone did not
correlate with target word identification reaction times ($F_{1,99}$=1.06, p=0.3050). Further, larger
wave V amplitudes were associated with better target word identification accuracy ($F_{1,99}$=9.52,
p=0.0026) and reaction times ($F_{1,99}$=8.82, p=0.0037) but not comprehension question accuracy
nor subjective task load ($F_{1,99}$=1.41, p=0.2377 & $F_{1,99}$=0.66, p=0.4172 respectively). Collectively,
these results indicate a strong relationship between wave V and measures of selective attention
(i.e., target word identification). Meanwhile, neural encoding speed (i.e., wave V latencies) but
not encoding strength (i.e., amplitudes) related to both narrative comprehension and perceived
effort.
596

597   **Table 1: Brainstem-behavior regression model results for wave V main effects**
598

Wave V predictor	Outcome	β estimate	$F_{1,99}$ statistic	P-value *value = p<0.05
Latencies	Target word identification accuracy	-0.0996	8.59	**0.0042***
	Target word identification reaction times	0.0283	1.06	0.3050
	Comprehension question accuracy	-0.0935	4.73	**0.0320***
	Subjective task load scores	9.7502	19.28	**<0.0001***

Amplitudes	Target word identification accuracy	0.4241	9.52	**0.0026***
	Target word identification reaction times	-0.3235	8.82	**0.0037***
	Comprehension question accuracy	0.2050	1.41	0.2377
	Subjective task load scores	-8.0859	0.66	0.4172

599
600
Discussion

We evaluated the effects of speech-in-noise (SiN) and selective attention on low-level
brainstem encoding. Our novel experimental paradigm permitted simultaneous measurement of
both ABRs and behavior in both single- and dual-talker conditions. ABRs recorded in the
presence of a competing talker were associated with smaller amplitudes and delayed latencies
compared to the mono-talker condition. Interestingly, the results also suggested a role of
directed attention on ABR wave V latencies. Faster wave V latencies were associated with
better accuracy of color word identification as well as higher comprehension question accuracy
and lower scores of subjective listening effort. Meanwhile, more robust wave V amplitudes
corresponded to better accuracy and faster reaction times for target word identification during
the selective attention listening task. These effects on directed attention were partially driven by
talker's sex. Collectively, the results suggest that faster, more robust neural encoding within the
auditory brainstem contributes to successful auditory selective attention and SiN listening
processes.
Selective auditory attention within the brainstem was characterized by wave V response
differences when participants were instructed to attend to or ignore short story narratives. These
results align with other reports of attention-driven influences on subcortical processing. For
example, Forte et al. (2017) and Saiz-Alía & Reichenbach (2020) noted larger computationally-
derived auditory brainstem amplitudes for attended over unattended speech. Other studies have
shown modulation of transient-evoked ABRs while attending to the auditory modality versus
another (typically visual) domain (Lukas, 1980; Bauer and Bayles, 1990; Ikeda and Campbell,
2021, 2024; Kumar et al., 2023) or with increasing cognitive interference (Sörqvist et al., 2012;
Brännström et al., 2020). Our results are also consistent with attentional modulation of
sustained subcortical frequency-following responses (Hoormann et al., 2000; e.g., Galbraith et
al., 2003; Hairston et al., 2013; Lehmann and Schönwiesner, 2014; cf. Varghese et al., 2015). It
is likely that corticofugal pathways—particularly those projecting to the inferior colliculus, a key
neural generator of wave V (Starr and Hamilton, 1976; Hashimoto et al., 1981; Huffman and
Henson, 1990)—contribute to directed attention effects on the ABR. Yet, the ABR has
historically been viewed as largely immune from attention effects (e.g., Amadeo and Shagass,
1973; Picton and Hillyard, 1974; Connolly et al., 1989; Gregory et al., 1989). These
discrepancies across studies likely result from differences in experimental paradigms. It is
possible that attention-related effects on subcortical processes are most effectively revealed

when another attention-demanding task is presented concurrently with the auditory stimuli
(Lukas, 1980; e.g., Brännström et al., 2020; Kumar et al., 2023) or when the stimuli are more
ecologically valid and relevant for real-world listening, as in the case of continuous speech (Ding
and Simon, 2012; Zion Golumbic et al., 2013; Forte et al., 2017; Backer et al., 2019; Etard et al.,
2019; Teoh and Lalor, 2019; Saiz-Alía and Reichenbach, 2020; Teoh et al., 2022; Carta et al.,
2024). Thus, Cheech demonstrates potential for reliable detection of top-down, selective
attention effects on subcortical processes.
Crucially, faster and/or larger Cheech-evoked wave V responses were associated with
more successful speech listening performance, including behavioral measures of selective
attention (i.e., identification accuracy, RTs) as well as comprehension questions and subjective
reports of listening effort. That is, more robust encoding in lower-level brainstem generators was
related to higher-level speech recognition abilities, with and without the presence of a competing
talker. Subcortical responses have been associated with other speech-listening behaviors
including SiN perception (Papakonstantinou et al., 2011; Bramhall et al., 2015; Saiz-Alía and
Reichenbach, 2020), self-reports of speech-listening abilities (Anderson et al., 2013), early
categorization of speech phonemes (Carter and Bidelman, 2023; Rizzi and Bidelman, 2023), or
even measures of auditory processing disorders (Omidvar et al., 2023). Collectively, our results
provide further support for the important role of the brainstem in speech perception, both in quiet
and the presence of a masking talker, as well as higher-level narrative comprehension and
subjective perception of task-related listening effort.
As expected, the presence of a competing talker was associated with weaker brainstem
encoding (i.e., wave V reduced amplitudes, delayed latencies), worse comprehension accuracy,
and higher ratings of subjective task load. SiN perception (the "cocktail party problem") is an
important ability for listening in real-world environments, and some speech-hearing difficulties
are only exposed when stimuli are presented in background noise. It remains possible that the
variability of SiN perceptual skills, even among normal hearing listeners, could be traced to low-
level neural encoding differences (e.g., Bharadwaj et al., 2015; Bramhall et al., 2015, 2017; Jain
et al., 2019) or even top-down corticofugal modulations (Price and Bidelman, 2021). Varied
contributions (or degeneration/selective loss) of auditory nerve fibers during speech encoding,
particularly those with low-spontaneous discharge rates, could also help explain individual SiN
differences (Mehraei et al., 2016, 2017). While the current study cannot differentiate between
these theories—since both top-down and/or bottom-up mechanisms could alter ABR wave V
and its relationship with speech perception—our results provide evidence of subcortical
involvement for continuous SiN and suggest a possible neural factor for individual perception
differences.
Contrasting with our findings, previous studies that compared male and female talkers
reported larger wave V response amplitudes for male voices compared to female voices (Forte
et al., 2017; Saiz-Alía and Reichenbach, 2020; Polonenko and Maddox, 2021). Yet, our
selective attention-talker sex interaction for wave V amplitudes did align with previous reports of
larger male-attended amplitudes compared to unattended speech, though a similar trend was
noted for the female voice in Forte et al. (2017). One explanation could be due to differences in
experimental stimulus design. Our Cheech and/or female-voice modulation processes could
have affected perceptual clarity and thus impacted brain and/or behavioral responses even
though average RMS values were equated across the stories. Our female-sounding voice was
modified from the original male voice (same talker for all stories) to maintain similar intonation,
phrasing, and other expressive characteristics across the stories; only the speaker's F0 and
vocal tract length parameters were resynthesizd using MATLAB STRAIGHT to create a female-
sounding voice (Kawahara and Morise, 2011). Additionally, higher rates of stimulation are
associated with reduced ABRs (e.g., Don et al., 1977; Weber and Fujikawa, 1977; Fowler and

Noffsinger, 1983). Higher F0s and thus faster glottal pulse rates could explain subcortical
encoding differences from speech stimuli spoken by different talkers (Polonenko and Maddox,
2021). While our stimuli had approximately 10% more chirps for female-voiced stories
compared to the male-voiced stories (i.e., faster glottal pluses creates more chirp-embedding
opportunities), we set a minimum inter-stimulus interval of 18.2 ms between chirps to reduce
overlapping activity and detrimental effects of high stimulation rates on neural encoding
processes. We also equated the number of chirp trials across conditions in our analyses to
avoid any systematic confounds produced by talker sex-based SNR differences due to
imbalanced chirp events. Finally, chirps are also designed to activate the entire cochlea in a
near-synchronous fashion which produces more robust ABRs in general (Elberling et al., 2007,
2010) which could explain neurophysiologic differences between our study and other
computationally-derived methods more generally.
Additionally, our male and female talkers were spatially separated using head-related
transfer functions (HRTFs), and the talker locations were dynamically swapped with each other
on average every 6 seconds to track spatially-driven selective attention similar to Teoh & Lalor
(2019). Pilot testing confirmed that the combined effects of pitch modulation (to produce a
female-sounding voice) and spatial separation led to better listening performance, consistent
with previous literature demonstrating talker-gender, spatially-separated release of masking
effects in multi-talker environments (Oh et al., 2021, 2022). Additionally, gaze direction (as when
participants were asked to follow the target cue across the two screens) could potentially
modulate neural or myogenic activity that could contaminate responses (e.g., O'Beirne and
Patuzzi, 1999; Patuzzi and O'Beirne, 1999; Bidelman et al., 2024). Thus, we cannot rule out
certain idiosyncratic effects of our experimental paradigm on the observed brainstem-behavior
relationships. Future studies, both involving Cheech and other (ideally continuous) speech
stimuli, may help elucidate the underlying neural encoding mechanisms involved in processing
individual talker characteristics and its impact on speech perception abilities.
708
Conclusion

Collectively, our results provide evidence for the role of brainstem encoding process in individual
speech perception abilities including SiN recognition, selective auditory attention, and
comprehension. These findings underscore the importance of lower-level auditory structures for
speech processing, even in audiometrically normal listeners. They furthermore have
implications for an improved understanding of other populations (e.g., older adults, children,
individuals with hearing loss or assistive hearing devices, etc.) and of certain speech processing
disorders. Our study highlights the advantage of a stimulus like chirped-speech (Cheech) and
more ecologically valid paradigms to uncover novel and impactful relationships between lower-
level neural processing and higher-level listening skills in real-world auditory environments.

Appendix A

<u>Short story stimulus details</u>

Story title	Author	Collection	Experimental condition	Male or female voice
The Master Cat, or Puss in Boots	Charles Perrault, translated by J. E. Mansion and Robert Samber	The Fairy Tales of Charles Perrault	Mono-talker	Male
Lily and the Lion	Jacob Grimm & Wilhelm Grimm	Grimms' Fairy Tales	Dual-talker	Male
The Four Clever Brothers	Jacob Grimm & Wilhelm Grimm	Grimms' Fairy Tales	Dual-talker	Female
The Wanderers	Edith Howes	Maoriland Fairy Tales	Practice only (mono-talker)	Male

724 **Appendix B**

725 <u>Cheech synthesis parameters</u>

Cheech Parameter	**Value**
Fs (sample rate; Hz)	48000
Speech filter bands (Hz)	0-250; 500-1000; 2000-4000; 11000+
Chirp filter bands (Hz)	250-500; 1000-2000; 4000-11000
Speech filter order	10
Chirp filter order	4
Voicing (speech envelope) filter, to identify voicing by power (Hz)	20-1000
Voicing threshold, for identifying sufficient glottal pulse energy (normalized amplitude)	0.03
Minimum voiced period to chirp (sec)	0.05
Minimum chirp ISI (sec)	0.0182
Minimum gap after first chirp (sec)	0.048
Maximum voicing duration, above which gaps will be inserted (intra-voicing gap) to prevent very long chirp trains especially during loud speech modulations (sec)	0.400
Intra-voicing gap (sec)	0.05
Normalized chirp amplitude	0.10

Female voice characteristics
    Resynthesized voice mean F0 pitch (Hz)
    Maximum frequency modulation (Hz)
    Vocal tract length warping factor
	
	180
	4000
	1.04

Resynthesized voice characteristics
    Score, to reduce thumping in the resynthesized audio
    Margin, to reduce buzzing in the resynthesized audio
	
	0.6
	
	0.6

References

Adler NE, Epel ES, Castellazzo G, Ickovics JR (2000) Relationship of subjective and objective
social status with psychological and physiological functioning: Preliminary data in
healthy, white women. Health Psychol 19:586–592.

Ahonniska J, Cantell M, Tolvanen A, Lyytinen H (1993) Speech perception and brain laterality:
The effect of ear advantage on auditory event-related potentials. Brain Lang 45:127–
146.

Amadeo M, Shagass C (1973) Brief latency click-evoked potentials during waking and sleep in
man. Psychophysiology 10:244–250.

Anderson S, Parbery-Clark A, White-Schwoch T, Kraus N (2013) Auditory brainstem response to
complex sounds predicts self-reported speech-in-noise performance. J Speech Lang
Hear Res JSLHR 56:31–43.

Armstrong C, Thresh L, Murphy D, Kearney G (2018) A perceptual evaluation of individual and
non-individual HRTFs: A case study of the SADIE II database. Appl Sci 8:2029.

Backer KC, Kessler AS, Lawyer LA, Corina DP, Miller LM (2019) A novel EEG paradigm to
simultaneously and rapidly assess the functioning of auditory and visual pathways. J
Neurophysiol 122:1312–1329.

Bauer LO, Bayles RL (1990) Precortical filtering and selective attention: An evoked potential
analysis. Biol Psychol 30:21–33.

Bharadwaj HM, Masud S, Mehraei G, Verhulst S, Shinn-Cunningham BG (2015) Individual
differences reveal correlates of hidden hearing deficits. J Neurosci 35:2161–2172.

Bidelman GM (2018) Subcortical sources dominate the neuroelectric auditory frequency-
following response to speech. NeuroImage 175:56–69.

Bidelman GM, Bhagat SP (2015) Right-ear advantage drives the link between olivocochlear
efferent 'antimasking' and speech-in-noise listening benefits. NeuroReport 26:483.

Bidelman GM, Sisson A, Rizzi R, MacLean J, Baer K (2024) Myogenic artifacts masquerade as
neuroplasticity in the auditory frequency-following response. Front Neurosci 18:1422903.

Brainard DH (1997) The Psychophysics Toolbox. Spat Vis 10:433–436.

Bramhall NF, Konrad-Martin D, McMillan GP, Griest SE (2017) Auditory brainstem response
altered in humans with noise exposure despite normal outer hair cell function. Ear Hear
38:e1–e12.

Bramhall NF, Ong B, Ko J, Parker M (2015) Speech perception ability in noise is correlated with
auditory brainstem response wave I amplitude. J Am Acad Audiol 26:509–517.

Brännström KJ, Wilson WJ, Waechter S (2020) Increasing cognitive interference modulates the
amplitude of the auditory brainstem response. J Am Acad Audiol 29:512–519.

Broderick MP, Anderson AJ, Di Liberto GM, Crosse MJ, Lalor EC (2018) Electrophysiological
correlates of semantic dissimilarity reflect the comprehension of natural, narrative
speech. Curr Biol CB 28:803-809.e3.

Campbell T, Kerlin JR, Bishop CW, Miller LM (2012) Methods to eliminate stimulus transduction
artifact from insert earphones during electroencephalography. Ear Hear 33:144–150.

Carta S, Aličković E, Zaar J, Valdés AL, Liberto GMD (2024) Cortical encoding of phonetic
onsets of both attended and ignored speech in hearing impaired individuals. PLOS ONE
19:e0308554.

Carter JA, Bidelman GM (2023) Perceptual warping exposes categorical representations for
speech in human brainstem responses. NeuroImage 269:119899.

Coffey EBJ, Herholz SC, Chepesiuk AMP, Baillet S, Zatorre RJ (2016) Cortical contributions to
the auditory frequency-following response revealed by MEG. Nat Commun 7:11070.

Connolly JF, Aubry K, McGillivary N, Scott DW (1989) Human brainstem auditory evoked
potentials fFail to provide evidence of efferent modulation of auditory input during
attentional tasks. Psychophysiology 26:292–303.

Delorme A, Makeig S (2004) EEGLAB: an open source toolbox for analysis of single-trial EEG
dynamics including independent component analysis. J Neurosci Methods 134:9–21.

Ding N, Simon JZ (2012) Emergence of neural encoding of auditory objects while listening to
competing speakers. Proc Natl Acad Sci 109:11854–11859.

Don M, Allen AR, Starr A (1977) Effect of click rate on the latency of auditory brain stem
responses in humans. Ann Otol Rhinol Laryngol 86:186–195.

Elberling C, Callø J, Don M (2010) Evaluating auditory brainstem responses to different chirp
stimuli at three levels of stimulation. J Acoust Soc Am 128:215–223.

Elberling C, Don M, Cebulla M, Stürzebecher E (2007) Auditory steady-state responses to chirp
stimuli based on cochlear traveling wave delay. J Acoust Soc Am 122:2772–2785.

Escabí MA, Miller LM, Read HL, Schreiner CE (2003) Naturalistic auditory contrast improves
spectrotemporal coding in the cat inferior colliculus. J Neurosci 23:11489–11504.

Etard O, Kegler M, Braiman C, Forte AE, Reichenbach T (2019) Decoding of selective attention
to continuous speech from the human auditory brainstem response. NeuroImage 200:1–
11.

Forte AE, Etard O, Reichenbach T (2017) The human auditory brainstem response to running
speech reveals a subcortical mechanism for selective attention Shinn-Cunningham BG,
ed. eLife 6:e27203.

Fowler CG, Noffsinger D (1983) Effects of stimulus repetition rate and frequency on the auditory
brainstem response in normal, cochlear-impaired, and VIII nerve/brainstem-impaired
subjects. J Speech Lang Hear Res 26:560–567.

Galbraith GC, Olfman DM, Huffman TM (2003) Selective attention affects human brain stem
frequency-following response. NeuroReport 14:735.

Gregory SD, Heath JA, Rosenberg ME (1989) Does selective attention influence the brain-stem
auditory evoked potential? Electroencephalogr Clin Neurophysiol 73:557–560.

Hairston WD, Letowski TR, McDowell K (2013) Task-related suppression of the brainstem
frequency following response. PLOS ONE 8:e55215.

Hamilton LS, Huth AG (2020) The revolution will not be controlled: natural stimuli in speech
neuroscience. Lang Cogn Neurosci 35:573–582.

Hart SG, Staveland LE (1988) Development of NASA-TLX (Task Load Index): Results of
empirical and theoretical research. In: Human Mental Workload (Hancock PA, Meshkati
N, eds). Amsterdam: North Holland Press.

Hashimoto I, Ishiyama Y, Yoshimoto T, Nemoto S (1981) Brain-stem auditory-evoked potentials
recorded directly from human brain-stem and thalamus. Brain J Neurol 104:841–859.

Hickok G, Poeppel D (2004) Dorsal and ventral streams: a framework for understanding aspects
of the functional anatomy of language. Cognition 92:67–99.

Hoormann J, Falkenstein M, Hohnsbein J (2000) Early attention effects in human auditory-
evoked potentials. Psychophysiology 37:29–42.

Hornickel J, Skoe E, Kraus N (2008) Subcortical laterality of speech encoding. Audiol Neurotol
14:198–207.

Huffman RF, Henson OW (1990) The descending auditory pathway and acousticomotor
systems: connections with the inferior colliculus. Brain Res Rev 15:295–323.

Ikeda K, Campbell TA (2021) Reinterpreting the human ABR binaural interaction component:
isolating attention from stimulus effects. Hear Res 410:108350.

Ikeda K, Campbell TA (2024) Binaural interaction in human auditory brainstem and middle-
latency responses affected by sound frequency band, lateralization predictability, and
attended modality. Hear Res 452:109089.

Jain C, Dwarakanath VM, G A (2019) Influence of subcortical auditory processing and cognitive
measures on cocktail party listening in younger and older adults. Int J Audiol 58:87–96.

Jerger J, Martin J (2004) Hemispheric asymmetry of the right ear advantage in dichotic listening.
Hear Res 198:125–136.

Jewett DL, Williston JS (1971) Auditory-evoked far fields averaged from the scalp of humans.
Brain J Neurol 94:681–696.

Kawahara H, Masuda-Katsuse I, de Cheveigné A (1999) Restructuring speech representations
using a pitch-adaptive time–frequency smoothing and an instantaneous-frequency-
based F0 extraction: Possible role of a repetitive structure in sounds. Speech Commun
27:187–207.

Kawahara H, Morise M (2011) Technical foundations of TANDEM-STRAIGHT, a speech
835   analysis, modification and synthesis framework. Sadhana 36:713–727.

836   Kawahara H, Morise M, Takahashi T, Nisimura R, Irino T, Banno H (2008) Tandem-STRAIGHT:
837   A temporally stable power spectral representation for periodic signals and applications to
838   interference-free spectrum, F0, and aperiodicity estimation. In: 2008 IEEE International
839   Conference on Acoustics, Speech and Signal Processing, pp 3933–3936.

840   Keefe DH, Gorga MP, Jesteadt W, Smith LM (2008) Ear asymmetries in middle-ear, cochlear,
841   and brainstem responses in human infantsa). J Acoust Soc Am 123:1504–1512.

842   Krishnan A, Gandour JT (2009) The role of the auditory brainstem in processing linguistically-
843   relevant pitch patterns. Brain Lang 110:135–148.

844   Krishnan A, Plack CJ (2011) Neural encoding in the human brainstem relevant to the pitch of
845   complex tones. Hear Res 275:110–119.

846   Kulasingham JP, Bachmann FL, Eskelund K, Enqvist M, Innes-Brown H, Alickovic E (2024)
847   Predictors for estimating subcortical EEG responses to continuous speech. PLOS ONE
848   19:e0297826.

849   Kumar S, Nayak S, Pitchai Muthu AN (2023) Effect of selective attention on auditory brainstem
850   response. Hear Balance Commun 21:139–147.

851   Lehmann A, Schönwiesner M (2014) Selective attention modulates human auditory brainstem
852   responses: Relative contributions of frequency and spatial cues. PLOS ONE 9:e85442.

853   Lopez-Calderon J, Luck SJ (2014) ERPLAB: an open-source toolbox for the analysis of event-
854   related potentials. Front Hum Neurosci 8.

855   Luck SJ (2022) Applied Event-Related Potential Data Analysis. LibreTexts.

856   Lukas JH (1980) Human auditory attention: The olivocochlear bundle may function as a
857   peripheral filter. Psychophysiology 17:444–452.

858   Maddox RK, Lee AKC (2018) Auditory brainstem responses to continuous natural speech in
859   human listeners. eNeuro 5:1–13.

860   Mehraei G, Gallardo AP, Shinn-Cunningham BG, Dau T (2017) Auditory brainstem response
861   latency in forward masking, a marker of sensory deficits in listeners with normal hearing
862   thresholds. Hear Res 346:34–44.

863   Mehraei G, Hickox AE, Bharadwaj HM, Goldberg H, Verhulst S, Liberman MC, Shinn-
864   Cunningham BG (2016) Auditory brainstem response latency in noise as a marker of
865   cochlear synaptopathy. J Neurosci 36:3755–3764.

866   Miller LM, Moore BD, Bishop CW (2020) Frequency-multiplexed speech-sound stimuli for
867   hierarchical neural characterization of speech processing. Available at:
868   https://patents.google.com/patent/WO2016011189A1/en.

869   Nasreddine ZS, Phillips NA, Bédirian V, Charbonneau S, Whitehead V, Collin I, Cummings JL,
870   Chertkow H (2005) The Montreal Cognitive Assessment, MoCA: A brief screening tool
871   for mild cognitive impairment. J Am Geriatr Soc 53:695–699.

872   O'Beirne GA, Patuzzi RB (1999) Basic properties of the sound-evoked post-auricular muscle
873   response (PAMR). Hear Res 138:115–132.

874   Oh Y, Bridges SE, Schoenfeld H, Layne AO, Eddins D (2021) Interaction between voice-gender
875   difference and spatial separation in release from masking in multi-talker listening
876   environments. JASA Express Lett 1:084404.

877   Oh Y, Hartling CL, Srinivasan NK, Diedesch AC, Gallun FJ, Reiss LAJ (2022) Factors underlying
878   masking release by voice-gender differences and spatial separation cues in multi-talker
879   listening environments in listeners with and without hearing loss. Front Neurosci 16.

880   Omidvar S, Mochiatti Guijo L, Duda V, Costa-Faidella J, Escera C, Koravand A (2023) Can
881   auditory evoked responses elicited to click and/or verbal sound identify children with or
882   at risk of central auditory processing disorder: A scoping review. Int J Pediatr
883   Otorhinolaryngol 171:111609.

884   Papakonstantinou A, Strelcyk O, Dau T (2011) Relations between perceptual measures of
885   temporal processing, auditory-evoked brainstem responses and speech intelligibility in
886   noise. Hear Res 280:30–37.

887   Patuzzi RB, O'Beirne GA (1999) Effects of eye rotation on the sound-evoked post-auricular
888   muscle response (PAMR). Hear Res 138:133–146.

889   Pelli DG (1997) The VideoToolbox software for visual psychophysics: transforming numbers into
890   movies. Spat Vis 10:437–442.

891   Picton TW, Hillyard SA (1974) Human auditory evoked potentials. II: Effects of attention.
892   Electroencephalogr Clin Neurophysiol 36:191–200.

893   Polonenko MJ, Maddox RK (2021) Exposing distinct subcortical components of the auditory
894   brainstem response evoked by continuous naturalistic speech Reichenbach T, King AJ,
895   Reichenbach T, Simon JZ, eds. eLife 10:e62329.

896   Price CN, Bidelman GM (2021) Attention reinforces human corticofugal system to aid speech
897   perception in noise. NeuroImage 235:118014.

898   Rizzi R, Bidelman GM (2023) Duplex perception reveals brainstem auditory representations are
899   modulated by listeners' ongoing percept for speech. Cereb Cortex 33:10076–10086.

900   Saiz-Alía M, Reichenbach T (2020) Computational modeling of the auditory brainstem response
901   to continuous speech. J Neural Eng 17:036035.

902   Skoe E, Kraus N (2010) Auditory brainstem response to complex sounds: A tutorial. Ear Hear
903   31:302–324.

904   Sörqvist P, Stenfelt S, Rönnberg J (2012) Working memory capacity and visual–verbal cognitive
905   load modulate auditory–sensory gating in the brainstem: Toward a unified view of
906   attention. J Cogn Neurosci 24:2147–2154.

907   Starr A, Hamilton AE (1976) Correlation between confirmed sites of neurological lesions and
908   abnormalities of far-field auditory brainstem responses. Electroencephalogr Clin
909   Neurophysiol 41:595–608.

910   Teoh ES, Ahmed F, Lalor EC (2022) Attention differentially affects acoustic and phonetic feature
911   encoding in a multispeaker environment. J Neurosci 42:682–691.

912   Teoh ES, Lalor EC (2019) EEG decoding of the target speaker in a cocktail party scenario:
913   considerations regarding dynamic switching of talker location. J Neural Eng 16:036017.

914   Varghese L, Bharadwaj HM, Shinn-Cunningham BG (2015) Evidence against attentional state
915   modulating scalp-recorded auditory brainstem steady-state responses. Brain Res
916   1626:146–164.

917   Weber BA, Fujikawa SM (1977) Brainstem evoked response (BER) audiometry at various
918   stimulus presentation rates. Ear Hear 3:59.

919   Zion Golumbic EM, Ding N, Bickel S, Lakatos P, Schevon CA, McKhann GM, Goodman RR,
920   Emerson R, Mehta AD, Simon JZ, Poeppel D, Schroeder CE (2013) Mechanisms
921   underlying selective neuronal tracking of attended speech at a "cocktail party." Neuron
922   77:980–991.

924