eDNA provides accurate and precise estimates of abundance by integrating bioenergetics and particle mass-balance

Beaulieu J[1,2], MC Yates[1,2], DJ Fraser[2,3], ME Cristescu[2,4] and AM Derry[1,2]

[1] Département des sciences biologiques, Université du Québec à Montréal, Montréal, QC, Canada

[2] Groupe interuniversitaire en limnologie (GRIL), Montréal, QC, Canada

[3] Department of Biology, Concordia University, Montreal, QC, Canada

[4] Department of Biology, McGill University, Montréal, QC, Canada

ORCID and emails:

Julien Beaulieu: **0000-0003-2597-8921**   beaulieu.julien.3@courrier.uqam.ca
Matthew C. Yates**: 0000-0002-9199-1078**   matthew.yates@outlook.com
Dylan J. Fraser**: 0000-0002-5686-7338**   dylan.fraser@concordia.ca
Melania E. Cristescu: **0000-0002-0854-4040**   melania.cristescu@mcgill.ca
Alison M. Derry: **0000-0001-5768-8027**   derry.alison@uqam.ca

Data availability statement

Data and codes will be available upon manuscript acceptance

Funding statement

This research was funded by a Fonds de Recherche du Québec—Nature et Technologie (FRQNT) team grant to AMD, MEC, and DJF, a grant from the Natural Sciences and Engineering Research Council of Canada (NSERC Strategic Project grant, AMD, DJF), and a NSERC discovery grant to AMD.

Ethics approval statement

All field work and fish handling protocols were approved in permits issued by Parks Canada, Banff-Yoho and Kootenay National Parks (KooNP- 2017- 24,999), and the Concordia University Animal Ethics Committee.

Conflict of interest disclosure

There are no competing interests to declare

Abstract

Anthropogenic activities have led to an unprecedented crisis in freshwater biodiversity loss. The capacity to monitor the abundance of wild populations is critical to conserving biodiversity, but conventional physical specimen collection methods are invasive, costly, and labour-intensive. Environmental DNA (eDNA) offers a promising alternative, being easy to sample, with studies under controlled laboratory conditions showing consistent correlations between eDNA concentration and abundance. However, applying eDNA to monitor abundance remains contentious, as eDNA particle dynamics and the ecology of eDNA production can decouple this relationship in natural ecosystems. To address this, we integrated bioenergetics and mass-balance frameworks to relate eDNA concentrations to freshwater fish population *abundance* estimated through conventional mark-recapture in Brook Trout (*Salvelinus fontinalis*) across nine Rocky Mountains lakes, five of which underwent size-selectively harvest over two years. Our integrated framework improved the variance explained in eDNA concentrations from 24% to 71%. The integrated model accurately distinguished most abundance estimates across populations and sampling periods, detecting both natural and harvest-induced reductions in abundance within several populations. This study is the first to empirically integrate the DNA production ecology and particle dynamics, demonstrating that properly accounting for 'the ecology of eDNA' enables rapid, accurate abundance quantification. We also discuss future research and applications of eDNA in light of these results.

Significance statement

Addressing the unprecedented freshwater biodiversity crisis requires cost- and time-effective

methods for estimating abundance and biomass. Our study empirically demonstrates the potential

of eDNA for rapid and accurate assessment of abundance in natural populations by integrating

hydrology, chemistry and bioenergetics into a mass-balance modelling framework. Through a

replicated whole-ecosystem experiment, we show that eDNA not only detects differences in fish

abundance and biomass across populations, but also captures natural and experimentally induced

abundance changes within populations. Our modeling approach enables eDNA to generate

abundance and biomass estimates comparable in precision and accuracy to high-quality mark-

recapture methods.

Introduction

Anthropogenic activities have triggered an unparalleled biodiversity crisis in freshwater ecosystems (1). Conserving remaining biodiversity, remediating degraded ecosystems, and halting the decline of endangered populations demand rigorous monitoring and proactive aquatic ecosystem management (2). However, freshwater ecosystems are notoriously difficult to monitor, with the decline in freshwater fish populations, often referred to as an 'invisible collapse' (3, 4). Monitoring the abundance and biomass of species in aquatic ecosystems can be particularly challenging, due to its labour- and resource-intensive nature. These challenges are magnified at larger geographic scale, were obtaining such data for remote or large freshwater waterbodies can be exceedingly difficult if not impossible. As a result, the availability of such critical data for conservation and management is often severely limited.

The analysis of environmental DNA (eDNA), which refers to DNA particles found in environmental mediums like air, soil, and water (5), has revolutionised biomonitoring in natural ecosystems. The analysis of eDNA has proven highly effective for quantifying ecosystem-level biodiversity as well as detecting cryptic, rare, and traditionally difficult-to-detect species in freshwater ecosystems (6, 7). However, numerous studies have demonstrated consistent and significant positive correlations with the concentration of eDNA in an aquatic environment and the abundance or biomass of organisms within that environment, demonstrating its strong potential for additionally providing estimates of population abundance and/or biomass in natural ecosystems (8, 9). Yet, the biological processes involved in the production of eDNA production are complex and, as a physical particle, numerous environmental parameters can impact its degradation and transportation in natural ecosystems (10, 11). Collectively, these processes can decouple the relationships between the concentration of eDNA in natural ecosystems and organism

abundance/biomass (12). This can be readily observed in the discrepancy between the strong correlation eDNA exhibits with abundance under controlled laboratory conditions relative to the moderate correlations observed in studies conducted within natural ecosystems (9).

Our understanding of the behaviour and dynamics of eDNA particles in natural ecosystems is constantly growing and improving through rigorous research occurring in labs across the globe. The decay rate of eDNA, for example, can depend on complex interactions among abiotic conditions (e.g., temperature, pH and UV), biotic conditions (e.g., bacterial activity), and eDNA properties (amplicon and particle size) (13–18). In both lentic and lotic systems, eDNA particles move within water masses at different rates, which can additionally affect their deposition. The concentration measured at one place can thus be influenced by eDNA production upstream and/or its movement within currents or within thermocline layers (19–22).

Metabolic and bioenergetic theory has also recently been integrated as a framework to relate eDNA production to population density and biomass by accounting for the physiological processes involved in its production (23). Metabolic processes, for example, scale allometrically with organism size (24, 25). The strong link between eDNA production and organism metabolism implies that eDNA production likely scales allometrically within and across species (23, 26). Similarly, temperature has long been recognized as having a strong impact upon physiological rates within poikilothermic fishes (27), which similarly extends to eDNA production (16, 28). As a result, eDNA production can be modelled as a function of allometrically scaled organismal mass weighted with a temperature-dependent function to adjust for the impact of the metabolic rate of the organism on eDNA production (12, 26, 29, 30), e.g.:

$$eDNA\ production = I_0 \sum_{i=1}^{N} (M_i^b) \times f(T) \qquad (1)$$

Where I_0 is the eDNA production coefficient, N is the number of individuals in the

population, M_i is the mass of the i[th] individual, b is the allometric scaling coefficient, and $f(T)$ is

the temperature-dependent function, which considers the maximal metabolic rate of the organism

at a specific temperature (12).

A substantial body of work describes the complex dynamics of eDNA production,

degradation, and transport, all of which interact to influence eDNA concentrations in natural

ecosystems. Integrating across these processes provides an understanding of particle dynamics that

affect particle input and output rates; as a result, the application of mass-balance approaches are

ideally suited to modelling eDNA steady-state concentrations, in a similar manner as classic models

used for estimating phosphorus concentration (16, 31, 32). If we assume the concentration of eDNA

to be constant in time, we can state:

$$eDNA_{produced} + eDNA_{imported} = eDNA_{degraded} + eDNA_{sedimented} + eDNA_{exported} \quad (2)$$

Where the eDNA production rate is given by the metabolic/bioenergetics function described above

(equation 1), and eDNA imported describes the rate eDNA enters the system from up-stream

sources. The rate at which eDNA is degraded is typically described by an exponential decay curve

as a function of temperature, pH, and other factors that can influence eDNA degradation rate, with

'k' being the degradation constant (e.g. Kagzi et al., 2022). No research has yet examined eDNA

sedimentation rates independent of particle decay, but sedimentation of particles in the water are

also typically modeled using exponential decay curves (31). The rate at which eDNA is exported

(i.e., transported) refers to the amount of eDNA exiting by the outflow of a water body, which can

be calculated as the outflow times the concentration at the surface (31). By developing equation 2

and isolating eDNA concentration, the formula can be expressed as:

$$134 \qquad [eDNA]_{water} = \frac{W}{Q + (k_{deg} + k_{sed})V} \qquad (3)$$

Where *[eDNA]$_{water}$* is the eDNA concentration in the water, *W* is the average eDNA production

(equation 1) plus the average eDNA importation from upstream, *V* is the volume of the water body,

Q is the water flow rate, *K$_{deg}$* is the degradation constant at these specific environmental conditions,

and *K$_{sed}$* is the sedimentation rate.

Applying a modeling technique that integrates both bioenergetics and mass balance

approaches to eDNA concentrations to estimate total population abundance and biomass in natural

ecosystems remains unexplored but holds considerable potential for improving the capacity to link

eDNA concentrations observed in natural ecosystems to wild population abundance. With refined

quantitative knowledge on eDNA production, decay, and transport rates, eDNA could become a

cost-effective tool for species quantification in aquatic systems (12, 34). To date, however, no study

has integrated across all these processes to generate total population abundance/biomass estimates

in natural ecosystems.

We investigated the application of an integrated bioenergetics and mass-balance approach

to estimate population abundance from eDNA concentrations observed in natural lentic

ecosystems, aiming to: i) differentiate total abundance *across* populations; and ii) detect changes

in abundance *within* populations. To accomplish these objectives, we quantified Brook Trout

(*Salvelinus fontinalis*) eDNA concentrations and total abundance (using mark-recapture

techniques) in nine Rocky Mountain lakes (CA) across two seasons for two years, while

simultaneously conducting an annual experimental size-selective harvest in five of the nine lakes

to manipulate total abundance and population density across the study period. We also investigated

the additional power of an integrated bioenergetics and mass-balance approach in explaining

observed eDNA concentrations relative to approaches previously taken in the literature, including

i) correlating eDNA concentrations as a linear function of population abundance; and ii) correlating

eDNA concentrations as a function of allometrically scaled population biomass. We predicted that

taking an integrated bioenergetic and mass-balance approach that accounts for the processes

involved in eDNA production, degradation, and transportation would increase the power of eDNA

to detect changes in the abundance and biomass of Brook Trout across space and time.

Methods

Study system

Our study included nine headwater mountain lakes in the Canadian Rockies, located within

Banff, Yoho, and Kootenay National Parks (Canada; Fig. 1). The nine study lakes contain fish

communities largely dominated by Brook Trout (*Salvelinus fontinalis*) that were introduced to the

lakes between the 1920s and the 1970s and are now invasive in the region (35). Six of the lakes

were solely inhabited by Brook Trout, while the three other had fish communities also containing

one to five other species. Five of the nine study lakes were subjected to a size-selective harvest

treatment where approximatively 64 % of each population was removed in the Fall of each year

from 2017 to 2019, targeting the largest-bodied individuals for harvest. At the beginning of the

experiment (2017), the Brook Trout populations differed in density and population size structure

between lakes, and the harvesting treatment was successful at inducing changes in population

demography over the duration of the three-year experimental manipulation (36). The other four

study lakes were monitored as references in which Brook Trout populations were not directly

manipulated (Appendix 1). All lakes are small, cold, oligotrophic headwater lakes, but they span a

1000 m elevation gradient over which there are differences between lakes in temperature, pH, and

nutrient concentrations (Appendix 1; Trépanier-Leroux et al., 2023).

Field sampling of eDNA

Environmental DNA samples were collected each year between 2018 and 2019 for two time

periods: i) in the summer before harvesting (June/July); and ii) in the summer/fall following harvest

activities. A systematic stratified survey design was used to collect eDNA samples, in which 4

samples each were collected from the pelagic and littoral zones of each study lake. In the littoral

zones, eDNA sample sites were distributed equidistantly around the perimeter of the lake. Littoral

samples were collected approximatively 1-3 m from the shore at a minimum depth of 30 cm from

the surface, but approximately 15 cm above the lake bottom. In the pelagic zones, four samples

were collected at equidistant intervals across a transect that followed the longest axis of the lake

and went through the lake center (defined as the midpoint of the longest axis). Pelagic samples

were collected at a depth of 0.5 to 1 m from the surface. Water samples were collected with 700

ml Whirl-Pak™ bags from a canoe assigned to sample a single specific lake, or from an inflatable

kayak, which was decontaminated 48 hours prior with a 2% regular strength household bleach

solution for 15 min (including paddle and lifejacket) and left to dry in the sun.

Water samples were immediately filtered on the lakeshore on a water filtering manifold

(Wildco, Florida, USA) that was decontaminated with a 20-30% bleach solution 2-12 h prior to

sampling. Manifold components were stored individually in plastic zippered bags after bleaching

and during transportation. All samples were stored in the shade prior to sampling, and all filtering

was conducted in the shade under a tarp. One litre of water from each of the eight water sampling

sites per lake was filtered through a 0.7 µm glass fibre filter (GE Healthcare Life Sciences, Ontario,

Canada) using a vacuum hand pump (Soil Moisture, California, USA). Vacuum pumps were

decontaminated between lakes by wiping with a 30% bleach solution and resting for 10 minutes,

after which they were wiped with distilled water to remove residual bleach. All littoral samples

were filtered on one manifold, and all pelagic samples were filtered on another to prevent cross-

contamination between samples from different lake zones. As a negative control, 500 mL of

distilled water was filtered prior to filtering lake samples. After filtration, filters were folded using

metal forceps (decontaminated using a 30% bleach solution for 15 minutes) and stored in a sterile

2 mL microcentrifuge tube containing 700 µL of ATL buffer that was then individually placed in

a plastic zippered bag and stored in a cooler bag that was decontaminated by wiping with a 30%

bleach solution. If a filter became clogged (i.e. < 1 L of water was filtered), the final volume of

water filtered was recorded and the sample was then stored in buffer. Samples were then

transported in a cooler bag with bleached frozen gel packs to Kootenay Crossing (BC, Canada),

where they were placed in a second zippered plastic bag and stored in a -20 ℃ freezer. At the end

of the sampling season, all samples were transported to Montreal on dry ice and placed in a -80 ℃

freezer until extraction. All protocols for contamination prevention in the field, as well as during

transport and storage can be found in Appendix 2.

Laboratory analysis of eDNA

DNA was extracted from each filter was extracted using Qiagen DNeasy Blood and

Tissue[TM] kit and Qiashredder[TM] columns following a modified extraction protocol (see Yates et al.

(2021a) for more details). Extracted DNA was eluted into 130 µL of AE buffer and stored in a

clean -20 °C freezer solely dedicated to the storage of extracted eDNA product (i.e. no post-PCR

products or tissue samples). An extraction blank of 700 µL of ATL buffer was included in all

batches as a control. Methods for contamination prevention are detailed in Supplementary

Information (Appendix 2).

Brook Trout eDNA concentration was quantified using quantitative PCR (qPCR) with the

cyt b TaqMan minor groove assay (BRK2) described in Wilcox et al. (2013). For details on PCR

reaction preparation, component concentrations, standard curve preparation, and cycling

conditions, see Yates et al., (2021a). All samples were run in triplicate in a 20 µL reaction volume

with 5 µL of template DNA on a Stratagene MX 3000P thermal cycler. Reactions were spiked with

an internal positive control (IPC) to test for inhibition, which was defined as a $1 > Ct$ shift in the

IPC relative to a reaction containing 5 µL of nuclease-free ultrapure water. If inhibition was

detected, samples were diluted down to 3 µL and re-run. Brook Trout eDNA concentrations at each

site were calculated by averaging replicate values, and final copy number data was converted to

eDNA concentration per L of water filtered (copies/L). Standard curve R^2 values for the qPCR

plates ranged from 0.98-0.99, and reaction efficiency varied between 83.1 and 98.6%. The

efficiency of two of the 11 plates was suboptimal ($< 85\%$). However, data generated from these

two plates were retained, as their inclusion did not impact results (Appendix 3). Samples associated

with negative controls that contained more than 5 copies/rx (on average) were removed from the

analysis due to contamination concerns (Cobb Lake samples from fall 2019). The majority of

samples collected across lakes per sampling period (26/36 sampling events) showed no

contamination, while the remaining contamination in field controls was low enough to not impact

the estimated lake eDNA concentrations.

Brook Trout mark-recapture and population size structure estimation

Mark-recapture studies of Brook Trout were conducted in all lakes during May and June in

both 2018 and 2019. The exception was Cobb Lake where, in addition to mark-recapture sampling

in May/June, a series of isolated marking events continued until September for both years. Fish

were captured using a combination of fyke nets, angling, and backpack electrofishing. Large (1 m

hoop diameter, 2 cm mesh) and small (0.7 m hoop diameter and 0.8 cm mesh) fyke nets were distributed around the perimeter of lakes with the lead attached to shore and the end of the trap facing the center of the lake. Nets were checked daily. Angling was used to supplement Brook Trout capture efforts at sites where fyke catchability was low (predominantly Cobb Lake). Marks were also assigned to Brook Trout captured by electrofishing the shore and inlets/outlets of lakes.

Captured Brook Trout were anesthetized using clove oil and measured for fork length ($\pm$ 1 mm) and mass ($\pm$ 0.1 g). Any unmarked Brook Trout were gastrically tagged with a BioMark HPT8 pre-loaded Passive Integrated Transponder (PIT) tag (Boise, Idaho, USA). Only Brook Trout greater than or equal to 80 mm were tagged to reduce tagging mortality. Other species, when present, were not tagged. The tag number of any recaptured Brook Trout was recorded. Recovered fish were released in the center of the lake to standardize release location and promote mixing. Marking ceased once recapture ratios approached twenty five percent for several consecutive days to standardize marking efforts across all populations and to ensure that enough Brook Trout were tagged to facilitate estimation of census size (Nc) with credible intervals within 10% to 25% of true values, following general methodologies reviewed in Krebs (2009).

To obtain representative snapshots of the size structure of each Brook Trout population before and following harvest, population size structure estimates were conducted in August 2018 and 2019. The exception was Cobb Lake where size structure assessments continued to October 12th. During mark-recapture, fyke-nets were deployed in littoral zone areas extending to the centre of the lake and, as a result, size-structure assessments obtained from fyke nets may be more biased towards small-medium bodied individuals that tend to prefer littoral habitats (41). To obtain a relatively unbiased estimate of population size structure, Brook Trout were captured in large and small sinking mixed mesh gillnets with clear monofilament distributed throughout the lake. Large

mixed-mesh gillnets were 15.6 m long, 1.8 m deep and had an equal area of 64-51-89-38-76 mm

mesh panels. Small mixed-mesh gillnets were 12.5 meters long, 1.8 meters deep, and consisted of

an equal area of 32-19-38-13-25 mm mesh panels. Such index nets are widely used in North

America for size structure assessments (42–46), as these tend to capture a representative size/age

structure of populations (47). Nets were checked daily and moved to different locations across the

lake, to capture a representative sample of Brook Trout in each lake. Sampling ceased when

approximately five to ten percent of the population was captured, apart from Cobb Lake where the

harvested fish were also included for size structure assessment such that approximately 71% of

individuals were used. Captured Brook Trout were euthanized with clove oil, PIT tags were

recorded, and length/mass data were collected as described for the marking period.

Schnabel population size estimates were used to estimate the abundance of Brook Trout in

a lake (48). All size structure assessment removals were pooled together into one final sampling

event for the population estimates which controlled for the removal of marks at large. Note that

population estimates only account for Brook Trout greater than the minimum tagging size (80 mm

fork length), which correspond to all individuals age 1 and older (36). All population estimates

were conducted in R (49) with the mrClosed function from the Fisheries Stock Assessment package

FSA (50). Credible intervals for Schnabel population estimates followed recommendations from

Seber (2002) as implemented in the FSA package.

Size-selective harvesting of Brook Trout

Populations at Cobb, Mud, Olive, Ross, and Temple lakes were harvested annually with

size- selective gillnets (25– 89 mm mesh) targeting the top 75% (but effectively harvesting the top

64% on average) of individual body sizes according to the size distributions estimated during stock

assessment (i.e. depleting the largest size classes sequentially until the only remaining individuals

were in the lowest 25% quartile of the original size structure). Details can be found in Matte et al.

(2023).

Lake temperature and its effect on the physiology of eDNA production

To adjust for maximal metabolic rate of fish at a given temperature, we used a temperature-

dependent function specific to Brook Trout published from previous empirical studies (29, 52).

Temperature data were constituted of both high frequency (15 min) surface temperature from data

loggers (Onset Hobo®; UA-002-64, Onset Computer Corporation, Cape Cod, Massachusetts) and

vertical profiles from a YSI multiparameter sonde (model 10,102,030; YSI Inc.). The dataloggers

were deployed at 0.5 m from the surface in June and retrieved in September. The YSI vertical

profiles were taken at the deepest point of the lake at the same time as eDNA sampling, except six

lakes at Fall 2019 where a YSI malfunction necessitated us to use data from the same time period

in 2018 as a proxy. There was approximately a +/- 2°C difference between fall 2018 and 2019 for

the lakes that had YSI data for both seasons, and as a result this likely had very little impact on

model estimates. When possible, mean surface temperature over the interval from two weeks

before to two weeks after each eDNA sampling event per lake was used as input in the temperature-

dependent function in the eDNA mass balance model. However, due to logistic constraints and

material failures, some lakes and sampling periods had temperature data for only a subset of the

time intervals or only the epilimnion mean temperature from the YSI vertical profiles. A table with

the lakes, dates, and temperature data can be found in the supplementary materials (Appendix 4).

Lake hydrology

To calculate the rate of lake outflow, lake watershed area was estimated using the flow

direction and flow accumulation hydrologic models from ArcGIS pro V.3.2. The input data were

15 m resolution Digital Elevation Models (DEM) for 7 out of the 9 study lakes. However, the 15

m resolution was too low to effectively determine the watershed for two of the lakes (Mud and

McNair). Fifteen m resolution DEM are publicly available form the Government of Canada and

under Open Government Licence (https://open.canada.ca/en/open-government-licence-canada).

The Mud and McNair lakes required high resolution data, but none were available for their area.

The Mud Lake watershed area was therefore estimated by process of elimination by estimating the

watershed area of the surrounding larger water courses. The remaining area that was not part of

any other watershed was assumed to be Mud Lake's watershed. We could not obtain watershed

area for McNair Lake. The lake outflow rate was estimated by multiplying each lake watershed

area with the flux per area of watershed from hydrological stations from the same area (Kootenay:

Kootenay River: 50.886944, -116.045833; Banff: Pipestone River: 51.433056, -116.174722;

Environment and Natural Resources Canada, 2024). The rate of outflow for each lake at each

sampling period was calculated as the mean outflow from two weeks before the sampling date to

two weeks after.

Lake volume and epilimnion volume

Detailed bathymetric maps and lake volume produced by high frequency sonar were

available for only four of our lakes (Temple, Olive, Helen, and Margaret lakes). For the remaining

five study lakes, we used older bathymetric maps from Parks Canada that were georeferenced and

digitised in arcGIS pro®. The depth between isobaths were interpolated using Bayesian Kriging

method. The epilimnion volume was taken as the input in the mass balance model (equation 5 and

8). Epilimnetic volume was chosen because eDNA samples were collected from the epilimnion of

each lake; this was done because most study lakes undergo thermal stratification during summer,

and because Brook Trout spend most of their time feeding and resting in surface waters (41).

Additionally, research has demonstrated that the transfer of eDNA across the thermocline is limited

during lake stratification (54). The epilimnion volume was calculated as the mean thermocline

depth times the lake surface area from the bathymetric maps. The mean thermocline depth was

calculated from two YSI® vertical temperature profiles per summer. For the study lakes that were

not thermally stratified (assessed from the YSI vertical temperature profile) at the time of eDNA

sampling period, total lake volume (estimated from the bathymetric maps) was used as input in

mass-balance models for that time point because eDNA signal is expected to become homogenous

throughout the water column during mixing (54).Unfortunately, the extent of thermal stratification

during the fall of 2019 could not be quantified due to the malfunction of the YSI. Lakes that were

mixed in the fall of 2018 were therefore also assumed to be mixing in fall 2019 except for Margaret

Lake, where stratification was very weak in fall 2018 and it was not fully mixed. As fall 2019 was

colder than fall 2018, we assumed that Margaret Lake was mixing in fall 2019; this increased model

fit for this lake.

Statistical analyses

Constructing and evaluating an integrated bioenergetics/mass-balance model

To obtain a representative mean eDNA concentration for each lake and sampling period,

the littoral and pelagic samples were weighted according to the littoral and pelagic zone area in

each lake. The littoral zone was defined as the area where the depth is bellow or equal to 2 m. To

parameterise the mass balance model (quation 3), a non-linear Bayesian model with informative

priors was used. The parameters estimated were I_0 (eDNA generation coefficient), b (allometric

scaling coefficient), and k (a combined degradation and sedimentation constant). Although

equation 3 has both the degradation constant k_{deg} and the sedimentation constant k_{sed}, the current

state of knowledge on eDNA degradation and sedimentation does not allow for a precise estimate

of the values for k_{deg} and k_{sed}, and so one "disappearance constant" k was parameterised in the

model as the sum of both (equation 4).

$$[eDNA]_{Lake} \sim \frac{I_0 \sum_{i=1}^{N}(M_i^b) \times f(T)}{Q + kV} \qquad (4)$$

The model was fitted with a gaussian distribution, four chains, 10 000 iterations, and *adapt_delta*

= 0.99. A lake specific k parameter was estimated to account for potential differences in

degradation rates between lakes, and a sampling period specific eDNA generation coefficient (I_0)

to account for season and sampling period effect on mean Brook Trout eDNA concentrations

observed across lakes (Gaudet-Boulay et al., 2023), either due to ecological differences in the

eDNA production coefficient (I_0) across seasons or potential (unintentional) methodological

differences in sampling. For all analysis, $\sum_{i=1}^{N}(M_i^b)$ was approximated using the Taylor expansion

for the first moment (55) such that equation 4 becomes:

$$[eDNA]_{Lake} \sim \frac{I_0 f(T) \times N(\bar{M} + \frac{\sigma_M^2}{2} b(b-1)\bar{M}^{b-2})}{Q + kV} \qquad (5)$$

Where $\bar{M}$ is the mean mass of the fish and σ_M^2 is the variance of the mass of the fish. The priors

used were:

$$\ln(I_0) \sim normal(10, 2)$$
$$b \sim beta(2.8, 1.2)$$
$$\ln(k) \sim normal(-0.7, 0.1)$$
$$Standard\ deviation_{K\ lake} \sim exponential(3)$$
$$\sigma \sim \theta + N^\alpha$$
$$\alpha \sim normal(0, 0.5),\ lb = 0$$
$$\theta \sim normal(0, 1),\ lb = 0$$

The prior on I_0 ensured posterior predictions were the correct order of magnitude, the prior on b

ensured values were between 0 and 1 with a higher probability density around 0.7, which is

approximately the *b* value estimated from Brook Trout bioenergetic models and expected based on

bioenergetic theory (29). The prior for *k* ensured values corresponding roughly to the expected

values based on Lamb et al. (2022). eDNA data were scaled to Kilocopies m^{-3} to avoid working

with very large numbers. The prior on sigma accounted for heteroscedasticity and increases

variance with total abundance (N), and α and θ have a lower bound (*lb*) at zero.

Comparing the mass balance eDNA approach to other existing models

Most previous studies examining relationships between eDNA concentrations and organism

abundance/biomass typically directly regress observed eDNA concentrations against estimates of

abundance and/or biomass (38). To evaluate the relative importance of accounting for both

bioenergetics and particle dynamics (i.e., mass balance modelling), we also fitted models in which

eDNA concentrations were directly related to both population biomass (equation 6) and

allometrically scaled biomass (equation 7, Yates et al. 2021a). Both models were fitted assuming

a Gaussian distribution:

391
$$[eDNA]_{Lake} \sim intercept + \beta \frac{N \times \bar{M}}{lake\ area\ (ha)} \tag{6}$$

392
$$[eDNA]_{Lake} \sim \frac{I_0 \sum_{i=1}^{N}(M_i^b)}{lake\ area\ (ha)} \approx \frac{I_0 \times N(\bar{M} + \frac{\sigma_M^2}{2} b(b-1)\bar{M}^{b-2})}{lake\ area\ (ha)} \tag{7}$$

393   With the priors:

394
$$
\begin{aligned}
\ln(I_0) &\sim normal(0,5)\\
b &\sim beta(2.8,1.2)\\
\sigma &\sim cauchy(0,5)
\end{aligned}
$$

395   The performance of the different models was compared using the bayes R^2 and the value of the

396   slope comparing the predicted and measured eDNA concentrations (the closer to one the better).

 Evaluating the efficacy of eDNA to estimate population abundance

We used our integrated bioenergetics/mass-balance approach to derive estimates of

population abundance to compare their precision and accuracy relative to estimates of abundance

derived from mark-recapture studies. Specifically, we are interested in evaluating the efficacy of

eDNA to both i) differentiate abundance *across* study populations; and ii) detect changes in

abundance *within* a population over time (i.e., population reductions due to harvesting). To

produce posterior estimates of total abundance for each of the populations during each sampling

period using the mass balance approach, *N* was isolated from equation 5 and the model was refitted

with the same priors. Additionally, season and sampling period can have a strong effect on mean

Brook Trout eDNA concentrations observed across lakes (Gaudet-Boulay et al., 2023), either due

to ecological differences in the eDNA production coefficient (I_0) across seasons or potential

(unintentional) methodological differences in sampling. A sampling-period specific I_0 was

therefore also fitted in the model. Equation 5 then becomes:

$$N \sim \frac{[eDNA]_{Lake} \times (Q + kV)}{I_0 f(T) \times \left(\bar{M} + \frac{\sigma_M^2}{2} b(b-1)\bar{M}^{b-2}\right)} \tag{8}$$

411   With the priors:

$$\ln(I_0) \sim normal(10, 2)$$
$$b \sim beta(2.8, 1.2)$$
$$\ln(k) \sim normal(-0.7, 0.1)$$
$$Standard\ deviation_{K\ lake} \sim exponential(3)$$
$$Standard\ deviation_{\log(I_0\ Sampling\ period)} \sim exponential(3)$$
$$\sigma \sim \theta + N^\alpha$$
$$\alpha \sim normal(0, 0.5),\ lb = 0$$
$$\theta \sim normal(0, 1),\ lb = 0$$

In this model, we also accounted for the measurement error on the mark-recapture total abundance

estimates (56). We also examined the efficacy of the integrated eDNA model to differentiate

population biomass, which can be estimated from the integrated eDNA model (equation 8) for each

population by multiplying the posterior distribution of N values by the mean mass of each

population. The performance of the model was assessed with the Bayesian R^2 and the number of

abundance estimates from eDNA that overlapped with mark-recapture (and their respective 95%

credible intervals).

Examining potential sources of bias

To assess the presence of systematic bias of the model, a Bayesian mixed model with non-

informative priors was fitted with the z-score of each eDNA prediction as the independent variable.

The fixed effects were mark-recapture population estimates, elevation, lake thermal mixing status

(yes = stratified or no = mixed), season (summer or fall), lake area and experimental treatment

(harvested or reference lake). We included interactions between thermal mixing status and

elevation, between season and treatment, and between season and elevation. Lake was included as

a random intercept to account for non-independence. This bias testing model was performed on the

outputs of the eDNA mass-balance models, including sampling period specific I_0.

Can our nine lakes calibrate a model to predict abundance in novel systems?

Due to repeated sampling within and across years, the model described in equation 8 can

estimate lake-specific degradation constants and sampling period specific I_0, and is thus well-suited

to monitoring abundance within and across this specific set of nine study lakes. However, this

model cannot be extended to estimate population abundance in a lake without pre-existing data

(e.g., where a lake specific k is unknown). To explore the efficacy of this approach to estimate

abundance in lakes (equation 8) without pre-existing baseline data, we additionally fitted two

models: i) a model with lake specific k but no sampling period specific I_0 to assess the importance

of accounting for local, lake-specific eDNA degradation rates (referred to as equation 8.1 in the

results) and ii) a model without a lake specific eDNA disappearance constant (k) nor sampling

period specific I_0 to assess how a generalizable model would perform as well as the importance of

a sampling period specific eDNA generation coefficient (referred to as equation 8.2 in results). The

performance of the different models was compared using Bayesian R^2 values. All analyses were

performed in *R* (49) using the *brms* package (version 2.18.0) (57).

Results

Fish abundance varied significantly between lakes with estimates ranging from 72 (46 –

117) to 3173 (2644 – 3965) individuals based on mark-recapture. Temporal variation in population

size was particularly pronounced in the size-selectively harvested lakes, where annual harvest

ranged from 46.8% to 92.7% of the population. Despite these high harvest levels, compensatory

effect allowed populations to partially recover between harvesting treatments (36). Natural

mortality also contributed to population changes. For instance, in Dog Lake population abundance

declined by 32% between the summers of 2018 and 2019, reflecting substantial non-harvest losses.

Comparing the mass balance eDNA approach to other existing models

The eDNA mass balance model explained more variability in eDNA concentration and was

more accurate than a linear regression with biomass or allometrically scaled biomass (Table 1, Fig.

2). While all three models had little bias, biomass alone explained only 24% of the variation in

predicted eDNA values (Bayes R^2). This increased to 44% after integrating the effect of allometry

on eDNA production (e.g., allometrically scaled mass). Integrating bioenergetics and mass-balance

modelling approaches, however, explained 66% of the variation in observed eDNA concentrations.

Evaluating the efficacy of eDNA to estimate population abundance

Total abundance estimates derived from eDNA concentration data (equation 8) effectively

differentiated populations across spatial and temporal scales (Fig. 3). The study populations

exhibited a substantial gradient in total abundance across lakes, with the smallest population

exhibiting a 32-fold difference in abundance relative to the largest population. At this scale, the

integrated eDNA model was very effective at differentiating abundance across populations (Fig 3).

Variations in abundance across populations were significantly greater than within

population changes, even in populations that exhibited substantial declines due to size selective

harvesting manipulations. Within a population, abundance levels were more consistent across

seasons and years (Fig. 3 and Fig. 4). Yet, even within populations, the accuracy and precision of

the integrated eDNA model was sufficient to detect changes across seasons and years in several of

the harvested lakes (e.g., McNair and Temple), as well as natural fluctuations in abundance across

control populations (e.g., Margaret and Dog lakes).

The majority (31/35 or 89%) of the total Brook Trout abundance estimates derived from

the eDNA mass balance model (equation 8) fell within the credible intervals for total abundance

estimated by mark-recapture (Fig. 3 and 4). Yet, five estimates showed no overlap between the

80% credible intervals for total abundance from both eDNA and mark-recapture, leaving 26 (74%)

estimates that exhibited a high correspondence. The integrated eDNA model similarly proved

effective at differentiating population biomass across space and time. The biomass estimates from

the integrated eDNA model were highly correlated with estimates derived from mark-recapture

(Bayes $R^2 = 0.80$) (Fig. 5) and, as with abundance, enabled the clear differentiation of biomass both

across and within populations (Fig. SM1).

Examining potential sources of bias

When testing for different bias of the eDNA total abundance estimates, there was no

significant systematic bias detected related to the total Brook Trout abundance nor lake size. There

was a non-significant tendency to generate larger estimates in lakes experiencing experimental

harvesting during the fall seasons (e.g., Ross and Mud at fall 2019) and smaller abundance

estimates in control lakes and some harvest lakes outside of harvest periods (e.g. Helen Lake at fall

2018 and 2019, Dog Lake at fall 2018, and Mud Lake at summer 2018) compared to mark-recapture

(Fig 4, Table 2, and Appendix 5). There was also a non-significant trend for the integrated eDNA

model to produce estimates of abundance lower than mark-recapture estimates for samples taken

during lakes that were experiencing thermal mixing, especially at high elevation. However, this

was largely driven by Helen Lake, for which total abundance estimates from the integrated eDNA

model were lower than mark-recapture while mixing occurred during fall 2018 and 2019.

Can our nine lakes calibrate a model to predict abundance in novel systems?

Incorporating a lake specific eDNA disappearance constant (k; the sum of the sedimentation

and degradation constant) significantly improved model performance, as demonstrated by the large

standard deviation on a general k estimated across lakes and the reduction in model performance

when removing lake specific k (equation 8.2)*,* with the Bayes R^2 dropping from 0.70 to 0.51 (Table

1 and Appendix 6). The effect of accounting for a sampling period specific eDNA generation

coefficient (I_0; equation 8.1) also reduced model performance, but to a substantially lesser extent

than including a lake-specific k. Removing the sampling period specific eDNA generation

coefficient (I_0) decreased the Bayes R^2 value from 0.72 to 0.70. This mainly impacted predictions

based on the summer 2018 samples, making the temporal trends in McNair and Olive less

congruent to the mark-recapture (Table 1 and Fig SM2). Brook Trout abundances predicted from

the eDNA mass balance model that lacked a sampling period specific I_0 (equation 8.1) also fell

within the 95% credible intervals for total abundance estimated by mark-recapture for 31/35

estimates (Appendix 5).

Discussion

Despite the increasing demand for abundance data to understand and manage ecosystems

(34), the application of eDNA to quantify the abundance or biomass of aquatic organisms remains

a controversial topic (8, 9, 58). This stems from the complex interplay of biotic and abiotic factors

that influence eDNA production, degradation, and transportation in natural aquatic environments

(12), which are considerably more variable across space and time in natural ecosystems relative to

controlled laboratory settings (9).

Herein we demonstrate, that eDNA can provide estimates of organism abundance and

biomass comparable in precision and accuracy to mark-recapture. However, achieving this

accuracy required accounting for biotic and abiotic factors that affect its distribution in natural

ecosystems, namely the ecology of its production and the dynamics of the physical eDNA particles.

Applying our integrated bioenergetics and mass balance approach to model quantitative eDNA

signals greatly improved the relationship between eDNA concentration and total abundance

estimates in the study ecosystems, compared to a basic linear correlation between either eDNA and

biomass or eDNA and allometrically scaled biomass. Our integrated approach allowed for the

accurate differentiation of population abundance and biomass across study populations (i.e., space)

based on lake eDNA concentrations. Quantifying the concentration of eDNA within the study lakes

also facilitated the prediction or monitoring of temporal changes in fish abundance/biomass over

time within several populations.

While previous research demonstrated the influence of several factors on eDNA

concentration and dynamics, including transportation (7, 11, 22), degradation (11, 13, 15, 16),

temperature (28, 59), and fish metabolic rate (12, 23, 26), this study is the first to integrate mass

balance processes with bioenergetics to relate eDNA concentration to absolute population

abundance in natural ecosystems. We empirically demonstrate that eDNA could represent a time-

and cost-effective tool for generating total abundance estimates with sufficient precision to detect

changes in abundance by integrating relatively simple-to-collect environmental and ecological

data.

Evaluating and comparing the efficacy of eDNA to estimate population abundance

eDNA abundance estimates corresponded to mark-recapture abundance estimates in 31 out

of 35 cases when considering the 95% credible interval, and in 26 out of 35 cases when considering

the 80% credible intervals. Most discrepancies (6 out of 9) were likely due to issues associated

with our unique study design causing the violation of assumptions in our mass-balance modelling

(i.e. experimental harvesting), and/or the violation of assumptions associated with the mark-

recapture estimates for populations in the fall sampling periods (e.g., no natural mortality or

emigration).

There was a non-significant trend wherein estimates of abundance derived from eDNA

tended to be higher than for mark-recapture in harvested lakes in the fall. The time necessary for

eDNA concentrations to stabilize depends on eDNA degradation rate, which can be impacted by

amplicon size, temperature, and potentially UV radiation, pH, and microbial activity (14). The

values estimated by the eDNA mass balance model for the k degradation parameter (the eDNA

"disappearance rate", or the sum of the sedimentation and degradation rates) corresponded to

expectations based on previous controlled experiments. Notably, a meta-analysis reported

degradation constant values between 0–0.2 h-1 or 0 – 2.5 day-1 in freshwater ecosystems (Lamb

et al., 2022), and we obtained an average k value estimate of 0.49 (0.41 – 0.6) day-1. Based on the

eDNA disappearance constant k estimated by the model, the outflow rate, and the summer eDNA

concentration in each lake and each year, we estimated that three to twenty days would be necessary

for eDNA turnover in this study system (Appendix 8). For logistical reasons (i.e., the harvesting

treatment took a substantial amount of time and extended late in the season), there was generally

less than seven days between the end of harvesting and eDNA sampling within the harvested lakes

which is less than the turnover rate in most cases. A tendency for the integrated eDNA model to

overestimate fish abundance following harvesting could also be explained by many eDNA copies

released during harvesting. Carcasses can be a significant source of eDNA (60) and often remained

in nets for 24 - 48h, thus increasing the initial amount of eDNA and time before stabilisation back

to a steady state concentration.

There were also four cases where eDNA estimates were lower than mark-recapture

estimates during fall sampling periods when looking at 80% credible intervals (two in Helen Lake

and two in Dog Lake). An instinctive response may be to attribute the discrepancies to error in the

integrated eDNA model. However, the discrepancies between the eDNA-derived and mark-

recapture abundance estimates could be attributable to violations of the assumptions of the mark-

recapture methodology employed. In Helen Lake, the discrepancies could be attributed to mixing,

but no other lakes experiencing mixing exhibited an underestimation of Brook Trout abundance

from the integrated eDNA model. Alternatively, Helen Lake also has an outflow stream where the

Brook Trout are known to spawn. The lower estimates of total fall abundance from the integrated

eDNA model in Helen Lake could therefore be due to fish leaving the lake to spawn, especially

since spawning is known to occur earlier in high-altitude alpine lakes such as Helen Lake (61).

Similarly, mark-recapture estimates demonstrated that Dog Lake exhibited a substantial natural

decline in abundance between 2018 and 2019. While mortality was assumed to be negligeable

between summer and fall, we do not actually know when this mortality event occurred because

mark-recapture studies occurred in the spring and summer of each year. Results from the eDNA

model could imply that the natural mortality event may have occurred between the Summar and

fall sampling periods both in 2018 and 2019, as the estimate of abundance derived from the

integrated eDNA model in fall 2018 corresponds well to the mark-recapture estimates of abundance

obtained during summer 2019. In these cases, eDNA may more accurately reflect the abundance

of fish in the study lakes than mark recapture estimates, for which we assumed there was no natural

mortality nor emigration for the fall estimates.

Generalisation of the integrated bioenergetic and mass balance model

While 25% (9 out of 35) of the estimates from the integrated bioenergetic and mass balance

model exhibit some discrepancy between eDNA-derived and mark-recapture estimates of

abundance when looking at the 80% CI, only 3 out of 35 (9%) of the discrepancies lack potential

explanations, which is reduced to 1 of 35 (3%) when looking at the 95% CI. Additionally, although

discrepancies between eDNA and mark-recapture derived estimates in abundance existed, these

discrepancies still reflect a relatively high level of precision *across* study populations, for which

eDNA permitted broad differentiation across high- and low-abundance populations. Although we

advise caution, these results imply that eDNA can potentially be used to reliably estimate

abundance and biomass in nature across populations, especially given that mark-recapture

estimates themselves can exhibit biases or fail when assumptions are violated. Our study design

does not allow us to identify which estimate is closer to the 'true' population size, and these issues

would only be compounded in larger natural ecosystems. Overall, we detected no systematic bias

in eDNA-derived estimates related to lake size, mixing, elevation, and population size. The

integrated bioenergetics and mass balance approach that we applied to the analysis of eDNA

concentrations produced mostly accurate population abundance estimates in small to medium-sized

oligotrophic lakes. With further validation, such models have the potential to be extended to other

aquatic ecosystems.

Accounting for lake mixing was an important component of the mass-balance modelling

because the volume of the waterbody is a critical parameter in mass-balance modelling approaches.

Previous research has demonstrated that eDNA disperses throughout a lake and is roughly

homogenous during thermal mixing (54). During lake stratification, we found that the volume of

the epilimnion is the appropriate volume input data, likely because Brook Trout remains close to

the surface during summer (41). In contrast, when a lake is undergoing thermal mixing, the volume

of the entire lake is the appropriate volume input data in the mass-balance model as the signal is

diluted in the entire lake. We detected no systematic bias related to lake thermal mixing when data

for Helen Lake was removed from the model; in all other lakes mixing explained no discrepancies

between eDNA and mark-recapture abundance estimates. The absence of overestimation of Brook

Trout abundance during mixing implies that the resuspension of sedimentary DNA during lake

mixing likely had a negligible impact on abundance estimates, despite sediment representing a

potentially potent source of eDNA (62). However, the amount of eDNA stored within sediments

and how it exchanges with the water column also depends on an organism's spatial distribution, as

well as biotic and abiotic conditions in water and at the water-sediment interface (63). Sedimentary

DNA re-suspension may be more important in other types of lakes or for other species, particularly

for systems inhabited by benthivorous fish species that disturb sediment during feeding (e.g. Carp;

Parkos et al., 2003).

Future directions: Can our nine lakes calibrate a model to predict abundance in novel systems?

Our perspective on the future application of eDNA to estimate population abundance and

biomass is not to replace conventional methods, but rather to complement and/or improve upon

estimates generated by them. Notably, the integrated bioenergetic and mass-balance model

developed herein requires population size structure data which can only be obtained using mixed

mesh index gillnetting. These methods, often used in state and provincial monitoring programs to

estimate relative abundance based on CPUE (e.g. Zimmerman & Palo, 2011), require fewer

resources than mark-recapture techniques. To broadly apply eDNA to estimate abundance using

the methods and models developed herein also requires the establishment of a 'calibration' system,

in which abundance estimated using conventional methods on a subset of ecosystems are used to

calibrate an eDNA/abundance model. Once calibrated, this model can be applied to estimate

abundance in other systems or time points lacking conventional estimates, by sampling eDNA and

the required physicochemical parameters.

Our findings imply that the capacity to identify local predictors of k would be particularly

critical for such applications, as baseline data to estimate k rates within an individual system

substantially improved our model performance. To increase the reliability of abundance estimates

in freshwater systems derived from eDNA, further research is needed on eDNA dynamics in lentic

systems, with a particular focus on drivers of sedimentation and degradation rates. A quantitative

understanding of these interactions would allow us to better predict sedimentation and degradation

rates specific to local and temporal conditions, which would in turn likely improve the integrated

eDNA bioenergetic and mass-balance model performance as such dynamics could be integrated

into the modelling process. Additionally, such knowledge would facilitate the estimation of the

disappearance k parameter for lakes and time points outside of any putative "calibration system"

lakes simply by measuring the relevant environmental variables.

A growing number of individual studies on eDNA degradation have found that eDNA

degradation is the result of a complex combination of many variables such as pH, temperature, UV

radiation and microbial activity, and eDNA properties such as amplicon size and whether the

targeted gene is nuclear or mitochondrial (13, 15–17, 66). Although temperature has emerged as a

consistently impactful parameter affecting eDNA degradation rates, the importance of other

parameters such as UV, microbial activity, eDNA source, and amplicon size remains to be

confirmed (Lamb et al. 2022, Jo and Minamoto 2021, Saito and Doi 2021). Collectively, we are

still far from producing effective eDNA degradation models, as few studies to date have examined

the interaction between these different environmental variables and eDNA properties (especially

in natural ecosystems) and even fewer have examined eDNA sedimentation. However, in the

absence of controlled empirical studies, further large-scale observational studies in natural

ecosystems could facilitate the identification of drivers of local eDNA decay rates. Experiments

studying degradation dynamics in natural ecosystems are thus crucial to facilitate and improve the

adoption of eDNA to estimate abundance in natural ecosystems.

Theoretical and empirical evidence suggests that eDNA production rate is related to fish

metabolic rate (23, 38, 67, 68). However, to our knowledge, only one study to date has

experimentally verified this relationship under controlled empirical conditions and found a positive

correlation between fish activity rate, energy use, and eDNA shedding rate (68). The mass balance

model assumes eDNA production scales with body mass and is influenced by temperature-

dependent metabolic rates in poikilothermic Brook Trout (Hervé et al., 2023; Thalinger et al., 2021;

Yates, Cristescu, et al., 2021; Yates et al., 2023). We used a temperature-dependent metabolic

function (Hartman & Cox, 2008) to adjust for temperature effects on eDNA, though this may not

fully reflect eDNA production dynamics. While many studies have documented an effect of

temperature on eDNA production (e.g., Caza-Allard et al., 2022; T. Jo et al., 2019), whether this

effect directly reflects metabolic responses to temperature remains to be tested. Investigating this

could improve model assumptions and performance.

Our study additionally included mostly only headwater lakes, so eDNA inputs from

upstream were considered equal to zero. However, this is rarely the case within a watershed. To

add eDNA transportation into the lake within the integrated model, data on both the inflow rate

and the average eDNA concentration in the inflow are required. In theory, these data can be

implemented in equation 4 (31) such that:

674
$$[eDNA]_{Lake} \sim \frac{I_0 \sum_{i=1}^{N}(M_i^b) \times f(T) + Q_{in}[inflow]}{Q_{out} + kV} \tag{9}$$

Where Q_{in} is the inflow rate, Q_{out} is the outflow rate, and [*inflow*] is the mean eDNA concentration

of the inflow.

Conclusion

Our study proposes an integrated bioenergetics and mass-balance approach to relate eDNA

concentration to abundance estimation. Our integrated eDNA model performed well at estimating

abundance, facilitating total abundance and biomass estimates comparable to conventional mark-

recapture methodologies, often considered a 'gold-standard' for population estimates. We show

that eDNA can detect both spatial and temporal differences in fish abundance, provided the

equilibrium assumptions of the model are respected. To respect these assumptions, we recommend

avoiding sampling ten to twenty days after known fluctuations in abundance (e.g. harvest),

depending on the eDNA degradation rate and outflow flux. Overall, results of this study represent

an important advancement in the use of eDNA to quantify total abundance, with strong implications

for the application of eDNA for abundance estimates in a context of ecosystem management. The

quantitative analysis of eDNA, applied with an integrated bioenergetics and mass-balance

approach, could provide a cost-effective, precise, and accurate tool for managers and researchers,

in an era where the acquisition of population abundance data is increasingly important for

biodiversity management (e.g. Canals et al., 2021; Fonseca et al., 2023; Kelly et al., 2024).

Acknowledgements

We thank Parks Canada for logistic support, Andrew Macdonald for statistical advice, Maria Gheta

for the help with the Taylor expansions, Yves Prairie for math validation, Jérôme Lemelin for the

map, and Mélanie Desrochers for GIS advice. We also express our gratitude to Louis Astorg, Dylan

Glaser, Brent Brooks, Shannon Clarke, Jean-Michel Matte, Justin Budyk, Mélia Lagacé, Mathilde

Salamon, Sydney Urschel, and Ahmad Yaghi for field work. This research was funded by a Fonds

de Recherche du Québec-Nature et Technologie (FRQNT) team grant to AMD, MEC, and DJF, a

grant from the Natural Sciences and Engineering Research Council of Canada (NSERC Strategic

Project grant, AMD, DJF), and a NSERC Discovery grant to AMD.

Authors contribution

JB designed the models and performed the analysis. JB and MCY contributed to the writing of the

manuscript. MCY and JB collected the eDNA samples. MCY extracted the eDNA samples and

conducted laboratory analyses. All authors contributed to the study design and subsequent edits to

the manuscript.

References

1. J. S. Albert, *et al.*, Scientists' warning to humanity on the freshwater biodiversity crisis. *Ambio* **50**, 85–94 (2021).

2. L. T. Ho, P. L. M. Goethals, Opportunities and Challenges for the Sustainability of Lakes and Reservoirs in Relation to the Sustainable Development Goals (SDGs). *Water (Basel)* **11**, 1462 (2019).

3. J. R. Post, *et al.*, Canada's Recreational Fisheries: The Invisible Collapse? *Fisheries (Bethesda)* **27**, 6–17 (2002).

4. A. Howarth, V. M. Nguyen, S. J. Cooke, Managing Canadian freshwater fisheries: persistent challenges and emerging opportunities. *Canadian Journal of Fisheries and Aquatic Sciences* **80**, 1436–1455 (2023).

5. G. F. Ficetola, C. Miaud, F. Pompanon, P. Taberlet, Species detection using environmental DNA from water samples. *Biol Lett* **4**, 423–425 (2008).

6. C. L. Jerde, Can we manage fisheries with the inherent uncertainty from eDNA? *J Fish Biol* **98**, 341353 (2019).

7. A. J. Sepulveda, N. M. Nelson, C. L. Jerde, G. Luikart, Are Environmental DNA Methods Ready for Aquatic Invasive Species Management? *Trends Ecol Evol* **35**, 668–678 (2020).

8.   M. L. Rourke, *et al.*, Environmental DNA (eDNA) as a tool for assessing fish biomass:

A review of approaches and future considerations for resource surveys. *Environmental*

DNA **4**, 9–33 (2022).

9.   M. C. Yates, D. J. Fraser, A. M. Derry, Meta-analysis supports further refinement of

eDNA for monitoring aquatic species-specific abundance in nature. *Environmental*

DNA **1**, 5–13 (2019).

10.   K. A. Stewart, Understanding the effects of biotic and abiotic factors on sources of

aquatic environmental DNA. *Biodivers Conserv* **28**, 983–1001 (2019).

11.   L. Carraro, H. Hartikainen, J. Jokela, E. Bertuzzo, A. Rinaldo, Estimating species

distribution and abundance in river networks using environmental DNA. *Proceedings*

of the National Academy of Sciences **115**, 11724–11729 (2018).

12.   M. C. Yates, M. E. Cristescu, A. M. Derry, Integrating physiology and environmental

dynamics to operationalize environmental DNA (eDNA) as a means to monitor

freshwater   macro-organism   abundance.   *Mol   Ecol*   1–20   (2021).

https://doi.org/10.1111/mec.16202.

13.   K. M. Strickler, A. K. Fremier, C. S. Goldberg, Quantifying effects of UV-B,

temperature, and pH on eDNA degradation in aquatic microcosms. *Biol Conserv* **183**,

85–92 (2015).

14.   P. D. Lamb, V. G. Fonseca, D. L. Maxwell, C. C. Nnanatu, Systematic review and

meta-analysis: Water type and temperature affect environmental DNA decay. *Mol Ecol*

Resour **22**, 2494–2505 (2022).

15.   T. Jo, T. Minamoto, Complex interactions between environmental DNA (eDNA) state
750   and water chemistries on eDNA persistence suggested by meta-analyses. *Mol Ecol*
751   *Resour* **21**, 1490–1503 (2021).

752   16.   T. Jo, H. Murakami, S. Yamamoto, R. Masuda, T. Minamoto, Effect of water
753   temperature and fish biomass on environmental DNA shedding, degradation, and size
754   distribution. *Ecol Evol* **9**, 1135–1146 (2019).

755   17.   T. Saito, H. Doi, A Model and Simulation of the Influence of Temperature and
756   Amplicon Length on Environmental DNA Degradation Rates: A Meta-Analysis
757   Approach. *Front Ecol Evol* **9**, 1–8 (2021).

758   18.   P. F. P. Brandão-Dias, *et al.*, Particle size influences decay rates of environmental DNA
759   in aquatic systems. *Mol Ecol Resour* **23**, 756–770 (2023).

760   19.   J. E. Littlefair, M. D. Rennie, M. E. Cristescu, Environmental nucleic acids: a field-
761   based comparison for monitoring freshwater habitats using eDNA and eRNA. *bioRxiv*
762   2021.12.01.469845 (2021).

763   20.   S. F. Jane, *et al.*, Distance, flow and PCR inhibition: EDNA dynamics in two headwater
764   streams. *Mol Ecol Resour* **15**, 216–227 (2015).

765   21.   D. Pont, Predicting downstream transport distance of fish eDNA in lotic environments.
766   *Mol Ecol Resour* **24**, 1–14 (2024).

767   22.   A. J. Shogren, *et al.*, Controls on eDNA movement in streams: Transport, Retention,
768   and Resuspension. *Sci Rep* **7**, 5065 (2017).

23.   M. C. Yates, T. M. Wilcox, M. Y. Stoeckle, D. D. Heath, Interspecific allometric

scaling in eDNA production among northwestern Atlantic bony fishes reflects

physiological allometric scaling. *Environmental DNA* **5**, 1105–1115 (2023).

24.   M. J. Vanni, P. B. McIntyre, Predicting nutrient excretion of aquatic animals with

metabolic ecology and ecological stoichiometry: A global synthesis. *Ecology* **97**, 3460–

3471 (2016).

25.   J. E. Allegier, S. J. Wenger, A. D. Rosemond, D. E. Schindler, C. A. Layman,

Metabolic theory and taxonomic identity predict nutrient recycling in a diverse food

web. *PNAS* **112 (20)** (2015).

26.   M. C. Yates, *et al.*, Allometric scaling of eDNA production in stream-dwelling brook

trout (Salvelinus fontinalis) inferred from population size structure. *Environmental*

DNA **3**, 553–560 (2021).

27.   K. W. Thornton, A. S. Lessem, A Temperature Algorithm for Modifying Biological

Rates. *Trans Am Fish Soc* **107**, 284–287 (1978).

28.   I. Caza-Allard, M. Laporte, G. Côté, J. April, L. Bernatchez, Effect of biotic and abiotic

factors on the production and degradation of fish environmental DNA: An experimental

evaluation. *Environmental DNA* **4**, 453–468 (2022).

29.   D. Deslauriers, S. R. Chimpps, J. E. Breck, J. A. Rice, C. P. Madenjian, Fiash

Bioenergetic 4.0: An R-based Modeling Application. *Fisheries (Bethesda)* **42**, 586–

596 (2017).

30.   P. C. Hanson, T. B. Johnson, D. E. Schindler, J. F. Kitchell, Fish Bioenergetics 3.0.

World Wide Web Internet And Web Information Systems (1997).

31.   S. C. Chapra, K. H. Reckhow, *Ingeneering Approaches for Lake Management. Volume*
792   *2: Mechanistic Modeling* (Butterworth, 1983).

793   32.   L. M. Sassoubre, K. M. Yamahara, L. D. Gardner, B. A. Block, A. B. Boehm,
794   Quantification of Environmental DNA (eDNA) Shedding and Decay Rates for Three
795   Marine Fish. *Environ Sci Technol* **50**, 10456–10464 (2016).

796   33.   K. Kagzi, R. M. Hechler, G. F. Fussmann, M. E. Cristescu, Environmental RNA
797   degrades more rapidly than environmental DNA across a broad range of pH conditions.
798   *Mol Ecol Resour* **22**, 2640–2650 (2022).

799   34.   S. Huang, K. Yoshitake, S. Watabe, S. Asakawa, Environmental DNA study on aquatic
800   ecosystem monitoring and management: Recent advances and prospects. *J Environ*
801   *Manage* **323**, 116310 (2022).

802   35.   J. S. Messner, M. M. Maclennan, R. D. Vinebrooke, Higher temperatures enhance the
803   effects of invasive sportfish on mountain zooplankton communities. *Freshw Biol* **58**,
804   354–364 (2013).

805   36.   J. M. O. Matte, D. M. Glaser, J. R. Post, D. J. Fraser, Contrasting demographic
806   responses to size-selective harvesting among neighbouring wild fish populations.
807   *Journal of Applied Ecology* (2023). https://doi.org/10.1111/1365-2664.14420.

808   37.   D. Trépanier-Leroux, *et al.*, Density-dependent effects of exotic brook trout on aquatic
809   communities in mountain lakes revealed by environmental DNA and morphological
810   taxonomy. *Hydrobiologia* **851**, 1489–1512 (2023).

38.   M. C. Yates, *et al.*, The relationship between eDNA particle concentration and organism abundance in nature is strengthened by allometric scaling. *Mol Ecol* **30**, 3068–3082 (2021).

39.   T. M. Wilcox, *et al.*, Robust detection of rare species using environmental DNA: the importance of primer specificity. *PLoS One* **8**, e59520 (2013).

40.   C. J. Krebs, "Estimating Abundance in Animal and Plant Populations" in *Ecology: The Experimental Analysis of Distribution and Abundance*, 6th ed., (Pearson Benjamin Cummings, 2009).

41.   R. Tiberti, L. Nelli, S. Brighenti, R. Iacobuzio, M. Rolla, Spatial distribution of introduced brook trout Salvelinus fontinalis (Salmonidae) within alpine lakes: Evidences from a fish eradication campaign. *European Zoological Journal* **84**, 73–88 (2017).

42.   S. A. Bonar, W. A. Hubert, D. W. Willis, *Standard methods for sampling North American freshwater fishes* (American Fisheries Society, 2009).

43.   W. A. Hubert, K. L. Pope, J. M. Dettmers, "Passive capture techniques" in *A. V. Zale, D. L. Parrish, & T. . Sutton (Eds.), Fisheries Techniques*, 3rd ed., (American Fisheries Society, 2012).

44.   L. Johnson, Homeostatic Characteristics of Single Species Fish Stocks in Arctic Lakes. *Canadian Journal of Fisheries and Aquatic Sciences* **40**, 987–1024 (1983).

45.   J. R. Post, E. A. Parkinson, N. T. Johnston, Density-dependent processes in structured fish populations: Interaction strengths in whole-lake experiments. *Ecol Monogr* **69**, 155–175 (1999).

46. H. G. M. Ward, P. J. Askey, J. R. Post, D. A. Varkey, M. K. McAllister, Basin
characteristics and temperature improve abundance estimates from standard index
netting of rainbow trout (Oncorhynchus mykiss) in small lakes. *Fish Res* **131–133**, 52–
59 (2012).

47. G. E. Morgan, Manual of instructions - fall walleye index netting (FWIN). (2002).

48. Z. E. Schnabel, The Estimation of the Total Fish Population of a Lake. *The American*
Mathematical Monthly **45**, 348–352 (1938).

49. R. CoreTeam, R: A language and environment for statistical computing. [Preprint]
(2020). Available at: https://www.r-project.org.

50. D. H. Ogle, *Introductory Fisheries Analyses with R* (Chapman & Hall, 2015).

51. G. A. F. Seber, *The Estimation of Animal Abundance and Related Parameters*, B.
Publishing, Ed., 2 nd (2002).

52. K. J. Hartman, M. K. Cox, Refinement and Testing of a Brook Trout Bioenergetics
Model. *Trans Am Fish Soc* **137**, 357–363 (2008).

53. Environment and Natural Ressources Canada, Real-Time Hydrometric Data. (2024).
Available at: https://wateroffice.ec.gc.ca/map/index_e.html.

54. J. E. Littlefair, L. E. Hrenchuk, P. J. Blanchfield, M. D. Rennie, M. E. Cristescu,
Thermal stratification and fish thermal preference explain vertical eDNA distributions
in lakes. *Mol Ecol* **30**, 3083–3096 (2021).

55. J. Beaulieu, M. Gheta, A. A. M. Macdonald, Using Taylor Expansions to fit power law
functions. (2024). Available at: https://julien-beaulieu.github.io/taylor-expansions-
stats.github.io/vignette_taylor_series.html.

56. R. McElreath, *Statistical rethinking: A Bayesian course with examples in R and Stan*,
Chapman an (2018).

57. P. C. Bürkner, Advanced Bayesian multilevel modeling with the R package brms. *R
Journal* **10**, 395–411 (2018).

58. M. B. Parsley, E. J. Crespi, T. A. G. Rittenhouse, J. L. Brunner, C. S. Goldberg,
Environmental DNA concentrations vary greatly across productive and degradative
conditions, with implications for the precision of population estimates. *Sci Rep* **14**,
17392 (2024).

59. T. S. Jo, Correlation between the number of eDNA particles and species abundance is
strengthened by warm temperature: simulation and meta-analysis. *Hydrobiologia* **850**,
39–50 (2023).

60. K. J. Dunker, *et al.*, Potential of environmental DNA to evaluate northern pike (Esox
lucius) eradication efforts: An experimental test and case study. *PLoS One* **11**, 1–21
(2016).

61. J. Beaulieu, *et al.*, Environmental DNA for assessing impact and recovery of aquatic
communities in an invaded mountain lake. *Lake Reserv Manag* **40**, 111–131 (2024).

62. C. R. Turner, K. L. Uy, R. C. Everhart, Fish environmental DNA is more concentrated
in aquatic sediments than surface water. *Biol Conserv* **183**, 93–102 (2015).

63.   E. Capo, *et al.*, Lake Sedimentary DNA Research on Past Terrestrial and Aquatic
874   Biodiversity: Overview and Recommendations. *Quaternary* **4**, 6 (2021).

875   64.   J. J. Parkos, V. J. Santucci, D. H. Wahl, Effects of adult common carp (Cyprinus carpio)
876   on multiple trophic levels in shallow mesocosms. *Canadian Journal of Fisheries and*
877   *Aquatic Sciences* **60**, 182–192 (2003).

878   65.   J. K. M. Zimmerman, R. T. Palo, Reliability of catch per unit effort (CPUE) for
879   evaluation of reintroduction programs – A comparison of the mark-recapture method
880   with   standardized   trapping.   *Knowl   Manag   Aquat   Ecosyst*   07   (2011).
881   https://doi.org/10.1051/kmae/2011016.

882   66.   T. Saito, H. Doi, Degradation modeling of water environmental DNA : Experiments on
883   multiple   DNA   sources   in   pond   and   seawater.   850–860   (2021).
884   https://doi.org/10.1002/edn3.192.

885   67.   A. Hervé, *et al.*, Do temperature and population size structure influence the quantity of
886   fish eDNA in water? *Front Environ Sci* **11**, 1–6 (2023).

887   68.   B. Thalinger, *et al.*, The Effect of Activity, Energy Use, and Species Identity on
888   Environmental DNA Shedding of Freshwater Fish. *Front Ecol Evol* **9**, 1–13 (2021).

889   69.   R. P. Kelly, *et al.*, Toward a national eDNA strategy for the United States.
890   *Environmental DNA* **6**, 1–10 (2024).

891   70.   V. G. Fonseca, *et al.*, The Application of eDNA for Monitoring Aquatic Non-
892   Indigenous Species: Practical and Policy Considerations. *Diversity (Basel)* **15**, 1–17
893   (2023).

894   71.   O. Canals, I. Mendibil, M. Santos, X. Irigoien, N. Rodríguez-Ezpeleta, Vertical

895   stratification of environmental DNA in the open ocean captures ecological patterns and

896   behavior of deep-sea fishes. *Limnol Oceanogr Lett* **6**, 339–347 (2021).

899 **Table 1**. Parameter estimates and Bayes R^2 for the linear regression, allometrically scaled biomass regression and the mass balance
900 model. The formula column refers to the formula of the model specified as equations in the method section. The numbers in parentheses
901 represent the 95 % credible interval on parameter estimates. The Bayes R^2 reported are the mean of the Bayes R^2 of each model iteration.
902 b is the allometric coefficient, k is the sum of the degradation and sedimentation constant, I_0 is the eDNA generation coefficient, β is the
903 slope of the linear model, sigma is the variance. Sd stands for standard deviation.

Model	Bayes R^2	b	k	Sd(ln(k))	Ln(I_0) or β	Sd(ln(I_0))	Sigma	Formula
Linear regression	0.24	-	-	-	0.13 (0.05-0.21)	-	2387 (1889-3037)	equation 6
Allometrically scaled biomass	0.44	0.58 (0.25-0.89)	-	-	0.03 (-1.4-1.43)	-	2162.93 (1707-2775)	equation 7
Mass balance eDNA	0.66	0.61 (0.31-0.92)	0.5 (0.41-0.6)	0.17 (0.01-0.59)	9.71 (8.18-11.11)	-	θ = 5.01 (4.35-5.77) α = 0.14 (0.09-0.18)	equation 5
Mass balance with N as dependent variable, with lake specific k and sampling period specific I_0	0.72	0.59 (0.18-0.93)	0.49 (0.40-0.6)	0.40 (0.14-0.82)	10.06 (8.33-11.98)	0.33 (0.03-0.83)	θ = 2.49 (1.26-3.75) α = 0.16 (0.10-0.21)	equation 8
Mass balance with N as dependent variable without sampling period specific I_0	0.70	0.65 (0.21-0.96)	0.49 (0.4-0.59)	0.5 (0.21-0.97)	9.7 (8.13-11.76)	-	θ = 2.84 (1.69-4.02) α = 0.15 (0.09-0.2)	equation 8.1
Mass balance with N as dependent variable without lake specific k nor sampling period specific I_0	0.51	0.47 (0.06-0.96)	0.49 (0.4-0.59)	-	10.60 (8.23-12.47)	-	θ = 2.75 (1.55-3.98) α = 0.17 (0.11-0.21)	equation 8.2

Table 2. Parameter estimates for bias testing. The bayes R^2 reported is the mean of the Bayes R^2
of each draw. The dependent variable is the z-score of each total Brook Trout population abundance
estimate, calculated as the estimate from eDNA minus the estimate from mark-recapture divided
by the standard deviation of the draws of the estimate from eDNA. After running the model with
all the lakes, the same model was fitted without Helen Lake to see if some of the results were driven
only by this one lake. CI stands for credible interval.

Variable	With Helen Lake			Without Helen Lake		
	Parameter estimate	CI (95 %)	Bayes R^2	Parameter estimate	CI (95 %)	Bayes R^2
Total abundance	-0.00045	-0.0087 – 0.0018	0.40	-0.00054	-0.0008 – 0.0018	0.31
Lake area	0.02	-0.18 – 0.21		-0.01	-0.21 – 0.19	
Mixing (yes)	0.20	-1.72 – 2.10		-0.14	-2.12 – 1.88	
Elevation	-0.00024	-0.0021 – 0.0026		-0.00008	-0.0031 – 0.0029	
Season (fall)	0.15	-1.67 – 1.98		-0.28	-2.17 – 1.62	
Treatment (Harvest)	-0.06	-1.38 – 1.30		-0.35	-1.76 – 1.12	
Elevation*Mixing	-0.001	-0.004 – 0.002		-0.00009	-0.0026 – 0.0024	
Elevation*Season	-0.0005	-0.0025 – 0.0021		0.0001	-0.002 – 0.003	
Season*Treatment	1.17	-0.39 – 2.70		0.80	-0.76 – 2.35	

List of figures

a) Linear regression   **b) Allometrically-scaled mass**   **c) Mass balance**