Sporophyte Directed Gametogenesis via the Ubiquitin Proteasome System

Prakash Sivakumar[1,2,†], Saurabh Pandey[1,†,‡], A Ramesha[1,†,§], Jayeshkumar Narsibhai Davda[1], Aparna Singh[1,¶], Chandan Kumar[1,#], Hardik Gala[1], Veeraputhiran Subbiah[1,††], Harikrishna Adicherla[1], Jyotsna Dhawan[1], L. Aravind[3], Imran Siddiqi[1,2,*]

Affiliations:

[1]CSIR–Centre for Cellular and Molecular Biology; Uppal Road, Hyderabad 500007, India.

[2]Academy of Scientific and Innovative Research (AcSIR), Ghaziabad - 201002, India

[3]National Center for Biotechnology Information, National Library of Medicine, National Institutes of Health; Bethesda, MD 20894, USA.

*Corresponding author. Email: imran@ccmb.res.in

[†] These authors contributed equally to the work

Present address:

[‡] databaum GmbH; Holsteinischer Kamp 80, 22081 Hamburg, Germany.

[§] Seri-Biotech Research Laboratory, Central Silk Board; Bangalore-560035, India.

[¶] Department of Botany, Banaras Hindu University; Varanasi 221005, India.

[#] University of Texas at Austin; Texas, USA.

[††] Deceased

Abstract

Plants alternate between diploid sporophyte and haploid gametophyte generations. In mosses which retain features of ancestral land plants, the gametophyte is dominant and has an independent existence. However, in flowering plants the gametophyte has undergone evolutionary reduction to just a few cells enclosed within the sporophyte. The gametophyte is thought to retain genetic control of its development even after reduction. Here we demonstrate that male gametophyte development in Arabidopsis, long considered to be autonomous, is also under genetic control of the sporophyte via a repressive mechanism involving large-scale regulation of protein turnover. We identify an Arabidopsis gene *SHUKR* as an inhibitor of male gametogenesis. SHUKR is unrelated to proteins of known function and acts sporophytically in meiosis to control gametophyte development by negatively regulating expression of a large set of ubiquitination genes specific to post-meiotic gametogenesis. This control is late-emerging as *SHUKR* homologs are found only in eudicots. We show that *SHUKR* is rapidly evolving under positive selection suggesting that variation in control of protein turnover during male gametogenesis has played an important role in evolution within eudicots.

Introduction

Alternation between two distinct generations: a diploid sporophyte and a haploid gametophyte is a central feature of the land plant life cycle[1–3]. Land plants likely evolved from an algal ancestor related to extant charophycean algae which comprise the closest relatives to land plants[4,5]. Charophycean algae have a predominantly haploid life cycle, the diploid phase being transient as the zygote directly undergoes meiosis to give haploid spores. The emergence of a multicellular diploid sporophyte contributed to reproductive success on land by increasing the number of spores produced from a single fertilization event, and also increased potential for genetic variability and evolutionary change[1]. From a simple ancestral body plan, land plants comprising bryophytes (mosses, liverworts, and hornworts) and vascular plants evolved progressively specialized cell types, organs, and architectures adapted for life on land. These acquisitions have been based on development and increase in complexity of a multicellular sporophyte[6,7]. In vascular plants the sporophyte evolved to become the dominant phase and the gametophyte has reduced. In flowering plants the gametophyte is reduced to just a few cells and develops enclosed within the sporophyte[8] (Fig. 1a). The differing trajectories taken by the sporophyte and gametophyte in land plant evolution reflects an uncoupling between their body plan evolution consistent with different genetic programs directing their body organization[7,9,10,11,12].

Formation of a germline is an early step in animal development[13,14], however in plants the formation of germ cell precursors takes place late in development[15]. The fact that bryophytes have a multicellular independent gametophyte which produces gametes following extensive mitotic divisions allows overall assignment of the germline to the

gametophyte generation[16]. In flowering plants the assignment of the germline is more

difficult because of extreme reduction of the gametophyte and its development

enclosed within the sporophyte[15,17,18]. Evidence from comparative evolution showing

functional conservation of transcription factors controlling early steps in germ cell

development between the liverwort *Marchantia polymorpha* and Arabidopsis argues

for the male germline residing in the gametophyte[16]. Asymmetric cell division of the

haploid microspore in Arabidopsis at pollen mitosis I forms a generative cell (GC) and

vegetative cell (VC)[16,18]. In Arabidopsis, specification of the GC requires bHLH

transcription factors, BNB1/BNB2 and DROP1/DROP2 which act in the gametophyte

and are orthologous to MpBNB and MpDROP respectively, that also act in the

gametophyte to direct gametangia specification and differentiation in Marchantia[19,20].

Sperm differentiation is controlled by gametophytic action of the bHLH transcription

factor DUO1 in both Arabidopsis and Marchantia[21,22]. On the female side

gametophytically acting transcription, splicing, and signaling related factors are

important for cell specification and cellularization in Arabidopsis female

gametophytes[23–29]. Therefore, the gametophyte in flowering plants has been

considered to largely retain genetic control of its development even though it lacks an

independent existence.

The extreme reduction of the gametophyte and its development enclosed within the

sporophyte provided a "functionally unitary individual" for natural selection[30],

addressing the sensitive requirements for fertilization and reproductive development

in a land environment. Formation and function of meiotic cells depends on a range of

cellular interactions in the sporophyte[8,31–33]. These interactions involve transcriptional

regulators acting nonautonomously[34–36], cell surface receptor signaling[37–39], and small

RNA pathways[36,40,41] together with phytohormone effectors[42]. Small RNAs[43] and the

phytohormone auxin[44,45] have also been shown to mediate developmental interactions

between cells within gametophytes. The tapetum is a sporophytic nurse tissue

surrounding male meiocytes and provides enzymes for release of microspores and

components for synthesis of the protective secondary outer coat of the pollen grain,

the exine[46,47]. Mutants deficient in tapetal function manifest defects in exine formation

and lead to pollen abortion[48–50]. The tapetum also produces small RNAs that direct

DNA methylation of transposons and selected genes in male meiosis contributing to

genome integrity and paternal epigenetic inheritance[51].

Studies have shown that isolated uninucleate microspores can form functional pollen

in vitro suggesting their development into pollen is autonomous[52,53]. It remains unclear

whether genetic controls during meiosis broadly regulate postmeiotic gametophyte

development. A challenge in uncovering such regulation has been the existence of

very many sporophytic mutants that affect core meiosis functions including meiotic

recombination and chromosome organization[54,55] or else development of somatic cells

within the anther or ovule[56–59]. These mutants have direct or indirect effects on

gametophyte development and show spore or gametophyte abortion, making it difficult

to distinguish from mutants affecting meiotic regulation of postmeiotic gametogenesis.

Here we report the discovery in Arabidopsis that following evolutionary reduction in

cell number of the gametophyte, the sporophyte within the eudicot lineage has gone

on to evolve a process established in meiosis under control of the *SHUKR* gene, by

which it directs male gametophyte development. The *shukr* mutant shows normal

meiosis but postmeiotic development is defective. This process controlled by SHUKR

involves large-scale control of regulated protein turnover in the developing

gametophyte and is under rapid evolution with positive selection. Our findings suggest

that the sporophyte in addition to providing supportive conditions for gametogenesis

and fertilization, also exerts genetic control in meiosis of postmeiotic male

gametogenesis. This control is part of a process that is capable of generating variation

in the gametophyte during evolution, which has adaptive value.

Results

***SHUKR* is required for male fertility**

We identified the *SHUKR* gene in a screen for novel meiotic functions based on

coexpression with known meiotic genes in Arabidopsis (see Methods). Three

independent insertion lines having insertions in *At2g38690* showed male sterility (Fig.

1b, Extended Data Fig. 1a). Allelism tests indicated the mutants to be allelic. Anthers

from plants homozygous for the insertion contained a highly reduced number of viable

pollen, resulting in sterility (Fig. 1b, c). Use of wild type pollen as the male donor in

crosses to mutant plants resulted in full seed set indicating that female fertility was

unaffected (Fig. 1c). The male sterile phenotype was inherited as a single gene

recessive trait (Fig. 1c, Extended Data Fig. 1c). We named the *At2g38690* gene

SHUKR (*SKR*) after the Sanskrit word Shukranu (/śukrāṇu/) for sperm. We confirmed

the phenotype was due to loss of function of *SKR* by complementation with a

transgenic copy. *skr/+* plants were transformed with wild type *At2g38690* genomic

DNA (*gSKR*) by in planta transformation and T1 *skr* progeny were examined for

fertility. The *gSKR* transgene rescued male sterility confirming the mutant phenotype

was due to disruption of *At2g38690* (Fig. 1b, Extended Data Fig. 1d). Examination of

SKR gene expression based on tissue RT-PCR, a proSKR:GFP-GUS reporter, and

analysis of publicly available expression profiling data showed that SKR gene

expression was specific to reproductive tissues and more specifically to male

meiocytes (Fig. 1d-g; Extended Data Fig. 2). Expression of SKR protein was also

meiosis-specific (Fig. 2c).

Early male gametogenesis defects in the *shukr* mutant

Analysis of meiosis and pollen development was performed to identify the stage at

which defects arose in *skr*. Pollen development is divided into two stages:

microsporogenesis comprising meiosis together with microspore development, and

microgametogenesis or male gametophyte development comprising development of

the haploid microspore into pollen[60]. No defects were observed in *skr* during meiosis

based on analysis of chromosome spreads (Fig. 2a). Light microscopic examination

of whole mount cleared anthers and plastic sections, showed that tetrad formation and

release of microspores took place in *skr* (Figs. 1h,2b; Extended Data Fig. 3), however,

microspores degenerated shortly after release from tetrads (Fig. 2b; Extended Data

Fig. 4). Newly released wild type microspores were angular in shape (Fig. 2b)[61]

whereas *skr* microspores were more rounded and variable in cell wall thickness (Fig.

2b), suggesting loss of rigidity and possible defects in cell wall development. Hence

sterility in *skr* appeared to be associated with defects in male gametogenesis between

tetrad formation and early gametophyte development.

Ultrastructural examination of tetrad and early microgametogenesis stages (Fig. 3a)

by transmission electron microscopy revealed that a significant proportion of *skr*

tetrads harbored microspores in which the plasma membrane (PM) was retracted from

the surrounding callose wall (Fig. 3b-d). PM retraction was associated with presence

of material between the PM and callose layer likely reflecting unincorporated cell wall

precursors or disrupted material arising from disturbance in cell wall development.

Hence onset of the mutant phenotype was detected at the tetrad stage. To test for

additional defects in *skr*, we examined expression of the LAT52:GFP reporter that

specifically marks the single vegetative cell after pollen mitosis 1[62]. Among *skr*

microspores in which expression of LAT52:GFP took place, we observed two signals

in 30% of cases (Fig. 3e). Therefore loss of *SKR* led to at least two distinct defects in

early gametogenesis: retraction of the PM with associated disruption of cell wall

development, and aberrant expression of an early differentiation marker.

Recessive gene action in the diploid sporophyte would give Mendelian 3:1 segregation

of the mutant allele. Gene requirement in the haploid male gametophyte would result

in reduced transmission of the mutant allele through the male and a deviation from this

ratio. The genetic analysis suggested that *SKR* acts sporophytically as we did not

observe a significant deviation from this ratio (Fig. 4a), whereas the mutant phenotype

is with respect to the early gametophytic phase. We tested whether *SKR* also has a

gametophytic component to its action by reciprocal crosses of *skr/+* to wild type and

examining transmission frequency of the mutant *skr* allele through the gametophyte in

comparison to the wild type allele. This test measures fitness of mutant relative to wild

type gametes from the same flowers. If there is a functional requirement for *SKR* gene

expression at the haploid stage in the gametophyte, then mutant pollen would show

reduced fitness and the mutant allele would be transmitted to the next generation at

lower frequency than the wild type allele. The results indicated no significant difference

in transmission frequency of the mutant *skr* allele from that of the wild type (Fig. 4a),

thus failing to provide evidence for *SKR* gene action within the gametophyte. Therefore

the genetic results together with ultrastructure and marker expression indicated that

SKR gene action is in the diploid sporophyte and is required for proper execution of

early stages of gametogenesis. Further tests confirmed this finding (see below).

Meiotic control of male gametophyte gene expression by *SHUKR*

The sporophyte is known to play an important role in formation of the pollen wall and

aperture[60,63]. Much of the material and machinery for building the outer wall (exine), is

synthesized and released by the tapetum, a layer of nurse cells adjacent to the

meiocytes[47,64]. Examination of *SKR* gene expression described above indicated that

SKR is expressed specifically in male meiocytes. Therefore unlike the sporophytic

genes which function in the tapetum, *SKR* appears to act in meiosis. We determined

the localization of SKR protein in the course of meiosis using a complemented line

carrying a *proSKR:GFP-gSKR* gene fusion. SKR protein was detected specifically in

male meiocytes from prophase to tetrad stage and extending to early microspores.

Subcellular localization of GFP-SKR varied during meiosis, from nuclear enrichment

at prophase, changing toward cytoplasmic in late prophase, and back to nuclear at the

tetrad stage. Signal intensity was highest in tetrads (Fig. 2c). Localization in early

microspores shifted from nuclear towards more of cytoplasmic, before declining to an

undetected level in later stage uninucleate microspores (UNM). Expression was not

detected in tapetal cells (Fig. 2c). Therefore SKR protein is mainly present during

meiosis and the onset of the *skr* mutant phenotype corresponds to the tetrad stage at

which SKR signal is highest. High SKR expression in tetrads followed by steep decline

in free microspores suggests a possible role in repressing gametogenesis or else in

ending the sporophytic program before initiation of gametogenesis.

To investigate the above possibilities we examined gene expression during meiosis

by RNAseq profiling of meiotic anthers (extending upto and not beyond tetrad stage)

for *skr* and wild type. 619 genes were differentially expressed in *skr* of which 379 genes

were upregulated (Supplementary Table 1). We analyzed upregulated genes based

on cell and tissue expression in publicly available datasets using the CoNekT

platform[65] and found a dramatic enrichment for male gametogenesis-enriched

genes[66,67] (Fig. 4b, Supplementary Table 2). 187/379 (49%) of upregulated genes

were male gametogenesis-enriched. Genes showing maximal enrichment in UNMs

represented 80%, followed by bicellular and tricellular pollen stages (16%).

Upregulated genes included *DML3* which has been suggested to be an early

gametogenesis marker[68]. Increased expression of a large set of gametogenesis-

enriched genes in meiotic anthers of *skr* suggests that SKR negatively regulates

expression of gametogenesis genes and represses gametogenesis.

228   SHUKR acts sporophytically and inhibits gametogenesis

To test for a role in repression of gametogenesis as well as the possibility that SKR

protein acts postmeiotically to promote pollen development, we adopted a

misexpression strategy by directing expression of SKR in microspores using the well

characterized microspore specific MSP1 promoter[69]. *proMSP1:SKR* in a *skr* mutant

failed to rescue the mutant phenotype, indicating that SKR expression in the

gametophyte is not sufficient to restore function. Furthermore, *proMSP1:SKR* in wild

type in fact caused a reduction in pollen viability (Fig. 4c). These results show that

expression of SKR during gametogenesis interferes with gametophyte development

and argue against the possibility that SKR protein present during postmeiotic stages

contributes to pollen development. Therefore SKR acts sporophytically and is an

inhibitor of male gametogenesis.

SHUKR represses the ubiquitin proteasome system

Gene ontology (GO)[70] together with further analysis of individual gene families

revealed a striking signature of regulated protein turnover among the 379 genes

upregulated in _skr_ RNAseq. 23% of upregulated genes were associated with ubiquitin-

mediated degradation of proteins[71] (Fig. 5a, Supplementary Table 3; p < 2.0e-33). At

the level of molecular function, there was enrichment for ubiquitin protein transferase

activity covering the Skp1-Cullin-Fbox (SCF) complex of E3 ubiquitin ligase as well as

other components of the ubiquitin protease system (UPS) including _RING/U-box_ and

ARIADNE genes which encode E3 ubiquitin ligases[72,73]. F-box genes encode the

substrate specificity component of SCF and comprised the largest family among

upregulated genes in _skr_. F-box genes in Arabidopsis make up a family of

approximately 740 genes[74,75] (Supplementary Table 3), several of which show

expression during male gametogenesis[76]. 69 F-box genes representing 9% of all F-

box genes in Arabidopsis were upregulated in _skr_ (R.F. = 6.6, p < 1.74e-36; Fig. 5a;

Supplementary Table 3). Hence a significant fraction of F-box genes are implicated in

pollen development. 75% of upregulated F-box genes were maximally enriched at the

UNM stage and 9% at later stages comprising bicellular and tricellular pollen. Among

downregulated genes, there was enrichment for stress and hypoxia related genes

(Supplementary Table 3). Upregulation of a large number of UPS genes suggested

involvement of SKR in general control of protein homeostasis and further experiments

were performed to test this hypothesis (see below). We also examined stage-wise

expression of a subset of upregulated UPS genes at the cellular level, by generating

lines carrying a GFP gene fusion for three different F-box genes (*At3g28330,*

At5g02980 and *At5g62510*) and segregating for *skr*. In each case, we observed F-

box:GFP expression in wild type only during postmeiotic development of microspores

into pollen, whereas *skr* showed precocious F-box:GFP expression in meiosis (Fig.

5b, Extended Data Fig. 5). Hence SKR can be seen to act in meiosis to bring about

repression of these F-box genes. Taken together, these observations led us to

hypothesize that SKR represses onset of male gametogenesis during meiosis through

timing of expression of a large set of male gametophyte-preferred F-box and other

UPS genes.

Control of protein homeostasis by *SHUKR*

We directly investigated the functional basis for the *skr* phenotype by performing a

suppressor screen on *skr* using EMS mutagenesis of seeds collected from a large

number of *skr* plants followed by screening of M2 plants for improved seed set. We

isolated a suppressor of *skr* which showed elongated siliques and increased pollen

viability, whereas *skr* shows low pollen viability. The suppressor line was backcrossed

to the parent *skr* strain and F1 plants were sterile. F2 seed were collected from multiple

F1 plants and grown to give F2 plants which showed recessive Mendelian inheritance

for fertility (Extended Data Fig. 6a). Whole genome sequencing of bulk segregant

fertile and sterile plants from the backcrossed F2 population led to identification of an

SNP at the *SKI2* locus which created a stop codon after 517 amino acids and showed

100% linkage to the suppressor phenotype (Fig. 5c, Extended Data Fig. 6b). *SKI2*

encodes the RNA helicase subunit of the SKI complex and prevents secondary siRNA

production in Arabidopsis[77]. *ski2* has also been isolated as a suppressor of

overexpression phenotypes in a different context[78]. We tested an independent allele

of *ski2* by crossing to *skr* and observed suppression of the *skr* mutant phenotype,

confirming *ski2* as a suppressor of *skr* (Extended Data Fig. 6c). We then performed

RNAseq on meiotic anthers of the *skr ski2* backcrossed suppressor line. Comparison

of differentially expressed genes with the *skr* parent revealed a prominent signature of

the proteasome and translation among downregulated genes: 56% of genes encoding

subunits of the proteasome and 29% of translation related genes were down-regulated

in *skr ski2* (Fig. 5d, Supplementary Table 4). Notably, F-box genes as a class were

not downregulated in *skr ski2* (R.F.= 0.3; $p < 7.7e-19$). The striking reduction in

transcript levels of the majority of proteasomal genes in *skr ski2* strongly supports the

hypothesis that the *skr* phenotype arises from uncoordinated degradation by the

proteasome, of regulator(s) of gametogenesis due to inappropriate expression of

gametogenesis specific F-box and possibly other ubiquitination genes. Reduction of

proteasome activity by downregulation of a major subset of proteasome genes

rescues the mutant phenotype.

We confirmed involvement of the proteasome in the *skr* phenotype using two different

approaches. First, we performed knockdown of PBE1, a core component of the

proteasome[79] in the *skr* mutant background using *pro*SKR to direct meiosis-specific

expression of PBE1 hairpin RNA and scored for pollen viability in RNAi plants.

Knockdown of PBE1 by RNAi has been used in a different context[80] and we used the

same region for hairpin RNA construction in our experiments. Second, we treated

inflorescences of *skr* mutant plants with two physiological inhibitors of the proteasome

(MG132 and MG115). In both regimes, we observed an increase in formation of viable

pollen that was significant for PBE1:RNAi and MG115 (Fig. 5e,f; $p < 0.05$). Taken

together, these results clearly establish deregulation of proteasome activity as a major

basis for the *skr* mutant phenotype. The large number of F-box genes showing

deregulation during meiosis in *skr*, most of which normally are preferentially expressed

only after meiosis during gametogenesis, highlights the scope of this control. Hence,

SKR plays a crucial role in meiosis to repress a proteasomal program that promotes

gametogenesis and which comes into play in free microspores following meiosis. SKR

therefore sporophytically regulates postmeiotic development of the gametophyte

through timing of regulated protein turnover by the proteasome (Fig. 5g). Notably, this

programme is distinct from the canonical meiotic processes of chromosome reduction

and spore formation, which remain unaffected in *skr*.

323   *SHUKR* is eudicot specific and rapidly evolving

A cDNA of the *SKR* coding region was amplified by RT-PCR, sequenced, and found

to encode a predicted protein of 191 amino acids in agreement with the annotated

CDS. Homologs were found only in plants and bore no similarity to any known protein

family. SHUKR is therefore the founding member of a new protein family (hereinafter

SHUKR). The only alignable region conserved across all SHUKR homologs is always

at the C-terminus and this is usually the only globular domain in the protein from

Shannon entropy-based compositional analysis. The domain has a central

compositionally biased, poorly structured variable region enriched in negatively

charged residues. Thus, the SHUKR domain varies in length from 67 to 119 residues.

It has a hallmark CxC motif and a dyad of aromatic amino acids (usually Y or F) close

to the end of the domain (Fig. 6a, Supplementary Figure 1). The SHUKR family is

present in both great divisions of eudicots: the Superrosids and Superasterids. It is

additionally present in Proteales (Nelumbo) and Ranunculales (Macleaya, Papaver),

which are the two basal-most branches of eudicots. It is absent in monocots and basal

angiosperms. Hence the origin of SHUKR is at the base of eudicots.

To understand evolution of the SHUKR family we used the SHUKR domain of 119

representatives from 85 species for tree construction. Branch length analysis of the

tree revealed two lineages displaying elevated evolutionary rates:

Brassicaceae+Cleomaceae and Asteraceae (Fig. 6b; Extended Data Fig. 7). This was

further tested using the maximum-likelihood distance matrix computed from the input

alignment for the tree. The mean pairwise distance was calculated between each

operational taxonomic unit (OTU) and rest of the OTUs, and centered using overall

mean distance for the matrix. These mean-centered distances showed a bimodal

distribution with the main mode between -1.25 to 0.5, a second mode from 0.5 to 1.5,

and a few outliers beyond that (Fig. 6c). The second mode corresponds to sequences

with higher values of mean-centered distance, indicating their more extensive

divergence. Brassicaceae+Cleomaceae ($p = 1.1472e{-}04$) and Asteraceae ($p = 1.6e{-}$

16) were significantly enriched among sequences with mean-centered distance

greater than 0.5 indicating they represent atypically fast-evolving groups. We note that

the high divergence of the SHUKR family outside the SHUKR domain posed

challenges in alignment across the entire coding sequence. Therefore the analysis

likely represents a conservative assessment of evolutionary rates within the *SHUKR*

gene family.

***SHUKR* is under positive selection**

We assessed rapid evolution in particular plant clades by computing column-wise

Shannon entropy from an alignment of the SHUKR domain for four plant families with

good representation in the Genbank nr database: Brassicaceae, Asteraceae,

Fabaceae and Rosaceae. The column-wise entropy was computed using both the 20

amino acid alphabet and a reduced 7 amino acid alphabet that groups together

residues as per their shared sidechain chemistry (see Methods). Elevation of entropy

in the reduced alphabet indicates changes in sidechain properties along a column and

is a proxy for diversifying selection[81]. Entropy plots in both alphabets indicated

significantly elevated entropy in Brassicaceae and Asteraceae relative to other

families, suggesting action of diversifying selection in these lineages (Fig. 6d). At least

30% of species examined had more than one paralog per genome most of which

appeared to represent recent lineage-specific duplications as they tended to group

together in the tree. However, Asteraceae appears to have experienced a duplication

early in their history. Likewise the duplication in Brassicaceae and Cleomaceae

predated their divergence. We further assessed diversifying selection at *SHUKR* within

Brassicaceae based on an alignment spanning the length of the gene, using branch-

site models[82] that consider proteins as a mosaic of conserved and divergent domains,.

The branch-site test results provided further support for existence of positive selection

(p = 0.0031) (Extended Data Table 1; Extended Data Fig. 8; Supplementary Figure 2),

demonstrating that gene duplication has been followed by diversifying selection at

SHUKR.

Adaptive value for variation in protein turnover during male gametogenesis

383   Plant F-box genes have undergone independent lineage-specific expansions as well

384   as reductions during land plant evolution[83,84]. The Brassicaceae family has an

385   expanded set of F-box genes, a majority of which are of short taxonomic scale

386   indicative of more recent emergence[84]. These recent members were significantly

enriched among upregulated F-box genes in *skr* (p < 1.5e-05) (Fig. 7a,c; Supplementary Table 5). Orthology analysis of the total set of F-box genes from 10 eudicot species[84] indicated that 58% (37/64) of F-box genes upregulated in *skr* belonged to orthogroups that included *Arabidopsis lyrata* but not 8 other eudicot species examined spanning 7 families including *Carica papaya* (family *Caricaceae*, order *Brassicales*), suggesting that these were likely specific to the Brassicaceae lineage (Fig. 7b; Supplementary Table 5). 8% (5/64) of genes belonged to orthogroups having only *A. thaliana* and not *A. lyrata* suggesting a more recent origin, and 23% (15/64) had at least one other species but not *A. lyrata* indicative of recent loss (Fig. 7b). We examined the possibility of diversifying selection at the set of *skr* upregulated genes by testing whether the ratio K_a/K_s of non-synonymous to synonymous substitutions per site was higher relative to the overall set of F-box genes. Plots of K_a/K_s gave significantly higher values for the set of F-box genes upregulated in *skr* relative to the overall set of F-box genes (p = 0.0075; Fig. 7d). Hence the set of *skr* upregulated F-box is itself undergoing diversifying selection. Interestingly, male gametes have been proposed as a source for emergence of new genes in both animals and plants[85–87], and pollen-preferred F-box genes also appear to belong to this set. Variation in protein turnover during male gametophyte development can therefore be seen to operate at two levels in evolution. One is through rapid evolution and positive selection in a regulator of UPS genes involved in male gametophyte development. The second is in changes and diversifying selection in components of the UPS system itself, the set of F-box genes that direct substrates for proteasomal degradation in gametogenesis. The large number of F-box genes involved and their distribution across the F-box phylogenetic tree (Fig. 7c) suggests the likelihood of levels of multiple proteins being subject to regulation by SKR. Variation in this control

provides a plausible route for the sporophyte to influence development and evolution of the gametophyte. These findings lead us to hypothesize that processes that provide for variation in levels of large sets of proteins rather than just a single target during male gametophyte development have adaptive value and have contributed significantly to evolution in Brassicaceae.

Discussion

Extensive functional and gene expression studies of gametophyte development[15,18,88] have supported the view that the male gametophyte retains control of its development even after evolutionary reduction to just a few cells enclosed within the sporophyte. Experiments in angiosperm species have also shown that isolated microspores can form pollen in culture[52,53,89], supporting this view. Here we have demonstrated in Arabidopsis that the sporophyte in fact plays an important role in control of gametogenesis. This direction is by a repressive mechanism involving large scale control of regulated protein turnover in male gametophyte development by a novel gene, *SHUKR*. *SHUKR* acts during meiosis and is an inhibitor of gametogenesis (Fig. 5g). Our findings in Arabidopsis suggest that significant aspects of the male gametogenesis programme are established in the diploid sporophyte during meiosis but held in a repressed condition until microspores are released. We suggest the term sporophyte directed gametogenesis (SDG) for this phenomenon.

Recent emergence of *SHUKR* at the base of eudicots suggests that SDG arose late in evolution following reduction of the gametophyte. Gametophyte reduction and its close proximity to the sporophyte may have provided conditions necessary for the emergence of SDG. Rapid evolution and diversifying selection at *SHUKR* and the set

of F-box genes under its control points to an adaptive value for variation in protein

turnover in male gametophyte evolution. This variation provides a way for the

sporophyte to generate diversity in the male gametophytes which face intense

selective pressure in the course of transmission and intraspecific competition[90]. The

location of this control in the sporophyte is also of interest because it raises the

possibility that gametophyte development can be modulated by the sporophyte as a

function of its own status and environment.

References

1. Graham, L. E. The Origin of the Life Cycle of Land Plants: A simple modification in the life cycle of an extinct green alga is the likely origin of the first land plants. *American Scientist* **73**, 178–186 (1985).

2. Kenrick, P. Alternation of Generations in Land Plants: New Phylogenetic and Palaeobotanical Evidence. *Biological Reviews* **69**, 293–330 (1994).

3. Niklas, K. J. & Kutschera, U. The evolution of the land plant life cycle. *New Phytologist* **185**, 27–41 (2010).

4. Graham, L. E. Origin of Land Plants. *Wiley.com* https://www.wiley.com/en-us/Origin+of+Land+Plants-p-9780471615279 (1993).

5. Delwiche, C. F. & Cooper, E. D. The Evolutionary Origin of a Terrestrial Flora. *Curr Biol* **25**, R899-910 (2015).

6. Kenrick, P. & Crane, P. R. The origin and early evolution of plants on land. *Nature* **389**, 33–39 (1997).

7. Graham, L. E., Cook, M. E. & Busse, J. S. The origin of plants: Body plan changes contributing to a major evolutionary radiation. *Proceedings of the National Academy of Sciences* **97**, 4535–4540 (2000).

8. Ma, H. & Sundaresan, V. Chapter Thirteen - Development of Flowering Plant Gametophytes. in *Current Topics in Developmental Biology* (ed. Timmermans, M. C. P.) vol. 91 379–412 (Academic Press, 2010).

9. Harrison, C. J. & Morris, J. L. The origin and early evolution of vascular plant shoots and leaves. *Philos Trans R Soc Lond B Biol Sci* **373**, 20160496 (2018).

10. Sakakibara, K. *et al.* KNOX2 genes regulate the haploid-to-diploid morphological transition in land plants. *Science* **339**, 1067–1070 (2013).

11. Mosquna, A. *et al.* Regulation of stem cell maintenance by the Polycomb protein

FIE has been conserved during land plant evolution. *Development* **136**, 2433–

2444 (2009).

12. Okano, Y. *et al.* A polycomb repressive complex 2 gene regulates apogamy and

gives evolutionary insights into early land plant evolution. *Proc Natl Acad Sci U S*

A **106**, 16321–16326 (2009).

13. Strome, S. & Lehmann, R. Germ versus soma decisions: lessons from flies and

worms. *Science* **316**, 392–393 (2007).

14. Hayashi, K. & Surani, M. A. Self-renewing epiblast stem cells exhibit continual

delineation of germ cells with epigenetic reprogramming in vitro. *Development*

136, 3549–3556 (2009).

15. Berger, F. & Twell, D. Germline Specification and Function in Plants. *Annual*

Review of Plant Biology **62**, 461–484 (2011).

16. Hisanaga, T. *et al.* Building new insights in plant gametogenesis from an

evolutionary perspective. *Nat Plants* **5**, 663–669 (2019).

17. Grossniklaus, U. Plant germline development: a tale of cross-talk, signaling, and

cellular interactions. *Sex Plant Reprod* **24**, 91–95 (2011).

18. Twell, D. Male gametogenesis and germline specification in flowering plants. *Sex*

Plant Reprod **24**, 149–160 (2011).

19. Yamaoka, S. *et al.* Generative Cell Specification Requires Transcription Factors

Evolutionarily Conserved in Land Plants. *Curr Biol* **28**, 479-486.e5 (2018).

20. Saito, M. *et al.* A bHLH heterodimer regulates germ cell differentiation in land

plant gametophytes. *Curr Biol* **33**, 4980-4987.e6 (2023).

21. Borg, M. *et al.* The R2R3 MYB Transcription Factor DUO1 Activates a Male

Germline-Specific Regulon Essential for Sperm Cell Differentiation in

Arabidopsis. *The Plant Cell* **23**, 534–549 (2011).

22. Higo, A. *et al.* Transcription factor DUO1 generated by neo-functionalization is

associated with evolution of sperm differentiation in plants. *Nat Commun* **9**, 5283

(2018).

23. Kasahara, R. D., Portereiko, M. F., Sandaklie-Nikolova, L., Rabiger, D. S. &

Drews, G. N. MYB98 is required for pollen tube guidance and synergid cell

differentiation in Arabidopsis. *Plant Cell* **17**, 2981–2992 (2005).

24. Gross-Hardt, R. *et al.* LACHESIS restricts gametic cell fate in the female

gametophyte of Arabidopsis. *PLoS Biol* **5**, e47 (2007).

25. Moll, C. *et al.* CLO/GFA1 and ATO are novel regulators of gametic cell fate in

plants. *Plant J* **56**, 913–921 (2008).

26. Rabiger, D. S. & Drews, G. N. MYB64 and MYB119 Are Required for

Cellularization and Differentiation during Female Gametogenesis in Arabidopsis

thaliana. *PLOS Genetics* **9**, e1003783 (2013).

27. Yuan, L. *et al.* The CKI1 Histidine Kinase Specifies the Female Gametic

Precursor of the Endosperm. *Developmental Cell* **37**, 34–46 (2016).

28. Portereiko, M. F. *et al.* AGL80 Is Required for Central Cell and Endosperm

Development in Arabidopsis. *The Plant Cell* **18**, 1862–1872 (2006).

29. Bemer, M., Wolters-Arts, M., Grossniklaus, U. & Angenent, G. C. The MADS

Domain Protein DIANA Acts Together with AGAMOUS-LIKE80 to Specify the

Central Cell in Arabidopsis Ovules. *The Plant Cell* **20**, 2088–2101 (2008).

30. Bateman, R. M. & DiMICHELE, W. A. Heterospory: The Most Iterative Key

Innovation in the Evolutionary History of the Plant Kingdom. *Biological Reviews*

69, 345–417 (1994).

31. Feng, X., Zilberman, D. & Dickinson, H. A conversation across generations:

soma-germ cell crosstalk in plants. *Dev Cell* **24**, 215–225 (2013).

32. Schmidt, A., Schmid, M. W. & Grossniklaus, U. Plant germline formation:

common concepts and developmental flexibility in sexual and asexual

reproduction. *Development* **142**, 229–241 (2015).

33. Petrella, R., Cucinotta, M., Mendes, M. A., Underwood, C. J. & Colombo, L. The

emerging role of small RNAs in ovule development, a kind of magic. *Plant*

Reprod **34**, 335–351 (2021).

34. Yang, W. C., Ye, D., Xu, J. & Sundaresan, V. The SPOROCYTELESS gene of

Arabidopsis is required for initiation of sporogenesis and encodes a novel nuclear

protein. *Genes Dev* **13**, 2108–2117 (1999).

35. Schiefthaler, U. *et al.* Molecular analysis of NOZZLE, a gene involved in pattern

formation and early sporogenesis during sex organ development in Arabidopsis

thaliana. *Proc Natl Acad Sci U S A* **96**, 11664–11669 (1999).

36. Mendes, M. A. *et al.* The RNA-dependent DNA methylation pathway is required

to restrict SPOROCYTELESS/NOZZLE expression to specify a single female

germ cell precursor in Arabidopsis. *Development* **147**, dev194274 (2020).

37. Zhao, D.-Z., Wang, G.-F., Speal, B. & Ma, H. The excess microsporocytes1 gene

encodes a putative leucine-rich repeat receptor protein kinase that controls

somatic and reproductive cell fates in the Arabidopsis anther. *Genes Dev* **16**,

2021–2031 (2002).

38. Canales, C., Bhatt, A. M., Scott, R. & Dickinson, H. EXS, a putative LRR receptor

kinase, regulates male germline cell number and tapetal identity and promotes

seed development in Arabidopsis. *Curr Biol* **12**, 1718–1727 (2002).

39. Jia, G., Liu, X., Owen, H. A. & Zhao, D. Signaling of cell fate determination by the

TPD1 small protein and EMS1 receptor kinase. *Proc Natl Acad Sci U S A* **105**,

2220–2225 (2008).

40. Nonomura, K.-I. *et al.* A germ cell specific gene of the ARGONAUTE family is

essential for the progression of premeiotic mitosis and meiosis during

sporogenesis in rice. *Plant Cell* **19**, 2583–2594 (2007).

41. Singh, M. *et al.* Production of viable gametes without meiosis in maize deficient

for an ARGONAUTE protein. *Plant Cell* **23**, 443–458 (2011).

42. Su, Z. *et al.* The THO Complex Non-Cell-Autonomously Represses Female

Germline Specification through the TAS3-ARF3 Module. *Curr Biol* **27**, 1597-

1609.e2 (2017).

43. Martínez, G., Panda, K., Köhler, C. & Slotkin, R. K. Silencing in sperm cells is

directed by RNA movement from the surrounding nurse cell. *Nat Plants* **2**, 16030

(2016).

44. Pagnussat, G. C., Alandete-Saez, M., Bowman, J. L. & Sundaresan, V. Auxin-

dependent patterning and gamete specification in the Arabidopsis female

gametophyte. *Science* **324**, 1684–1689 (2009).

45. Panoli, A. *et al.* Auxin Import and Local Auxin Biosynthesis Are Required for

Mitotic Divisions, Cell Expansion and Cell Specification during Female

Gametophyte Development in Arabidopsis thaliana. *PLoS One* **10**, e0126164

(2015).

46. Gómez, J. F., Talle, B. & Wilson, Z. A. Anther and pollen development: A

conserved developmental pathway. *Journal of Integrative Plant Biology* **57**, 876–

891 (2015).

47. Ariizumi, T. & Toriyama, K. Genetic regulation of sporopollenin synthesis and

pollen exine development. *Annu Rev Plant Biol* **62**, 437–460 (2011).

48. Wilson, Z. A., Morroll, S. M., Dawson, J., Swarup, R. & Tighe, P. J. The

Arabidopsis MALE STERILITY1 (MS1) gene is a transcriptional regulator of male

gametogenesis, with homology to the PHD-finger family of transcription factors.

Plant J **28**, 27–39 (2001).

49. Sorensen, A.-M. *et al.* The Arabidopsis ABORTED MICROSPORES (AMS) gene

encodes a MYC class transcription factor. *Plant J* **33**, 413–423 (2003).

50. Zhang, W. *et al.* Regulation of Arabidopsis tapetum development and function by

DYSFUNCTIONAL TAPETUM1 (DYT1) encoding a putative bHLH transcription

factor. *Development* **133**, 3085–3095 (2006).

51. Long, J. *et al.* Nurse cell--derived small RNAs define paternal epigenetic

inheritance in Arabidopsis. *Science* **373**, eabh0556 (2021).

52. Tanaka, I. & Ito, M. Induction of typical cell division in isolated microspores of

Lilium longiflorum and Tulipa gesneriana. *Plant Science Letters* **17**, 279–285

(1980).

53. Moreno, R. M., Macke, F., Alwen, A. & Heberle-Bors, E. In-situ seed production

after pollination with in-vitro-matured, isolated pollen. *Planta* **176**, 145–148

(1988).

54. Chaudhury, A. M. Nuclear Genes Controlling Male Fertility. *Plant Cell* **5**, 1277–

1283 (1993).

55. Mercier, R. & Grelon, M. Meiosis in plants: ten years of gene discovery.

Cytogenet Genome Res **120**, 281–290 (2008).

56. Blackmore, S., Wortley, A. H., Skvarla, J. J. & Rowley, J. R. Pollen wall

development in flowering plants. *New Phytol* **174**, 483–498 (2007).

57. Scott, R. J., Spielman, M. & Dickinson, H. G. Stamen structure and function.

Plant Cell **16 Suppl**, S46-60 (2004).

58. Bencivenga, S., Colombo, L. & Masiero, S. Cross talk between the sporophyte

and the megagametophyte during ovule development. *Sex Plant Reprod* **24**,

113–121 (2011).

59. Figueiredo, D. D. & Köhler, C. Bridging the generation gap: communication

between maternal sporophyte, female gametophyte and fertilization products.

Curr Opin Plant Biol **29**, 16–20 (2016).

60. McCormick, S. Male Gametophyte Development. *Plant Cell* **5**, 1265–1275

(1993).

61. Owen, H. A. & Makaroff, C. A. Ultrastructure of microsporogenesis and

microgametogenesis inArabidopsis thaliana (L.) Heynh. ecotype Wassilewskija

(Brassicaceae). *Protoplasma* **185**, 7–21 (1995).

62. Twell, D. Use of a nuclear-targeted β-glucuronidase fusion protein to

demonstrate vegetative cell-specific gene expression in developing pollen. *The*

Plant Journal https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1365-

313X.1992.00887.x (1992).

63. Dobritsa, A. A. & Coerper, D. The novel plant protein INAPERTURATE

POLLEN1 marks distinct cellular domains and controls formation of apertures in

the Arabidopsis pollen exine. *Plant Cell* **24**, 4452–4464 (2012).

64. Ma, H. Molecular genetic analyses of microsporogenesis and

microgametogenesis in flowering plants. *Annu Rev Plant Biol* **56**, 393–434

(2005).

65. Proost, S. & Mutwil, M. CoNekT: an open-source framework for comparative

genomic and transcriptomic network analyses. *Nucleic Acids Res* **46**, W133–

W140 (2018).

66. Honys, D. & Twell, D. Comparative analysis of the Arabidopsis pollen

transcriptome. *Plant Physiol* **132**, 640–652 (2003).

67. Pina, C., Pinto, F., Feijó, J. A. & Becker, J. D. Gene Family Analysis of the

Arabidopsis Pollen Transcriptome Reveals Biological Implications for Cell

Growth, Division Control, and Gene Expression Regulation. *Plant Physiol* **138**,

744–756 (2005).

68. Ichino, L. *et al.* Single-nucleus RNA-seq reveals that MBD5, MBD6, and

SILENZIO maintain silencing in the vegetative cell of developing pollen. *Cell Rep*

41, 111699 (2022).

69. Honys, D. *et al.* Identification of microspore-active promoters that allow targeted

manipulation of gene expression at early stages of microgametogenesis in

Arabidopsis. *BMC Plant Biol* **6**, 31 (2006).

70. Raudvere, U. *et al.* g:Profiler: a web server for functional enrichment analysis and

conversions of gene lists (2019 update). *Nucleic Acids Res* **47**, W191–W198

(2019).

71. Moon, J., Parry, G. & Estelle, M. The ubiquitin-proteasome pathway and plant

development. *Plant Cell* **16**, 3181–3195 (2004).

72. Smalle, J. & Vierstra, R. D. The ubiquitin 26S proteasome proteolytic pathway.

Annu Rev Plant Biol **55**, 555–590 (2004).

73. Mladek, C., Guger, K. & Hauser, M.-T. Identification and characterization of the

ARIADNE gene family in Arabidopsis. A group of putative E3 ligases. *Plant*

Physiol **131**, 27–40 (2003).

74. Gagne, J. M., Downes, B. P., Shiu, S.-H., Durski, A. M. & Vierstra, R. D. The F-

box subunit of the SCF E3 complex is encoded by a diverse superfamily of genes

in Arabidopsis. *Proc Natl Acad Sci U S A* **99**, 11519–11524 (2002).

75. Kuroda, H. *et al.* Classification and expression analysis of Arabidopsis F-box-

containing protein genes. *Plant Cell Physiol* **43**, 1073–1085 (2002).

76. Ikram, S. *et al.* Functional redundancy and/or ongoing pseudogenization among

F-box protein genes expressed in Arabidopsis male gametophyte. *Plant Reprod*

27, 95–107 (2014).

77. Branscheid, A. *et al.* SKI2 mediates degradation of RISC 5'-cleavage fragments

and prevents secondary siRNA production from miRNA targets in Arabidopsis.

Nucleic Acids Res **43**, 10975–10988 (2015).

78. Zhang, X. *et al.* Suppression of endogenous gene silencing by bidirectional

cytoplasmic RNA decay in Arabidopsis. *Science* **348**, 120–123 (2015).

79. Fu, H., Doelling, J. H., Arendt, C. S., Hochstrasser, M. & Vierstra, R. D.

Molecular organization of the 20S proteasome gene family from Arabidopsis

thaliana. *Genetics* **149**, 677–692 (1998).

80. Hatsugai, N. *et al.* A novel membrane fusion-mediated plant immunity against

bacterial pathogens. *Genes Dev* **23**, 2496–2506 (2009).

81. Krishnan, A., Iyer, L. M., Holland, S. J., Boehm, T. & Aravind, L. Diversification of

AID/APOBEC-like deaminases in metazoa: multiplicity of clades and widespread

roles in immunity. *Proc Natl Acad Sci U S A* **115**, E3201–E3210 (2018).

82. Yang, Z. PAML 4: phylogenetic analysis by maximum likelihood. *Mol Biol Evol*

24, 1586–1591 (2007).

83. Xu, G., Ma, H., Nei, M. & Kong, H. Evolution of F-box genes in plants: Different

modes of sequence divergence and their relationships with functional

diversification. *Proc Natl Acad Sci U S A* **106**, 835–840 (2009).

84. Hua, Z., Zou, C., Shiu, S.-H. & Vierstra, R. D. Phylogenetic Comparison of F-Box

(FBX) Gene Superfamily within the Plant Kingdom Reveals Divergent

Evolutionary Histories Indicative of Genomic Drift. *PLOS ONE* **6**, e16219 (2011).

85. Kaessmann, H. Origins, evolution, and phenotypic impact of new genes.

Genome Res **20**, 1313–1326 (2010).

86. Wu, D.-D. *et al.* "Out of Pollen" Hypothesis for Origin of New Genes in Flowering

Plants: Study from Arabidopsis thaliana. *Genome Biology and Evolution* **6**, 2822–

2829 (2014).

87. Cui, X. *et al.* Young Genes out of the Male: An Insight from Evolutionary Age

Analysis of the Pollen Transcriptome. *Mol Plant* **8**, 935–945 (2015).

88. Nelms, B. & Walbot, V. Gametophyte genome activation occurs at pollen mitosis

I in maize. *Science* **375**, 424–429 (2022).

89. Barinova, I. *et al.* Antirrhinum majus microspore maturation and transient

transformation in vitro. *J Exp Bot* **53**, 1119–1129 (2002).

90. Delph, L. F. Pollen competition is the mechanism underlying a variety of

evolutionary phenomena in dioecious plants. *New Phytol* **224**, 1075–1079

(2019).

91. Zhao, L. & Kunst, L. SUPERKILLER Complex Components Are Required for the

RNA Exosome-Mediated Control of Cuticular Wax Biosynthesis in Arabidopsis

Inflorescence Stems. *Plant Physiol* **171**, 960–973 (2016).

92. Toufighi, K., Brady, S. M., Austin, R., Ly, E. & Provart, N. J. The Botany Array

Resource: e-Northerns, Expression Angling, and promoter analyses. *Plant J* **43**,

153–163 (2005).

93. Schmid, M. *et al.* A gene expression map of Arabidopsis thaliana development.

Nat Genet **37**, 501–506 (2005).

94. Alexander, M. P. Differential staining of aborted and nonaborted pollen. *Stain*

Technol **44**, 117–122 (1969).

95. Clough, S. J. & Bent, A. F. Floral dip: a simplified method for Agrobacterium-

mediated transformation of Arabidopsis thaliana. *Plant J* **16**, 735–743 (1998).

96. Karimi, M., Depicker, A. & Hilson, P. Recombinational cloning with plant gateway

vectors. *Plant Physiol* **145**, 1144–1154 (2007).

97. Reddy, T. V., Kaur, J., Agashe, B., Sundaresan, V. & Siddiqi, I. The DUET gene

is necessary for chromosome organization and progression during male meiosis

in Arabidopsis and encodes a PHD finger protein. *Development* **130**, 5975–5987

(2003).

98. Sanders, P. M. *et al.* Anther developmental defects in Arabidopsis thaliana male-

sterile mutants. *Sex Plant Reprod* **11**, 297–322 (1999).

99. Schneitz, K., Hülskamp, M. & Pruitt, R. E. Wild-type ovule development in

Arabidopsis thaliana: a light microscope study of cleared whole-mount tissue.

The Plant Journal **7**, 731–749 (1995).

100.   Andreuzza, S., Nishal, B., Singh, A. & Siddiqi, I. The Chromatin Protein

DUET/MMD1 Controls Expression of the Meiotic Gene TDM1 during Male

Meiosis in Arabidopsis. *PLoS Genet* **11**, e1005396 (2015).

101. Singh, D. K., Spillane, C. & Siddiqi, I. PATRONUS1 is expressed in meiotic

prophase I to regulate centromeric cohesion in Arabidopsis and shows synthetic

lethality with OSD1. *BMC Plant Biol* **15**, 201 (2015).

102. Siddiqi, I., Ganesh, G., Grossniklaus, U. & Subbiah, V. The dyad gene is

required for progression through female meiosis in Arabidopsis. *Development*

127, 197–207 (2000).

103. Schwartz, J. *et al.* Removing Stripes, Scratches, and Curtaining with

Nonrecoverable Compressed Sensing. *Microscopy and Microanalysis* **25**, 705–

710 (2019).

104. Jiao, W.-B. & Schneeberger, K. Chromosome-level assemblies of multiple

Arabidopsis genomes reveal hotspots of rearrangements with altered

evolutionary dynamics. *Nat Commun* **11**, 989 (2020).

105. Quinlan, A. R. BEDTools: The Swiss-Army Tool for Genome Feature

Analysis. *Curr Protoc Bioinformatics* **47**, 11.12.1-34 (2014).

106. Garrison, E. & Marth, G. Haplotype-based variant detection from short-read

sequencing. Preprint at https://doi.org/10.48550/arXiv.1207.3907 (2012).

107. Mansfeld, B. N. & Grumet, R. QTLseqr: An R Package for Bulk Segregant

Analysis with Next-Generation Sequencing. *Plant Genome* **11**, (2018).

108. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

109. Walker, J. *et al.* Lineage-specific DNA methylation regulates Arabidopsis

sexual development. *Nat Genet* **50**, 130–137 (2018).

110. Xu, D. *et al.* BBX21, an Arabidopsis B-box protein, directly activates HY5 and

is targeted by COP1 for 26S proteasome-mediated degradation. *Proc Natl Acad*

Sci U S A **113**, 7655–7660 (2016).

111.   Altschul, S. Gapped BLAST and PSI-BLAST: a new generation of protein

733   database search programs. *Nucleic Acids Research* **25**, 3389–3402 (1997).

734   112.   Potter, S. C. *et al.* HMMER web server: 2018 update. *Nucleic Acids Res* **46**,

735   W200–W204 (2018).

736   113.   Zimmermann, L. *et al.* A Completely Reimplemented MPI Bioinformatics

737   Toolkit with a New HHpred Server at its Core. *J Mol Biol* **430**, 2237–2243 (2018).

738   114.   Mistry, J. *et al.* Pfam: The protein families database in 2021. *Nucleic Acids*

739   *Res* **49**, D412–D419 (2021).

740   115.   Berman, H. M. *et al.* The Protein Data Bank. *Nucleic Acids Res* **28**, 235–242

741   (2000).

742   116.   Drozdetskiy, A., Cole, C., Procter, J. & Barton, G. J. JPred4: a protein

743   secondary structure prediction server. *Nucleic Acids Res* **43**, W389-394 (2015).

744   117.   Baek, M. *et al.* Accurate prediction of protein structures and interactions using

745   a three-track neural network. *Science* **373**, 871–876 (2021).

746   118.   Goodstein, D. M. *et al.* Phytozome: a comparative platform for green plant

747   genomics. *Nucleic Acids Res* **40**, D1178-1186 (2012).

748   119.   Lyons, E. *et al.* Finding and comparing syntenic regions among Arabidopsis

749   and the outgroups papaya, poplar, and grape: CoGe with rosids. *Plant Physiol*

750   **148**, 1772–1781 (2008).

751   120.   Numa, H. & Itoh, T. MEGANTE: A Web-Based System for Integrated Plant

752   Genome Annotation. *Plant Cell Physiol* **55**, e2 (2014).

753   121.   Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software

754   version 7: improvements in performance and usability. *Mol Biol Evol* **30**, 772–780

755   (2013).

122.   Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-

analysis of large phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).

123.   Gao, F. *et al.* EasyCodeML: A visual tool for analysis of selection using

CodeML. *Ecol Evol* **9**, 3891–3898 (2019).

124.   Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for

Phylogenetic Inference in the Genomic Era. *Mol Biol Evol* **37**, 1530–1534 (2020).

125.   Guindon, S. *et al.* New algorithms and methods to estimate maximum-

likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst Biol* **59**,

307–321 (2010).

126.   Lechner, M. *et al.* Proteinortho: Detection of (Co-)orthologs in large-scale

analysis. *BMC Bioinformatics* **12**, 124 (2011).

Figure Legends

Fig. 1: *SHUKR* is required for pollen development. (a) Schematic showing

reduction of the gametophyte in flowering plants. (b) Sterility (short siliques) and

reduced pollen viability (green staining) in *skr*, and complementation by a genomic

SKR transgene. (c) Tables showing reduced seed set, male-specific sterility, and

recessive Mendelian segregation for *skr*; numbers in brackets are those expected for

a 3:1 ratio and the χ^2 test p value. (d) RT-PCR analysis of SKR expression in

vegetative and reproductive tissues. (e) GUS staining for *proSKR:GFP-GUS*

transgene expression in successive unopened buds. (f) GUS staining for

proSKR:GFP-GUS expression in meiotic stage anthers. (g) anti-GFP antibody staining

for *proSKR:GFP-GUS* expression in male meiosis and microspores. (h) Normal tetrad

formation in *skr*. Scale bar: (e) 0.5 mm, (f) 20 µm.

Fig. 2: Staging of *skr* phenotype and SKR protein expression. (a) Meiotic chromosome development and segregation in wild type (a-j) and corresponding stages in *skr* (a'-j'). a-d,a'-d' prophase I; e,e' metaphase I; f,f' telophase I; g,g' metaphase II; h,h' anaphase II; i,i' telophase II; j,j' tetrad. 5846 meiocytes were scored for *skr* and 2281 for WT. (b) Whole mount DIC images of cleared anthers from successive stage buds; arrowheads indicate degenerating microspores starting in anthers containing early microspores. (c) Anti-GFP staining of male meiosis stages in a *proSKR:GFP-gSKR* complementing line; asterisks: somatic/tapetal cells; arrowheads: older uninucleate microspores with low GFP signal. Scale bar: (a) 5 µm, (b,c) 20 µm.

Fig. 3: Ultrastructure and marker expression reveal early gametogenesis defects in *skr*. (a-c) Ultrastructure of tetrad and early uninucleate microspores showing onset of mutant phenotype at tetrad stage in *skr*. (a) WT: Normal tetrad and microspore; *skr*: normal tetrad, degenerating microspore with retracted PM. (b) *skr* tetrad in which one spore shows PM retraction from callose wall. (c) Magnified image of boxed region from (b) showing PM retraction from callose wall (Cal) and disrupted cell wall material (arrowhead). (d) Quantitation of normal spores and spores showing PM retraction and broken cell wall material (R&B) in tetrads (Fisher's exact test p < 0.001). (e) Presence of two LAT52:GFP signals in a proportion of *skr* microspores (Fisher's exact test p < 0.001). Scale bar (a): Tetrad images 5 µm; microspore images 2 µm; (b) 5 µm; (C) 1 µm.

Fig. 4: *SHUKR* is a gametogenesis inhibitor. (a) Reciprocal crosses comparing transmission of *skr* and wild type alleles; numbers in brackets are those expected for equal transmission and χ^2 test p-value. (b) Tissue/cell enrichment categories for

upregulated genes in *skr*. (c) Pollen viability in WT and *skr* plants transgenic for

proMSP1:SKRcds or control. Data are pools of 4-10 anthers from 5 independent

transgenic lines in each case. Scale bar (c) 100 µm.

Fig. 5: *SHUKR* regulates the proteasome in pollen development. (a) Gene family

categories for upregulated genes in *skr*. (b) Stagewise expression of an

At3g28330:2xnlsGFP transgene during meiosis and in microspores for *skr* and

wildtype. Frames: a,f meiocytes; b,g tetrad; c-e,h microspores; upper half: GFP, lower

half: DIC. (c) Suppression of *skr* male sterility by *ski2*. Bar graph shows viable pollen

counts in single anthers from 2-4 plants. (d) Downregulation of proteasome genes in

skr ski2. (e) Mean viable pollen counts/anther in independent *skr; proSKR:PBE1RNAi*

and control *T1* transgenic plants. (f) Pollen viability increase in *skr* following

proteasome inhibitor treatment in 3 replicate experiments. (g) Model for sporophytic

control of pollen development by *SHUKR*. During meiosis *SHUKR* establishes a

programme that determines timing of expression of F-box and other UPS genes

following meiosis. Scale bar: (b) 5 µm; (c) plant 1 cm, anthers 100 µm. Abbreviations:

UNM, uninucleate microspore; BCP, bicellular pollen; TCP, tricellular pollen; MP,

mature pollen; SGT, sporophyte-gametophyte transition.

Fig. 6: Sequence and evolutionary analysis of the SHUKR domain. (a) Multiple

sequence alignment of the SHUKR domain. Lines mark CxC and aromatic dyad. (b)

Phylogenetic tree of the SHUKR domain from 119 species. Dark green,

Brassicaceae+Cleomaceae; Light green, Asteraceae. (c) Distribution of mean-

centered Maximum Likelihood (ML) distance differences. (d) Columnwise Shannon

entropy distributions for four plant families. P-value calculated using Kruskal-Wallis

test for Brassicaceae and Asteraceae relative to the other two families.

Fig. 7: Diversifying selection in SHUKR regulated F-box genes. (a) Enrichment for

short taxonomic scale (STS) F-box genes (R.F. = 1.2) and underenrichment for long

taxonomic scale (LTS) genes (R.F. = 0.1) among *skr* upregulated set (p < 1.49e-05).

(b) 37 genes were unique to the Brassicaceae lineage of which 5 were found only in

Arabidopsis thaliana. 15 genes were present in *Arabidopsis thaliana* and at least one

other species but absent in *Arabidopsis lyrata*. (c) Phylogenetic tree of Arabidopsis F-

box genes showing *skr* upregulated, LTS, and STS sets. (d) Violin plot of K_a/K_s for *skr*

upregulated F-box and total F-box genes. Other eudicots: *Cucumis sativus*, *Glycine*

max, *Manihot esculenta*, *Mimulus guttatus*, *Populus trichocarpa*, *Ricinus communis*,

Vitis vinifera. Abbreviations: Al *Arabidopsis lyrata*, At *Arabidopsis thaliana*, Cp *Carica*

papaya.

Extended Data Figure Legends

Extended Data Fig. 1:

(a) Gene diagram of *SKR* showing location of insertions: *skr-1* (GT_5_101780), *skr-2*

(CSHL_GT5061), *skr-3* (SALKSeq_034497.1). (b) Toluidine blue stained tetrads. (c)

Inflorescence stem and Alexander staining showing recessive male sterility of *skr.* (d)

Complementation of *skr-3* by a *gSKR* transgene. Scale bar: 1 cm for inflorescence,

100 μm for anthers.

Extended Data Fig. 2:

Profiling of SKR expression in male germline development.

TPM values for SKR were collected from the CoNekt database, except for meiocytes.

Meiocyte TPM value were taken from (Walker 2018)[21]. UNM – Uninucleate

microspore; BCP – Bicellular pollen; TCP – Tricellular pollen; MP – Mature pollen.

Extended Data Fig. 3:

Sections of anther development stages.

Corresponding stages are shown for wild type (A-D, I-L) and *skr* (E-H, M-P). Stages

are numbered according to (Sanders 1999)[9]. Abbreviations: E epidermis, En

endothecium, ML middle layer, T tapetum, Td tetrad, Msp microspore, DM

degenerating microspore, PG pollen grain, St stomium, MD microspore debris. Scale

bar 20 µm.

Extended Data Fig. 4:

Staging of microspore degeneration in *skr*.

Microspore stage and phenotypes are shown together with stage of ovules from the

same bud (Schneitz 1995)[10] as a reference for comparison to wild type

Extended Data Fig. 5:

Precocious expression of F-box gene:nlsGFP fusions in meiosis in *skr*.

(a) At5g02980. A-C, A'-C': wild type; D-F, D'-F': skr; A,A',D,D' meiocytes; B,B',E,E':

tetrad; C,C',F,F': uninucleate microspores. (b) At5g62510. A-E, A'-E': wild type; F-H,

F'-H': skr; A,A',F,F' meiocytes; B,B',G,G': tetrad; C-E, C'-E',H,H': uninucleate

microspores. A-H: GFP fluorescence; A'-H': corresponding DIC. GFP expression in

skr is first detected in meiocytes (At5g02980) or tetrad stage At5g62510) whereas in

wild type earliest GFP signal was in released microspores. Scale bar: 5 µm

Extended Data Fig. 6:

ski2 is a suppressor of *skr*.

(a) Mendelian segregation of the suppressor mutation in the 08-10 *skr* suppressor line

(b) A C>T transition in the *ski2* allele (named *ski2-7*) present in the 08-10 *skr* suppressor line results in a TGA termination codon at position 518. (c) Restoration of fertility and pollen viability in *skr-3* by the independent allele *ski2-6*. Scale bar: inflorescence 1 cm; anthers 100 µm.

Extended Data Fig. 7:

Phylogenetic tree of the SHUKR family from eudicots.

119 representatives from 85 species were used for tree construction using only the SHUKR domain that is shared by all representatives of the family. Genbank ID and species are indicated. Brassicaceae+Cleomaceae (dark green) and Asteraceae (light green)

Extended Data Fig. 8:

Phylogenetic tree of Brassicales *SHUKR* and *SHUKR-like* (*SKL*) genes used for branch-site test.

The *SHUKR* gene family is represented by a single gene in *Carica papaya*, whereas Brassicaceae and Cleomaceae experienced a gene duplication event prior to their divergence. CDS were aligned using MAFFT, and RAxML was used to build the phylogenetic tree with 1000 bootstraps. The red branch was selected as foreground for *SKR*. Branch-site test for positive selection was performed using the CODEML package implemented in EASYCoDEML. Likelihood Ratio Test (LRT) results of Model

A vs Model A null (Extended Data Table 1), gave a p-value of 0.0031 for *SKR*. The tree was unrooted.

Extended Data Table 1:

Branch-site test results for positive selection in the SKR branch in the Brassicaceae. Branch site models postulate three classes of sites: those under purifying selection ($0< \omega <1$), neutral ($\omega=1$), and those under positive selection ($\omega>1$) in the foreground branch being tested (Yang 2007)[53]. Model A is compared to Model A null in which the third class is set to $\omega=1$. Positive sites are indicated together with posterior probabilities. P-value for likelihood ratio test (LRT) results are calculated from the χ^2 distribution taking as critical value $\chi^2= 2(LnL_{MA}-LnL_{MAnull})$ and dividing by 2 (Yang 2007)[53].

Methods

Plant Material and Growth Conditions

Arabidopsis mutant strains used in the study were *skr-1* (GT_5_101780), *skr-2* (CSHL_GT5061) and *skr-3* (SALKSeq_034497.1), and *ski2-6* (SALK_122393)[91]. *skr-1* and *skr-2* alleles are in Ler ecotype and have a Ds insertion in AT2G38690; *skr-3* is in Col-0 ecotype. Plants were grown at 22 °C under 16 hours of light and 8 hours of dark in a 1:1:1 proportion of peat, perlite, and vermiculite soaked with 0.5x Murashige Skoog (MS) solution initially followed by irrigation during growth with water. For growth on plates and selection of transformants, seeds were surface sterilized and plated on MS plates supplemented with 2% sucrose and appropriate antibiotics.

Coexpression screen for meiotic genes

We selected 23 known meiotic genes (*DUET/MMD1, MS5/TDM1, DMC1, TAM, SDS,*

AML5, RAD51, RAD50, ASY1, ZYP1, REC8, SPO11-1, SPO11-2, PHS1, MRE11,

MND1, OSD1, NBS1, MSH4, MSH5, HOP2, MER3, PRD1), and each of these genes

was used as a query in the Expression Angler program[92] to identify co-expressed

genes. For each query, the top 100 hits were selected from each of three data sets,

NASCArrays, Bio-analytic resource, and AtGenExpress Plus-Extended tissue

compendium. We selected those genes common to three data sets and which

appeared at least twice in independent queries. Single copy genes were selected

using the Arabidopsis single copy genes list from the University of California, Davis

(http://www.cgpdb.ucdavis.edu/COS_Arabidopsis/). We manually selected 29 genes

that showed floral stage 9 peak/specific expression corresponding to meiosis stages

from AtGenExpress Visualization Tool[93]. We tested one to two insertion alleles for

each of the selected genes and for each of these alleles about 20 plants were

genotyped by PCR to identify homozygous mutant plants and these were phenotyped

by examining silique length and pollen viability by Alexander stain[94].

Plasmid constructs and transgenics

For complementation, a genomic fragment of 2220 bp (gSKR) comprising the *SKR*

gene including 3'UTR was amplified from Ler ecotype and cloned into a modified

binary vector pZP222 by Gateway cloning (Invitrogen). The binary construct was

introduced into Agrobacterium strain AGL1 and transformed into *skr-1/+* and *skr-3/+*

plants by in planta transformation[95]. *skr* mutant T1 plants were genotyped by PCR and

tested for complementation. Plant vectors were ordered from

(https://gatewayvectors.vib.be/index.php/gateway-vectors). The pro*SKR*:*GFP-gSKR*

gene fusion construct was made by assembling the SKR promoter, GFP, and SKR

genomic coding region plus 3'UTR using three-way Gateway technology in the binary destination vector pB7m34GW,0[96]. The pro*SKR:nlsGFP-GUS* construct comprised 1170 bp genomic DNA from 690 bp upstream of ATG upto 30 bp of the second exon fused in frame with GFP-GUS in the pBGWFS7 vector. F-box gene nlsGFP fusions were assembled in the pBnRGW_Redseed vector and introduced into plants. Single locus insert transgenic skr/+ lines were made homozygous for the transgene and examined in T4 for GFP fluorescence. The hairpin RNAi construct for PBE1[80] was expressed under control of SKR regulatory elements comprising promoter and 3'UTR and contained the PDK intron from pBm42GWIWG8_1. The construct and a noRNAi control were introduced into *skr/+* plants and transgenics that were *skr* mutant were analysed for pollen viability. The *skr-1* mutant allele was used for all studies unless indicated otherwise. proMSP1:SKRcds constructs were made via overlap extension PCR, cloned into pENTR D TOPO vector using NotI and AscI restriction mediated cloning, and further cloned into pZP222 via Gateway cloning.

Cytology and imaging

For Alexander staining, anthers from unopened buds were dissected out on a slide, placed under a coverslip, and Alexander stain was added from the side to fill the entire coverslip followed by sealing using nail-polish. The slides were incubated at room temperature for 3-4 days and visualized under 10x /20 x objective.Whole anther and ovule clearing was performed as in[97], anther staging of plastic sections was performed as in[98] and ovule staging as in[99]. Immunostaining for GFP:SKR in meiosis was performed as in[100]. Acid chromosomal spreads were prepared as described in[101]. GFP / DAPI fluorescence was visualized after mounting of meiotic anthers under the coverslip with 30-50 % glycerol / Antifade. ImageJ and Photoshop were used to edit

and assemble images. Electron microscopy sample preparation and imaging was

done as in[102] and stripe artifacts in the EM images were removed using the method

mentioned in[103].

Suppressor screening and analysis

skr-1 mutant seeds (M0) were bulked from multiple plants and 0.3 % EMS solution

was used to mutagenise the M0 seeds for 5 hours followed by extensive washing with

water. Seeds from 934 M1 plants were collected individually and M2 families were

screened for fertility restoration. One suppressor line 08-10 was identified and

backcrossed to *skr* mutants to get BC1 F1 seeds. Seeds from BC1 F1 plants were

collected and bulk segregant analysis on BC1 F2 plants was performed on fertile and

sterile pools. Genomic DNA was isolated from the fertile and sterile pools (50 plants

each) and used for whole genome sequencing.

To map the suppressor, the parental un-mutagenised *skr* genome was sequenced and

skr specific reference genome was made by replacing the SNPs of *skr* to Ler-0

genome[104] sequence using the *bedtools – consensus*[105]. Variants were called for the

fertile and sterile bulk reads from the *skr* reference genome using *freebayes*[106] and

indels were removed. SNPindex was calculated using the R package QTLSeqr[107] for

the fertile bulk and sterile bulk. SNPs with a deltaSNP of fertile to sterile score greater

than 0.5 were analysed for genic perturbations. The 08-10 line had an in frame stop

codon in the SKI2 gene, at amino acid position 518. The suppressor phenotype was

confirmed by generating the double mutant using the independent alleles *ski2-6* and

skr-3. The *ski2* allele in line 08-10 was named *ski2-7.*

1004

Expression profiling

RNA from meiotic stage anthers was isolated using Qiagen kits for *skr-1,* Ler and *skr-1 ski2-7*. RNA sequencing was performed on Illumina platforms following the manufacturer's instructions. Differentially expressed genes were found using DESeq2 keeping a FDR adjusted *P* value < 0.05[108]. Validation for eight upregulated F-box genes was performed by RT-qPCR (Supplementary Figure 3) which confirmed the RNAseq results. The tissue-specific expression analysis of deregulated genes was performed by downloading TPM values for each gene from CoNekT (https://evorepro.sbs.ntu.edu.sg) database[65]. Meiocyte RNAseq TPM value[109] was combined with the CoNekT dataset. The tissue enrichment score was calculated by row normalization using the largest TPM score. Enrichment score of 1, in a tissue/cell type was taken to represent maximal expression in that type. The number of non-overlapping genes with enrichment score 1, was calculated and plotted for meiocytes, UNM, BCP, TCP/MP and others.

Proteasome inhibitor treatment

MG132 and MG115 stocks at 20 mM concentration were made in DMSO. Inhibitor solutions were used at 100 μM concentration[110] by dissolving in half strength MS where 0.5 % DMSO was present. Control solution was 0.5 % DMSO in half strength MS. Sections of freshly cut *skr* inflorescence stems were placed in 200 μl inhibitor or control solution in Eppendorf tubes for 3 hrs, followed by 60 hrs of recovery in half strength MS solution. Anthers were dissected and viable pollens were identified by Alexander staining. At least 100 anthers were analysed per sample (treatment and control) per experiment. Number of anthers with 10 or more viable pollen were counted per 100 anthers for treatment and control and the ratio was calculated. The experiment

1030 was conducted blind and repeated thrice. The ratio paired t test (two-tailed) on the

1031 logarithm of ratios was used to calculate p values (GraphPad).

1032

Sequence analysis

Sequence searches to detect homologs of the SHUKR domain were conducted using

the PSI-BLAST (RRID:SCR_001010)[111] and JACKHMMER (RRID:SCR_005305)[112]

programs with profile-inclusion threshold of expect (e)-value at 0.01 against the non-

redundant database of National Center for Biotechnology Information (NCBI) clustered

down to 50%. Clustering of proteins based on bit score density and length of aligned

sequence was performed using the MMseqs program. Remote homology searches

were performed using profile-profile comparisons with HHpred program

(RRID:SCR_010276)[113] against profile libraries from the PFAM (RRID:

SCR_004726)[114] and PDB (RRID:SCR_012820)[115] databases. Multiple sequence

alignments were built using Mafft program with maxiterate parameter set to 3000

followed by manual adjustments based on secondary structure prediction. Secondary

structures were predicted using the JPred (RRID:SCR_016504)[116] and

RoseTTaFold[117] programs. The globular domains in the protein were objectively

identified using Shannon entropy-based compositional analysis with the SEG program

with a window size of 45, low-cut of 3.4, high-cut of 3.75, and minimal length of the

high-complexity segment of 15 residues.

1050

The analysis of the Shannon entropy (H_j) for a given multiple sequence alignment was

performed using the equation:

$$H_j = -\sum_{i=1}^{M} P_i(log_2(P_i))$$

P is the fraction of residues of amino acid type i and M is the number of amino acid

types. The Shannon entropy for the jth position in the alignment can range from 0 (only

one residue at that position) to 4.32 (all 20 residues equally represented at that

position). The reduced alphabet was created by the following grouping: h= L, I, M, V,

A; a= F, Y, W; b=K, R, H; n=E, D; p=Q, N; o=T, C, S; t=P, G, d=- (gap).

1059

Phylogenetic analysis

SKR and SKL coding sequences of different species were extracted from Genbank,

Phytozome, and CoGeBLAST[118,119] (Supplementary File 1). Genomic fragments were

annotated to CDS by the online tool MEGANTE[120]. For the branch site analysis,

MAFFT was used to build the alignment and RAxML was used to build a phylogenetic

tree with 1000 bootstraps[121,122]. Branch site model tests of the CODEML package[82]

were performed as implemented in EasyCoDEML[123] by selecting the branch of choice.

The tree was unrooted. Protein phylogenetic analysis was performed using the

maximum likelihood method implemented in the IQtree program (RRID:

SCR_017254)[124] under multiple parameter regimes using: 1) the Q.plant substitution

matrix derived from alignments specific to plant nuclear genes and 1 invariant site

category with 8 gamma distributed sites; 2) the LG substitution matrix with 1 invariant

site category with 8 gamma distributed sites or 3) with a 20-profile mixture model.

Bootstrap values were calculated using the Shimodaira-Hasegawa-like approximate

likelihood ratio (SH-aLRT) [125].

1075

1076   F-box gene sequences of 710 Arabidopsis genes used for ortholog analysis were

1077   obtained from Ref. 84 of which 64/69 were common to *skr* upregulated. Classification

1078 of LTS and STS genes was according to Ref. 84. Identification of orthologs was done

1079 using Proteinortho[126].

1080

Acknowledgments

We thank David Twell for kindly providing the LAT52:GFP reporter line, ABRC and

CSHL for seeds. Kathakali Bannerjee and Manoj Kamble for technical support, Mekala

Sai Kiran for maintenance of plants, the NGS and Advanced Microscopy facilities at

CCMB, and Priyanka Pant for help in illustrations. We are grateful to Elliot Meyerowitz,

Utpal Nath, and Manjula Reddy for valuable comments on the manuscript.

1087

Funding

This work was supported by a CSIR Network Project Grant MLP0120 and a

Department of Biotechnology Centre of Excellence Grant

BT/PR21305/COE/34/44/2016 to IS. SP and PS were supported by CSIR Fellowships.

AR was supported by a Department of Biotechnology Research Associate Fellowship.

IS acknowledges a JC Bose fellowship from the Department of Science and

Technology and a Senior Scientist fellowship from the Indian National Science

Academy.

1096

1097 **Author contributions**

1098 PS, SP, JND, JD and IS conceived and planned experiments. SP, PS, AR, JND, VS,

1099 HA, AS, and IS performed experiments. SP, PS, AR, AS, CK, HG, JD, and IS provided

1100 analysis of the experimental results. LA contributed the evolutionary analysis of Figure

1101 6. IS and PS wrote the manuscript with contributions from LA and JD.

1102

1103 **Competing interests**

1104 Authors declare no competing interests.

1105

1106 **Additional information**

1107 Supplementary information is available for this paper.

1108

1109 **Materials and correspondence**

1110 Correspondence and requests for materials to be addressed to IS. Sequencing data

1111 will be deposited at the European Nucleotide Archive.

1112

Supplementary information

Supplementary Table 1: Differentially expressed gene lists.

Supplementary Table 2: Male gametogenesis enrichment analysis of *skr* upregulated

genes.

Supplementary Table 3: Analysis of genes differentially expressed in *skr.*

Supplementary Table 4: Differentially expressed translation-related genes in *skr*

ski2.

Supplementary Table 5: Enrichment for short taxonomic scale (STS) F-box genes

upregulated in *skr*.

Supplementary Table 6: List of primers

Supplementary Figure 1: Multiple sequence alignment of the SHUKR domain.

Supplementary Figure 2: Multiple sequence alignment of SKR protein sequences

used for branch-site analysis.

Supplementary Figure 3: Validation for eight *skr* upregulated F-box genes by RT-

qPCR.

Supplementary File 1: Sequences of *SHUKR* family genes used in branch-site

analysis.

Fig. 1: *SHUKR* is required for pollen development.

Fig. 2: Staging of *skr* phenotype and SKR protein expression

Fig. 3: Ultrastructure and marker expression reveal early gametogenesis defects in *skr*

Fig. 4: *SHUKR* is a gametogenesis inhibitor

Fig. 5: *SHUKR* regulates the proteasome in pollen development

Fig. 6: Sequence and evolutionary analysis of the SHUKR domain

Fig. 7: Diversifying selection in SHUKR regulated F-box genes

Extended Data Fig. 1:
(a) Gene diagram of *SKR* showing location of insertions: *skr-1* (GT_5_101780), *skr-2* (CSHL_GT5061), *skr-3* (SALKSeq_034497.1). (b) Toluidine blue stained tetrads. (c) Inflorescence stem and Alexander staining showing recessive male sterility of *skr.* (d) Complementation of *skr-3* by a *gSKR* transgene. Scale bar: 1 cm for inflorescence, 100 μm for anthers.

Extended Data Fig. 2:

Profiling of SKR expression in male germline development.

TPM values for SKR were collected from the CoNekt database, except for meiocytes. Meiocyte TPM value were taken from (Walker 2018)[21]. UNM – Uninucleate microspore; BCP – Bicellular pollen; TCP – Tricellular pollen; MP – Mature pollen.

Extended Data Fig. 3:

Sections of anther development stages.

Corresponding stages are shown for wild type (A-D, I-L) and *skr* (E-H, M-P). Stages are numbered according to (Sanders 1999)[9]. Abbreviations: E epidermis, En endothecium, ML middle layer, T tapetum, Td tetrad, Msp microspore, DM degenerating microspore, PG pollen grain, St stomium, MD microspore debris. Scale bar 20 µm.

shukr

| Ovule stg 1-2 | tetrad | Ovule stg 2-1 | degenerating microspore | Ovule stg 2-2 | degenerating microspore | Ovule stg 2-3 | degenerating microspore |

Wild type

| Ovule stg 1-2 | tetrad | Ovule stg 2-1 | vacuolated microspore | Ovule stg 2-2 | late vacuolated microspore | Ovule stg 2-3 | post-UNM |

Ovule stage	Pollen development stage	
	Wild type	***shukr***
1-2	Tetrad	Tetrad
2-1	Vacuolated microspores	Degenerating microspores
2-2	Late vacuolated microspores	Degenerating microspores
2-3	Post-mitotic	Degenerating microspores

Extended Data Fig. 4:

Staging of microspore degeneration in *skr*.

Microspore stage and phenotypes are shown together with stage of ovules from the same bud (Schneitz 1995)[10] as a reference for comparison to wild type.

Extended Data Fig. 5:

Precocious expression of F-box gene:nlsGFP fusions in meiosis in *skr*.

(a) At5g02980. A-C, A'-C': wild type; D-F, D'-F': skr; A,A',D,D' meiocytes; B,B',E,E': tetrad; C,C',F,F': uninucleate microspores. (b) At5g62510. A-E, A'-E': wild type; F-H, F'-H': skr; A,A',F,F' meiocytes; B,B',G,G': tetrad; C-E, C'-E',H,H': uninucleate microspores. A-H: GFP fluorescence; A'-H': corresponding DIC. GFP expression in *skr* is first detected in meiocytes (At5g02980) or tetrad stage At5g62510) whereas in wild type earliest GFP signal was in released microspores. Scale bar: 5 μm

a

Cross	Tested Generation	Plants tested	Sterile	Fertile	Sterile : Fertile
skr X 08-10 suppressor line	Backcrossed F1	20	20	0	1:0
skr X 08-10 suppressor line	Backcrossed F2	180	129	51	2.53:1

b

c

skr-3   *skr-3 ski2-6*   *skr-3*

Extended Data Fig. 6:

ski2 is a suppressor of *skr*.

(a) Mendelian segregation of the suppressor mutation in the 08-10 *skr* suppressor line (b) A C>T transition in the *ski2* allele (named *ski2-7*) present in the 08-10 *skr* suppressor line results in a TGA termination codon at position 518. (c) Restoration of fertility and pollen viability in *skr-3* by the independent allele *ski2-6*. Scale bar: inflorescence 1 cm; anthers 100 µm.

Extended Data Fig. 7:

Phylogenetic tree of the SHUKR family from eudicots.

119 representatives from 85 species were used for tree construction using only the SHUKR domain that is shared by all representatives of the family. Genbank ID and species are indicated. Brassicaceae+Cleomaceae (dark green) and Asteraceae (light green)

Scale 0.1 ⊢⊣

ThSKR *Tarenaya hassleriana*
EsSKR *Eutrema salsugineum*
SpSKR *Schrenkiella parvula*
SiSKR *Sisymbrium irio*
RsSKR *Raphanus sativus*
BnSKR *Brassica napus*
BrSKR *Brassica rapa*
BvSKR *Barbarea vulgaris*
ChSKR *Cardamine hirsuta*
AlSKR *Arabidopsis lyrata*
AtSKR *Arabidopsis thaliana*
CsSKR *Camelina sativa*
CgSKR *Capsella grandiflora*
CrSKR *Capsella rubella*
ThSKL *Tarenaya hassleriana*
SiSKL *Sisymbrium irio*
SpSKL *Schrenkiella parvula*
BnSKL *Brassica napus*
RsSKL *Raphanus sativus*
BrSKL *Brassica rapa*
EsSKL *Eutrema salsugineum*
BvSKL *Barbarea vulgaris*
ChSKL *Cardamine hirsuta*
AlSKL *Arabidopsis lyrata*
AtSKL *Arabidopsis thaliana*
CsSKL *Camelina sativa*
CrSKL *Caspella rubella*
CgSKL *Capsella grandiflora*
CpSKL *Carica papaya*

Extended Data Fig. 8:

Phylogenetic tree of Brassicales *SHUKR* and *SHUKR-like* (*SKL*) genes used for branch-site test.

The *SHUKR* gene family is represented by a single gene in *Carica papaya*, whereas Brassicaceae and Cleomaceae experienced a gene duplication event prior to their divergence. CDS were aligned using MAFFT, and RAxML was used to build the phylogenetic tree with 1000 bootstraps. The red branch was selected as foreground for *SKR*. Branch-site test for positive selection was performed using the CODEML package implemented in EASYCoDEML. Likelihood Ratio Test (LRT) results of Model A vs Model A null (Extended Data Table 1), gave a p-value of 0.0031 for *SKR*. The tree was unrooted.

Model	LnL	Estimates of parameters					Model compared	LRT P-value	Positive sites
Model A	-4394.302739	Site class	0	1	2a	2b	Model A vs.Model A null	0.003120857	7 E 0.932,22 F 0.672,25 V 0.529,31 E 0.613,43 F 0.553,57 E 0.962*,63 L 0.974*,82 K 0.594,100 P 0.747,112 R 0.668,114 C 0.643
		f	0.66753	0.21979	0.08477	0.02791			
		ω0	0.20115	1.00000	0.20115	1.00000			
		ω1	0.20115	1.00000	22.04175	22.04175			
Model A null	-4398.670454	1							Not Allowed

Extended Data Table 1: Branch-site test results for positive selection in the SKR branch in the Brassicaceae. Branch site models postulate three classes of sites: those under purifying selection (0< ω<1), neutral (ω=1), and those under positive selection (ω>1) in the foreground branch being tested (Yang, 2007)[82]. Model A is compared to Model A null in which the third class is set to ω=1. Positive sites are indicated together with posterior probabilities. P-value for likelihood ratio test (LRT) results are calculated from the χ^2 distribution taking as critical value χ^2= 2(LnL$_{MA}$-LnL$_{MAnull}$) and dividing by 2 (Yang, 2007)[82].