Actin polymerization by WASp limits nuclear envelope rupture and inflammation in macrophages

Roberto Amadio[1,2], Giulia Bracchetti[1], Zahraa Alraies[3,#], Giulia Maria Piperno[1], Lucia Lopez Rodriguez[1], Mathieu Maurin[3], Martina Conti[4], Laura Andolfi[4], Maria Carmina Castiello[5,6], Francesca Ferrua[5], Anna Villa[5,6], Alessandro Aiuti[5], Ana-Maria Lennon-Dumenil[3] and Federica Benvenuti[1,*]

[1]Cellular Immunology, International Centre for Genetic Engineering and Biotechnology (ICGEB), Trieste, Italy

[2]University of Trieste, Department of Life Sciences (DSV), 34127 Trieste, Italy

[3]INSERM-U932 Immunité et Cancer, Institut Curie, PLS Research University, 75005 Paris, France

[4]Istituto Officina dei Materiali - National Research Council (CNR-IOM) - Basovizza, Trieste, Italy

[5]San Raffaele Telethon Institute for Gene Therapy (SR-TIGET), IRCSS San Raffaele Scientific Institute, 20132 Milan, Italy

[6]Institute for Biomedical Technologies - National Research Council (CNR-ITB), 20054 Segrate, Milan, Italy

[#]Present address: Laboratory of Cell and Molecular Biology, Experimental Research (C1), 005E - NYU Abu Dhabi, Saadiyat Campus, Abu Dhabi, United Arab Emirates

*Corresponding author, Federica Benvenuti, benvenut@icgeb.org

Abstract

Mutations in the immune-specific actin regulator WASp induce a proinflammatory state in myeloid cells, whose underlying causes remain poorly defined. Here, we applied microfabricated tools that mimic tissue mechanical forces to explore the role of WASp in connecting mechano-sensing to the activation of inflammatory responses in macrophages. We show that WASp-deficient macrophages carry nuclear structure alterations and undergo increased blebbing and nuclear rupture when exposed to mechanical confinement. High-resolution imaging indicates that WASp drives the formation of protective perinuclear actin structures in response to confinement. Functionally, WASp null macrophages respond to mechanical confinement by inducing a transcriptional profile consistent with the release of immunogenic DNA in the cytosol. The proinflammatory state of mechanically confined WASp-deficient macrophages depends, in part, on the cGAS-STING pathway of cytosolic DNA sensing. Together, these data uncover a WASp-dependent mechanism to restrict activation of inflammatory signalling in tissue macrophages and provide hints to target unabated inflammation in Wiskott-Aldrich Syndrome.

Running title: Branched actin polymerization in control of nuclear integrity and inflammation

Keywords: Wiskott-Aldrich Syndrome, actin, nuclear mechanics, nuclear rupture, inflammation

ORCIDs and e-mails:

R.A: 0000-0003-0861-0703, roberto.amadio@icgeb.org

Introduction

Misplaced cytosolic DNA is emerging as a critical cause of inflammation in various monogenic and systemic autoimmune diseases (Decout et al., 2021). Mitochondrial DNA is a well-established immunogenic trigger, engaging TLR9, cGAS or NLRP3, depending on the context (Rodríguez-Nuevo et al., 2018; Maekawa et al., 2019; Xian et al., 2022). In addition, nuclear alterations and genomic DNA damage leads to chromosomal DNA extrusion into the cytosol (Härtlova et al., 2015; Dou et al., 2017; Dunphy et al., 2018). DNA molecules that access the cytosol are detected by sensors such as cGAS and STING and trigger activation of cytokine genes, senescence or cell death, depending on the severity of damage (Dou et al., 2017; Mu et al., 2020; Sladitschek-Martens et al., 2022). Hence, maintaining nuclear integrity is critical to restrict unwanted activation, a process that is particularly relevant in cells that navigate complex environments (Denais et al., 2016), or are continuously exposed to mechanical compression (De Silva et al., 2023). Nuclear stability relies on chromatin mechanics (Nava et al., 2020) and cytoskeletal forces. Within cytoskeletal components, microtubules limit nuclear deformations in dendritic cells (Thiam et al., 2016), while intermediate filaments protect nuclei of murine embryonic fibroblast during constricted migration (Patteson et al., 2019). Arp2/3-dependent branched actin polymerization deforms the nuclei of migrating dendritic cells (Thiam et al., 2016) and maintains nuclear integrity in stromal cells (Sladitschek-Martens et al., 2022). Recently, genetic ablation of the Cdc42 GTPase DOCK8 showed that branched actin polymerization preserves nuclear integrity in T and dendritic cells during confined migration (Reis-rodrigues et al., 2024; Shen et al., 2024), suggesting broader nuclear defects in actinopathies.

The Wiskott-Aldrich Syndrome (WAS) is a primary immunodeficiency characterized by frequent autoimmune and autoinflammatory manifestations. WAS is caused by mutations in Wiskott-Aldrich Syndrome protein (WASp), a critical activator of the Arp2/3 complex which controls branched actin polymerization specifically in non-erythroid immune cells (Blundell et al., 2010; Naseem et al., 2022). Altered platelets generation, autoreactive B cells and impaired Tregs function predispose to autoimmunity in Wiskott-Aldrich Syndrome (WAS) (Sereni et al., 2018; Calixto Vieira et al., 2023). Our group identified a link between the release of immunogenic DNA by neutrophil extracellular traps and excessive production of type-I interferon by WASp KO (WKO) plasmacytoid dendritic cells (pDCs) (Prete et al., 2013; Cervantes-Luevano et al., 2018). In addition, WKO macrophages activates the inflammasome due to impaired bacterial clearance (Lee et al., 2017), and defective maturation of the endolysosomal network upon engulfment of circulating immuno-complexes triggers TLR9 and cGAS pathways in DCs (Piperno et al., 2020; Amadio et al., 2021). WASp has been observed to

polymerize actin patches in dendritic cells migrating inside confined environments, which push perpendicularly to the direction of movement and maintain vertical space for the cell to crawl (Gaertner et al., 2022). Despite these observations, a direct contribution of WASp to nuclear stability and inflammation in myeloid cells remains elusive.

In this work, we studied the role of WASp in preserving nuclear stability under mechanical pressure. By combining microfabricated tools that mimic tissues' mechanical forces, imaging and transcriptional profiling, we found that WASp-driven actin polymerization contributes to nuclear envelope integrity and counteracts mechanical challenge in macrophages. Consequently, lack of WASp causes inflammatory activation by extruded genomic DNA, which is in part mediated by STING.

Results

Inflammatory phenotype in mouse and human WKO macrophages

Macrophages are the most abundant myeloid cells subset in tissues and critically contribute to inflammation in pathological conditions (Ginhoux and Guilliams, 2016). WASp null macrophages were recently shown to express an inflammatory signature at steady state (Liang et al., 2023). To investigate whether mechanical compression further induces inflammation in WKO macrophages, we focused on lung alveolar macrophages (AMO), a representative population of long-lasting resident macrophages constantly exposed to physical constrains and subjected to almost 2D confinements (De Silva et al., 2023). Flow cytometry profiling of AMO (Supplementary Fig.1A), showed higher levels of the activation marker CD86 in WKO mice (10 weeks of age), as compared to WT (Fig.1A). We next isolated AMO from 10 weeks and 25 weeks old mice. The absolute number of AMO was higher in WKO as compared to controls, especially at 25 weeks of age (Fig.1B). Notably, two key inflammatory genes were significantly induced in AMO from 25 weeks old WKO mice (Fig.1C). Similarly, bone marrow-derived macrophages expressed higher activation of inflammatory genes (Fig.1D). We conclude that, consistently with previous findings in the spleen (Prete et al., 2013; Liang et al., 2023), WASp deficient macrophages are constitutively activated and that this phenotype is exacerbated by tissue residency *in-vivo*.

To translate the importance of macrophage activation to humans, we designed an inflammatory signature (Supplementary Table 1) and used it to interrogate two different data sets. First, we re-analysed a publicly available dataset of WKO iPSCs-derived macrophages (GSE107963, TPM normalized reads), and their isogenic counterparts that have been corrected to re-express WASp (Yuan et al., 2022). The inflammatory score, as well as relevant inflammatory genes, were consistently reduced by restoring WASp expression in iPSCs (Fig.1E and F). We then performed a bulk RNA-seq on a cohort of PBMCs from 6 WAS patients, before and after WASp reconstitution by gene therapy. Major WAS defects (i.e. BCR signalling and genes related to platelets function) were normalized upon re-expression of WASp, without inducing alteration in other Arp2/3 activators (Supplementary Fig.1B and C). Interestingly, despite the complex composition of PBMCs that comprises several subsets (mainly B, T, NK and monocytes) and patient heterogeneity, we found higher inflammation in WAS samples, which was diminished by WASp restoration (Fig.1G and H). Selected inflammatory genes emerging from RNA-seq, were validated by RT-qPCR, confirming that

re-introducing WASp is sufficient to normalize expression to the level of healthy donors (Supplementary Fig.1D). Intriguingly, the overall correction index was greater in samples showing a higher basal level of inflammation (Supplementary Fig.1E). Altogether, our data indicate that WASp restrain expression of inflammatory genes in mouse macrophages and human cells.

Nuclear defects in WKO macrophages

Having established that WASp deficiency correlates with elevated expression of inflammatory genes, we next sought to explore the underlying molecular basis. For this purpose, we moved to bone marrow-derived macrophages (BMDM) that recapitulate inflammatory features and are easy to obtain in large numbers. We initially assessed simple nuclear parameters in 2D, such as area, roundness and micronuclear occurrence (Fig.2A). We found that the nuclei of WKO macrophages had an increased mean projected area, were more irregular and showed a higher incidence of micronuclei (Fig.2B), indicating general alterations in nuclear homeostasis. Previous findings in other systems linked defects in Lmna (encoding for Lamin A/C) and nuclear envelope folding to impaired mechanical tension and reduced protection from deformations (Kim et al., 2017; Tang et al., 2023). We thus focused on Lamin A/C and the nuclear envelope protein Emerin in our model. Total Lamin A/C levels were reduced in WKO, both at the mRNA and protein level, whereas mRNA expression of type B lamins and Emerin were not different (Fig.2C and D). Importantly, we noticed higher degree of Lamin A/C wrinkles and irregularities in nuclei of WASp null macrophages (Fig.2E), which coincided with Emerin accumulation (Fig.2F). Altogether these data suggest defective nuclear envelope homeostasis and increased folding of the nuclear envelope in the absence of WASp.

Next, as the nucleus is highly connected to chromatin and cellular mechanics, we explored chromatin parameters and global cellular stiffness. We did not observe substantial differences in global heterochromatin content and peripheral heterochromatin between the two genotypes by chromocenters analysis (Supplemental Fig.2A). Similarly, no changes in heterochromatin distribution were detected by H3K9me2 staining, but an increase in the total intensity of this marker (Supplemental Fig.2B). Cellular mechanics, as measured by Atomic Force Microscopy (AFM) indentation (Supplementary Fig.2C) showed that WKO cells were stiffer than WT ones (Supplementary Fig.2D). Of note, a stiffer cytoplasm has been linked to reduced nuclear stability under 3D physical constrains (Bastianello et al., 2023). Together, these initial data indicate a global defect in nuclear-cytoskeletal mechanics in macrophages derived from WASp null animals, which may affect nuclear stability under 3D tissues forces.

Increased constricted migration and nuclear deformation in WKO macrophages

To test the hypothesis of reduced nuclear stability in WKO, we first used an assay to measure nuclear deformations in macrophages inside narrow constrictions by means of microchannels (Vargas et al., 2016). We adopted different channel dimensions, ranging from 2 μm to 8 μm in width and having a constant height of 5 μm. Macrophages were seeded in the pre-channel chamber and recorded overnight. The number of cells that entered the channel was calculated at the end of the recording period by counting the number of cells inside channels and normalizing by the number of approaching cells (Supplementary Fig.3A; materials and methods). As expected, reducing the size of the channel, limited the number of cells entering inside constrictions (Fig.3A). Interestingly, the entrance rate of WKO macrophages was consistently higher, especially at smaller constrictions (Fig.3B). By carefully inspecting nuclear shape in cells inside channels we observed an increased rate of nuclear herniations in WKO (Fig.3C). The ratio between front and back herniations was not different between genotypes and skewed to the front (Fig.3D), as already expected based on observations in other cell types (Denais et al., 2016). Since herniations predispose to nuclear rupture, we next transduced BMDM with a lentiviral construct expressing nuclear localization signal (NLS)-tagged GFP (NLS–GFP), which is retained inside nuclei and released only upon nuclear envelope break (Fig.3E). Interestingly, we observed a larger fraction of WKO macrophages undergoing nuclear envelope (NE) rupture and cytosolic release of fluorescent signal than WT (Fig.3F and G). Moreover, and consistent with increased nuclear fragility, we detected multiple rupture events on the same cell exclusively in WKO (Fig.3H).

A parallel set of experiments confirmed the superior ability of WASp null cells to translocate through small pores in transwell assays (Supplementary Fig.3B). Notably, migration induced an increase in nuclear deformation as well as γH2AX staining, particularly in WKO cells (Supplementary Fig.3C). Expression of the relevant chemokine receptor targeted to promote transmigration was equal in the two genotypes, ruling out that defective sensing of the chemotactic gradient affected the observed migration rate (Supplementary Fig.3D). Overall, these data indicate that WASp-deficient macrophages have an enhanced ability to navigate through small constriction, in line with a previous report on dendritic cells (Gaertner et al., 2022), suggesting reduced resistance to nuclear deformation. Importantly, the nuclear envelope of WKO cells ruptures during constricted migration, releasing nuclear material into the cytosol.

WASp limits nuclear ruptures under mechanical confinement

To more faithfully recapitulate deformation events experienced by macrophages in tissues, we next applied a vertical confiner device (Alraies et al., 2024) based on PDMS micropillars (Supplementary Fig.4A). The heights of BMDM nuclei plated on fibronectin-coated glass slides without confinement range between 5 and 6 µm (Supplementary Fig.4B), therefore we selected a height of 6 µm and 3 µm as mild and strong confinement respectively (Supplementary Fig.4C). Confocal images were acquired after one hour of confinement to examine nuclear parameters. We classified nuclei into two categories: 1) intact, when the nuclear boundaries were regular and continuous; 2) damaged, when blebs or herniations were found to interrupt the nuclear perimeter. (Fig.4A). The fraction of damaged nuclei in WASp null cells was slightly higher under mild confinement and significantly higher under strong confinement (Fig.4B), suggesting loss of nuclear integrity. Indeed, measuring NLS-GFP localization after confinement at 3 µm, we observed an increased frequency of NE rupture and NLS-GFP cytosolic release in WKO cells (Fig.4C and D). Live tracking of NLS-GFP leakage revealed that this event was often preceded by blebbing/herniation of the nuclear surface (Fig.4E). To causally link WASp to nuclear deformation and loss of NE integrity, we next transduced WASp null macrophages with a lentiviral construct expressing WASp-GFP (Supplementary Fig.4D) (Blundell et al., 2008). This resulted in around 50% of cells expressing a clear GFP signal and WASp-GFP fusion protein of the expected size (Supplementary Fig. 4E and F). The resulting mixed population of GFP-positive (rescued) and negative (control) cells was subjected to confinement at 3 µm. In this highly controlled setting, we found that GFP-positive cells showed significantly fewer nuclear blebs than GFP-negative ones (Fig. 4F and G). Moreover, the fraction of cells showing Lamin A/C rupture was reduced in WASp-GFP positive cells (Fig.4H and I), confirming a direct impact of WASp in preserving nuclear stability under confinement.

WASp directs perinuclear actin polymerization in confined macrophages

We then aimed to investigate a potential mechanism by which WASp and actin polymerization support nuclear stability during mechanical challenge. We took advantage of a fixation-under-confinement protocol, which maintains intact cytoskeletal structures once confinement is released (Alraies et al., 2024). Accumulation of dense actin structures in the central body of the cell were induced by confinement in WT but not in WKO macrophages (Fig.5A). These actin patches closely resemble the Arp2/3-dependent structures described in mechanically confined dendritic cells with a role in nuclear envelope unfolding (Alraies et al., 2024). By high-resolution imaging, we further

classified confined cells into four categories according to actin distribution: 1) perinuclear actin patches; 2) intranuclear actin accumulation; 3) filamentous perinuclear actin ring and 4) diffuse cytosolic actin (Fig.5B). Among central body actin structures, perinuclear actin patches were observed in a substantial proportion of WT cells but were virtually absent in WKO cells (Fig.5C and D). Intranuclear actin accumulation was present in a small fraction of cells, in similar frequencies between the two genotypes. Additionally, a small fraction of WKO cells showed thin filamentous actin surrounding the nucleus (Fig.5C and D), a phenomenon not detected in WT. To explore the role of WASp in driving the polymerization of perinuclear actin structures, we leveraged WKO macrophages transduced with WASp-GFP for colocalization analysis (Fig.5E). Of note, the Mander's coefficient between phalloidin and GFP was high in the perinuclear actin patches induced by confinement (Fig.5F), strongly suggesting an active role of WASp in directing the formation of these structures. Consistently, also the distribution of WASp-GFP in macrophages inside microchannels was enriched in the central part of the cell body and around the nucleus (Supplementary Fig.5A and B). We conclude that WASp drives the nucleation of perinuclear actin patches in macrophages exposed to mechanical confinement. Together with the previous observations, this result suggests that a local perinuclear actin response may help to preserve nuclear integrity.

Confinement-induced nuclear ruptures exacerbate inflammation in WKO macrophages

To investigate the functional consequences of increased nuclear fragility in WKO, we analysed the transcriptional profiles of macrophages upon confinement (Fig.6A). WT and WKO macrophages at steady state (unconfined) express a common transcriptional program and some genotype-specific genes (Supplementary Fig.6A). Remarkably, WKO cells were enriched in immunoglobulin coding genes, an observation that deserves future investigations. Moreover, transcripts related to extracellular matrix organization, vascular remodelling and cellular activation/migration were higher in unconfined WKO macrophages than in WT. In contrast, genes controlling cellular metabolism were less expressed in WKO (Supplementary Fig.6A and Supplementary Table 2). Confinement induced profound transcriptional remodelling in both genotypes (Supplementary Fig.6B). Analysis of differentially expressed genes (DEGs) between unconfined and confined cells unveiled a core of upregulated genes shared by WT and WKO (Fig.6B), including some inflammatory cytokines and chemokines (Fig.6C). Interestingly, we observed several genes controlling cell activation and tissue remodelling selectively upregulated in confined WKO cells (Fig.6B), paralleled by reduced expression of metabolic pathways (Supplementary Fig.6C), indicating increased activation in WASp KO samples.

To explore the molecular mechanisms underlying cellular responses to confinement, we evaluated the expression of genes implicated in cytoskeletal dynamics and control of nuclear stability (Supplementary Table 4) across all the conditions. A signature including genes that control actin branching (branched actin score) was significantly higher in WT than in WKO macrophages, suggesting that WASp deficiency broadly impacts the process. Confinement induced transcriptional activation of this set of genes in both genotypes, indicating a fast cellular adaptation to mechanical inputs (Fig.6D). However, expression in WKO remained consistently lower despite a mild induction by confinement. This trend was even more prominent when selecting specific WASp effectors that have been directly linked to NE integrity (*Actr2*, *Actr3*, *Arpc4)* (Sladitschek-Martens et al., 2022; Delgado, 2024); showing a drastic reduction of expression in both unconfined and confined WASp deficient cells. Of note, a second gene module linked to nuclear mechanics including several genes previously implicated in NE stability, such as Lamin A/C, Emerin and Lap2β (Tmpo) (Chen et al., 2021; Lavenus et al., 2022; De Silva et al., 2023) was significantly upregulated in WT cells, but not in WKO, upon confinement (Fig.6E). Conversely, SUN2 and RhoA/ROCK, whose overexpression has been linked to nuclear blebbing and innate activation (Mu et al., 2020), were much more induced in WKO, together with RAC2-related genes (Supplementary Fig.6D). All together, these data suggest a defective branched actin-based response to confinement in WKO macrophages, concomitant to induction of a compensatory RhoA/RAC-mediated pathway, with a potentially negative impact on nuclear stability.

Defective mechanotransduction and reduction in the expression of genes linked to actin-dependent nuclear integrity were previously shown to induce activation of a proinflammatory cell state in Arp2/3 deficient fibroblast (Sladitschek-Martens et al., 2022). Hence, we examined the expression of this signature, defined as "nuclear envelope rupture (NER) inflammatory score" in our dataset. Remarkably, the NER inflammatory score was selectively activated in confined WASp null macrophages, causally connecting WASp to the control of proinflammatory activation in cells exposed to mechanical inputs (Fig.6F). Among the genes included in NER inflammatory score, Il6 was selected to perform further functional analyses. qPCR confirmed that Il6 transcripts are preferentially induced by confinement in WKO (Fig.6G). To exclude any potential bias in macrophages derived from constitutive WASp KO animals, we induced acute WASp deletion in WT BMDM by CRISPR-Cas9 gene editing (Supplementary Fig.6E). Importantly, WASp-depleted macrophages responded to confinement by augmented Il6 induction as compared to control cells (Fig.6H),

indicating that WASp directly restricts confinement-induced activation. Given the widespread implications of the cGAS-STING pathway of cytosolic DNA sensing (Ablasser and Gulen, 2016) to innate activation and specifically in driving the NER inflammatory score (Sladitschek-Martens et al., 2022), we assessed its contribution to confinement-induced activation of WKO cells. Efficient STING ablation in WKO macrophages (Supplementary Fig.6F) blunted Il6 production upon direct stimulation by DMXAA, confirming the functional disruption of the pathway (Supplementary Fig.6D). We then exposed WKO macrophages carrying or not STING depletion to confinement at 3 μm of height. Induction of Il6 transcripts was significantly reduced in WKO STING KO macrophages as compared to STING-proficient cells, causally connecting the pathway to the proinflammatory state of WKO macrophages (Fig.6I). The partial ablation of Il6 responses in STING-depleted cells, suggests that other sensors/mechanisms contribute to the inflammatory activation downstream of NE ruptures in WASp-deficient cells.

Collectively, these data uncover a differential transcriptional response by WASp null macrophages to mechanical confinement, showing that WASp limits NE ruptures and the firing of cytokine signalling in myeloid cells exposed to physiologically relevant constrains (Fig.6J).

Discussion

In this study we identified an unprecedented role for WASp in preserving nuclear integrity, thus restraining inflammatory responses in macrophages exposed to tissue forces.

The direct connection between WASp-mediated actin polymerization and nuclear integrity has emerged by studying macrophages in confined environments that recapitulate tissue forces experienced *in-vivo*. Hence, the approach used in this study uncovers an essential supplementary function for the hematopoietic-specific actin nucleation-promoting factors WASp, beyond the well-established role at the cell cortex and in intracellular trafficking (Linder et al., 1999; Piperno et al., 2020; Gaertner et al., 2022). These findings are particularly relevant given the intrinsically migratory nature of immune cells and the nucleus as the main rate-limiting factor for constricted migration.

Previous studies showed that WASp null dendritic cells migrate faster inside microchannels or through constriction. However, the findings were not interpreted as dependent on altered nuclear properties (Gaertner et al., 2022; Oliveira et al., 2022). Our analysis confirms that WASp null macrophages, likewise dendritic cells, are facilitated in entering small constrictions. Additionally, we show that this phenomenon is linked to the enhanced ability of the nucleus to deform and a predisposition to undergo nuclear rupture. Cells harbouring defects in Lamin A/C or a disrupted nuclear-cytoskeleton coupling show similar nuclear deformations, increased migration and high rates of nuclear ruptures when exposed to mechanical confinement (Davidson et al., 2014; Harada et al., 2014; Kim et al., 2017; Bell et al., 2022). In line with our findings, a parallel study by our collaborators observed increased nuclear deformation and ruptures in confined dendritic cells that lack the WASp effector Arp2/3 (Delgado, 2024). More than one mechanism, not mutually exclusive, may explain reduced nuclear integrity in WASp null cells. The dense F-actin patches formed upon confinement and the immediate transcriptional response to supply factors controlling actin branching in WT, but not in WKO cells, suggest that WASp induces cytoskeletal remodelling downstream of mechanical stimuli, creating a protective perinuclear actin barrier. Alternatively, as indicated by the thin actin rings surrounding the nucleus in a fraction of WKO cells and by gene expression data, lack of WASp may drive compensatory actin polymerization by different factors. As documented in a previous report, the resulting abnormal actin network may cause rupture by exerting excessive pressure on the NE under confinement (Mu et al., 2020). Interestingly, similar perinuclear actin rings were also observed in epithelial cells upon mechanical stretching, which has a role in promoting chromatin compaction (Le et al., 2016). Moreover, we cannot exclude that WASp

limits nuclear rupture by acting at the DNA level, protecting it from damage (Kovacs et al., 2023), as it occurs in the case of Arpc4 (Delgado et al. 2024). Of note, despite mild Lamin A/C defects and slight accumulation of DNA damage in WASp null macrophages, the number of tissue-resident cells *in-vivo* is not affected, as opposed to the phenotype in Arpc4 and LMNA/C deficient animals where Langerhans cells and alveolar macrophages, respectively, are depleted in tissues (De Silva et al., 2023; Delgado, 2024). Therefore, WASp ablation induces a milder phenotype than complete blockade of Arp2/3 or Lamin A/C deletion, and cells can cope with nuclear damage by inducing inflammation but without undergoing death.

The unscheduled release of genomic DNA into the cytosol has been documented consequently to DNA damage (Dunphy et al., 2018; Mukherjee et al., 2019; Song et al., 2021), defective Lamin A/C (De Silva et al., 2023; En et al., 2024) and disrupted mechano-signalling due to reduced expression of Arp2/3 regulators, a direct WASp effector (Sladitschek-Martens et al., 2022). The latter study associated nuclear instability and rupture to cGAS-STING activation and senescence induction in fibroblast. We observed a similar inflammatory activation downstream of NE rupture in WKO, raising many interesting points. While STING predominantly activates IRF3 upon exogenous introduction of DNA in the cytosol by transfection, leakage of genomic DNA following DNA damage leads to unconventional STING signalling, predominantly engaging NF-kB rather than IRF3 (Dunphy et al., 2018). Our NER inflammatory activation skewed towards cytokines rather than interferon supports the latter mechanism of STING engagement. Importantly, our work does not exclude the potential involvement of other DNA sensors nor the role of additional mechanisms connecting nuclear deformation and rupture to proinflammatory signalling in WASp null cells. Remarkably, the NER inflammatory signature induced in confined WASp null macrophages has been linked to the induction of senescence in fibroblasts (Sladitschek-Martens et al., 2022). Even if we have not explored senescence in WASp null cells or tissues, the expression of this signature under confinement, together with Lamin A/C defects and nuclear rupture, lead us to speculate that lack of WASp may predispose to accelerated ageing of the myeloid compartment *in-vivo*. Beyond the specific role of WASp described here, our findings suggest that general actin defects may be linked to inflammatory conditions due to nuclear alterations, as proposed by others (Sladitschek-Martens et al., 2022), underscoring the relevance of investigating mechanical control of inflammation in immune cells. From a clinical perspective, our study has implications for understanding the molecular basis of WAS autoimmunity. Transcriptional data in human iPSCs-derived macrophages and total cells from the peripheral blood of gene therapy-treated WAS patients consistently indicate

that WASp restricts the accrual of inflammation. However, inflammatory relapses are still frequent after treatment in WAS, including hematopoietic stem cell transplantation (Sudhakar et al., 2021). According to recent evidence, including the data presented here, gene therapy offers better control of inflammation in treated patients (Sereni et al., 2019; Magnani et al., 2022). Nevertheless, defining the precise origin and cellular basis of WAS inflammation is important to designing the management of residual inflammation, especially for those patients who do not have access to gene therapy.

In summary, our results unveil a cell-intrinsic WASP-mediated mechanism to control nuclear stability and inflammation under physiologically relevant settings, expanding the role of WASp-driven Arp2/3 actin polymerization in preserving myeloid cell homeostasis in tissues.

Materials and methods

Reagents and tools table

Reagent/Resource	Reference or Source	Identifier
Experimental Models		
WASp[-/-] mice - B6.129S6-Wastm1Sbs/J	Jackson #:019458	IMSR_JAX:019458
HC and WAS PBMCs	San Raffaele Institute	Supplementary Table 5
Recombinant DNA		
pCDH-NLS-GFP	Addgene	#132772
pMD2.G	Addgene	#12259
psPAX2	Addgene	#12260
SEWW-eGFP-WASp	(Blundell et al., 2008)	/
Antibodies Flow cytometry		
CD45 - APC-Cy7	BioLegend	103114
CD11c - Bv786	BioLegend	117335
F4/80 - Bv421	eBioscience	25-5870-80
MHC-II - AF700R	BioLegend	151507
LY6C - PE	eBioscience	11-0541-81
CD64 - BV605	BioLegend	108923
SiglecF - PerCP-Cy5.5	BioLegend	107105
CD80 - APC	BioLegend	139323
CD86 - Bv650	BioLegend	104713
CD3 - FITC	BioLegend	100305
LY6G - FITC	Bio-Rad	MCA6077GA
B220 - FITC	BioLegend	103208
CD19 - FITC	BioLegend	115503
NK1.1 - FITC	BioLegend	116410
Antibodies FACS		
CD11c - Bv786	BioLegend	117308
SiglecF - PerCP-Cy5.5	BioLegend	121406
Antibodies immunofluorescence		
Anti-Emerin	Thermo Fischer	PA5-29731
Anti-H3K9me2	Abcam	ab1220
Anti-LMNA/C	Cell Signalling	mAb#4777 (4C11)

Anti-yH2AX	Sigma Aldrich	05-636 (JBW301)
Anti-Mouse-488, Donkey	Invitrogen	A21202
Anti-Mouse-555, Goat	Invitrogen	A21424
Anti-Mouse-647, Chicken	Invitrogen	A21463
Anti-Rabbit-488, Goat	Invitrogen	A21441
Anti-Rabbit-594, Donkey	Invitrogen	A21207
Antibodies western blot		
Anti-GAPDH-HRP	Sigma Aldrich	G9295 (GAPDH-71.1)
Anti-WASp	Santa Cruz Biotech	sc-13139 (B-9)
Anti-LMNA/C	Cell Signalling	mAb#4777 (4C11)
Anti-STING	Cell Signalling	13647 (D2P2F)
Anti-TUBα-HRP	Cell Signalling	#9099 (11H10)
Anti-mouse-HRP	Jackson	115-035-003
Anti-Rabbit-HRP	Jackson	711-035-152
Chemicals, Enzymes and other reagents		
2-mercaptoethanol	Gibco	31350010
ACK lysis buffer	Gibco	A1049201
Alexa Fluor™ 555 Phalloidin	Thermo Fischer	A34055
Alt-R® Cas9 Electroporation Enhancer	IDT	1075916
Alt-R® CRISPR-Cas9 tracrRNA	IDT	1072533
Alt-R® Negative Control crRNA #2	IDT	1072545
Alt-R™ crRNA STING	IDT	Mm.Cas9.TMEM173.1.AA
Alt-R™ crRNA WASp	IDT	Mm.Cas9.WAS.1.AC
Alt-R™ spCas9 Nuclease V3	IDT	1081058
Bovine Serum Albumine (BSA)	Sigma Aldrich	A8806
Clarity ECL substrate	Bio-Rad	170-5060
Collagenase type 2	Worthington	LS004176
Counting Bright beads	Miltenyi	C36950
DMXAA	Invivogen	tlrl-dmx
DNAse I from bovine pancreas	Millipore-Sigma	11284932001
EDTA	Thermo Fischer	15575020
FBS - low endotoxin	Euroclone	ECS0186D - lot. EUS037123
Fibronectin Bovine protein	Thermo Fischer	33010-018
Gentamicin	Gibco	15710064
Glycine	Sigma Aldrich	G7126
Hepes	Invitrogen	15630080

Hoechst 33342	Invitrogen	H3570
IMDM medium	Thermo Fischer	31980030
Lenti-X concentrator	Takara	631231
Live and Dead staining –-Bv510	Invitrogen	L34966
LPS	Invivogen	tlrl-pb5lps
Mouse CD11c microbeads	Miltenyi	130-125-835
Mowiol 40-88	Sigma Aldrich	324590
PDMS; SYLGARD 184	Dow	1673921
PFA	Thermo Fischer	28908
Poly-Ethylene-Glycol (PEG)	Sigma Aldrich	81260-1KG
PrimeScript™ RT Reagent Kit	Takara	RR036A
Protein inhibitor cocktail	Sigma Aldrich	P8340
Recombinant mouse CCL19 (MIP3-β)	Gibco - Peprotech	250-27B-1MG
Recombinant mouse MSC-F	Miltenyi	130-101-706
RIPA buffer	/	/
RNA Later™	Invitrogen	AM7021
RPMI 1640 medium	Thermo Fischer	72400047
Sodium Azide	Sigma Aldrich	13412
Sodium pyruvate	Gibco	11360070
Supernatant-containing GM-CSF	Produced by J558 cells	/
TB Green® Premix kit	Takara	RR420A
Triton-X-100	Sigma Aldrich	X100-500ML
TRIzol™ reagent	Thermo Fischer	15596026
Tween-20	Sigma Aldrich	02286
Kits		
BCA assay kit	Thermo Fischer	23225
RNA high-sensitivity kit	Agilent	5067-5579
RNeasy micro kit	Qiagen	74004
Universal Index Twist kit 96	Diatech-Labline	101307
WatchMaker - mRNA Library Prep Kit	Diatech-Labline	7BK0001-024
Software		
Fiji (ImageJ)	v.1.54f	/
FlowJo	Tree Star, Inc. v.10.8.2	/
Graphpad	Prism v.8.0.1	/
R software	R version 4.3.1	/
Other		

| J-558 cell line (GM-CSF producers) | / | / |

Methods and protocols

Mice

WASp$^{-/-}$ mice (IMSR_JAX:019458) on a C57BL/6 genetic background were a gift from S. Snapper (Massachusetts General Hospital, Boston, Massachusetts, USA) (Snapper et al., 1998). Animal care and experimentation were conducted under national and international laws and policies (Italian D.L.gs. no. 26/2014 art.3.1A; European Directive 63/2010/UE). Mice were bred and maintained at the ICGEB animal Bio-experimentation facility in sterile isolators (12h/12h light/dark cycle, T 21 °C ± 2 °C, RH 55% ± 10%) and received a standard chow diet and water *ad libitum*. Experiments involving *ex-vivo* bone marrow culture were performed using WKO males' mice and WT littermates as controls of 8-12 weeks of age. In case of *in-vivo* experiments WASp$^{-/-}$ and heterozygous WASp$^{+/-}$ female mice of 10 and 25 weeks of age were employed.

Primary murine cell culture and human PBMCs

Mouse BM-derived macrophages (BMDMs) were generated in vitro from BM of C57BL/6 WT or WKO mice using recombinant M-CSF (rMCSF, Miltenyi Biotech). BM-derived cells were cultured at a concentration of 1×10^6 cells/mL in non-treated 10 cm Petri dishes (Greiner) for 7 days using RPMI 1640 supplemented with 20 ng/mL rMCSF, 10% low endotoxin FBS (Euroclone), plus 50 µg/mL gentamicin (Gibco), 50 µM 2-mercaptoethanol (Gibco) and 1% Sodium Pyruvate (Gibco). Mouse BM-derived DCs (BMDCs) were generated in vitro from bone marrow of C57BL/6 WT or WKO mice using supernatant containing GM-CSF produced from J558 cell line. BMDCs were cultured at a concentration of 1.5×10^6 cells/mL in Nunc-nontreated 6 well-plates (Thermo Fisher Scientific) for 7 days using IMDM supplemented with 30% GM-CSF supernatant, 10% low endotoxin FBS (Euroclone), 50 µg/mL gentamicin (Gibco) and 50 µM 2-mercaptoethanol (Gibco). All mice used for BM culture were males of 7 to 12 weeks of age.

Human PBMCs were gently provided by Dr. Anna Villa from San Raffaele Institute and shipped in RNA Later™ (Thermo Fischer Scientific). RNA extraction and library preparation for sequencing are

481 described in the respective paragraphs. Information about patients is provided in supplementary

482 table 5.

Flow cytometry and sorting of alveolar macrophages

Alveolar macrophages (AMO) were obtained from homozygous WASp$^{-/-}$ or heterozygous WASp$^{+/-}$

female mice of 10 or 25 weeks old. After animal sacrifice, lungs were perfused in PBS to wash away

red blood cells and digested with collagenase type 2 (265 U/mL; Worthington) and DNase I

(250 U/mL; Millipore-Sigma) in complete media at 37°C for 30'. Collagenase activity was then

stopped by EDTA 10 mM, and cell suspension was filtered using 100 μm cell strainer (Corning). The

obtained single cell suspension was enriched for CD11c+ population using the mouse CD11c

microbeads (Miltenyi) and stained with anti-CD11c, anti-CD45 and anti-SiglecF. 5×10^6 cells/mL were

resuspended in PBS 2% FBS for sorting and at least 100.000 AMO were sorted with a BD FACSAria II

in PBS 20% FBS. Sorted AMO were centrifuged and frozen at -70°C in 350 μL of RLT lysis buffer prior

to RNA extraction. Flow cytometry analysis was performed in a BD FACSCelesta™ starting from 1/5

of single cell suspension before CD11c enrichment, using antibodies reported in supplementary

table 6 and Counting Bright beads (Miltenyi). All antibodies were incubated 1 h at 4°C in dark

conditions. After antibody staining, samples were incubated 20' at 4°C in dark conditions with Live

and Dead fixable aqua (Invitrogen) 1:1000 in PBS and fixed over-night in PBS 1% PFA before a final

wash and acquisition. Data analysis was performed on FlowJo software.

RNA extraction and gene expression analysis

RNA from BMDM at steady state was obtained using of TRIzol^{TM} reagent (Thermo Fischer Scientific)

and treated with DNaseI before retro transcription. RNA extraction and purification of sorted AMO,

confined BMDM and human PBMCs was performed by RNeasy micro kit (Qiagen). Retro

transcription and qPCR were done by PrimeScript™ RT Reagent Kit (Takara) and TB Green® Premix

(Takara). Real time machine used for qPCR was a CFX96 Touch (Bio-Rad). Specific primers used for

amplification were the following (5'-> 3'):

Mouse: Gapdh fwd AGAAGGTGGTGAAGCAGGCAT, Gapdh rev CGAAGGTGGAAGAGTGGGAGT; Ifnb1

fwd GCACTGGGTGGAATGAGTCT, Ifnb1 rev AGTGGAGAGCAGTTGAGGACA; Il1b fwd

TGTAATGAAAGACGGCACACC, Il1b rev TCTTCTTTGGGTATTGCTTGG; Il6 fwd

GAGGATACCACTCCCAACAGA, Il6 rev AAGTGCATCATCGTTGTTCAT; Isg15 fwd

AGCAATGGCCTGGGACCTAAA, Isg15 rev CAGACCCAGACTGGAAAGGG; Mx1 fwd

TCATCAGAGTGCAAGCGAGG, Mx1 rev TCTGATACGGTTTCCTGTGCT; Tnfa fwd

CTGTAGCCCACGTCGTAGC, Tnfa rev TTAAGATCCATGCCGTTG.

<u>Human</u>: Hprt1 fwd AATTATGGACAGGACTGAACGTCTTGCT, rev

TCCAGCAGGTCAGCAAAGAATTTATAGC; Mxa fwd ACAGAACCGCCAAGTCCAAA, rev

GCGGATCAGCTTCTCACCTT; Rsad2 fwd ATCCTTTGTGCTGCCCCTTG, rev

TTGATCTTCTCCATACCAGCTTCC; Il6 fwd AGACAGCCACTCACCTCTTCAG, rev

TTCTGCCAGTGCCTCTTTGCTG; Il1b fwd ATGATGGCTTATTACAGTGGCAA, rev

GTCGGAGATTCGTAGCTGGA.

Lentiviral preparation and BMDM infection

Second generation lentiviral particles were prepared by co-transfecting HEK-293T cells with plasmid

of interest, relative packaging and envelope. 48 hours after transfection, supernatant-containing

virus was collected, centrifuged 300g for 5' to remove cell debris and filtered through 0.45 µm filter.

Cleared supernatants were mixed with Lenti-X concentrator (Takara) and concentrated 100x,

following manufacturer instructions. 20 µL of concentrated lentiviral particles were used to infect 2

x 10^6 BMDM at day 4 in a 6-well plate. At day 6, viral-containing supernatant was removed and

attached cells were washed 2x in sterile PBS. Infected BMDM were used at day 7 after two more

washes in PBS to get rid of any remaining viral particle. Plasmid used for generation of lentivectors

were the following: pMD2.G (Addgene #12259), psPAX2 (Addgene #12260), SEWW-eGFP-WASp

(Blundell et al., 2008), pCDH-NLS-GFP (Addgene #132772), pTRIP-mCherry-FLAG-cGAS

E225A/D227A (Addgene #127657).

CRISPR-Cas9 RNPs electroporation

CRISPR-Cas9-mediated gene knock-out was achieved through electroporation of ribonucleoprotein

(RNP) particles containing crRNA, tracRNA and Cas9 protein complexes. spCas9 and RNA oligos were

purchased from IDT, annealed and mixed following manufacturer instructions. At day 4 of

differentiation, 1.5 x 10^6 cells were electroporated in 100 µL of media in a NEPA electroporator with

the following parameters: poring pulse 275 V, 4 x 1 ms, interval 50 ms, D. Rate 10 %; transfer pulse

20 V, 5 x 50 ms, interval 50 ms, D. Rate 40 %, polarity +/-. After 24 h media was changed and cells

were used for experiments at day 7 of differentiation.

6-well plate confiner

Mechanical confinement was performed in 6-well plates (Liu et al., 2015). 12 mm glass coverslips were washed with hand soap, rinsed in ddH$_2$O, then in isopropanol and dried with a pressurized air gun. The clean coverslips were treated in a plasma cleaner (Diener Zepto) at 40W for 30 s. To make pillars of desired height, polydimethylsiloxane (PDMS; SYLGARD 184, Dow) was mixed at a ratio of PDMS A:cross-linker B 1:10 (wt/wt) and poured on top of wafer molds-containing pillars of 3 µm or 6 µm. The wafer molds, containing pillars of 3 µm or 6 µm, were fabricated by conventional micromachining from square pieces (25x25 mm) of a 500 µm thick silicon wafer. After cleaning the wafer with acetone and isopropanol, a positive photoresist MEGAPOSIT™ SPR™220 3.0 or 7.0 was spinned on. After the soft bake, they were exposed to a UV lamp for 30 seconds with a suitable mask and subjected a post-bake process. The development was carried out with MF24A. To functionalized with micropillars, glass coverslips were pressed to the wafers and baked at 85°C for 20 min before being carefully detached with a blade. After a wash in isopropanol, coverslips with micropillars were dried and plasma treated for 30 more seconds. Finally, these coverslips were washed in PBS and incubated with non-adhesive PEG solution (Sigma) 500 µg/mL in PBS for 30 min at RT and UV-sterilized for 20'. Before confinement, functionalized glass coverslips were washed in H$_2$O and PBS and kept in sterile PBS at 4°C for a maximum of 48 h. To perform confinement experiments, we used large PDMS pistons placed on the plate lid to press the coverslips on the bottom of the plate. A drop of 250 µL containing 300.000 cells for immunofluorescent assays or 500.000 cells for RNA collection was confined for each well. 6-well plates with a glass bottom were used for cell imaging (MatTek, P06G-1.5-20-F); whereas non-treated 6-well plates (Corning) were adopted for RNA experiments. Nuclear blebbing quantification was performed right after confinement on live cells based on Hoechst 33342 staining (1:10.000 in PBS). Both types of confinements were kept for 1 h before fixation or RNA extraction. Fixation for IF was performed directly on confined cells, by leaving 4% PFA for 45 min before removing the lid, quenching PFA with 0,1 M glycine in PBS and washing cells 3x with PBS before proceeding with IF staining.

Microchannel preparation and live imaging

Microchannels MC004 were purchased from 4DCell. And prepared following manufactures instructions. Coating was performed using 10 µg/mL fibronectin for 1 h at RT. Before utilization, channels were incubated from 30 min to overnight in 2 mL of BMDM media. Timelapse imaging was

carried out overnight on a Nikon Eclipse TiE inverted microscope equipped with a 20x/0.75 NA dry objective, a motorized stage and a humidified chamber at 37°C and 5% CO_2. Live imaging of migrating macrophages in microchannels was recorded with a frame every 5 min, based on NLS-GFP (FITC channel) and DIC (transmitted light). After overnight migration, cells inside channels were fixed by adding 10 µL of 4% PFA to each accession port and waiting for 1 h at RT before washing in PBS. Channels were kept in PBS 0,02% Sodium Azide until further imaging. Hoechst 1:1000 in PBS was added for at least 1 h before imaging fixed microchannels to facilitate detection and segmentation of nuclear morphology and herniations Fixed samples were imaged on a Zeiss LSM880 confocal microscope with a 63x/1.4 NA oil objective for individual nuclei and a 20x/0.60 NA dry objective for large field acquisitions and quantification of cell entry.

Immunofluorescence imaging

Immunofluorescence staining at steady state was done with 50.000 BMDM seeded on fibronectin-coated (10ug/mL, Millipore Sigma) 12 mm microscope slides (Epredia, Thickness value #1). Cells were left 30' at 37°C and 5% CO_2 to adhere. Fixation was performed in 4% PFA for 7 min at RT, followed by quenching in 0,1 M glycine in PBS for 3 min. After 3x washes in PBS, cells were permeabilized in PBS 1% BSA, 0,1% Triton-X100, blocked in PBS 3% BSA, 10 mM Hepes and stained ON at 4°C in primary antibodies diluted in PBS 0,2% BSA, 0,1% Triton-X100. Secondary antibodies were incubated at RT for 2h, followed by eventual phalloidin staining (Thermo Fischer Scientific, used 1:500) and 10 min nuclear staining with Hoechst 33342 (Invitrogen) 1:1000 in PBS. Slides were mounted with Mowiol and imaged at Zeiss LSM880 confocal microscope with a 63x/1.4 NA oil objective, or a 20x/0.60 NA dry objective. A full list of antibodies, dyes and probes used is presented in supplementary table 6.

AFM based indentation measurements

AFM-based indentation measurements were performed by NanoWizard III AFM (JPK Instruments, Bruker) integrated with an inverted optical microscope (Axiovert 200; Zeiss) for sample observation. Cells were analysed in petri-dish and in culture medium at room temperature. The status of cells was constantly monitored by optical microscope. Indentation of the whole cell body was performed using MLCT-SPH-5um probes (Bruker), which have a cylindrical tip shank ending in a spherical tip with a 5 µm radius and a nominal spring constant of 0.06 N/m. Once localized the cell by optical microscope, the probe was brought over the nucleus area and pressed down to indent the cell. The

motion of the z-piezo and the force were recorded. On each cell three to five force–displacement (F-D) curves were acquired with a force load of 0.5-1nN and at a rate of at 5μm/s in closed-loop feedback mode. Cell elastic properties were assessed by determining the Young's modulus (E), which can be evaluated by fitting the force (F) indentation (δ) curves with the Hertz model.

$$F = \frac{4}{3}\frac{E}{(1-\nu^2)}\sqrt{R}\delta^{3/2}$$

where F is the force, R is the radius of the spherical indenter and ν is the sample's Poisson ratio (set to 0.5 for the cell) and δ is the indentation depth.

Data fitting was performed on the approaching F–D curves. JPK Data Processing software was used to convert the F-D curves into the force–indentation curves by subtracting the cantilever bending from the signal height to calculate indentation. The resulting force-indentation curves were fitted with the Hertz model for indentation depth of 500 nm.

Transwell migration assay

24-well plates bearing transwell inserts of 5 μm and 3 μm. 5 μm plates were from Corning; 3 μm plates from Thermo Fischer Scientific. At day 7 of differentiation, cells were detached using MACS buffer (0,5% BSA, 2 mM EDTA in PBS), counted and resuspended in media without FBS containing 200 ng/mL of LPS (Invivogen) for 30 min in incubator. After LPS washout, cells were plated at a concentration of 200.000 cells in 200 μL for each insert in complete media. On the bottom of each well, 600 μL of complete media supplemented with 250 ng/mL of CCL19 (Peprotech) as chemoattractant were added to promote transmigration. Cells were kept 5 h in incubator to allow migration, following by collection of migrated cells for flow cytometry or IF analysis. Flow cytometry quantification of migrating cells was performed at BD Accuri™ C6 flow cytometer, using Counting Bright beads and normalizing the number of transmigrating cells to the total number of plated cells.

Protein extraction and western blot

Cells were lysed in an appropriated volume of RIPA buffer (50 mM Tris/HCl pH 7.5, 150 mM NaCl, 1% NP-40, 0,1% SDS, 0,1% DOC in ddH$_2$O) supplemented 1:30 with protease inhibitor cocktail (Sigma P8340). After centrifugation at 17000 g for 20 min, supernatant-containing protein extracts were kept at -20°C until further processing. Total protein concentration was quantified by means of BCA assay (Thermo Fisher), following manufacturer instructions. At least 20 μg of total protein lysates

were loaded on a 10% custom made polyacrylamide gel and transferred on PVDF membranes upon

SDS-page electrophoresis. Membranes were blocked at least 1 h at RT in 1x TBS, 0,1% Tween 20

(TBST), 3% dry fat milk. Primary antibodies were incubated ON at 4°C on a shaker. After 3x washes

in TBST, secondary HRP-conjugated antibodies were kept 1-2 h at RT. Chemiluminescence signal was

developed by Clarity ECL substrate (Bio-Rad) on a ChemiDoc™ Touch (Bio-Rad). A list of antibodies

used is reported in supplementary table 6.

Library preparation and RNA-sequencing

For mouse and human samples, total RNA was extracted using RNeasy Micro kit (Qiagen), according

to the manufacturer's protocol. RNA purity, integrity and quantity was checked at TapeStation 2200

(Agilent Technologies Inc., Santa Clara, CA, USA). RNA-seq libraries were prepared from 3 ng (human

PBMCs) or 5 ng (mouse BMDM) of total RNA using mRNA Library Prep Kit (WATCHMAKER) following

manufacture instruction. Briefly, we selected poly(A) using magnetic beads to focus sequencing on

polyadenylated transcripts. After RNA fragmentation, a step of cDNA synthesis was performed. The

resulting fragments were used for A-tailing and were ligated to the adapters and Index Anchors

(TWIST bioscience, Twist universal system). PCR amplification (21 cycles) was performed to generate

the final indexed cDNA libraries. Individual library quantification and quality assessment was

performed using TapeStation 2200. Libraries were sent to Macrogen to be sequences on Novaseq

(Human PBMCs) or NovaseqX (Mouse BMDM) instrument on Illumina platform as paired end run

150 bp reads. A total of 6 PBMCs samples were sequenced before as well after gene therapy. Four

biological replicates of BMDM confined per genotype and three replicates for unconfined control

were sequenced.

RNA-sequencing analysis

Raw FastQ files were downloaded from Macrogen. Quality control, trimming and alignment were

performed using standard pipelines for FastQC, Trimgalore and STAR.

<u>Confiner data</u>: Raw reads count matrix outputs were imported and analysed in R software (R version

4.3.1). Normalized reads were obtained by standard DEseq2 functions and used for heatmaps

visualization. DESeq2 package (Love et al., 2014) was further used for differential gene expression

analysis. GSEA was performed on gseGO function of ClusterProfiler package on Log2FC ordered

genes with p-value cutoff of 0.01 and Benjamini-Hochberg p-value correction. DEGs were considered

as genes bearing p-value< 0.05 and a Log2FC> 1.2 or Log2FC< -1.2. GO analysis was performed on

DEGs using enrichGO function, based on biological processes, with a p-value cutoff of 0.05 and Benjamini-Hochberg statistical correction. Visualizations of functional enrichment results were obtained by ClusterProfiler and ggplot2 packages.

PBMCs data: DESeq2 analysis was performed starting on unnormalized reads comparing WAS pre- vs post-gene therapy samples. gseGO from ClusterProfiler have been used to perform GSEA on Log2FC ordered genes, using Entrez ID as key type, "BP" as ontology and 0.05 as p-value cutoff, without further statistical correction methods. Full list of GSEA is available as Supplementary Table 7.

Analysis of publicly available dataset from GSE107963: Data were downloaded as TPM normalized count matrix from GSE107963 (www.ncbi.nlm.nih.gov/geo/download/?acc=GSE107963, data deposited on 15.12.2022 and downloaded on 13.08.2024). Samples used for analysis refer to 3 clones of a WT iPSC cell line (WASp +) and 3 isogenic CRISPR-Cas9-mediated Was KO iPSCs clones (WASp -) differentiated into macrophages. TPM matrix was filtered and further processed to keep only genes with non-zero reads. Log2(TPM+1) was used for heatmap visualization.

Image analysis

All raw images were analysed in Fiji ImageJ (Schindelin et al., 2012), using custom-written macros.

Lamin A/C alteration: Lamin A/C alteration score has been calculated based on standard deviation (σ) of Lamin A/C signal. Nuclear segmentation has been based on Hoechst 33342 channel of z-projected nuclei, with a size threshold > 10 μm^2, to exclude potential micronuclei from the analysis. Alteration score has been plotted as σ/10. Masks of Lamin A/C wrinkles for representative pictures has been obtained by an adaptive threshold method.

Chromocenters and H3K9me2 quantification: to quantify the number of chromocenters and the area of peripheral heterochromatin, nuclei were first segmented based on Hoechst 33342 signal. Chromocenters were detected based on an adaptive threshold and a size cutoff. Area of peripheral heterochromatin has been calculated as the sum of chromocenters' area in close proximity to nuclear edges, as those within 1 μm from nuclear borders. The same cutoff has been used for H3K9me2 peripheral heterochromatin, measuring fluorescent intensity instead of chromocenters area.

Microchannels analysis and WASp-GFP distribution: quantification of cells inside microchannels an nuclear rupture events have been conducted manually. Kymographs of NLS-GFP migrating cells were generated using a semi-automated macro in ImageJ. WASp-GFP distribution has been obtained in a

semi-automated way as follow. Briefly individually cropped cells inside channels were subjected to automatic nuclear detection. Nuclear centre of mass was then moved to the centre of the image and all individual cells were cropped and horizontally aligned. The size of the cells was normalized at 600 pixels and expanded 50 times before saving at a resolution size of 1200x200 pixels (w x h). Size normalized cells were then merged by Stack>Z project>Average Intensity to obtain an average intensity map profile of WASp distribution.

Statistics

Primary data were collected in Microsoft Excel and statistical analyses were performed using GraphPad Prism version 9. For comparison between two or more groups with normally distributed data two-tailed Student's T-test, one-way ANOVA or two-way ANOVA were used as appropriate. The non-parametric Kruskal-Walli's test with Dunn's multiple comparison was performed to compare three or more unmatched groups. In case of normalized values to 1 or 0, statistics has been calculated by a one-sample t-test. P values <= 0.05 were considered significant, if not stated differently.

Supplementary information

Description of supplementary data

- Supplementary table 1: Gene list of inflammatory signatures used in Fig.1
- Supplementary table 2: RNA-seq data normalized reads BMDM
- Supplementary table 3: RNA-seq data normalized reads PBMCs
- Supplementary table 4: Gene lists for signatures used in Fig.6
- Supplementary table 5: WAS patients' information
- Supplementary table 6: List of antibodies used in the work
- Supplementary Table 7: GSEA on WAS PBMCs dataset

Data availability

Sequencing data have been deposited at GEO database: BMDM confiner data (accession: GSE285024), publicly available as of the 24/12/2024 and PBMCs WAS data (accession: GSE285174), publicly available as of the 25/12/2024. Source data are provided in supplementary files. Any

additional information required to reanalyse the data reported in this paper is available from the lead contact upon formal request.

Author contributions

R.A, **G.M.P** and **F.B** conceived the study and designed the experiments. **R.A** performed most of the experiments, data analysis, and prepared the figures. **R.A** and **F.B** wrote the manuscript. **G.B** helped in the execution of experiments and performed WASp-GFP confinement. **Z.A** provided training and support for the confiner assays and helped in performing confiner experiments in Fig.4 (A, B and C). **M.M** assisted in image analysis. **G.M.P** and **L.L.R** assisted in the experimental part and critical discussion of the data. **G.M.P** prepared library for sequencing. **M.C** prepared photolithographic molds used for confiner slides fabrication. **L.A** performed and analysed atomic force microscopy. **M.C.C**, **F.F**, **A.A** and **A.V** provided human PBMCs samples. **A.-M.L.-D** critically discussed the data and the manuscript. **F.B** supervised the whole study. All authors read and approved the final version of the manuscript.

Disclosure and competing interest's statement

The authors declare no competing interests.

Acknowledgements

This work was supported by Telethon grant GGP20102 to **F.B**. **G.B**, **G.M.P** and **L.L.R**- were supported by ICGEB Arturo Falaschi pre- and post-doctoral fellowships. **R.A** was supported by Italian Telethon. **R.A** received an EMBO scientific exchange grant (10017) to perform part of the work in the lab of **A.-M.L.-D**. We Thank Luca Triboli from ICGEB Trieste to help discussing bioinformatic data analysis.

References

Ablasser, A., and Gulen, M. F. (2016). The role of cGAS in innate immunity and beyond. *J. Mol. Med.* 94, 1085–1093. doi:10.1007/s00109-016-1423-2.

Alraies, Z., Rivera, C. A., Delgado, M. G., Sanséau, D., Maurin, M., Amadio, R., et al. (2024). Cell shape sensing licenses dendritic cells for homeostatic migration to lymph nodes. *Nat. Immunol.* 25. doi:10.1038/S41590-024-01856-3.

Amadio, R., Piperno, G. M., and Benvenuti, F. (2021). Self-DNA Sensing by cGAS-STING and TLR9 in Autoimmunity : Is the Cytoskeleton in Control ? *Front. Immunol.* 12. doi:10.3389/fimmu.2021.657344.

Bastianello, G., Porcella, G., Beznoussenko, G. V., Kidiyoor, G., Ascione, F., Li, Q., et al. (2023). Cell stretching activates an ATM mechano-transduction pathway that remodels cytoskeleton and chromatin. *Cell Rep.* 42, 113555. doi:10.1016/j.celrep.2023.113555.

Bell, E. S., Shah, P., Zuela-Sopilniak, N., Kim, D., Varlet, A.-A. A., Morival, J. L. P. P., et al. (2022). Low lamin A levels enhance confined cell migration and metastatic capacity in breast cancer. *Oncogene* 41, 1–20. doi:10.1038/S41388-022-02420-9.

Blundell, M. P., Bouma, G., Calle, Y., Jones, G. E., Kinnon, C., and Thrasher, A. J. (2008). Improvement of Migratory Defects in a Murine Model of Wiskott–Aldrich Syndrome Gene Therapy. *Mol. Ther.* 16, 836–844. doi:10.1038/MT.2008.43.

Blundell, M. P., Worth, A., Bouma, G., and Thrasher, A. J. (2010). The Wiskott-Aldrich syndrome: The actin cytoskeleton and immune cell function. *Dis. Markers* 29, 157–175. doi:10.3233/DMA-2010-0735.

Calixto Vieira, R., Goncalves, L., Westerberg, L. S., Rhaissa, C., Vieira, C., and Goncalves Pinho, L. (2023). Understanding immunoactinopathies: A decade of research on WAS gene defects. *Pediatr. Allergy Immunol.* 34, e13951. doi:10.1111/PAI.13951.

Cervantes-Luevano, K. E., Caronni, N., Castiello, M. C., Fontana, E., Piperno, G. M., Naseem, A., et al. (2018). Neutrophils drive type I interferon production and autoantibodies in patients with Wiskott-Aldrich syndrome. *J. Allergy Clin. Immunol.* 142, 1605-1617.e4. doi:10.1016/j.jaci.2017.11.063.

Davidson, P. M., Denais, C., Bakshi, M. C., and Lammerding, J. (2014). Nuclear deformability constitutes a rate-limiting step during cell migration in 3-D environments. *Cell. Mol. Bioeng.* 7, 293. doi:10.1007/S12195-014-0342-Y.

De Silva, N. S., Siewiera, J., Alkhoury, C., Nader, G. P. F., Nadalin, F., de Azevedo, K., et al. (2023). Nuclear envelope disruption triggers hallmarks of aging in lung alveolar macrophages. *Nat. Aging* 3, 1251–1268. doi:10.1038/s43587-023-00488-w.

Decout, A., Katz, J. D., Venkatraman, S., and Ablasser, A. (2021). The cGAS–STING pathway as a therapeutic target in inflammatory diseases. *Nat. Rev. Immunol.* doi:10.1038/s41577-021-00524-z.

Delgado, M.-G. et al. (2024). The Arp2/3 complex maintains the epidermal Langerhans cell network and DNA integrity. *Under Submiss.*

Denais, C. M., Gilbert, R. M., Isermann, P., McGregor, A. L., Te Lindert, M., Weigelin, B., et al. (2016).
Nuclear envelope rupture and repair during cancer cell migration. *Science (80-.).* 352. Available at:
/pmc/articles/PMC4833568/ [Accessed March 24, 2023].
Dou, Z., Ghosh, K., Vizioli, M. G., Zhu, J., Sen, P., Wangensteen, K. J., et al. (2017). Cytoplasmic chromatin
triggers inflammation in senescence and cancer. *Nature* 550. doi:10.1038/nature24050.
Dunphy, G., Flannery, S. M., Almine, J. F., Connolly, D. J., Paulus, C., Jønsson, K. L., et al. (2018). Non-
canonical Activation of the DNA Sensing Adaptor STING by ATM and IFI16 Mediates NF-κB Signaling
after Nuclear DNA Damage. *Mol. Cell*. doi:10.1016/j.molcel.2018.07.034.
En, A., Bogireddi, H., Thomas, B., Stutzman, A. V., Ikegami, S., LaForest, B., et al. (2024). Pervasive nuclear
envelope ruptures precede ECM signaling and disease onset without activating cGAS-STING in Lamin-
cardiomyopathy mice. *Cell Rep.* 43, 114284. doi:10.1016/j.celrep.2024.114284.
Gaertner, F., Reis-Rodrigues, P., de Vries, I., Hons, M., Aguilera, J., Riedl, M., et al. (2022). WASp triggers
mechanosensitive actin patches to facilitate immune cell migration in dense tissues. *Dev. Cell* 57, 47-
62.e9. doi:10.1016/j.devcel.2021.11.024.
Ginhoux, F., and Guilliams, M. (2016). Tissue-Resident Macrophage Ontogeny and Homeostasis. *Immunity*
44, 439–449. doi:10.1016/j.immuni.2016.02.024.
Harada, T., Swift, J., Irianto, J., Shin, J. W., Spinler, K. R., Athirasala, A., et al. (2014). Nuclear lamin stiffness
is a barrier to 3D migration, but softness can limit survival. *J. Cell Biol.* 204, 669.
doi:10.1083/JCB.201308029.
Härtlova, A., Erttmann, S. F., Raffi, F. A. M., Schmalz, A. M., Resch, U., Anugula, S., et al. (2015). DNA
Damage Primes the Type I Interferon System via the Cytosolic DNA Sensor STING to Promote Anti-
Microbial Innate Immunity. *Immunity* 42, 332–343. doi:10.1016/j.immuni.2015.01.012.
Kim, J. K., Louhghalam, A., Lee, G., Schafer, B. W., Wirtz, D., and Kim, D. H. (2017). Nuclear lamin A/C
harnesses the perinuclear apical actin cables to protect nuclear morphology. *Nat. Commun. 2017 81* 8,
1–13. doi:10.1038/s41467-017-02217-5.
Kovacs, M. T., Vallette, M., Wiertsema, P., Dingli, F., Loew, D., Nader, G. P. de F., et al. (2023). DNA damage
induces nuclear envelope rupture through ATR-mediated phosphorylation of lamin A/C. *Mol. Cell* 83,
3659. doi:10.1016/J.MOLCEL.2023.09.023.
Le, H. Q., Ghatak, S., Yeung, C. Y. C., Tellkamp, F., Günschmann, C., Dieterich, C., et al. (2016). Mechanical
regulation of transcription controls Polycomb-mediated gene silencing during lineage commitment.
Nat. Cell Biol. 18, 864–875. doi:10.1038/ncb3387.
Lee, P. P., Lobato-Márquez, D., Pramanik, N., Sirianni, A., Daza-Cajigal, V., Rivers, E., et al. (2017). Wiskott-
Aldrich syndrome protein regulates autophagy and inflammasome activity in innate immune cells.
Nat. Commun. 8, 1–13. doi:10.1038/s41467-017-01676-0.
Liang, F., Peng, C., Luo, X., Wang, L., Huang, Y., Yin, L., et al. (2023). A single-cell atlas of immunocytes in the

spleen of a mouse model of Wiskott-Aldrich syndrome. *Cell. Immunol.*, 104783.

doi:10.1016/j.cellimm.2023.104783.

Linder, S., Nelson, D., Weiss, M., and Aepfelbacher, M. (1999). Wiskott-Aldrich syndrome protein regulates

podosomes in primary human macrophages. *Cell Biol. Commun. by John A. Glomset* 96, 9648–9653.

Available at: www.pnas.org. [Accessed March 9, 2022].

Liu, Y. J., Le Berre, M., Lautenschlaeger, F., Maiuri, P., Callan-Jones, A., Heuzé, M., et al. (2015).

Confinement and low adhesion induce fast amoeboid migration of slow mesenchymal cells. *Cell* 160,

659–672. doi:10.1016/j.cell.2015.01.007.

Love, M. I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-

seq data with DESeq2. *Genome Biol.* 15, 1–21. doi:10.1186/s13059-014-0550-8.

Maekawa, H., Inoue, T., Ouchi, H., Jao, T. M., Inoue, R., Nishi, H., et al. (2019). Mitochondrial Damage

Causes Inflammation via cGAS-STING Signaling in Acute Kidney Injury. *Cell Rep.* 29, 1261-1273.e6.

doi:10.1016/j.celrep.2019.09.050.

Magnani, A., Semeraro, M., Adam, F., Booth, C., Dupré, L., Morris, E. C., et al. (2022). Long-term safety and

efficacy of lentiviral hematopoietic stem/progenitor cell gene therapy for Wiskott-Aldrich syndrome.

Nat. Med. 28. doi:10.1038/s41591-021-01641-x.

Mu, X., Tseng, C., Hambright, W. S., Matre, P., Lin, C. Y., Chanda, P., et al. (2020). Cytoskeleton stiffness

regulates cellular senescence and innate immune response in Hutchinson–Gilford Progeria Syndrome.

Aging Cell 19, 1–16. doi:10.1111/acel.13152.

Mukherjee, S., Abdisalaam, S., Bhattacharya, S., Srinivasan, K., Sinha, D., and Asaithamby, A. (2019).

"Mechanistic link between DNA damage sensing, repairing and signaling factors and immune

signaling," in *Advances in Protein Chemistry and Structural Biology* (Academic Press Inc.), 297–324.

doi:10.1016/bs.apcsb.2018.11.004.

Naseem, A., Steinberg, Z., and Cavazza, A. (2022). Genome editing for primary immunodeficiencies: A

therapeutic perspective on Wiskott-Aldrich syndrome. *Front. Immunol.* 0, 4553.

doi:10.3389/FIMMU.2022.966084.

Nava, M. M., Miroshnikova, Y. A., Biggs, L. C., Whitefield, D. B., Metge, F., Boucas, J., et al. (2020).

Heterochromatin-Driven Nuclear Softening Protects the Genome against Mechanical Stress-Induced

Damage. *Cell* 181, 800. doi:10.1016/J.CELL.2020.03.052.

Oliveira, M. M. S., Kung, S. Y., Moreau, H. D., Maurin, M., Record, J., Sanséau, D., et al. (2022). The WASp

L272P gain-of-function mutation alters dendritic cell coordination of actin dynamics for migration and

adhesion. *J. Leukoc. Biol.* 111, 793–803. doi:10.1002/JLB.1AB0821-013RR.

Patteson, A. E., Vahabikashi, A., Pogoda, K., Adam, S. A., Mandal, K., Kittisopikul, M., et al. (2019). Vimentin

protects cells against nuclear rupture and DNA damage during migration. *J. Cell Biol.* 218, 4079–4092.

doi:10.1083/JCB.201902046.

Piperno, G. M., Naseem, A., Silvestrelli, G., Amadio, R., Caronni, N., Cervantes-Luevano, K. E., et al. (2020).
Wiskott-Aldrich syndrome protein restricts cGAS/STING activation by dsDNA immune complexes. *JCI*
Insight 5. doi:10.1172/jci.insight.132857.

Prete, F., Catucci, M., Labrada, M., Gobessi, S., Castiello, M. C., Bonomi, E., et al. (2013). Wiskott-Aldrich
syndrome protein-mediated actin dynamics control type-I interferon production in plasmacytoid
dendritic cells. *J. Exp. Med.* 210, 355–374. doi:10.1084/jem.20120363.

Reis-rodrigues, P., Canigova, N., Avellaneda, M. J., Gaertner, F., Vaahtomeri, K., Riedl, M., et al. (2024).
Global coordination of protrusive forces in migrating immune cells. *bioRxiv*.

Rodríguez-Nuevo, A., Díaz-Ramos, A., Noguera, E., Díaz-Sáez, F., Duran, X., Muñoz, J. P., et al. (2018).
Mitochondrial DNA and TLR9 drive muscle inflammation upon Opa1 deficiency. *EMBO J.* 37.
doi:10.15252/embj.201796553.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., et al. (2012). Fiji: An open-
source platform for biological-image analysis. *Nat. Methods* 9, 676–682. doi:10.1038/nmeth.2019.

Sereni, L., Castiello, M. C., Di Silvestre, D., Della Valle, P., Brombin, C., Ferrua, F., et al. (2019). Lentiviral
gene therapy corrects platelet phenotype and function in patients with Wiskott-Aldrich syndrome. *J.*
Allergy Clin. Immunol. 144, 825–838. doi:10.1016/j.jaci.2019.03.012.

Sereni, L., Castiello, M. C., Marangoni, F., Anselmo, A., di Silvestre, D., Motta, S., et al. (2018). Autonomous
role of Wiskott-Aldrich syndrome platelet deficiency in inducing autoimmunity and inflammation. *J.*
Allergy Clin. Immunol. 142, 1272–1284. doi:10.1016/j.jaci.2017.12.1000.

Shen, C., Postat, J., Cerf, A., Merino, M., Bhagrath, A., Patel, D., et al. (2024). DOCK8 regulates a
mechanosensitive actin redistribution that maintains immune cell cohesion and protects the nucleus
during migration. *bioRxiv*, 2024.07.26.605273. Available at:
https://www.biorxiv.org/content/10.1101/2024.07.26.605273v1%0Ahttps://www.biorxiv.org/content
/10.1101/2024.07.26.605273v1.abstract.

Sladitschek-Martens, H. L., Guarnieri, A., Brumana, G., Zanconato, F., Battilana, G., Xiccato, R. L., et al.
(2022). YAP/TAZ activity in stromal cells prevents ageing by controlling cGAS–STING. *Nature* 607, 790–
798. doi:10.1038/s41586-022-04924-6.

Snapper, S. B., Rosen, F. S., Mizoguchi, E., Cohen, P., Khan, W., Liu, C. H., et al. (1998). Wiskoff-Aldrich
syndrome protein-deficient mice reveal a role for WASP in T but not B cell activation. *Immunity* 9, 81–
91. doi:10.1016/S1074-7613(00)80590-7.

Song, X., Aw, J. T. M., Ma, F., Cheung, M. F., Leung, D., and Herrup, K. (2021). DNA repair inhibition leads to
active export of repetitive sequences to the cytoplasm triggering an inflammatory response. *J.*
Neurosci. 41, 9286–9307. doi:10.1523/JNEUROSCI.0845-21.2021.

Sudhakar, M., Rikhi, R., Loganathan, S. K., Suri, D., and Singh, S. (2021). Autoimmunity in wiskott–aldrich
syndrome: Updated perspectives. *Appl. Clin. Genet.* 14, 363–388. doi:10.2147/TACG.S213920.

Tang, W., Chen, X., Wang, X., Zhu, M., Shan, G., Wang, T., et al. (2023). Indentation induces instantaneous
nuclear stiffening and unfolding of nuclear envelope wrinkles. *Proc. Natl. Acad. Sci. U. S. A.* 120,
e2307356120. Available at: https://www.pnas.org/doi/abs/10.1073/pnas.2307356120 [Accessed
November 7, 2023].

Thiam, H. R., Vargas, P., Carpi, N., Crespo, C. L., Raab, M., Terriac, E., et al. (2016). Perinuclear Arp2/3-driven
actin polymerization enables nuclear deformation to facilitate cell migration through complex
environments. *Nat. Commun.* 7, 1–14. doi:10.1038/ncomms10997.

Vargas, P., Chabaud, M., Thiam, H. R., Lankar, D., Piel, M., and Lennon-Dumenil, A. M. (2016). Study of
dendritic cell migration using micro-fabrication. *J. Immunol. Methods* 432, 30–34.
doi:10.1016/J.JIM.2015.12.005.

Xian, H., Watari, K., Sanchez-Lopez, E., Offenberger, J., Onyuru, J., Sampath, H., et al. (2022). Oxidized DNA
fragments exit mitochondria via mPTP- and VDAC-dependent channels to activate NLRP3
inflammasome and interferon signaling. *Immunity* 55, 1370-1385.e8.
doi:10.1016/j.immuni.2022.06.007.

Yuan, B., Zhou, X., Suzuki, K., Ramos-Mandujano, G., Wang, M., Tehseen, M., et al. (2022). Wiskott-Aldrich
syndrome protein forms nuclear condensates and regulates alternative splicing. *Nat. Commun. 2022*
131 13, 1–20. doi:10.1038/s41467-022-31220-8.

  Figures and Figure legends

  Figure 1 – WASp deficiency correlates to enhanced expression of inflammatory genes in

  macrophages. (**A**) Alveolar macrophages (AMO) from 10 weeks old mice were harvested and

  processed for flow cytometry. Data show mean fluorescent intensity (MFI) of CD86. Data are from

  one representative of two experiments (n=3 mice per genotype). Two-tailed unpaired t-test;

  P<=0.01. (B**) Quantification of AMO total numbers in 10- and 25-weeks-old adult mice (n=3 mice,

  data are from one representative of two experiments). One-way ANOVA test with multiple

  comparisons, *P<=0.05. (**C**) Inflammatory genes expression in AMO. mRNA expression (RT-qPCR,

  plotted as fold to WT) of the indicated genes in sorted AMO from 10-and 25-weeks-old WKO mice.

  n=3 mice for each condition, one sample t-test, *P<=0.05. (**D**) Inflammatory genes in primary bone

  marrow-derived macrophages (BMDM). RT-qPCR values in WKO are plotted as fold to WT. Number

  of biological replicates are: Tnfa and Il1b (n=3); Ifnb1 (n=4); Mx1 (n=5); Il6 and Isg15 (n=8). One

  sample t-test. *P<=0.05, **P<=0.01. (**E**) Expression of an inflammatory signature (list of genes used

  to calculate inflammatory z-score can be found in Supplementary Table 1) on the GSE107963

  dataset. Control (WASp +) or gene-edited (WASp -) human iPSCs-derived macrophages. (**F**) Heatmap

  of selected inflammatory genes expression (Log2+1 TPM normalized expression, z-score gene-wise

  scaled) of WT and WASp KO iPSCs-derived macrophages from GSE107963. (**G**) Inflammatory score

  in PBMC of a cohort of 6 WAS patients before (WASp -) and after restoration of WASp by gene

  therapy (WASp +). (**H**) Heatmap of selected inflammatory genes expression (Log2+1 normalized

  reads, z-score gene-wise scaled) in human PBMCs from WAS patients pre- and post- gene therapy.

  Supplementary Figure 1 - WASp reduces steady state inflammatory activation of macrophages in-

  vivo and in-vitro. (**A**) Flow cytometry gating strategy to identify AMO in mouse lung tissues. (**B**) GSEA

  analysis of relevant modulated pathways in WAS PBMCs before and after gene therapy correction.

  Values are expressed as normalized enrichment score. (**C**) Expression of the indicated genes in

  individual PBMC samples pre- and post-gene therapy. Data are plotted as Log2 FC post vs pre gene

  therapy, each dot is a patient. One sample t-test; *P<=0.05 and **P<=0.01. (**D**) RT-qPCR validation

  of selected inflammatory genes in 6 WAS patients pre- and post-gene therapy. The dotted line

  represents the mean gene expression of 3 healthy donor controls (HC). One sample t-test: *P<=0.05.

  (**E**) Correlation between inflammatory score (IS) before WASp restoration (y-axis) and IS reduction

(Δ IS) upon WASp re-expression (x-axis). Data combines 3 human iPSCs-derived macrophages and 6

human PBMCs. Data were analysed by linear regression; $P=0.0095$.

Figure 2 –Nuclear parameters are modified in WASp null macrophages. (**A**) WT and WKO BMDM

were plated on glass and labelled with Hoechst to visualise nuclei. Representative images show

examples of the three parameters analysed: area (top), roundness (middle) and micronuclei

(bottom). Scale bar 10 μm. (**B**) Quantification of the three nuclear parameters depicted in A, each

dot is an independent experiment (*N*=6 experiments with *n*>2000 nuclei analysed for each

genotype). Two-tailed unpaired t-test; *$P<=0.05$. (**C**) Representatives immunoblot (top left) and

relative quantification of LMNA/C protein expression relative to GAPDH (top right); *n*=3 biological

replicates. Data were analysed by one sample t-test. *$P<0.05$. (**D**) Expression of lamin isoforms from

RNA-seq analysis of WT and WKO BMDM. Normalized reads for Lamin A/C, LMNB isoforms 1 and 2

and Emerin. Representative data from *n*=2 replicates. Data were analysed by two-way ANOVA test

with multiple comparisons and Sidak's correction method. ****$P<=0.0001$. (**E**) Representatives

maximum z-projection of Lamin A/C staining (top) and the corresponding segmentation mask for

wrinkles detection (bottom). Quantification of LMNA/C alterations as defined in methods. A higher

score indicates less homogeneity and more wrinkles. *n*=405 WT and *n*=308 WKO individual nuclei

measured, in 3 independent experiments. Scale bar 5 μm. Two-tailed unpaired t-test, **** =

$P<0.0001$. (**F**) Representative images of Lamin A/C (grey) and Emerin (red) co-staining (top) and line

intensity profiles (bottom). Histograms depict areas of colocalization at nuclear boundaries (blue

area) and in nuclear wrinkles (red area). Three independent co-labelling experiments were

performed. Scale bar 5 μm.

1000

Supplementary Figure 2 – Chromatin features and cellular mechanics in WT and WASp KO

macrophages. (**A**) Representative images (left) and quantification (right) of the number of

chromocenters (heterochromatin foci) in WT and WKO BMDM nuclei and the mean area of

peripheral heterochromatin foci. *n*=65 individual nuclei measured from *N*=3 independent

experiments. Scale bar 5 μm. Statistics was calculated by two-tailed unpaired t-test. ns, not

significant. (**B**) Representative images (left) and quantification (right) of mean fluorescent intensity

and ratio of peripheral vs central H3K9me2 marker of facultative heterochromatin. *N*=3 independent

experiments, with *n*=65 individual nuclei measured in total. Data were analysed by two-tailed unpaired t-test. ***$P<=0.001$; ns, not significant. Scale bar 5 μm. (**C**) Scheme of AFM-indentation measurement of a single BMDM cell. Bright field image scale bar 50 μm (**D**) Quantification of Young's modulus of *n*=71 WT and *n*=62 WKO BMDM cells, from three independent measurements. Data were analysed by two-tailed unpaired t-test. ***$P<=0.001$.

Figure 3 – WASp controls the deformation and integrity of the nucleus during migration in microchannels. (**A**) Analysis of nuclear parameters in macrophages migrating through 2-8 μm x 5 μm fibronectin coated micro-channels. Representative images of nuclei within channels after overnight migration (applying a nuclear mask). (**B**) Entrance rate of the two genotypes inside channels of different dimensions. Data are from *N*=3 and *N*=4 independent experiments for WT and WKO, respectively. Data were analysed by two-way ANOVA test, *$P<=0.05$. (**C**) Representative examples of WT and WKO macrophage nuclei inside 6 μmx5μm channels (left). Corresponding quantification of the fraction of cells showing herniated nuclei (right). *n*=40 WT and 57 WKO cells in *N*=3 independent experiment, two-tailed unpaired t-test. (**D**) Quantification of the position of the herniation relative to the direction of migration, two-way ANOVA with Sidak's correction method. ns, not significant. (**E**) WT and WKO BMDM were transduced with NLS-GFP to track nuclear rupture during migration. (**F**) Representative time frames (top) and relative kymographs (bottom) showing a nuclear rupture event in WKO migrating inside 6 μm x 5 μm channels, detected as cytosolic leakage of NLS-GFP. Scale bar 10 μm. (**G**) Quantification of the fraction of cells undergoing nuclear rupture during live imaging. *N*=3 independent experiments, with at least 20 individual cells analysed per genotype for each experiment. Two-tails unpaired t-test. *$P<=0.05$. (**H**) Quantification of multiple nuclear ruptures detected during live imaging experiments expressed as fraction of total ruptures recorded.

Supplementary Figure 3 – Constricted migration augments DNA damage in WKO cells. (**A**) Schematic overview of nuclear mask segmentation from raw channel images and quantification of entrance rate as percent of cells inside channels and approaching cells. Approaching cells were defined as those present within 50 μm from channel beginning (yellow dashed line). (**B**) Schematic overview of transwell assay used for BMDCs migration (top) and relative quantification of

transmigrating cells (bottom), in transwell of indicated pore size. N=3 independent replicates are plotted as normalized fold to respective WT (value=1). Statistics was calculated using a one-sample t-test. (**C**) Representative pictures of yH2AX staining pre and post 3 μm transwell migration in BMDCs and relative quantification of number of yH2AX foci per cell in WT and WKO. Data are from N=3 independent experiments with n=178, 202 (WT and WKO pre transwell) and n=13, 33 (WT and WKO post transwell). *P<=0.05 in a standard one-way ANOVA test. (**D**) Flow cytometry quantification of CCR7 median expression in WT and WKO BMDCs before 3 μm transwell migration and following 200 ng/mL of LPS stimulation for 30' (N=3) or in unstimulated control (N=2). Statistics was performed comparing WT and WKO after LPS stimulation by an unpaired t-test. ns, not significant.

Figure 4 – Vertical confiners induce higher deformation and rupture in WASp-deficient macrophages.

(**A**) Representative images of WT and WKO macrophage's nuclei under 3 or 6 μm confinement height. Insets show nuclear deformation. On the right, representative examples of intact or damaged nuclei used for quantification. Scale bar 10 μm. (**B**) Quantification of the fraction of cells carrying a damaged nucleus after 1 h confinement at the indicated height. At least 200 cells per conditions were analysed in N=3 independent experiments. Two-tailed unpaired t-test. ***P<=0.001. (**C**) Assay on NLS-GFP cytosolic release under confinement. Representative images of NLS-GFP$^+$ macrophages after 1h of confinement at 3 μm. Scale bar 20 μm. (**D**) Corresponding quantification of nuclear ruptures upon confinement based on NLS-GFP cytosolic leakage. Each dot is an independent experiment (N=3). n=173 WT and n=102 WKO cells. Data are expressed as fold to WT mean. *P<=0.05 in a two-tailed unpaired t-test. (**E**) Individual frames from supplementary movie xx showing nuclear envelope rupture during live imaging. The timeframe is indicated by f; yellow arrow highlights bleb formation and subsequent rupture in WKO. (**F**) WKO cells transduced with WASP-GFP were subjected to confinement. Representative images of WASp-GFP$^+$ and WASp-GFP$^-$ macrophages after 1 h confinement at 3 μm. Dashed lines mark nuclear boundaries. Scale bar 20 μm; 5 μm for insets. (**G**) Corresponding quantification of nuclear blebbing in control or WASp-GFP reconstituted cells. Data are expressed as fold on WKO ctrl, n=83 WT and n=99 WKO cells from N=3 independent experiments, one sample t-test. **P<=0.01. (**H**). Representative images of confined (3μm) WASp-GFP$^+$ and WASp-GFP$^-$ macrophages stained for LMNA/C (yellow). Nuclear insets show LMNA/C and Hoechst signal in WKO ctrl (1) and WKO WASp-GFP (2). Scale bar 20 μm. (**I**) Quantification of LMNA/C

rupture events in WKO ctrl and WKO WASp-GFP based on **H**. Data are from n=135 WKO ctrl and n=22

WKO WASp-GFP in *N*=2 experiments.

Supplementary Figure 4 – WASp limits nuclear rupture under confinement. (**A**) Cells were

subjected to vertical confinement, as schematically represented. (**B**) Nuclear height quantification in

2D without confinement in BMDM. Data from *N*=1 experiment with a total of 10 z-stacks nuclei

measured for genotype. (**C**) Representative images of BMDM nuclear height from live-imaging

acquisition of cell under confinement at 6 µm and 3 µm. Vertical scale bar 6 µm, with a tick every 1

µm. (**D**) WKO BMDM were transduced with lentiviral particles encoding WASp-GFP. (**E**) Flow

cytometry shows the fraction of WKO cells expressing WASp-GFP to visualize WASp-GFP expression.

(**F**) Representatives immunoblot of WASp expression in WT, WKO ctrl and WKO WASp-GFP

macrophages after 7 days of differentiation.

Figure 5 – WASp triggers perinuclear actin polymerization under confinement. (**A**) Representative

images of actin distribution in WT and WKO BMDM confined at 3 µm height. Scale bars 20 µm. (**B**)

Quantification of the fraction of cells with central actin accumulation. n=number of cells, as indicated

in figure. *N*=2 experiments for unconfined controls and *N*=4 for confined samples. Two-tailed

unpaired t-test. **$P<=0.01$. (**C**) Representative images to define patterns of actin distribution in

macrophages confined at 3µm. Scale bar 5 µm. (**D**) Corresponding quantification of the fraction of

WT and WKO macrophages in each category. n=number of cells, as indicated in figure; from *N*=3

individual experiments. (**E**) WKO cells transduced with WASp-GFP were confined at 3 µm and

analysed to visualize the relative distribution of WASp and F-actin. Scale bar 5 µm. Line intensity

profile of selected region showing the intensity distribution of WASp-GFP and phalloidin (bottom).

(**F**) Quantification of WASp-GFP and F-actin Mander's coefficient of colocalization on perinuclear

actin patches. Data are from **E**, with a total of *n*=20 cells analysed from *N*=2 independent

experiments.

Supplementary Figure 5 – WASp perinuclear localization under confinement.

(**A**) WASp-GFP expressing cells were recorded during migration in 5 x 6 μm microchannels. (**B**) After fixation at the end of the recording period, WASp distribution was calculated by dividing the cell volume into three equal parts (top left). Average intensity map of WASp-GFP was obtained by normalizing and merging different cells from a fixed end time point analysis (bottom left). Data are from *n*=42 cells obtained in *N*=2 independent experiments. Representative line intensity profile of WASp-GFP signal (right).

Figure 6 – Mechanical confinement triggers innate activation in WASp KO macrophages. (**A**) BMDM were subjected to 1 h confinement at 3 μm or kept in unconfined conditions and processed for RNA-seq and gene qPCR expression analysis. (**B**) Venn diagram of common and unique upregulated DEGs in WT and WKO BMDM confined at 3 μm as compared to their respective unconfined controls. Relevant processes and genes are highlighted. (**C**) Log2FC of commonly upregulated inflammatory genes by confinement in the two genotypes. (**D**) Z-score of branched actin signature across the two genotypes in unconfined and confined conditions (left). Normalized expression of relevant individual genes is showed as heatmap (right). (**E**) Z-score of nuclear mechanic signature across the two genotypes in unconfined and confined conditions (left). Normalized expression of relevant individual genes is showed as heatmap (right). (**F**) Z-score of NER induced inflammatory signature across the two genotypes in unconfined and confined conditions. *P* in **D**, **E**, and **F** were calculated by one-way ANOVA test. *P<=0.05, **P<=0.01, **P<=0.001. (**G**) Quantification of Il-6 mRNA in confined WT and WKO BMDM after 1 h at indicated heights. *n*=6 for unconfined and 3 μm confined samples, *n*=4 for 6 μm confined samples. **P<=0.01 on a one-way ANOVA test. (**H**) Normalized expression of Il-6 mRNA in WT ctrl and WT WASp KO BMDM. Data from 6 individual confinement replicates. (**I**) Normalized expression of Il-6 mRNA in WKO ctrl and WKO STING KO BMDM from 4 individual confinement experiments. *P<=0.05 on a one-sample t-test. (**J**) Model of WT and WKO macrophage response to confinement.

Supplementary Figure 6 – Mechanical confinement triggers innate activation in WASp KO macrophages. (**A**) Venn diagram of differentially expressed genes between WT and WKO in unconfined resting condition. Relevant genes and processes are highlighted. (**B**) PCA analysis of BMDM unconfined controls and 3 μm confined samples from RNA-seq. (**C**) Venn diagram of DEGs

downregulated by confinement in WT and WKO at 3 μm. Relevant genes are highlighted. (**D**) Z-score

of Rac2 signature across the two genotypes in unconfined and confined conditions (left). Normalized

expression of Sun2/RhoA/ROCK genes is showed as heatmap (right). (**E**) Representative western blot

of KO efficiency control in WT macrophages after 72 h from sgCtrl or sgWASp CRISPR-Cas9

electroporation. (**F**) Representative western blot for KO efficiency in WKO macrophages after sgCtrl

or sgSTING CRISPR-Cas9. (**G**) Il6 quantification in WKO and WKO STING KO at steady state and upon

direct STING stimulation by 3 μg/mL of DMXAA.

1142

1143

Figure 1

Suppl. Figure 1

A

B

C

D

E

Figure 2

Suppl. Figure 2

Figure 3

A

| | 8 um | 6 um | 5 um | 4 um | 3 um | 2 um |

WT

WKO

Nuclear mask

Direction of migration →

B

Entrance rate (%)

WT / WKO

8um 6um 5um 4um 3um 2um

C

Nuclear herniation 6µm

Representative nuclear masks

WT

WKO

Direction of migration →

Herniation (% of total)

WT WKO

D

Herniation (% of total)

ns ns

WT / WKO

Front Back

E

NLS-GFP

16h migration + fixation and staining

F

WT WKO

Time

WT WKO

Time

Herniation

Rupture

G

Nuclear rupture 6um

NE rupture (%)

*

WT WKO

H

WT / WKO

Multiple rupture events (% of ruptures)

WT WKO

Suppl. Figure 3

A

Raw image

50um

Nuclear mask

B

C

D

Figure 4

A
WT
Intact
WKO
Damaged

B
WT WKO
6μm
3μm

6 μm
ns
Blebbing nuclei (% of tot)
WT WKO

3 μm
***
Blebbing nuclei (% of tot)
WT WKO

C
NLS-GFP Hoechst Phalloidin Merge
WT
WKO

D
3 um
*
NE rupture (fold to WT mean)
WT WKO

E
NLS-GFP live imaging
f=0 f=30 f=90 f=150
WT
WKO

F
Phalloidin Hoechst GFP
WKO ctrl
WKO WASp-GFP

G
3 um
**
Blebbing nuclei (fold to WKO ctrl)
WASp-GFP: - +

H
GFP Lamin A/C
Phalloidin Hoechst
LMNA/C Hoechst
1
2

I
Lamin A/C:
Intact
Ruptured
Fraction of cells
31% 21%
WASp-GFP: - +

Suppl. Figure 4

Figure 5

A 3um

WT
Merge | Phalloidin | Hoechst | Inset

WKO
Merge | Phalloidin | Hoechst | Inset

B

Central actin patches

Percent of cells

*** ***

n=155 n=126 n=315 n=387
WT WKO WT WKO
nc   3 um

C 3 µm

Phalloidin | Hoechst | Merge | Phalloidin LUT

Perinuclear
Nuclear
Ring
Diffuse

D

Actin distribution
- Perinuclear patch
- Nuclear accumulation
- Filamentous ring
- Diffuse cytosolic

Fraction of cells

WT
n=52

WKO
n=58

E 3 µm

WASp-GFP | Phalloidin | Hoechst | Merge

F

Mander's coefficient

n=20 cells

WASp-GFP
Phalloidin

WASp-GFP-phalloidin coloc

Distance (um)

Suppl. Figure 5

A

B

WASp distribution

Back   Centre   Front

5 um   Avg WASp intensity

Back   Direction of migration →   Front

Figure 6

A unconfined · 3 μm confined

1h confinement → RNA extraction → RT-qPCR + RNA-seq

B Upregulated by confinement

WT — 36, 130, 151 — WKO

Chemotaxis: Ccl5, Tnfsf14

Cell activation: CD40, CD86, Il12b, Nod2

Tissue remodelling: Vega, Suco, Cav1, Ptger

Stress-activated MAPK cascade: Per1, Dusp10, Fas,, Il1a

Acute inflammatory response: Il1b, Ptgs2, Cxcl1, Nlrp3, Tnf

Response to chemokines: Cxcr4, Cxcl2, Cxcl10, Cxcl3

C Common inflammatory genes

Log2FC confined vs unconfined — Cytokines (Cxcl3, Cxcl2, Cxcl10, Il1b, Il1a), Cell activation (Tnf, Tnfsf9, Nfkbia, Nr4a3, Nlrp3) — WT, WKO

D Branched actin score — Z-score — WT, WT 3um, WKO, WKO 3um — Actr2, Actr3, Arpc4, Was

E Nuclear mechanics score — Z-score — WT, WT 3um, WKO, WKO 3um — Emd, Lemd2, Lemd3, Lmna, Plec, Sun1, Tmpo

F NER inflammatory score — Z-score — WT, WT 3um, WKO, WKO 3um

G Il6 mRNA (relative) — nc, 6, 3 WT; nc, 6, 3 WKO

H 3 um — Il-6 mRNA (fold to Ctrl) — p=0.12 — WT: Ctrl, Was ko

I 3 um — Il-6 mRNA (fold to Ctrl) — WKO: Ctrl, Sting ko

J WT — Actin patch, Nuclear stability. WKO — Nuclear rupture, cGAS-STING activation

Suppl. Figure 6

A

Steady state WKO vs WT

Immunoglobulin variable regions
Igkv4-59, Igkv4-61, Igkv14-2

Extracellular matrix organization
Ptx3, Col2a1, Adamts9, Hspg2

Vascular remodelling
Flt4, Adam12, Aldh1a2

Cellular activation/migration
MS4A4A, Ephb2, Lilra5, Cxcl1

12598 Unmodulated

253 Upregulated   66 Downregulated

Cell metabolism
Gpi, Pfkl

Prostaglandin metabolic process
Pdpn, Sphk1, Cyp2s1

B

group
WT_ctrl   WT_conf
WKO_ctrl   WKO_conf

PC2: 23% variance
PC1: 59% variance

C

Downregulated by confinement

85   12   91

Extracellular matrix
Col3a1, Col5a1, Ccn2

Metabolism and mitochondria
Aldh1a1, Afmid, Diablo, Timm10b

D

Rac2 score

Z-score

WT   WT 3um   WKO   WKO 3um

*   ***

RhoA/ROCK genes
WT   WKO   WT 3um   WKO 3um
Sun2
Rhoa
Rock1
Rock2

E

WT

sgRNA: Ctrl   WASp

WASp

STING

αTUB

WASp ratio: 1   0.17

F

WKO

sgRNA: Ctrl   STING

WASp

STING

αTUB

STING ratio: 1   0.13

G

WKO ctrl
WKO STING KO

Il6 mRNA (fold to WKO DMSO mean)

Sting KO: -   +   -   +
DMSO   DMXAA