1 **FRONT MATTER**
2
3 **Title**
4 Main title: Emergence of Pre-movement Beta Activity with Stable Sensorimotor Predictions to
5 Facilitate Motor Adjustments
6 Short title: Beta Oscillations and Sensorimotor Adaptation
7
8
9 **Authors**

10 Martina Bracco,[1,2]* Varsha Vasudevan,[1] Vridhi Rohira,[1] Quentin Welniarz ,[1] Mihoby
11 Razafinimanana,[3] Alienor Richard,[1] Christophe Gitton,[4] Sabine Meunier,[1] Antoni Valero-
12 Cabré,[2,5,6] Denis Schwartz,[7]† Traian Popa,[8,9]† and Cécile Gallea[1]*†
13
14
15 **Affiliations**
1. Sorbonne Université, Institut du Cerveau - Paris Brain Institute - ICM, Mov'it Team, Inserm, CNRS, APHP, Hôpital de la Pitié Salpêtrière, Paris, France
2. Sorbonne Université, Institut du Cerveau - Paris Brain Institute - ICM, Dynamics, Plasticity and Rehabilitation Group, Frontlab Team, Inserm, CNRS, APHP, Hôpital de la Pitié Salpêtrière, Paris, France
3. Sorbonne Université, CNRS, Inserm, IBPS, Neurosciences Paris Seine, Paris, France
4. Sorbonne Université, Institut du Cerveau - Paris Brain Institute - ICM, CENIR, Inserm, CNRS, APHP, Hôpital de la Pitié Salpêtrière, Paris, France
5. Cognitive Neuroscience and Information Tech. Research Program, Open University of Catalonia (UOC), Barcelona, Spain
6. Dept. Anatomy and Neurobiology, Laboratory of Cerebral Dynamics, Boston University School of Medicine, Boston, USA
7. CERMEP-Imagerie du Vivant, MEG Departement, Lyon, France
8. Lausanne University Hospital (CHUV), Department of Clinical Neurosciences, Lausanne, Switzerland
9. Lake Lucerne Institute (LLUI), Vitznau, Switzerland

* co-corresponding authors: martina.bracco@icm-institute.org, cecile.gallea.icm@gmail.com
† These authors contributed equally to this work.

Abstract

Daily, we adapt to perturbations to maintain efficient performance, especially when transitioning between environments with different features. Such an adaptative process relies on updating motor programs based on past errors, a process associated with beta oscillations (13-35 Hz). Yet, the spatio-temporal dynamics underlying the transition between different environments, and the updating of motor programs in response to specific environmental perturbations remain elusive. Here, we show for the first time that pre-movement beta activity in a cerebello-cortical network emerges as the features of a new environment and motor outcomes become more stable and predictable. Notably, the cerebellum and parietal cortex drive prefrontal activity near movement initiation to update these predictions. Using a single-trial approach, we reveal that pre-movement beta bursts predict the evolution of trial-by-trial motor adjustments within a stable environment. These findings demonstrate that pre-movement beta activity within cerebellar-cortical network encodes essential information for achieving efficient sensorimotor predictions following environmental changes.

Teaser

Pre-movement beta activity in cerebello-cortical network predicts how efficiently we adapt our movements to environmental changes based on prior experience.

59 **MAIN TEXT**
60
61 **Introduction**
62
Successful adaptation to environmental perturbations requires rapid corrections to maintain
optimal performance. This is the case in tennis when players adjust their forehands to compensate
for a headwind. This adaptive process requires detecting changes in the environment and updating
motor commands accordingly. Such updates depend on the formation of internal models, essentially
sensorimotor predictions informed by prior experience of the environment. These internal models
are based on *prediction errors,* arising when expected events fail to occur, and on an *efference copy*,
a copy of the motor command used to predict the expected sensory outcomes of the action. Over
the course of motor adaptation, prediction errors are largest early on, when the environment's
characteristics are unknown, and gradually decrease as the brain integrates the environmental
changes. Trial-by-trial, the brain uses past errors to update its sensorimotor predictions,
progressively consisting of a more predictable representation of the environment and, thus,
promoting motor adjustments in response to these environmental changes (*1*). Brain and behavior,
therefore, continuously oscillate between *stable* states, characterized by small kinematic
fluctuations, and *transition* states, marked by large fluctuations in response to perturbation before
returning to a stable state. In this framework, brain and behavior states *emerge* from the interaction
between the body's biomechanics, environmental constraints and task goal (*2*). But how does the
brain process state-dependent information to transition from an unknown to a stable and predictable
environment, hence supporting behavioral adaptation?
Beta oscillations (13-35Hz) are known to play a crucial role in adaptation processes (*3–9*).
In sensorimotor adaptation, pre-movement beta synchronization is modulated by prior errors (*10–*
12). Specifically, up to two seconds before movement onset, beta synchronization decreases
following erroneous movements caused by unexpected perturbations and increases after correct
movements (*10, 11*). These findings suggest that pre-movement beta oscillations encode high-level
sensorimotor predictions, integrating past performance and supporting motor adjustments when
needed (*13, 14*). Yet, the precise spatio-temporal dynamics of pre-movement beta oscillations
during adaptive behavior remain largely unexplored.
Beta oscillations are ubiquitous in the brain and have been identified in several areas,
including the motor, parietal, and frontal cortices, as well as the cerebellum and the basal ganglia
(*15–25*). Such a widespread distribution suggests the role of beta oscillation in potentially
facilitating the flow of sensorimotor predictions across different brain areas (*26*). In the context of
motor adaptation, the cerebellum and the parietal cortex have been characterized as critical for
predictive processes (*27*). The cerebellum acts as a 'comparator' between the predicted sensory
feedback derived from the efference copy and the actual sensory feedback, to adjust motor plans in
the next trial (*28–30*). Furthermore, recent work has shown that cerebellar activity increases well
before movement execution (*31–33*), suggesting an involvement in motor preparation and planning
(*34, 35*). Like the cerebellum, the parietal cortex is active at the early stages of motor planning,
providing information to compensate for motor and target errors (*19, 36*). The parietal cortex acts
as an integrator of somatosensory information and maintains predictions of the sensory
consequences of movements (*37*). Additionally, it supplies the prefrontal cortex with essential
contextual information for goal-generation (*38*). This suggests that the parietal cortex, and possibly
the cerebellum, may communicate with frontal and prefrontal areas to encode sensory predictions
used to update motor plans during the pre-movement period (*39, 40*).
Here, we aimed to characterize the brain network dynamics that mediate adaptive behavior
through pre-movement beta oscillations. Specifically, we sought to determine how sensorimotor
predictions and the emergence of specific brain activity patterns evolve over time to facilitate
progressive motor adjustments. To this aim, we recorded magnetoencephalography (MEG) from

healthy human participants as they performed blocks of 50 consecutive trials of a reaching task
under different environmental characteristics; i.e., either normal (no rotation) or rotated visual
feedback (rotation). Using a joystick, participants controlled the position of a cursor on a screen. In
'normal' visual feedback blocks, the cursor's trajectory matched the joystick position, resulting in
minimal trajectory errors and stable movement outcomes. In contrast, rotated visual feedback
introduced a perceptual bias, leading to significant trajectory errors during early trials. These errors
decreased and stabilized over the 50 trials as participants adapted to the perturbation (*41*, *42*).
Our analysis involved two complementary approaches. First, we conducted conventional
data-driven analyses of average pre-movement beta oscillations to identify their global evolution
and spatial distribution, across different states (transition *vs.* stable) and environmental
perturbations (no rotation *vs.* rotation). Dynamic effective connectivity analysis during the pre-
movement period further revealed the spatio-temporal features of information processing before
target appearance, i.e., before movement planning and efference copy generation had occurred.
Conventional average analyses on the functional interactions of brain networks have enabled
researchers to pinpoint statistically significant neural "puzzle pieces". While these analyses provide
insight into the functional role that a specific network may play, they do not reveal how this process
unfolds through time, nor do they clarify how the coordinated interactions among a network of
brain regions translate into behavior (*43*). This is particularly important for sensorimotor adaptation,
where trial-by-trial variations of beta activity could reveal important dynamics as individual trials
carry critical information for progressively improving motor performance. Furthermore, single-trial
analysis approaches have recently shown beta oscillations within cortical and subcortical brain
areas emerge as brief events or bursts during the pre-movement period (*22*, *44–48*). For example,
in single trials, beta bursts before the onset of a simple movement correlate with the kinematics
during its execution (*44*) and predict movement initiation timing in decision tasks guided by visual
cues (*45*, *48*). On such a basis, we integrated the conventional analysis with a single-trial analysis
to determine whether pre-movement beta bursts carry information that supports motor adjustments
on a trial-by-trial basis. We used a predictive model that simultaneously captured the influence of
previous errors on current motor performance and the effect of beta bursts on subsequent motor
performance.
138
139 **Results**
140
141 ***Pre-movement beta activity supports the update and maintenance of sensorimotor predictions***
142
143 *Behavior*
144
145   Participants performed blocks of 50 consecutive trials of a reaching task with different
environmental characteristics, *i.e.*, either with normal (0°; no rotation) or rotated visual feedback
(25°, -30°; rotation; Figure 1). These blocks were randomly interleaved across participants,
inducing a transition from one environment to another every 50 trials. This transition ensured that
while participants' performance stabilized within a block, the alternation between different
environments hindered their ability to consolidate adaptive performance across blocks (*43*), and
induced a sudden switch between environments with different characteristics. This distinction is
crucial for dissociating the effect of transition between environments from simple error adjustments
over trials.
The goal of the task was to reach a peripheral target, as fast as possible. Starting from a
central position, participants controlled a cursor using a joystick with the right dominant hand.
Participants' motor performance was quantified by estimating trajectory errors on each trial. This
was quantified as the area under the curve (AUC) representing the cursor's trajectory deviating
from an ideal straight line between the starting and the final target position (higher AUC values
indicate greater errors). To characterize changes in participants' performance, we divided each

block with the same environmental characteristics (*Perturbation*) into two *States*: the transition state (first 16 trials), in which participants showed an overall 80% performance improvement, and a stable state (last 16 trials) in which performance stabilized (figure 2A). This division was based on the observed population's errors within blocks with rotated feedback (see Methods for details).

As expected, errors decreased significantly from the transition to the stable state (2x2 repeated measures ANOVA; Main Effect of *State* (transition *vs.* stable): $F(1,16) = 560.27$, $p < 0.001$; $\eta_p^2 = 0.97$). However, participants consistently made more errors in blocks with rotation compared to those with no rotation (2x2 repeated measures ANOVA; Main Effect of *Perturbation* (rotation *vs.* no rotation): $F(1,16) = 155.56$, $p < 0.001$; $\eta_p^2 = 0.67$). This difference persisted across all states, with substantially more errors in stable ones (no rotation *vs.* rotation, transition state: $t = -15.66$, $p < 0.001$, Cohen's $d = -4.57$; no rotation *vs.* rotation, stable state: $t = -7.13$, $p < 0.001$, Cohen's $d = -2.08$). Notably, the reduction in errors was mainly observed in the environment when rotation was applied (2x2 repeated measures ANOVA; *State* x *Perturbation* Interaction: $F(1,16) = 110.536$, $p < 0.001$; $\eta_p^2 = 0.09$; transition *vs.* stable, no rotation: $t = 2.28$, $p = 0.19$, Cohen's $d = 0.31$; transition *vs.* stable, rotation: $t = 20.67$, $p < 0.001$, Cohen's $d = 2.80$; Figure 2B).

Averaged pre-movement beta oscillatory activity (MEG): sensor and source space

We recorded participants' brain activity during the reaching task to determine the spatio-temporal changes of pre-movement beta synchronization (14-26 Hz) in relation to changes across different states and environmental perturbations. To achieve this, for each participant, we transformed the MEG data into a time-frequency domain, averaging across trials to obtain separate event-related, average, power spectra corresponding to *State* (transition and stable) and *Perturbation* (rotation and no rotation; Figure 3), as we did for the behavioral analysis. Individual data were entered into a 2x2 non-parametric cluster-based approach and paired t-tests.

We found a significant overall increase in pre-movement beta synchronization from transition to stable states (2.5-1.35 s before movement onset; cluster analysis; Main Effect of *State* (transition *vs.* stable): $p < 0.001$) in both perturbation conditions (rotation (transition *vs.* stable): $p = 0.009$; no rotation (transition *vs.* stable): $p = 0.003$; Figure 4A). Source localization revealed that this change primarily originated from the cerebellum and temporal cortices bilaterally (Figure 4A, Table S1). Interestingly, when passing from a transition to a stable state, pre-movement beta synchronization showed a smaller increase during rotation than in no rotation blocks (2.35-2 s before movement onset; cluster analysis; *State* x *Perturbation* Interaction: $p = 0.014$; Figure 4B). This interaction between states and perturbations was not attributable to a difference in beta power during the transition state (cluster analysis; transition (rotation *vs.* no rotation); all p values > 0.229) or to an overall difference across perturbations (rotation *vs.* no rotation; all p values > 0.073). The source estimates revealed that this interaction originated from a network of regions including the left parietal cortex, the right frontal and prefrontal cortices, and the right cerebellum (Figure 4B, Table S1).

Dynamics of pre-movement beta connectivity (MEG)

We performed an effective dynamic connectivity analysis in the beta range (14-26 Hz) during the 2.5 s preceding movement initiation. We computed connectivity using nonparametric conditional Granger causality during the transition and stable states, across both perturbations (rotation, no rotation). We specifically focused on those brain areas highlighted in our previous source space analysis and known to be relevant for sensorimotor adaptation and prediction processes: the posterior right cerebellum (rCB), the left parietal cortex (lPC), and the right prefrontal cortex (rPFC; (*35*)). To assess the temporal dynamics of such effective connectivity patterns, we examined five overlapping 0.45 s time windows prior to target appearance (from -2.5

to -2.05 s; from -2.15 to -1.7 s; from -1.8 to -1.35 s; from -1.45 to -0.9 s; from -1.0 to -0.65 s),
capturing at least six cycles of beta oscillation at its lowest frequency (14 Hz).
Our results showed that, like beta power, connectivity increased from transition to stable
states within the same block. This increase was reciprocal among the right cerebellum, the left
parietal cortex, and the right prefrontal cortex (5x2x2 ANOVA (*Time Window x State x*
Perturbation), Main Effect of *State*: Cerebellum to Prefrontal, $F(1,16) = 6.57$, $p = 0.02$, $\eta_p^2 = 0.29$;
Cerebellum to Parietal, $F(1,16) = 3.88$, $p = 0.06$, $\eta_p^2 = 0.19$; Parietal to Cerebellum, $F(1,16) = 5.53$,
$p = 0.03$, $\eta_p^2 = 0.26$; Parietal to Prefrontal, $F(1,16) = 11.97$, $p = 0.003$, $\eta_p^2 = 0.43$; Prefrontal to
Cerebellum, $F(1,16) = 6.61$, $p = 0.02$, $\eta_p^2 = 0.29$; Prefrontal to Parietal, $F(1,16) = 5.36$, $p = 0.03$,
$\eta_p^2 = 0.25$; Figure 5A). However, an interesting pattern emerged in the five time windows
considered in the pre-movement period. As the time window approached the movement initiation,
connectivity specifically increased from both the cerebellum (5x2x2 ANOVA, Main Effect of *Time*
Window: $F(2.28,36.54) = 4.32$, $p = 0.02$, $\eta_p^2 = 0.20$, Huynh-Feldt sphericity correction; -2.5/-2.05
vs. -1.45/-0.9: $t = -3.51$, $p = 0.008$, Cohen's $d = -0.36$; -2.5/-2.05 *vs.* -1.0/-0.65: $t = -3.03$, $p = 0.04$,
Cohen's $d = -0.31$; Figure 5B) and the parietal cortex (5x2x2 ANOVA, Main Effect of *Time*
Window: $F(2.65,42.36) = 3.16$, $p = 0.04$, $\eta_p^2 = 0.16$, Huynh-Feldt sphericity correction; -2.5/-2.05
vs. -1.0/-0.65: $t = -3.07$, $p = 0.02$, Cohen's $d = -0.31$; Figure 5B) to the prefrontal cortex. Notably,
this increase was not bidirectional (5x2x2 ANOVA, Main Effect of *Time Window*, Prefrontal to
Cerebellum: $F(2.58,41.35) = 1.15$, $p = 0.34$, $\eta_p^2 = 0.07$; Prefrontal to Parietal: $F(1.55,24.75) = 0.87$,
$p = 0.40$, $\eta_p^2 = 0.05$). There was no significant effect of *Perturbation* (5x2x2 ANOVA, Main Effect
of *Perturbation* Cerebellum to Prefrontal: $F(1,16) = 2.21$, $p = 0.16$, $\eta_p^2 = 0.12$; Parietal to Prefrontal:
$F(1,16) = 0.86$, $p = 0.37$, $\eta_p^2 = 0.05$), indicating that the increase in connectivity from cerebellar
and parietal areas to the prefrontal cortex was independent of the error magnitude.

Trial-by-trial beta burst dynamics encode motor adjustments over consecutive trials of the same
environment

Beta burst characterization

We sought to determine the trial-by-trial relationship between pre-movement beta bursts
and trajectory errors in our task. To this aim, we isolated beta bursts in each trial of each block
(Figure 6A). Beta bursts were defined as local maxima power exceeding a set power cut-off in the
2.5-second pre-movement epoch, at frequencies ranging from 14 to 26 Hz, and from sensors
included in the significant clusters from the average beta power analysis (*State* x *Perturbation*
Interaction; sensors shown in Figure 4B). The power cut-off was fixed to 3 times the median power,
which provided the strongest correlation between the average mean power and the area covered by
beta bursts (3X; all r values > 0.96; all p values < 0.001 Figure 6B), thus best representing the
average beta power. This cut-off allowed us to detect ~7 bursts per trial, each lasting less than 150
ms (Figure 6C).
Among potentially quantifiable beta features relevant to behavior (*i.e.*, bursts mean power,
burst number, last burst power; (*49*)), the mean power of beta bursts showed the highest correlation
with the average beta power (repeated measures ANOVA; Main effect of Beta *Feature* (beta bursts
mean power *vs.* bursts number *vs.* last burst power): $F(2,32) = 481.30$, $p < 0.001$; $\eta_p^2 = 0.97$; bursts
mean power *vs.* bursts number: $t = 5.04$, $p < 0.001$, Cohen's $d = 1.54$; bursts mean power *vs.* last
bursts power: $t = 29.03$, $p < 0.001$, Cohen's $d = 8.84$; bursts number *vs.* last bursts power: $t = 23.99$,
$p < 0.001$, Cohen's $d = 7.31$; Figure 6D). Additionally, the power of beta bursts contributed
significantly to error reduction along the different states (transition *vs.* stable), compared to
background beta activity (3x2x2 repeated measures ANOVA; *Feature* x *States* interaction: $F(1,16)$
$= 13.14$, $p = 0.002$, $\eta_p^2 = 0.45$; Figure 6E). More specifically, the power of the beta bursts increased
across states (beta bursts mean power; transition *vs.* stable state: $t = -4.96$, $p < 0.001$, Cohen's $d =$
-1.53), reflecting the average beta power trends, while the background activity remained stable

(background beta activity; transition *vs*. stable state: t = 0.48, p = 1.00, Cohen's d = 0.11).
Altogether these results support the use of beta burst power as a potentially relevant variable to
model the relationship between beta activity and motor adjustments over trials.
260
Trial-by-trial relationship between beta bursts and trajectory errors: predictive model

To investigate whether beta burst dynamics during the pre-movement period could predict
trajectory error across consecutive trials, we used a time-series modeling approach. Specifically,
we employed a regression framework, the Auto Regressive Moving Average (regARMA),
modeling the dynamics of trajectory errors over consecutive trials, and including the trial-wise beta
burst power as an external predictor. The optimal model was selected based on the lowest Akaike
Information Criteria (AIC) values, tested across all feasible model combinations on population-
averaged data from the 17 participants, as detailed in the Methods section. The regARMA (1,1)
model with a Gaussian distribution and no intercept was identified as the best-fitting model for all
perturbations (Table S2).
We validated the selected regARMA (1,1) model using a leave-one-out prediction approach.
For each participant, model parameters were estimated using the average data from the other 16
participants. These parameters were then used to predict the left-out participant's error evolution
across trials. Specifically, participant's error in the first trial and the trial-by-trial beta burst
evolution over 50 trials served to predict the error across the remaining 49 trials (Figures 7A; see
Figures S1, S2, and S3 for individual simulations). This enabled the assessment of the model's
ability to generalize and accurately capture the relationship between beta burst evolution and motor
performance across individual participants.
We then used the coefficient of the regARMA model's external predictor as a measure of
beta burst power contribution to the motor adjustments over trials. Beta contribution defines the
strength of the influence that pre-movement beta bursts have on motor adjustments, with higher
values indicating a stronger role of beta oscillations in facilitating the adjustment of motor
performance and reducing errors across trials. Our analysis revealed that the contribution of beta
burst power varied across perturbations (repeated measures ANOVA, Main Effect of *Perturbation*
(0° *vs.* 25° *vs.* -30°): F(1.4,22.55) = 6244.68, p < 0.001, η_p^2= 0.99, Huynh-Feldt sphericity
correction; Figure 7B). Specifically, the beta contribution increased with the increase of degrees of
rotation (0° *vs.* 25°: t = -73.55, p < 0.001, Cohen's d = -20.20; 0° *vs.* 30°: t = -109.64, p < 0.001,
Cohen's d = -30.10; 25° *vs.* 30°: t = -36.09, p < 0.001, Cohen's d = -9.91; Figure 7B). Importantly,
the model performed consistently across all perturbations, as indicated by non-significant
differences in model prediction error (normalized mean squared error, N-PMSE; repeated measures
ANOVA; Main Effect of *Perturbation* (0° *vs.* 25° *vs.* -30°): F (2,32) = 2.50, p = 0.10, η_p^2= 0.14;
Figure 7C). This suggests that the difference in beta contribution between different perturbations
was not due to overall differences in the models' generalizability.
294
Model specificity

We also evaluated whether the relationship between MEG activity and motor performance
was specific to the beta frequency. To this aim, we tested the same regARMA model with an
external predictor derived from an alternative frequency. We specifically selected alpha (8-12 Hz)
as it had also shown pre-movement synchronization. However, unlike beta, alpha synchronization
exhibited a less robust burst-like behavior, with less than 2 bursts per trial on average (Figure S4A,
B, C). As a consequence, we characterized alpha dynamics across trials with the mean alpha power.
This trial-wise averaged alpha was tested for its predictability of trajectory error evolution using
the regARMA model. The optimal model selection procedure was repeated using the same steps
carried out for beta bursts, which led to the choice of regARMA(1,1) as the one with the least AIC
value for each perturbation (Table S3). Further, we tested the model using population leave-one-

out cross-validation and compared the prediction errors of these models using alpha and beta bursts as external predictors. Here, we found that the model prediction error was consistently larger for the model using alpha mean power independent of the induced perturbation (N-PMSE; 2x2 repeated measures ANOVA; Main Effect of Model (beta bursts *vs.* mean alpha): $F_{(2,32)} = 5.04$, $p = 0.039$, $\eta_p^2 = 0.24$, Figure S4D). Because we were forced to use for each trial mean alpha power (rather than alpha bursts), we performed a control analysis replacing beta bursts with average beta power in each trial and found that the results remained similar (N-PMSE; 2x2 repeated measures ANOVA; Model (mean beta *vs.* mean alpha): $F_{(2,32)} = 6.84$, $p = 0.019$, $\eta_p^2 = 0.30$, Figure S4E).

Discussion

Our findings offer a comprehensive view of how pre-movement beta activity emerges from the transition to stable states, enabling adaptation to varying environmental perturbations. By integrating both conventional and dynamic analytical approaches, we provide novel insights into the physiological relevance of pre-movement beta activity on at least three levels. Anatomically, by identifying cerebello-cortical networks as key in this process; connectivity-wise, by showing directional and dynamical network interactions mediated by beta oscillations from cerebellar and parietal to prefrontal areas; and behaviorally, by pinpointing the trial-by-trial contribution of pre-movement beta burst activity to motor adjustments.

Using a conventional analysis of averaged beta activity, we first showed that state-dependent sensorimotor predictions impacted pre-movement beta activity. Pre-movement beta synchronization increased during stable states when the characteristic of environmental perturbations was known and motor errors were minimized, compared to the transition state, when a switch from one perturbation to another had just occurred. We argue that the increase in pre-movement beta synchronization occurs regardless of performance improvement based on two facts. First, beta power amplitude was equally low in the transition state for both perturbation conditions (rotation and no rotation) despite large differences in trajectory error magnitude. Second, pre-movement beta synchronization during the stable state was more pronounced in the absence of rotation (during which motor performance did not improve) compared to when rotation was applied. This modulation of pre-movement beta synchronization up to 2.5 s before movement onset aligns with previous studies using transient and unpredictable perturbations (*10, 11*) which show systematic attenuation of beta synchronization following trials with larger trajectory errors. Here, with our design consisting of consecutive trials with the same perturbation, we elegantly demonstrate that high trajectory errors are not a necessary condition for a subsequent attenuation of pre-movement beta activity. Instead, it is the switch to a yet unknown set of rules governing a specific environment, leading to lower outcome predictability. The originality of our contribution lies in the investigation of pre-movement beta synchronization when, following the switch of environment, outcome predictability is built over several trials of an environment with the same characteristics. This outcome predictability is independent of *prediction errors* per se, as the timing of pre-movement beta changes occur before target appearance when motor planning and efference copy are not yet available. We hypothesize that pre-movement beta activity encodes sensorimotor predictions based on past experience, reflecting the response of the sensorimotor system to the switch between environments with different characteristics. As participants build a stable representation of the environment and motor outcomes become predictable, beta power increases up to 2.5 seconds before movement onset, reflecting updated sensorimotor predictions informed by previous trials. Importantly, building such sensorimotor predictions is easier in an environment with minimal perturbation (no rotation) where motor performance is overall more stable.

Using MEG, we further identified for the first time two distinct brain networks associated with beta oscillations: one in detecting the environmental characteristics to optimize motor adjustments, and another subtending the interaction between the detection of environmental changes and the building of sensorimotor predictions based on motor outcome stability. The first

network included the posterior cerebellum bilaterally and the right temporal cortex. The cerebellum holds well-established credits in sensorimotor adaptation, with bilateral activations frequently observed during motor adaptation tasks (*50*). Conversely, the role of the temporal lobes in such a process remains debated (*29*). Even if not strictly necessary for the motor component of adaptation, the temporal cortex plays a significant role in the acquisition, implementation, and retrieval of action selection strategies. In this context, the concurrent involvement of the cerebellum and the temporal lobe may reflect their contribution to the formation and acquisition of sensorimotor predictions useful to motor adjustment over trials. Instead, the second network included a broader set of regions, involving this time the right posterior cerebellum, the left parietal cortex, and the right prefrontal cortex. Such asymmetry of cerebellar activity identified within the two networks may be explained by two possible factors. It may simply reflect statistical power: as participants performed the reaching task with their right hand, the right cerebellum, ipsilateral to the movement and contralateral to the motor cortices involved in task performance, was comparably more active than the left cerebellum. Alternatively, the right posterior cerebellum may be more involved in cognitive processes related to adaptation, such as in the formation of sensorimotor predictions (*50*, *51*), as compared to the left cerebellum (*52*, *53*). Interestingly, the right cerebellum has been shown to contribute to cognitive aspects of implicit motor sequence learning (*51*), its activity being modulated by environmental uncertainty. Prediction errors within the cerebellum allow building inference to account for uncertainty, as well as sensorimotor predictions on a trial-by-trial basis to estimate future motor states (*54*, *55*). We, therefore, hypothesize that this second network contributes to accumulating evidence to update the internal model based on environmental characteristics, reduce uncertainty, and stabilize motor outcomes (*56–58*). Unlike with prediction errors, increased beta synchronization may reflect the refining of sensorimotor predictions and the success (rather than failure) of expected events to occur. As these sensorimotor predictions are refined, humans can engage in metacognitive processes, evaluating errors from previous trials, and informing motor plans of the subsequent movement.

The effective connectivity further revealed how this information, supported by beta oscillations, flows in the cerebello-cortical network and unfolds under two different time scales. First, we observed a global bidirectional connectivity increase between the cerebellum, parietal, and prefrontal cortices from the transition to the stable state. Second, within a single trial, connectivity dynamically increased between the caudal regions (cerebellum, parietal) toward the prefrontal cortex in later time windows closer to motor planning. We hypothesize that these three areas coordinate operations to identify the rules governing a set environment, by operating jointly, albeit with varying degrees of specialization. While the cerebellum and the parietal cortex are frequently implicated in sensorimotor prediction leading to adaptation (*27*), the involvement of the prefrontal cortex is generally associated with strategic processes of error correction (*35*). Our results suggest that the prefrontal cortex receives cumulative information from both the cerebellum and parietal cortex, potentially aiding in the implementation of appropriate strategies to favor the stability of motor outcomes.

Our finding associating pre-movement beta activity in the cerebellum due to changes in states and perturbation characteristics has paramount implications for our field. First, for many decades the cerebellum was conceived as a mere motor structure, involved in the correction of ongoing motor activity. In recent years, studies have suggested a role in non-motor aspects of movement execution, such as motor planning and event anticipation (*31–33*). Our study further supports the role of the cerebellum in more cognitive aspects of sensorimotor adaptation (*53*). Additionally, while beta activity within the cerebellum has been shown before (*15*, *19*, *21*), we bring an important brick of knowledge on how this signal is timely used to progressively improve motor performance. Finally, localizing cerebellar sources has long remained controversial (*59*, *60*). Our MEG data repeatedly converge with independent data-driven analyses, reinforcing the reliability of our findings. This was likely facilitated by careful positioning of participants' heads

to optimize sensor caudal coverage, the use of individualized head models including the cerebellum, and time-frequency analyses capturing non-phase-locked activity, thus mitigating signal cancellation due to the complexity of cerebellar anatomy (*60*).

While the results from averaged beta synchronization highlighted the functional role of the cerebello-cortical network in generating sensorimotor predictions tied to environmental characteristics and motor outcome stability, trial-by-trial analyses offered complementary insights into adaptive behavior. The evolution of beta burst suggests the spontaneous emergence of order from what initially appears as a disorganized system in the early trials. According to complex system theory, the brain can change and learn from experience through emergent activity, with brain states and behavior connected by non-linear relationships (*2, 61*). Specifically, beta bursts likely originating from the collective interaction of brain components within the cerebello-cortical network may support the emergence of a stable state. Through predictive modeling of trial-by-trial changes in both motor performance and beta burst power, we provided, for the first time, evidence for the non-linear relationship between pre-movement beta bursts and motor adjustments. In the case of small perturbation (no rotation), beta burst amplitude explained the stability of behaviour. In the case of large perturbation (rotation), beta burst amplitude contributed to explaining the drastic changes in behavior. Indeed, the brain-behavior relationship was strongest in blocks with larger trajectory errors (e.g., rotation; 25°–30°) compared to conditions with more stable error amplitudes (e.g., no rotation; 0°). These observations are not in contrast with the overall increase in pre-movement beta synchronization in blocks without perturbation. Instead, they suggest that pre-movement beta bursts encode information critical for engaging trial-specific motor adjustments based on past experience, highlighting their role in encoding and updating sensorimotor predictions. Trial-by-trial, this iterative process the emergence of beta supports the stabilization and or improvement of motor outcomes in specific contexts. Although speculative, we propose that beta bursts may reflect the activity of an expanding pool of neurons responding more synchronously to increase the system stability, recruiting a set of structures within the network to fine-tune motor plans. Alternatively, the progressive increase in beta burst amplitude could be related to action inhibition: as participants learn the rules governing the environment trial by trial, their knowledge of what to do may lead to increased eagerness to initiate the movement. However, we consider this unlikely, as action initiation is not linked to the mean beta burst power over a large time window (up to 2.5 seconds before movement onset). Instead, it is associated with the timing and amplitude of the final burst occurring closer to movement onset (*44, 45*). It is closer to movement onset that beta activity serves as an indicator of movement preparation, as final motor plans are enacted (*5, 62–64*), and it is more closely related to whether or not to initiate a movement (*20, 63*).

In conclusion, we provide important insights into the anatomical structures and coding mechanisms supporting both predictive and emergent processes during motor adaptation. We propose a unifying perspective in which sensorimotor predictions and emergent properties of brain activity are tied to the same physiological assumption: that, following a perturbation, beta oscillations are neural signals that allow recovery of behavioral stability. Our findings also open the field to new conception of cerebellar function, with the idea that beta burst signals in cerebellar networks could perform the same predictive function across diverse domains, including language, spatial memory, reward learning (*65*). Our study, though, is also hindered by some current limitations that will be addressed in future work. First, the specific contributions of the cerebellum and the parietal cortex on sensorimotor prediction and motor adjustments could not be disentangled. These regions may operate in tandem, but future work, using non-invasive brain stimulation could help causally delineate their distinctive roles. Second, while MEG improved the localization of cerebellar beta activity, its yet limited spatial resolution compared to functional Magnetic Resonance Imaging (fMRI), calls for caution when pinpointing the role of deep brain sources. Future work assessing changes of beta activity in other forms of learning, or clinical populations with cerebellar dysfunction (e.g., cerebellar ataxia) or diseases with altered beta activity (*e.g.*,

Parkinson's disease (*64*), essential tremor (*21*)) will shed further light on such questions and confirm their relevance in sensorimotor control.

Materials and Methods

Population

We recruited 17 right-handed (as defined by the Edinburgh Questionnaire(*66*)) healthy volunteers (12 females, 45.7±14.1 years; mean ± std) with no history of medical, neurological, or psychiatric conditions, and with normal or corrected-to-normal vision. All participants met the additional MRI safety criteria (*67*) and provided written informed consent. The study was approved by the ethics committee "Ile de France VI - Groupe Hospitalier Pitié-Salpêtrière" (INSERM C13-45) and conducted in accordance with the Declaration of Helsinki.

Experimental paradigm and task description

Participants performed a modified version of a classic sensorimotor reaching task (*42*), consisting of blocks of 50 trials under either normal or rotated visual feedback. Participants controlled a joystick (Thetix, Current Designs, Philadelphia, USA) with their dominant right hand. A gray cursor on the screen indicated the joystick's position. Upon target presentation, participants were instructed to move the cursor towards the visual target swiftly (taking the straightest path). In the normal visual feedback blocks, the cursor's trajectory matched the joystick position and the expected visual feedback (0° rotation), while in the rotated visual feedback the trajectory was rotated either 25° clockwise (25° rotation) or 30° counterclockwise (-30° rotation) relative to the actual joystick trajectory. Rotation amplitudes were selected to induce noticeable trajectory errors and elicit explicit motor adjustments. Stimuli were displayed on a screen with a white background using MATLAB (The MathWorks, Natick, Massachusetts, United States of America (USA)) and its Psychophysics Toolbox. Each trial began with the presentation of a solid gray cursor (0.36° radius, viewed from 80 cm) within a 0.72° radius fixation circle at the bottom center of the screen. After 4.2 s (jittered by ±1.15 s) a circular black target (0.57° radius) appeared at one of five possible locations (9.3° radially from the fixation circle; Figure 1). Target locations were pseudorandomly determined, ensuring equal probability for each location across trials. Each trial ended when the cursor remained within the target for 0.3 s, causing the target to turn green and disappear (Figure 1). Participants were then instructed to return the cursor to the central fixation circle. The maximum reaching movement duration was 2.5 seconds; if participants exceeded this time limit, the target disappeared, prompting participants to return the joystick to the central location. All participants completed two separate experimental sessions on different days. The two sessions occurred at least one week apart but no more than 6 weeks apart. Both sessions were identical, each comprising six blocks of 50 trials of the reaching task, with two blocks per rotation (0°, 25°, or -30° rotation), for a total of 4 blocks per rotation. Blocks with different rotations were randomly interleaved to prevent retention effects across different sessions. The interleaved rotation blocks required participants to continuously update their sensorimotor predictions from one block to another, even in those blocks in which no rotation was applied (*42*).

Behavioral data

We quantified 'trajectory errors' as the Area Under the Curve (AUC) representing the deviation of the actual trajectory from an ideal straight line between the start and target positions (in cm²). The start position was defined as the cursor's location at the time point when it left the fixation circle for at least three consecutive time frames, whereas the endpoint was determined as

the cursor's position when it entered the target boundary. For trials in which the reach movement exceeded 2.5 seconds, resulting in missing data (0 to 6.5% of the total amount of trials, with an average of 3.6 % of trials for the whole population), we used interpolation to estimate these values. Specifically, we used a power law function to fit the collected data (*68*), defined as follows:

$$y = a \cdot x^{-b}$$

Where *y* is the AUC, *x* is the trial number, *a* is the scaling constant, and *b* is the exponent. We used nonlinear least squares fitting to estimate the parameters *a* and *b* from the observed data, and these coefficients were then used to interpolate missing AUC values. To differentiate between transition and stable states, we identified the trial where on average participants reached 80% of their error adjustments within a block (*69*). This threshold captured the transition point in which error reduction stabilized. To do so, for each participant, we calculated a power fit curve for the AUC values and used its first derivative to determine when 80% of the cumulative change had been achieved. On average, this corresponded to the first 16 trials within each block. Thus, the first 16 trials of each block included errors which were categorized as the transition state, whereas for consistency purposes, the last 16 trials of each block represented the stable state (Figure 2A). AUC values were then averaged across these states and blocks with similar perturbations (rotation and no rotation). We assessed motor performance improvement using a 2x2 repeated measures ANOVA with *State* (transition *vs.* stable) and *Perturbation* (rotation *vs.* no rotation) as within-subjects factors. The significance level was set at α=0.05, with the Huynh-Feldt correction applied when necessary to account for violations of sphericity. When appropriate, Bonferroni-corrected post-hoc comparisons were conducted.

Brain data

T1-anatomical image acquisition and preprocessing

We first acquired individual T1-weighted MRI scans (MP-RAGE sequence) using an 8-channel head coil within a 3T MRI system (Verio system, Siemens, Germany) to enable accurate spatial co-registration with the MEG sensors and compute individual head models. The MRI scans were then segmented with Freesurfer (*70*), resulting in a cortical mesh with 15,000 vertices and a cerebellar mesh with 2,000 vertices (Brainstorm; (*71*)).

MEG data acquisition and preprocessing

Participants' oscillatory brain activity was recorded using a 306-sensors whole-head MEG system (Megin Triux), comprising 102 radially oriented magnetometers and 204 horizontal gradiometers. Only gradiometers were used in the analysis due to their higher sensitivity to local cortical sources. Participants' heads were carefully positioned to optimize sensor caudal coverage. Data were digitized at 1000 Hz with a band pass filter of 0-300 Hz. Fiducial markers and other anatomical boundaries were recorded with a 3D-Polhemus FASTRAK® digitizer, and vitamin patches were placed at the fiducial points for accurate registration with the MRI. Head movements were monitored using four coils positioned on the skull, while horizontal and vertical ocular movements were tracked through bipolar electrooculogram (EOG). Additionally, the electrocardiogram (ECG) was recorded using two electrodes placed on the left side of the abdomen and the right clavicle.

To reduce ambient noise, we used the temporal Signal Space Separation (tSSS) algorithm, a version of MaxFilter, in its standard configuration. Unlike the typical approach of segmenting the data into 20-second windows, the analysis was performed on the entire recording to avoid introducing discontinuities in the resulting signal. Further preprocessing was conducted using MATLAB and its open-source Fieldtrip toolbox ((*72*); www.fieldtriptoolbox.org). Specifically, the

data was band-pass filtered (0.5–250 Hz, Butterworth filter), band-stop filtered (49.5–50.5 Hz) and low-pass filtered (40Hz). We then segmented the data into 5-second epochs centered around movement onset (from -3s to 2s). Trials containing artifacts such as SQUID jumps, eye blinks, head movements, and facial muscle activity, were excluded, leading to the removal of approximately 5.8% of trials.

Analysis of the average pre-movement beta activity in sensor space (MEG)

To analyze the average pre-movement beta synchronization, we transformed the MEG data into the time-frequency domain. We followed the Fieldtrip FLUX pipeline for low-frequency brain oscillations ((*73*); https://neuosc.com/flux/), consisting of a fast Fourier transformation for frequencies ranging from 2 to 40 Hz with a 500 ms sliding window using a single Hanning taper. The resulting power was then normalized relative to a common baseline (*7*, *11*). Specifically, power at each sensor, frequency, and time point was divided by the average power across all trials, independent of behavioral condition, and log-transformed. Time-frequency representations were, then, averaged across trials corresponding to the same behavioral state (transition and stable) and perturbations (0°, 25°, -30°). Data from the rotated visual feedback blocks (25° and -30°) were averaged, yielding four conditions per participant: rotation (transition and stable states) and no-rotation (transition and stable state). We performed statistical comparisons using a 2x2 non-parametric cluster-based approach (*74*) with *State* (transition *vs.* stable) and *Perturbation* (rotation *vs.* no rotation) as within-subject factors. We included all gradiometers (204) and individual time points between -2.5 s to 0 (relative to movement onset). We focused on the beta oscillations (14-26 Hz), avoiding potential overlap with the alpha range and aligning with prior studies associating beta modulation with sensorimotor adaptation (*11*, *12*).

To assess pre-movement beta synchronization changes across states and perturbations, we first computed differences between the transition and stable state within each perturbation condition (Δ(stable, transition)). We then run a two-tailed paired t-test at each sample to compare power changes between perturbations (Interaction: rotation, Δ(stable, transition) *vs.* no rotation Δ(stable, transition)). Adjacent time points and sensors were organized into clusters (threshold of p < 0.025). These resulting clusters were then evaluated for statistical significance against a permutation distribution, generated through 100,000 permutations involving random shuffling of conditions across all participants (significance level p < 0.025). We also ran a cluster analysis to test the main effects of *State* and *Perturbation* on pre-movement beta synchronization. To assess the main effect of *State*, we combined the data from both perturbations (rotation and no rotation) and statistically compared the two states (stable *vs.* transition). Similarly, to test the main effect of *Perturbation*, we combined the data from both states (transition and stable) for both perturbations and then statistically compared the two perturbations (rotation *vs.* no rotation). When appropriate, we performed post-hoc cluster tests using the same statistical approach.

Analysis of the average pre-movement beta activity in source space (MEG)

To localize pre-movement beta sources, participants' MRIs and MEG sensor positions were first aligned using the fiducial landmarks and the scalp markers. The individual volume conduction models were then constructed using single-shell head model (*75*), derived from MRI segmentations. The lead field matrix was computed based on each participant's source and volume conduction models, encompassing the cortex and cerebellum (3D grid 1 cm resolution). To identify sources of pre-movement beta activity indicated by the cluster analysis, we used the Dynamical Imaging of Coherent Sources (DICS) approach implemented in Fieldtrip (*76*). We opted for this approach because it is particularly suited for localizing sources in the time-frequency domain (*77*). Source estimates generated by beamforming approaches, including DICS, are prone to a noise bias toward the center of the head (*76*, *77*). This bias was corrected by contrasting conditions where possible

(for example, Interaction: rotation, Δ(stable, transition) *vs.* no rotation, Δ(stable, transition)) or by
applying the neural activity index (NAI; (*78*)) when direct contrasts were unavailable (for example,
Main Effect of *State*: stable *vs.* transition). To estimate the power at each specific brain location
(grid), we used participants' lead fields and computed individual cross-spectral density (CSD)
matrices. CSD matrices were computed using the multitaper method on the beta band (20 Hz, ±6
Hz, covering 14-26 Hz). We focused on sensors and time points identified as significant in the
cluster analysis (*State* x *Perturbation* Interaction). Although cluster-based permutation tests on
time-frequency representations do not directly establish the significance of specific sensor and time-
latency differences (*79*), we could reasonably assume that the identified sensors and time points
corresponded to regions where the effects were strongest. Finally, to highlight the specific regions
contributing the most to pre-movement beta activity, we normalized the individual source signals
using an MNI template with a regularly spaced (8 mm) three-dimensional grid of locations covering
the brain volume. These normalized sources were then imported into SPM12 (Statistical Parametric
Wellcome Department of Cognitive Neurology, London, UK), for further analysis. We performed
paired t-tests comparing the relevant conditions identified by the sensor-space cluster-based results.
Source locations were identified by identifying power peak-maxima (threshold: $p < 0.001$) to
anatomical regions defined by the AAL atlas (*80*).
620
Dynamic connectivity analysis: nonparametric Granger causality (MEG)
We assessed the dynamic connectivity between relevant areas identified by the source
analysis. Specifically, we generated a source model by specifying regions of interest (ROIs) using
predefined MNI coordinates for parietal, frontal, and cerebellar as defined by the AAL atlas (left
parietal inferior, right frontal inferior, and right Crus II; Table S1). To extract source signals, we
applied a linearly constrained minimum variance (LCMV) beamformer, which is well-suited for
localizing sources in the time domain (*23*). As we wanted to determine the temporal evolution of
this connectivity, we analyzed five overlapping 0.45 s time windows before target appearance (0.1
s overlap; from -2.5 to -2.05; from -2.15 to -1.7; from -1.8 to -1.35; from -1.45 to -0.9; from -1.0 to
-0.65). These time-windows were selected to capture at least six cycles of beta oscillations at the
lowest frequency (14 Hz). For each grid location corresponding to the three ROIs, we computed
the CSD matrix of source signals using a fast Fourier transform with 5 Hz frequency smoothing.
The resulting power was then averaged across grid locations belonging to the same ROI. For the
connectivity computation, we applied a multivariate nonparametric Granger causality approach.
Specifically, we used a blockwise method to estimate directional influences between pairs of
sources conditional on the remaining source (*81*). This analysis yielded six directional connectivity
values (two for each pair of ROIs) across the five time-windows.
Finally, to assess how connectivity evolves during the pre-movement period, we performed
six separate 5x2x2 repeated measures ANOVAs with *Time Window* (-2.5 to -2.05, -2.15 to -1.7, -
1.8 to -1.35, -1.45 to -0.9, and -1.0 to -0.65 s), *State* (transition *vs.* stable), and *Perturbation* (rotation
vs. no rotation) as within-subject factors. Significance was set at $\alpha = 0.05$, with Huynh-Feldt
corrections applied when necessary, and Bonferroni post-hoc tests conducted where applicable.
643
Analysis of the single trial-based pre-movement beta activity
To analyze pre-movement beta activity at the single-trial level, we adopted the approach by
Shin et al. (*49*). We focused on sensors showing significant changes from transition to stable states
when comparing the different perturbations. We analyzed a time window from -2.5 to 0 s relative
to the movement onset, and frequencies between 2 and 40 Hz. The time-frequency transformation
used a complex Morlet wavelet characterized by 7 wavelet cycles. The obtained power was
normalized by calculating each frequency's median power across each trial, separately for each
participant and block (factors of median; FOM). Beta bursts were defined as regions within the beta

range (14-26 Hz) where the power exceeded 3 times the median power (3X). We chose this cut-off because it yielded the best representation of the trial-by-trial variability in pre-movement mean power. We determined this threshold empirically as it provided the best representation of trial-by-trial variability in mean pre-movement beta power. Specifically, we quantified the percentage of the spectrogram area exceeding thresholds ranging from 1 to 7 times the median power and correlated this percentage with the mean beta power for each trial (Pearson's test). The 3X threshold yielded the strongest correlation (please see Results; Figure 6B).

Beta bursts can be quantified in different ways. We, therefore, needed to choose the beta feature that best represented the process we aimed to characterize. To do this, we first selected the beta features identified by previous work as relevant to behavior (i.e., bursts mean power, burst number, last burst power; (44, 49)). We then correlated these features with the average beta power (Pearson's test) and used repeated measure ANOVA with beta features as a within-subject factor (beta bursts mean power *vs.* bursts number *vs.* last burst power) to identify the feature with the strongest correlation (i.e., events mean power; Figure 6D). Finally, to verify the relevance of the selected beta feature to motor adjustments, we compared the temporal dynamics of beta burst power to background beta activity. Background activity was defined as the power outside burst periods within each trial. For trials removed due to artifacts, missing values were estimated using a moving median interpolation (8-trial window). A 2x2x2 repeated measures ANOVA was conducted with *Feature* (beta burst power *vs.* background beta activity), *State* (transition *vs.* stable), and *Perturbation* (rotation *vs.* no rotation) as within-subject factors. For all ANOVAs, the significance was set at $\alpha = 0.05$, with Huynh-Feldt correction applied when necessary, and Bonferroni post-hoc tests where applicable.

Brain-behavior relationship: predictive modeling

Model framework

To quantify the relationship between participants' trajectory error evolution and beta burst dynamics, we used a regARMA model, treating the data across trials as a time-series. The regARMA(p,q) model is given by the following equation:

$$Y_t = c + \beta X_t + u_t$$
$$where$$
$$u_t = \sum_{i=1}^{p} a_i Y_{t-i} + \varepsilon_t + \sum_{j=1}^{q} b_i \varepsilon_{t-j}$$

Where Y_t is the dependent variable (trajectory error amplitude at trial t), X_t is the external predictor (beta burst power at trial *t*), β represents the influence of X_t on Y_t (beta bursts contribution to AUC), *p* is the AR order, and *q* is the MA order.

Because regARMA model architecture assumes stationarity, before model implementation, we tested for stationarity of the time-series involved (*Y* and *X*). This analysis was performed across all perturbations on the average data across 17 participants, using the Augmented Dickey-Fuller test. Both trajectory errors and beta burst power time-series were found to be stationary (all p values < 0.05, Table S4).

Optimal model selection

To specify the regARMA model architecture, we determined the autoregressive (AR) order, moving average (MA) order, noise distribution (Gaussian *vs.* t-distribution), and intercept term. To

select the optimal AR and MA orders we used the population average trajectory errors (Y_t) and beta bursts mean power (X_t) across all participants. We examined the Autocorrelation Function (ACF) and Partial Autocorrelation Function (PACF) and noted the lag (m for ACF and n for PACF) at which these sharply dropped below two standard deviations of the data while using a trend stationary model. We then determined the optimal model architecture by comparing all feasible ARMA models by fixing AR and MA orders at n and m respectively, and varying noise distributions (Gaussian *vs.* t-distribution) and the inclusion of an intercept term. Model selection was based on the lowest AIC value among models where β significantly contributed to explaining variance. After fixing the noise distribution and deciding on the inclusion or exclusion of an intercept term based on the AIC values, we also tested models with AR and MA orders increasing and decreasing from (n,m) to check if the AIC values improved. A summary of AIC values for all feasible tested ARMA models is provided in Table S2. We selected regARMA (1,1) as the optimal candidate because it was consistently the best model for all rotations with the least AIC values. Only rotation -30° had two possible candidates, i.e., ARMA (1,1) and ARMA (1,2). However, as the difference between their AIC values was lower than 2, we opted for the model consistent with the other rotations.

Model parameter estimation and validation

We validated the optimal model selected across the population of 17 participants using a leave-one-out approach. For each participant and perturbation, we estimated the model parameters using data from the population average of the remaining 16 participants. The stationarity was reconfirmed for each of the averages calculated from the different combinations of 16/17 participants before the parameter estimation. These parameters were then used to predict the left-out participant's trajectory error time-series. Specifically, we used the participant's error from the first trial, along with the trial-by-trial beta burst dynamics over the 50 trials of a block, to predict the trajectory error across the remaining 49 trials (Figures 7A; see Figures S1, S2 and S3 for individual simulations). To quantify prediction errors, we calculated the prediction mean squared error (PMSE) between the model's predictions and observed data, from the 2nd to 50th trials.

$$PMSE = \sqrt{\frac{1}{49}\sum_{t=2}^{50}\left(Y_t^{sim} - Y_t^{\exp}\right)^2}$$

Given that PMSE can vary with the scale of the data (e.g., trajectory errors amplitude at 0° rotation are typically smaller than those at 25° or -30°), we min-max normalized the PMSE (N-PMSE) to ensure fair comparisons across different conditions, e.g., perturbations or different models. In the min-max normalization was done as

$$N - PMSE = \frac{PMSE - \min\left(PMSE_{condition}\right)}{\min\left(PMSE_{condition}\right) - \max\left(PMSE_{condition}\right)}$$

To test whether beta burst power differently influenced error amplitude evolution across perturbations, we conducted repeated-measures ANOVA with *Perturbation* (0° *vs.* 25° *vs.* -30°) as a within-subject factor using beta contribution (*β*) as the dependent variable. Significance was set at α = 0.05, with Huynh-Feldt corrections and Bonferroni post-hoc tests applied where necessary. Similar analyses were conducted to compare the model prediction accuracy (N-PMSE) across perturbations.

Model specificity

We also tested whether the relationship between trajectory errors and beta burst power was specific to frequency. To this aim, we replaced beta burst power with alpha mean power while predicting reach errors. Alpha mean power was used instead of alpha burst power due to the absence of reliable pre-movement alpha bursts in our data (Figure S5). We employed the same model selection, estimation, and validation processes described in the previous section (for details on alpha mean power time-series stationarity and model selection, please see tables S4 and S5). To determine the specificity of the original model, we compared model fit (AIC) using a 2x3 repeated-measures ANOVA with *Model* (beta bursts *vs.* alpha mean power) and *Perturbations* (0° *vs.* 25° *vs.* -30°) as within-subject factors. Significance was set at $\alpha = 0.05$, with Huynh-Feldt corrections applied when necessary, and Bonferroni corrected post-hoc tests conducted where applicable. The same approach was used for a control analysis, where we compared models using mean alpha power with those using mean beta power as external predictors (for details on alpha and beta mean power time-series stationarity and model selection, please see tables S3, S5 and S6).

References

1. J. Izawa, T. Rane, O. Donchin, R. Shadmehr, Behavioral/Systems/Cognitive Motor Adaptation as a Process of Reoptimization. doi: 10.1523/JNEUROSCI.5359-07.2008 (2008).
2. A. Fuchs, V. K. Jirsa, J. A. S. Kelso, Theory of the Relation between Human Brain Activity (MEG) and Hand Movements. *Neuroimage* **11**, 359–369 (2000).
3. E. M. Reuter, A. Booms, L. A. Leow, Using EEG to study sensorimotor adaptation. *Neurosci Biobehav Rev* **134**, 104520 (2022).
4. L. Struber, M. Baumont, P. A. Barraud, V. Nougier, F. Cignetti, Brain oscillatory correlates of visuomotor adaptive learning. *Neuroimage* **245**, 118645 (2021).
5. M. Bracco, D. Veniero, M. Oliveri, G. Thut, Prismatic adaptation modulates oscillatory EEG correlates of motor preparation but not visual attention in healthy participants. *Journal of Neuroscience* **38** (2018).
6. S. Espenhahn, B. C. M. van Wijk, H. E. Rossiter, A. O. de Berker, N. D. Redman, J. Rondina, J. Diedrichsen, N. S. Ward, Cortical beta oscillations are associated with motor performance following visuomotor learning. *Neuroimage* **195**, 340–353 (2019).
7. H. Tan, C. Wade, X. P. Brown, Post-Movement Beta Activity in Sensorimotor Cortex Indexes Confidence in the Estimations from Internal Models. doi: 10.1523/JNEUROSCI.3204-15.2016 (2016).
8. A. Schwey, D. Battaglia, J. Bahuguna, N. Malfait, Behavioral/Cognitive Different Faces of Medial Beta-Band Activity Reflect Distinct Visuomotor Feedback Signals. doi: 10.1523/JNEUROSCI.2238-22.2023 (2023).
9. H. Tan, N. Jenkinson, P. Brown, Systems/Circuits Dynamic Neural Correlates of Motor Error Monitoring and Adaptation during Trial-to-Trial Learning. doi: 10.1523/JNEUROSCI.4739-13.2014 (2014).
10. J. Alayrangues, F. Torrecillos, A. Jahani, N. Malfait, Error-related modulations of the sensorimotor post-movement and foreperiod beta-band activities arise from distinct neural substrates and do not reflect efferent signal processing. *Neuroimage* **184**, 10–24 (2019).
11. F. Torrecillos, J. Alayrangues, B. E. Kilavik, N. Malfait, Distinct Modulations in Sensorimotor Postmovement and Foreperiod β-Band Activities Related to Error Salience Processing and Sensorimotor Adaptation. *Journal of Neuroscience* **35**, 12753–12765 (2015).
12. H. T. Darch, N. L. Cerminara, I. D. Gilchrist, R. Apps, Pre-movement changes in sensorimotor beta oscillations predict motor adaptation drive. *Scientific Reports 2020 10:1* **10**, 1–12 (2020).

13. C. E. Palmer, R. Auksztulewicz, S. Ondobaka, J. M. Kilner, Sensorimotor beta power reflects
the precision-weighting afforded to sensory prediction errors. *Neuroimage* **200**, 59–71
(2019).
14. A. K. Engel, P. Fries, Beta-band oscillations — signalling the status quo? *Curr Opin*
Neurobiol **20**, 156–165 (2010).
15. B. Pollok, M. Butz, J. Gross, A. Schnitzler, Intercerebellar Coupling Contributes to Bimanual
Coordination. *J Cogn Neurosci* **19**, 704–719 (2007).
16. D. S. Soteropoulos, S. N. Baker, Cortico-cerebellar coherence during a precision grip task in
the monkey. *J Neurophysiol* **95**, 1194–1206 (2006).
17. J. P. Pellerin, Y. Lamarre, Local field potential oscillations in primate cerebellar cortex
during voluntary movement. *J Neurophysiol* **78**, 3502–3507 (1997).
18. R. Courtemanche, J. P. Pellerin, Y. Lamarre, Local field potential oscillations in primate
cerebellar cortex: Modulation during active and passive expectancy. *J Neurophysiol* **88**, 771–
782 (2002).
19. B. Pollok, J. Gross, D. Kamp, A. Schnitzler, Evidence for Anticipatory Motor Control within
a Cerebello-Diencephalic-Parietal Network. (2007).
20. S. Picazio, D. Veniero, V. Ponzo, C. Caltagirone, J. Gross, G. Thut, G. Koch, Prefrontal
control over motor cortex cycles at beta frequency during movement inhibition. *Current*
Biology **24**, 2940–2945 (2014).
21. M. K. Pan, Y. S. Li, S. B. Wong, C. L. Ni, Y. M. Wang, W. C. Liu, L. Y. Lu, J. C. Lee, E.
P. Cortes, J. P. G. Vonsattel, Q. Sun, E. D. Louis, P. L. Faust, S. H. Kuo, Cerebellar
oscillations driven by synaptic pruning deficits of cerebellar climbing fibers contribute to
tremor pathophysiology. *Sci Transl Med* **12**, 1769 (2020).
22. P. Khanna, J. M. Carmena, Beta band oscillations in motor cortex reflect neural population
signals that delay movement onset. *Elife* **6** (2017).
23. W.-J. Neumann, A. Jha, A. Bock, J. Huebl, A. Horn, G.-H. Schneider, T. H. Sander, V.
Litvak, A. A. Kü Hn, A. A. Kü, Cortico-pallidal oscillatory connectivity in patients with
dystonia. doi: 10.1093/brain/awv109.
24. J. S. Brittain, A. Sharott, P. Brown, The highs and lows of beta activity in cortico-basal
ganglia loops. *European Journal of Neuroscience* **39**, 1951–1959 (2014).
25. J. W. Chung, E. Ofori, G. Misra, C. W. Hess, D. E. Vaillancourt, Beta-band activity and
connectivity in sensorimotor and parietal cortex are important for accurate motor
performance. *Neuroimage* **144**, 164–173 (2017).
26. R. Courtemanche, D. Popa, C. Léna, Exploring oscillations in expert sensorimotor
anticipation: The tennis return of serve. *Neuronal Oscillations of Wakefulness and Sleep:*
Windows on Spontaneous Activity of the Brain, 3–44 (2020).
27. S. J. Blakemore, A. Sirigu, Action prediction in the cerebellum and in the parietal lobe. *Exp*
Brain Res **153**, 239–245 (2003).
28. R. C. Miall, D. M. Wolpert, Forward Models for Physiological Motor Control. *Neural*
Networks **9**, 1265–1279 (1996).
29. R. Shadmehr, J. W. Krakauer, A computational neuroanatomy for motor control. doi:
10.1007/s00221-008-1280-5.
30. H. Imamizu, S. Miyauchi, T. Tamada, Y. Sasaki, R. Takino, B. Pütz, T. Yoshioka, M.
Kawato, Human cerebellar activity reflecting an acquired internal model of a new tool.
Nature 2000 403:6766 **403**, 192–195 (2000).
31. J. Jankowski, S. Paus, L. Scheef, M. Bewersdorff, H. H. Schild, T. Klockgether, H. Boecker,
Abnormal Movement Preparation in Task-Specific Focal Hand Dystonia. *PLoS One* **8**,
e78234 (2013).
32. F. P. Chabrol, A. Blot, T. D. Mrsic-Flogel, Cerebellar Contribution to Preparatory Activity
in Motor Neocortex. *Neuron* **103**, 506-519.e4 (2019).

33.   Z. Gao, C. Davis, A. M. Thomas, M. N. Economo, A. M. Abrego, K. Svoboda, C. I. De
841   Zeeuw, N. Li, A cortico-cerebellar loop for motor planning. *Nature 2018 563:7729* **563**,
842   113–116 (2018).
843   34.   E. Courchesne, G. Allen, Prediction and preparation, fundamental functions of the
844   cerebellum, *Learning & Memory 4(1)* (1997)pp. 1–35.
845   35.   R. Ivry, Exploring the role of the cerebellum in sensory anticipation and timing: Commentary
846   on Tesche and Karhu. *Hum Brain Mapp* **9**, 115 (2000).
847   36.   A. Pilacinski, A. Lindner, Distinct contributions of human posterior parietal and dorsal
848   premotor cortex to reach trajectory planning. *Scientific Reports 2019 9:1* **9**, 1–16 (2019).
849   37.   D. M. Wolpert, S. J. Goodbody, M. Husain, Maintaining internal representations: the role of
850   the human superior parietal lobe. *Nature Neuroscience 1998 1:6* **1**, 529–533 (1998).
851   38.   A. Genovesio, S. P. Wise, R. E. Passingham, Prefrontal–parietal function: from foraging to
852   foresight. *Trends Cogn Sci* **18**, 72–81 (2014).
853   39.   S. Kuang, P. Morel, A. Gail, Planning Movements in Visual and Physical Space in Monkey
854   Posterior Parietal Cortex. *Cerebral Cortex* **26**, 731–747 (2016).
855   40.   R. A. Andersen, C. A. Buneo, INTENTIONAL MAPS IN POSTERIOR PARIETAL
856   CORTEX. *Annu. Rev. Neurosci* **25**, 189–220 (2002).
857   41.   R. Shadmehr, M. A. Smith, J. W. Krakauer, Error Correction, Sensory Prediction, and
858   Adaptation in Motor Control. doi: 10.1146/annurev-neuro-060909-153135 (2010).
859   42.   J. W. Krakauer, C. Ghez, M. F. Ghilardi, Adaptation to Visuomotor Transformations:
860   Consolidation, Interference, and Forgetting. *The Journal of Neuroscience* **25**, 473–478
861   (2005).
862   43.   Y. J. John, K. S. Sawyer, K. Srinivasan, E. J. Müller, B. R. Munn, J. M. Shine, It's about
863   time: Linking dynamical systems with human neuroimaging to understand the brain.
864   *Network Neuroscience* **6**, 960–979 (2022).
865   44.   F. Torrecillos, G. Tinkhauser, P. Fischer, A. L. Green, T. Z. Aziz, T. Foltynie, P. Limousin,
866   L. Zrinzo, K. Ashkan, P. Brown, H. Tan, Modulation of Beta Bursts in the Subthalamic
867   Nucleus Predicts Motor Performance. *Journal of Neuroscience* **38**, 8905–8917 (2018).
868   45.   S. Little, J. Bonaiuto, G. Barnes, S. Bestmann, Human motor cortical beta bursts relate to
869   movement planning and response errors. *PLoS Biol* **17**, e3000479 (2019).
870   46.   T. O. West, B. Duchet, S. F. Farmer, K. J. Friston, H. Cagnan, When do bursts matter in the
871   primary motor cortex? Investigating changes in the intermittencies of beta rhythms
872   associated with movement states. *Prog Neurobiol* **221** (2023).
873   47.   J. J. Bonaiuto, S. Little, S. A. Neymotin, S. R. Jones, G. R. Barnes, S. Bestmann, Laminar
874   dynamics of high amplitude beta bursts in human motor cortex. *Neuroimage* **242**, 118479
875   (2021).
876   48.   J. R. Wessel, Behavioral/Cognitive-Bursts Reveal the Trial-to-Trial Dynamics of Movement
877   Initiation and Cancellation. doi: 10.1523/JNEUROSCI.1887-19.2019 (2020).
878   49.   H. Shin, R. Law, S. Tsutsui, C. I. Moore, S. R. Jones, The rate of transient beta frequency
879   events predicts behavior across tasks and species. *Elife* **6** (2017).
880   50.   R. M. Hardwick, C. Rottschy, R. C. Miall, S. B. Eickhoff, A quantitative meta-analysis and
881   review of motor learning in the human brain. *Neuroimage* **67**, 283–297 (2013).
882   51.   F. Gheysen, G. Lasne, M. Pélégrini-Issac, G. Albouy, S. Meunier, H. Benali, J. Doyon, T.
883   Popa, Taking the brakes off the learning curve. *Hum Brain Mapp* **38**, 1676 (2016).
884   52.   L. S. Popa, T. J. Ebner, Cerebellum, Predictions and Errors. *Front Cell Neurosci* **12** (2018).
885   53.   N. Sendhilnathan, A. C. Bostan, P. L. Strick, M. E. Goldberg, A cerebro-cerebellar network
886   for learning visuomotor associations. *Nature Communications 2024 15:1* **15**, 1–14 (2024).
887   54.   E. R. Palacios, C. Houghton, P. Chadderton, Accounting for uncertainty: Inhibition for
888   neural inference in the cerebellum. *Proceedings of the Royal Society B: Biological Sciences*
889   **288** (2021).

55. M. Manto, J. M. Bower, A. B. Conforto, J. M. Delgado-García, S. N. F. Da Guarda, M. Gerwig, C. Habas, N. Hagura, R. B. Ivry, P. Marien, M. Molinari, E. Naito, D. A. Nowak, N. O. Ben Taib, D. Pelisson, C. D. Tesche, C. Tilikete, D. Timmann, Consensus Paper: Roles of the Cerebellum in Motor Control—The Diversity of Ideas on Cerebellar Involvement in Movement. *Cerebellum* **11**, 457 (2012).

56. K. Friston, R. Adams, R. Montague, What is value—accumulated reward or evidence? *Front Neurorobot* **6** (2012).

57. P. A. Butcher, R. B. Ivry, S.-H. Kuo, D. Rydz, J. W. Krakauer, J. A. Taylor, The cerebellum does more than sensory prediction error-based learning in sensorimotor adaptation tasks. *J Neurophysiol* **118**, 1622–1636 (2017).

58. J. Doyon, H. Benali, Reorganization and plasticity in the adult brain during learning of motor skills. *Curr Opin Neurobiol* **15**, 161–167 (2005).

59. J. G. Samuelsson, P. Sundaram, S. Khan, M. I. Sereno, M. S. Hämäläinen, Detectability of cerebellar activity with magnetoencephalography and electroencephalography. *Hum Brain Mapp* **41**, 2357–2372 (2020).

60. L. M. Andersen, K. Jerbi, S. S. Dalal, Can EEG and MEG detect signals from the human cerebellum? *Neuroimage* **215**, 116817 (2020).

61. J. H. Holland, Complexity: A Very Short Introduction. *Complexity: A Very Short Introduction*, doi: 10.1093/ACTRADE/9780199662548.001.0001 (2014).

62. H. Shibasaki, M. Hallett, What is the Bereitschaftspotential? doi: 10.1016/j.clinph.2006.04.025 (2006).

63. Y. Zhang, Y. Chen, S. L. Bressler, M. Ding, Response preparation and inhibition: The role of the cortical sensorimotor beta rhythm. *Neuroscience* **156**, 238 (2008).

64. S. Little, P. Brown, The functional role of beta oscillations in Parkinson's disease. *Parkinsonism Relat Disord* **20**, S44–S48 (2014).

65. J. Diedrichsen, M. King, C. Hernandez-Castillo, M. Sereno, R. B. Ivry, Universal Transform or Multiple Functionality? Understanding the Contribution of the Human Cerebellum across Task Domains. *Neuron* **102**, 918–928 (2019).

66. R. C. Oldfield, The assessment and analysis of handedness: The Edinburgh inventory. *Neuropsychologia* **9**, 97–113 (1971).

67. F. G. Shellock, A. Spinazzi, MRI safety update 2008: Part 2, screening patients for MRI. *American Journal of Roentgenology* **191**, 1140–1149 (2008).

68. V. Della-Maggiore, J. Scholz, H. Johansen-Berg, T. Paus, The rate of visuomotor adaptation correlates with cerebellar white-matter microstructure. *Hum Brain Mapp* **30**, 4048–4053 (2009).

69. A. G. Feldman, M. F. Levin, The Equilibrium-Point Hypothesis – Past, Present and Future. *Adv Exp Med Biol* **629**, 699–726 (2009).

70. B. Fischl, FreeSurfer. *Neuroimage* **62**, 774–781 (2012).

71. F. Tadel, S. Baillet, J. C. Mosher, D. Pantazis, R. M. Leahy, Brainstorm: A User-Friendly Application for MEG/EEG Analysis. *Comput Intell Neurosci* **2011**, 13 (2011).

72. R. Oostenveld, P. Fries, E. Maris, J. M. Schoffelen, FieldTrip: Open Source Software for Advanced Analysis of MEG, EEG, and Invasive Electrophysiological Data. *Comput Intell Neurosci* **2011**, 156869 (2011).

73. O. Ferrante, L. Liu, T. Minarik, U. Gorska, T. Ghafari, H. Luo, O. Jensen, FLUX: A pipeline for MEG analysis. *Neuroimage* **253**, 119047 (2022).

74. E. Maris, R. Oostenveld, Nonparametric statistical testing of EEG- and MEG-data. *J Neurosci Methods* **164**, 177–190 (2007).

75. G. Nolte, The magnetic lead field theorem in the quasi-static approximation and its use for magnetoencephalography forward calculation in realistic volume conductors. *Phys Med Biol* **48**, 3637 (2003).

76. J. Gross, J. Kujala, M. Hä Mä Lä Inen, L. Timmermann, A. Schnitzler, R. Salmelin, Dynamic imaging of coherent sources: Studying neural interactions in the human brain. *PNAS* **98**, 694–699 (2001).

77. B. U. Westner, S. S. Dalal, A. Gramfort, V. Litvak, J. C. Mosher, R. Oostenveld, J. M. Schoffelen, A unified view on beamformers for M/EEG source reconstruction. *Neuroimage* **246**, 118789 (2022).

78. B. D. Van Veen, W. Van Drongelen, M. Yuchtman, A. Suzuki, Localization of brain electrical activity via linearly constrained minimum variance spatial filtering. *IEEE Trans Biomed Eng* **44**, 867–880 (1997).

79. J. Sassenhagen, D. Draschkow, Cluster-based permutation tests of MEG/EEG data do not establish significance of effect latency or location. *Psychophysiology* **56**, e13335 (2019).

80. N. Tzourio-Mazoyer, B. Landeau, D. Papathanassiou, F. Crivello, O. Etard, N. Delcroix, B. Mazoyer, M. Joliot, Automated Anatomical Labeling of Activations in SPM Using a Macroscopic Anatomical Parcellation of the MNI MRI Single-Subject Brain. *Neuroimage* **15**, 273–289 (2002).

81. Y. Chen, S. L. Bressler, M. Ding, Frequency decomposition of conditional Granger causality and application to multivariate neural field potential data. (2006).

Acknowledgments

This work was carried out in the MRI and MEG platform of the CENIR within the Paris Brain Institute (ICM).

Fundings

This study was supported by:
Areva (SM and DS)
Fondation pour la Recherche Médicale (CGa)
European Union's Horizon 2020 research and innovation program; the Marie Słodowska-Curie grant agreement 897941 (MB)

Authors contributions

Conceptualization: MB, SM, DS, TP, CG. Methodology: MB, VV, CGa. Data Collection: QW, AR, CGa. Data Analysis: MB, VV, VR, QW, MR. Visualization: MB, VV, CGa. Supervision: AVC, DS, TP, CGa. Writing—original draft: MB, VV, CGa. Writing—review & editing: MB, VV, VR, WQ, MR, AR, CGi, SM, AVC, DS, TP, CGa.

Competing interests

The authors declare that they have no competing interests.

Data and materials availability

The data will be publicly available upon publication. Data analysis was performed using publicly available software and toolboxes as described in the Methods section. Any additional information may be requested from the corresponding authors (MB, CG).

Supplementary Materials

990   Supplementary Materials contain:
991   Figures S1 to S4
992   Tables S1 to S6
993

994
Figures and Tables

Fig. 1. Reaching task description.
Each trial began with the presentation of a solid gray cursor within a fixation circle at the bottom
center of the screen. After 4.2 s (jittered by ±1.15 s), a circular black target appeared at one of five
possible locations. We instructed participants to swiftly move the cursor toward the visual target,
aiming to follow the straightest path (ideal trajectory). Each reaching movement had a maximum
duration of 2.5 s; if this limit was exceeded, the target disappeared, and participants were prompted
to return the joystick to the central position. A trial ended when the cursor remained within the
target for 0.3 s, causing the target to turn green and disappear. Participants then returned the cursor
to the central fixation circle. Participants performed blocks of consecutive trials of the same
perturbation. In the "No rotation" blocks, the cursor's trajectory aligned with the joystick movement
and expected visual feedback. In the "Rotation" blocks, the visual feedback of the trajectory was
rotated either 25° clockwise (25° rotation) or 30° counterclockwise (-30° rotation) relative to the
real trajectory. The red line indicates an example of trajectory during the initial states of both "No
rotation" and "Rotation" blocks, while the black dashed line represents the ideal trajectory.
1011

1012

Fig. 2. Behavioral performance results.

(**A**) Evolution of trajectory error amplitude (AUC) along 50 consecutive trials of a block, averaged across participants relative to *Perturbation* (Rotation, red; No rotation black). Shading indicates SEM. Vertical dashed lines divide the blocks into transition (first 16 trials) and stable state(last 16 trials). (**B**) Results of the 2x2 repeated measures ANOVA comparing the trajectory errors categorized by *State* (transition *vs.* stable) and *Perturbation* (rotation *vs.* no rotation). Box plots show the median (horizontal line), the lower and upper quartiles (box), the minimum and maximum values that are not outliers (whiskers), and the outliers, computed using the interquartile range (dots). *** indicates p < 0.001; ns indicates p > 0.05.

1023

Fig. 3. Time-frequency power spectra across different States and Perturbations.
The time-frequency plots illustrate the power spectrum for frequencies ranging from 12 to 40 Hz, spanning -2.5 to 0 seconds relative to movement onset. Power was normalized by dividing each time-frequency-sensor value by the average power across all trials, independent of behavioral conditions, which was then log-transformed. For visualization, data was averaged across all 204 horizontal gradiometers and all participants. The black vertical dashed line indicates the median target appearance (i.e., -0.5 s); the black vertical solid line indicates the movement onset (*i.e.*, 0 s); the white horizontal dashed lines highlight the beta frequency range analyzed (14-26 Hz).

1033

Fig. 4. Averaged pre-movement beta oscillatory activity results.

(**A**) Main effect of *State*, obtained by combining the data from both *Perturbations* (rotation and no rotation). (**B**) Interaction effect between *State* and *Perturbation*, obtained by comparing the difference in beta activity between transition and stable states within each perturbation (Δ(stable, transition)). For each effect, the results of the cluster-based analysis are represented with power spectra plots (top row), scalp topographies (central row), and source space representations (bottom row). For the power spectra plots, the two-dimensional clusters (time and sensors) are collapsed across the significant sensors (shown within the topography plots) to display the temporal evolution of the analyzed beta power (14-26 Hz). The vertical dashed line indicates median target appearance (at -0.5 s), and shaded areas mark time windows significant in the cluster-based analysis (Main effect of *State*: -2.5 to -1.35 seconds; Interaction of *State* × *Perturbation*: -2.35 to -2 seconds). For scalp topographies, data are collapsed across significant time points to highlight spatial distributions. Only significant t-values are shown, and black dots indicate significant sensors (threshold: $p < 0.05$). The source space results (threshold: $p < 0.001$) are overlaid on a 3D brain surface template (bottom panel).

1049

1050

Fig. 5. Conditional Granger causality results.

(**A**) Main effect of *State* for the 5x2x2 repeated measures ANOVA (comparing transition *vs.* stable states independent of the other factors). (**B**) Main effect of *Time Window* from the same ANOVA, comparing the five time-windows (-2.5/-2.05; -2.15/-1.7; -1.8/-1.35; -1.4/-0.9; -1.0/-0.65). (**C**) Graphical representation of the Conditional Granger Causality analysis showing the effect of *State* (left) and *Pre-movement window* (right). rCB: right cerebellum (Crus II; purple); rPFC: right prefrontal cortex (frontal inferior triangularis; organge); lPC: left parietal cortex (parietal inferior, green). Box plots show the median (horizontal line), the lower and upper quartiles (box), the minimum and maximum values that are not outliers (whiskers), and the outliers, computed using the interquartile range (dots). Dashed arrows reflect the connectivity direction. ** indicates $p < 0.025$; * indicates $p < 0.05$.

1064

Fig. 6. Beta bursts features.

(**A**) Beta bursts in the 14–26 Hz range are shown from -2.5 seconds to movement onset for the first three and last three trials of an example block in a representative participant. Data were normalized as factors of the median (FOM) for each frequency, calculated separately for each participant and condition. (**B**) The threshold for data normalization was selected based on the highest correlation between the percentage of the spectrogram area exceeding thresholds ranging from 1 to 7 times the median power and the mean beta power for each trial. Colored lines represent the averages of trials and blocks for individual participants, while the black line shows the group average. The dashed vertical black line indicates the threshold with the highest correlation (3X median power). (**C**) Probability distribution of events number (top) and events duration (bottom), where an event is a beta burst exceeding the 3X median cut-off. Distributions are plotted as the average across trials and blocks for individual participants (colored lines), with the population average in black. (**D**) Correlations between selected beta burst features (mean beta burst power, number of bursts, and power of the last burst) and average beta power. (**E**) Control analysis results verifying the relevance of the selected beta feature (mean burst power) to error reduction. Box plots depict the results of the 2×2×2 repeated measures ANOVA, with *Feature* (beta burst power *vs.* background beta activity), *State* (transition *vs.* stable), and *Perturbation* (rotation *vs.* no rotation) as factors. Box plots show the median (horizontal line), the lower and upper quartiles (box), the minimum and maximum values that are not outliers (whiskers), and the outliers, computed using the interquartile range (dots). *** indicate $p < 0.001$.

1085

1086

Fig. 7. Predictive model results.
(**A**) Real and predicted trajectory error amplitude along 50 consecutive trials of a block, averaged
across participants for each perturbation (Rotation with -30° in red and 25° in blue; No rotation,
black). The amplitude of real trajectory errors is quantified as the area under the curve (AUC),
representing the deviation from the ideal straight-line trajectory between the start and target
positions. Higher AUC values indicate greater trajectory error amplitude. Predicted errors are
estimated by validating the chosen regARMA(1,1) model using a leave-one-out cross-validation
approach. Shading indicates SEM. (**B**) Results of a repeated measures ANOVA comparing the
coefficient of the regARMA model's external predictor, representing the contribution of beta burst
power to trajectory error evolution, across perturbations (0° *vs.* 25° *vs.* -30°). (**C**) Results of a
repeated measures ANOVA comparing the normalized prediction mean squared error (N-PMSE)
of the regARMA models across different perturbations (0° *vs.* 25° *vs.* -30°). Box plots show the
median (horizontal line), the lower and upper quartiles (box), the minimum and maximum values
that are not outliers (whiskers), and the outliers, computed using the interquartile range (dots). ***
indicates $p < 0.001$; ns indicates $p > 0.05$.
1102
1103