Critical Scaling of Novelty in the Cortex

Tiago L. Ribeiro, Ali Vakili, Bridgette Gifford, Raiyyan Siddiqui, Vincent Sinfuego, Sinisa Pajevic, and Dietmar Plenz[†]

Section on Critical Brain Dynamics, National Institute of Mental Health, Bethesda, MD, USA

[†] Correspondence: plenzd@mail.nih.gov

Running Title: Critical scaling of novel cortical action potentials *in vivo*

Correspondence: Dietmar Plenz, Ph.D., Section on Critical Brain Dynamics, National
                 Institute of Mental Health, Porter Neuroscience Research Center, Rm 3A-
                 1000, 35 Convent Drive, Bethesda, MD 20892. office: (301) 675-5311

Keywords: Perturbation, population coding, criticality, scaling, neuronal avalanches, visual
             cortex, *in vivo*, 2-photon imaging, holography, all-optical

Abstract

The ability to detect and transmit novel events is essential for adaptive behavior in uncertain environments. Here, we investigate how holographically triggered, unanticipated action potentials propagate through the primary visual cortex of resting mice, focusing on pyramidal neuron communication. We find that these novel spikes — uncorrelated with ongoing activity — exert a disproportionately large influence on neighboring neurons, whose response scales as a power law (exponent ~0.2–0.3). Even a few such spikes can recruit a large fraction of the local network, enabling robust decoding of perturbation origin despite high trial-by-trial variability and ongoing activity dominated by large activity fluctuations in the form of scale-invariant, parabolic neuronal avalanches. Simulations confirm this scaling to small, local perturbations aligns with the high susceptibility of complex systems near criticality. These results suggest that critical dynamics facilitate efficient transmission of novel signals, revealing a fundamental mechanism for cortical novelty detection.

Introduction

Thriving in uncertain environments depends on the brain's ability to detect and respond to unexpected events. In mammals, such events must be rapidly and effectively communicated across brain regions to support accurate evaluation and adaptive behavior[1]. In the cortex, the fundamental unit of this communication is the action potential generated by a single pyramidal neuron. Therefore, understanding how the brain processes unexpected events requires examining how novel action potentials — those that deviate from near-future predictions based on ongoing activity — are transmitted through the surrounding cortical network.

Neuronal spiking has been found to shape cortical activity and behavior. A spike in a pyramidal neuron from superficial layers (L2/3) of cortex of mice has been estimated to influence fewer than 1% of nearby neurons, to evoke strong local inhibition[2-5], and in reptilian cortex to reliably recruit activity sequences[6]. Coordinated activity in small groups of L2/3 pyramidal neurons elicits spatially extended variable responses[7] and can drive trained behavior via pattern completion[8,9]. Similarly, spike bursts in deep layers (L5/6) can elicit sensitized behavioral responses[10,11] and alter global brain states[12,13]. These holographic perturbation studies in mice are in line with previous findings from cortical micro stimulation in primates highlighting the capacity of few spikes to significantly alter brain activity under specific conditions (for review[14]). However, a comprehensive framework to quantify the impact of action potentials on the spatiotemporal dynamics of cortical networks remains elusive.

In principle, novel spikes representing unexpected events should induce propagation

across the cortex both broadly (to reach multiple functional domains for evaluation given their

uncertain nature) and rapidly (to be able to act quickly). However, this dissemination of

information is constrained by several factors. First, even in the resting state, the cortex exhibits

substantial activity, with each neuron receiving thousands of action potentials every second.

Accordingly, individual neurons operate in a fluctuation-dominated[15], high-dimensional[16] regime

that affects their ability to respond to and encode information from novel spikes[17] or sensory

inputs[18]. Second, the sparse synaptic connectivity between L2/3 pyramidal neurons —estimated

at only 5–8% locally and diminishing sharply within 200 µm[19-23] — further restricts the

transmission of novel spikes[24]. Additionally, the weak and transient synaptic currents of these

failure-prone connections in vivo[2,22], coupled with strong local inhibition[25,26], further reduce the

potential influence of novel spikes introduced randomly into a pyramidal neuron in the awake

state. Given these limitations, understanding how novel spikes are encoded requires a focus on

the network's broader dynamics rather than the activity of individual neurons or microcircuits

within a particular experimental constraint. However, adopting this broader perspective presents

a significant challenge due to the inherent complexity of cortical activity.

Research suggests that brain networks operating near a critical state exhibit heightened

sensitivity to small perturbations[27,28], which might enable the cortex to detect and process new

information despite its complex internal activity and sparse topology. In this state, the system

undergoes internal fluctuations while remaining responsive to small, local events that can

propagate broadly, facilitating the recognition and dissemination of novel information. In the

superficial cortical layers, these fluctuations manifest as "neuronal avalanches" — scale-

invariant patterns of synchronized neuron activity characteristic of the awake state[29-31]. These

avalanches align with theoretical predictions of critical dynamics. However, direct experimental

evidence demonstrating that novel, localized information propagates within the neuronal

avalanche regime and scales in line with predictions from critical dynamics is lacking, leaving a

gap in understanding whether and how criticality supports the spread of novel information in the

brain.

Using an all-optical approach, we holographically triggered spikes in individual L2/3

pyramidal neurons while monitoring activity of hundreds of nearby neurons in the primary visual

cortex of awake mice. Our results demonstrate robust scaling of network responses during

spontaneous, ongoing avalanche activity, with novel spikes rapidly generating widespread, yet

transient information-rich activity across large cortical distances. Simulations show that this

pattern of response emerges naturally in networks operating near a critical state, where small

inputs can lead to large effects. Together, our findings suggest that critical dynamics play a key

role in enabling the brain to efficiently process unexpected information.

Results

89 Holographically triggered spikes in a pyramidal neuron induce rapid, widespread

90 excitation in superficial cortical layers

In awake, quietly resting adult mice, we holographically targeted single L2/3 pyramidal neurons

(Target Cells, TC) in primary visual cortex (V1) that chronically co-expressed the calcium

indicator jGCaMP7s and the opsin ChrimsonR (~5 – 10 mW, 100 ms; ~100 – 150 trials per TC)

while simultaneously monitoring spiking activity from ~100 – 300 nearby neurons using 2-

photon imaging (Fig. 1a; 2PI, field of view: 450 μm × 450 μm; ~50 – 140 μm cortical depth;

temporal resolution: 22 ms; n = 6 mice; 16 experiments). Sparse co-expression, stimulation

beam profiling, motion correction analysis, and anatomical response analysis confirmed select

single neuron stimulation (Supplementary Figs. 1–5).

Holographic stimulation of a single pyramidal neuron caused wide-spread, transient

effects across the surrounding network (Fig. 1b – f; see also Supplementary Fig. 4). Within 132

ms (see Materials and Methods), approximately 15% of non-targeted neurons increased spiking

(Positive Responders, PosR), while fewer than 1% showed decreased spiking (Negative

Responders, NegR), despite high trial-by-trial variability (α = 2.5% significance threshold, one-

tailed Welch's *t*-test; see Materials and Methods). Ongoing, baseline activity was consistent

across all groups (TC, PosR, NegR, and non-responders, NonR), with neurons exhibiting spike

count variability well above Poisson expectations (Fig. 1g, base; n = 54 TC; n = 2,182 unique

neurons, classifying into n = 1,221 PosR + 69 NegR + 7,687 NonR = 8,977 neurons due to

multi-target experiments). Upon stimulation, PosR firing rates and spike count variance

increased markedly remaining well above expectations for a Poisson process as reported

previously[32] (Fig. 1g; response).

112 Novel spikes scale network responses without change in overall network variability

113 Holographically triggered spikes in TC neurons were classified as "novel" because they were

114 not predicted by ongoing activity, as demonstrated by autocorrelation analysis. During the

115 baseline period, ongoing spiking exhibited significant temporal order, with high autocorrelation

116 across all groups (r ≈ 0.2, Fig. 2a, b; base' *vs* base). However, during the response, the

117 autocorrelation for TC neurons (base *vs* resp) dropped to nearly zero (r ≈ 0; p < 10^{-3}, Wilcoxon

Fig. 1 | Single-neuron holographic stimulation drives widespread excitation in primary visual cortex of awake, resting mice. a Schematic of all-optical setup combining holographic stimulation and 2-photon imaging (2PI) in awake, resting mice (V1). (DM: dichroic mirror, M: Mirror, F: filter, PMT: photomultiplier, L: lens, SLM: spatial light modulator). Inset: Mean 2P image in L2/3 showing co-expression of jGCaMP7s and ChrimsonR; one pyramidal neuron (*circle*) was targeted (*flash*). **b** Stimulation (100 ms) of the target cell (TC) evokes relative fluorescence changes (ΔF/F) in soma, dendrites, and nearby neurons (*arrows*). Averaged over n = 150 trials. *Inset*: Mean across 54 TCs. **c** Variable ΔF/F responses to 100 stimulations (*red*) in a single TC with average in black. **d** Example ΔF/F traces (*lines*) and deconvolved spikes (*bars*) from the TC (r*ed*) and 2 responders (nonTC; *black*). **e** Z-scored spike counts in TC and surrounding neurons (n = 166) across 50 trials. *Bottom*: summed trial-wise z-scores showing variable network recruitment (*arrows*). **f** Average z-scored responses in TCs (n = 54), Positive Responders (PosR, n = 1,221), Negative Responders (NegR, n = 69) and Non-significant Responders (NonR, n = 7,687) from 16 experiments (6 mice). *Purple/Red bar*: response/stimulation window. Reconstructed protracted responses are due to deep interpolation (see also Supplementary Figs. 7 & 10). *Base', Base* and *Resp* indicate respective time windows of analyses. **g** The response to single TC stimulation is largely excitatory, while maintaining variability above Poisson-level expectation (*dashed line*). Summary scatterplot of variance vs. mean spike count/132 ms for baseline and response (n = 6 mice, n = 16 experiments). Approximately 15% of neurons exhibit a significant increase in spike count from baseline (*PosR*), while fewer than 1% show decreased firing (*NegR*). For *NonR*, only CPF are shown. *CPF*: cumulative probability function.

rank sum), indicating that stimulation introduced spikes independently from the preceding

baseline, disrupting the temporal organization of ongoing TC spiking. In contrast, temporal

correlations remained stable for PosR and NonR, while the observed decrease in NegR

autocorrelation reflected floor effects on suppressed low baseline activity (Fig. 2b).

Spike count changes in TC in relation to other neuron categories exhibited distinct

differences for baseline and response which we quantified using cross-correlations and scaling

functions. During baseline, TC spike counts (TC_{SC}^{base}) were significantly correlated with all groups

(average cross-correlation <cc> ≈ 0.05; Fig. 2c) and followed a consistent scaling relationship:

$$PosR_{SC}^{base}, NegR_{SC}^{base}, NonR_{SC}^{base} \sim \left(TC_{SC}^{base}\right)^{h^{base}}, \tag{1}$$

with $h^{base} \cong 0.5$ (linear regression; Fig. 2d). This scaling, which covered the majority of baseline

fluctuations from 1 to ~7 spikes/132 ms (transient firing rate increases of 8 – 53 Hz) was found

both for the total spike count per neuron category and per-neuron spike count (Supplementary

Fig. 6a).

During stimulation, only the excitatory response in PosR$_{SC}$ showed scaling,

$$PosR_{SC}^{stim} \sim \left(TC_{SC}^{stim}\right)^{h^{stim}}, \tag{2}$$

with a corresponding exponent of $h^{stim} \cong 0.3$ significantly lower compared to baseline (Fig. 2e, f;

neuron: 0.26 ± 0.09; network: 0.28 ± 0.10; fit ± 95% confidence interval; see Materials and

Methods; see also Supplementary Table 1). In contrast, NonR$_{SC}$ exhibited no scaling, while

NegR$_{SC}$ showed negative scaling (see also Supplementary Fig. 6b). These holographic

manipulations stayed within ongoing fluctuations in cortical firing as our 1–7 added spikes per

132 ms closely match the 1–7 spikes/132 ms seen during baseline (<20% of trials exceeded 7

spikes/132 ms). We also observed qualitatively similar scaling when measuring responses

directly from the calcium transients (see Supplementary Fig. 6c).

The distinct scaling relationships between TC stimulation and the excitatory network

response was mirrored in corresponding positive, zero, and negative cross-correlations between

TC and the other neuron categories (Fig. 2c, right). Novel spikes preferentially scale with

neighboring sub-networks composed of PosR that demonstrate a significantly higher baseline

correlation compared to NonR and NegR (Fig. 2c, left). On average though, positive cross-

correlation found for TC_{SC}^{stim} and $PosR_{SC}^{stim}$ exhibited a relatively weak overlap in evoked and

functional connectivity (see Supplementary Fig. 7; R^2 = 0.018, linear regression; $p < 10^{-5}$, for

the *t*-statistic of the two-sided hypothesis test). Thus, scaling to novel spikes engages both

established and alternative functional connectivities.

Scaling to novel spikes was robust to the significance level (α) employed. At ~10 times

higher α (less strict significance criteria), the scaling exponent h^{stim} for $PosR_{SC}^{stim}$ was still close to

0.2 (Fig. 2g). Despite our partial control of holographical spike generation due to variability from

external (e.g., head movement) and internal (e.g., network inhibition) factors, these scaling

Fig. 2 | Scaling of cortical network responses to novel, holographically triggered spikes in individual pyramidal neurons in awake mice. a Schematic view of activity windows to calculate auto- and cross-correlations. **b** Holographically triggered TC spikes are novel, as evidenced by their zero autocorrelation with ongoing activity. In contrast, autocorrelations remain high for other neuron categories, except NegR, due to spike failure during the response. **c** Ongoing TC spike fluctuations positively correlate with all neuron groups. However, evoked TC spikes selectively correlate positively with $PosR_{SC}^{stim}$, show no correlation with $NonR_{SC}^{stim}$, and correlate negatively with $NegR_{SC}^{stim}$. **d** Scaling for ongoing fluctuations in TC with respect to all other groups. Note limited range in spontaneous TC count fluctuations of <10 spikes/132 ms. Same color code as in panel e. **e** Robust scaling for novel spikes in TC_{SC}^{stim} with respect to $PosR_{SC}^{stim}$ but not $NonR_{SC}^{stim}$ or $NegR_{SC}^{stim}$. **f** Scaling of population spike count for each neuron category separately and combined (*black*). **g** Strong positive scaling for $PosR_{SC}^{stim}$ remains even when significance level α is greatly reduced (larger α). Change in scaling exponent as a function of α per neuron (*solid lines*) and population (*dashed lines*) for each group. **h** PosR are predominantly located within a 100 μm radius of the TC but also distribute throughout the network as α increases. **i** Relative number of neurons per category changes with α. Note that PosR and NegR make up the majority of the observed network at the lowest significance level at which robust scaling is still observed. **j** Fraction of significant responders as function of α. *Solid/dashed lines*: original/FDR-corrected data (see Materials and Methods). Vertical line indicates the level of α used for most analysis. *Shaded areas* (d – f) and error bars (b, c): SD across neurons. When comparing correlations (b, c), 1, 2 & 3 stars indicate $p < 0.05$, 10^{-3} & 10^{-6}, respectively, using Wilcoxon rank sum test.

relationships, nevertheless, derived a posteriori by averaging trials, confirm that driving a single

pyramidal neuron distinctly affects neuronal populations compared to baseline. The scaling is

also robust when running non-denoised (no Deep Interpolation) datasets (see Supplementary

Fig. 8a).

The sparse synaptic connections between local L2/3 pyramidal neurons — estimated at

only 5–8% and largely absent beyond 200 µm[19-23,33] — suggest that among 200 neurons

observed, less than 16 neurons are synaptically connected to a given TC. Indeed, PosR were

consistently found near TCs within a ~200 µm radius as reported previously[4], in line with

expectations for locally connected neighbors providing the strongest spike count increases. On

the other hand, a substantial portion of PosR was distributed throughout the network (Fig. 2h)

and this proportion increased as the significance threshold (α) was lowered (higher α). At the

lowest significance level examined, nearly half of all recorded neurons showed significant

positive responses, with an average scaling exponent of h^{stim} ~ 0.22.

Recruitment of both PosR and NegR from the NonR pool (Fig. 2g–i) indicates that novel

spikes are transmitted through the network via multi-synaptic pathways, triggering both

increases and decreases in spiking, similar to what has been observed when stimulating groups

of pyramidal neurons together[7]. When correcting for multiple comparisons[34,35], we find that our

significance threshold (α = 2.5) underestimates the number of significant PosR neurons by a

factor of ~1.7 times, while NegR become statistically non-significant. This demonstrates that

novel spikes primarily drive a net excitatory response across the network (Fig. 2j). Notably,

activating a single pyramidal neuron engages a large number of neurons spread across a wide

area – an outcome that contrasts with what one would expect based on anatomical connections

and synaptic physiology in activity regimes dominated by spontaneous fluctuations.

The robust scaling seen during TC stimulation may be partly due to a reduction in

spiking variability, known to occur during sensory stimulation[36] and often measured using the

Fano factor (FF), which compares the variance to the mean of spike counts. We found that

repeated holographic stimulation did lead to lower FF in TC neurons, consistent with more

regular input. But for PosR and NonR neurons (defined at the original α = 2.5% threshold), FF

did not change, even when we adjusted for differences in their average mean spike count[37] (Fig.

3a, b). Likewise, there was no change in FF observed for PosR and NonR population responses

(Fig. 3c). To test whether variability in TC spike triggering masked changes in FF for PosR and

NonR, we analyzed subsets of trials with high or low TC spike count, for overall network activity

and response timing. Yet, FF stayed consistently high and unchanged in both subsets under all

conditions (Fig. 3d; Supplementary Fig. 9).

Our findings show that novel spikes from a single pyramidal neuron trigger strong,

scalable responses in many nearby neurons, even though overall network activity remains

dominated by fluctuations.

Fig. 3 | Network variability remains unchanged despite holographically triggered spikes. a Example TC with trials split by low vs. high spike count. *Top*: Base spike distribution defines 91st percentile and 2× threshold. *Bottom*: These thresholds are applied to classify stimulation trials. **b** The Fano Factor (FF) remains unchanged for PosR and NonR despite significant scaling in $PosR_{SC}^{stim}$ (α = 2.5%). *Transparent bars*: Mean spike count uncorrected FF (see Materials and Methods). **c** Population FF also shows no change across categories. **d** Summary statistics of mean TC trials separated by low and high target counts. Note the remaining high variance of PosR despite the large drop in TC variance. *Error bars* (b–d): SD across neurons. 1 star indicates p < 0.05, Wilcoxon rank sum test.

Precise origin information from novel spikes spreads transiently across the cortical

network

We next examined whether the broad network responses to novel spikes also encoded

information about their point of origin. Across seven experiments, we stimulated up to 10 TCs in

a semi-random sequence (each for 100 ms with 2–10 s between trials; Fig. 4a). We found that

the origin of each novel spike event could be decoded with high accuracy from the activity of

non-stimulated neurons (nonTC) using two machine learning approaches: XGBoost[38] (XGB)

and Random Forest[39,40] (RF) (see Materials and Methods).

Both algorithms achieved ~60% decoding accuracy and F1-scores exceeded 90% for

both algorithms when utilizing the full population of nonTC (Fig. 4b, left; Supplementary Fig.

10a). Decoding performance positively correlated with the number of trials in which the TCs

were successfully activated (Supplementary Fig. 10b). For comparison, spontaneous spikes in

TC occurring within 132 ms during ongoing activity (base) could also be decoded although with

significantly lower accuracy (Fig. 4b, right; see Materials and Methods). Remarkably, it took the

removal of 10–70% of the top-performing decoding neurons for accuracy to drop to chance

levels (Fig. 4c). In a representative experiment (Fig. 4d), F1-scores of TCs dropped to chance

levels after removing >70% of the best-encoding neurons. On average, 20–30% of the best-

encoding neurons were sufficient to achieve high decoding accuracy, regardless of the

algorithm used (Fig. 4c; Supplementary Fig. 10c).

Positively responsive neurons (PosR, defined at $\alpha = 2.5\%$) played a major role,

contributing 30–60% of the decoding power (Fig. 4d). To confirm that decoding wasn't driven by

direct stimulation of nearby cells, we performed a spatial exclusion analysis. Excluding neurons

within 10 μm of TCs had no effect on performance, but gradually removing more distant

neighbors led to a steady drop in accuracy (Fig. 4e), confirming a spatially distributed response

in the network to stimulation.

Individual neurons could significantly respond to stimulation from up to five different TCs,

indicating overlapping functional subnetworks capable of encoding multiple spike origins (Fig.

4d, inset). Importantly, this information was short-lived — decoding accuracy dropped rapidly

after the stimulation ended, in line with the brief activity profiles observed throughout the

network (Fig. 4f, cf. Fig. 1f; analysis performed on a single frame – see Materials and Methods).

We note that elevated decoding before the stimulus onset was attributed to temporal averaging

(132 ms window) and Deep-Interpolation, which enhances signal quality but slightly blurs

response timing (see Supplementary Fig. 10d). Qualitatively similar results were obtained using

the calcium transient directly (Supplementary Fig. 10e).

Together, our results show that superficial layers of cortex can briefly and reliably

encode when and where a novel spike originates. This supports the existence of a robust,

distributed coding mechanism for processing unexpected inputs in real time across the network.

Novel spike encoding persists across fluctuation-dominated, network state of parabolic

avalanches

Our findings so far demonstrate the remarkable impact of just a few action potentials amid the

thousands of spikes generated every second in the awake brain's cortical neighborhood. Given

our 0.45×0.45 mm field of view and ~170 imaged neurons (*cf*. Fig. 2i), we estimate our sampling

to be ~5% of this neighborhood[19]]. On average, we observed ~23 PosR per TC ($\alpha = 2.5\%$),

Fig. 4 | Origin of novel spikes propagates transiently and broadly in L2/3 of awake, resting mouse cortex. a Example field of view showing targeted TCs (circled) and surrounding network (mean baseline activity). **b** *Left*: Decoding TC identity using XGBoost (XGB) or Random Forest (RF) achieves highly above-chance accuracy and F1 scores (n = 7 experiments; chance: 10–25%). *Right*: Accuracy when decoding from Base compared to Resp (see Materials and Methods; paired *t*-test, p = 0.012). **c** Distributed coding: a large fraction of the top-ranked neurons (10–70% or 10–100, by Shapley value) must be removed to reduce accuracy and F1 to chance (dashed lines; A–G experiments with number of TC indicated). **d** Likelihood (*L*) that dropped neurons are PosR. *Top*: L (*blue shade*) and F1 scores (*grey*: single TC; *black*: mean) for a single experiment. *Bottom*: average ± SD across experiments. *Inset*: Neurons can respond to multiple TCs—distribution of PosR counts per neuron. **e** Spatial restriction: excluding neurons by radius around TCs confirms localized stimulation. *Top*: Drop in overall accuracy when excluding all neurons within a certain distance from all TCs. *Bottom*: Number of neurons removed by this procedure. **f** Novel spike information is transient: Decoding accuracy rises above chance only when the response analysis window overlaps with stimulation (0–100 ms; *red shaded area*). *Top*: Single experiment ± Deep Interpolation (22-ms window). *Bottom*: All experiments (22-ms window, Deep-Interpolated).

which would translate to more than 450 neighboring neurons significantly *increasing* their firing

within tens of milliseconds in response to our holographic stimulation. This estimate, however, is

likely highly conservative as increasing α (lowering the significance criteria) further increases the

number of neighboring neurons with detectable changes in spiking, as confirmed by our

decoding results. This adjustment increasingly recruits both positive and negative responses

from the pool of previously non-significant responders (it is important to note that we are likely

heavily underestimating the number of NegR[25,26], given the constraints to detect significantly

lower spike counts from neurons that fire very sparsely; see Materials and Methods).

Furthermore, for many neurons, the spike count variability during novel spike encoding

exceeded what would be expected from a simple Poisson process (Fig. 1), with Fano factors

comparable to those observed during ongoing spontaneous activity (Fig. 3). This high variability

extended across neuronal subtypes, including positively responsive neurons (PosR). To better

characterize this fluctuation-dominated regime, we analyzed its statistical properties in the

context of novel spike encoding.

Holographic stimulation frequently occurred during periods of heightened ongoing

activity (Fig. 5a), defined by continuous population spiking above a low threshold. These active

periods followed heavy-tailed distributions in both size (total spike count) and duration,

consistent with previously described neuronal avalanches[29,41] (Fig. 5b). After correcting for

subsampling using temporal coarse graining, we found that the average avalanche size scaled

quadratically with duration — a key feature of parabolic avalanches[29,41,42]. These parabolic

events included avalanches lasting up to ~0.7 seconds, which accounted for 87% of all

avalanches and about 20% of spikes during spontaneous activity (Fig. 5c, d). Longer

avalanches (>1 s) showed near-linear scaling, suggesting activity too widespread to be fully

captured within the imaging field of view[29,41,43].

Despite these dynamic and large-scale fluctuations, decoding the origin of novel spikes

remained highly accurate across avalanche activity regimes. When baseline activity was low (θ

< 0.3 spikes/132 ms), stimulus-evoked spike counts scaled modestly; when baseline activity

was higher (θ > 0.3), the scaling approached a value of 1, indicating strong coupling between

pre-stimulus and evoked activity (Fig. 5e). This allowed us to classify activity into low- and high-

baseline regimes, both of which prominently featured parabolic avalanche dynamics (Fig. 5f).

Importantly, decoding accuracy differed by less than 4% on average between these

regimes, regardless of the threshold used (Fig. 5g, h), and response scaling remained

consistent across both states (Supplementary Fig. 11a).

Fig. 5 | High decoding accuracy of novel spikes occur during scale-free avalanche dynamics despite highly fluctuating levels of baseline activity. a *Top*: Population activity in visual cortex of the awake mouse exhibits large fluctuations during quiet resting. *Bottom*: Zoomed in view of population activity in response to 3 (i – iii) holographic stimulations of a single TC. *Vertical lines*: TC stimulation. *Red line*: TC spike count. *Grey stripe*: Population threshold for avalanche detection. Shown is the population spike count (excluding TCs) per 22-ms frame. **b** Suprathreshold contiguous fluctuations in population activity exhibit the characteristic hallmarks of neuronal avalanches. Avalanche size (*left*) and duration (*right*) distributions as a function of the temporal coarse-graining factor *k*. **c** Scaling of avalanche dynamics as a function of temporal coarse-graining *k*. **d** Approximately 87% of avalanches are parabolic in nature, with durations <0.7 s, containing about 18% of thresholded spikes. Summary scaling of parabolic avalanches is shown for durations up to ~0.7 s (*k* = 4). **e** Mean response spike count scaling as a function of mean baseline spike count for non-TC neurons. This analysis reveals a LOW baseline activity regime (mean baseline spike count < ~0.3), in which responses are mostly independent from base activity, distinct from a HIGH baseline activity regime in which response activity can be predicted from base. **f** Mean avalanche duration is more than double for avalanches that start during LOW base trials compared to HIGH. 86% of all LOW avalanches are parabolic (scaling of $\chi = 2$), whereas ~54% of HIGH avalanches are parabolic. **g** Decoding accuracy remains largely unchanged when obtained using only LOW or HIGH baseline activity regimes, regardless of the threshold used to define the two regimes. Overall accuracy for LOW base trials (*solid lines*) and HIGH base trials (*dashed lines*) for each dataset (same color code as in Fig. 4), as function of the threshold used to define LOW and HIGH base trials. **h** Difference in accuracy for LOW and HIGH trials is small (with LOW trials having slightly higher accuracy), regardless of the threshold used to separate them.

These results suggest that cortical networks in vivo operate within a fluctuation-

dominated regime where variable responses to local perturbations scale in line with

expectations from critical dynamics[28,44], which we further explore in the following section.

Model simulations reveal observed experimental perturbation responses to be in line

with those from critical networks

To study how networks with varying internal excitability respond to local stimulation, we

simulated a two-dimensional network of 90,000 excitatory units with local, radial connectivity

P_{conn}(r) and three-state dynamics (rest, active, refractory) (Fig. 6a). Analysis focused on a

central field of view of 456×456 µm^2 with 1,440 neurons, including 10 target cells (TCs), to

mimic experimental access and reduce edge effects.

Each resting unit could be activated either by an external Poisson input at constant rate

P_{Poiss}, or by an active neighbor with transmission probability P_{trans}. The system exhibits dynamics

of the directed percolation universality class, with a phase transition centered around a critical

state — between a subcritical phase (where activity fades without input) and a supercritical

phase (where activity can self-sustain).

Near criticality, the branching parameter σ — the average number of near future

downstream spikes per originating spike — is close to 1. We define $P_{trans} = \sigma/\bar{k}$, where $\bar{k} \approx 43$

is the average number of connections per unit. Adjusting σ, moves the system in and out of the

critical state. Empirically, the critical point was near σ = 1.0066, slightly above 1 to compensate

for propagation loss (e.g. spike collision) in the network.

Figure 6b shows example traces of ongoing network activity under subcritical, critical,

and slightly supercritical conditions, highlighting both the increase in overall activity and the

emergence of larger fluctuations as the network approaches criticality. This shift is quantified in

Fig. 6c, where the relationship between network activity and external drive (P_{Poiss}) reveals the

critical transition. In the subcritical regime (σ << 1), inputs spread weakly, and network activity

increases linearly with external drive, yielding a slope $\gamma \approx 1$ (see Refs.[45,46]). In the supercritical

regime (σ > 1.0066), even low input leads to network saturation, resulting in $\gamma \approx 0$. At the critical

point, the system produces scale-invariant responses – external spikes can trigger cascades of

all sizes – producing a power-law relationship with an intermediate slope $0 < \gamma < 1$. For our 2D

network, the predicted critical exponent is $\gamma = 0.285$ (see Ref[28]), as shown in Fig. 6c (magenta

curve). To match experimental conditions, we adjusted P_{Poiss} in each simulation to reproduce

average baseline activity observed in our in vivo experiments.

Fig. 6 | Simulated networks near criticality replicate experimental response scaling and decoding. a Simulation setup: A 2D network of 90,000 neurons with local spatial connectivity ($\bar{k} \approx$ 43 connections/neuron) receives external Poisson input. A central 1,440-neuron region (FoV; 456 x 456 μm^2) with 10 TCs was analyzed to mimic experimental conditions and reduce edge effects. An external Poisson drive with fixed rate and active neighbors with fixed probability allow neurons to sequence through quiet, active, and refractory states. **b** Network activity under subcritical, critical, and slightly supercritical conditions shows TC (*red*) and non-TC (*black; 50 neurons randomly selected*) spiking during repeated TC stimulation (*red bars*). *Bottom*: Color coded summed activity of from the whole FoV. **c** Response scaling as a function of spike transmission probability ($P_{trans} = \sigma/\langle k \rangle$; σ is the branching parameter) reveals a power law (exponent ~0.285) at the critical point (*magenta*). Subcritical (*blue*) and supercritical (*green*) regimes show linear and self-sustaining behavior, respectively. **d** Distribution of mean spike counts during baseline. Note decrease in activity fluctuation range during baseline for subcritical systems (*blue*). **e** Mean response of PosR as function of TC spike count. Close to criticality, response curves follow a power law profile similar to the one observed in experiments (cp. Fig. 2e). **f** Power law exponents for the response curves of PosR for varying branching parameter. Note that the exponents approach the value observed in experiments (~0.28) close to criticality. **g** Decoding accuracy is highest close to criticality and drops to chance only after removing a very large fraction of the best predictors. Accuracy as function of number of best decoding neurons dropped for varying branching parameter. **h** F1 scores increases with TC's (*circles*) output strength. Output strength is measured by the sum of all probabilities that a single spike in the TC, in a quiet network, would propagate to each of its direct neighbors, secondary neighbors, etc. up to order 5. It estimates the expected number of spikes that each spike in the TC would generate on average, ignoring refractory period and potential collisions (see Materials and Methods).

As shown in Fig. 6d, the distribution of ongoing firing rates is narrow in subcritical

conditions but broadens near the critical point, reflecting a fluctuation-dominated regime. When

measuring the response on PosR as function of TC spike count, we observed power-law

scaling, with the slope approaching the one obtained experimentally ($h^{stim} \sim 0.285$) at criticality

(Fig. 6e, f). Interestingly, when considering only trials in which the network had a low amount of

activity preceding the stimulation, the slope of the scaling responses crosses the value 0.285

close to criticality, whereas considering only trials with a high amount of activity we observe that

the highest slope happens at criticality, albeit at a smaller value, as in the experiments

(Supplementary Fig. 11b, c). Importantly, classification accuracy improved sharply from

subcritical to critical regimes and decayed gradually as increasing numbers of neurons were

removed (Fig. 6g), again matching experimental observations. Furthermore, decoding success

was higher for targets that were more well-connected to the network (output strength estimates

the expected number of spikes each TC spike would generate in a quiet network; see Materials

and Methods) and was maximized at criticality (Fig. 6h).

Despite higher activity fluctuations, this critical regime maximized the number of positive

responders (PosR) to TC stimulation (Fig. 7a). As in the experimental results, the fraction of

cells that are PosR is maximal around TCs and decay over larger distances (Fig. 7b). At

criticality, about 93% of neurons receiving direct input from TCs were classified as PosR.

However, only about 16% of all PosR had a direct connection from TCs, showing the

importance of multi-synaptic pathways in the critical network. This reveals a unique "mixing"

effect at criticality: while a TC's average output strength is higher, its functional spread—the

range of PosRs it engages—becomes much broader, reflecting the greater diversity seen in

critical states (Fig. 7c).

To further examine how the network's dynamical regime affects information spread, we

measured the number of PosR for each TC in the subcritical network ($\sigma = 0.5$) and compared it

to that of the critical network ($\sigma = 1.0066$). Even though the two networks have the same

structure, the number of PosR for each TC is uncorrelated between them (Supplementary Fig.

12). This means that measuring a TC's effect under subcritical conditions (lower excitability)

gives little or no indication of its impact near criticality.

Together, these results strongly support the idea that novel inputs are most effectively

propagated in networks operating near criticality, where activity is governed by fluctuations and

scale-invariant dynamics.

Fig. 7 | PosR are close and relate to output strength of TCs. a The mean number of PosR per target peaks close to the critical point of the network. **b** Average fraction of neurons that are classified as PosR as function of distance from TC. As expected, the fraction is highest close to TCs and decreases as the network deviates from criticality. **c** Number of PosR of each TC (*circles*) as a function of the TC's output strength. *Right*: Covariance matrix ellipses (*shaded area*; 2.5 s.d.) and center of mass (*lines*; mean ± s.d.) for the scatter plot on the left.

Discussion

Efficient communication of novel neural signals is vital for processing surprises and guiding future actions. Using an all-optical approach allowed us to precisely induce action potentials holographically in identified neurons and simultaneously measure its effect in the network at cellular resolution and high fractional sampling of the neuronal population in the awake, quietly resting mouse. Alternative perturbation approaches, such as extracellular microstimulation lack spatial selectivity[47], or as in the case of intracellular patching[2], are limited in capturing the network response. Here, we demonstrated in awake animals that introducing spikes in a single L2/3 pyramidal neuron rapidly activates widespread, net excitatory responses in the surrounding cortical network, encoding the novel stimulus' origin. Remarkably, these responses scaled robustly and were efficiently decoded amidst ongoing, large fluctuations in network activity organized as scale-invariant, parabolic neuronal avalanches. Our scaling exponent of ~0.2–0.3 in response to novel spikes was captured in a 2-dimensional network with

critical dynamics allowing for highest decoding accuracy and engaging large neighborhoods in

encoding stimulus origin. This high susceptibility and scaling to local perturbations[27,28], suggests

critical dynamics as a cornerstone of cortical processing, offering key insights into brain

function.

Our findings challenge a longstanding discussion on cortical coding fueled by the

intricate statistical anatomy of the cortex and its synaptic connections. Pyramidal neurons in

associative layers establish few synaptic connections with their neighbors[19-23]. These excitatory

connections are relatively weak, prone to failure, and decay within milliseconds[22,48]. The

combination of sparse and weak connectivity makes it unlikely that a spike in a single neuron

reliably triggers activity propagation in the surround network. To address this, most cortical

codes rely on activity patterns that amplify synaptic impact on postsynaptic pyramidal neurons.

For instance, coincident activation of numerous synapses from diverse neurons

converging onto a postsynaptic target enhances the likelihood of activity propagation, favoring

coding models based on local synchrony, such as synfire chains[49,50], waves[51,52] or phase-locked

oscillations[53,54]. Alternatively, repeated or burst firing of individual neurons[12,55], strong synaptic

connections[56], subthreshold priming by background activity[15,57] supported by subcellular

amplification mechanisms such as short-term synaptic facilitation[58,59] and non-linear dendritic

processing[60] can dynamically increase individual connection strength supporting rate-based

coding.

Our findings do not support a dichotomic separation into synchrony vs. firing rate

separation to secure propagation of neuronal activity (see also[24,61]). We show robust scaling of

novel spikes within the cortical neighborhood from single to multiple action potentials delivered

within 100 ms without an obvious threshold above which propagation is facilitated. This

suggests that neither tight synchrony nor repeat firing is strictly necessary for effective encoding

of novel spikes[62]. This conclusion is in line with findings that spike counts over 100 ms predict

behavior independent of considerations of synchrony at the millisecond level[63].

By emphasizing the sufficiency of a few action potentials in cortical codes, our results

thus challenge existing paradigms and highlight a minimalist yet robust mechanism for cortical

information processing supporting the prominence of sparse, single action potential firing found

in vivo for associative layers[64]. Furthermore, the recruitment of significant responders is most

likely composed of a mixture of direct (excitation, inhibition) as well as indirect (disfacilitation,

disinhibition) local circuit interactions[4,7], while preserving the general scaling relationship (cf. Fig.

2). Simulations have found that in such a fluctuation-dominated but relatively balanced regime,

a reduction in membrane conductance due to increased synaptic bombardment still enables

neurons to generate few spikes in response to rapid changes in synaptic input[17].

The importance of critical dynamics for sensory cortical networks has been argued from

the point of view of optimal decoding of stimulus intensities through maximization of the dynamic

range[46]. In that context, a critical network is expected to produce response-input curves that

scale as a power law, with exponents that are typically much smaller than one[28]. These low

exponents allow the network to present distinct responses for a wide range of stimuli. On the

other hand, the difference between these distinct responses gets smaller as exponents

approach zero, indicating the possibility of an optimal value depending on the necessity of

having a larger dynamic range versus the need for a more precise differentiation between

responses. The study of psychophysics has demonstrated that psychological value scales with

stimulus intensity as a power law with <1 exponents, a result known as Stevens' law of

psychophysics[65]. Therefore, it has been argued that the origins of Steven's low exponents for

response curves might originate precisely at this critical networks' ability to amplify low-intensity

stimuli and thus respond to a wide range of stimulus intensities[66]. We were able to show that *in*

vivo sensory cortical networks do display low-exponent (see Supplementary Tables 1 & 2),

power-law behavior for their response to varying stimuli intensity (as proxied by the number of

spikes evoked in a target cell during stimulation). This is in line with predictions from critical

dynamics, with the observed exponents closely matching the expected value (0.285) for a two-

dimensional directed percolation system[28]. The expected exponent for the response scaling of a

3D network is 0.45, whereas higher dimensional systems would show an exponent of 0.5. Note

that our observed ~0.5 slope during baseline activity does not represent a response of the

network: since neuronal firing during ongoing activity is correlated, that relationship is more

indicative of autocorrelation in the network as a whole. Our measured scaling exponent thus

emphasizes the intriguing finding that superficial layers 2/3, defined anatomically as an

extended 2-dimensional sheet, functionally also operate more closely to a 2-dimensional critical

network.

Our findings of maintained high Fano factor in evoked responses is in line with the

expectation for a critical system, though, squares with the quenched variability demonstrated in

response to sensory stimuli[36]. This difference may arise because single-neuron stimulation

introduces a smaller perturbation than sensory stimulation, which in the awake state is

dominated by inhibition[25], affecting response variability. Early studies in anesthetized animals[18]

(e.g. in the visual cortex of cats) demonstrated that trial-by-trial variability in stimulus responses

can be approximated as average response overlaid on a correlated, variable background. In the

awake state, our observation of consistently high FF values for responses in the presence of

scale-invariant parabolic avalanches indicates the need for a conceptual approach that

bypasses the reliance on averaging. Avalanches emerge in superficial layers of cortex when

animals recover from anesthesia[67,68], exhibit high neuronal selectivity[29,69] and spatiotemporal

order[70-74] characterized by power law statistics with exponents ≤ 2 that is, averages are not

defined and variability is high. Yet, we demonstrate that decoding of novel spikes is similar

across the fluctuation dominated avalanche regime.

Our decoding analysis provided important insights about the fine-grained nature of this

communication of novelty. First, strongly decoding neurons were found in the immediate

neighborhood of TCs supporting a spatial weighing in the response, in line with expectations

from the local, synaptic connectivity statistics of pyramidal neurons. On the other hand, a large

fraction of neurons with high decoding contribution did not fall into the PosR category or were

located at long distances from the TC, supporting the contribution of indirect pathways.

Similarly, in our simulations the number of identified PosR greatly surpassed that of direct

postsynaptic neurons of each TC, further suggesting the importance of indirect pathways to the

coding of novel information. In fact, only about 16% of PosR had a direct connection from TCs.

This number could be even higher for the brain, since the presence of inhibitory connections

(absent in our simulations) could facilitate certain pathways in detriment of others, while also

requiring stronger excitatory-to-excitatory connections in order to stay in a balanced propagation

regime. This also de-emphasizes reliance on distinct, non-overlapping network connectivity.

The ability of PosR neurons to contribute significantly to the decoding of multiple TCs is

in line with the spatial overlap of neuronal populations encoding different stimuli and their

temporal separation. Our findings align with the notion that cortical networks are organized to

maximize coding efficiency and redundancy, where overlapping populations can fluctuate in

their response patterns to encode different stimuli[75,76]. The rapid decay found for the encoding

of novel spikes further underscores this dynamic, ensuring that novel signals do not disrupt

ongoing network activity but still convey novel information effectively within the constraints of the

temporal and spatial coding architecture described for e.g. visual stimuli of the primary visual

cortex in the awake mouse[77].

Our results reveal the communication of single spikes within the scale-invariant, critical-

state dynamics of cortical networks, where synchronized fluctuations are neither overly

constrained nor entirely random. We propose this remarkable sensitivity of the mammalian

cortex, to efficiently disseminate novel information from individual neurons, underscores the

fundamental role of critical dynamics in cortical communication and computation.

Materials and Methods

Animal surgery

All procedures were approved by the NIH Animal Care and Use Committee (ACUC) and experiments followed the NIH *Guide for the Care and Use of Laboratory Animals*. Mice were obtained from Jackson labs, bred inhouse with C57BL/6 backgrounds (Jackson Laboratory) under a reversed 12:12 hr light/dark cycle. Chronic 2PI imaging in adult (>6 weeks) mice was enabled by using a head bar in combination with a cranial window placed centered at ~2.5 mm from the midline (right hemisphere) and ~1 mm rostral to the lambdoid suture and consisting of a stack of circular glass cover slips using established protocols[78]. Mice were injected with a 7:1 mixture of a viral construct to express jGCaMP7s[79] and ChrimsonR[80] in pyramidal neurons using the CaMKII promotor. A total of 2 – 3 injections of virus (100 – 400 nL; <1 µL in total; 10^{13} vg/mL; pGP-AAV-syn-jGCaMP7s-WPRE AAV9 & pAAV-CamKIIa-ChrimsonR-mScarlet-KV2.1, Addgene) were administered into the right hemisphere. Chronic 2PI started after >2 weeks in identified V1 at a depth of ~100–200 µm.

Identification of V1 maps

Retinotopic maps of V1 and higher visual areas (HVAs) were generated for all mice prior to recording using published protocols[81,82]. Briefly, awake, head-fixed mice faced with their left eye a 19" LCD monitor placed at 10 cm distance and tilted 30º towards the mouse's midline. Using Psychophysics toolbox[83], contrast-reversing, spherically corrected checkerboard bars were drifted across the screen vertically (altitude) and horizontally (azimuth) for each of the four directions (30 repeats per direction). Simultaneous wide-field imaging (Quantalux, Thorlabs) captured jGCaMP7s fluorescence, which was averaged for each direction. Altitude and azimuth phase maps were calculated by phase-wrapping the first harmonics of the 1D Fourier transform for each of the four averages and subsequently subtracting the maps of the opposite directions[82]. Sign maps were generated by taking the sine of the angle between the gradients in the altitude and azimuth maps and processed[81]. Borders were drawn around visual area patches and overlaid onto anatomical reference images to identify V1.

Holographic stimulation of single neurons

Holographic stimulation[84] was done by manually selecting individual pyramidal neurons as target from the two-photon image of the field of view in both red and green channels. Neurons that express the opsin (ChrimsonR, with mScarlet tag) were selected for stimulation. Based on

the selected targets, a map of top-hat patterns with diameter of 10 µm was used to generate the

hologram on the Spatial Light Modulator (SLM, Meadowlark, LCoS high resolution 1920×1152

phase modulator). Light from the laser/OPA system (Carbide/Orpheus-F – Light Conversion, λ =

1064 nm, repetition rate = 600 kHz, pulse width <100 fs) was expanded (4x) to fill the area of

the SLM (8 mm × 8 mm). The hologram on the pupil plane was shrunk to fit the size of the

Galvo-Galvo scanners (~3 mm diameter) and form the desired pattern of top hats after the

objective lens (Nikon 16X, working distance = 3mm and effective focal length = 12.5 mm) on the

selected targets. Each target neuron was continuously stimulated by the top-hat beam shape for

100 ms for a total of ~100—150 trials per experiment, with an interval of 2 s (10 s in some

recordings) between trials. Power at target neurons ranged from ~2.5 – 10 mW. It is important

to highlight that no artifact from this stimulation was detectable on our imaging with our

combination of GECI/opsin, stimulation wavelength and the low power required to stimulate

single targets (see Supplementary Fig. 4).

2PI imaging, pre-processing pipeline and meta data collection

For standard 2PI, images were acquired by a scanning microscope (Bergamo II series, B248,

Thorlabs Inc.) coupled to a pulsed femtosecond Ti:Sapphire 2-photon laser with dispersion

compensation (Chameleon Discovery NX, Coherent Inc.). The microscope was controlled by

ThorImageLS and ThorSync software (Thorlabs Inc.). The wavelength was tuned to 940 nm to

excite jGCaMP7s. Signals were collected through a 16× 0.8 NA microscope objective (Nikon).

Emitted photons were collected through a 525/50 nm band filter using GaAsP photomultiplier

tubes. The field of view was ~450×450 µm. Imaging frames of 512×512 pixels were acquired at

45.527 Hz by bidirectional scanning of a 12 kHz Galvo-resonant scanner. Beam turnarounds at

the edges of the image were blanked with a Pockels cell. The average power for imaging was

<70 mW, measured at the sample.

The obtained tif-movies in uint16 format were rigid motion-corrected via the python-

based software package '*suite2p*'[85]. Registered images were further denoised using machine-

learning based, deep interpolation[86] (see below) and then semi-automatically processed by

suite2p for ROI selection (we performed visual curation of the automatically-selected ROIs to

ensure only neurons were included in the analysis; this is done so processes, such as the

dendrites identified in Fig. 1b, are not part of the analysis and therefore cannot be classified as

significant responders or contribute spikes to response curves or other measurements) and

fluorescence signal extraction. For each labeled neuron, raw soma and neuropil fluorescence

signals over time were extracted for each ROI. Spiking probabilities were obtained from

neuropil-corrected fluorescence traces ($F_{corrected} = F_{ROI} - 0.7*F_{neuropil}$) via MLspike

(https://github.com/MLspike), utilizing its autocalibration feature to obtain unitary spike event

amplitude, decay time, and channel noise for individual ROIs.

Deep-interpolation

Deep-interpolation[86] (Deep-IP; https://github.com/AllenInstitute/deepinterpolation) removes

independent noise by using local spatiotemporal data across a noisy image stack of $N_{pre} + N_{post}$

frames to predict, or interpolate, pixel intensity values throughout a single withheld central

frame. The deep neural network is a nonlinear interpolation model based on a UNet inspired

encoder-decoder architecture with 2D convolutional layers where training and validation are

performed on noisy images without the need for ground truth data. As described previously in

detail[29], after rigid motion correction, individual denoised frames were obtained by streaming

one 60-frame ($N_{pre} = N_{post} = 30$ frames) registered, image stack through the provided Ai-93

pretrained model[86] for each frame to be interpolated. At an imaging rate of ~45 Hz, these 60

frames correspond to a combined ~1.3 seconds of data surrounding the frame to be

interpolated. This process did not alter qualitatively our main results (see Fig. 4f &

Supplementary Fig. 8).

Postprocessing pipeline

Our postprocessing pipelines were custom-written in Matlab (Mathworks) and Python

(www.python.org). Some routines utilized NumPy (https://numpy.org/) and Matplotlib

(https://matplotlib.org/).

Stimulus response measurement

Unless stated otherwise, spike count over a window of 6 imaging frames (~132 ms) starting at

the stimulation onset was used as a measure of the response to holographic stimulus of each

target cell for all cells. That window was chosen because it includes the full stimulus duration

(100 ms) and one extra frame after the stimulus offset, for which the increased activity in TC can

still be observed (see Fig. 1f). This was compared to the baseline count, defined as a 6-frame

window starting ~264 ms before stimulus onset. Due to difficulties and limitations of spike

deconvolution of calcium traces, some parts of the time series resulted in artificially high spike

rates. To avoid artificial increase of variability measures due to these artifacts, we introduced an

outlier removal procedure in which trials with spike count above 10 standard deviations over the

mean are removed from the analysis. This procedure resulted in <0.1% trials removed, across
all neurons for all experiments.

Target selection and exclusion

Up to 25 target cells were visually selected based on expression and location in the field of view
(cells close to the edges were avoided) for each experiment. Not all targets were responsive to
stimulation. Targets in which fewer than 20% of the trials evoked a spike count above the 91%
percentile of its baseline count were excluded from the analysis. This procedure resulted in 54
target cells across 16 experiments from 6 mice being considered out of 193 targets attempted
across 33 experiments from 11 mice. We estimate an average of 46.3 ± 17.9 ChrimsonR-
labeled cells per field of view, which translates to an average density of labeled cells of ~$2.3×10^{-2}$
cells per 100 μm^2. This sparse labeling facilitates the holographic stimulation of individual
targets (see examples at Supplementary Fig. 1).

Definition of significant responders

Cells with significant response to each target's stimulation were defined as those whose spike
count distribution was significantly different from its baseline count distribution. Specifically, we
used a z-test to compare the stimulus response spike count to the average and standard
deviation of the baseline count for each non-target cell. A cell was defined as a positive
responder if the z statistic is above the highest α = 2.5% (unless stated otherwise) of the
distribution and as a negative responder if the z statistic is below the lowest α = 2.5%.
Otherwise, the cell is classified as a non-significant responder. We have also corrected for false-
discovery rate (FDR), by employing the matlab function mafdr [34,35], with an FDR of 2.5%. After
FDR correction, the number of positive responders increased in relation to the one obtained at α
= 2.5%, whereas the number of negative responders dropped to nearly zero. It is important to
highlight that, given a low baseline activity rate for most cells, with an average count in the
relevant window for our analysis near zero, it is technically challenging to define negative
responders in this scenario using spike data. Other groups have shown suppressed neurons
upon select pyramidal neuron stimulation using fluorescent traces, for which detection of
significant decreases during stimulation might be easier[7,87].
579   This procedure is done for each target stimulated in an experiment separately.
Therefore, in experiments where multiple targets were selected, each non-target cell is tested
for significance of its response to each target's stimulation, possibly being a significant
responder for some targets and non-significant responder to others (see Fig. 4d).

Fano Factor calculation and significance

Fano Factor was calculated as usual, for each cell separately, by dividing the variance of their

spike count by the mean of their spike count across trials. To correct for changes in firing rate

across conditions[37], we introduced a procedure in which the window used to measure the spike

count during stimulation varied from 1 to 9 frames (~22 ms to 198 ms), starting at the onset. The

window length was determined by minimizing the normalized difference between the mean

baseline and stimulus counts. Cells for which this difference was above 20% were excluded

from the analysis. This procedure resulted in less than 30% of cells removed, on average,

across the non-target cells and an average of 65% of targets removed (minimum of 13% or 7

targets considered). Values were reported as mean ± error over the mean across cells, unless

stated otherwise. Significance of Fano Factor changes across conditions was obtained by

employing the Wilcoxon rank sum test at a significance level of 0.05 (p-values reported in

Results section and figure legends).

Trial separation and controls

To control for different conditions affecting the variability, the Fano Factor was recalculated and

compared using a subset of trials. To control for target cell response variability affecting non-

target cells, we separated trials into low target count trials and high target count trials. We

defined low count trials as those for which the target cell produced a spike count during

stimulation that was above the 91st percentile of the baseline count distribution and up to twice

that number of spikes, whereas high count trials were any trial with count above that limit. To

control for the influence of ongoing fluctuation of the network activity we separated trials into low

and high population baseline count, defined as trials for which the population count during

baseline was lower than the 25th percentile of its distribution or higher than 75th percentile,

respectively. To control for temporal influence of the responses to the stimulus, we defined two

consecutive 6-frame windows, centered at the peak of the target's average response to the

stimulus (2—3 frames after onset), as early and late response windows.

Correlation analysis

Spike counts across trials were compared by computing correlations across time

(autocorrelation, calculated using the spike counts over pairs of windows separated by a

temporal lag of 132 ms) and across cells (comparing spike counts of targets *vs* non-targets).

Values are reported as mean ± error over the mean. Significance of correlation changes across

conditions was obtained by employing the Wilcoxon rank sum test at a significance level of 0.05

(p-values reported in Results section and figure legends).

Decoding Analysis

Classification model selection, performance metrics, and data preprocessing

In our prediction and classification scheme, we used target neuron ensembles as "classes" and

the activity of the responding/non-target neurons as the "features". As described in Stimulus

response measurement section, we use spike count over a window of 6 imaging frames as a

feature value for each neuron. For the temporal analysis, this window is offset relative to

stimulus onset, as described below. In a preliminary analysis, we compared several machine

learning strategies and found that the ensemble methods of boosting (XGBoost[38]) and bagging

(scikit-learn[88] Random Forest classifier) performed best to several other common classification

schemes such as Logistic Regression, Naive Bayes, Neural Networks, and Support Vector

Machines. These two approaches yielded the highest accuracy and were then applied to all of

our analyses presented in this work. In all cases we split the data samples randomly so that

80% was used for training and 20% for testing. We repeat this split 100 times and from these

report the mean and standard errors of our metrics. As metrics we use overall accuracy, per-

class recall and precision, as well as F1-scores (harmonic mean of recall and precision). For the

results in Fig. 4f and Supplementary Fig. 10d we use all data and all trials, but for others we

performed additional pre-processing before running classification, as detailed below.

Since not all trials were equally successful in eliciting spikes from target cells, for all

other decoding analysis we selectively filtered trials and targets, with a criterion that the target

neuron had to produce at least one spike across any of the six frames after the onset of

stimulation. We then selected valid targets as those that spiked on at least half of their trials.

Thus, non-significantly driven TC (based on conservative response spike count) excluded some

TC in the spike count analysis which nevertheless had significantly F1-score in the decoding

task (gray TC in Supplementary Fig. 10a). To characterize feature importances and interpret our

classification schemes we used Shapley analysis[89], as described below.

Decoding from baseline

To examine whether TCs are decodable during spontaneous activity, we applied a similar

analysis to the background activity in the periods between stimulation trials. We segmented this

background activity using the same bin size (6 frames) as in the stimulation condition and

identified pseudo-trials in which only one of the previously used TCs spiked within a bin. To

avoid potential artifacts introduced by Deep Interpolation, we excluded windows immediately

preceding or following stimulation. For each of these background spike datasets, we restricted

the analysis to the subset of TCs used during decoding in the stimulation condition and

downsampled the data to match the sample distribution of the stimulation trials (Fig. 4b, *right*).

Radius Analysis

For each target neuron, we implemented an exclusion zone based on the distance from the

target neuron's soma, *r*. We progressively increased *r*, and the neurons/features falling within

this distance were systematically eliminated from the feature matrix, enabling quantification of

the relationship between F1-score degradation and exclusion zone radius for individual target

neurons. We computed the probability distribution of any feature neuron being excluded, as a

function of distance, $P_{ex}(r)$. Specifically, we aggregated the frequency of neuron eliminations at

each radius across all datasets and normalized by the total number of exclusions.

The spatial dependency of predictive performance was characterized by analyzing the

differential changes in F1-score across the range of exclusion radii. For each target neuron

across all datasets, we identified the radius at which the decrease in F1-score first exceeded

three standard deviations below the mean rate of change. These radii were averaged across

datasets and smoothed using a Gaussian kernel ($\sigma = 1$) to obtain an estimate of the spatial

scale of predictive influence.

Neuron Drop Analysis and Positive Responder Probability

To evaluate the distributed nature of neural information and assess the robustness of our

decoding framework, we conducted an iterative feature elimination analysis for each target

neuron across all datasets. For each iteration, including initial model fitting, we use Shapley

values (employed using SHAP package: SHapley Additive exPlanations[90]) to quantitatively rank

feature neurons based on their predictive contribution for a given target neuron. Through

sequential elimination of the highest-ranked feature neurons and subsequent model retraining,

we systematically assessed how the removal of informative neurons impacted decoding

performance. This iterative process of SHAP-guided feature elimination and model

reconstruction was repeated until a single neuron remained. At each iteration we recorded F1-

score, precision, and overall accuracy and the entire procedure was replicated one hundred

times, utilizing different out-of-bag random selection of training and testing sets.

To characterize the relationship between feature importance and significant responders,

we quantified the probability that eliminated neurons exhibited significant responses. For each

target neuron, we constructed a binary vector corresponding to the sequential elimination order,

where entries were assigned values of 1 or 0 based on whether the eliminated neuron was a

significant responder. These binary sequences were averaged across all iterations of the

elimination procedure to obtain target-specific probabilities and then averaged across all target

neurons.

Temporal profiles of the prediction accuracy

To further characterize our holographic stimulation data we explored the accuracy of our

prediction scheme as a function of time. This procedure allowed us to study how long the

information about the origin of stimulation persists in network. Note that the prediction scheme is

a more stringent requirement than studying only the causal effects of stimulation, since even

though the activity after the stimulation can reverberate in a network for a long time, identifying

the node that initiated it is more challenging. Furthermore, for the temporal analysis we disabled

our filtering of inactive targets/trials, utilizing all data. This simplifies the interpretation of these

results, with an expected overall decrease in accuracy compared to the other decoding

analyses.

We assessed the temporal course of the prediction accuracy across datasets, each with

different levels of responsiveness of the targets. We varied the size of our response window (1

or 6 frames) and then moved it in steps of 1. The main results are present in Figure 4f (single-

frame analysis) & Supplementary Fig. 10d (six-frame analysis). For the six-frame analysis, the

fading "pink" rectangles indicate the amount of data used from the stimulation region (100 ms)

window, so that the fully white regions use data strictly before or after stimulation is over. Note

that the bins and the stimulation region are not perfectly aligned, as the trigger for the

stimulation was independent from the image scanning raster. We define the reference bin 0 as

the first bin which fully overlaps with the stimulation.

Avalanche analysis

Continuous epochs of suprathreshold population activity

Continuous periods of population activity were identified by applying a threshold Θ on the

population activity $p(t)$, the sum of the spikes from all neurons at a given time t, such that:

715
$$p_{\Theta}(t) = \begin{cases} p(t) - \Theta & p(t) > \Theta \\ 0 & p(t) \leq \Theta \end{cases} \tag{1}$$

This procedure is known as soft-thresholding and was employed for all analysis unless

otherwise stated. For a given recording $p(t)$ and coarse-graining value k, the dependence of the

number N of epochs on the threshold Θ, N(Θ) was obtained for a range of thresholds $\Theta \in [\Theta_1,$

$\Theta_2]$ such that Θ_1 was low enough that it removed no population activity from the time course and

Θ_2 was high enough that it would remove all population activity from the time course. The

function N(Θ) was typically well-approximated by a log-normal distribution and a corresponding

fit yielded shape parameters μ and σ. The threshold used in the analysis for all recordings and

coarse-graining factors was chosen such that $\Theta = \mu$ and estimated for each k. See Capek &

Ribeiro et al.[29] for more details.

Temporal coarse-graining

A temporal coarse-graining operation was applied to the thresholded population activity $p_\Theta(t)$.

For a given temporal coarse-graining factor k an ensemble of K different coarse-grained time

series $p_k^0(\tau), p_k^1(\tau), ..., p_k^{K-1}(\tau)$ was arrived at through the following method:

730
$$p_k^j(\tau) = \sum_{i=k\tau+j}^{k(\tau+1)+j-1} p_\Theta(i) \quad for \ \tau \in \left[0, 1, ..., \left\lfloor \frac{T-j}{k} \right\rfloor\right] \tag{2}$$

For each time series $p_k^j(\tau)$, epochs were extracted by finding pairs (τ_1, τ_2) such that $p_k^j(\tau_1) = 0$,

$p_k^j(\tau_2) = 0$ and $p_k^j(\tau') > 0$ for all $\tau' \in [\tau_1 + 1, ..., \tau_2 - 1]$. The size of the epoch is given by $S =$

$\sum_{i=\tau_1}^{\tau_2} p_k^j(i)$ and its corresponding duration given by $\tau_2 - \tau_1 - 1$. For a given ensemble of coarse-

grained time series, all epochs were pooled.

Scaling curve fit

For more precise evaluation of the scaling curve, mean avalanche size *vs* duration, we

introduced the following fitting function:

739
$$S(d) = \frac{C d^{\chi_{sh}}}{\left(1+(d/\Phi)^\gamma\right)^{(\chi_{sh}+\chi_{lg})/\gamma}} \tag{3}$$

This function is a double power law with initial slope χ_{sh}, transitioning to a second slope χ_{lg} at

around the point $d = \Phi$. The parameter γ controls how abruptly that transition happens and has

been fixed at 4 for all the curves presented. The other parameters were free to adjust to the

data, and all fits were performed in log-space, i.e., the log(S(log(d)) was fit to the log of data

(taking the log of both average sizes as well as durations).

Simulations

We created a 300x300 two-dimensional lattice of excitable cellular automata[91]. We

defined the spacing of the lattice as 12 μm, to mimic the high density of cells in brain tissue. We

connected cells by a probability function that changes as function of the distance between cell

pairs, with a short (gaussian, 100 μm characteristic decay) and a long (exponential, 290 μm

characteristic decay) component: $P_{conn} = 0.05e^{-r^2/2\times100^2} + 0.03e^{-r/290}$. At each time step (set

at 2 ms to mimic average spike transmission times in the brain), each cell can be in one of three

states: 1) resting, 2) active, or 3) refractory. A resting cell can become active by either receiving

an external Poisson drive (with probability P_{poiss}) or by an active presynaptic neighbor (with

probability P_{trans}). We set $P_{trans} = \sigma/\bar{k}$, where $\bar{k}$ is the average number of connections per cell

(~43 with the numbers described above) and σ is the branching parameter, a variable we can

adjust to move the system in and out of criticality and is expected to be slightly above 1 at the

critical point (to compensate for energy loss/collisions in spike propagation in the network –

~1.0066 on our simulations). We set $P_{poiss}(\sigma)$, so that the average amount of activity during

baseline for each σ value matches the average baseline activity in the experiments (see Fig.

6d).

762   We then selected for each "experiment" (10 total), a non-overlapping set of 10 cells in

the center of the network (inner 38x38 cells) to be the target of extra external drive (as in the

mice experiments, for 100 ms or 50 time steps, with 2 s inter-trial interval), which was set so the

average activity of each target in the simulations during stimulation matches the average activity

of targets during stimulation in the experimental data. All analyzes are then performed exactly

as done for the experimental data, but only using the inner 38x38 cells, to avoid border effects

(periodic boundary conditions were used to simplify the simulations).

Output Strength

To evaluate the relationship between the structure of the network and each target neuron's

ability to spread information regarding their firing, we introduced the measure of output strength:

$OS(i) = \sum_{o=1}^{5}\left(\sum_{j=1}^{N}(P_{trans}A(i,j))^o\right)$, where *A* is the network's adjacency matrix (*A(i,j)* = 1 when

there is a connection from neuron *i* to neuron *j*, and 0 otherwise), *N* is the number of neurons

and *P_{trans}* is the spike transmission probability. By taking *P_{trans}* *A* to the power *o*, we evaluate the

likelihood for a spike to transmit from *i* to *j* via an *o*-order path (e.g., order 1 would be a direct

connection, order 2 would be a two-step connection – neuron *i*, to neuron *k*, to neuron *j*).

Therefore, after summing over all other neurons, *OS(i)* estimates the number of spikes

generated in the network for each spike neuron *i* produced, excluding dynamics (collisions,

refractory periods, etc.) and only considering the network's structure.

Data availability

The preprocessed imaging data used in this study are available in the general repository Zenodo using the following access DOI:TBD. The source data for all figures and supplemental figures in this study are provided for this paper.

Code availability

Computer code used in this study is available at the following github link: TBD.

Acknowledgements

We thank Craig C. Stewart and members of the Plenz lab for help with animal surgery, animal care. This research was supported by the Division of the Intramural Research Program (DIRP) of the National Institute of Mental Health (NIMH), USA, ZIAMH002797, ZIAMH002971, and the BRAIN initiative Grant U19 NS107464-01.

This research was supported by the Intramural Research Program of the National Institutes of Health (NIH), USA, ZIAMH002797, ZIAMH002971, and the BRAIN initiative Grant U19 NS107464-01. This research utilized the supercomputing resources of the National Institutes of Health (NIH, USA; Biowulf, http://hpc.nih.gov).

The contributions of the authors were made as part of their official duties as NIH federal employees, are in compliance with agency policy requirements, and are considered Works of the United States Government. However, the findings and conclusions presented in this paper are those of the author(s) and do not necessarily reflect the views of the NIH or the U.S. Department of Health and Human Services.

Author Contributions

TLR and DP conceived and planned the study; TLR, BG, and VS performed experiments; AV took the lead in holographic setup design. TLR took the lead in data analysis with support from BG and VS. SP and RS took the lead on machine learning based decoding of experimental data. All

807   authors contributed to the analyses. TLR and DP wrote the manuscript.

Competing interests

810   The authors declare no competing interests.

Supplementary Figures

812   **Supplementary Figure 1**

Supplementary Fig. 1 | a Widefield image showing GCaMP7s (*green*) and ChrimsonR (*red*) expression, with the merged image indicating co-expression (single mouse). **b** Optimal co-expression achieved with a virus titer ratio of 14:1 (n = 23 mice). **c** Examples of individual neurons with significant GCaMP7s (*green*) and ChrimsonR (*red*) expression from 3 mice. **d** Mean expression density showing twice as many neurons positive for GCaMP7s compared to ChrimsonR (from c).

814   **Supplementary Figure 2**

Supplementary Fig. 2 | Schematic design and evaluation of the stimulation path using spatial light modulation. a Laser light (Light Conversion, 1064 nm, 600 kHz, <100 fs) is expanded to fill the area of the spatial light modulator (SLM; Meadowlark, 1920×1152 resolution). The hologram is resized to fit the Galvo-Galvo scanners (~3 mm diameter), which guide the stimulation beam in the focal image plane (Nikon 16X, 3 mm working distance, 12.5 mm effective focal length). Target cells (TC) expressing both GCaMP7s and ChrimsonR (opsin with tdTomato tag) are selected from the mean 2PI image of the field of view from both red and green channels for stimulation and simultaneous estimation of the evoked spike count. Map of TC informs a hologram of top-hat patterns with diameter of 10 µm for the SLM. Optical power was adjusted to 5—10 mW per TC, depending on the depth and expression level. **b** Spiking above base line increases with stimulation power (N = 3 TC; 100 ms stimulation duration). **c** No difference in evoked spike count for 5 and 10 mW stimulation across all TC. **d** Schematics of the stimulation xy-profile reconstruction using fluorescence microbeads. Galvo-Galvo (GG) controlled stimulation of gel-embedded microbeads (3 µm diameter) and simultaneous Galvo-Resonance (GR) 2PI of elicited response. *Circles*: Individual microbeads before and during stimulation. **e** Experimentally reconstructed and smoothed xy-profile of a flat-hat stimulation (10 µm diameter; 5 mW) at 4 different positions within the field of view at center (0,0), off-x (+100 µm, 0), off-y (0, +100 µm), and off-xy (+100 µm, +100 µm). *Inset*: Non-smoothed reconstruction.

816   **Supplementary Figure 3**

Supplementary Fig. 3 | Analysis of motion correction from processed recordings reveals low amount of motion artifact affecting stimulation. a Distribution of motion offsets observed on each frame over all recordings. *Blue*: all data. *Orange*: maximum displacement observed during stimulation for each trial. **b** Spatial profile of stimulation response from targets measured using motion offset in trials during stimulation. Note that responses drop to approximately zero in frames with an offset above ~4.4 μm.

817
818

819   **Supplementary Figure 4**

Supplementary Fig. 4 | Example recording with activity from two example targets and some of their significant responders highlighted. a Example field of view with target 1 highlighted in *red* and its positive (*blue*) and negative (*purple*) responders indicated. **b** Same recording as in a, but now highlighting target 9 and its significant responders. **c** *Left*: time average fluorescent response to stimulation for the significant responders labeled in a. *Right*: example raw (*dark*) and denoised (*light*) fluorescent traces (*lines*) and deconvolved spikes (*bars*) for the labeled neurons in a. **d** *Left*: time average fluorescent response to stimulation for the significant responders labeled in b. *Right*: example raw (*dark*) and denoised (*light*) fluorescent traces (*lines*) and deconvolved spikes (*bars*) for the labeled neurons in b. *Bottom*: zoom in from the right panel above.

821 **Supplementary Figure 5**

Supplementary Fig. 5 | Summary of isolating individual pyramidal neurons in layer 2/3 of the primary visual cortex during rest using holographic stimulation and 2PI. For each target cell (TC), mean luminance after motion correction and denoising (*left*) and the ΔF/F in response to stimulation in a 100×100-pixel area (~88×88 μm) centered on the TC (*right*) are shown. *Asterisks* are visual guides for TC position.

823   **Supplementary Figure 6**

Supplementary Fig. 6 | Spike count in all different subpopulations scale with spike count of target cells during baseline activity, but only Positive Responders scale during stimulation. a Mean spike count on PosR (*blue*), NonR (*grey*) and NegR (*purple*) as function of spike count of TC, in log-log, during baseline. Power law fits are indicated by solid lines, with obtained exponents shown. *Left*: mean calculated per neuron. *Right*: mean calculated after summing over the population. **b** Same as in a, but during holographic stimulation. **c** Response scaling calculated using the fluorescent traces (ΔF/F).

825   **Supplementary Figure 7**

Supplementary Fig. 7 | Correlations between PosR and TC change during stimulation in comparison with baseline. For each PosR-TC pair, their spike count correlation during stimulation is plotted against baseline. Note the shift towards higher values during Resp compared to Base (cp. Fig. 2c). A linear regression (*orange line*) shows that these measures are very weakly related ($R^2 = 0.018$; linear regression; $p < 10^{-5}$, for the *t*-statistic of the two-sided hypothesis test), indicating that the stimulation of TCs engages both existing but also new networks.

827 **Supplementary Figure 8**

Supplementary Fig. 8 | Response scaling and avalanche analysis are qualitatively unchanged without denoising from Deep Interpolation. a Average response from PosR (*blue*), NonR (*grey*) and NegR (*purple*) as function of TC spike count during stimulation, for datasets processed without denoising (no Deep Interpolation). **b** Avalanche analysis for datasets processed without denoising. *Left*: Distribution of avalanche sizes for different coarse graining levels *k* (*color bar*). *Right*: Mean avalanche size as function of avalanche duration at *k* = 6 showing an exponent below 2 for the short avalanches regime, in line with previous results obtained from noisy datasets (Ribeiro, Capek et al., 2023).

829 **Supplementary Figure 9**

Supplementary Fig. 9 | High response variability in PosR and NonR does not originate from variability in evoked spikes of TC. a Summary statistics of mean TC trials separated by low and high target counts. Note the remaining high variance of PosR despite the large drop in TC variance. No change in Fano Factor (FF) when subdividing into TC trials with high or low preceding population spike count (**b**) or early (first half of the response window) vs. late (second half of the response window) spiking responses (**c**). When comparing FF (b, c), 1 & 2 stars indicate $p < 0.05$ & 10^{-3}, respectively, using Wilcoxon rank sum test.

830
831

832 **Supplementary Figure 10**

Supplementary Fig. 10 | Decoding accuracy, stimulation efficacy and impact of deep interpolation. a Overview of F1-scores for TCs (*green*) and TCs that did not reach 20% of spike success (*gray*) upon stimulation. F1-scores are compared to chance level (*red bars*) and overall accuracy (*red circles*). XGBoost (XGB; *left*) and Random Forest (RF; *right*) results are very similar across all datasets. Potential multi-cell targets (cp. Supplementary Fig. 5; TCs 27-29, 31-32, 35, 39, 45 and 49) are labeled and marked in *yellow*, and overall do not show above-average decoding success. **b** For each data set, stimulation success (at least 1 spike) varied among TC. Target stimulation success positively correlated with decoding power, i.e. F1-score, of a TC in XGB (*left*) and RF (*right*). **c** Z-scored F1-scores calculated with the Random Forest method for each TC and then averaged, for each dataset (same color code as in Fig. 4), as function of number of best predictors dropped (cp. XGB in Fig. 4c, *right*). *Inset*: Average number of best predictors dropped before F1-scores reach chance levels. **d** Deep Interpolation slightly advances stimulus information and significantly increases accuracy thereby slightly prolonging transient response time. Analysis done over 132-ms window (cp. Fig. 4f, 22-ms window). The enhanced accuracy achieved with Deep Interpolation demonstrates the method's noise-reducing benefits, which improve the model's ability to capture relevant neural signals. However, it also shows that it undermines our ability to study temporal progression of activations, as its window of smoothing ranges across 30 frames. **e** Low (but still above chance) F1 and accuracy when XGB and RF are trained on the raw ΔF/F traces.

834 **Supplementary Figure 11**

Supplementary Fig. 11 | Spike count in subpopulations scale similar with spike count of target cells during low and high avalanche activity states. a Mean spike count on PosR (*blue*), NonR (*grey*) and NegR (*purple*) as function of spike count of TC, in log-log, during holographic stimulation. Power law fits are indicated by solid lines, with obtained exponents shown. Separating trials with preceding low (*left*) or high (*right*) baseline network activity. Experimental data. **b** Mean response of PosR as function of TC spike count during trials with low base activity for the simulations. *Bottom*: Response scaling exponents as function of branching parameter σ. **c** Same as in b, but for trials with high base activity prior to stimulation.

836   **Supplementary Figure 12**

Supplementary Fig. 12 | No correlation between number of PosR for TCs between sub- and critical networks with identical connection structure. Number of PosR for each TC (*circles*) at the critical network (σ = 1.0066) as function of the same measure from a subcritical network (σ = 0.5). From the *t*-statistics of the two-sided hypothesis test we obtained p = 0.27 for the slope of the linear regression (*red line*), with R^2 = 0.002.

838 **Supplementary Table 1**

Experimental response scaling exponents (± 95% C.I.)				
Condition	Figure	PosR	NonR	NegR
Base (neuron)	S6a *left*	0.46 ± 0.07	0.52 ± 0.27	0.44 ± 0.02
Base (network)	2d, S6a *right*	0.55 ± 0.22	0.41 ± 0.27	0.76 ± 0.24
Resp (neuron)	2e, S6b *left*	0.26 ± 0.09	0.01 ± 0.10	-0.30 ± 0.18
Resp (network)	2f, S6b *right*	0.28 ± 0.10	0.04 ± 0.09	-0.07 ± 0.26
Calcium	S6c	0.18 ± 0.04	0.05 ± 0.04	-0.18 ± 0.10
Low	S9 *left*	0.29 ± 0.14	0.06 ± 0.07	-0.33 ± 0.50
High	S9 *right*	0.22 ± 0.21	-0.04 ± 0.18	-0.31 ± 0.66
No DeepIP	S10a	0.35 ± 0.18	0.08 ± 0.09	-0.78 ± 1.48

840 **Supplementary Table 2**

Simulation response scaling exponents (± 95% C.I.)			
Branching parameter	Response (Fig. 7b)	Low (Fig. 7c, left)	High (Fig. 7c, right)
0.3	0.015 ± 0.048	N/A	N/A
0.5	0.045 ± 0.031	N/A	N/A
0.9	0.105 ± 0.038	N/A	N/A
0.95	0.133 ± 0.040	N/A	N/A
1	0.176 ± 0.112	N/A	N/A
1.003	0.179 ± 0.087	0.144 ± 0.150	0.124 ± 0.015
1.005	0.166 ± 0.044	0.182 ± 0.127	0.112 ± 0.043
1.0061	0.174 ± 0.032	0.201 ± 0.115	0.116 ± 0.019
1.0066	0.247 ± 0.086	0.275 ± 0.080	0.129 ± 0.039
1.0076	0.186 ± 0.082	0.469 ± 0.111	0.109 ± 0.028
1.009	0.198 ± 0.070	0.672 ± 0.108	0.112 ± 0.038
1.01	0.189 ± 0.072	0.920 ± 0.098	0.091 ± 0.013

References

1   Allen, W. E. *et al.* Global Representations of Goal-Directed Behavior in Distinct Cell Types of Mouse Neocortex. *Neuron* **94**, 891–907.e896 (2017). https://doi.org/10.1016/j.neuron.2017.04.017

2   London, M., Roth, A., Beeren, L., Häusser, M. & Latham, P. E. Sensitivity to perturbations in vivo implies high noise and suggests rate coding in cortex. *Nature* **466**, 123–127 (2010).

3   Meyer, J. F., Golshani, P. & Smirnakis, S. M. The Effect of Single Pyramidal Neuron Firing Within Layer 2/3 and Layer 4 in Mouse V1. *Frontiers in Neural Circuits* **12** (2018). https://doi.org/10.3389/fncir.2018.00029

4   Chettih, S. N. & Harvey, C. D. Single-neuron perturbations reveal feature-specific competition in V1. *Nature* **567**, 334–340 (2019). https://doi.org/10.1038/s41586-019-0997-6

5   Jouhanneau, J.-S., Kremkow, J. & Poulet, J. F. A. Single synaptic inputs drive high-precision action potentials in parvalbumin expressing GABA-ergic cortical neurons in vivo. *Nature communications* **9**, 1540 (2018). https://doi.org/10.1038/s41467-018-03995-2

6   Hemberger, M., Shein-Idelson, M., Pammer, L. & Laurent, G. Reliable sequential activation of neural assemblies by single pyramidal cells in a three-layered cortex. *Neuron* **104**, 353–369. e355 (2019).

7   O'Rawe, J. F. *et al.* Excitation creates a distributed pattern of cortical suppression due to varied recurrent input. *Neuron* **111**, 4086–4101.e4085 (2023). https://doi.org/10.1016/j.neuron.2023.09.010

8   Marshel, J. H. *et al.* Cortical layer–specific critical dynamics triggering perception. *Science*, eaaw5202 (2019). https://doi.org/10.1126/science.aaw5202

9   Carrillo-Reid, L., Han, S., Yang, W., Akrouh, A. & Yuste, R. Controlling Visually Guided Behavior by Holographic Recalling of Cortical Ensembles. *Cell* **178**, 447–457.e445 (2019). https://doi.org/10.1016/j.cell.2019.05.045

10   Houweling, A. R. & Brecht, M. Behavioural report of single neuron stimulation in somatosensory cortex. *Nature* **451**, 65–68 (2008). https://doi.org/10.1038/nature06447

11   Wolfe, J., Houweling, A. R. & Brecht, M. Sparse and powerful cortical spikes. *Curr. Opin. Neurobiol.* **20**, 306–312 (2010). https://doi.org/https://doi.org/10.1016/j.conb.2010.03.006

12   Cheng-yu, T. L., Poo, M.-m. & Dan, Y. Burst spiking of a single cortical neuron modifies global brain state. *Science* **324**, 643–646 (2009).

13   Zucca, S. *et al.* An inhibitory gate for state transition in cortex. *Elife* **6**, e26177 (2017).

14   Histed, M. H., Ni, A. M. & Maunsell, J. H. Insights into cortical mechanisms of behavior from microstimulation experiments. *Prog. Neurobiol.* **103**, 115–130 (2013). https://doi.org/10.1016/j.pneurobio.2012.01.006

15   Vaadia, E. *et al.* Dynamics of neuronal interactions in monkey cortex in relation to behavioural events. *Nature* **373**, 515–518 (1995).

16   Stringer, C. *et al.* Spontaneous behaviors drive multidimensional, brainwide activity. *Science* **364**, eaav7893 (2019). https://doi.org/doi:10.1126/science.aav7893

17   Kuhn, A., Aertsen, A. & Rotter, S. Neuronal integration of synaptic input in the
fluctuation-driven regime. *J. Neurosci.* **24**, 2345–2356 (2004).
https://doi.org/10.1523/JNEUROSCI.3349-03.2004
18   Arieli, A., Sterkin, A., Grinvald, A. & Aertsen, A. Dynamics of ongoing activity:
Explanation of the large variability in evoked cortical responses. *Science* **273**, 1868–1871
(1996).
19   Braitenberg, V. & Schüz, A. *Anatomy of the Cortex: Statistics and Geometry.* (Springer-
Verlag, 1991).
20   Thomson, A. M., West, D. C., Wang, Y. & Bannister, A. P. Synaptic connections and
small circuits involving excitatory and inhibitory neurons in layers 2-5 of adult rat and cat
neocortex: triple intracellular recordings and biocytin labelling in vitro. *Cereb. Cortex* **12**,
936–953 (2002).
21   Levy, R. B. & Reyes, A. D. Spatial Profile of Excitatory and Inhibitory Synaptic
Connectivity in Mouse Primary Auditory Cortex. *J. Neurosci.* **32**, 5609 (2012).
https://doi.org/10.1523/JNEUROSCI.5158-11.2012
22   Jouhanneau, J.-S., Kremkow, J., Dorrn, A. L. & Poulet, J. F. In vivo monosynaptic
excitatory transmission between layer 2 cortical pyramidal neurons. *Cell reports* **13**,
2098–2106 (2015).
23   Campagnola, L. *et al.* Local connectivity and synaptic dynamics in mouse and human
neocortex. *Science* **375**, eabj5861 (2022). https://doi.org/doi:10.1126/science.abj5861
24   Kumar, A., Rotter, S. & Aertsen, A. Spiking activity propagation in neuronal networks:
reconciling different perspectives on neural coding. *Nat. Rev. Neurosci.* **11**, 615–627
(2010). https://doi.org/10.1038/nrn2886
25   Haider, B., Hausser, M. & Carandini, M. Inhibition dominates sensory responses in the
awake cortex. *Nature* **493**, 97–100 (2013).
https://doi.org/http://www.nature.com/nature/journal/v493/n7430/abs/nature11665.html#s
upplementary-information
26   Tremblay, R., Lee, S. & Rudy, B. GABAergic interneurons in the neocortex: From
cellular properties to circuits. *Neuron* **91**, 260–292 (2016).
https://doi.org/10.1016/j.neuron.2016.06.033
27   Chialvo, D. R., Cannas, S. A., Grigera, T. S., Martin, D. A. & Plenz, D. Controlling a
complex system near its critical point via temporal correlations. *Scientific Reports* **10**,
12145 (2020). https://doi.org/10.1038/s41598-020-69154-0
28   Muñoz, M. A., Dickman, R., Vespignani, A. & Zapperi, S. Avalanche and spreading
exponents in systems with absorbing states. *Phys. Rev. E* **59**, 6175–6179 (1999).
29   Capek, E. *et al.* Parabolic avalanche scaling in the synchronization of cortical cell
assemblies. *Nature communications* **14**, 2555 (2023).
https://doi.org/10.1101/2022.11.02.514938
30   Plenz, D. *et al.* Self-Organized Criticality in the Brain. *FrPhy* **9** (2021).
https://doi.org/10.3389/fphy.2021.639389
31   Beggs, J. M. & Plenz, D. Neuronal avalanches in neocortical circuits. *J. Neurosci.* **23**,
11167–11177 (2003). https://doi.org/10.1523/JNEUROSCI.23-35-11167.2003
32   Softky, W. R. & Koch, C. The highly irregular firing of cortical cells is inconsistent with
temporal integration of random EPSPs. *J. Neurosci.* **13**, 334–350 (1993).

| 930 | 33 | Holmgren, C., Harkany, T., Svennenfors, B. & Zilberter, Y. Pyramidal cell communication within local networks in layer 2/3 of rat neocortex. *The Journal of physiology* **551**, 139–153 (2003). |

33 Holmgren, C., Harkany, T., Svennenfors, B. & Zilberter, Y. Pyramidal cell
communication within local networks in layer 2/3 of rat neocortex. *The Journal of*
physiology **551**, 139–153 (2003).
34 Storey, J. D. A Direct Approach to False Discovery Rates. *Journal of the Royal*
Statistical Society. Series B (Statistical Methodology) **64**, 479–498 (2002).
35 Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society. Series B*
(Methodological) **57**, 289–300 (1995).
36 Churchland, M. M. *et al.* Stimulus onset quenches neural variability: a widespread
cortical phenomenon. *Nat. Neurosci.* **13**, 369–378 (2010).
37 Rajdl, K., Lansky, P. & Kostal, L. Fano factor: a potentially useful information. *Frontiers*
in computational neuroscience **14**, 569049 (2020).
38 Chen, T. & Guestrin, C. in *Proceedings of the 22nd ACM SIGKDD International*
Conference on Knowledge Discovery and Data Mining   785–794 (Association for
Computing Machinery, San Francisco, California, USA, 2016).
39 Ho, T. K. in *Proceedings of 3rd international conference on document analysis and*
recognition. 278–282 (IEEE).
40 Breiman, L. Random forests. *MLear* **45**, 5–32 (2001).
41 Srinivasan, K., Ribeiro, T. L., Kells, P. & Plenz, D. The recovery of parabolic avalanches
in spatially subsampled neuronal networks at criticality. *Scientific Reports* **14**, 19329
(2024). https://doi.org/10.1038/s41598-024-70014-4
42 Miller, S. R., Yu, S. & Plenz, D. The scale-invariant, temporal profile of neuronal
avalanches in relation to cortical γ–oscillations. *Scientific Reports* **9**, 16403 (2019).
https://doi.org/10.1101/757278
43 Yu, S., Klaus, A., Yang, H. & Plenz, D. Scale-invariant neuronal avalanche dynamics
and the cut-off in size distributions. *PLoS One* **9**, e99761 (2014).
https://doi.org/10.1371/journal.pone.0099761
44 Marro, J. & Dickman, R. Nonequilibrium phase transitions in lattice models.
Nonequilibrium Phase Transitions in Lattice Models (2005).
45 Ribeiro, T. L., Ribeiro, S., Belchior, H., Caixeta, F. & Copelli, M. Undersampled critical
branching processes on small-world and random networks fail to reproduce the statistics
of spike avalanches. *PloS one* **9**, e94992 (2014).
https://doi.org/10.1371/journal.pone.0094992
46 Kinouchi, O. & Copelli, M. Optimal dynamical range of excitable networks at criticality.
Nat. Phys. **2** 348–351 (2006).
47 Histed, M. H., Bonin, V. & Reid, R. C. Direct activation of sparse, distributed
populations of cortical neurons by electrical microstimulation. *Neuron* **63**, 508–522
(2009). https://doi.org/10.1016/j.neuron.2009.07.016
48 Seeman, S. C. *et al.* Sparse recurrent excitatory connectivity in the microcircuit of the
adult mouse and human cortex. *eLife* **7**, e37349 (2018).
https://doi.org/10.7554/eLife.37349
49 Abeles, M. *Corticonics: Neural Circuits of the Cerebral Cortex*. Vol. 1 (Cambridge
University Press, 1991).
50 Diesmann, M., Gewaltig, M. O. & Aertsen, A. Stable propagation of synchronous spiking
in cortical neural networks. *Nature* **402**, 529–533 (1999). https://doi.org/10.1038/990101

Prechtl, J. C., Cohen, L. B., Pesaran, B., Mitra, P. P. & Kleinfeld, D. Visual stimuli induce waves of electrical activity in turtle cortex. *Proc.Natl.Acad.Sci.U.S.A* **94**, 7621–7626 (1997).

Rubino, D., Robbins, K. A. & Hatsopoulos, N. G. Propagating waves mediate information transfer in the motor cortex. *Nat. Neurosci.* **9**, 1549–1557 (2006).

Singer, W. Cortical dynamics revisited. *Trends Cogn Sci* (2013).

Gray, C. M., Konig, P., Engel, A. K. & Singer, W. Oscillatory responses in cat visual cortex exhibit inter-columnar synchronization which reflects global stimulus properties. *Nature* **338**, 334–337 (1989).

Lisman, J. E. Bursts as a unit of neural information: making unreliable synapses reliable. *Trends Neurosci.* **20**, 38–43 (1997).

Riquelme, J. L., Hemberger, M., Laurent, G. & Gjorgjieva, J. Single spikes drive sequential propagation and routing of activity in a cortical network. *eLife* **12**, e79928 (2023). https://doi.org/10.7554/eLife.79928

Shu, Y., Hasenstaub, A., Badoual, M., Bal, T. & McCormick, D. A. Barrages of synaptic activity control the gain and sensitivity of cortical neurons. *J. Neurosci.* **23**, 10388–10401 (2003).

Tsodyks, M., Uziel, A. & Markram, H. Synchrony generation in recurrent networks with frequency-dependent synapses. *J. Neurosci.* **20**, RC50 (2000).

Molnár, G. *et al.* Complex Events Initiated by Individual Spikes in the Human Cerebral Cortex. *PLoS Biol.* **6**, e222 (2008). https://doi.org/10.1371/journal.pbio.0060222

Ferrarese, L. *et al.* Dendrite-Specific Amplification of Weak Synaptic Input during Network Activity In Vivo. *Cell Reports* **24**, 3455–3465.e3455 (2018). https://doi.org/https://doi.org/10.1016/j.celrep.2018.08.088

Harris, K. D. & Mrsic-Flogel, T. D. Cortical connectivity and sensory coding. *Nature* **503**, 51–58 (2013). https://doi.org/10.1038/nature12654

Shadlen, M. N. & Newsome, W. T. The variable discharge of cortical neurons: implications for connectivity, computation, and information coding. *J. Neurosci.* **18**, 3870–3896 (1998).

Histed, M. H. & Maunsell, J. H. Cortical neural populations can guide behavior by integrating inputs linearly, independent of synchrony. *Proc. Natl. Acad. Sci. U. S. A.* **111**, E178–187 (2014). https://doi.org/10.1073/pnas.1318750111

Barth, A. L. & Poulet, J. F. A. Experimental evidence for sparse firing in the neocortex. *Trends Neurosci.* **35**, 345–355 (2012).

Stevens, S. S. *Psychophysics: Introduction to its perceptual, neural and social prospects*. (Routledge, 2017).

Assis, V. R. & Copelli, M. Dynamic range of hypercubic stochastic excitable media. *Phys Rev E Stat Nonlin Soft Matter Phys* **77**, 011923 (2008). https://doi.org/10.1103/PhysRevE.77.011923

Scott, G. *et al.* Voltage imaging of waking mouse cortex reveals emergence of critical neuronal dynamics. *J. Neurosci.* **34**, 16611 – 16620 (2014). https://doi.org/10.1523/JNEUROSCI.3474-14.2014

Bellay, T., Klaus, A., Seshadri, S. & Plenz, D. Irregular spiking of pyramidal neurons organizes as scale-invariant neuronal avalanches in the awake state. *eLife* **4**, e07224 (2015). https://doi.org/10.7554/eLife.07224

69   Bellay, T., Shew, W. L., Yu, S., Falco-Walter, J. J. & Plenz, D. Selective participation of
single cortical neurons in neuronal avalanches. *Frontiers in Neural Circuits* **14** (2021).
https://doi.org/10.3389/fncir.2020.620052
70   Petermann, T. *et al.* Spontaneous cortical activity in awake monkeys composed of
neuronal avalanches. *Proc. Natl. Acad. Sci. U.S.A.* **106**, 15921–15926 (2009).
https://doi.org/10.1073/pnas.0904089106
71   Beggs, J. M. & Plenz, D. Neuronal avalanches are diverse and precise activity patterns
that are stable for many hours in cortical slice cultures. *J. Neurosci.* **24**, 5216–5229
(2004).
72   Stewart, C. V. & Plenz, D. Inverted-U profile of dopamine-NMDA-mediated
spontaneous avalanche recurrence in superficial layers of rat prefrontal cortex. *J.*
Neurosci. **26**, 8148–8159 (2006). https://doi.org/10.1523/JNEUROSCI.0723-06.2006
73   Lombardi, F., Herrmann, H. J., Plenz, D. & de Arcangelis, L. Temporal correlations in
neuronal avalanche occurrence. *Scientific Reports* **6**, 24690 (2016).
https://doi.org/10.1038/srep24690
74   Lombardi, F., Herrmann, H., Plenz, D. & de Arcangelis, L. On the temporal organization
of neuronal avalanches. *Front. Syst. Neurosci.* **8**, 204 (2014).
https://doi.org/10.3389/fnsys.2014.00204
75   Jun, N. Y. *et al.* Coordinated multiplexing of information about separate objects in visual
cortex. *eLife* **11**, e76452 (2022). https://doi.org/10.7554/eLife.76452
76   Caruso, V. C. *et al.* Single neurons may encode simultaneous stimuli by switching
between activity patterns. *Nature communications* **9**, 2715 (2018).
https://doi.org/10.1038/s41467-018-05121-8
77   Levy, M., Sporns, O. & MacLean, J. N. Network analysis of murine cortical dynamics
implicates untuned neurons in visual stimulus coding. *Cell Reports* **31** (2020).
https://doi.org/10.1016/j.celrep.2020.03.047
78   Goldey, G. J. *et al.* Removable cranial windows for long-term imaging in awake mice.
Nature Protocols **9**, 2515 (2014). https://doi.org/10.1038/nprot.2014.165
79   Dana, H. *et al.* High-performance GFP-based calcium indicators for imaging activity in
neuronal populations and microcompartments. *bioRxiv* (2018).
80   Klapoetke, N. C. *et al.* Independent optical excitation of distinct neural populations. *Nat.*
Methods **11**, 338 (2014). https://doi.org/10.1038/nmeth.2836
https://www.nature.com/articles/nmeth.2836#supplementary-information
81   Juavinett, A. L., Nauhaus, I., Garrett, M. E., Zhuang, J. & Callaway, E. M. Automated
identification of mouse visual areas with intrinsic signal imaging. *Nature Protocols* **12**,
32–43 (2017). https://doi.org/10.1038/nprot.2016.158
82   Sit, K. K. & Goard, M. J. Distributed and retinotopically asymmetric processing of
coherent motion in mouse visual cortex. *Nature communications* **11**, 1–14 (2020).
83   Kleiner, M., Brainard, D. & Pelli, D. What's new in Psychtoolbox-3? (2007).
84   Adesnik, H. & Abdeladim, L. Probing neural codes with two-photon holographic
optogenetics. *Nat. Neurosci.* **24**, 1356–1366 (2021). https://doi.org/10.1038/s41593-021-
00902-9
85   Pachitariu, M. *et al.* Suite2p: beyond 10,000 neurons with standard two-photon
microscopy. *bioRxiv*, 061507 (2017). https://doi.org/10.1101/061507
86   Lecoq, J. *et al.* Removing independent noise in systems neuroscience data using
DeepInterpolation. *Nat. Methods* (2021). https://doi.org/10.1038/s41592-021-01285-2

87 Dalgleish, H. W. *et al.* How many neurons are sufficient for perception of cortical
activity? *eLife* **9**, e58889 (2020). https://doi.org/10.7554/eLife.58889
88 Pedregosa, F. Scikit-learn: Machine learning in python Fabian. *Journal of machine*
learning research **12**, 2825 (2011).
89 Shapley, L. S. Notes on the n-person game—ii: The value of an n-person game. (RAND
Coorperation, Santa Monica, CA, USA, 1951).
90 Scott, M. & Su-In, L. A unified approach to interpreting model predictions. *Advances in*
neural information processing systems **30**, 4765–4774 (2017).
91 Greenberg, J. M. & Hastings, S. P. Spatial Patterns for Discrete Models of Diffusion in
Excitable Media. *SIAM Journal on Applied Mathematics* **34**, 515–523 (1978).

1076