Title: Integrative analysis of angiogenic signaling in obesity: capillary features and VEGF binding kinetics

Yunjeong Lee[1], Keith Lionel Tukei[1], Yingye Fang[1], Shobhan Kuila[1], Xinming Liu[1], Princess I. Imoukhuede[1*]

[1]Department of Bioengineering, University of Washington, Seattle, WA, USA
[*]Corresponding author: pii@uw.edu; +1 (206) 685-2002

ORCID

Yunjeong Lee: https://orcid.org/0000-0001-7824-1339
Keith Lionel Tukei: https://orcid.org/0000-0003-2249-3347
Yingye Fang: https://orcid.org/0000-0003-4137-866X
Shobhan Kuila: https://orcid.org/0000-0001-5679-2105
Xinming Liu: https://orcid.org/0000-0002-4683-8779
Princess I Imoukhuede: https://orcid.org/0000-0002-4257-1085

Abstract

Obesity is a global health crisis, with its prevalence particularly severe in the United States, where over 42% of adults are classified as obese. Obesity is driven by complex molecular and tissue-level mechanisms that remain poorly understood. Among these, angiogenesis—primarily mediated by vascular endothelial growth factor (VEGF-A)—is critical for adipose tissue expansion but presents unique challenges for therapeutic targeting due to its intricate regulation. Systems biology approaches have advanced our understanding of VEGF-A signaling in vascular diseases, but their application to obesity is limited by scattered and sometimes contradictory data. To address this gap, we performed a comprehensive analysis of the existing literature to synthesize key findings, standardize data, and provide a holistic perspective on the adipose vascular microenvironment. The data mining revealed five key findings: (1) obesity increases adipocyte size by 78%; (2) vessel density in adipose tissue decreases by 51% in obese mice, with vessels being 47–58% smaller and 4–9 times denser in comparison with tumor vessels; (3) capillary basement membrane thickness remains similar regardless of obesity; (4) VEGF-A shows the strongest binding affinity for VEGFR1, with four times stronger affinity for VEGFR2 than for NRP1; and (5) binding affinities measured by radioligand binding assay and surface plasmon resonance (SPR) are significantly different. These consolidated findings provide essential parameters for systems biology modeling, new insights into obesity-induced changes in adipose tissue, and a foundation for developing angiogenesis-targeting therapies for obesity.

Keywords

Data mining, data analysis, obesity, adipose tissue, vascular endothelial growth factor

Introduction

Angiogenesis, primarily driven by vascular endothelial growth factor-A (VEGF-A, i.e., VEGF-A165), is essential for healthy adipose tissue expansion and remodeling[1]. Insufficient vascularization causes unhealthy white adipose tissue expansion in obesity, leading to fibrosis, inflammation, and systemic insulin resistance in preclinical models[2–5]. Therapeutic modulation of angiogenesis remains controversial because targeting the primary molecular drivers of vascularization, the VEGF family, does not fully control obesity[6,7]. This is despite the numerous studies establishing the relationship between VEGF signaling and adipose tissue development, which have found that (1) VEGF-A mRNA expression is upregulated during adipocyte differentiation in vitro[8], (2) overexpression of VEGF-A in adipose tissue not only promotes angiogenesis but also improves the metabolic system[9], and (3) anti-VEGF suppresses both angiogenesis and the formation of differentiating adipocytes[10]. Despite these discoveries, how to control adipogenesis via VEGF-A signaling is still unclear. This knowledge gap can be addressed by using systems biology to describe the fundamental processes governing vascularization in obesity with mathematical models and experimental data, simulating them with computational approaches, and analyzing the system to identify the best approaches to addressing insufficient vascularization in obesity.

Although several mathematical models have been developed to explain VEGF-A signaling in cancer or peripheral arterial disease, they are hard to employ in developing obesity-specific models because of the following distinctive features of adipose tissue: (1) an extremely high proportion of adipocytes in adipose tissue volume[11] that is not present in the established muscle[12] or tumor models[13–15] (Figure 1A), (2) phenotypically different capillaries in adipose tissue compared with unorganized tumor capillaries, and (3) different levels of the molecular drivers of angiogenesis, VEGFRs, in adipose tissue (Figure 1B)[13,16]. Thus, there is a need to develop adipose tissue–specific computational models, and these require adipose tissue–specific data and mathematical representations.

To address these needs for adipose tissue–specific data on vascular and adipocyte properties, we performed an in-depth analysis of the experimental literature, identifying data governing adipocyte and vascular morphology and data that standardize VEGFR binding kinetics. These data offer new insights into the adipose-tissue vascular microenvironment, reflecting the differential changes occurring as a result of obesity. They will accelerate the use of systems biology to mathematically represent, model, and predict approaches for treating obesity.

Methods

Search Strategy

We mined the literature using PubMed, Google Scholar, and the University of Washington Libraries. Search keywords included adipocyte size, high-fat diet mice, gonadal adipose tissue, vessel size, vessel diameter, capillary diameter, vessel density, tumor, retina capillary basement membrane thickness, muscle capillary basement membrane thickness, brain capillary basement membrane thickness, thickness of renal glomerular capillary basement membrane, VEGF, VEGF-A, VEGF165, VEGF-A165, VEGFR1, Flt-1, VEGFR2, KDR, NRP1, surface plasmon resonance, SPR, radioligand, radioactivity, 123I, 125I, binding affinity, and K_d.

Study selection and eligibility criteria

We included *ex vivo* studies that measured adipocyte diameter or the cross-sectional area of an adipocyte,
the cross-sectional area of vessels (vessel size), and vessel densities per mm^2 in gonadal adipose tissue of
lean and obese mice. Since most computational models for VEGF signaling have been developed in tumors,
identifying potential differences between tumors and adipose tissue is an important goal. Thus, data for
vessel size and vessel density in mouse tumors were included in our analysis. We also included *ex vivo*
studies reporting the capillary basement membrane thickness in retina, muscle, heart, brain, and kidney
from mice and rats. To analyze the binding kinetics of VEGF-A to its receptors, we included *in vitro* studies
that measured binding affinities (K_d), association rate constants (k_{on}), and dissociation rate constants (k_{off})
using radioligand assays and surface plasmon resonance. We excluded papers if they (1) did not provide
enough information to extract average values and standard errors (e.g., the standard deviation was
reported without sample size or only the average value was reported) or (2) reported very large values
considered as outliers (e.g., the vessel size in tumors reported by Koyama *et al.*[17] is 10 times larger than
values in other studies). We did not include non-English papers.
We did not analyze VEGF-B and VEGF-C, although they influence lipid metabolism and lymphangiogenesis,
which are important mechanisms in obesity, because of a lack of information about their binding affinities.
Instead, we gathered and reported a list of binding data for VEGF-B and VEGF-C to VEGFRs regardless of
measurement method.
110
111 ### Data extraction
112
The data were extracted from the full text or supplementary materials of the selected papers. We used
minor data extraction techniques in the following cases. (1) When information about average values and
standard errors was provided by an image, we used ImageJ V1.53k (https://imagej.net/) to extract that
information[18]. (2) When papers provided a graph showing association and dissociation phases of VEGF-A
binding without providing values, we estimated kinetic data by fitting 1:1 Langmuir equations to the
provided graph. This data extraction process was performed by using an in-house code written in Python
Programming Language V3.11.0 (http://www.python.org/). The detailed approaches for data extraction,
data fitting, and ImageJ analysis are provided in the *Supplementary information*.
All data sets include the following information: the first author's name, year published, data collection
techniques, measurements (i.e., means), and standard errors. For geometric data, we additionally
extracted the following: species, strain, sample size, sex, age, diet or status (e.g., healthy or diabetic mice),
duration of diet, body weight, and location of tissue. For tumor data, we included tumor cell lines, the
location in the mouse body where the tumors were injected, and antibodies used for vessel staining. For
binding affinity data, we extracted the ligands used, the receptors, and their sources. The data extraction
was done independently by two authors (Yunjeong Lee and Keith Lionel Tukei), and any disagreement was
resolved by discussion among three authors (Yunjeong Lee, Keith Lionel Tukei, and Shobhan Kuila).
129
130 ### Data analysis
131
The weighted average and standard deviation of each group were calculated with a random-effects model.
We assumed that the analyzed data followed normal distributions. The weight of the i^{th} study was defined
by $1/(SE_i^2 + T^2)$, where SE_i is the standard error of the i^{th} study and T^2 is the between-study variance.
Cochran's Q-test, the most common way to assess the presence of heterogeneity between studies, was
performed. Two statistics evaluated the level of heterogeneity: (1) I^2, which represents the proportion of
variation between studies among the total variation, and (2) prediction intervals, which show the amount
of dispersion of the observed measurements[19].
139

After calculating the weighted average and standard deviations of groups, we used Welch's t-test to
compare a pair of two groups because (1) the groups are unpaired, (2) the groups have unequal variances,
and (3) the analyzed data were assumed to follow the normal distribution. When we performed a one-
tailed test to identify inequality between more than two groups, we used a one-tailed Welch's t-test with
the Benjamini-Hochberg method for multiple testing. The Benjamini-Hochberg method was chosen since
it is balanced between Type I and Type II errors and its statistical power is stronger than that of the
Bonferroni method, a commonly used multiple-testing correction. The Bonferroni method is more
conservative because it divides the p-value by the number of tests, and it is recommended when the cost
of Type I errors is more expensive than Type II errors[20]. Since the cost of Type I errors in our analysis is
similar to that of Type II errors, we chose the Benjamini-Hochberg method. When we compared more than
two groups, the homogeneity of variances was assessed by Bartlett's test since it (1) is appropriate when
the sample sizes are unequal, and (2) is generally powerful for various variance ratios between groups.
After testing the homogeneity of variance, Welch's ANOVA followed by Dunnett's T3 test was used to
compare means of multiple groups. Dunnett's T3 test was chosen because it is appropriate for a small
sample dataset. All statistical analysis was performed using the "metafor", "weights", "stats", and
"PMCMRplus" libraries in R Project for Statistical Computing, RStudio V4.3.0 (http://www.r-project.org/).
The full codes and Excel files are available on the Open Science Framework (doi: 10.17605/OSF.IO/N9CEY).
157
158 ### Surface Plasmon Resonance (In Vitro)
159
Surface plasmon resonance (SPR) experiments were performed at 25°C using the Reichert 4SPR system
(Reichert, Inc., USA) with PEG-coated gold sensor chips containing 10% COOH (Reichert, Inc., USA
#13206061). The chip was divided into four flow cells: growth factors were immobilized in channels 1 or
3, leaving channels 2 or 4 blank as references. The running buffer was 1x HBS-EP pH 7.4 (10 mM HEPES,
3 mM EDTA, 150 mM NaCl, 0.005% Tween-20). The ligands NRP1 (Cat. #3870-N1-025/CF, R&D Systems)
and VEGFR2 (Cat. #357-KD-050/CF, R&D Systems) were immobilized using an amine coupling method. EDC
(1-ethyl-3-[3-dimethylaminopropyl]carbodiimide, 40 mg/mL), and NHS (N-hydroxysuccinimide, 10 mg/mL)
were dissolved in water, mixed, and injected at 10 µL/min for 7 minutes to activate the surface. Proteins
were diluted to 30 µg/mL in 10 mM acetate buffer (pH 4.0) and injected at 10 µL/min until the
immobilization level reached ≥2000 RU, based on an Rmax target of less than 200 RU to minimize mass
transfer effects. The surface was deactivated by injecting 1M ethanolamine hydrochloride-NaOH (pH 8.5)
for 7 minutes at 10 µL/min.
172
For kinetic analysis, analyte VEGF-A (Cat. #293-VE-010, R&D Systems) was injected at concentrations of 50,
25, 12.5, 6.25, and 3.12 nM, and the association and dissociation curves were fitted using a 1:1 Langmuir
binding model in TraceDrawer ver.1.8.1. Sensorgrams were visually inspected, and the fitting was validated
by the $\chi2$-to-Rmax ratio (<0.10), ensuring a reliable 1:1 interaction model. Raw sensorgrams (3.12–50 nM)
were aligned, and nonspecific binding was subtracted using reference channel sensorgrams. Global fitting,
considered more accurate than single-curve fitting, was applied using nonlinear least-squares analysis in
TraceDrawer to determine association (k_{on}) and dissociation (k_{off}) rates across multiple response curves.
The results are presented as mean ± standard error.
181
182 ## Results
183
184 ### Study selection
185
186 A total of 76 studies were analyzed (Figure 2; n = the number of studies): n = 10 for adipocyte size; n = 11
187 for vessel size; n = 12 for vessel density; n = 34 for capillary basement membrane thickness; n = 8 for VEGF-

A binding affinity to VEGFR1; $n = 12$ for VEGF-A binding affinity to VEGFR2; $n = 10$ for VEGF-A binding affinity NRP1; $n = 3$ for VEGF-A to VEGFR1 association and disassociation rates; $n = 6$ for VEGF-A to VEGFR2 association and disassociation rates; and $n = 3$ for VEGF-A to NRP1 association and dissociation rates. After assessing full-text articles, we included only studies that provided information enabling the extraction of average values and standard errors, since those values were essential in the analysis. A total of 21 studies were excluded from the final selection. The rationales for their exclusion were as follows:

1) *Lack of data to extract mean ± standard error (15 studies)*: Lang et al.[21], Sletta et al.[22], Cai et al.[23], Bloodworth et al.[24], Welt et al.[25], Colotti et al.[26], Geretti et al.[27], Guerrin et al.[28], Mamluk et al.[29], Papo et al.[30], Shibuya et al.[31], Soker et al.[32], Tillo et al.[33], Pan et al.[34], and Vintonenko et al.[35] were excluded because the mean or standard error could not be extracted.

2) *Outliers (3 studies)*: Koyama et al.[17] was excluded because the reported vessel size was 10 times larger than those in other studies. Fuh et al.[36] and Teran and Nugent[37] were also excluded because their surface plasmon resonance analysis reported too large VEGF-A binding affinities (about 30 times and 700 times larger than other studies, respectively). In Teran and Nugent, especially, the rationale for this discrepancy was not reasonable. They claimed two possibilities as the reason for this discrepancy: (1) experimental settings different from those in the cell-based binding assay, and (2) the properties of the Fc-receptor chimeras used for surface plasmon resonance. However, we could not find such a noticeable difference in binding affinities across other studies using surface plasmon resonance and cell-based assays (please refer to the supplementary tables). Also, studies using surface plasmon resonance (e.g., Mamer et al.[38]and Papadopoulos et al.[39]) used Fc-receptors from the same source (R&D systems) as Teran and Nugent and yielded binding affinities similar to those reported in other studies. The discrepancy might be caused by the use of fibroblast growth factor receptors for nonspecific binding.

3) *Non-murine geometrical data (1 study)*: Belligoli et al.[40] was excluded because they reported human capillary basement membrane thickness and we focused on murine tissues.

4) *Lack of tissue data (2 studies)*: Fraselle-Jacobs et al.[41], which measured capillary basement membrane thickness in adipose tissue, was excluded from the analysis because we found only one paper that provided data for capillary basement membrane thickness in adipose tissue. Smith et al.[42], which measured capillary basement membrane thickness in the inner ear, was excluded for the same reason.

Adipocyte diameter is larger in the adipose tissue of diet-induced obese mice than in lean mice

After selecting papers on the basis of eligibility criteria and extracting data, we first investigated adipocyte size in the most well-studied adipose tissue, the gonadal adipose tissue of lean and diet-induced obese mice. We analyzed this data because of the exceptionally large occupation of intracellular space in adipose tissue compared with tumors. Indeed, while tumor cells occupy about 40% of tumor[13–15] volume, the adipocytes' volume is about 90% of the total adipose tissue volume[11] (Figure 1A). The large diameter of adipocytes contributes to their large volume percentage and the smaller interstitial space available for VEGF. This is a distinctive feature of adipose tissue compared with tumors. We analyzed 22 measurements from 10 studies (Table S1 and Figure 3). The main result is that adipocytes of obese mice were significantly larger than those of lean mice: the adipocytes of diet-induced obese mice were about 78% larger than those of lean mice (Table 1; 71 ± 5.3 µm vs. 40 ± 4.3 µm, p < 0.001).

Smaller vessel size and higher vessel density in adipose tissue compared with tumors

Adipose tissue is known to be highly vascularized[1]. As vessel size and vessel density are useful measures
of vascular morphology, we compared vessel size and vessel density between obese and lean mice (Table
1, Figure 4, Tables S2, S4, Figures S1, S2, S4, and S5). We found that obese mice, in comparison with lean
mice, had half the vessel density but similar vessel size (vessel density: 350 ± 68 /mm^2 vs. 720 ± 120 /mm^2,
$p = 0.011 < 0.05$; vessel size: 64 ± 10 μm^2 vs. 51 ± 6.2 μm^2, $p = 0.159 > 0.05$).
Tumor tissue is also known for its high vascularity[43], and tumor vessels in lean mice were two times larger
than in lean and obese adipose tissue (Figures 4A and S3, and Table S3; lean adipose vs. tumor: 51 ± 6.2
μm^2 vs. 120 ± 18 μm^2, $p = 0.006 < 0.01$; obese adipose vs. tumor: 64 ± 10 μm^2 vs. 120 ± 18 μm^2, $p = 0.012$
< 0.05). However, the tumor was four and nine times less vascularized than were lean and obese adipose
tissue, respectively (Figures 4B and S6, and Table S5; lean adipose vs. tumor: 720 ± 120 /mm^2 vs. 84 ± 26
/mm^2, $p = 0.002 < 0.01$; obese adipose vs. tumor: 350 ± 68 /mm^2 vs. 84 ± 26 /mm^2, $p = 0.003 < 0.01$).
Overall, our data show that obesity reduces vascular density but not vessel size in adipose tissue and that
adipose tissue has higher vascular density but smaller vessel size than tumors.

Table 1. Analysis of adipocyte size, vessel size, and vessel densities in adipose tissues and tumors

	Mean ± Standard Error (no. measurements)	Alternative hypothesis	Difference between means (p-value)
Adipocyte size			
Lean adipose tissue	40 ± 4.3 μm ($n = 9$)	Lean ≤ obese	**Significant**
Obese adipose tissue	71 ± 5.3 μm ($n = 13$)		**(p = 9.02 × 10^{-5}<0.01)**
Vessel size			
Lean adipose tissue	51 ± 6.2 μm^2 ($n = 6$)	Lean ≤ obese	Not significant (p = 0.159 > 0.05)
Obese adipose tissue	64 ± 10 μm^2 ($n = 9$)	Lean ≤ tumor	**Significant (p = 0.006 < 0.01)**
Tumor	120 ± 18 μm^2 ($n = 9$)	Obese ≤ tumor	**Significant (p = 0.012 < 0.05)**
Vessel density			
Lean adipose tissue	720 ± 120 /mm^2 ($n = 7$)	Lean ≥ obese	**Significant (p = 0.011 < 0.05)**
Obese adipose tissue	350 ± 68 /mm^2 ($n = 9$)	Lean ≥ tumor	**Significant (p = 0.002 < 0.01)**
Tumor	84 ± 26 /mm^2 ($n = 11$)	Obese ≥ tumor	**Significant (p = 0.003 < 0.01)**

Effect of obesity on capillary basement membrane thickness

The capillary basement membrane is a component of the extracellular matrix in tissue. Since the thicker
capillary basement membrane occupies a larger volume fraction in adipose tissue, its thickness affects the
interstitial space volume in the tissue. Additionally, it is known that capillary basement membrane
thickening is associated with diabetes, and diabetes is one of the common comorbidities of obesity. Thus,

to determine the relationship between capillary basement membrane thickness and obesity, we analyzed
34 studies that measured capillary basement membrane thickness in lean and fat mice and rats (Table 2,
Figure 5, Table S6, and Figures S7–S8). The measurements from the retina, muscle, and heart were
examined in both categories of mice and rats since measurements from other tissues (e.g., brain and
kidney) were not found for obese mice and rats. Interestingly, capillary basement membrane thickness in
tissues of obese mice and rats was similar to that of lean mice and rats, with no significant difference
(obese vs. lean: 104 ± 14 nm vs. 94 ± 5 nm, $p = 0.536 > 0.05$). Our result indicates that obesity does not
affect capillary basement membrane thickness.
266
In order to examine if capillary basement membrane thickness varies across tissues, we examined capillary
basement membrane measurements from multiple tissues: retina, muscle, heart, brain, and kidney (Table
2, Table S6, Figures S9–S14). These tissues were chosen because they are the most well-studied tissues in
the field in regard to capillary basement membrane thickness. The obese mouse and rat data were
excluded from the retina, muscle, and heart datasets for fair comparison, since they did not include brain
and kidney measurements. The test for homogeneity of variance showed significantly different variances
across tissues ($p = 1.30 \times 10^{-5} < 0.001$). From the following Welch's ANOVA and Dunnett's T3 test, the retina
had similar capillary basement membrane thickness as muscle (101 ± 6 nm vs. 79 ± 10 nm, $p = 0.555 > 0.05$;
Table S10), but the retina had significantly different thickness compared with heart and brain (retina:
101 ± 6 nm vs. heart: 69 ± 7 nm, $p = 0.031$; retina vs. brain: 101 ± 6 nm vs. 75 ± 5 nm, $p = 0.019$). Capillary
basement membrane thickness in muscle was similar to that in heart and brain (muscle vs. heart: 79 ± 10
nm vs. 69 ± 7 nm, $p = 0.990 > 0.05$; muscle vs. brain: 79 ± 10 nm vs. 75 ± 5 nm, $p = 1.000 > 0.05$). We did
not analyze this in adipose tissue, because of the single datum (109 ± 11 nm from Fraselle-Jacobs *et al.*[41]).
Capillary basement membrane thickness was greatest in the kidney (181 ± 17 nm), and it was about 2–3
times thicker than in other tissues (retina vs. kidney: $p < 0.01$; muscle vs. kidney: $p < 0.001$; heart vs. kidney:
$p < 0.001$; brain vs. kidney: $p < 0.001$). Our result suggests that capillary basement membrane thickness
varies across tissues.
284
285 **Table 2. Capillary basement membrane thickness in mouse and rat tissues**

	Mean ± Standard Error (no. measurements)	Null hypothesis	Difference between means (p-value)
All tissues			
Lean mice	94 ± 5 nm ($n = 51$)	Lean = obese	Not significant ($p = 0.536 > 0.05$)
Obese mice	104 ± 14 nm ($n = 7$)		
Tissue-specific			
Retina	101 ± 6 nm ($n = 38$)	Retina = muscle = heart = brain = kidney	[ANOVA] $F(4, 22) = 10.8$ $p = 5.42 \times 10^{-5} < 0.001$
Muscle	79 ± 10 nm ($n = 7$)		
Heart	69 ± 7 nm ($n = 6$)		
Brain	75 ± 5 nm ($n = 14$)		
Kidney	181 ± 17 nm ($n = 16$)		

286
287
288 ***Binding data for VEGF-A:VEGFR2 and VEGF-A:NRP1 measured by surface plasmon resonance***
289
We measured the binding rates and affinities for VEGF-A with NRP1 (Figure S15A), as kinetics and affinity
studies were lacking. The VEGF-A:VEGFR2 binding rates were also measured as a positive control. The
VEGF-A:NRP1 binding affinity (K_d) was measured as 6.36 ± 1.07 nM, with an association rate (k_{on}) of

$7.96 \pm 2.15 \times 10^5$ M^{-1} s^{-1} and dissociation rate (k_{off}) of $1.56 \pm 0.55 \times 10^{-3}$ s^{-1}. The kinetics rate for VEGF-A
and VEGFR2 was determined with binding affinity $K_d = 520 \pm 250$ pM, which is in a pM range similar to
those reported in other studies (Figure S15B)[44–46]. The association rate (k_{on}) for VEGF-A:VEGFR2 was
calculated to be $6.24 \pm 0.46 \times 10^5$ M^{-1}s^{-1}, with the dissociation rate (k_{off}) of $3.18 \pm 1.98 \times 10^{-4}$ s^{-1}. These
measurements were included in our analysis. We measured a lower binding affinity and slower
dissociation rate in VEGF-A's binding to NRP1 in comparison with its binding to VEGFR2.
299
300 ***Stronger binding of VEGF-A to VEGFR1 than to VEGFR2 and NRP1***
301
VEGF receptors have distinct functions in angiogenesis; thus, comparing their binding affinities offers
receptor-level insights into differential VEGFR signaling in obesity. We analyzed 21 studies that measured
VEGF-A binding affinity to VEGFR1, VEGFR2, and the co-receptor NRP1, using either radioligand assays or
surface plasmon resonance (Table 3, Figure 6A, Tables S7–S9, Figures S16–S18). As expected, the binding
affinity of VEGF-A to VEGFR1 was significantly stronger than to VEGFR2, being about six times stronger
(34 ± 12 pM vs. 210 ± 60 pM, $p = 0.017 < 0.05$). On the other hand, the binding affinity of VEGF-A to NRP1
was not significantly weaker than the VEGF-A:VEGFR1 affinity, although it was about 24 times weaker
(34 ± 12 pM vs. 820 ± 350 pM, $p = 0.065 > 0.05$). Similarly, VEGF-A binding affinity to NRP1 was not
significantly weaker than to VEGFR2, although it was four times weaker (210 ± 60 pM vs. 820 ± 350 pM,
$p = 0.088 > 0.05$). Overall, our study shows the strongest binding affinity of VEGF-A to VEGFR1 and its
weakest binding affinity to NRP1.
313
314 **Table 3. Analytic K_d values for VEGF-A binding to its receptors**

Receptors	Mean ± Standard Error (no. measurements)	Alternative hypothesis	Difference between means (p-value)
VEGFR1	34 ± 12 pM ($n = 10$)	VEGFR1 ≤ VEGFR2	**Statistical difference (p = 0.017 < 0.05)**
VEGFR2	210 ± 60 pM ($n = 17$)	VEGFR1 ≤ NRP1	No difference (p = 0.065 > 0.05)
NRP1	820 ± 350 pM ($n = 12$)	VEGFR2 ≤ NRP1	No difference (p = 0.088 > 0.05)

315
Association (k_{on}) and dissociation (k_{off}) rate constants allow us to predict not only the equilibrium state of
a ligand-receptor interaction, but also how fast the system responds to changes in the concentration of
the ligand or to another competitor. We analyzed data for VEGFR1, VEGFR2, and NRP1 binding kinetics
with VEGF-A. The association rates of VEGF-A and these receptors were not significantly different,
although the association rate of VEGFR1 was larger than that of VEGFR2, as expected (Table 4, Figure 6B,
Table S7–S9, Figure S19–S21; $11 \pm 9.3 \times 10^6$ M^{-1}s^{-1} vs. $4.6 \pm 1.7 \times 10^6$ M^{-1}s^{-1}, $p = 0.274 > 0.05$). The
association rate of NRP1 ($1.3 \pm 0.54 \times 10^6$ M^{-1}s^{-1}) was four to eight times smaller than those of VEGFR1
and VEGFR2 ($p = 0.274 > 0.05$ and $p = 0.238 > 0.05$, respectively). The dissociation rate constants for all
receptors were also similar, although the mean for VEGFR1 was eight times smaller than the mean for
VEGFR2 (Table 4, Figure 6C, Table S7–S9, Figure S22–S24; $1.0 \pm 0.89 \times 10^{-4}$ s^{-1} vs. $7.7 \pm 4.3 \times 10^{-4}$ s^{-1}), and
that of NRP1 was larger by an order of magnitude ($3.5 \pm 1.6 \times 10^{-3}$ s^{-1}). Overall, our data suggest that
association and dissociation rates of VEGF-A with these receptors do not differ significantly, while its
binding affinities for VEGFR1 and VEGFR2 do differ significantly.
329

Table 4. Analytic association and dissociation rate values of VEGF-A binding to VEGFR1, VEGFR2, and NRP1

	Mean ± Standard Error (no. measurements)	Alternative hypothesis	Difference between means (p-value)
Association rates			
VEGFR1	$11 \pm 9.3 \times 10^6$ M^{-1}s^{-1} ($n = 3$)	VEGFR1 ≥ VEGFR2	No difference (p = 0.274 > 0.05)
VEGFR2	$4.6 \pm 1.7 \times 10^6$ M^{-1}s^{-1} ($n = 9$)	VEGFR1 ≥ NRP1	No difference (p = 0.274 > 0.05)
NRP1	$1.3 \pm 0.54 \times 10^6$ M^{-1}s^{-1} ($n = 3$)	VEGFR2 ≥ NRP1	No difference (p = 0.238 > 0.05)
Dissociation rates			
VEGFR1	$1.0 \pm 0.89 \times 10^{-4}$ s^{-1} ($n = 3$)	VEGFR2 ≥ VEGFR1	No difference (p = 0.123 > 0.05)
VEGFR2	$7.7 \pm 4.3 \times 10^{-4}$ s^{-1} ($n = 9$)	NRP1 ≥ VEGFR1	No difference (p = 0.123 > 0.05)
NRP1	$3.5 \pm 1.6 \times 10^{-3}$ s^{-1} ($n = 3$)	NRP1 ≥ VEGFR2	No difference (p = 0.123 > 0.05)

We also examined the binding affinities of VEGF-B and VEGF-C for VEGF receptors (Table 5). VEGF-B binds to VEGFR1 and NRP1, while VEGF-C binds to VEGFR2 and VEGFR3. VEGF-B includes two isoforms, VEGF-B167 and VEGF-B186, with VEGF-B167 predominantly expressed in most normal tissues[47]. VEGF-B186 must be proteolytically processed to bind to NRP1, while VEGF-B167 binds to NRP1 without cleavage[48]. From the data mining, we found that VEGF-B binds to VEGFR1 and the b1 domain of NRP1 with binding affinities of 114 pM and 36 µM, respectively. VEGF-B peptides bind to the b1 domain of NRP1 with binding affinities of 0.39–9.55 µM. Most studies show that the mature VEGF-C and its mutants bind to VEGFR2 and VEGFR3 with binding affinities in the nM range, while Joukov *et al*.[49] reported a pM range of K$_d$ for VEGF-C:VEGFR2 and VEGFR3. These different binding affinities are possibly due to different measurement methods or cells used to produce recombinant VEGF-C. Due to the lack of data, we did not analyze them using a random-effects model.

Table 5. Binding affinities of VEGF-B and VEGF-C to VEGFRs and NRP1

Ligand	Receptor	Method	Measurements	Reference
Mouse VEGF-B186	VEGFR1[†]	Radioligand assay (Competitive binding with human VEGF-A165)	114 pM	50
Full length human VEGF-B176	Human NRP1-b1[‡]	Surface plasmon resonance (VEGF-B167 was immobilized)	36 µM	51
Human VEGF-B167 peptide		Surface plasmon resonance (NRP1-b1 was immobilized)	0.39 µM	
VEGF-B186 peptide[†]			9.55 µM	
Human △N△C[§]	VEGFR2[†]	Radioligand assay (Saturating binding assay; Scatchard analysis)	410 pM	49
	VEGFR3[†]		135 pM	

Recombinant human VEGF-C	Human VEGFR2	Surface plasmon resonance	22 nM	52
	Human VEGFR3		4.4 nM	
Human VEGF-C156[¶]	Human VEGFR3		115 nM	
Human D123A/Q130A double mutant VEGF-C	Human monomeric VEGFR3 ECD (D1–7) [††]	Isothermal titration calorimetry	34 nM	53
Human C137A mutant VEGF-C	Human monomeric VEGFR3 ECD (D1–2)		250 nM	
	Human monomeric VEGFR3 ECD (D1–3)		140 nM	
	Human monomeric VEGFR3 ECD (D1–5)		3.7 nM	
	Human monomeric VEGFR3 ECD (D1–7)		5.6 nM	
	Human D5 double mutant 5EA of VEGFR3 (D1–5)		12 nM	

[†]Species were not specified.

[‡]NRP1-b1: b1 domain of NRP1.

[§]△N△C: a VEGF-C mutant mimicking mature VEGF-C. The N- and C-terminal propeptides were deleted at
or near the proteolytic processing sites.

[¶]VEGF-C156: a mutant generated by replacing the second conserved Cys (Cys[156]) residue of the
recombinant processed VEGF-C (△N△C) by a Ser residue

[††]Monomeric VEGFR3 ECD (D1–X): a VEGFR3 extracellular domain that is C-terminal truncated, soluble,
and monomeric. It has domains from the membrane proximal domains, D1 to DX, where X is a number
from 2 to 7.

Significant difference in binding affinities between radioligand and SPR assays, except for VEGF-A:NRP1

The binding affinity is a specific type of equilibrium constant, calculated by dividing the dissociation rate
"constant" by the association "constant". Thus, theoretically, the value should be consistent under similar
observation conditions. Nevertheless, different techniques to detect ligand–receptor binding may yield
variations in binding kinetic measurements due to different experimental settings. In order to assess
variability in VEGF-A binding affinity measurements, we analyzed data from radioligand assays and SPR.
The analysis of 21 studies showed that the VEGF-A binding affinity for VEGFR1 measured by radioligand
assay was about eight times weaker than that measured by SPR: 46 ± 15 pM vs. 5.9 ± 2.6 pM, respectively
(Table 6, Figure S25–S27). Also, the binding affinity of VEGF-A to VEGFR2 measured by radioligand assay
was about six times weaker than that measured by SPR (Figure S28 and S29; radioligand assay vs. SPR;
310 ± 110 pM vs. 51 ± 15 pM) while radioligand assay showed seven-times stronger binding affinities for

NRP1 (Figure S30 and S31; 0.36 ± 0.08 nM vs. 2.69 ± 1.96 nM). We applied Welch's t-test to compare K_d
values measured by radioligand assay and SPR for each receptor. All results of t-tests for VEGF receptors
except NRP1 were significant (p = 0.036 < 0.05 for VEGFR1; p = 0.048 < 0.05 for VEGFR2; p = 0.346 > 0.05
for NRP1). Our results indicate that radioligand assay and SPR report significantly different VEGFR binding
affinities.
373
374 **Table 6. Analytic K_d values for VEGF-A binding to its receptors measured by radioligand and SPR (surface**
375 **plasmon resonance) assays**

Receptors	Mean ± Standard Error (no. measurements)		Alternative hypothesis	Difference between means (p-value)
VEGFR1	Radioligand	46 ± 15 pM (n = 7)	Radioligand K_d ≠ SPR K_d	**Significant (p = 0.036 < 0.05)**
	SPR	5.9 ± 2.6 pM (n = 3)		
VEGFR2	Radioligand	310 ± 110 pM (n = 8)		**Significant (p = 0.048 < 0.05)**
	SPR	51 ± 15 pM (n = 9)		
NRP1	Radioligand	0.36 ± 0.08 nM (n = 9)		No difference (p = 0.346 > 0.05)
	SPR	2.69 ± 1.96 nM (n = 3)		

376
377
378 **Discussion**
379
The prevalence of obesity in the United States reached 42% in 2021[54], and obesity causes severe
cardiovascular diseases. Adipose tissue expansion in obesity requires angiogenesis, but how to control
angiogenesis in obesity is still unknown. Systems biology can provide new insight into this problem by
incorporating biological data and computational models. Several studies have constructed computational
models to understand pathological angiogenesis in cancer or peripheral artery diseases[13,55]. However,
applying the same models to adipose tissue requires adaptation to its unique microenvironment,
specifically by incorporating adipose tissue-specific data. Thus, prior to developing obesity models, it is
important to identify and analyze data affecting angiogenic signaling in the targeted tissue. The work in
this paper yielded the following five key findings: (1) obese mice have 78% larger adipocytes than lean
mice; (2) obesity reduces vessel density but does not affect vessel size, and adipose tissue has smaller
vessel size but higher vessel density than tumors; (3) obesity does not affect capillary basement membrane
thickness, as opposed to what has been reported in diabetes; (4) by standardizing the binding rates, we
confirmed that VEGF-A's strongest binding is to VEGFR1 rather than to VEGFR2 or NRP1; and (5) the
binding affinities measured by radioligand assay and SPR are significantly different. From these key findings,
our study will enable the development of adipose tissue-specific computational models.
395
396 ***Obesity increases adipocyte size and reduces vessel density but not vessel size***
397
Our analysis determined that in obese mice, adipocyte diameter is increased by 78% and vessel density is
reduced by 51%, consistent with findings from a previous meta-analysis of human adipose tissue[56].
However, vessel size may not be affected by obesity, as suggested by our analysis. While we did not identify
studies statistically comparing vessel size between lean and obese human subjects, representative
histological images with stained vessels appear to show that the vessels in their adipose tissues are of
similar size[57]. In obesogenic conditions, adipocytes undergo hypertrophy (cell enlargement), which

reduces vascular density and creates hypoxic conditions within the tissue, making these processes crucial for inclusion in a computational model of obesity. Overall, our study confirms the association of obesity with adipocyte size and vessel density and suggests its non-association with vessel size.

Contribution of our study to the successful development of adipose-tissue models; difference between adipose tissue and tumor vasculature

Because both tumors and obesity are affected by hypoxia, one may hypothesize that vessels in the two conditions may be similar. Hence, our work aimed to identify (1) any similarities that would enable computational tumor models to be applied to obesity, or (2) any differences that would justify novel obesity-specific computational models. To address this, we examined the differences in vascular morphologies between adipose tissue and tumors, demonstrating that adipose tissues exhibit a higher vessel density but a smaller vessel size than tumors. Specifically, the vessel size in adipose tissue is approximately half that in tumors, while vessel density is four to nine times higher. Consequently, the surface area of vessels per unit volume in adipose tissue may be 3–6 times greater than in tumors.

These differences that we observed between tumors and obesity may be understood through their differing mechanisms of hypoxia and angiogenesis. In obesity, hypoxia is induced as vessel densities are reduced by the enlargement of adipocytes, while in tumors hypoxia results from poor oxygen diffusion from abnormal vasculature[58,59]. Tumors exhibit chaotic vessel formation due to their continuous pro-angiogenic state and rapid neovascularization. This rapid angiogenesis limits the development of mature vessel structures and results in larger, irregular vessels often concentrated at the periphery rather than throughout the tissue[60,61]. In contrast, adipose tissue has unique features. The stromal vascular fraction in the adipose tissue supports a well-organized, hierarchical vasculature where each adipocyte is adjacent to at least one vessel, creating an evenly distributed network across the tissue[62,63].

If a tumor model is adopted for obesity, this significant difference would lead to incorrect predictions of VEGF-A distributions in adipose tissue (e.g., concentrations of free VEGF-A, interstitial matrix-bound VEGF-A, and receptor-bound VEGF-A) and potentially affect predictions of anti-VEGF drug efficacy. Instead, the development of adipose-specific models, with parameters defined here, would enable researchers to identify the most effective obesity treatment.

Why does obesity not affect capillary basement membrane thickness?

We sought to understand whether obesity alters capillary basement membrane (CBM) thickness in adipose tissue, but could not because there were no data available. We instead examined three organs from obese mice and rats—retina, muscle, and heart—and compared them with their correlates in lean mice and rats. This analysis showed that CBM thickness is not altered by obesity in non-adipose tissue. Thus, in computational modeling, the CBM thickness could be assumed to have the same value in lean and obese conditions. Experimental studies should clarify whether obese and lean adipose tissue also have similar CBM.

The insensitivity of CBM thickness to obesity was unexpected, given that 80% of the studies we analyzed used prediabetic and diabetic mice and rats (40% for each group; only 20% used obese, non-diabetic mice) and that diabetes, a common comorbidity of obesity, has been associated with CBM thickening in diabetic models, particularly in the retina, muscles, brain, and kidneys[64–69]. In fact, CBM thickening has been recognized as a hallmark of diabetic retinopathy, nephropathy, and cardiomyopathy, and in the brain, it is

also associated with Alzheimer's disease[70]. This raises the question: what factors might account for the

lack of a significant effect of obesity on CBM thickness in our study? We suggest two possible explanations

for our findings:

1) **Duration of diabetes**: The duration of diabetes rather than diabetes itself may determine the CBM

thickness[71]. For example, 6–7-month-old Zucker diabetic fatty rats develop diabetes after 3–5 months of

age[72], suggesting that CBM thickening may not have progressed enough to reach significance. Similarly,

C57BL/6J mice develop obesity and diabetes after 4 months on a diet high in fat and simple carbohydrate[73].

However, the mice in Williams *et al.* had undergone only a 4-month high-fat diet and exhibited CBM

thickness comparable to that of lean mice[74].

2) **Obese rodent strains**: The strain of obese rats included in our analysis may also explain our findings.

Studies by Lash *et al.* and Dosso *et al.* on obese Zucker rats[75,76], which are a model for prediabetes[77],

support this possibility. Considering that humans with prediabetes exhibit CBM thickness similar to that

of lean, healthy humans (103 nm vs. 117 nm)[40], it is reasonable to expect that CBM thickness will not differ

significantly between the prediabetic Zucker rats and lean rats.

We sought to determine if CBM thickness varies across tissues, in order to decide whether computational

models should consider this aspect of the targeted tissue. Importantly, tissues have phenotypically

different capillaries depending on their function. For example, kidneys have continuous fenestrated

capillaries, with continuous basement membrane and fenestrated endothelium that allows large

molecules to pass across the wall[78]. On the other hand, heart and muscle have continuous basement

membranes and non-fenestrated endothelium, which allow only small molecules to pass[78,79]. The liver has

discontinuous capillaries with a fragmented basement membrane, allowing the movement of large

molecules for liver metabolism[80]. All tissues we examined in this study have continuous CBMs, so we aimed

to identify any differences in CBM thickness between these tissues of lean mice and rats. The retina and

kidney have higher CBM thicknesses than other tissues, possibly due to different distributions of molecular

components (collagen, nidogen, etc.), cell types, or physical factors (e.g., hydrostatic pressure)[81,82]. For

example, the kidney CBM is composed of two basement membranes: one from the endothelium and

another from the epithelium[83].

Standardized VEGF-A binding rates accelerate the development of a more feasible computational model

Our analysis confirms that VEGF-A binding to VEGFR1 is the strongest and establishes that VEGF-A binding

to NRP1 is the weakest of its binding to receptors. This strong VEGF:VEGFR1 binding is attributed to its

highest association rate (k_{on}) and lowest dissociation rate (k_{off}), whereas VEGF:NRP1 binding displayed the

opposite kinetic profile. The binding affinity (K_d) is calculated by $K_d = k_{off}/k_{on}$. Our standardized values of

VEGF-A binding affinities to VEGFR1 and VEGFR2 are consistent with values used in previous

computational studies, while VEGF-A binding affinity to NRP1 from our study was three times weaker than

the previously used values[13,84,85]. On the other hand, our association and dissociation rate constants

differed by two to four times and 1.3 to 3.5 times from the previously used values, respectively. This

difference might have affected the simulation outcomes of the previous studies[13,84,85] such as the

proportion of VEGF-A among other ligands bound to the same VEGF receptors, because the rate constants

are key factors determining how fast VEGF-A binds to its receptors in competing with other ligands. To our

best knowledge, ours is the first analysis to standardize the VEGF-A binding rates, because only a few

studies have gathered and analyzed VEGF-A binding data[86]. Thus, our analysis enhances the understanding

of complex VEGF-A signaling mechanisms and enables the development of more feasible computational

models by providing standardized binding affinities.

Variation in measurements from different ligand-binding assays: computational modelers should be aware of this when choosing kinetic values

Our analysis revealed that measurement techniques can significantly impact binding affinity values. Specifically, binding affinities are commonly measured using two techniques: the radioligand assay and surface plasmon resonance (SPR). When comparing data from these methods, we found that radioligand assays yield higher binding affinity measurements for VEGF-A:VEGFR1 and VEGF-A:VEGFR2, while no difference was observed for VEGF-A:NRP1. Several factors may explain these notable differences. First, heparin has been shown to reduce the VEGF-A:VEGFR1 binding affinity in porcine aortic endothelial cells[87]. Specifically, K_d was 54 pM in the absence of heparin but increased to 77–118 pM when heparin was present. This suggests that heparan sulfate proteoglycan (HSPG) on the cell membrane likely influences the binding affinity observed in the radioligand assay, which is a cell-based assay. In contrast, SPR is a chip-based assay and may not fully replicate the membrane-associated effects. Second, the different structures of receptors used in the radioligand assay and SPR may also affect the binding affinity. While the radioligand assays that we cited in our analysis used full-length receptors on cells, in SPR reconstituted receptors with no transmembrane and intracellular domains are bound to chips. The absence of these domains and associated membrane stabilization may contribute to differences in VEGF-A binding. For example, previous studies showed that mutation of the transmembrane domain of NRP1 significantly reduced VEGF-A binding[88]. Thus, computational modelers should consider these variations in the measurements of ligand-binding assays to more consciously choose kinetic values and investigate their impact on model outcomes through a sensitivity analysis across the pM to nM range that we identified across these assays.

Study limitations

This study has three limitations: (1) **Small sample sizes in datasets**: The small sample size may overestimate the confidence interval or prediction interval and result in questionable standardized values. The overestimated confidence interval also affects the results of statistical tests in which we compare group means. Indeed, one of our datasets, VEGF-A:NRP1 binding affinities measured by SPR, included only three studies, which yielded wide prediction intervals and insignificant differences in measurements between SPR and the radioligand assay. Additional related studies should be accumulated to reduce the possible overestimation of pooled variance. (2) **Lack of moderator analysis**: Moderator analysis is usually performed to identify the moderators that affect heterogeneity in datasets. Our analysis showed high heterogeneity between studies for each dataset. For example, the lower bound of the 95% prediction intervals for some datasets was smaller than 0 (e.g., −4.7 to 10.0 nM for binding affinities of VEGF-A:NRP1 measured by SPR). Most values of I^2, which indicates the proportion of total variability attributed to between-study variability, exceeded 90%, categorizing the standardized values as having high heterogeneity (I^2 >75%)[89]. However, we lacked the data to perform moderator analysis. We expect that the high heterogeneity between studies in datasets would be caused by different mouse models, cell lines, different ligand–receptor interaction conditions, etc. (3) **Not meta-analysis**: We performed an extensive literature search (*n* = 75 studies) using free-text search-phrase approaches to find targeting measurements. However, we acknowledge that this is not a meta-analysis. A meta-analysis requires an exhaustive literature search using both controlled vocabulary (e.g., MeSH) and text word search. Another best practice for meta-analysis is registration in a database such as PROSPERO[90]. Future studies should build upon the principles that we have established here to elevate the analysis to the level of a meta-analysis.

Conclusions

Our findings will enable advances in three important areas: (1) **Adipose tissue microenvironment and obesity**: Our results show that obesity does not affect capillary basement membrane thickness even though there is increased capillary basement thickness associated with diabetes. Also, obesity decreases vessel density in adipose tissue while vessel size remains the same. These findings help to identify which structures within the vascular microenvironment are affected by obesity. Further, our results align with previous studies reporting that obese mice have larger adipocytes than lean mice[91,92]. (2) **Understanding of vascular dysfunction**: Our observation that adipose tissue has smaller vessels and higher vessel density compared with tumors underlies the importance of investigating tissue-specific vascular phenotypes. These different phenotypes indicate that tumor and adipose tissue angiogenesis may be differently regulated—despite theories to the contrary, which focus on the hypoxia in both environments. Further, by identifying vessel morphology and considering it alongside VEGF-A kinetic parameters, researchers can examine crosstalk between adipose tissue and tumors via VEGF-A signaling. Indeed, there is evidence that adipocyte-induced VEGF–mTOR signaling increases tumor cell growth and that obesity upregulates the signal[93]. Thus, the enhanced knowledge provided by our analysis should be helpful for the development of angiogenesis-targeting treatment in both tumor and obesity research. 3) **Computational modeling**: With the consolidated data that we provide on adipose tissue and adipose vascular characteristics, systems biology researchers can develop much-needed computational models of obesity. **Therefore, this work has broad potential impact on** biological and biomedical research.

Acknowledgments

We thank Dr. Vincenza Cifarelli for providing invaluable support through discussions. This material is based upon work supported by the National Science Foundation under Grant No. 1923151. Research reported in this publication was also supported by the National Institute of Heart, Lung, and Blood of the National Institutes of Health under Award Number 7R01HL159946-02. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation or the National Institutes of Health.

Data availability statement

The data that support the findings of this study are available in the methods and/or supplementary material of this article.

Conflict of interests

The authors declare no conflicts of interest.

Author contribution

Yunjeong Lee, Yingye Fang, and Princess I Imoukhuede designed the research. Yunjeong Lee and Keith Lionel Tukei acquired, analyzed, and interpreted the data. Shobhan Kuila acquired and interpreted the SPR data. Xinming Liu acquired the capillary basement membrane thickness and VEGF-B binding affinity data. Yunjeong Lee, Keith Lionel Tukei, Shobhan Kuila, Yingye Fang, and Princess I Imoukhuede were involved in drafting and revising the manuscript.

References

1. Herold, J. & Kalucka, J. Angiogenesis in Adipose Tissue: The Interplay Between Adipose and Endothelial Cells. *Frontiers in Physiology* vol. 11 Preprint at https://doi.org/10.3389/fphys.2020.624903 (2021).

2. Rausch, M. E., Weisberg, S., Vardhana, P. & Tortoriello, D. V. Obesity in C57BL/6J mice is characterized by adipose tissue hypoxia and cytotoxic T-cell infiltration. *Int J Obes* **32**, (2008).

3. Meister, B. M. *et al.* Healthy versus Unhealthy Adipose Tissue Expansion: the Role of Exercise. *Journal of Obesity and Metabolic Syndrome* vol. 31 Preprint at https://doi.org/10.7570/JOMES21096 (2022).

4. Khan, T. *et al.* Metabolic Dysregulation and Adipose Tissue Fibrosis: Role of Collagen VI. *Mol Cell Biol* **29**, (2009).

5. Sung, H. K. *et al.* Adipose vascular endothelial growth factor regulates metabolic homeostasis through angiogenesis. *Cell Metab* **17**, (2013).

6. Wu, L. E. *et al.* Systemic VEGF-A neutralization ameliorates diet-induced metabolic dysfunction. *Diabetes* **63**, (2014).

7. Fang, Y., Kaszuba, T. & Imoukhuede, P. I. Systems Biology Will Direct Vascular-Targeted Therapy for Obesity. *Front Physiol* **11**, (2020).

8. Claffey, K. P., Wilkison, W. O. & Spiegelman, B. M. Vascular endothelial growth factor. Regulation by cell differentiation and activated second messenger pathways. *Journal of Biological Chemistry* **267**, (1992).

9. Sun, K. *et al.* Dichotomous effects of VEGF-A on adipose tissue dysfunction. *Proc Natl Acad Sci U S A* **109**, (2012).

10. Nishimura, S. *et al.* Adipogenesis in obesity requires close interplay between differentiating adipocytes, stromal cells, and blood vessels. *Diabetes* **56**, (2007).

11. Arner, P. & Rydén, M. Human white adipose tissue: A highly dynamic metabolic organ. *J Intern Med* **291**, (2022).

12. Yen, P., Finley, S. D., Engel-Stefanini, M. O. & Popel, A. S. A two-compartment model of VEGF distribution in the mouse. *PLoS One* **6**, (2011).

13. Finley, S. D., Engel-Stefanini, M. O., Imoukhuede, P. I. & Popel, A. S. Pharmacokinetics and pharmacodynamics of VEGF-neutralizing antibodies. *BMC Syst Biol* **5**, (2011).

14. Hao, S. J., Wan, Y., Xia, Y. Q., Zou, X. & Zheng, S. Y. Size-based separation methods of circulating tumor cells. *Advanced Drug Delivery Reviews* vol. 125 Preprint at https://doi.org/10.1016/j.addr.2018.01.002 (2018).

15. Del Monte, U. Does the cell number 10⁹ still really fit one gram of tumor tissue? *Cell Cycle* **8**, (2009).

16. Fang, Y., Cifarelli, V. & Imoukhuede, P. I. Sex differences in VEGF/PDGF ligand and receptor profiles in adipose tissue during weight gain and weight loss. *Under review*.

17. Koyama, S. *et al.* Tumour blood vessel normalisation by prolyl hydroxylase inhibitor repaired sensitivity to chemotherapy in a tumour mouse model. *Sci Rep* **7**, (2017).

18. Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of image analysis. *Nature Methods* vol. 9 Preprint at https://doi.org/10.1038/nmeth.2089 (2012).

19. Borenstein, M. How to understand and report heterogeneity in a meta-analysis: The
difference between I-squared and prediction intervals. *Integr Med Res* **12**, (2023).
20. Krzywinski, M. & Altman, N. Comparing samples—part II. *Nat Methods* **11**, 355–356
(2014).
21. Lang, S. *et al.* Three-dimensional quantification of capillary networks in healthy and
cancerous tissues of two mice. *Microvasc Res* **84**, (2012).
22. Sletta, K. Y. *et al.* Oxygen-dependent regulation of tumor growth and metastasis in human
breast cancer xenografts. *PLoS One* **12**, (2017).
23. Cai, W. *et al.* PET of vascular endothelial growth factor receptor expression. *Journal of*
Nuclear Medicine **47**, (2006).
24. Bloodworth, J. M., Engerman, R. L., Camerini-Dávalos, R. A. & Powers, K. L. Variations in
capillary basement membrane width produced by aging and diabetes mellitus. *Adv*
Metab Disord **1**, (1970).
25. Welt, K., Weiss, J., Koch, S. & Fitzl, G. Protective effects of Ginkgo biloba extract EGb 761
on the myocardium of experimentally diabetic rats. II. Ultrastructural and
immunohistochemical investigation on microvessels and interstitium. *Experimental and*
Toxicologic Pathology **51**, (1999).
26. Colotti, G. *et al.* Neuropilin-1 is required for endothelial cell adhesion to soluble vascular
endothelial growth factor receptor 1. *FEBS Journal* **289**, (2022).
27. Geretti, E., Shimizu, A., Kurschat, P. & Klagsbrun, M. Site-directed mutagenesis in the B-
neuropilin-2 domain selectively enhances its affinity to VEGF165, but not to semaphorin
3F. *Journal of Biological Chemistry* **282**, (2007).
28. Guerrin, M. *et al.* Vasculotropin/Vascular endothelial growth factor is an autocrine
growth factor for human retinal pigment epithelial cells cultued in vitro. *J Cell Physiol* **164**,
(1995).
29. Mamluk, R. *et al.* Neuropilin-1 binds vascular endothelial growth factor 165, placenta
growth factor-2, and heparin via its b1b2 domain. *Journal of Biological Chemistry* **277**,
(2002).
30. Papo, N., Silverman, A. P., Lahti, J. L. & Cochran, J. R. Antagonistic VEGF variants
engineered to simultaneously bind to and inhibit VEGFR2 and $\alpha v\beta 3$ integrin. *Proc Natl*
Acad Sci U S A **108**, (2011).
31. Shibuya, M. *et al.* Nucleotide sequence and expression of a novel human receptor-type
tyrosine kinase gene (flt) closely related to the fms family. *Oncogene* **5**, (1990).
32. Soker, S., Miao, H. Q., Nomi, M., Takashima, S. & Klagsbrun, M. VEGF165 mediates
formation of complexes containing VEGFR-2 and neuropilin-1 that enhance VEGF165-
receptor binding. *J Cell Biochem* **85**, (2002).
33. Tillo, M. *et al.* VEGF189 binds NRP1 and is sufficient for VEGF/NRP1-dependent neuronal
patterning in the developing brain. *Development* **142**, (2015).
34. Pan, Q. *et al.* Neuropilin-1 binds to VEGF121 and regulates endothelial cell migration and
sprouting. *Journal of Biological Chemistry* **282**, (2007).
35. Vintonenko, N. *et al.* Overexpression of VEGF189 in breast cancer cells induces apoptosis
via NRP1 under stress conditions. *Cell Adh Migr* **5**, (2011).

36. Fuh, G., Garcia, K. C. & De Vos, A. M. The interaction of neuropilin-1 with vascular
endothelial growth factor and its receptor Flt-1. *Journal of Biological Chemistry* **275**,
(2000).
37. Teran, M. & Nugent, M. A. Characterization of receptor binding kinetics for vascular
endothelial growth factor-A using SPR. *Anal Biochem* **564–565**, (2019).
38. Mamer, S. B., Wittenkeller, A. & Imoukhuede, P. I. VEGF-A splice variants bind VEGFRs
with differential affinities. *Sci Rep* **10**, (2020).
39. Papadopoulos, N. *et al.* Binding and neutralization of vascular endothelial growth factor
(VEGF) and related ligands by VEGF Trap, ranibizumab and bevacizumab. *Angiogenesis* **15**,
(2012).
40. Belligoli, A. *et al.* Characterization of subcutaneous and omental adipose tissue in
patients with obesity and with different degrees of glucose impairment. *Sci Rep* **9**, (2019).
41. Fraselle-Jacobs, A., Jeanjean, M., Hauser, N. & Remacle, C. Effect of aging on the
morphology of epididymal adipose tissue in the rat. *Exp Gerontol* **22**, (1987).
42. Smith, T. L., Raynor, E., Prazma, J., Buenting, J. E. & Pillsbury, H. C. Insulin-dependent
diabetic microangiopathy in the inner ear. *Laryngoscope* **105**, (1995).
43. Forster, J., Harriss-Phillips, W., Douglass, M. & Bezak, E. A review of the development of
tumor vasculature and its effects on the tumor microenvironment. *Hypoxia* **Volume 5**,
21–32 (2017).
44. Whitaker, G. B., Limberg, B. J. & Rosenbaum, J. S. Vascular Endothelial Growth Factor
Receptor-2 and Neuropilin-1 Form a Receptor Complex that is Responsible for the
Differential Signaling Potency of VEGF165 and VEGF121. *Journal of Biological Chemistry*
276, (2001).
45. Waltenberger, J., Claesson-Welsh, L., Siegbahn, A., Shibuya, M. & Heldin, C. H. Different
signal transduction properties of KDR and Flt1, two receptors for vascular endothelial
growth factor. *Journal of Biological Chemistry* **269**, (1994).
46. Huang, X., Gottstein, C., Brekken, R. A. & Thorpe, P. E. Expression of soluble VEGF
receptor 2 and characterization of its binding by surface plasmon resonance. *Biochem
Biophys Res Commun* **252**, (1998).
47. Li, X., Aase, K., Li, H., Von Euler, G. & Eriksson, U. Isoform-specific expression of VEGF-B in
normal tissues and tumors. *Growth Factors* **19**, (2001).
48. Makinen, T. *et al.* Differential binding of vascular endothelial growth factor B splice and
proteolytic isoforms to Neuropilin-1. *Journal of Biological Chemistry* **274**, (1999).
49. Joukov, V. *et al.* Proteolytic processing regulates receptor specificity and activity of VEGF-
C. *EMBO Journal* **16**, (1997).
50. Olofsson, B. *et al.* Vascular endothelial growth factor B (VEGF-B) binds to VEGF receptor-1
and regulates plasminogen activator activity in endothelial cells. *Proc Natl Acad Sci U S A*
95, (1998).
51. Mota, F. *et al.* Peptides Derived from Vascular Endothelial Growth Factor B Show Potent
Binding to Neuropilin-1. *ChemBioChem* **23**, (2022).
52. Mäkinen, T. *et al.* Isolated lymphatic endothelial cells transduce growth, survival and
migratory signals via the VEGF-C/D receptor VEGFR-3. *EMBO Journal* **20**, (2001).
53. Leppänen, V. M. *et al.* Structural and mechanistic insights into VEGF receptor 3 ligand
binding and activation. *Proc Natl Acad Sci U S A* **110**, (2013).

54. Stierman, B. *et al.* National health and nutrition examination survey 2017–march 2020
prepandemic data files-development of files and prevalence estimates for selected health
outcomes. *Natl Health Stat Report* **2021**, (2021).
55. Wu, F. T. H. *et al.* VEGF and soluble VEGF receptor-1 (sFlt-1) distributions in peripheral
arterial disease: An in silico model. *Am J Physiol Heart Circ Physiol* **298**, (2010).
56. Ye, R. Z., Richard, G., Gévry, N., Tchernof, A. & Carpentier, A. C. Fat Cell Size: Measurement
Methods, Pathophysiological Origins, and Relationships With Metabolic Dysregulations.
Endocrine Reviews vol. 43 Preprint at https://doi.org/10.1210/endrev/bnab018 (2022).
57. Spencer, M. *et al.* Adipose tissue extracellular matrix and vascular abnormalities in
obesity and insulin resistance. *Journal of Clinical Endocrinology and Metabolism* **96**,
(2011).
58. Höckel, M. & Vaupel, P. Tumor hypoxia: Definitions and current clinical, biologic, and
molecular aspects. *Journal of the National Cancer Institute* vol. 93 Preprint at
https://doi.org/10.1093/jnci/93.4.266 (2001).
59. Sørensen, B. S. & Horsman, M. R. Tumor Hypoxia: Impact on Radiation Therapy and
Molecular Pathways. *Frontiers in Oncology* vol. 10 Preprint at
https://doi.org/10.3389/fonc.2020.00562 (2020).
60. Lugano, R., Ramachandran, M. & Dimberg, A. Tumor angiogenesis: causes, consequences,
challenges and opportunities. *Cellular and Molecular Life Sciences* **77**, 1745–1770 (2020).
61. Nagy, J. A., Chang, S. H., Dvorak, A. M. & Dvorak, H. F. Why are tumour blood vessels
abnormal and why is it important to know? *British Journal of Cancer* vol. 100 Preprint at
https://doi.org/10.1038/sj.bjc.6604929 (2009).
62. Ramakrishnan, V. M. & Boyd, N. L. The Adipose Stromal Vascular Fraction as a Complex
Cellular Source for Tissue Engineering Applications. *Tissue Engineering - Part B: Reviews*
vol. 24 Preprint at https://doi.org/10.1089/ten.teb.2017.0061 (2018).
63. Cao, Y. Adipose tissue angiogenesis as a therapeutic target for obesity and metabolic
diseases. *Nature Reviews Drug Discovery* vol. 9 Preprint at
https://doi.org/10.1038/nrd3055 (2010).
64. Carlson, E. C. *et al.* Ultrastructural morphometry of capillary basement membrane
thickness in normal and transgenic diabetic mice. *Anatomical Record - Part A Discoveries
in Molecular, Cellular, and Evolutionary Biology* **271**, (2003).
65. Rodrigues, M., Currier, C. & Yoon, J. Electron microscopy of renal and ocular changes in
virus-induced diabetes mellitus in mice. *Diabetologia* **24**, (1983).
66. Cherian, S., Roy, S., Pinheiro, A. & Roy, S. Tight glycemic control regulates fibronectin
expression and basement membrane thickening in retinal and glomerular capillaries of
diabetic rats. *Invest Ophthalmol Vis Sci* **50**, (2009).
67. Gambaro, G. *et al.* Glycosaminoglycans prevent morphological renal alterations and
albuminuria in diabetic rats. *Kidney Int* **42**, (1992).
68. Yagihashi, S., Goto, Y., Kakizaki, M. & Kaseda, N. Thickening of glomerular basement
membrane in spontaneously diabetic rats. *Diabetologia* **15**, (1978).
69. Johnson, P. C., Brendel, K. & Meezan, E. Thickened cerebral cortical capillary basement
membranes in diabetes. *Arch Pathol Lab Med* **106**, (1982).

70. Mancardi, G. L., Perdelli, F., Rivano, C., Leonardi, A. & Bugiani, O. Thickening of the
basement membrane of cortical capillaries in Alzheimer's disease. *Acta Neuropathol* **49**,
(1980).
71. Campbell, M. J. A. The effect of age and the duration of diabetes mellitus on the width of
the basement membrane of small vessels. *Aust Dent J* **19**, (1974).
72. Yokoi, N. *et al.* A novel rat model of type 2 diabetes: The zucker fatty diabetes mellitus
ZFDM rat. *J Diabetes Res* **2013**, (2013).
73. Surwit, R. S., Kuhn, C. M., Cochrane, C., McCubbin, J. A. & Feinglos, M. N. Diet-induced
type II diabetes in C57BL/6J mice. *Diabetes* **37**, (1988).
74. Williams, I. M. *et al.* Transendothelial Insulin Transport is Impaired in Skeletal Muscle
Capillaries of Obese Male Mice. *Obesity* **28**, (2020).
75. Lash, J. M., Sherman, W. M. & Hamlin, R. L. Capillary basement membrane thickness and
capillary density in sedentary and trained obese Zucker rats. *Diabetes* **38**, (1989).
76. Dosso, A. *et al.* Ocular complications in the old and glucose-intolerant genetically obese
(fa/fa) rat. *Diabetologia* **33**, (1990).
77. Pinhas-Hamiel, O. & Zeitler, P. Prediabetes among Obese Youth. in *Global Perspectives on*
Childhood Obesity: Current Status, Consequences and Prevention (2010).
doi:10.1016/B978-0-12-374995-6.10008-8.
78. Sarin, H. Physiologic upper limits of pore size of different blood capillary types and
another perspective on the dual pore theory of microvascular permeability. *J Angiogenes*
Res **2**, (2010).
79. Maleszewski, J. J., Lai, C. K., Nair, V. & Veinot, J. P. Anatomic considerations and
examination of cardiovascular specimens (excluding devices). in *Cardiovascular Pathology*
27–84 (Elsevier, 2022). doi:10.1016/B978-0-12-822224-9.00013-X.
80. Griffin, C. T. & Gao, S. Building discontinuous liver sinusoidal vessels. *Journal of Clinical*
Investigation vol. 127 Preprint at https://doi.org/10.1172/JCI92823 (2017).
81. Leclech, C., Natale, C. F. & Barakat, A. I. The basement membrane as a structured surface
– role in vascular health and disease. *Journal of Cell Science* vol. 133 Preprint at
https://doi.org/10.1242/jcs.239889 (2021).
82. Baum, O. & Bigler, M. Pericapillary basement membrane thickening in human skeletal
muscles. *American Journal of Physiology - Heart and Circulatory Physiology* vol. 311
Preprint at https://doi.org/10.1152/ajpheart.00048.2016 (2016).
83. Scott, R. P. & Quaggin, S. E. The cell biology of renal filtration. *Journal of Cell Biology* vol.
209 Preprint at https://doi.org/10.1083/jcb.201410017 (2015).
84. Mac Gabhann, F. & Popel, A. S. Model of competitive binding of vascular endothelial
growth factor and placental growth factor to VEGF receptors on endothelial cells. *Am J*
Physiol Heart Circ Physiol **286**, (2004).
85. Wu, F. T. H., Stefanini, M. O., Mac Gabhann, F. & Popel, A. S. A compartment model of
VEGF distribution in humans in the presence of soluble VEGF receptor-1 acting as a ligand
trap. *PLoS One* **4**, (2009).
86. Rusnati, M. *et al.* Exploiting surface plasmon resonance (SPR) technology for the
identification of fibroblast growth factor-2 (FGF2) antagonists endowed with
antiangiogenic activity. *Sensors* **9**, (2009).

87.   Ito, N. & Claesson-Welsh, L. Dual effects of heparin on VEGF binding to VEGF receptor-1
and transduction of biological responses. *Angiogenesis* **3**, (1999).
88.   Roth, L. *et al.* Transmembrane domain interactions control biological functions of
neuropilin-1. *Mol Biol Cell* **19**, (2008).
89.   Higgins, J. P. T., Thompson, S. G., Deeks, J. J. & Altman, D. G. Measuring inconsistency in
meta-analyses. *British Medical Journal* vol. 327 Preprint at
https://doi.org/10.1136/bmj.327.7414.557 (2003).
90.   Pieper, D. & Rombey, T. Where to prospectively register a systematic review. *Systematic
Reviews* vol. 11 Preprint at https://doi.org/10.1186/s13643-021-01877-1 (2022).
91.   Murano, I. *et al.* Dead adipocytes, detected as crown-like structures, are prevalent in
visceral fat depots of genetically obese mice. *J Lipid Res* **49**, (2008).
92.   Vangoitsenhoven, R. *et al.* At similar weight loss, dietary composition determines the
degree of glycemic improvement in diet-induced obese C57BL/6 mice. *PLoS One* **13**,
(2018).
93.   Sahoo, S. S. *et al.* Adipose-Derived VEGF–mTOR Signaling Promotes Endometrial
Hyperplasia and Cancer: Implications for Obese Women. *Molecular Cancer Research* **16**,
(2018).

822
823

Figure legends

Figure 1. Distinctive characteristics of adipose tissue compared with tumor and muscle. (A)
Proportions of tissue volume that are intracellular vs. extracellular in tumor, muscle, and adipose
tissue (ref. from Arner & Rydén[11], Yen et al.[12], Finley et al.[13], Hao et al.[14], and Del Monte[15]). The
adipose tissue has a significantly large proportion of intracellular volume compared with other
tissues. (B) The number of receptors on endothelial cells (EC) measured in tumor, muscle, and
adipose tissue (ref. from Finley et al.[13] and Fang et al.[16]). Tumor has a high level of VEGFR1, muscle
has similar levels of VEGFR1 and VEGFR2, and adipose tissue has a high level of VEGFR2.

Figure 2. Flow chart of the analysis

**Figure 3. Adipocyte diameter (μm) in gonadal adipose tissue in lean and diet-induced obese
mice.** Forest plots of the analysis of adipocyte diameter (μm) measured in adipose tissue of lean
mice (A) and obese mice (B) across studies are shown. The first, third, and last columns represent
the list of references, their weights in the analysis, and measurements with a 95% confidence
interval, respectively. The size of the blue diamond for each study is the visualization of its
statistical weight in the analysis. The two diamonds below the table and the black dashed line
show the combined measurements across studies. The upper diamond represents the mean and
95% confidence interval, and the lower diamond represents the mean and 95% prediction interval.
The results of the heterogeneity test are represented by the estimate of between-studies variance
(τ^2), Cochran's Q-test statistic for heterogeneity (χ^2), degrees of freedom (df), p-value (p), and
proportion of heterogeneity-induced variability among the total variability (I^2). In the scatter plot
(C), the error bar shows weighted mean ± standard error of each group. A one-tailed Welch's t-
test was used for the comparison. Abbreviation: week (wk).

Figure 4. Characteristics of vessels in gonadal adipose tissue in lean and diet-induced obese mice and mouse tumors. The vessel size (A) and the vessel density (B) in adipose tissues and tumors are plotted by study. The error bar represents a weighted mean ± standard error of each group. A one-tailed Welch's t-test (p-value < 0.05) with multiple testing correction was used to compare two tissues. The reference names in tumor data include mouse strain and tumor lines. Abbreviation: week (wk), experiment (exp), and dimethyl sulfoxide (DMSO).

Figure 5. Capillary basement membrane thickness (nm) in lean and obese mouse and rat tissues. The capillary basement membranes in retina, muscle, and heart were compared between lean and obese mice and rats. A two-tailed Welch's t-test (p-value < 0.05) was used to compare CBM thicknesses.

Figure 6. Binding rates of VEGF to VEGFR1, VEGFR2, and NRP1. The respective binding affinities (A; K_d), association rate constants (B; k_{on}, M^{-1} s^{-1}), and dissociation rate constants (C; k_{off}, s^{-1}) of VEGF for VEGFR1, VEGFR2, and NRP1 are plotted. The error bar represents the weighted mean ± standard error of the rate for each receptor. A one-tailed Welch's t-test with multiple testing corrections (p-value < 0.05) was used to compare a pair of groups. Abbreviation: surface plasmon resonance (SPR), human umbilical vein endothelial cell (HUVEC), and porcine aortic endothelial (PAE).

Figure S1. Forest plots of the analysis of vessel size (μm^2) measured in adipose tissue of lean mice across studies. The first, third, and last columns represent the list of references, their weights in the analysis, and measurements with a 95% confidence interval, respectively. The size of the blue diamond for each study represents its statistical weight in the analysis. The two diamonds below the table and the black dashed line show the combined measurements across studies. The upper diamond represents the mean and 95% confidence interval, and the lower diamond represents the mean and 95% prediction interval. The results of the heterogeneity test are represented by the estimate of between-studies variance (τ^2), Cochran's Q-test statistic for heterogeneity (χ^2), degrees of freedom (df), p-value (p), and proportion of heterogeneity-induced variability among the total variability (I^2). Abbreviation: week (wk).

Figure S2. Forest plots of the analysis of vessel size (μm^2) measured in adipose tissue of diet-induced obese mice across studies. The first, third, and last columns represent the list of references, their weights in the analysis, and measurements with a 95% confidence interval, respectively. The size of the blue diamond for each study represents its statistical weight in the analysis. The two diamonds below the table and the black dashed line show the combined measurements across studies. The upper diamond represents the mean and 95% confidence interval, and the lower diamond represents the mean and 95% prediction interval. The results of the heterogeneity test are represented by the estimate of between-studies variance (τ^2), Cochran's Q-test statistic for heterogeneity (χ^2), degrees of freedom (df), p-value (p), and proportion of heterogeneity-induced variability among the total variability (I^2). Abbreviation: week (wk).

Figure S3. Forest plots of the analysis of vessel size (μm²) measured in mouse tumors across
studies. The first, third, and last columns represent the list of references, their weights in the
analysis, and measurements with a 95% confidence interval, respectively. The size of the blue
diamond for each study represents its statistical weight in the analysis. The two diamonds below
the table and the black dashed line show the combined measurements across studies. The upper
diamond represents the mean and 95% confidence interval, and the lower diamond represents
the mean and 95% prediction interval. The results of the heterogeneity test are represented by
the estimate of between-studies variance (τ^2), Cochran's Q-test statistic for heterogeneity (χ^2),
degrees of freedom (df), p-value (p), and proportion of heterogeneity-induced variability among
the total variability (I²). The reference names include mouse strain and tumor lines. Abbreviation:
experiment (exp).
904
Figure S4. Forest plots of the analysis of vessel density (no./mm²) measured in adipose tissue
of lean mice across studies. The first, third, and last columns represent the list of references, their
weights in the analysis, and measurements with a 95% confidence interval, respectively. The size
of the blue diamond for each study represents its statistical weight in the analysis. The two
diamonds below the table and the black dashed line show the combined measurements across
studies. The upper diamond represents the mean and 95% confidence interval, and the lower
diamond represents the mean and 95% prediction interval. The results of the heterogeneity test
are represented by the estimate of between-studies variance (τ^2), Cochran's Q-test statistic for
heterogeneity (χ^2), degrees of freedom (df), p-value (p), and proportion of heterogeneity-induced
variability among the total variability (I²). Abbreviation: week (wk).
915
Figure S5. Forest plots of the analysis of vessel density (no./mm²) measured in adipose tissue
of diet-induced obese mice across studies. The first, third, and last columns represent the list of
references, their weights in the analysis, and measurements with a 95% confidence interval,
respectively. The size of the blue diamond for each study represents its statistical weight in the
analysis. The two diamonds below the table and the black dashed line show the combined
measurements across studies. The upper diamond represents the mean and 95% confidence
interval, and the lower diamond represents the mean and 95% prediction interval. The results of
the heterogeneity test are represented by the estimate of between-studies variance (τ^2),
Cochran's Q-test statistic for heterogeneity (χ^2), degrees of freedom (df), p-value (p), and
proportion of heterogeneity-induced variability among the total variability (I²). Abbreviation:
week (wk).
927
Figure S6. Forest plots of the analysis of vessel density (no./mm²) measured in mouse tumors
across studies. The first, third, and last columns represent the list of references, their weights in
the analysis, and measurements with a 95% confidence interval, respectively. The size of the blue
diamond for each study represents its statistical weight in the analysis. The two diamonds below
the table and the black dashed line show the combined measurements across studies. The upper
diamond represents the mean and 95% confidence interval, and the lower diamond represents
the mean and 95% prediction interval. The results of the heterogeneity test are represented by
the estimate of between-studies variance (τ^2), Cochran's Q-test statistic for heterogeneity (χ^2),
degrees of freedom (df), p-value (p), and proportion of heterogeneity-induced variability among

the total variability (I^2). The reference names include mouse strain and tumor lines. Abbreviation:
experiment (exp) and dimethyl sulfoxide (DMSO).

Figure S7. Forest plots of the analysis of capillary basement membrane thickness (nm)
measured in tissues of lean mice and rats across studies. The first, third, and last columns
represent the list of references, their weights in the analysis, and measurements with a 95%
confidence interval, respectively. The size of the blue diamond for each study represents its
statistical weight in the analysis. The two diamonds below the table and the black dashed line
show the combined measurements across studies. The upper diamond represents the mean and
95% confidence interval, and the lower diamond represents the mean and 95% prediction interval.
The results of the heterogeneity test are represented by the estimate of between-studies variance
(τ^2), Cochran's Q-test statistic for heterogeneity (χ^2), degrees of freedom (df), p-value (p), and
proportion of heterogeneity-induced variability among the total variability (I^2). Abbreviation:
month (mo), interventricular septal (IVS), and left ventricular (LV).

Figure S8. Forest plots of the analysis of capillary basement membrane thickness (nm)
measured in tissues of obese mice and rats across studies. The first, third, and last columns
represent the list of references, their weights in the analysis, and measurements with a 95%
confidence interval, respectively. The size of the blue diamond for each study represents its
statistical weight in the analysis. The two diamonds below the table and the black dashed line
show the combined measurements across studies. The upper diamond represents the mean and
95% confidence interval, and the lower diamond represents the mean and 95% prediction interval.
The results of the heterogeneity test are represented by the estimate of between-studies variance
(τ^2), Cochran's Q-test statistic for heterogeneity (χ^2), degrees of freedom (df), p-value (p), and
proportion of heterogeneity-induced variability among the total variability (I^2). Abbreviation:
month (mo).

Figure S9. Capillary basement membrane thickness (nm) in tissues of lean mice and rats. The
capillary basement membrane thickness in retina, muscle, heart, brain, and kidney in lean mice
and rats were compared. Welch's ANOVA followed by Dunnett's T3 test was used to compare a
pair of tissues. The error bar represents the weighted mean ± standard error of each group.

Figure S10. Forest plots of the analysis of capillary basement membrane thickness (nm)
measured in retina of lean mice and rats across studies. The first, third, and last columns
represent the list of references, their weights in the analysis, and measurements with a 95%
confidence interval, respectively. The size of the blue diamond for each study represents its
statistical weight in the analysis. The two diamonds below the table and the black dashed line
show the combined measurements across studies. The upper diamond represents the mean and
95% confidence interval, and the lower diamond represents the mean and 95% prediction interval.
The results of the heterogeneity test are represented by the estimate of between-studies variance
(τ^2), Cochran's Q-test statistic for heterogeneity (χ^2), degrees of freedom (df), p-value (p), and
proportion of heterogeneity-induced variability among the total variability (I^2). Abbreviation:
month (mo).

Figure S11. Forest plots of the analysis of capillary basement membrane thickness (nm) measured in muscle of lean mice and rats across studies. The first, third, and last columns represent the list of references, their weights in the analysis, and measurements with a 95% confidence interval, respectively. The size of the blue diamond for each study represents its statistical weight in the analysis. The two diamonds below the table and the black dashed line show the combined measurements across studies. The upper diamond represents the mean and 95% confidence interval, and the lower diamond represents the mean and 95% prediction interval. The results of the heterogeneity test are represented by the estimate of between-studies variance (τ^2), Cochran's Q-test statistic for heterogeneity (χ^2), degrees of freedom (df), p-value (p), and proportion of heterogeneity-induced variability among the total variability (I^2). Abbreviation: month (mo).

Figure S12. Forest plots of the analysis of capillary basement membrane thickness (nm) measured in heart of lean mice and rats across studies. The first, third, and last columns represent the list of references, their weights in the analysis, and measurements with 95% confidence interval, respectively. The size of the blue diamond for each study represents its statistical weight in the analysis. The two diamonds below the table and the black dashed line show the combined measurements across studies. The upper diamond represents the mean and 95% confidence interval, and the lower diamond represents the mean and 95% prediction interval. The results of the heterogeneity test are represented by the estimate of between-studies variance (τ^2), Cochran's Q-test statistic for heterogeneity (χ^2), degrees of freedom (df), p-value (p), and proportion of heterogeneity-induced variability among the total variability (I^2). Abbreviation: month (mo), interventricular septal (IVS), and left ventricular (LV).

Figure S13. Forest plots of the analysis of capillary basement membrane thickness (nm) measured in brain of lean mice and rats across studies. The first, third, and last columns represent the list of references, their weights in the analysis, and measurements with a 95% confidence interval, respectively. The size of the blue diamond for each study represents its statistical weight in the analysis. The two diamonds below the table and the black dashed line show the combined measurements across studies. The upper diamond represents the mean and 95% confidence interval, and the lower diamond represents the mean and 95% prediction interval. The results of the heterogeneity test are represented by the estimate of between-studies variance (τ^2), Cochran's Q-test statistic for heterogeneity (χ^2), degrees of freedom (df), p-value (p), and proportion of heterogeneity-induced variability among the total variability (I^2). Abbreviation: month (mo).

Figure S14. Forest plots of the analysis of capillary basement membrane thickness (nm) measured in kidney of lean mice and rats across studies. The first, third, and last columns represent the list of references, their weights in the analysis, and measurements with a 95% confidence interval, respectively. The size of the blue diamond for each study represents its statistical weight in the analysis. The two diamonds below the table and the black dashed line show the combined measurements across studies. The upper diamond represents the mean and 95% confidence interval, and the lower diamond represents the mean and 95% prediction interval. The results of the heterogeneity test are represented by the estimate of between-studies variance

1025   (τ^2), Cochran's Q-test statistic for heterogeneity (χ^2), degrees of freedom (df), p-value (p), and
1026   proportion of heterogeneity-induced variability among the total variability (I^2). Abbreviation:
1027   month (mo).

1028

1029   **Figure S15. Interaction kinetics of VEGF-A:NRP1 and VEGF-A:VEGFR2.** NRP1 (A) and VEGFR2 (B)
1030   were immobilized and VEGF-A as analyte was passed over them at different concentrations: 50
1031   nM (yellow), 25 nM (green), 12.5 nM (blue), 6.25 nM (red), and 3.125 nM (black). Note: the thin
1032   black overlapping lines are fitted curves of a 1:1 Langmuir model drawn with TraceDrawer ver.
1033   1.8.1 software**.**

1034

Figure S16. Forest plots of the analysis of binding affinity (pM) of VEGF to VEGFR1 across studies.
The first, third, and last columns represent the list of references, their weights in the analysis, and
measurements with a 95% confidence interval, respectively. The size of the blue diamond for each
study represents its statistical weight in the analysis. The two diamonds below the table and the
black dashed line show the combined measurements across studies. The upper diamond
represents the mean and 95% confidence interval, and the lower diamond represents the mean
and 95% prediction interval. The results of the heterogeneity test are represented by the estimate
of between-studies variance (τ^2), Cochran's Q-test statistic for heterogeneity (χ^2), degrees of
freedom (df), p-value (p), and proportion of heterogeneity-induced variability among the total
variability (I^2). Abbreviation: porcine aortic endothelial (PAE), human umbilical vein endothelial
cell (HUVEC), and surface plasmon resonance (SPR).

1046

Figure S17. Forest plots of the analysis of binding affinity (nM) of VEGF to VEGFR2 across studies.
The first, third, and last columns represent the list of references, their weights in the analysis, and
measurements with a 95% confidence interval, respectively. The size of the blue diamond for each
study represents its statistical weight in the analysis. The two diamonds below the table and the
black dashed line show the combined measurements across studies. The upper diamond
represents the mean and 95% confidence interval, and the lower diamond represents the mean
and 95% prediction interval. The results of the heterogeneity test are represented by the estimate
of between-studies variance (τ^2), Cochran's Q-test statistic for heterogeneity (χ^2), degrees of
freedom (df), p-value (p), and proportion of heterogeneity-induced variability among the total
variability (I^2). Abbreviation: porcine aortic endothelial (PAE) and human umbilical vein
endothelial cell (HUVEC).

1058

Figure S18. Forest plots of the analysis of binding affinity (nM) of VEGF to NRP1 across studies.
The first, third, and last columns represent the list of references, their weights in the analysis, and
measurements with a 95% confidence interval, respectively. The size of the blue diamond for each
study represents its statistical weight in the analysis. The two diamonds below the table and the
black dashed line show the combined measurements across studies. The upper diamond
represents the mean and 95% confidence interval, and the lower diamond represents the mean
and 95% prediction interval. The results of the heterogeneity test are represented by the estimate
of between-studies variance (τ^2), Cochran's Q-test statistic for heterogeneity (χ^2), degrees of
freedom (df), p-value (p), and proportion of heterogeneity-induced variability among the total
variability (I^2). Abbreviation: human umbilical vein endothelial cell (HUVEC).

1069

Figure S19. Forest plots of the analysis of association rates, k_{on} ($\mu M^{-1}s^{-1}$) of VEGF to VEGFR1 across studies. The first, third, and last columns represent the list of references, their weights in the analysis, and measurements with a 95% confidence interval, respectively. The size of the blue diamond for each study represents its statistical weight in the analysis. The two diamonds below the table and the black dashed line show the combined measurements across studies. The upper diamond represents the mean and 95% confidence interval, and the lower diamond represents the mean and 95% prediction interval. The results of the heterogeneity test are represented by the estimate of between-studies variance (τ^2), Cochran's Q-test statistic for heterogeneity (χ^2), degrees of freedom (df), p-value (p), and proportion of heterogeneity-induced variability among the total variability (I^2).

Figure S20. Forest plots of the analysis of association rates, k_{on} ($\mu M^{-1}s^{-1}$) of VEGF to VEGFR2 across studies. The first, third, and last columns represent the list of references, their weights in the analysis, and measurements with a 95% confidence interval, respectively. The size of the blue diamond for each study represents its statistical weight in the analysis. The two diamonds below the table and the black dashed line show the combined measurements across studies. The upper diamond represents the mean and 95% confidence interval, and the lower diamond represents the mean and 95% prediction interval. The results of the heterogeneity test are represented by the estimate of between-studies variance (τ^2), Cochran's Q-test statistic for heterogeneity (χ^2), degrees of freedom (df), p-value (p), and proportion of heterogeneity-induced variability among the total variability (I^2).

Figure S21. Forest plots of the analysis of association rates, k_{on} ($\mu M^{-1}s^{-1}$) of VEGF to NRP1 across studies. The first, third, and last columns represent the list of references, their weights in the analysis, and measurements with a 95% confidence interval, respectively. The size of the blue diamond for each study represents its statistical weight in the analysis. The two diamonds below the table and the black dashed line show the combined measurements across studies. The upper diamond represents the mean and 95% confidence interval, and the lower diamond represents the mean and 95% prediction interval. The results of the heterogeneity test are represented by the estimate of between-studies variance (τ^2), Cochran's Q-test statistic for heterogeneity (χ^2), degrees of freedom (df), p-value (p), and proportion of heterogeneity-induced variability among the total variability (I^2).

Figure S22. Forest plots of the analysis of dissociation rates, k_{off} (s^{-1}) of VEGF to VEGFR1 across studies. The first, third, and last columns represent the list of references, their weights in the analysis, and measurements with a 95% confidence interval, respectively. The size of the blue diamond for each study represents its statistical weight in the analysis. The two diamonds below the table and the black dashed line show the combined measurements across studies. The upper diamond represents the mean and 95% confidence interval, and the lower diamond represents the mean and 95% prediction interval. The results of the heterogeneity test are represented by the estimate of between-studies variance (τ^2), Cochran's Q-test statistic for heterogeneity (χ^2), degrees of freedom (df), p-value (p), and proportion of heterogeneity-induced variability among the total variability (I^2).

1113

Figure S23. Forest plots of the analysis of dissociation rates, k_{off} (s^{-1}) of VEGF to VEGFR2 across
studies. The first, third, and last columns represent the list of references, their weights in the
analysis, and measurements with a 95% confidence interval, respectively. The size of the blue
diamond for each study represents its statistical weight in the analysis. The two diamonds below
the table and the black dashed line show the combined measurements across studies. The upper
diamond represents the mean and 95% confidence interval, and the lower diamond represents
the mean and 95% prediction interval. The results of the heterogeneity test are represented by
the estimate of between-studies variance (τ^2), Cochran's Q-test statistic for heterogeneity (χ^2),
degrees of freedom (df), p-value (p), and proportion of heterogeneity-induced variability among
the total variability (I^2).

1124

Figure S24. Forest plots of the analysis of dissociation rates, k_{off} (s^{-1}) of VEGF to NRP1 across
studies. The first, third, and last columns represent the list of references, their weights in the
analysis, and measurements with a 95% confidence interval, respectively. The size of the blue
diamond for each study represents its statistical weight in the analysis. The two diamonds below
the table and the black dashed line show the combined measurements across studies. The upper
diamond represents the mean and 95% confidence interval, and the lower diamond represents
the mean and 95% prediction interval. The results of the heterogeneity test are represented by
the estimate of between-studies variance (τ^2), Cochran's Q-test statistic for heterogeneity (χ^2),
degrees of freedom (df), p-value (p), and proportion of heterogeneity-induced variability among
the total variability (I^2).

1135

1136 **Figure S25. Binding affinities of VEGF to VEGFR1, VEGFR2, and NRP1 measured by cell-based**
1137 **(radioligand) assay and chip-based (surface plasmon resonance) assay.** The error bar represents
1138 the weighted mean ± standard error of VEGF binding affinities for VEGFR1, VEGFR2, and NRP1. A
1139 one-tailed Welch's t-test (p-value < 0.05) was used to compare a pair of groups.

1140

Figure S26. Forest plots of the analysis of binding affinity (pM) of VEGF to VEGFR1 measured by
radioligand assay across studies. The first, third, and last columns represent the list of references,
their weights in the analysis, and measurements with a 95% confidence interval, respectively. The
size of the blue diamond for each study represents its statistical weight in the analysis. The two
diamonds below the table and the black dashed line show the combined measurements across
studies. The upper diamond represents the mean and 95% confidence interval, and the lower
diamond represents the mean and 95% prediction interval. The results of the heterogeneity test
are represented by the estimate of between-studies variance (τ^2), Cochran's Q-test statistic for
heterogeneity (χ^2), degrees of freedom (df), p-value (p), and proportion of heterogeneity-induced
variability among the total variability (I^2). Abbreviation: porcine aortic endothelial (PAE) and
human umbilical vein endothelial cell (HUVEC).

1152

Figure S27. Forest plots of the analysis of binding affinity (pM) of VEGF to VEGFR1 measured by
surface plasmon resonance across studies. The first, third, and last columns represent the list of
references, their weights in the analysis, and measurements with a 95% confidence interval,
respectively. The size of the blue diamond for each study represents its statistical weight in the

analysis. The two diamonds below the table and the black dashed line show the combined
measurements across studies. The upper diamond represents the mean and 95% confidence
interval, and the lower diamond represents the mean and 95% prediction interval. The results of
the heterogeneity test are represented by the estimate of between-studies variance (τ^2),
Cochran's Q-test statistic for heterogeneity (χ^2), degrees of freedom (df), p-value (p), and
proportion of heterogeneity-induced variability among the total variability (I^2).

1163

Figure S28. Forest plots of the analysis of binding affinity (pM) of VEGF to VEGFR2 measured by
radioligand assay across studies. The first, third, and last columns represent the list of references,
their weights in the analysis, and measurements with a 95% confidence interval, respectively. The
size of the blue diamond for each study represents its statistical weight in the analysis. The two
diamonds below the table and the black dashed line show the combined measurements across
studies. The upper diamond represents the mean and 95% confidence interval, and the lower
diamond represents the mean and 95% prediction interval. The results of the heterogeneity test
are represented by the estimate of between-studies variance (τ^2), Cochran's Q-test statistic for
heterogeneity (χ^2), degrees of freedom (df), p-value (p), and proportion of heterogeneity-induced
variability among the total variability (I^2). Abbreviation: porcine aortic endothelial (PAE) and
human umbilical vein endothelial cell (HUVEC).

1175

Figure S29. Forest plots of the analysis of binding affinity (pM) of VEGF to VEGFR2 measured by
surface plasmon resonance across studies. The first, third, and last columns represent the list of
references, their weights in the analysis, and measurements with a 95% confidence interval,
respectively. The size of the blue diamond for each study represents its statistical weight in the
analysis. The two diamonds below the table and the black dashed line show the combined
measurements across studies. The upper diamond represents the mean and 95% confidence
interval, and the lower diamond represents the mean and 95% prediction interval. The results of
the heterogeneity test are represented by the estimate of between-studies variance (τ^2),
Cochran's Q-test statistic for heterogeneity (χ^2), degrees of freedom (df), p-value (p), and
proportion of heterogeneity-induced variability among the total variability (I^2).

1186

Figure S30. Forest plots of the analysis of binding affinity (nM) of VEGF to NRP1 measured by
radioligand assay across studies. The first, third, and last columns represent the list of references,
their weights in the analysis, and measurements with a 95% confidence interval, respectively. The
size of the blue diamond for each study represents its statistical weight in the analysis. The two
diamonds below the table and the black dashed line show the combined measurements across
studies. The upper diamond represents the mean and 95% confidence interval, and the lower
diamond represents the mean and 95% prediction interval. The results of the heterogeneity test
are represented by the estimate of between-studies variance (τ^2), Cochran's Q-test statistic for
heterogeneity (χ^2), degrees of freedom (df), p-value (p), and proportion of heterogeneity-induced
variability among the total variability (I^2). Inside the parentheses of reference names, the cell lines
used in the assay are included. Abbreviation: human umbilical vein endothelial cell (HUVEC).

1198

Figure S31. Forest plots of the analysis of binding affinity (nM) of VEGF to NRP1 measured by
surface plasmon resonance across studies. The first, third, and last columns represent the list of

references, their weights in the analysis, and measurements with a 95% confidence interval,
respectively. The size of the blue diamond for each study represents its statistical weight in the
analysis. The two diamonds below the table and the black dashed line show the combined
measurements across studies. The upper diamond represents the mean and 95% confidence
interval, and the lower diamond represents the mean and 95% prediction interval. The results of
the heterogeneity test are represented by the estimate of between-studies variance (τ^2),
Cochran's Q-test statistic for heterogeneity (χ^2), degrees of freedom (df), p-value (p), and
proportion of heterogeneity-induced variability among the total variability (I^2).

Author(s) and Year	Weight (%)	Measurements (μm) [95% CI]
Lijnen et al., 2006	7.8	66 [62, 70]
Lijnen et al., 2001 (17-wk diet)	7.3	80 [70, 90]
Lijnen et al., 2001 (32-wk diet)	7.8	86 [83, 89]
Maquoi et al., 2002	7.7	83 [77, 89]
Morange et al., 2000	7.6	82 [75, 89]
Voros et al., 2005 (2-wk diet)	7.8	35 [34, 36]
Voros et al., 2005 (5-wk diet)	7.8	41 [37, 45]
Voros et al., 2005 (15-wk diet)	7.8	53 [52, 54]
Lijnen, Maquoi et al., 2003	7.7	85 [80, 90]
Lijnen et al., 2003	7.4	95 [86, 104]
Maquoi et al., 2003	7.7	76 [72, 81]
Lijnen et al., 2007	7.8	89 [86, 92]
Van Hul et al., 2012	7.7	58 [54, 63]
Random-effects model		71 [61, 82]
Prediction interval (95%)		71 [33, 110]

Adipocyte diameter (μm)

Test for heterogeneity: $\tau^2 = 362.13$; $\chi^2 = 2304.07$, df = 12 , $p < 0.001$; $I^2 = 99\%$

Lean adipose
- Lijnen et al., 2006
- Lijnen, Maquoi et al., 2003
- Van Hul et al., 2012
- Voros et al., 2005 (15-wk diet)
- Voros et al., 2005 (2-wk diet)
- Voros et al., 2005 (5-wk diet)
- Voros et al., 2005 (Littermate of ob/ob mice)

Obese adipose
- Lijnen et al., 2003
- Lijnen et al., 2006
- Lijnen et al., 2007
- Lijnen, Maquoi et al., 2003
- Maquoi et al., 2003
- Van Hul et al., 2012
- Voros et al., 2005 (15-wk diet)
- Voros et al., 2005 (2-wk diet)
- Voros et al., 2005 (5-wk diet)

Tumor
- Fernandez-Rodriguez et al., 2016 (Exp 1- C57BL/6 & B16F1)
- Fernandez-Rodriguez et al., 2016 (Exp 2- C57BL/6 & B16F1)
- Goel et al., 2013 (C57BL/6J & E0771 tumor)
- Goel et al., 2013 (Nude & 4T1 primary tumor)
- Jones et al., 2013 (C57BL/6 & LLC cell)
- Kostourou et al., 2013 (B6;129 & B16F0 cell & DMSO)
- Kostourou et al., 2013 (B6;129 & B16F0 cell)
- Kostourou et al., 2013 (B6;129 & CMT19T cell)
- Tufto & Rofstad, 1999 (Balb/c nu/nu & D-12 cell)
- Tufto & Rofstad, 1999 (Balb/c nu/nu & R-18 cell)
- Tufto & Rofstad, 1999 (Balb/c nu/nu & U-25 cell)

Author(s) and Year	Weight (%)	Measurements (μm²) [95% CI]
Lijnen et al., 2006	17.3	27 [24, 30]
Voros et al., 2005 (2-wk diet)	17.3	52 [48, 56]
Voros et al., 2005 (5-wk diet)	17.0	50 [45, 55]
Voros et al., 2005 (15-wk diet)	16.7	49 [42, 56]
Lijnen, Maquoi et al., 2003	15.9	74 [65, 83]
Van Hul et al., 2012	15.7	59 [49, 69]
Random-effects model		51 [39, 64]
Prediction interval (95%)		51 [20, 83]

Vessel size (μm^2)

Test for heterogeneity: $\tau^2 = 222.05$; $\chi^2 = 172.70$, df = 5 , $p < 0.001$; $I^2 = 97$ %

Author(s) and Year	Weight (%)	Measurements (µm²) [95% CI]
Lijnen et al., 2006	11.5	41 [35, 47]
Voros et al., 2005 (2-wk diet)	11.6	48 [44, 52]
Voros et al., 2005 (5-wk diet)	11.6	41 [37, 45]
Voros et al., 2005 (15-wk diet)	11.5	54 [48, 60]
Lijnen, Maquoi et al., 2003	8.2	140 [103, 177]
Lijnen et al., 2003	11.6	47 [42, 52]
Maquoi et al., 2003	11.5	76 [68, 84]
Van Hul et al., 2012	11.6	49 [44, 54]
Lijnen et al., 2007	10.9	108 [93, 123]
Random-effects model		64 [45, 84]
Prediction interval (95%)		64 [3, 126]

Vessel size (µm²)

Test for heterogeneity: $\tau^2 = 871.34$; $\chi^2 = 158.99$, df = 8 , $p < 0.001$; $I^2 = 99\%$

Author(s) and Year	Weight (%)	Measurements (µm²) [95% CI]
Mesquita et al., 2012 (C3H & fibrosarcoma)	11.2	53 [34, 72]
Mesquita et al., 2012 (Nude & fibrosarcoma)	10.7	130 [101, 159]
Goel et al., 2013 (Nude & 4T1 primary tumor)	11.5	87 [81, 92]
Goel et al., 2013 (C57BL/6J & E0771 tumor)	11.6	89 [87, 91]
Tufto & Rofstad, 1999 (Balb/c nu/nu & D-12 cell)	11.3	135 [119, 151]
Tufto & Rofstad, 1999 (Balb/c nu/nu & R-18 cell)	11.5	93 [85, 102]
Tufto & Rofstad, 1999 (Balb/c nu/nu & U-25 cell)	11.4	113 [102, 124]
Fernandez-Rodriguez et al., 2016 (Exp 1- C57BL/6 & B16F1)	10.2	216 [179, 254]
Fernandez-Rodriguez et al., 2016 (Exp 2- C57BL/6 & B16F1)	10.6	210 [179, 240]
Random-effects model		123 [88, 158]
Prediction interval (95%)		123 [14, 232]

Vessel size (µm²)

Test for heterogeneity: $\tau^2 = 2780.53$; $\chi^2 = 174.02$, df = 8 , $p < 0.001$; $I^2 = 99$ %

Author(s) and Year	Weight (%)	Measurements (no./mm²) [95% CI]
Lijnen et al., 2006	11.2	290 [245, 335]
Voros et al., 2005 (2-wk diet)	11.1	790 [731, 849]
Voros et al., 2005 (5-wk diet)	11.2	410 [371, 449]
Voros et al., 2005 (15-wk diet)	11.2	490 [453, 527]
Lijnen, Maquoi et al., 2003	11.0	200 [133, 267]
Lijnen et al., 2003	11.3	120 [108, 132]
Maquoi et al., 2003	11.2	210 [177, 243]
Van Hul et al., 2012	10.5	400 [292, 508]
Lijnen et al., 2007	11.2	238 [207, 269]
Random-effects model		349 [216, 482]
Prediction interval (95%)		349 [-68, 766]

Vessel density (no./mm²)

Test for heterogeneity: $\tau^2 = 40684.94$; $\chi^2 = 928.59$, df = 8 , p < 0.001 ; $I^2 = 99$ %

Author(s) and Year	Weight (%)	Measurements (no./mm²) [95% CI]
Tufto & Rofstad, 1999 (Balb/c nu/nu & D-12 cell)	9.3	20 [15, 24]
Tufto & Rofstad, 1999 (Balb/c nu/nu & R-18 cell)	9.3	19 [16, 23]
Tufto & Rofstad, 1999 (Balb/c nu/nu & U-25 cell)	9.3	18 [13, 22]
Fernandez-Rodriguez et al., 2016 (Exp 1- C57BL/6 & B16F1)	9.2	71 [52, 90]
Fernandez-Rodriguez et al., 2016 (Exp 2- C57BL/6 & B16F1)	9.3	49 [40, 58]
Kostourou et al., 2013 (B6;129 & B16F0 cell)	9.3	32 [21, 43]
Kostourou et al., 2013 (B6;129 & CMT19T cell)	9.3	30 [24, 35]
Kostourou et al., 2013 (B6;129 & B16F0 cell & DMSO)	9.2	65 [51, 79]
Jones et al., 2013 (C57BL/6 & LLC cell)	8.9	148 [113, 183]
Goel et al., 2013 (Nude & 4T1 primary tumor)	8.4	292 [236, 348]
Goel et al., 2013 (C57BL/6J & E0771 tumor)	8.6	212 [162, 262]
Random-effects model		84 [32, 135]
Prediction interval (95%)		84 [-93, 261]

Vessel density (no./mm²)

Test for heterogeneity: $\tau^2 = 7457.07$; $\chi^2 = 298.33$, df = 10 , p < 0.001 ; $I^2 = 100\%$

Author(s) and Year	Weight (%)	Measurements (nm) [95% CI]
Cuthbertson & Mandel, 1986 (1.5-mo CBA mice, retina, mid-zone)	2.0	50 [41, 59]
Cuthbertson & Mandel, 1986 (4-mo CBA mice, retina, mid-zone)	2.0	59 [46, 72]
Cuthbertson & Mandel, 1986 (8-mo CBA mice, retina, mid-zone)	2.0	75 [65, 85]
Cuthbertson & Mandel, 1986 (12-mo CBA mice, retina, mid-zone)	1.9	108 [91, 125]
Cuthbertson & Mandel, 1986 (20-mo CBA mice, retina, mid-zone)	1.8	154 [128, 180]
Cuthbertson & Mandel, 1986 (1.5-mo Balb/c mice, retina, mid-zone)	2.0	55 [42, 68]
Cuthbertson & Mandel, 1986 (8-mo Balb/c mice, retina, mid-zone)	2.0	77 [64, 90]
Cuthbertson & Mandel, 1986 (20-mo Balb/c mice, retina, mid-zone)	1.4	158 [110, 206]
Cuthbertson & Mandel, 1986 (1.5-mo Balb/c mice, retina, center zone)	2.0	41 [36, 46]
Cuthbertson & Mandel, 1986 (1.5-mo Balb/c mice, retina, periphery zone)	2.0	64 [56, 72]
Creutzfeldt et al., 1970 (7-mo Spiny mice, muscle)	2.0	73 [62, 84]
Carlson et al., 2003 (11-mo FVB/N mice, retina)	2.0	93 [80, 106]
Carlson et al., 2003 (11-mo FVB/N mice, muscle, extensor digitorum)	2.0	77 [68, 86]
Carlson et al., 2003 (11-mo FVB/N mice, muscle, diaphragm)	2.0	55 [51, 58]
Carlson et al., 2003 (11-mo FVB/N mice, heart, IVS)	2.0	63 [60, 66]
Carlson et al., 2003 (11-mo FVB/N mice, heart, LV)	2.0	57 [55, 59]
Williams et al., 2020 (5.5-mo C57BL/6 mice, muscle)	2.0	108 [97, 118]
Velic et al., 2013 (12-mo FVB mice, heart)	2.0	48 [43, 54]
Lash et al., 1989 (2.8-mo Zucker rats, muscle)	2.0	62 [59, 64]
Lash et al., 1989 (4.5-mo Zucker rats, muscle)	2.0	56 [54, 58]
Dosso et al., 1990 (17-mo Zucker rats, retina)	2.0	94 [88, 99]
Danis & Yang, 1993 (6.5-mo Zucker rats, retina)	2.0	89 [85, 93]
Cherian et al., 2009 (6-mo Sprague-Dawley rats, retina)	2.0	51 [47, 55]
Saito et al., 2003 (5.5-mo LETO rats, heart)	2.0	90 [87, 93]
Saito et al., 2003 (15.5-mo LETO rats, heart)	2.0	87 [84, 90]
Begieneman et al., 2009 (Rats, heart)	2.0	69 [60, 77]
Das et al., 1990 (10.5-mo Wistar rats, retina, inner nuclear)	1.6	167 [128, 206]
Das et al., 1990 (10.5-mo Wistar rats, retina, nerve fiber)	1.6	206 [169, 244]
Robison et al., 1983 (7-mo Sprague-Dawley rats, retina)	1.9	97 [81, 113]
Robison et al., 1983 (11-mo Sprague-Dawley rats, retina)	2.0	94 [80, 108]
Roy et al., 2003 (7-mo Sprague-Dawley rats, retina)	1.9	85 [68, 102]
Evans et al., 2000 (6-mo Sprague-Dawley rats, retina)	2.0	66 [61, 72]
Zheng et al., 2007 (8-mo Sprague-Dawley rats, retina)	2.0	76 [68, 83]
Dosso et al., 2004 (7-mo Wistar rats, retina)	2.0	112 [101, 122]
Frank et al., 1983 (6-mo Wistar rats, retina)	1.9	82 [66, 97]
Frank et al., 1983 (9-mo Wistar rats, retina)	1.9	95 [79, 110]
Frank et al., 1983 (12-mo Wistar rats, retina)	1.6	121 [83, 159]
Frank et al., 1983 (19.5-mo Wistar rats, retina)	1.8	160 [137, 184]
Chakrabarti & Sima, 1989 (7-mo BB rats, retina, superficial capillary)	2.0	135 [120, 150]
Chakrabarti & Sima, 1989 (7-mo BB rats, retina, deep capillary)	2.0	95 [89, 102]
McCaleb et al., 1991 (10.5-mo Wistar rats, retina)	2.0	84 [82, 87]
Robison et al., 1986 (7-mo Sprague-Dawley rats, retina)	1.9	123 [104, 142]
Clements et al., 1998 (5.5-mo db/m mice, retina)	2.0	79 [76, 83]
Sima et al., 1988 (4-mo Lewis rats, retina, superficial capillary)	2.0	146 [140, 152]
Sima et al., 1988 (4-mo Lewis rats, retina, deep capillary)	2.0	97 [91, 104]
Kern & Engerman, 1994 (20-mo Sprague-Dawley rats, retina)	2.0	159 [146, 172]
Chakrabarti et al., 1991 (6-mo BB rats, retina, superficial capillary)	2.0	162 [155, 169]
Chakrabarti et al., 1991 (6-mo BB rats, retina, deep capillary)	2.0	143 [136, 149]
Chakrabarti et al., 1991 (6-mo BB rats, muscle)	2.0	125 [119, 131]
Roy et al., 2011 (6-mo Sprague-Dawley rats, retina)	2.0	52 [48, 55]
Li et al., 2018 (8-mo Sprague-Dawley rats, retina)	2.0	95 [88, 101]
Random-effects model		94 [84, 104]
Prediction interval (95%)		94 [23, 164]

Capillary basement membrane thickness (nm)

Test for heterogeneity: $\tau^2 = 1260.74$; $\chi^2 = 4239.49$, df = 50 , $p < 0.001$; $I^2 = 99$ %

Author(s) and Year	Weight (%)	Measurements (nm) [95% CI]
Cuthbertson & Mandel, 1986 (1.5-mo CBA mice, mid-zone)	2.7	50 [41, 59]
Cuthbertson & Mandel, 1986 (4-mo CBA mice, mid-zone)	2.7	59 [46, 72]
Cuthbertson & Mandel, 1986 (8-mo CBA mice, mid-zone)	2.7	75 [65, 85]
Cuthbertson & Mandel, 1986 (12-mo CBA mice, mid-zone)	2.6	108 [91, 125]
Cuthbertson & Mandel, 1986 (20-mo CBA mice, mid-zone)	2.4	154 [128, 180]
Cuthbertson & Mandel, 1986 (1.5-mo Balb/c mice, mid-zone)	2.7	55 [42, 68]
Cuthbertson & Mandel, 1986 (8-mo Balb/c mice, mid-zone)	2.7	77 [64, 90]
Cuthbertson & Mandel, 1986 (20-mo Balb/c mice, mid-zone)	1.9	158 [110, 206]
Cuthbertson & Mandel, 1986 (1.5-mo Balb/c mice, center zone)	2.7	41 [36, 46]
Cuthbertson & Mandel, 1986 (1.5-mo Balb/c mice, periphery zone)	2.7	64 [56, 72]
Carlson et al., 2003 (11-mo FVB/N mice)	2.7	93 [80, 106]
Dosso et al., 1990 (17-mo Zucker rats)	2.7	94 [88, 99]
Danis & Yang, 1993 (6.5-mo Zucker rats)	2.8	89 [85, 93]
Cherian et al., 2009 (6-mo Sprague-Dawley rats)	2.8	51 [47, 55]
Das et al., 1990 (10.5-mo Wistar rats, inner nuclear)	2.2	167 [128, 206]
Das et al., 1990 (10.5-mo Wistar rats, nerve fiber)	2.2	206 [169, 244]
Robison et al., 1983 (7-mo Sprague-Dawley rats)	2.6	97 [81, 113]
Robison et al., 1983 (11-mo Sprague-Dawley rats)	2.7	94 [80, 108]
Roy et al., 2003 (7-mo Sprague-Dawley rats)	2.6	85 [68, 102]
Evans et al., 2000 (6-mo Sprague-Dawley rats)	2.7	66 [61, 72]
Zheng et al., 2007 (8-mo Sprague-Dawley rats)	2.7	76 [68, 83]
Dosso et al., 2004 (7-mo Wistar rats)	2.7	112 [101, 122]
Frank et al., 1983 (6-mo Wistar rats)	2.6	82 [66, 97]
Frank et al., 1983 (9-mo Wistar rats)	2.6	95 [79, 110]
Frank et al., 1983 (12-mo Wistar rats)	2.2	121 [83, 159]
Frank et al., 1983 (19.5-mo Wistar rats)	2.5	160 [137, 184]
Chakrabarti & Sima, 1989 (7-mo BB rats, superficial capillary)	2.7	135 [120, 150]
Chakrabarti & Sima, 1989 (7-mo BB rats, deep capillary)	2.7	95 [89, 102]
McCaleb et al., 1991 (10.5-mo Wistar rats)	2.8	84 [82, 87]
Robison et al., 1986 (7-mo Sprague-Dawley rats)	2.6	123 [104, 142]
Clements et al., 1998 (5.5-mo db/m mice)	2.8	79 [76, 83]
Sima et al., 1988 (4-mo Lewis rats, superficial capillary)	2.7	146 [140, 152]
Sima et al., 1988 (4-mo Lewis rats, deep capillary)	2.7	97 [91, 104]
Kern & Engerman, 1994 (20-mo Sprague-Dawley rats)	2.7	159 [146, 172]
Chakrabarti et al., 1991 (6-mo BB rats, superficial capillary)	2.7	162 [155, 169]
Chakrabarti et al., 1991 (6-mo BB rats, deep capillary)	2.7	143 [136, 149]
Roy et al., 2011 (6-mo Sprague-Dawley rats)	2.8	52 [48, 55]
Li et al., 2018 (8-mo Sprague-Dawley rats)	2.7	95 [88, 101]
Random-effects model		101 [89, 113]
Prediction interval (95%)		101 [26, 175]

Capillary basement membrane thickness (nm)

0   75   150   225   300

Test for heterogeneity: $\tau^2 = 1408.02$; $\chi^2 = 2656.05$, df = 37 , $p < 0.001$; $I^2 = 99\%$

Author(s) and Year		Weight (%)	Measurements (nm) [95% CI]
Creutzfeldt et al., 1970 (7-mo Spiny mice)		13.9	73 [62, 84]
Carlson et al., 2003 (11-mo FVB/N mice, extensor digitorum)		14.2	77 [68, 86]
Carlson et al., 2003 (11-mo FVB/N mice, diaphragm)		14.5	55 [51, 58]
Williams et al., 2020 (5.5-mo C57BL/6 mice)		14.0	108 [97, 118]
Lash et al., 1989 (2.8-mo Zucker rats)		14.5	62 [59, 64]
Lash et al., 1989 (4.5-mo Zucker rats)		14.5	56 [54, 58]
Chakrabarti et al., 1991 (6-mo BB rats)		14.4	125 [119, 131]
Random-effects model			79 [59, 99]
Prediction interval (95%)			79 [23, 136]

Capillary basement membrane thickness (nm)

Test for heterogeneity: $\tau^2 = 724.88$; $\chi^2 = 525.10$, df = 6 , $p < 0.001$; $I^2 = 99$ %

Author(s) and Year	Weight (%)	Measurements (nm) [95% CI]
Carlson et al., 2003 (11-mo FVB/N mice, IVS)	16.8	63 [60, 66]
Carlson et al., 2003 (11-mo FVB/N mice, LV)	16.9	57 [55, 59]
Velic et al., 2013 (12-mo FVB mice)	16.6	48 [43, 54]
Saito et al., 2003 (5.5-mo LETO rats)	16.8	90 [87, 93]
Saito et al., 2003 (15.5-mo LETO rats)	16.8	87 [84, 90]
Begieneman et al., 2009 (Rats)	16.0	69 [60, 77]
Random-effects model		69 [56, 82]
Prediction interval (95%)		69 [34, 104]

Capillary basement membrane thickness (nm)

Test for heterogeneity: $\tau^2 = 270.62$; $\chi^2 = 440.02$, df = 5 , $p < 0.001$; $I^2 = 99\%$

Author(s) and Year	Weight (%)	Measurements (nm) [95% CI]
Ceafalan et al., 2019 (6-mo C57BL/6J mice)	8.1	57 [54, 59]
Ceafalan et al., 2019 (24-mo C57BL/6J mice)	8.0	108 [103, 112]
Hawkes et al., 2013 (2-mo C57BL/6 mice, striatum)	7.6	67 [59, 76]
Hawkes et al., 2013 (2-mo C57BL/6 mice, cortex)	7.4	66 [55, 77]
Hawkes et al., 2013 (2-mo C57BL/6 mice, hippocampus)	7.9	63 [57, 68]
Hawkes et al., 2013 (2-mo C57BL/6 mice, thalamus)	7.8	70 [63, 77]
Hawkes et al., 2013 (7-mo C57BL/6 mice, striatum)	7.1	75 [62, 88]
Hawkes et al., 2013 (7-mo C57BL/6 mice, cortex)	7.6	57 [49, 66]
Hawkes et al., 2013 (7-mo C57BL/6 mice, hippocampus)	7.5	64 [54, 73]
Hawkes et al., 2013 (7-mo C57BL/6 mice, thalamus)	7.5	59 [48, 69]
Hawkes et al., 2013 (23-mo C57BL/6 mice, striatum)	7.9	74 [69, 80]
Hawkes et al., 2013 (23-mo C57BL/6 mice, cortex)	5.4	97 [73, 122]
Hawkes et al., 2013 (23-mo C57BL/6 mice, hippocampus)	5.5	110 [86, 133]
Hawkes et al., 2013 (23-mo C57BL/6 mice, thalamus)	4.8	108 [80, 137]
Random-effects model		75 [65, 84]
Prediction interval (95%)		75 [40, 110]

Capillary basement membrane thickness (nm)

Test for heterogeneity: $\tau^2 = 297.55$; $\chi^2 = 403.76$, df = 13 , $p < 0.001$; $I^2 = 96$ %

Author(s) and Year	Weight (%)	Measurements (nm) [95% CI]
Rodrigues et al., 1983 (7-mo SJL/J mice)	6.4	91 [89, 93]
Carlson et al., 2003 (11-mo FVB/N mice)	6.3	178 [162, 194]
Chang et al., 2012 (7.5-mo FVB/NJ mice)	5.5	224 [170, 278]
Cherian et al., 2009 (6-mo Sprague-Dawley rats)	6.3	81 [73, 89]
Gambaro et al., 1992 (9.5-mo Sprague-Dawley rats)	6.4	236 [234, 238]
Yagihashi, 1978 (1-mo Wistar rats)	6.3	119 [112, 126]
Yagihashi, 1978 (2-mo Wistar rats)	6.4	129 [125, 134]
Yagihashi, 1978 (3-mo Wistar rats)	6.4	135 [132, 139]
Yagihashi, 1978 (4-mo Wistar rats)	6.4	147 [146, 149]
Yagihashi, 1978 (5-mo Wistar rats)	6.4	154 [151, 158]
Yagihashi, 1978 (6-mo Wistar rats)	6.4	160 [157, 164]
Yagihashi, 1978 (7-mo Wistar rats)	6.3	185 [178, 191]
Yagihashi, 1978 (8-mo Wistar rats)	6.3	205 [198, 212]
Yagihashi, 1978 (9-mo Wistar rats)	6.3	238 [227, 249]
Fox et al., 1977 (17-mo Wistar rats)	6.2	305 [285, 325]
Chakrabarti et al., 1991 (6-mo BB rats)	5.9	326 [287, 365]
Random-effects model		181 [147, 215]
Prediction interval (95%)		181 [42, 320]

Capillary basement membrane thickness (nm)

Test for heterogeneity: $\tau^2 = 4729.17$; $\chi^2 = 12747.86$, df = 15 , $p < 0.001$; $I^2 = 100$ %

Author(s) and Year	Weight (%)	Measurements (pM) [95% CI]
Waltenberger et al., 1994 (PAE cell)	10.4	16 [13, 19]
Waltenberger et al., 1994 (HUVEC)	10.4	9 [7, 11]
Mamer et al., 2020	10.4	1 [0, 2]
von Tiedemann & Bilitewski, 2002 (SPR)	10.4	8 [6, 9]
von Tiedemann & Bilitewski, 2002 (Radioligand)	10.0	74 [59, 89]
Papadopoulos et al., 2012	10.4	9 [7, 11]
Jin et al., 2007	10.4	7 [5, 9]
Breier et al., 1995	9.5	114 [92, 136]
Ito & Claesson-Welsh, 1999	10.2	54 [43, 65]
Rouet et al., 2005	8.1	59 [21, 98]
Random-effects model		34 [10, 57]
Prediction interval (95%)		34 [-42, 109]

Binding affinity, K_d (pM)

Test for heterogeneity: $\tau^2 = 1343.99$; $\chi^2 = 514.99$, df = 9 , p < 0.001 ; $I^2 = 100\%$

Author(s) and Year	Weight (%)	Measurements (nM) [95% CI]
Huang et al., 1998 (VEGF-A165)	5.2	0.23 [-0.01, 0.47]
Huang et al., 1998 (VEGF-A164)	5.6	0.24 [0.04, 0.44]
Terman et al., 1992	6.6	0.07 [0.06, 0.09]
Waltenberger et al., 1994 (PAE cell)	5.9	0.76 [0.61, 0.91]
Waltenberger et al., 1994 (HUVEC)	5.9	0.77 [0.62, 0.92]
Whitaker et al., 2001 (COS-1 cell)	5.2	0.34 [0.10, 0.57]
Whitaker et al., 2001 (Balb/c cell)	6.2	0.29 [0.18, 0.40]
Cunningham et al., 1999 (full-length predimer)	6.6	0.04 [0.03, 0.05]
Cunningham et al., 1999 (full-length monomer)	6.6	0.05 [0.04, 0.06]
Cunningham et al., 1999 (short-length predimer)	6.6	0.01 [0.01, 0.02]
Mamer et al., 2020	6.6	0.01 [0.01, 0.01]
In-house data, 2023	3.1	0.52 [0.03, 1.01]
Rouet et al., 2005	4.4	0.29 [-0.03, 0.61]
Lu et al., 2023	6.0	0.12 [-0.03, 0.26]
Papadopoulos et al., 2012	6.6	0.09 [0.08, 0.10]
Soker et al., 1996	6.6	0.01 [0.01, 0.01]
Tessler et al., 1994	6.6	0.02 [0.02, 0.02]
Random-effects model		0.21 [0.09, 0.33]
Prediction interval (95%)		0.21 [-0.27, 0.68]

Binding affinity, K_d (nM)

Test for heterogeneity: $\tau^2 = 0.06$; $\chi^2 = 584.73$, df = 16 , $p < 0.001$; $I^2 = 100$ %

Author(s) and Year	Weight (%)	Measurements (pM) [95% CI]
Huang et al., 1998 (VEGF-A165)	1.5	230 [-5, 465]
Huang et al., 1998 (VEGF-A164)	2.1	240 [44, 436]
Cunningham et al., 1999 (full-length predimer)	18.5	37 [27, 47]
Cunningham et al., 1999 (full-length monomer)	18.3	52 [40, 63]
Cunningham et al., 1999 (short-length predimer)	18.8	14 [13, 16]
Mamer et al., 2020	18.8	10 [9, 11]
In-house data, 2023	0.4	520 [30, 1010]
Lu et al., 2023	3.5	115 [-29, 259]
Papadopoulos et al., 2012	18.1	89 [75, 102]
Random-effects model		51 [21, 81]
Prediction interval (95%)		51 [-24, 127]

Binding affinity, K_d (pM)

Test for heterogeneity: $\tau^2 = 1238.13$; $\chi^2 = 241.85$, df = 8 , p < 0.001 ; $I^2 = 100$ %

Author(s) and Year	Weight (%)	Measurements (nM) [95% CI]
Whitaker et al., 2001 (COS-1 cell)	0.8	2.09 [0.49, 3.69]
Whitaker et al., 2001 (Balb/c cell)	11.1	0.42 [0.17, 0.66]
Soker et al., 1996 (HUVEC)	15.8	0.20 [0.16, 0.24]
Soker et al., 1996 (MDA-MB-231)	15.6	0.28 [0.23, 0.33]
Soker et al., 1998	15.5	0.32 [0.26, 0.38]
Gu et al., 2002	1.1	0.93 [-0.46, 2.32]
Rouet et al., 2005	10.6	0.25 [-0.02, 0.51]
Gluzman-Poltorak et al., 2000	15.8	0.18 [0.14, 0.22]
von Wronski et al., 2006	13.8	0.75 [0.60, 0.90]
Random-effects model		0.36 [0.21, 0.50]
Prediction interval (95%)		0.36 [-0.04, 0.76]

Binding affinity, K_d (nM)

Test for heterogeneity: $\tau^2 = 0.04$; $\chi^2 = 77.41$, df = 8 , p < 0.001 ; $I^2 = 96\ \%$

Author(s) and Year	Weight (%)	Measurements (nM) [95% CI]
In-house data, 2023	33.9	6.4 [4.3, 8.5]
Herve et al., 2008	28.4	1.7 [-1.9, 5.3]
Lu et al., 2023	37.7	0.1 [0.0, 0.3]
Random-effects model		2.7 [-1.2, 6.5]
Prediction interval (95%)		2.7 [-4.7, 10.0]

Binding affinity, K_d (nM)

Test for heterogeneity: $\tau^2 = 10.21$; $\chi^2 = 34.33$, df = 2 , p < 0.001 ; $I^2 = 92$ %