Title: The β integrin modulates serotonin sensitivity via NPxY motifs to regulate egg

laying and mechanosensation behaviors in *C. elegans*.

Josh Haram Bumm*, Daniel Dogeon Lee*, Zhongqiang Qiu, Angela Haeun Choi,

Meghana Kolluri, Michael J. Olsen, Lianzijun Wang, and Myeongwoo Lee[#]

Department of Biology, Baylor University, Waco, Texas 76798, USA.

Keywords: phosphomimetic, serotonin transporter, fluoxetine, reserpine, talin, MEC-4.

Running title: Role of integrin NPxY motifs in egg laying and mechanosensation.

* These authors contributed equally to the work.

#To whom all correspondence should be addressed,

Myeongwoo Lee, Ph.D.

Department of Biology

Baylor University, Waco, TX 76798

Email) Myeongwoo_Lee@baylor.edu

ABSTRACT

Background: Integrin is an αβ heterodimeric receptor to the extracellular matrix; its

binding to the matrix recruits focal adhesions to two NPxY motifs, the tyrosine

phosphorylation sites in the cytoplasmic domain. Studies found that replacing tyrosines

(Y) with phenylalanines (F) in the motif of β1 integrin displayed little developmental or

behavioral defects. However, the tyrosine-to-alanine (A) caused embryonic lethality.

Results: Here we report novel functions of the NPxY motifs in *C. elegans pat-3* β

integrin. The membrane-proximal non-phosphorylation *pat-3(Y792F)* mutation caused

hypersensitive egg laying in serotonin, which is more prominent than the membrane-

distal NPxY^{804}. The double non-phosphorylatable *pat-3(YYFF)* mutant exhibited

serotonin hypersensitivity and defective egg retention. The phosphomimetic NPxY, *pat-*

3(Y804E), mutant displayed a reduced egg laying in response to serotonin and

fluoxetine, suggesting that the NPxY phosphorylation is associated with vulval

contraction and serotonin sensitivity. Additionally, *pat-3(Y792A)*, *pat-3(Y792F)*, *pat-*

3(Y804E), and pat-3(YYFF) mutants exhibited mechanosensation defects,

demonstrating that NPxY phosphorylation regulates sensory neuron activity. Further

revealed that exogenous serotonin reduced mechanosensation, while blocking

serotonin secretion rescued the mechanosensation of *pat-3* NPxY mutants, suggesting

that integrin NPxY modulates serotonin levels in *C. elegans*.

Conclusion: Our results underscore the functional importance of *pat-3* NPxY motifs in

muscle and neurons, potentially linking integrin NPxY motifs to neurotransmitter

response and mechanosensory functions.

INTRODUCTION

Integrin is an αβ heterodimeric cell surface receptor to the extracellular matrix (ECM),

controlling cell behaviors through bidirectional signaling. Inside-out signaling activates

the integrin extracellular domain to facilitate ECM binding, while outside-in recruits

signaling molecules and links cytoskeletons to the cell-ECM contacts, known as focal

adhesions (FAs) [1]. In this context, the cytoplasmic domain (CTD) of β integrin interacts

with linker proteins such as talins and kindlins to transduce mechanical information from

the ECM to the cell [2]. Two conserved NPxY (Asn-Pro-x-Tyr) motifs, potential tyrosine

phosphorylation sites, exist in the CTD of all metazoan β integrins [3] (Figure 1). The

membrane-proximal NPxY binds to the FERM domain of talin, while the membrane

distal motif interacts with kindlin [4]. Sakai *et al.* reported overexpression of *src* kinase in a

mouse fibroblast phosphorylated the NPxY motifs of β1A integrin [5]. In mice, knock-in β1

integrin YYFF (Tyrosine-to-Phenylalanine) failed to exhibit discernible phenotypes [6],

although integrin NPxY mutants showed resistance to tumorigenesis [7] and defects in

blood clotting [8]. Conversely, disruptive (tyrosine removal) β1 YYAA, Y783A, or Y795A

(Tyrosine-to-Alanine) mutations caused severe phenotypes such as embryonic lethality

[6]. These findings suggested that the Y-to-A transition generally brings severe

phenotypes to the animal, while the Y-to-F transition produces no overt developmental

phenotypes.

In the nematode *C. elegans*, PAT-3/β integrin is ubiquitously localized in tissues,

including body wall muscles, gonads, neurons, vulva, and the excretory canal [9]. PAT-3

integrin also contains two NPxY motifs [10]. Previous analyses by Itoh *et al.* suggested

that SRC family tyrosine kinases (SFK) could phosphorylate the *pat-3* CTD. For

instance, *C. elegans src-1* mutant exhibited gonad phenotypes similar to those caused

by the integrin HA (hemagglutin)-βtail dominant negative allele [10,11]. Our recent study

demonstrated that mutations in the membrane distal Y to A, E (glutamic acid), or F

mutations resulted in gonad migration and mild motility impairments [12]. These findings

suggest that the membrane-distal NPxY is dispensable for development and motility.

However, given the widespread distribution of integrins in muscle, neurons, and gonads,

we hypothesized that the NPxY motifs in PAT-3 CTD are crucial in undescribed tissue-
specific functions.
3
In *C. elegans* hermaphrodites, vulva development begins at the L1 larval stage and
culminates at the late larval stages. In the early adult stages, fertilized eggs accumulate
in the uterus. Within eight hours of post-fertilization, the fertilized eggs are expelled from
the uterus through vulva contractions, known as 'egg laying,' which is regulated by a
pair of serotonergic hermaphrodite-specific motor neurons (HSNs) connected to vulva
muscles. During egg laying, the HSN releases a neurotransmitter, serotonin, onto the
vulva muscle, inducing its contractions. When the serotonin production or its receptor is
defective, the vulva muscles fail to contract, accumulating unlaid eggs in the uterus and
causing the Egl (egg-laying defective) phenotype. Treating worms with exogenous
serotonin or selective serotonin reuptake inhibitors (SSRIs) stimulates vulva muscle
contraction and increases egg laying [13]. Mutations in the genes associated with muscle
attachment [14], such as βPAT-3 RNAi [13,15], perlecan/UNC-52 [16,17], and syndecan/SDN-1
[18], have been reported to cause defective egg laying, suggesting a link between the Egl
phenotype and impaired cell-ECM interactions.
18
In *C. elegans*, gentle touch on the body induces backward or forward movement. The
touch stimulus is detected by six touch receptor neurons (TRNs): ALMR, ALML, AVM,
PLMR, PLML, and PVM [19]. These TRNs express the MEC-4 DEG/ENaC channel
complex on their cell surfaces, transmitting touch stimuli to the neurons. Extensive
studies have identified several *mec* (mechanosensation abnormal) genes essential for
processing touch-induced behaviors [20]. Integrins and focal adhesion proteins transmit
mechanical forces and trigger cellular signaling responses [21]. Focal adhesion proteins
are expressed in mechanosensory cells, including TRNs [22]. Studies on focal adhesion
genes—such as *unc-97/PINCH* [23] mutant and removal of *pat-2*/α integrin [24], *pat-3*/β
integrin [10], *pat-6*/actopaxin [25], and *unc-112*/kindlin [26]—in ALM and PLM touch neurons
of genetic mosaic animals demonstrated the involvement of these genes in
mechanosensation [22].
31

In this study, our results revealed that the phosphomimetic *pat-3 (kq8042, Y804E)*

mutation exhibited defective egg-laying in response to serotonin or fluoxetine treatment.

Conversely, the membrane-proximal non-phosphorylation *pat-3 (kq7923, Y792F)*

mutant was hypersensitive to serotonin, while the membrane-distal NPxY mutations

showed varied serotonin hypersensitivity. In contrast, the double non-phosphorylation

pat-3(kq24, YYFF) mutant displayed a wild-type response to serotonin hypersensitivity

and increased egg retention in the uterus. Further investigation into the role of *pat-3*

NPxY motifs revealed that *pat-3(kq7921, Y792A)*, *pat-3(kq7923, Y792F)*, *pat-3(kq24,*

YYFF), and *pat-3(kq8042, Y804E)* mutants exhibited touch insensitivity in the anterior or

posterior regions of the body. Subsequent analysis revealed that exogenous serotonin

reduced mechanosensation, while inhibiting neurotransmitter secretion by reserpine, a

vesicular monoamine transporter (VMAT) inhibitor [27], rescued the mechanosensation of

pat-3 NPxY mutants. These findings suggest that phosphorylation of the NPxY motif is

critical for modulating serotonin response in both egg-laying machinery and

mechanosensation.

RESULTS

Mutations in the βpat-3 NPxY motifs.

To characterize the phosphorylation of the NPxY motif in the β*pat-3* integrin CTD, we

introduced mutations into the membrane-distal NPxY in the β*pat-3* CTD (Figure 1).

Previous analyses revealed that CRISPR edits in the membrane-distal NPxY of β*pat-3*

led to motility and gonad migration defects [12]. Additionally, we generated *pat-3(kq24,*

YYFF), a non-phosphorylatable mutation. Table 1 summarizes the behavioral

phenotypes of *pat-3* NPxY mutants described in this manuscript.

Most of the *pat-3* NPxY mutants exhibited regular muscle filament organization. Some

mutants were crossed with AH3437 *GFP::tln-1*, a GFP inserted in the N-terminus of

TLN-1 using the CRISPR-Cas9 system [28], to visualize the structure of dense bodies and

M lines (Figure 2, panels A to F). TLN-1 is expressed in body wall muscle and gonad

epithelial cells, displaying typical dense bodies and M-line localization (arrows in Figure

2). Except for *pat-3(kq7921, Y792A)* in Figure 2E, all other mutants displayed intact

dense bodies along the body wall, which is similar to that of AH3437 GFP::TLN-1

(Figure 2A), suggesting that the mutations were dispensable for organizing cell

attachment structures to the ECM. In addition to GFP::TLN-1, the NPxY mutants were

stained with rhodamine-conjugated phalloidin (Figure S1). Other than the deformity of

muscle filaments caused by the staining procedures, we failed to detect an abnormal

pattern of actin filaments.

The membrane-proximal and distal NPxY motifs affect egg-laying behavior.

Although the non-phosphorylatable NPxY lacks distinct phenotypes in mammalian β

integrins [6], the importance and conservation of the β integrin NPxY led us to investigate

the undescribed role of NPxY in integrin functions. Therefore, we sought to discover

functional phenotypes associated with the NPxY mutations. When the hermaphrodite is

submerged in an aqueous solution, it pauses egg laying because worms recognize the

environment as a lack of food [29]. Adding exogenous serotonin would stimulate vulva

contraction, which results in egg laying [13]. Before egg-laying assays, the fecundity of

viable NPxY mutants, N2, and other control mutants was assessed to determine the

optimal stages for measuring egg laying (Figure S2). All worm strains showed the peak

number of eggs on Day 2 (see Experimental Procedures). Therefore, the egg-laying

behavior was assessed with Day 2 hermaphrodites.

To monitor the egg laying of *pat-3* NPxY mutants, we incubated Day 2 adult

hermaphrodites in 3.0 mg/ml (17 mM) serotonin dissolved in M9 buffer. After one hour

of incubation, *pat-3(kq8042, Y804E)*, the phosphomimetic mutant, laid significantly

fewer eggs in the serotonin solution than N2 ($p < 0.0001$). The *pat-3(kq8043, Y804F),*

the distal non-phosphorylatable, and *pat-3(kq8041, Y804A)*, the distal tyrosine removal,

mutants showed comparable egg laying to N2 ($p < 0.8$) (Figure 3, Supplementary Table

3).

Active egg laying is stimulated by a pair of serotonergic HSN motor neurons, which

secrete serotonin (5-hydroxytryptamine, 5-HT) to induce vulva muscle contraction.

Secreted serotonin is then removed by the serotonin transporter (SERT), which is

expressed on the vulval tissue, as it enters inactive egg laying [30]. Previous studies

demonstrated that selective serotonin reuptake inhibitors (SSRIs) such as fluoxetine

block the function of SERT. It is suggested that fluoxetine binds to MOD-5/CeSERT and

stimulates *C. elegans* egg laying by blocking serotonin removal in the vulval tissue [31].

To test whether the Egl of *pat-3* NPxY mutants was due to abnormal 5-HT secretion,

wild-type N2 and the mutants were treated with 0.5 mg/ml (1.6 mM) of fluoxetine [32]. In

N2, fluoxetine treatment would yield comparable egg-laying results to serotonin

treatment (Figure 3). However, the egg laying of *pat-3(Y804E)* in fluoxetine was

defective ($p < 0.0001$), suggesting that the reduced egg laying of the mutant was due to

defective vulva contractions (Figure 3, Supplementary Table 3). The membrane-

proximal *pat-3(kq7923, Y792F)* was also generated to compare with the membrane-

distal NPxY mutants. The *pat-3(Y792F)* worms showed a significantly increased egg-

laying response to serotonin compared to N2 (mean = 12.8 eggs/hour, *p < 0.0001,*

Figure 3).

The membrane-proximal NPxY mutant showed hypersensitivity to serotonin and

fluoxetine.

Our results suggested that the proximal non-phosphorylation, *pat-3(Y792F),* increased

the response to exogenous serotonin, resulting in hyperactive egg laying (Figure 3). To

further analyze the increased egg laying in *pat-3(Y792F),* we diluted serotonin to a

concentration that resulted in no N2 egg laying. In Figure 4, N2 animals incubated in 0.1

mg/ml (565 µM) serotonin could not stimulate egg laying. In contrast, *pat-3(Y792F)*

showed a significant egg-laying increase at the diluted serotonin concentration (565 µM).

At the same time, the number of eggs laid was comparable to N2 in 3 mg/ml serotonin

(Figure 3), suggesting that the proximal non-phosphorylation would cause the

hypersensitive egg laying in serotonin and that the proximal NPxY regulates the

serotonin sensitivity. *pat-3(Y792F)* also displayed egg laying in 0.1 mg/ml fluoxetine, a

concentration that failed to stimulate egg laying in N2 (Figure 4), suggesting this mutant

is hypersensitive to fluoxetine treatment, although the response was significantly lower

than that of serotonin at the diluted concentration.

The membrane-proximal NPxY motif is known as a binding site for talin, a molecular

bridge connecting the actin cytoskeleton to integrin [33]. Our previous study created a

tryptophan (W) to alanine (A) mutation at position 387 in the F3 FERM domain of TLN-

1/talin, denoted as *tln-1(kq387, W387A).* Structural analysis of the talin-integrin

interaction demonstrated that W359 of mammalian talin, homologous to W387 in *tln-1,*

binds to the proximal NPxY of β3 integrin tail. However, *tln-1(kq387)* showed mild egg-

laying defects or delays in motility [34]. The *tln-1(kq387)* was treated with 0.1 mg/ml

serotonin and fluoxetine for egg-laying assays (Figure 4, supplementary table 4). Each

treatment showed increased egg laying during the incubation, suggesting that the focal

adhesion protein TLN-1 is also linked to serotonin hypersensitivity.

Egg retention of NPxY mutants was compared to N2.

Due to the delayed egg release, Egl mutants harbor more unlaid eggs in the uterus than

N2 [35]. Fertilized eggs are released from the uterus before reaching the 26-cell stage

embryo [36]. We scored the number of eggs in the uterus (Figure 5, Supplementary Table

5) from the Day 2 egg-laying worms. The number of unlaid eggs in most NPxY mutants

was comparable to N2 egg retention. However, *pat-3(Y792F)* displayed fewer unlaid

eggs in the uterus than N2 (*p < 0.0019*, n = 50) or the other mutants. This analysis

showed that *pat-3(Y792F)* retained fewer eggs in the uterus while laying more eggs in

serotonin stimulations, suggesting that egg laying was accelerated. However, *pat-*

3(Y792F) maintained egg retention (average 10 eggs per animal), which is considered

within a typical range of egg bearing [37]. Therefore, the statistical significance failed to

support the argument that the *pat-3(Y792F)* mutation accelerated egg release in the

uterus.

Double non-phosphorylatable NPxY showed a distinct Egl phenotype.

In mammalian studies, the same knock-in of YY to FF in mouse β1 and β3 integrins

showed no overt behavioral phenotypes [6]. The *pat-3(kq24, YYFF)* laid a comparable

number of eggs in serotonin or fluoxetine during the sixty-minute incubation, suggesting

that the egg-laying appeared similar to N2 (for serotonin, *p* < 0.99, and fluoxetine, *p* <

0.88) (Figure 3, Supplementary Table 3). However, the egg retention of *pat-3(YYFF)*

showed a higher number of unlaid eggs in the uterus than in N2 (*p* < 0.0001). The *pat-*

3(YYFF) carried about thirty percent more unlaid eggs than N2 animals (Figure 5). This

result suggested that *pat-3(YYFF)* animals also delayed egg-laying while capable of

vulva contraction in response to serotonin and fluoxetine (Figure 3). The *pat-3(YYFF)*

was also tested for serotonin sensitivity because it includes the proximal non-

phosphorylation Y792F. Likewise, *pat-3(YYFF)* displayed a substantial egg-laying

increase by the reduced serotonin and fluoxetine concentration, 0.1 mg/ml (Figure 4).

The serotonin hypersensitivity of Y792F and YYFF mutations prompted us to test the

'distal' non-phosphorylation, *pat-3(Y804F)*, which displayed only a slight increase

(average 1.4 eggs per hour) in egg laying and failed to show the same level of

hypersensitivity as *pat-3(Y792F)* or *pat-3(YYFF)* (Figure 4).

The mechanosensation of βpat-3 integrin NPxY mutants.

C. elegans egg laying is a complex behavior that requires the coordination of muscles

and neurons, including vulva and uterine muscles and HSN and VC motor neurons [37].

While the current study primarily focuses on contractile tissues, our results imply the

potential involvement of integrin in neurons associated with egg laying.

In *pat-3(Y804E)*, serotonin treatment failed to improve egg laying to N2 levels, although

egg retention fell within the normal range of 10 to 15 [37] (Figures 3 and 4). The *pat-*

3(Y792F) and *pat-3(Y804A)* mutants were more or equally sensitive to serotonin

compared to N2 ($p < 0.0001$ and $p < 0.87$, respectively), but both were less sensitive to

fluoxetine than N2 ($p < 0.0001$ and $p < 0.01$). In contrast, *pat-3(YYFF)* displayed egg

laying comparable to N2 in both treatments while exhibiting a higher accumulation of

unlaid eggs than N2 ($p < 0.0001$). This suggests that their altered behavior may result

from impaired ability to respond to serotonin or to initiate egg laying due to decreased

sensitivity to stimuli from surroundings (Figure 3).

Several studies demonstrated the connections among mechanosensation, egg laying,

and integrin. Chalfie *et al.* (1985) identified a pair of PLM touch neurons extending along

the posterior lateral side and synapse with the HSN motor neurons [38]. Vibrational

stimulation inhibited egg laying, and this response was dependent on the ALM and PLM

touch neurons [39]. Ravie *et al.* (2018) proposed that mechanical pressure from egg

accumulation in the uterus triggers egg laying through a potential homeostatic

mechanism, mediated by a touch neuron [40]. Chen and Chalfie (2014) demonstrated that

pat-3 mosaic animals exhibited mechanosensation defects involved in ALM [22]. These

possible links among mechanosensation, egg laying, and integrin led us to hypothesize

that the phosphorylation of integrin NPxY is necessary for the touch response, and that

mutations in PAT-3 NPxY phosphorylation would result in a defective response to

mechanical stimuli [41].

Therefore, we evaluated the NPxY mutants using gentle touch analyses [42] (Figure 6,

Supplementary Table 6). Well-fed adult animals were touched (gentle prodding) on the

anterior or posterior half of the body [43,44]. The scoring involved assessing backing

(anterior touch) or sprinting forward movement (posterior touch) after the gentle touch

stimuli (see Experimental Procedures) [23]. Our results revealed that most NPxY mutants

exhibited significantly decreased touch responses on the anterior or posterior side of the

body (Figure 6). However, *pat-3(kq8041, Y804A)* and *pat-3(kq8043, Y804F)* appeared

wild-type in response to gentle touch stimuli.

Two mutants, *pat-3(kq7923, Y792F)* and *pat-3(kq8042, Y804E),* exhibited significant

insensitivity to gentle touch (Figure 6, *p < 0.0001* compared to N2 posterior), with the

defect predominantly affecting the posterior ends. To strengthen the analysis of the role

of membrane-proximal NPxY in mechanosensation, *pat-3(kq7921, Y792A)*, a tyrosine

removal mutation, was generated (see Experimental Procedures). The *pat-3(Y792A)*

mutation caused a disruptive change, abolishing the intact membrane-proximal NPxY

structure. Although *pat-3(Y792A)* animals displayed severely uncoordinated movement

with irregular and patchy dense bodies (Figure 2E), they were viable and fertile. As

expected, the touch response in *pat-3(Y792A)* was severely defective in the anterior

and posterior bodies, suggesting that the proximal NPxY motif is also essential for

motility and mechanosensation (Figure 6). The non-phosphorylatable NPxY mutants,

pat-3(kq7923, Y792F) and *pat-3(kq24, YYFF),* displayed a severe posterior touch

insensitivity ($p < 0.0001$). The mechanosensation defects coincided with phenylalanine

(F) or alanine (A) in the proximal motif, suggesting that phosphorylated proximal NPxY

is required for mechanosensation.

The connection between mechanosensation and egg-laying behaviors.

Our current study revealed the involvement of the *pat-3* NPxY motif in egg laying, which

modulates serotonin sensitivity, while identifying the role of NPxY in mechanosensation.

Therefore, we examined a mechanosensation mutant for egg laying to explore the

linkage between touch sensitivity and egg-laying behaviors. The *mec-4(u253)*, a null

allele of the MEC-4/ENaC mechanoreceptor [45], demonstrated severe touch defects in

both ends (Figure 6). While the egg laying of *mec-4(u253)* in serotonin was not different

from the wild type (*p < 0.995*, compared to N2), it failed to respond to fluoxetine (*p <*

0.0001) (Figure 3), suggesting that *mec-4(u253)* has a lowered serotonin level or

defective CeSERT/MOD-5 serotonin transporter. Therefore, the sensitivity to serotonin

or fluoxetine was also tested by treating *mec-4(u253)* with the reduced concentrations

of serotonin and fluoxetine (Figure 4). The hypersensitivity experiment showed *mec-*

4(u253) laid eggs in 0.1 mg/ml serotonin, comparable to *mec-4* in 3.0 mg/ml. Our data

suggested that the mechanosensation channel on TRN is linked to serotonin sensitivity.

Young adult N2 were grown on serotonin-containing (0.5 mg/ml) NGM plates to address

the linkage between serotonin sensitivity and mechanosensation for 24 hours. The

touch sensitivity of the serotonin-grown N2 showed significantly decreased touch

response in the posterior body ($p > 0.0001$) (Figure 6). In another experiment, *pat-*

3(kq7923, Y792F) and *pat-3(kq24, YYFF)* were grown on NGM plates containing 60 µM

reserpine, which is a VMAT (vesicular monoamine transporter) inhibitor previously

known for depleting monoamine neurotransmitters, serotonin and dopamine, at the

presynaptic ends [27]. In *C. elegans*, reserpine, an inhibitor of CAT-1/VMAT, blocks

dopamine and serotonin uptake into synaptic vesicles [27], extends its lifespan, and

confers thermotolerance[46] at a concentration of 60 µM. We expected that the reserpine

treatment would reduce serotonin release in the neurons. The *pat-3(YYFF)* and *pat-*

3(Y792F) mutants were grown on plates containing 60 µM reserpine for 24 hours. The

reserpine treatment in the *pat-3* alleles significantly suppressed their defective

mechanosensation (Figure 6), suggesting that serotonin negatively regulates the

mechanosensation and that serotonin depletion rescues the NPxY mutation-induced

mechanosensation defects.

DISCUSSION

Our study reports new functions of integrin NPxY motifs. We observed varied egg-laying

responses to serotonin and fluoxetine among *pat-3* alleles: *pat-3(kq8042, Y804E), pat-*

3(kq7923, Y792F), and *pat-3(kq24, YYFF)*. Functional analysis of the membrane-distal

NPxY motif reveals that the phosphomimetic *pat-3(Y804E)* mutant significantly reduces

egg laying under both treatments. The double non-phosphorylatable *pat-3(YYFF)* allele

shows regular egg laying to serotonin and fluoxetine but retains more unlaid eggs in the

uterus. In contrast, the proximal non-phosphorylatable *pat-3(Y792F)* mutant exhibits

reduced fluoxetine sensitivity, increased egg laying in serotonin. Additionally, *pat-*

3(Y792F) and *pat-3(YYFF)* mutants display hypersensitivity to serotonin and fluoxetine.

Beyond egg-laying behavior, these NPxY mutants, including *pat-3(kq7921, Y792A)*,

exhibit defective mechanosensation, dependent on serotonin levels. This suggests that

timely phosphorylation of the NPxY motif is critical for regulating vulval muscle

contraction and touch receptor neuron function.

The phosphomimetic *pat-3(kq8042, Y804E)*

Our data demonstrate that *pat-3* NPxY is essential for responding to serotonin and

fluoxetine. Our findings reveal that tyrosine phosphorylation disrupts vulva contraction,

as observed in *pat-3(Y804E)*, suggesting that the membrane-distal Y^{804} residue is a

critical switch for serotonin and fluoxetine responses. This result suggests that the distal

NPxY^{804} potentially interacts with MOD-5/CeSERT or serotonin receptors.

Phosphorylation of distal NPxY^{804} may impair the function of MOD-5 or multiple

serotonin receptors, leading to impaired contraction of the vulva muscles. Notably,

Carneiro *et al.* (2008) reported that αIIbβ3 platelet integrin binds to the C-terminal

domain of platelet SERT, where active β3 integrin enhances the serotonin transporter

activity of the SERT [47]. Interestingly, the egg retention of the phosphomimetic *pat-*

3(kq8042, Y804E) is similar to that of N2, indicating that constitutive phosphorylation of

the distal motif specifically reduces vulva muscle contraction. Fluorescence microscopy

(Figure 2) reveals a wild-type distribution of TLN-1 in the body wall muscle of the *pat-*

3(Y804E) mutant, demonstrating that phosphorylation of the distal NPxY is dispensable

for muscle organization. We conclude that an unknown factor, parallel to or downstream

of *pat-3(Y804E)*, such as *goa-1/Goα* [48], likely influences serotonin sensitivity and

muscle contraction. Segalat *et al.* show that the activated GOA-1, a Goα protein

homolog expressed in muscle or HSN, inhibits egg laying [48]. The PAT-3 distal NPxY

phosphorylation potentially activates GOA-1 or other Gα subunits and, consequently,

inhibits the contraction of the vulva muscle.

pat-3(kq24, YYFF), the double non-phosphorylation.

The *pat-3(YYFF)* exhibits hypersensitivity to serotonin and higher uterine egg retention.

This observation suggests that *pat-3(YYFF)* is defective in initiating egg laying, possibly

due to impaired ability to detect mechanical pressure from the oviparous uterus. This

defect could arise from an abnormal serotonin level to trigger vulva contraction [35] or the

failure of the utero-vulva feed-forward mechanism to stimulate egg-laying via serotonin

secretion [49]. Our results on *pat-3(YYFF)* indicate that the egg retention of the YYFF

mutation is a combined effect created by an interaction between Y^{792} and Y^{804}. The

Y792F or Y804F mutation appears normal in egg retention (Figure 4), which supports a

potential interaction between the two NPxY motifs. Further studies will elucidate the

molecular nature of any possible interactions between the two NPxY motifs in β-integrin.

The hypersensitivity of membrane-proximal non-phosphorylation, *pat-3(kq7923, Y792F)*.

In contrast, the *pat-3(Y792F)* mutant strongly responds to serotonin, but has a reduced

reaction to fluoxetine (Figure 4). In *C. elegans*, the endogenous serotonin is produced

and secreted from NSM (neurosecretory motor in the head), ADF (amphid dual ciliated

ending F sensory), and HSN (hermaphrodite-specific motor) neurons [50]. There is no

evidence of serotonin production in muscle cells, suggesting that the NPxY

phosphorylation regulates serotonin production or secretion in non-muscle cells. Our

results also indicate the importance of the membrane-proximal NPxY in serotonin

sensitivity. The hypersensitivity to serotonin in *pat-3(Y792F)* and *pat-3(YYFF)* suggests

that Y^{792} regulates the sensitivity to serotonin. Walser *et al.* (2017) report that density-

dependent phosphatase (*dep-1*) unphosphorylates the membrane-proximal NPxY. Our

finding suggests that the proximal NPxY is kept unphosphorylated during active egg

laying.

Serotonin hypersensitivity shows that integrin regulates egg laying in a serotonin-

dependent or independent manner.

The molecular nature of hypersensitivity to serotonin and fluoxetine is unknown and

requires further studies. However, there are several possible explanations. Our search

finds that the serotonin hypersensitivity is similar to the *mod-5/CeSERT* loss-of-function

phenotypes. Ranganathan *et al.* show the serotonin-specific hypersensitivity of *mod-5*

mutants in hyper-enhanced slowing behavior [31]. In contrast, fluoxetine stimulation of

egg laying in *tph-1(gm280)* mutants indicates that egg laying could be induced without

the endogenous serotonin. Our analysis suggests that *pat-3(Y792F)* and *pat-3(YYFF)*

are serotonin hypersensitive, indicating that the proximal NPxY-dependent egg laying

can be serotonin-independent. Kullyev et al. (2010) describe that fluoxetine binds to *ser-*

5 and *ser-7* receptors and causes serotonin-independent behaviors. We speculate that

fluoxetine independently stimulates serotonin receptors on muscles, which are

enhanced by Y792F and YYFF mutations. However, our data indicate that *pat-3(Y792F)*

and *pat-3(YYFF)* showed less sensitivity to fluoxetine than serotonin. This result

suggests an alternative scenario that the level of serotonin might be increased in *pat-*

3(Y792F) and *pat-3(YYFF)* due to potential inhibitory effects of the NPxY mutations in

mod-5/CeSERT functions.

Several NPxY mutants display defective mechanosensation.

The *pat-3(kq7921, Y792A)* exhibits mechanosensation defects in both the anterior and

posterior regions, whereas *pat-3(kq7923, Y792F)*, *pat-3(kq8042, Y804E)*, and *pat-*

3(kq24, YYFF) show the defect primarily in the posterior half. Our analysis suggests that

NPxY phosphorylation modulates mechanosensation (see Figure 5). While Chen *et al.*

(2014) demonstrated that *pat-3(-)* mosaic in ALM neurons exhibits touch insensitivity [22],

many NPxY mutants exhibit posterior touch defects, suggesting that the role of the

NPxY motif is biased to the PLM neurons. This aligns with the βpat-3 localization of

ALM and PLM neurons in Gettner *et al.* [9] and Amezquita *et al.* [51].

Overall, our data suggest that touch stimuli induce phosphorylation of the NPxY motifs.

The touch insensitivity at the posterior body in *pat-3(YYFF)* and *pat-3(Y792F)* indicates

that phosphorylation of NPxY motifs may contribute to the mechanosensation.

Mechanical stimuli might simultaneously or sequentially phosphorylate each NPxY,

transmitting mechanical information to mechanotransduction targets within the cell. The

NPVY[804] motif appears to bind a downstream molecule, UNC-112/kindlin. The

phosphorylated NPVY[804] may fail the interaction between UNC-112/kindlin and the

membrane distal NPxY [52]. We propose that the *pat-3(Y804E)* mechanosensation defect

arises from impaired interaction with UNC-112/kindlin.

Likewise, phosphorylation of the proximal NPIY[792] inhibits binding to TLN-1/talin [53]. In

our study, however, the mechanosensation of *pat-3(Y792F)* is defective in the posterior

end, suggesting phosphorylation of proximal NPIY[792] transiently occurs in PLM and

affects its ability to sense mechanical stimuli. In contrast, tyrosine removal or single

non-phosphorylatable mutations in the distal motif, such as *pat-3(Y804A)* and *pat-*

3(Y804F), show wild-type touch sensitivity in the anterior or posterior end. This

suggests that these mutations have minimal impact on the involvement of distal NPxY in

mechanosensation.

How does βPAT-3 integrin contribute to mechanosensation? One straightforward

interpretation is that the NPxY mutants cause unknown muscle anomalies, thereby

reducing the motor activity of muscles in response to touch stimuli. However, the βPAT-

3 protein is expressed in all tissues, particularly muscles and neurons [9]. Therefore, one

cannot rule out the potential role of βPAT-3 NPxY in neurons. Previous analysis shows

that the *pat-3* integrin directs the posterior-to-anterior migration of PLM axons during

development [54]. This suggests that the mechanosensation defect observed in NPxY

mutants stems from impaired PLM axon outgrowth. Consequently, stranded PLM axons

fail to detect touch stimuli in the posterior body. Another possibility is that integrin may

act as a direct mediator of mechanotransduction, potentially interacting with MEC-4 or

other touch receptors [55]. Tyrosine phosphorylation of the proximal NPIY[792] transiently

occurs, facilitating interaction with MEC-4. Upon touch stimulation, the MEC-4 channel

may induce phosphorylation of the proximal motif, subsequently triggering the release of

neurotransmitters at the PLM synapse to inhibit motor neurons such as HSN [42,49].

Our analysis shows that exogenous serotonin reduces the touch sensitivity of N2, while

inhibition of neurotransmitter secretion rescues the mechanosensation of NPxY mutants.

This demonstrates that serotonin reduces sensory response while promoting egg laying.

This result suggests that integrin NPxY phosphorylation potentially orchestrates the

secretion and production of serotonin. Integrin maintains serotonin at a level,

presumably, in neurons. This means that the phosphorylation or dephosphorylation of

pat-3 NPxY modulates serotonin levels in the tissues. Thus, non-phosphorylation of the

proximal NPxY causes a general increase in serotonin, which attenuates

mechanosensation.

In mammalian studies, β3 integrin YYFF knock-in mice exhibited a mild defect in clot

retraction and a tendency to rebleed [8]. However, other studies with β1 YYFF knock-in

mice failed to detect any developmental defects in the mutant animals [6]. Our study

presents hypersensitive egg laying and defective mechanosensation phenotypes

associated with the Y792F, Y804E, and YYFF mutations. This suggests that the

membrane-proximal and distal NPxY motifs in mice β1 or β3 integrin play a role in

reproductive and sensory behaviors similar to those of βPAT-3 *C. elegans* integrin.

EXPERIMENTAL PROCEDURES

Animals and Genetics

All animals were fed with *E. coli*, OP50, and cultured at room temperature (21^{o}C to 23^{o}C)

at all times. *C. elegans* N2 worm, AH3437 GFP::*tln-1*, and *mec-4(u253),* were obtained

from Caenorhabditis Genetics Center at St. Paul, MN. The CRISPR-Cas9 edited worms,

pat-3(kq8041, Y804A), pat-3(kq8042, Y804E), and *pat-3(kq8043, Y804F)* were

previously generated and described [12,56]. All *pat-3* NPxY mutants and other lines are

listed in Supplementary Table 1 (Table S1).

Genetic double mutant lines (Table S1) were generated by crossing homozygote males

from AH3437 GFP::*tln-1* [28] to each NPxY edited line hermaphrodite. From the F2

generation, a worm with 100% green (F3) progeny and the *pat-3* NPxY allele

homozygosity (by PCR genotyping) was selected for further studies.

Phenotype Analysis and Fluorescence Imaging.

Fecundity analyses of N2, *pat-3 NPxY*, and other mutants were conducted as follows.

Eggs were collected from worms grown on NGM plates (Day -2). After 48 hours (Day 0),

the hatched larvae were transferred to individual NGM plates, with one worm per plate.

At 72 hours (Day 1), each worm was transferred daily to a new NGM plate for four

additional days. After the worm was removed and placed onto a new plate, the progeny

on the previous plate were allowed to grow for one more day. The viable progeny were

then stored at 4°C, and the number of progeny on each plate was subsequently counted

and recorded for analysis.

For egg-laying assays, an adult worm was placed in 100 µl of M9 buffer per well in a 96-

well microtiter plate containing fluoxetine (0.5 mg/ml, Sigma-Aldrich) or serotonin (3

mg/ml). After a 15-minute acclimation period, the number of eggs laid was counted 60

minutes later. For hypersensitivity egg-laying assays, adult worms were placed in M9

buffer containing 0.1 mg/ml fluoxetine or serotonin. For egg retention assays, eggs were

transferred to an OP50-seeded plate and grown for four days at room temperature.

Adult worms were then placed on a microscope slide containing 50 to 100 µl of M9

buffer, and eggs were extruded using a 26-gauge needle. The extruded eggs were

counted and categorized by developmental stage, with eggs at the bean-shaped or later

embryonic stages scored as late-stage embryos.

For mechanosensation assays, following the protocol by Hobert *et al.* (1997), young

adult animals that had been fed *E. coli* OP50 for two generations without starvation

were gently stroked with eyebrow hairs ten times—five times on the anterior body and

five times on the posterior body [23]. Responses such as backing movement to anterior

touch and sprinting or twisting to posterior touch were scored as 1, while no responses

to touch were scored as 0. For serotonin mechanosensation assays, worms were grown

on NGM plates containing 0.5 mg/ml serotonin or 60 µM reserpine (Sigma-Aldrich) for

24 hours before the touch assays. Sample images were taken using a Nikon Eclipse Ni-

U fluorescence microscope and a Coolsnap Dyno digital monochrome camera

(Photomerics, Tucson, AZ). Photograph images were analyzed using NIS elements

(Nikon, Melville, NY).

The generation of CRISPR-Cas9 knock-in C. elegans

The CRISPR-Cas9 system was used to edit the *pat-3* locus to create the *kq24*,

YY792/804FF (YYFF) mutation, and other alleles. For *kq24*, the two crRNA,

ZQPATY1(CGAGAACCCAATCTACAAAC) and ZQPAT3B

(TTTAAAAATCCAGTATACGC) were obtained from the exon 8 of the *pat-3* gene

covering the membrane-proximal and distal NPxY and purchased from IDT Inc.,

Coralville, IA. The homologous directed repair template

(taaatttatcaaattatcatttttcagAACGAGAATCCAATTTTTAAGCAAGCCACGACAACATTCA

AGAACCCGGTTTTTGCAGGAAAAGCCAACTAAatagtttttatccttatatt) spans 48 bases

upstream and 38 bases downstream of the target site, tyrosine (Y^{792} and Y^{804}).

Mutations modified the third position of codons in the crRNA target sequence to a

synonymous codon. The repair DNA templates, crRNA, tracrRNA, and Cas9 nuclease

were custom-made by IDT Inc., Coralville, IA. A mixture of template DNA, two crRNA

(ZQAPTY1 and ZQPAT3B), tracrRNA (cat. no. 1073190), and Cas9 protein (cat. no.

1081058) was annealed at room temperature. The mixture was micro-injected into the

syncytial gonad of N2 animals (P0) with *dpy-10* co-CRISPR [57,58]. Homozygous mutants

were crossed back to N2 two times. The mutation was confirmed by sequencing DNA

molecules extracted from homozygous animals [12]. The generation of other edits was

carried out similarly to the techniques mentioned above. The nucleotide sequences of

crRNA and templates are listed in Supplementary Table S2 (Table S2).

Statistical Analysis

Statistical analysis of multiple *C. elegans* strains was conducted using All Pairwise

Comparisons - Tukey HSD (Honestly Significant Difference) method [59]. The egg-laying

assay utilized a Poisson regression model, while the regular and late-stage egg

retention assays employed the Comparison of Means. A significance level was set at *p*

< 0.05. Touch sensing assays were subjected to logistic regression and all pairwise

comparisons for Tukey HSD. All data were analyzed using JMP Pro (version 17.0, SAS

Institute, Cary, NC) software. Pairwise comparison data with *p > 0.05* can be found in

Supplementary Tables 3, 4, 5, and 6.

Acknowledgments

The Office of Engaged Learning at Baylor University URSA Research Fund supported

this work. The *pat-3* mutants were generated for the BIO4108 Laboratory Class. The

authors acknowledge Catherine Ravikumar, Arlyn Alcid, and Jeancarlo Gutierrez for

their help during the initial stage of the project. The authors also appreciate Drs. Erin

Cram and Ilya Ruvinsky for their comments on the initial draft.

REFERENCES

1.   Hynes RO. Integrins: bidirectional, allosteric signaling machines. *Cell*. Sep 20
3   2002;110(6):673-87.
4

2.   Arnaout MA, Mahalingam B, Xiong JP. Integrin structure, allostery, and bidirectional
5   signaling. *Annu Rev Cell Dev Biol*. 2005;21:381-410.
6   https://doi.org10.1146/annurev.cellbio.21.090704.151217.
7

3.   Qiu Z, Sheesley P, Ahn JH, Yu EJ, Lee M. A Novel Mutation in an NPXY Motif of beta
8   Integrin Reveals Phenotypes Similar to him-4/hemicentin. *Front Cell Dev Biol*.
9   2019;7:247. https://doi.org10.3389/fcell.2019.00247.
10

4.   Theodosiou M, Widmaier M, Bottcher RT, et al. Kindlin-2 cooperates with talin to
11   activate integrins and induces cell spreading by directly binding paxillin. *Elife*. Jan 27
12   2016;5:e10130. https://doi.org10.7554/eLife.10130.
13

5.   Sakai T, Jove R, Fassler R, Mosher DF. Role of the cytoplasmic tyrosines of b1A integrins
14   in transformation by v-src. *Proceedings of the National Academy of Science, USA*.
15   2001;98(7):3808-3813.
16

6.   Czuchra A, Meyer H, Legate KR, Brakebusch C, Fassler R. Genetic analysis of beta1
17   integrin "activation motifs" in mice. *J Cell Biol*. Sep 11 2006;174(6):889-99.
18

7.   Meves A, Geiger T, Zanivan S, DiGiovanni J, Mann M, Fassler R. Beta1 integrin
19   cytoplasmic tyrosines promote skin tumorigenesis independent of their phosphorylation.
20   *Proc Natl Acad Sci U S A*. Sep 13 2011;108(37):15213-8.
21   https://doi.org10.1073/pnas.1105689108.
22

8.   Law DA, DeGuzman FR, Heiser P, Ministri-Madrid K, Killeen H, Phillips DR. Integrin
23   cytoplasmic tyrosine motif is required for outside-in aIIbb3 signalling and platelet
24   function. *Nature*. 1999;401:808-811.
25

9.   Gettner SN, Kenyon C, Reichardt LF. Characterization of bpat-3 heterodimers, a family of
26   essential integrin receptors in *C. elegans*. *Journal of Cell Biology*. 1995;129:1127-1141.
27

10.   Lee M, Cram EJ, Shen B, Schwarzbauer JE. Roles for beta(pat-3) integrins in development
28   and function of Caenorhabditis elegans muscles and gonads. *J Biol Chem*. Sep 28
29   2001;276(39):36404-10. https://doi.org10.1074/jbc.M105795200
30   M105795200 [pii].
31

11.   Itoh B, Hirose T, Takata N, et al. SRC-1, a non-receptor type of protein tyrosine kinase,
32   controls the direction of cell and growth cone migration in C. elegans. *Development*. Dec
33   2005;132(23):5161-72. https://doi.org10.1242/dev.02103.
34

12.   Hanna J, Ramani S, Williams T, et al. Three alleles in the pat-3 locus of Caenorhabditis
35   elegans: mutations in the membrane-distal NPxY phosphotyrosine motif. *MicroPubl Biol*.
36   Aug 15 2020;2020. https://doi.org10.17912/micropub.biology.000291.
37

13.   Trent C, Tsung N, Horvitz HR. Egg-laying defective mutants of the nematode *C. elegans*.
38   *Genetics*. 1983;104:619-647.
39

14.   Moerman DG, Fire A. Muscle: Structure, function, and development. In: Riddle DL, ed. *C.
40   elegans II*. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press; 1997:417-470.

15. Lee M, Shen B, Schwarzbauer JE, Ahn J, Kwon J. Connections between integrins and Rac GTPase pathways control gonad formation and function in C. elegans. *Biochim Biophys Acta*. May 25 2005;1723(1-3):248-55. https://doi.orgS0304-4165(05)00009-7 [pii] 10.1016/j.bbagen.2005.01.003.

16. Qiu Z, Park A, Wang L, Wilsey R, Lee M. The RGD (Arg-Gly-Asp) is a potential cell-binding motif of UNC-52/PERLECAN. *Biochem Biophys Res Commun*. Jan 1 2022;586:143-149. https://doi.org10.1016/j.bbrc.2021.11.083.

17. Wilsey R, Hodge S, Kenney K, et al. Two alleles of unc-52 locus disrupting potential cell-binding motif of UNC-52. *MicroPubl Biol*. May 17 2020;2020. https://doi.org10.17912/micropub.biology.000250.

18. Minniti AN, Labarca M, Hurtado C, Brandan E. Caenorhabditis elegans syndecan (SDN-1) is required for normal egg laying and associates with the nervous system and the vulva. *J Cell Sci*. Oct 1 2004;117(Pt 21):5179-90. https://doi.org10.1242/jcs.01394.

19. Goodman MB. Mechanosensation. *WormBook*. 2006:1-14. https://doi.org10.1895/wormbook.1.62.1.

20. Chalfie M, Hart AC, Rankin CH, Goodman MB. Assaying mechanosensation. *WormBook*. Jul 31 2014. https://doi.org10.1895/wormbook.1.172.1.

21. Zuidema A, Wang W, Sonnenberg A. Crosstalk between Cell Adhesion Complexes in Regulation of Mechanotransduction. *Bioessays*. Nov 2020;42(11):e2000119. https://doi.org10.1002/bies.202000119.

22. Chen X, Chalfie M. Modulation of C. elegans touch sensitivity is integrated at multiple levels. *J Neurosci*. May 7 2014;34(19):6522-36. https://doi.org10.1523/JNEUROSCI.0022-14.2014.

23. Hobert O, Moerman DG, Clark KA, Beckerle MC, Ruvkun G. A conserved LIM protein that affects muscular adherens junction integrity and mechanosensory function in Caenorhabditis elegans. *J Cell Biol*. Jan 11 1999;144(1):45-57.

24. Williams BD, Waterston RH. Genes critical for muscle development and function in *Caenorhabditis elegans* identified through lethal mutations. *Journal of Cell Biology*. 1994;124:475-490.

25. Lin X, Qadota H, Moerman DG, Williams BD. *C. elegans* PAT-6/Actopaxin plays a critical role in the assembly of integrin adhesion complexes *in vivo*. *Current Biology*. 2003/5/27 2003;13(11):922-932.

26. Rogalski TM, Mullen GP, Gilbert MM, Williams BD, Moerman DB. The UNC-112 gene in *Caenorhabditis elegans* encodes a novel component of cell-matrix adhesion structure required for integrin localization in the muscle cell membrane. *Journal of Cell Biology*. 2000;150(1):253-264.

27. Duerr JS, Frisby DL, Gaskin J, et al. The cat-1 gene of Caenorhabditis elegans encodes a vesicular monoamine transporter required for specific monoamine-dependent behaviors. *J Neurosci*. Jan 1 1999;19(1):72-84. https://doi.org10.1523/JNEUROSCI.19-01-00072.1999.

28. Walser M, Umbricht CA, Frohli E, Nanni P, Hajnal A. beta-Integrin de-phosphorylation by the Density-Enhanced Phosphatase DEP-1 attenuates EGFR signaling in C. elegans. *PLoS Genet*. Jan 2017;13(1):e1006592. https://doi.org10.1371/journal.pgen.1006592.

29. Trent C. *Genetic and behavioral studies of the egg-laying system in Caenorhabditis elegans* [Dissertation]. Cambridge, MA.: Massachusetts Institute of Technology.; 1983.

30. Waggoner LE, Zhou GT, Schafer RW, Schafer WR. Control of alternative behavioral states by serotonin in Caenorhabditis elegans. *Neuron*. 1998;21(1):203-214.

31. Ranganathan R, Sawin ER, Trent C, Horvitz HR. Mutations in the Caenorhabditis elegans serotonin reuptake transporter MOD-5 reveal serotonin-dependent and -independent activities of fluoxetine. *J Neurosci*. Aug 15 2001;21(16):5871-84. https://doi.org10.1523/JNEUROSCI.21-16-05871.2001.

32. Choy RK, Thomas JH. Fluoxetine-resistant mutants in C. elegans define a novel family of transmembrane proteins. *Mol Cell*. Aug 1999;4(2):143-52. https://doi.org10.1016/s1097-2765(00)80362-7.

33. Garcia-Alvarez B, de Pereda JM, Calderwood DA, et al. Structural determinants of integrin recognition by talin. *Mol Cell*. Jan 2003;11(1):49-58.

34. Sadeghian F, Ibrahim I, Ravichandran L, et al. An integrin binding motif in TLN-1/talin plays a minor role in motility and ovulation. *MicroPubl Biol*. 2023;2023. https://doi.org10.17912/micropub.biology.000726.

35. Tanis JE, Moresco JJ, Lindquist RA, Koelle MR. Regulation of serotonin biosynthesis by the G proteins Galphao and Galphaq controls serotonin signaling in Caenorhabditis elegans. *Genetics*. Jan 2008;178(1):157-69. https://doi.org10.1534/genetics.107.079780.

36. Wood WB. The Nematode. *Caenorhabditis elegans*. *Cold Spring Harbor Laboratory Press*. 1988.

37. Schafer WF. Genetics of egg-laying in worms. *Annu Rev Genet*. 2006;40:487-509. https://doi.org10.1146/annurev.genet.40.110405.090527.

38. Chalfie M, Sulston JE, White JG, Southgate E, Thomson JN, Brenner S. The neural circuit for touch sensitivity in Caenorhabditis elegans. *J Neurosci*. Apr 1985;5(4):956-64.

39. Zhang M, Chung SH, Fang-Yen C, et al. A self-regulating feed-forward circuit controlling C. elegans egg-laying behavior. *Curr Biol*. Oct 14 2008;18(19):1445-55. https://doi.org10.1016/j.cub.2008.08.047.

40. Ravi B, Garcia J, Collins KM. Homeostatic Feedback Modulates the Development of Two-State Patterned Activity in a Model Serotonin Motor Circuit in Caenorhabditis elegans. *J Neurosci*. Jul 11 2018;38(28):6283-6298. https://doi.org10.1523/JNEUROSCI.3658-17.2018.

41. Goodman MB, Schwarz EM. Transducing touch in Caenorhabditis elegans. *Annu Rev Physiol*. 2003;65:429-52. https://doi.org10.1146/annurev.physiol.65.092101.142659 092101.142659 [pii].

42. Li W, Kang L, Piggott BJ, Feng Z, Xu XZ. The neural circuits and sensory channels mediating harsh touch sensation in Caenorhabditis elegans. *Nat Commun*. 2011;2:315. https://doi.org10.1038/ncomms1308.

43. Chalfie M. Touch receptor development and function in Caenorhabditis elegans. *J Neurobiol*. Oct 1993;24(10):1433-41. https://doi.org10.1002/neu.480241013.

44. Chalfie M, Au M. Genetic control of differentiation of the Caenorhabditis elegans touch receptor neurons. *Science*. Feb 24 1989;243(4894 Pt 1):1027-33.

45. O'Hagan R, Chalfie M, Goodman MB. The MEC-4 DEG/ENaC channel of Caenorhabditis elegans touch receptor neurons transduces mechanical signals. *Nat Neurosci*. Jan 2005;8(1):43-50. https://doi.orgnn1362 [pii] 10.1038/nn1362.

46. Srivastava D, Arya U, SoundaraRajan T, Dwivedi H, Kumar S, Subramaniam JR. Reserpine can confer stress tolerance and lifespan extension in the nematode C. elegans. *Biogerontology*. Oct 2008;9(5):309-16. https://doi.org10.1007/s10522-008-9139-5.

47. Carneiro AM, Cook EH, Murphy DL, Blakely RD. Interactions between integrin alphaIIbbeta3 and the serotonin transporter regulate serotonin transport and platelet aggregation in mice and humans. *J Clin Invest*. Apr 2008;118(4):1544-52. https://doi.org10.1172/JCI33374.

48. Segalat L, Elkes DA, Kaplan JM. Modulation of serotonin-controlled behaviors by Go in Caenorhabditis elegans. *Science*. Mar 17 1995;267(5204):1648-51.

49. Ravi B, Nassar LM, Kopchock RJ, 3rd, Dhakal P, Scheetz M, Collins KM. Ratiometric Calcium Imaging of Individual Neurons in Behaving Caenorhabditis Elegans. *J Vis Exp*. Feb 7 2018;(132). https://doi.org10.3791/56911.

50. Dag U, Nwabudike I, Kang D, et al. Dissecting the functional organization of the C. elegans serotonergic system at whole-brain scale. *Cell*. Jun 8 2023;186(12):2574-2592 e20. https://doi.org10.1016/j.cell.2023.04.023.

51. Amezquita J, Desbois M, Opperman KJ, et al. Integrin adhesome axis inhibits the RPM-1 ubiquitin ligase signaling hub to regulate growth cone and axon development. *PLoS Genet*. Dec 2024;20(12):e1011496. https://doi.org10.1371/journal.pgen.1011496.

52. Bledzka K, Bialkowska K, Nie H, et al. Tyrosine phosphorylation of integrin beta3 regulates kindlin-2 binding and integrin activation. *J Biol Chem*. Oct 1 2010;285(40):30370-4. https://doi.org10.1074/jbc.C110.134247.

53. Anthis NJ, Haling JR, Oxley CL, et al. Beta integrin tyrosine phosphorylation is a conserved mechanism for regulating talin-induced integrin activation. *J Biol Chem*. Dec 25 2009;284(52):36700-10. https://doi.org10.1074/jbc.M109.061275.

54. Gettner SN. *Isolation and functional analysis of Caenorhabditis elegans integrin bpat-3* [Dissertation]. San Francisco, CA: University of California; 1994.

55. Shakibaei M, Mobasheri A. Beta1-integrins co-localize with Na, K-ATPase, epithelial sodium channels (ENaC) and voltage activated calcium channels (VACC) in mechanoreceptor complexes of mouse limb-bud chondrocytes. *Histol Histopathol*. Apr 2003;18(2):343-51. https://doi.org10.14670/HH-18.343.

56. Park A, Qiu Z, Bumm J, Lee M. A novel mutation in unc-112/kindlin locus causes distal tip cell migration defects. *MicroPubl Biol*. Jun 8 2020;2020. https://doi.org10.17912/micropub.biology.000265.

57. Paix A, Folkmann A, Rasoloson D, Seydoux G. High Efficiency, Homology-Directed Genome Editing in Caenorhabditis elegans Using CRISPR-Cas9 Ribonucleoprotein Complexes. *Genetics*. Sep 2015;201(1):47-54. https://doi.org10.1534/genetics.115.179382.

58. Dickinson DJ, Goldstein B. CRISPR-Based Methods for Caenorhabditis elegans Genome Engineering. *Genetics*. Mar 2016;202(3):885-901. https://doi.org10.1534/genetics.115.182162.

1   59.   Tukey JW. Comparing individual means in the analysis of variance. *Biometrics*. Jun
2   1949;5(2):99-114.
3   60.   Francis R, Waterston RH. Muscle cell attachment in *Caenorhabditis elegans*. *Journal of
4   Cell Biology*. 1991;114:465-479.
5

1 Figure 1. Amino acid sequences of wild-type (+) and mutant cytoplasmic domain (CTD) of *C.*

2 *elegans* βpat-3 integrin.

```
Homo sapiens β1A   775WDTGENPIYKSAVTTVVNPKYEGK798
βpat-3(+)          784WDTNENPIYKQATTTFKNPVYAGKAN809
βpat-3(Y792A)      784WDTNENPIAKQATTTFKNPVYAGKAN809
βpat-3(Y792F)      784WDTNENPIFKQATTTFKNPVYAGKAN809
βpat-3(YYFF)       784WDTNENPIFKQATTTFKNPVFAGKAN809
βpat-3(Y804A)      784WDTNENPIYKQATTTFKNPVAAGKAN809
βpat-3(Y804E)      784WDTNENPIYKQATTTFKNPVEAGKAN809
βpat-3(Y804F)      784WDTNENPIYKQATTTFKNPVFAGKAN809
```

6 Amino acid residues in *C. elegans* βPAT-3 cytoplasmic tail sequence and their functions.

7 The amino acid sequence of the *Homo sapiens* β1A integrin CTD is aligned

8 with *C. elegans pat-3 (+),* wild-type CTD, and mutant CTD such as *pat-3(kq7921,*

9 *Y792A), pat-3(kq7923, Y792F), pat-3(kq24, YYFF), pat-3(kq8041, Y804A), pat-*

10 *3(kq8042, Y804E), and pat-3(kq8043, Y804F).* Reds indicate mutations.

Figure 2. *GFP::tln-1/talin* localization in *pat-3* NPxY mutants

The five double mutants were generated by crossing GFP::*tln-1* to *pat-3* NPxY mutants.

In every picture, horizontal dots represent dense bodies (arrows), and thin lines

between the dense bodies represent M lines [60]. All images are taken under a 60X

objective lens, except panel F (40X). Homozygous mutants are shown here. A. AH3437

GFP::tln-1/talin body wall muscle of adult worm (control), B. *GFP::tln-1/talin; pat-*

3(kq8042, Y804E), C. *GFP::tln-1/talin; pat-3(kq8041, Y804A),* D. *GFP::tln-1/talin;pat-*

3(kq7923, Y792F), E. *GFP::tln-1/talin;pat-3(kq7921, Y792A),* F. *GFP::tln-1/talin ;pat-*

3(kq24, YYFF). Bars = 10 µm.

Figure 3. Egg-laying assays of integrin mutants. Results significantly different from N2 were marked with * ($p < 0.05$). Brackets indicate significant differences between the treatments. Multiple adult worms (n=40 or 50) were used for each experiment. *All Pairwise Comparisons - Tukey HSD.

2
3

Figure 4. *pat-3* NPxY mutants display hypersensitivity to serotonin and fluoxetine. Egg-laying assays were repeated in *pat-3* NPxY mutants at a concentration of serotonin and fluoxetine, which does not stimulate egg laying in N2 (0.1 mg/ml serotonin or fluoxetine). Multiple adult worms (n=40) were used for each experiment. Brackets indicate significant differences between the treatments. Results significantly different from N2 were marked with * ($p < 0.05$). *All Pairwise Comparisons - Tukey HSD.

3   Figure 5. Egg retention assays of integrin mutants. Results significantly different from

4   N2 were marked with * ($p < 0.05$). Multiple worms (n = 35 or 50) were used for each

5   scoring. Bars represent total eggs present at the time of assay. * All Pairwise

6   Comparisons – One-way analysis of variance.

Figure 6. Mechanosensation analysis of *pat-3* NPxY mutants and *mec-4*.

The single asterisk (*) mark on *pat-3(kq24)* anterior indicates $p = 0.048$, while all other

results significantly different from N2 are marked with ** ($p < 0.05$). Brackets indicate

significant differences between the treatment with serotonin (5-HT) or reserpine (RS)

and no treatment. All Pairwise Comparisons – Tukey HSD analyzed the averages of

each test (n = 50). 5-HT: serotonin treatment (0.5 mg/ml); RS: reserpine (60 µM).

Worms were treated with the drugs for 24 hours before touch assays.

1   Table 1. Summary of *pat-3* NPxY mutant phenotypes.

Location in the CTD	Allele Designation	Mutations	Phenotypes
Y792 series: **Membrane-proximal** NPxY	*pat-3(kq7921, Y792A)*	Y792 to alanine (A), Y removal, disruptive mutation	Viable, severely uncoordinated movement, unable to lay eggs, and touch insensitive on both ends of the body.
	pat-3(kq7923, Y792F)	Y792 to phenylalanine (F), non-phosphorylatable mutation	Viable, normal movement, normal egg laying, touch insensitivity on posterior half of the body
Y804 series: **Membrane-distal** NPxY	*pat-3(kq8041, Y804A)*	Y804 to alanine (A), Y removal, disruptive mutation	Viable, normal movement, normal egg laying and mechanosensation
	pat-3(kq8042, Y804E)	Y804 to glutamic acid (E), phosphomimetic mutation	Viable, less egg laying, touch insensitive
	pat-3(kq8043, Y804F)	Y804 to phenylalanine (F), non-phosphorylatable mutation	Viable, normal movement, normal egg laying and mechanosensation
YYFF: Double mutant	*pat-3(kq24, YYFF)*	Both Y792 and 804 to phenylalnine (F), double non-phosphorylatable mutation	Viable, normal movement, normal egg laying, retain unlaid eggs in the uterus, touch insensitivity on posterior half of the body.