Assessing the impact of character evolution models on phylogenetic and macroevolutionary inferences from fossil data

David F. Wright[1,2,3,4] **and Melanie J. Hopkins**[4]

[1] *Sam Noble Oklahoma Museum of Natural History, University of Oklahoma, 2401 Chautauqua Avenue, Norman, OK, 73072, USA*

[2] *School of Geosciences, University of Oklahoma, 100 E Boyd Street, Norman, OK, 73019, USA*

[3] *Department of Paleobiology, National Museum of Natural History, Smithsonian Institution, Washington, DC, USA*

[4] *American Museum of Natural History, Division of Paleontology (Invertebrates), Central Park West at 79th St., New York, NY 10024, USA*

Corresponding author: David F. Wright (wrightdf@ou.edu)

Key words: paleobiology, morphology, phylogenetics, Trilobita

Abstract

Understanding the evolution and phylogenetic distribution of morphologic traits is fundamental to macroevolutionary research. Despite decades of major advances and key insights from molecular systematics, organismal anatomical features remain a key source of biological data for both inferring phylogenies and investigating patterns of trait evolution among fossil and extant species. In paleobiology, morphologic characters are typically the only source of information available for reconstructing evolutionary trees. Systematists working with fossil data must make decisions regarding how morphological characters are modeled, whether they are continuous or categorical, and how to address biological sources of rate variation. To determine the impact of how different models of morphological evolution influence phylogenetic inferences and downstream comparative analyses of fossil data, we competed a series of increasingly complex model configurations of character evolution to a dataset of Cambrian-Ordovician trilobites containing both discrete morphological characters and continuous traits. Competed models vary in complexity, ranging from simple constant rate scenarios with only discrete categorical traits, to complex evolutionary models including both discrete and quantitative traits across multiple ecological partitions while accounting for multiple sources of rate variation. We compared topological distributions across competed models by visualizing their distances in multidimensional treespace. Results indicate the addition of continuous traits dramatically improves support for complex evolutionary scenarios. Remarkably, we find model complexity also has a major impact on which regions of treespace are explored, which suggests a critical need to carefully model morphologic evolution even when systematists are only interested in patterns of phylogenetic relationships alone.

1. Introduction

Investigating patterns of anatomical character evolution is a major component of macroevolutionary biology (Simpson, 1944; Harmon et al., 2010). Character evolution models are fundamental to documenting evolutionary trends (Hopkins and Lidgard, 2012), interpreting patterns of morphological disparity in the fossil record (Foote, 1997; Erwin, 2007), and testing alternative hypotheses of trait evolution using time-calibrated phylogenetic trees (Revell and Harmon, 2022). Moreover, despite decades of major advances and key insights from molecular systematics, organismal anatomical features remain a key source of biological data for both inferring phylogenies and studying patterns of trait evolution among fossil and extant species (Ronquist et al., 2012; Wright and Hillis, 2014; Bapst et al., 2018; A. Wright, 2019). In some disciplines, such as paleobiology, anatomical characters are frequently the most common and perhaps only source of information available for reconstructing phylogenies (Smith, 1994; Wiley and Lieberman, 2011). Most phylogenies of fossil taxa have historically been inferred using parsimony methods. However, over the last several years, systematists working with fossil data have increasingly begun using Bayesian hierarchical modeling approaches to simultaneously infer a tree topology, date divergences, and estimate macroevolutionary parameters (Warnock and Wright, 2020; Wright et al., 2021). In particular, Bayesian "tip-dating" phylogenetic methods that implement the fossilized birth-death process (FBD) (Stadler, 2010; Heath et al., 2014; Gavryushkina et al., 2014; Stadler et al., 2018) are especially appealing to paleobiologists because they can leverage both morphological and stratigraphic data from the fossil record to estimate a probability distribution of evolutionary histories (i.e., relationships and macroevolutionary dynamics) (Wright, 2017a; Paterson et al., 2019; King and Beck, 2020).

Simulations show the inclusion of fossil age information under the FBD process can improve inferred tree topologies for tip-dated analyses relative to undated approaches (Barido-Sottani et al., 2020; Mongiardino Koch et al., 2021). However, the inclusion of fossil age data in tip-dating studies predominantly impacts the probability of tree shapes and diversification histories (May and Rothfels, 2023), whereas morphological character data remain the primary source of information regarding the probability of alternative topologies (Warnock and Wright, 2020; Wright et al., 2021). Since appropriately modeling fossil age data can improve phylogenetic inferences (Barido-Sottani et al., 2020), it seems even more prudent to determine the extent to which inferences may also be influenced by different approaches to handling assumptions about character evolution, especially in empirical studies where the tree topology is often the parameter of greatest interest. Unfortunately, when one considers the complexity of information captured by different kinds of morphological characters and the vast number of ways in which they could be modeled, determining how to appropriately model character evolution remains a challenging task.

Discrete characters comprise the majority of morphological matrices in the literature. The simplest implementation of a discrete character evolution model is the Mk model of Lewis

(2001), which assumes characters are conditionally independent, evolve at uniform rates, and have symmetrical transition frequencies between character states. Although the Mk model has been shown to outperform parsimony under a variety of scenarios (Wright and Hillis, 2014), many empirical matrices exhibit dramatic heterogeneity in morphologic rates among characters, across lineages, and through time (Lloyd et al., 2012; Hopkins and Smith, 2015; Wright, 2017b). Consequently, a variety of approaches have been developed to relax assumptions about character evolution and accommodate different sources of rate variation among characters and lineages (Wagner and Marcot, 2010; Drummond and Stadler, 2016; Wright et al., 2016). For example, rate variation among characters is often modeled using a parametric distribution, such as a gamma or lognormal (Wagner, 2012; Harrison and Larsson, 2015; Wright, 2019). The variance of the rate distribution, which determines its shape, is a single parameter estimated from the data. Because per-character rates are independent draws from the same distribution, character evolution is implicitly modeled as having a common generating process but with variable rates. However, not all characters might be expected to evolve according to the same evolutionary "rules" or generating mechanisms. In such instances, characters may be partitioned into categorically different groups, each having unique parameters to describe their dynamics. For example, different anatomical regions may vary in morphologic rates or involve developmentally-linked correlated character changes (Clark and Middleton, 2008; Wagner, 2018). Alternatively, characters with ecological or functional significance may evolve at higher rates than non-ecologic traits due to elevated selective pressures that fluctuate in direction over time (Blomberg et al. 2013).

Another source of morphologic data comes from continuous characters, which can be modeled using diffusion processes such as Brownian motion (BM) or related models (Felsenstein, 1988). Although less frequently used to infer phylogenies, continuous characters may be used to infer both tree topology (Parins-Fukuchi, 2018a) and divergence times (Álvarez-Carretero et al., 2019). When modeled directly, continuous characters have been shown to contain at least as much, if not greater, phylogenetic signal than discrete characters (Parins-Fukuchi, 2018b). However, in many empirical matrices, continuous variation is binned into categories and treated as a discrete character (Rae, 1998), and the extent to which this process of discretization results in the loss of phylogenetic information is unclear (Rae, 1998).

The wide variety of ways to model morphological evolution leads to a large number of possible model configurations, and computational tractability prevents evaluation of all possible candidate models using conventional model selection techniques, such as Bayes Factors (Wright et al., 2021). Unlike substitution models for molecular sequence data, few frameworks exist for determining which character model configuration provides a best fit to morphologic data in advance (Lanfear et al., 2017). Remarkably, the practice of fitting molecular substitution models to sequence data prior to phylogeny reconstruction has previously been argued to be unnecessary if the purpose is solely to infer the tree topology or conduct ancestral state reconstruction, as both simulated and empirical results tend to be similar to the most parameter-rich substitution model

regardless of the generating process or best fit to empirical data (Abadi et al., 2019). If these results extend to morphological matrices of fossil taxa, then the time-consuming steps involved in model selection may be similarly unnecessary, thereby liberating paleobiologists from the need to carry out a series of difficult and computationally expensive analyses prior to phylogeny reconstruction. In this study, we investigate this question empirically using a morphological dataset of olenid trilobites, a major clade of extinct arthropods. Specifically, we (1.) compete a series of increasingly complex character model configurations to these data, (2.) conduct model selection to rank configurations by their relative fit, and (3.) compare the evolutionary implications associated with inferences made under each configuration. The goal of this study is to assess the degree to which different character model configurations alone may impact phylogenetic and macroevolutionary inferences in empirical analyses of fossil data.

2. Materials and methods

(a) Data
We focus our analyses on trilobites, a species-rich and morphologically diverse clade of Paleozoic arthropods, as a model system for evaluating character evolution models in phylogenetic paleobiology. We selected a morphological character matrix from a previously published parsimony-based analysis of the Cambrian – Ordovician trilobite family Olenidae that sampled both discrete and continuous characters (Monti and Confalonieri, 2019), and subsequently updated and modified the character matrix in two ways. First, we coded two additional species from the subfamily Balnibarbiinae, which were unrepresented in the original matrix (*Balnibarbi pulvurea* and *Cloacaspis senilis*, see Fortey 1974; Hopkins, 2019). Next, we noted the original matrix contained a substantial number of cells with missing entries for continuous characters. Since we are primarily interested in how different character models may impact downstream inferences rather than reevaluating previous authors' results, for computational tractability (Barido-Sotanni et al., 2023), we removed a combination of taxa and continuous characters in an effort to retain the maximum number of entries with no missing cells. However, all discrete morphologic characters from the original matrix were retained. The final dataset includes 38 species, 62 discrete characters, and 12 continuous traits, which is a relatively large character matrix for fossil marine invertebrates (Barido-Sottani et al., 2020), and we believe likely representative of many published morphologic datasets of similar size and taxonomic scope. Stratigraphic occurrence data for each fossil species were compiled from the literature and species were assigned ages corresponding to geologic intervals bounding their first appearance intervals using Gradstein et al. (2012). The character matrix and data are provided in the Supplemental Information.

(b) Character evolution models
We competed five morphologic model configurations that range from simple to complex, with more parameter-rich models accounting for increasingly more biologically complex evolutionary

dynamics (Figure 1). In the first three model configurations, all morphologic characters analyzed are discrete traits (e.g., "absence" or "presence" of a character), whereas the two most complex configurations also include continuous characters and implement separate character evolution models for discrete versus quantitative traits.

The simplest model configuration (**S**) corresponds to a constant rates model assuming a "strict" morphological clock and an equal rates implementation of the Mk model of discrete character evolution (Lewis, 2001). In contrast, the "relaxed" rates model (**R**) allows for discrete character rates to vary among characters and through time. Specifically, rates vary among characters following a lognormal distribution (Wagner, 2012), and across the phylogeny according to an uncorrelated lognormal clock model (Heled and Drummond, 2012). Next, the configuration of the **R+E** model modifies the **R** model configuration by partitioning "ecologic" characters and estimating separate lognormal rate distributions for traits with presumed ecological significance vs. non-ecological traits (**R+E**). Ecological significance was defined as any character relevant to sensory systems (e.g. size and placement of eyes) or behavior (e.g. molting or defense). A total of 13 characters (20%) were assigned to the "ecological" partition. Next, the **RE+C** model configuration uses the **R+E** configuration for discrete traits but includes continuous characters and models them as quantitative traits evolving according to a single rate Brownian motion process (Felsenstein, 1985). Finally, the most complex model implements a multivariate BM model for continuous traits (**mREC**), which accommodates rate variation among quantitative characters and accounts for evolutionary integration among correlated traits (Caetano and Harmon, 2019).

(c) Phylogenetic and macroevolutionary methods

All analyses were conducted using "tip-dating" Bayesian phylogenetic methods incorporating the fossilized birth-death (FBD) process (Stadler, 2010; Heath et al., 2014; Wright, 2017). The tree prior is an important component of Bayesian phylogenetic inference, especially for datasets including many (or only) fossil taxa (Barrido-Sotanni et al., 2020; Warnock and Wright, 2020). However, fossil occurrence times are not just priors for node ages and/or tip occurrences, they are also data, and changing our assumptions about how the tree prior is handled may inadvertently prevent us from discerning which differences are due to treatment of character data vs. tree priors (May and Rothfels, 2023). Thus, to focus solely on the impact of alternative character evolution models, we applied a simple time-homogeneous, sampled ancestor implementation of the FBD process across all model configurations.

The relative fit of alternative model configurations were assessed using Bayes factors (BF) following the stepwise model selection process outlined by Wright et al. (2021) for paleontological datasets, a common practice in phylogenetic paleobiology. Calculating Bayes factors involves estimating and comparing marginal likelihoods for the set of competed models, where the marginal likelihood is the probability of the data integrated over a model's parameter

space. Marginal likelihood estimation was conducted using Stepping-Stone (SS) sampling in RevBayes (Xie et al., 2011; Höhna et al., 2016) following the algorithm and procedures for generating power posteriors using SS sampling described in Höhna et al. (2024). Model-fitting using SS sampling estimates marginal likelihoods by iteratively sampling parameter space from the prior to the posterior, which approximates the sum of the probability of the data across all possible values of parameter space. Once marginal likelihoods were obtained for each model configuration, Bayes factors were calculated and sequentially compared among each successively more complex pair of models. Interpretation of BFs follow guidelines of the natural log posterior probability scale in Kass and Raftery (1995).

In addition to comparing the relative fit among models, we also wanted to determine the extent to which different model configurations might lead to different phylogenetic hypotheses and/or macroevolutionary inferences. To compare the posterior distributions for tree topologies and macroevolutionary parameters inferred under different models, we conducted Markov-chain Monte Carlo (MCMC) simulation in RevBayes (Höhna et al., 2016) for each of the character model configurations (Figure 1). Macroevolutionary parameters of interest include FBD parameters for speciation rate, extinction rate, and sampling rate, as well as the number of sampled ancestors in each tree of the posterior distribution. To evaluate whether character models impact node ages and/or branch durations, we also calculated the minimum implied gap (MIG) (Wills, 1999), which is equal to the sum of the ghost ranges implied by a timescaled phylogeny when terminal stratigraphic ranges are excluded. To determine whether clade support values were influenced by character model configurations, we inferred a Maximum Clade Credibility (MCC) tree for each character model configuration and compared their posterior probabilities of clades. Finally, to compare the samples of trees inferred by different character model configurations, we mapped the posterior distribution of phylogenies inferred by each model configuration into a "treespace" (Hillis et al., 2005; Wright and Lloyd, 2020; Pates et al., 2022). Treespaces were constructed via multivariate ordination of pairwise dissimilarity matrices using Principal Coordinate Analysis (PCoA). Pairwise dissimilarity among trees was calculated using a variety of tree distance metrics, including the simple Robinson-Foulds (RF) distance, which measures the number of splits that differ between trees (Robinson and Foulds, 1981), as well as information theoretic methods (Smith, 2020) and quartet distances (Estabrook et al., 1985).

Convergence of SS sampling and MCMC results were evaluated using standard approaches to MCMC diagnostics, including visual inspection of trace plots, comparing independent runs, and checking estimated sample sizes. All treespace analyses were performed in R using original scripts and functions from the ape package (Paradis and Schliep, 2019), the RF.dist function from the *phangorn* package (Schliep, 2011), the ClusteringInfoDist function from the *TreeDist* package (Smith, 2022), the TQDist function from the *Quartet* package (Sand et al., 2014; Smith 2019), and the cmdscale function from the stats base R package. Additional information about

treespace construction, pairwise dissimilarity metrics, and custom R scripts to carry out treespace analyses are provided in the Supplementary Information.

3. Results

The log marginal likelihood values for character model configurations and BFs for each sequential model comparison are shown in Table 1. Bayes Factors among competed models generally indicate increased support for increasingly complex character models with additional layers of biological complexity. However, the explanatory data available limits this trend, as BFs naturally penalize more parameter-rich models (Kass and Raftery, 1995), and our results decisively indicate the most parameter-rich model is not supported compared to a simpler model configuration.

Among models analyzing only discrete traits, BFs indicate the **R+E** configuration was the best-fitting model, which accounted for multiple patterns of rate variation and partitioned characters into categories comprising "ecologic" vs. non-ecologic features of the trilobite body plan. The inclusion of continuous traits dramatically increased log marginal likelihood values compared to model configurations with only discrete traits. Among the two model configurations analyzing both discrete and continuous traits, BFs indicate the **RE+C** model was favored over **mREC**, and had the largest log marginal likelihood value and decisive BF support among all model configurations evaluated.

The posterior distributions of macroevolutionary and sampling parameters estimated under different character evolution models show considerable variation (Figure 2). For example, speciation and extinction rates were inferred to be considerably higher for rate-variable models, whereas fossil sampling rates are estimated to be lower on average compared to a constant rate model. The best-fitting character model, **RE+C**, has the lowest mean fossil sampling rate among models. According to MIG values, models incorporating continuous trait evolution imply longer durations of "ghost" lineages than other rate-variable models that sample only discrete characters, and on average contain fewer sampled ancestors per tree than models without continuous trait data (Figure 2). Maximum clade credibility (MCC) trees inferred across model configurations show substantial variation in posterior probability values (Figure 2). Since MCC trees are fully bifurcating, support values for clades corresponding to their N-1 internal nodes can be compared. Interestingly, the **S** model, which has the lowest marginal likelihood and BF support (Table 1), has the highest mean posterior probability per clade across the 37 internal nodes. However, we find no statistical differences in clade posterior probabilities recovered in MCC trees across model configurations (one-way anova: $F = 1.17$, $df = 4$, $p = 0.326$). The MCC trees inferred under each character model are presented in Supplemental Information.

Results of treespace visualization and analysis of tree-to-tree properties (e.g., tree balance) were strongly concordant with one another across a variety of alternative distance metrics (see Supplemental Information). For simplicity, we focus our main results on pairwise distances calculated using the simple RF metric. Scree plots, projections of tree space onto other principal coordinates, and comparisons with tree spaces based on other tree samples and other distance measures are reported in the Supplemental Information.

A visualization of treespace projected along the first two PCoA eigenvectors is shown in Figure 3. Although trees inferred under different character model configurations show some overlap in the two-dimensional space of PCoA axes 1 and 2, the treespace is highly structured with respect to the density of points and their locations along PCO1, which shows a striking correspondence with character model complexity (Figures 1 and 3). Moreover, the MCC trees inferred under different models are highly dispersed throughout the treespace, and clusters of trees form "islands" within treespace such that more complex model configurations are almost non-overlapping with the simplest character evolution model. On average, the posterior distributions of trees for model configurations including continuous characters contain trees that are less balanced, and the least complex model returned a larger proportion of balanced trees compared to other model configurations (Supplemental Information).

4. Discussion

(a) Reconstructing evolutionary history: a place where all the different kinds of truths fit together

In comparative biology, knowledge of evolutionary relationships among species makes predictions regarding patterns of character change (Felsenstein, 1985; Harmon, 2019; Soul and Wright, 2021). But in morphological systematics, especially phylogenetic paleobiology, character evolution models are themselves used to infer phylogenetic hypotheses—a seemingly catch-22 situation. As a further complication, we note even simple models of character evolution across a phylogeny, such as constant rate Brownian motion, can generate highly complex patterns of trait evolution when speciation and extinction rates vary over a clade's lifetime (Foote, 1996; O'Meara et al., 2006). However, Bayesian phylogenetic methods naturally present a solution to these apparent dilemmas by providing a framework "where all the different kinds of truths fit together" (see Vonnegut, 1959 for a literary use of this concept). In other words, phylogenetic inferences cannot be made in a vacuum. Indeed, reconstructing evolutionary relationships is just one of the necessary ingredients for reconstructing a clade's *evolutionary history* (i.e., tree topology, divergences, speciation and extinction rates, etc.). Using this framework, all data concerning phenotypes, fossil occurrences, and for clades with extant taxa, molecular sequences, jointly make contributions to estimating relevant evolutionary parameters and determining the probability distribution of a clade's evolutionary history (Wagner and Marcot, 2010).

In this study, we focused our attention on how our assumptions about morphologic evolution, reflected in character models and interpretation of their parameters, may impact our understanding of phylogenetic relationships and macroevolutionary inferences. All character evolution models are wrong. Some are useful. And some are more wrong (or more useful) than others. To determine a best-fitting model, we used a stepwise approach to model selection commonly used in phylogenetic paleobiology to statistically compare an increasingly complex series of character models (Wright et al., 2021). In our empirical example of a family of Cambrian–Ordovician trilobites, the best-fitting character evolution model is consistent with highly heterogeneous rates of morphologic evolution among characters, across the phylogeny, and between fundamentally different types of characters.

(b) Impacts of character partitions and inclusion of continuous trait data
There has been conflicting evidence in the literature concerning whether biologically-based character partitions impact phylogenetic results (Clarke and Middleton, 2008; Porto et al., 2021; Wright et al., 2021; Casali et al., 2022). Discussion of biologically-based partitions (e.g., based on anatomical region, ecology, developmental features, etc.) is distinct from character partitions based on the number of states (Mulvey et al., 2024), which is an important but separate issue concerning how morphologic information is coded into characters and character states (Khakurel et al., 2024), and may (or may not) involve unobserved states (Tarasov, 2019).

The majority of models we analyzed partitioned the discrete morphologic characters into ecological vs. non-ecological categories and assigned them independent lognormal distributions to account for rate variation among characters. Interestingly, trees inferred under the simplest model including character partitions (**R+E**) show substantial overlap in treespace with simpler models (**S, R**) (Figure 3). However, Bayes factors strongly and consistently supported models including ecology-based character partitions. Aside from how partitions may impact tree topology, we wondered if the data may be pointing to an evolutionary explanation for these results. After all, the characters in our partitions differ with respect to whether or not they have a presumed functional or ecologic significance, and basic comparative biology predicts such traits may be more labile than non-ecologic traits (Blomberg et al., 2003). Recall the mean of a rate distribution is set equal to one in order for relative rates to be interpretable (e.g., Yang, 1994; Revell and Harmon, 2024), so the shape of the distribution describing rate variation among characters is often determined by a single parameter describing its variance, which is estimated from data. In Figure 4, we show the marginal distributions estimated for the rate variance parameters estimated for ecologic vs. non-ecologic character partitions under the best-fitting model. Remarkably, the distribution of rate variance parameters differ significantly between partitions (Kolmogorov-Smirnov test, $D = 0.467$, $p < 0.001$), with values estimated for characters in the ecologic partition to be larger than and outside the range of estimated values for non-ecologic characters. This suggests ecologic traits in olenid trilobites may have evolved more

rapidly than their general body plan characters, which confirms biological intuition and perhaps explains why character partitions were supported by Bayes Factors.

Our results indicate that modeling continuous characters as quantitative traits dramatically improved the log marginal likelihood values (Table 1). Although this may in part reflect the increased information contained in the trait data, it is nevertheless consistent with our intuition that it is best to include as much character data and information as possible in phylogenetic analyses, provided that data reliably reflect the researcher's understanding of their study system. Further, our results may be seen to corroborate other studies highlighting use of continuous traits to estimate phylogenies and/or divergence times (Parins-Fukuchi, 2018a; Álvarez-Carretero et al., 2019). However, the inclusion of continuous traits impacted both the macroevolutionary parameters estimated and tree topologies inferred (Figures 2-3). On average, phylogenies estimated under models including continuous traits inferred fewer sampled ancestors per tree. And perhaps for related reasons, they also have larger MIG values, indicating longer, unsampled "ghost lineage" durations across trees. We note that similar differences in node age distributions and/or branch durations have been shown to have a significant impact on downstream comparative analyses of fossil data (Bapst, 2014; Bapst et al., 2016; Bapst and Hopkins, 2017; Soul and Wright, 2019), and may be at least as influential as the overall tree topology (Soul and Friedman, 2015). Because we are dealing with empirical data, we cannot say with certainty which model (or inference) reflects the "truth". Nevertheless, we caution researchers that decisions made about characters when constructing a phylogenetic character matrix may have far-reaching consequences beyond inferring relationships alone.

(c) Character models influence the probability of tree topologies and the exploration of treespace
In addition to their impacts on macroevolutionary inferences, the character models we analyzed had a striking influence on which regions of treespace were explored during MCMC analysis (Figure 3). Regardless of whether we randomly sample 300 trees or select the 300 trees with the highest probabilities from each model's posterior, the treespaces show similar visualizations and properties (Supplemental Information). These patterns are persistent across treespaces constructed using a variety of alternative metrics to calculate pairwise distances between trees (Supplemental Information), and we are unaware of any bias that could lead to our treespace ordination to show a correspondence between PCoA axes and character model complexity. Thus, a more complex character model configuration may not solely estimate more evolutionary parameters than a simpler model, it may also influence which tree topologies are considered more probable than others. As these are empirical data, we cannot know the "true" tree, but we emphasize that our results indicate that our assumptions about character evolution make a difference when it comes to estimating tree topologies. Again, these results highlight how phylogenetic inference cannot be made in a vacuum (Wagner and Marcot, 2010; Wright et al., 2016).

Although the patterns in our treespace visualizations were robust to choice of tree distance metrics (Supplemental Information), treespaces are complex and multidimensional. Ordination methods can cause distortion in the observed distances when projected in lower dimensional space, and it is possible that variation is spread across many axes (Hillis et al., 2005; Wright and Lloyd, 2020). To determine the fidelity of distances represented in our treespaces and assess the impact of dimension reduction, we selected two reference trees from the **S** and **mREC** models (i.e., simplest to most complex models) and compared the relationship between RF distances with Euclidean distance in the ordination using increasingly reduced sets of axes (Figure 5). In general, trees that are closer to the reference trees in Euclidean space have low RF distances, whereas trees that are further away from the reference trees have higher RF distances. We find a positive correlation between RF distances between trees and their Euclidean distances, even when only a few axes are evaluated (Figure 5). Further, similar positive correlations were found using other tree distance metrics and were not sensitive to whether treespaces were constructed using randomly selected trees or trees with the highest posterior probabilities (Supplemental Information).

To the extent that different character models influence tree topology probabilities such that they preferentially explore different regions of treespace, we strongly recommend researchers carefully consider how they model morphological evolution and engage with model-fitting procedures even if their primary interest is solely to infer the phylogenetic relationships of their favorite clade.

(d) Limitations, recommendations, and future directions
We evaluated only a subset of possible model configurations, and it is interesting to speculate how outcomes may vary if different decisions were made regarding the initial model choices. For example, one could have compared a model equivalent to the simplistic "strict" rates configuration (model **S**) for discrete characters, but also include the quantitative trait data and model them according to a Brownian motion process. Of course, it is neither computationally feasible nor desirable to estimate marginal likelihoods and/or compare BFs across all possible model configurations, so we chose a subset of five character models we believed may represent an alternative set of plausible evolutionary dynamics for our study system. Since it is impossible to evaluate all possible models, we recommend researchers choose a subset of models that enable them to test specific evolutionary hypotheses about their clade.

Although Bayesian model selection using SS sampling is a popular approach, it is not without several shortcomings. One possible concern of the stepwise model selection approach surrounds the order in which more complex models are evaluated, which may inadvertently bias which model is ultimately favored (Wright et al., 2021). A promising alternative to the stepwise approach to Bayesian model selection is reversible-jump MCMC (rjMCMC) (Green, 1995). In a

Bayesian context, rjMCMC places priors on competing models and treats them as random variables, where models are sampled by MCMC according to their posterior probabilities. This approach is theoretically equivalent to SS sampling in the sense that it computes identical Bayes Factors, at least when tree priors do not vary among the set of competed models (May and Rothfels, 2023). Reversible-jump MCMC has the advantage of accounting for uncertainty in model choice when estimating evolutionary parameters (i.e., tree topology, etc.), can accurately test alternative models involving variation in tree priors (May and Rothfels, 2023), and is far more computationally efficient than SS sampling, which requires many replicates. Nevertheless, it faces similar challenges to SS sampling regarding practical limits to the number of scientifically interesting models evaluated.

Regardless of which approach is used for model selection, the results of a model-fitting procedure only provide information about the relative fit of a model. It does not necessarily imply the model is adequately capturing the dynamics of the biological system being investigated. In fact, a best-fitting character model from a statistical standpoint of relative fit may poorly describe the underlying data. Determining whether a given model provides a reasonable explanation for data involves assessing a model's adequacy. A variety of approaches to assessing model adequacy, including posterior predictive simulation (Höhna et al., 2018; Mulvey et al., 2024), are available for evaluating phylogenetic models of character evolution. We recommend researchers explore these newly developed tools to carefully interpret the implications and limits of their data.

Rather than exhaustively evaluate all possible alternative character models, investigate their adequacy, or infer the one "true" tree of olenid trilobites, we have focused on the far more modest goal of assessing whether different model-based assumptions about character evolution impact phylogenetic and macroevolutionary inferences using a dataset representative of many empirical studies. This is an important practical question, as the computational and analytical demands of model comparison is time consuming and a potential barrier to researchers who would prefer spending their time using phylogenies to solve taxonomic problems, describe new taxa, or investigate evolutionary patterns in their favorite clade. As systematic (paleo)biologists ourselves, we fully empathize with this sentiment occasionally expressed by colleagues. However, our results suggest inferences of phylogenetic relationships are highly sensitive to our assumptions about characters and how they evolved, and are not generally independent of other aspects of reconstructing a clade's macroevolutionary history. We find that character evolution models have a significant impact on the probability of alternative tree topologies and influence which regions of treespace are explored. We recommend researchers think carefully about character evolution not only when interested in testing evolutionary hypotheses, but also when their primary research question solely revolves around inferring phylogenetic relationships.

Acknowledgements

We thank K. St. John and S.R. Cole for advice during the early stages of this project. D.F.W. thanks Kurt Vonneget for the concept of the chronosynclastic infundibulum, which inspired his perspectives about Bayesian approaches to simultaneous inference in phylogenetic paleobiology.

Funding

This research was supported in part by the National Science Foundation (EAR-GEO 1848145) to M.J.H. D.F.W. acknowledges funding by the American Museum of Natural History's Lerner Gray and Gerstner Scholar fund during the initial stages of this research, and was also supported by the National Museum of Natural History (Smithsonian Institution) and the Sam Noble Oklahoma Museum of Natural History.

References

Abadi, S., Azouri, D., Pupko, T. and Mayrose, I., 2019. Model selection may not be a mandatory step for phylogeny reconstruction. *Nature communications*, *10*(1), p.934.

Álvarez-Carretero, S., Goswami, A., Yang, Z. and Dos Reis, M., 2019. Bayesian estimation of species divergence times using correlated quantitative characters. *Systematic Biology*, *68*(6), pp.967-986.

Bapst, D.W., 2014. Assessing the effect of time-scaling methods on phylogeny-based analyses in the fossil record. *Paleobiology*, *40*(3), pp.331-351.

Bapst, D.W., Wright, A.M., Matzke, N.J. and Lloyd, G.T., 2016. Topology, divergence dates, and macroevolutionary inferences vary between different tip-dating approaches applied to fossil theropods (Dinosauria). *Biology Letters*, *12*(7), p.20160237.

Bapst, D.W. and Hopkins, M.J., 2017. Comparing cal3 and other a posteriori time-scaling approaches in a case study with the pterocephaliid trilobites. *Paleobiology*, *43*(1), pp.49-67.

Bapst, D.W., Schreiber, H.A. and Carlson, S.J., 2018. Combined analysis of extant Rhynchonellida (Brachiopoda) using morphological and molecular data. *Systematic Biology*, *67*(1), pp.32-48.

Barido-Sottani, J., Van Tiel, N.M., Hopkins, M.J., Wright, D.F., Stadler, T. and Warnock, R.C., 2020. Ignoring fossil age uncertainty leads to inaccurate topology and divergence time estimates in time calibrated tree inference. *Frontiers in Ecology and Evolution*, *8*, p.183.

Barido-Sottani, J., Pohle, A., De Baets, K., Murdock, D. and Warnock, R.C., 2023. Putting the F into FBD analysis: tree constraints or morphological data?. *Palaeontology*, *66*(6), p.e12679.

Blomberg, S.P., Garland Jr, T. and Ives, A.R., 2003. Testing for phylogenetic signal in comparative data: behavioral traits are more labile. *Evolution*, *57*(4), pp.717-745.

Caetano, D.S. and Harmon, L.J., 2019. Estimating correlated rates of trait evolution with uncertainty. *Systematic Biology*, *68*(3), pp.412-429.

Casali, D.M., Freitas, F.V. and Perini, F.A., 2023. Evaluating the impact of anatomical partitioning on summary topologies obtained with Bayesian phylogenetic analyses of morphological data. *Systematic Biology*, *72*(1), pp.62-77.

Clarke, J.A. and Middleton, K.M., 2008. Mosaicism, modules, and the evolution of birds: results from a Bayesian approach to the study of morphological evolution using discrete character data. *Systematic biology*, *57*(2), pp.185-201.

Drummond, A.J. and Stadler, T., 2016. Bayesian phylogenetic estimation of fossil ages. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *371*(1699), p.20150129.

Erwin, D.H., 2007. Disparity: morphological pattern and developmental context. *Palaeontology*, *50*(1), pp.57-73.

Estabrook, G.F., McMorris, F.R. and Meacham, C.A., 1985. Comparison of undirected phylogenetic trees based on subtrees of four evolutionary units. *Systematic Zoology*, *34*(2), pp.193-200.

Felsenstein, J., 1985. Phylogenies and the comparative method. *The American Naturalist*, *125*(1), pp.1-15.

Felsenstein, J., 1988. Phylogenies and quantitative characters. *Annual Review of Ecology and Systematics*, pp.445-471.

Foote, M., 1997. The evolution of morphological diversity. *Annual Review of Ecology and Systematics*, *28*(1), pp.129-152.

Foote, M.1996. Models of morphological diversification. Pp.62-86 in Jablonski, D., Erwin, D.H. and Lipps, J.H, eds., *Evolutionary paleobiology. University of Chicago Press, Chicago*, pp.62-86.

Fortey, R.A., 1974. A new pelagic trilobite from the Ordovician of Spitsbergen, Ireland and Utah. *Palaeontology*, *17*(1), pp.111-124.

Gavryushkina, A., Welch, D., Stadler, T. and Drummond, A.J., 2014. Bayesian inference of sampled ancestor trees for epidemiology and fossil calibration. *PLoS computational biology*, *10*(12), p.e1003919.

Gradstein, F.M. ed., 2012. *The geologic time scale 2012*. Elsevier.

Green, P.J., 1995. Reversible jump Markov chain Monte Carlo computation and Bayesian model determination. *Biometrika*, *82*(4), pp.711-732.

Harmon, L.J., Losos, J.B., Jonathan Davies, T., Gillespie, R.G., Gittleman, J.L., Bryan Jennings, W., Kozak, K.H., McPeek, M.A., Moreno-Roark, F., Near, T.J. and Purvis, A., 2010. Early bursts of body size and shape evolution are rare in comparative data. *Evolution*, *64*(8), pp.2385-2396.

Harmon, L.J., 2019. *Phylogenetic comparative methods*. Traverse City, MI, USA: Independent.

Harrison, L.B. and Larsson, H.C., 2015. Among-character rate variation distributions in phylogenetic analysis of discrete morphological characters. *Systematic Biology*, *64*(2), pp.307-324.

Heath, T.A., Huelsenbeck, J.P. and Stadler, T., 2014. The fossilized birth–death process for coherent calibration of divergence-time estimates. *Proceedings of the National Academy of Sciences*, *111*(29), pp.E2957-E2966.

Heled, J. and Drummond, A.J., 2012. Calibrated tree priors for relaxed phylogenetics and divergence time estimation. *Systematic biology*, *61*(1), pp.138-149.

Hillis, D.M., Heath, T.A. and John, K.S., 2005. Analysis and visualization of tree space. *Systematic biology*, *54*(3), pp.471-482.

Höhna, S., Landis, M.J., Heath, T.A., Boussau, B., Lartillot, N., Moore, B.R., Huelsenbeck, J.P. and Ronquist, F., 2016. RevBayes: Bayesian phylogenetic inference using graphical models and an interactive model-specification language. *Systematic biology*, *65*(4), pp.726-736.

Höhna, S., Coghill, L.M., Mount, G.G., Thomson, R.C. and Brown, J.M., 2018. P3: Phylogenetic posterior prediction in RevBayes. *Molecular biology and evolution*, *35*(4), pp.1028-1034.

Höhna, S., Landis, M.J., Heath, T.A., and Moore, B.R. 2024. General introduction to model selection: comparing relative model fit with Bayes Factors. *RevBayes*. https://revbayes.github.io/tutorials/model_selection_bayes_factors/bf_intro.html

Hopkins, M.J., 2019. Phylogenetic analysis and revision of the trilobite subfamily Balnibarbiinae (Olenidae). *American Museum Novitates*, *2019*(3928), pp.1-20.

Hopkins, M.J. and Lidgard, S., 2012. Evolutionary mode routinely varies among morphological traits within fossil species lineages. *Proceedings of the National Academy of Sciences*, *109*(50), pp.20520-20525.

Hopkins, M.J. and Smith, A.B., 2015. Dynamic evolutionary change in post-Paleozoic echinoids and the importance of scale when interpreting changes in rates of evolution. *Proceedings of the National Academy of Sciences*, *112*(12), pp.3758-3763.

Kass, R.E. and Raftery, A.E., 1995. Bayes factors. *Journal of the american statistical association*, *90*(430), pp.773-795.

Khakurel, B., Grigsby, C., Tran, T.D., Zariwala, J., Höhna, S. and Wright, A.M., 2024. The fundamental role of character coding in Bayesian morphological phylogenetics. *Systematic biology*, *73*(5), pp.861-871

King, B. and Beck, R.M., 2020. Tip dating supports novel resolutions of controversial relationships among early mammals. *Proceedings of the Royal Society B*, *287*(1928), p.20200943.

Lanfear, R., Frandsen, P.B., Wright, A.M., Senfeld, T. and Calcott, B., 2017. PartitionFinder 2: new methods for selecting partitioned models of evolution for molecular and morphological phylogenetic analyses. *Molecular biology and evolution*, *34*(3), pp.772-773.

Lewis, P.O., 2001. A likelihood approach to estimating phylogeny from discrete morphological character data. *Systematic biology*, *50*(6), pp.913-925.

Lloyd, G.T., Wang, S.C. and Brusatte, S.L., 2012. Identifying heterogeneity in rates of morphological evolution: discrete character change in the evolution of lungfish (Sarcopterygii; Dipnoi). *Evolution*, *66*(2), pp.330-348.

May, M.R. and Rothfels, C.J., 2023. Diversification models conflate likelihood and prior, and cannot be compared using conventional model-comparison tools. *Systematic Biology*, *72*(3), pp.713-722.

Mongiardino Koch, N., Garwood, R.J. and Parry, L.A., 2021. Fossils improve phylogenetic analyses of morphological characters. *Proceedings of the Royal Society B*, *288*(1950), p.20210044.

Monti, D.S. and Confalonieri, V.A., 2019. First cladistic analysis of the trilobite family Olenidae from the Furongian and Ordovician. *Lethaia*, *52*(3), pp.304-322.

Mulvey, L.P., May, M.R., Brown, J.M., Höhna, S., Wright, A.M. and Warnock, R.C., 2024. Assessing the Adequacy of Morphological Models using posterior predictive simulations. *Systematic Biology*, p.syae055.

O'meara, B.C., Ané, C., Sanderson, M.J. and Wainwright, P.C., 2006. Testing for different rates of continuous trait evolution using likelihood. *Evolution*, *60*(5), pp.922-933.

Paradis, E. and Schliep, K., 2019. ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics*, *35*(3), pp.526-528.

Parins-Fukuchi, C., 2018a. Use of continuous traits can improve morphological phylogenetics. *Systematic Biology*, *67*(2), pp.328-339.

Parins-Fukuchi, C., 2018b. Bayesian placement of fossils on phylogenies using quantitative morphometric data. *Evolution*, *72*(9), pp.1801-1814.

Paterson, J.R., Edgecombe, G.D. and Lee, M.S., 2019. Trilobite evolutionary rates constrain the duration of the Cambrian explosion. *Proceedings of the National Academy of Sciences*, *116*(10), pp.4394-4399.

Pates, S., Wolfe, J.M., Lerosey-Aubril, R., Daley, A.C. and Ortega-Hernández, J., 2022. New opabiniid diversifies the weirdest wonders of the euarthropod stem group. *Proceedings of the Royal Society B*, *289*(1968), p.20212093.

Porto, D.S., Almeida, E.A. and Pennell, M.W., 2021. Investigating morphological complexes using informational dissonance and Bayes factors: a case study in corbiculate bees. *Systematic Biology*, *70*(2), pp.295-306.

Rae, T.C., 1998. The logical basis for the use of continuous characters in phylogenetic systematics. *Cladistics*, *14*(3), pp.221-228.

Revell, L.J. and Harmon, L.J., 2024. A discrete character evolution model for phylogenetic comparative biology with Γ-distributed rate heterogeneity among branches of the tree. *bioRxiv*, pp.2024-05.

Robinson, D.F. and Foulds, L.R., 1981. Comparison of phylogenetic trees. *Mathematical biosciences*, *53*(1-2), pp.131-147.

Ronquist, F., Klopfstein, S., Vilhelmsen, L., Schulmeister, S., Murray, D.L. and Rasnitsyn, A.P., 2012. A total-evidence approach to dating with fossils, applied to the early radiation of the Hymenoptera. *Systematic biology*, *61*(6), pp.973-999.

Revell, L.J. and Harmon, L.J., 2022. *Phylogenetic comparative methods in R*. Princeton University Press.

Sand, A., Holt, M.K., Johansen, J., Brodal, G.S., Mailund, T. and Pedersen, C.N., 2014. tqDist: a library for computing the quartet and triplet distances between binary or general trees. *Bioinformatics*, *30*(14), pp.2079-2080.

Schliep, K.P., 2011. phangorn: phylogenetic analysis in R. *Bioinformatics*, *27*(4), pp.592-593.

Simpson, G.G., 1944. *Tempo and mode in evolution*. Columbia University Press.

Smith, A.B., 2009. *Systematics and the fossil record: documenting evolutionary patterns*. John Wiley & Sons.

Smith, M.R., 2019. Quartet: comparison of phylogenetic trees using quartet and bipartition measures. *Comprehensive R Archive Network*, *10*.

Smith, M.R., 2020. Information theoretic generalized Robinson–Foulds metrics for comparing phylogenetic trees. *Bioinformatics*, *36*(20), pp.5007-5013.

Smith, M.R., 2022. Robust analysis of phylogenetic tree space. *Systematic Biology*, *71*(5), pp.1255-1270.

Soul, L.C. and Friedman, M., 2015. Taxonomy and phylogeny can yield comparable results in comparative paleontological analyses. *Systematic Biology*, *64*(4), pp.608-620.

Soul, L.C. and Wright, D.F., 2021. *Phylogenetic comparative methods: a user's guide for paleontologists*. Cambridge University Press.

Stadler, T., 2010. Sampling-through-time in birth–death trees. *Journal of theoretical biology*, *267*(3), pp.396-404.

Stadler, T., Gavryushkina, A., Warnock, R.C., Drummond, A.J. and Heath, T.A., 2018. The fossilized birth-death model for the analysis of stratigraphic range data under different speciation modes. *Journal of theoretical biology*, *447*, pp.41-55.

Tarasov, S., 2019. Integration of anatomy ontologies and evo-devo using structured Markov models suggests a new framework for modeling discrete phenotypic traits. *Systematic biology*, *68*(5), pp.698-716.

Vonnegut, K., 1959. *The Sirens of Titan*. Dial Press Trade Paperback.

Wagner, P.J., 2012. Modelling rate distributions using character compatibility: implications for morphological evolution among fossil invertebrates. *Biology Letters*, *8*(1), pp.143-146.

Wagner, P.J., 2018. Early bursts of disparity and the reorganization of character integration. *Proceedings of the Royal Society B*, *285*(1891), p.20181604.

Wagner, P.J. and Marcot, J.D., 2010. Probabilistic phylogenetic inference in the fossil record: current and future applications. *The Paleontological Society Papers*, *16*, pp.189-211.

Warnock, R.C. and Wright, A.M., 2020. *Understanding the tripartite approach to Bayesian divergence time estimation*. Cambridge University Press.

Wiley, E.O. and Lieberman, B.S., 2011. *Phylogenetics: theory and practice of phylogenetic systematics*. John Wiley & Sons.

Wills, M.A., 1999. Congruence between phylogeny and stratigraphy: randomization tests and the gap excess ratio. *Systematic Biology*, *48*(3), pp.559-580.

Wright, A.M., 2019. A systematist's guide to estimating Bayesian phylogenies from morphological data. *Insect Systematics and Diversity*, *3*(3), p.2.

Wright, A.M. and Hillis, D.M., 2014. Bayesian analysis using a simple likelihood model outperforms parsimony for estimation of phylogeny from discrete morphological data. *PLoS One*, *9*(10), p.e109210.

Wright, A.M., Lloyd, G.T. and Hillis, D.M., 2016. Modeling character change heterogeneity in phylogenetic analyses of morphology through the use of priors. *Systematic Biology*, *65*(4), pp.602-611.

Wright, A.M. and Lloyd, G.T., 2020. Bayesian analyses in phylogenetic palaeontology: interpreting the posterior sample. *Palaeontology*, *63*(6), pp.997-1006.

Wright, A.M, Wagner, P.J. and Wright, D.F., 2021. *Testing character evolution models in phylogenetic paleobiology: a case study with Cambrian echinoderms*. Cambridge University Press.

Wright, D.F., 2017a. Bayesian estimation of fossil phylogenies and the evolution of early to middle Paleozoic crinoids (Echinodermata). *Journal of Paleontology*, *91*(4), pp.799-814.

Wright, D.F., 2017b. Phenotypic innovation and adaptive constraints in the evolutionary radiation of Palaeozoic crinoids. *Scientific reports*, *7*(1), p.13745.

Xie, W., Lewis, P.O., Fan, Y., Kuo, L. and Chen, M.H., 2011. Improving marginal likelihood estimation for Bayesian phylogenetic model selection. *Systematic biology*, *60*(2), pp.150-160.

Yang, Z., 1994. Maximum likelihood phylogenetic estimation from DNA sequences with variable rates over sites: approximate methods. *Journal of Molecular evolution*, *39*, pp.306-314.

Tables and Figures.

Table 1. Results of model selection comparison. The log Bayes factor scale and interpretation follows Kass and Raftery (1995).

Model configuration	Marginal likelihood	log Bayes factor	Evidence against H_0
S	-1827.485	-	-
R	-1812.795	29.38	Very strong
R+E	-1808.086	9.42	Strong
RE+C	**-1646.190**	**323.79**	**Very strong**
mREC	-1666.600	- 40.82	Favors RE+C model

Figure 1.

Figure 1. Character evolution model configurations. Model configurations correspond to a range of simple to increasingly complex and parameter-rich models of morphological evolution; (*a*) model **S**, strict morphological clock of discrete morphological data with equal rates of change among characters; (*b*) model **R**, allows for rate variation among characters and across phylogeny; (*c*) model **R+E**, builds on model **R** with traits identified as ecologic vs. non-ecologic characters assigned to different character partitions; (*d*) model **RE+C**, builds on **R+E** but with continuous characters evolving according to a single-rate Brownian motion process. (*e*) model **mREC**, modifies model **RE+C** such that continuous characters evolve according to a multi-rate Brownian motion process.

Figure 2.

Figure 2. Impact of character models on macroevolutionary and phylogenetic inferences. Except for clade support values (lower right), all boxplots show variation in parameter estimates across the posterior distribution resulting from MCMC analysis of each model configuration. Clade support indicates variation in tree-wide posterior probability of clades in the MCC tree. Abbreviations for model configurations as for Figure 1. MIG indicates the minimum implied gaps. The character evolution model configuration with the highest model support, **RE+C**, is indicated in **bold** with a gray background.

Figure 3.

Figure 3. Tree space of 300 randomly selected trees from each posterior distribution. The tree space is a PCoA of a distance matrix based on RF distances between all pairs of trees; shown here is a projection of tree space onto the first two principal coordinates (PCO 1 and PCO 2). Black shapes correspond to the MCC trees for each model configuration. See Figure 1 caption for model configuration abbreviations.

Figure 4.

Figure 4. Marginal distributions of the rate variance parameters estimated for ecologic vs. non-ecologic discrete character partitions under the best-fitting character model configuration.

Figure 5.

Figure 5.

Assessment of the fidelity of projected pairwise distances and calculated pairwise distances based on the RF metric. Upper panels: tree space color coded by pairwise RF distance from a single selected tree. Tree 1 (upper left) is a randomly selected tree from the **S** model configuration's posterior distribution. Tree 1505 is the MCC tree from the **mREC** model configuration's posterior distribution. Cooler colors indicate smaller RF distance; warmer colors indicate larger RF distances. Lower panels: Euclidean distances calculated from subsets of PCoA scores plotted against original RF distance.