Habitat-specificity in SAR11 is associated with a handful of genes under high selection

Sarah J. Tucker[1,2,3,4,5], Kelle C. Freel[1], A. Murat Eren[3,4,5,6,7,#,*], Michael S. Rappé[1,#,*]

[1]Hawai'i Institute of Marine Biology, School of Ocean and Earth Science and Technology, University of Hawai'i at Mānoa, Kāne'ohe, HI 96744, USA; [2]Marine Biology Graduate Program, University of Hawai'i at Mānoa, Honolulu, Hawai'i, United States; [3]Josephine Bay Paul Center for Comparative Molecular Biology and Evolution, Marine Biological Laboratory, Woods Hole, MA 02543, USA; [4]Helmholtz Institute for Functional Marine Biodiversity, 26129 Oldenburg, Germany; [5]Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, 27570 Bremerhaven, Germany; [6]Institute for Chemistry and Biology of the Marine Environment, University of Oldenburg, 26129 Oldenburg, Germany; [7]Max Planck Institute for Marine Microbiology, 28359 Bremen, Germany

[#]Co-senior Authors

*Corresponding Authors. Email: meren@hifmb.de, rappe@hawaii.edu

Running Title: Ecotype differentiation in SAR11

Abstract

The order *Pelagibacterales* (SAR11) is the most abundant group of heterotrophic bacteria in the global surface ocean, where individual sublineages likely play distinct roles in oceanic biogeochemical cycles. Yet, understanding the determinants of niche partitioning within SAR11 has been a formidable challenge due to the high genetic diversity within individual SAR11 sublineages and the limited availability of high-quality genomes from both cultivation and metagenomic reconstruction. Here, we take advantage of 71 new SAR11 genomes from strains we isolated from the tropical Pacific Ocean to evaluate the distribution of metabolic traits across the *Pelagibacteraceae,* a recently classified family within the order *Pelagibacterales* encompassing subgroups Ia and Ib. Our analyses of metagenomes generated from stations where the strains were isolated reveals distinct habitat preferences across SAR11 genera for coastal or offshore environments, and subtle but systematic differences in metabolic potential that support these observations. We also observe higher levels of selective forces acting on habitat-specific metabolic genes linked to SAR11 fitness and polyphyletic distributions of habitat preferences and metabolic traits across SAR11 subclades, suggesting that contrasting lifestyles have emerged across multiple lineages independently. Together, these insights reveal niche-partitioning within sympatric and parapatric populations of SAR11 and demonstrate that the immense genomic diversity of SAR11 bacteria naturally segregates into ecologically and genetically cohesive units, or ecotypes, that vary in spatial distributions in the tropical Pacific.

Main

Bacterial populations typically harbor extensive variation in gene content that can be difficult to delineate into discrete ecological or evolutionary units, despite the significant impact such differences may have on the biogeochemical roles, ecological interactions, and fitness of these microorganisms[1–3]. Comparative genomics has enabled the identification of gene content variation that underpins distinct ecologies and contributes to the process of niche diversification[4–6]. However, because these analyses most often use a collection of genomes isolated from diverse environments where genes have evolved under a range of selective pressures[7], associating variations in gene content with specific forces of selection and identifying crucial genes that likely serve as determinants of fitness under specific conditions remains a formidable challenge.

The bacterial order *Pelagibacterales* (SAR11) is one of the most abundant and diverse bacterial groups in the surface oceans[8–10], where even a single SAR11 population can exhibit substantial sequence diversity in the environment[11]. Defining the genomic variation underlying distinct spatial distributions of SAR11 populations and how gene content differs across environmental conditions is vital to further understanding ocean biogeochemical cycling, eco-evolutionary relationships, and the biology of a ubiquitous and dominant lineage in the surface oceans. This task remains difficult because assessing the loss or gain of ecologically-relevant genes or metabolic pathways in SAR11 necessitates a robust collection of high quality genome sequences that represent a spectrum of both ecological and genomic diversity. SAR11 cells are historically recalcitrant to laboratory cultivation, and as such only a small portion of the total diversity of naturally occurring SAR11 populations can be interrogated

via high quality genomes such as those from isolated strains[12]. Yet even with a small set of isolate genomes, SAR11 has become a key model to help elucidate fundamental processes of ecology and evolution such as genome streamlining[13], marine oligotrophy [14], co-evolution[15,16], drivers of genetic diversity and evolution within organisms of large population sizes[11,17], structure-aware investigations of microbial population genetics[18], and ocean biogeochemical cycling[19–21].

In our companion work[22], we address the gap in the available genomic resources with novel SAR11 isolates that originate from a geographically constrained environment within the tropical Pacific, and develop a phylogenomic backbone that unites evolution with ecology to understand this diverse clade. Freel *et al.*,[22] delineated four families within SAR11, including the family *Pelagibacteraceae* which encompasses the historical SAR11 subgroups Ia and Ib. Here, we illuminate metabolic diversity found within *Pelagibacteraceae* and identify the determinants of niche differentiation within co-occurring *Pelagibacteraceae* populations by combining 71 of the new isolates with 21 previously sequenced *Pelagibacteraceae* isolate genomes. By taking advantage of metagenomes from the prominent source of isolation, we apply an integrated 'omics analysis to unveil differences in *Pelagibacteraceae* metabolic features across coastal and offshore habitats and identify metabolic genes under high selective pressures.

Results

Ecology and evolution of *Pelagibacteraceae* reveals strong habitat-specificity

By analyzing 92 high quality genomes from isolates obtained predominantly from coastal Kāneʻohe Bay, Oʻahu, Hawaiʻi, and the adjacent offshore (**Fig. 1, Supplementary file 1a**), we reveal that the genomic diversity of cultured *Pelagibacteraceae* partitions into eleven discrete monophyletic clusters, seven of which include isolates from the Kāneʻohe Bay Time-series (KByT) (**Fig. 1, Extended Data Fig. 1a**). The 92 genomes belong to both previously-described and new clades, which we recently described as eleven genera[22] (**Fig. 1**, **Supplementary file 1a**). The within-genus genome-wide average nucleotide identity (gANI) differed among genera, with the lowest within-genus average gANI found in *Proprepelagibacter* (87.1±0.0%, mean ± sd) and the highest within-genus average gANI found in *Littoralipelagibacter* (95.8±0.0; **Supplementary file 1b**). The average of all within-genus genome-wide average nucleotide identities was 92.3±2.7%.

We investigated the biogeography of *Pelagibacteraceae* isolate genomes through read recruitment with metagenomes from KByT and Station ALOHA in the adjacent North Pacific Subtropical Gyre. This revealed that the seven genera within the *Pelagibacteraceae* that contained KByT isolate genomes were commonly detected within the KByT system, while the three genera not containing KByT isolates were rarely detected, if at all (**Extended Data Fig. 1b**). Congruent with k-means cluster analysis that grouped the biogeochemical parameters from the metagenomic samples into coastal and offshore clusters (**Extended Data Fig. 3, Supplementary file 1c, Supplementary Information Note 1**), the patterns of genome distribution revealed a clear dichotomy where each genus was either highly present in the offshore or in the coastal environment, but not both (**Fig. 1c, Supplementary file 1d**). Genera with higher abundances in the coastal environment, hereafter referred to as 'coastal genera', did

not occur within a single monophyletic clade and instead appeared to have evolved multiple times across the evolutionary history of SAR11 (**Fig. 1c**).

For the most part, the patterns of distribution were consistent for all genomes within the same genus (**Extended Data Fig. 3**), and genomes that belonged to the same genus generally recruited similar proportions of reads from the environment. One exception was the genome HIMB1412 from the genus *Coralipelagibacter* which had the highest relative abundance among all *Pelagibacteraceae* genomes in the coastal environment (19.9±9.4%, mean ± sd, **Extended Data Fig. 3**). While we consistently detected all isolate genomes of *Xanthipelagibacter* in the offshore environment, three of eight genomes (HIMB83, FZCC0015, HIMB1456) were also frequently detected in the coastal environment (**Extended Data Fig. 3**). Regardless, their relative abundances were low in coastal samples (0.4±0.5%, mean ± sd) and sharply increased offshore (3.5±2.0%, mean ± sd; **Extended Data Fig. 3**), further supporting the characterization of *Xanthipelagibacter* as an offshore genus.

Overall, our metagenomic read recruitment reveals that *Pelagibacteraceae* communities partition between coastal and offshore habitats in the tropical Pacific, that these distinct communities are driven by differences in the distribution of individual *Pelagibacteraceae* genera, and that coastal genera (*Ampluspelagibacter*, *Coralipelagibacter*, *Littoralipelagibacter*) are distributed polyphyletically across the *Pelagibacteraceae*.

Fig. 1. *Pelagibacteraceae* **show polyphyletic habitat preferences for coastal or offshore environments across KByT. a**) The location of the Kāneʻohe Bay Time-series (KByT) sampling stations and the Hawaii Ocean Time-series at Station ALOHA. KByT spans a steep biogeochemical gradient, as noted by the dramatic increase in *Prochlorococcus* cellular abundances immediately offshore. **b**) Eleven genera are represented by genome-sequenced *Pelagibacteraceae* isolates, with the majority isolated from coastal Kāneʻohe Bay and the adjacent offshore. Nodes with circles represent ≥90% bootstrap support. Historically referenced clade names for *Pelagibacteraceae* genera are provided in parentheses. **c**) The relative read recruitment of *Pelagibacteraceae* isolate genomes summed at the level of genera for metagenomes from KByT and Station ALOHA. Metagenomes grouped into two clusters by environmental parameters, coastal and offshore, coinciding with the gradient in *Prochlorococcus* cellular abundances. *Pelagibacteraceae* genera showed distinct distribution patterns in the coastal and offshore waters that are distributed polyphyletically. Read recruitment <1% not shown.

Pelagibacteraceae pangenome includes differentially enriched genes and functions within coastal and offshore genera

Next, we sought to characterize the genomic diversity and functional gene content of

Pelagibacteraceae isolates in order to investigate the potential determinants of fitness that may

explain the distinct ecological patterns of distribution. We utilized a pangenomic approach that

partitioned all genes across all *Pelagibacteraceae* genomes into *de novo* gene families, or 'gene

clusters', based on amino acid sequence homology. This analysis resulted in 8,242 gene clusters

across all 92 genomes, of which 784 were core among all *Pelagibacteraceae* and made up the

majority (52-63%) of gene content found in any individual genome (**Fig. 2a; Supplementary**

file 2a). Half of the pangenome was composed of gene clusters that were singletons (4,201 of

8,242; **Fig. 2a**). Between the extremes of gene clusters that occur in every genome and those that

occur only in one, we found 206 accessory gene clusters that were 'genus-specific' (21±27,

mean±sd per genus; **Fig. 2a**; **Supplementary file 2b**) as they were present among all

representatives of a single genus, but absent from genomes belonging to any other genus. The

206 genus-specific gene clusters were not distributed uniformly across genera, as 95 were unique

to the genus *Superopelagibacter* alone (**Supplementary file 2b**). However, this number is likely an overestimation of the true *Superopelagibacter* genus-specific traits since this particular genus was represented by only two genomes, and the number of genus-specific gene-clusters negatively correlated with the number of genomes in a genus (r^2adj = 0.37, F=6.29, p = 0.037; Pearson's correlation). Gene clusters that were core to multiple genera (e.g., present in every genome for two or more genera, but absent in every other) were also not a dominant feature in the pangenome (Fig. 2a). They only represented 182 gene clusters, and were generally uniformly distributed across many different groupings of genera (**Fig. 2a**; **Supplementary file 2b**). We could further assign a subset of the gene clusters to coastal or offshore categories based on the environmental preference of the genomes in which they occurred, finding 32 and 125 gene clusters that were specific to coastal genera and offshore genera, respectively. (**Fig. 2a, Supplementary file 2b**).

We next focused on genes with well-characterized roles in known metabolisms by examining those that share orthology with genes described in the Kyoto Encyclopedia of Genes and Genomes (KEGG)[23] through KOfam models[24] and generating a 'functional pangenome' (**Fig. 2b**). The *Pelagibacteraceae* functional pangenome grouped genes based on functional identity rather than amino acid sequence similarity and consisted of 1,276 KOfam functions, with a large core (612 KOfams; **Fig. 2b**). Many of the remaining non-core functions were still shared widely across all eleven genera with a relatively uniform distribution (n=136; **Fig. 2b**) or were singletons (n=70). Similar to the conventional pangenome, the functional pangenome contained very few genus-specific functions (14 total among all genera; 0-5 KOfams per genus; **Fig. 2b**) and a limited number of functions that were core to multiple genera (e.g. present in every genome for two or more genera, but absent in every other; 32 KOfams total; **Fig. 2b;**

Supplementary file 2c). Only three KOfam functions were specific to coastal genera, while 13

KOfam functions were specific to offshore genera (**Fig. 2a; Supplementary file 2c**).

Pangenomic analysis using conventional and functional approaches revealed a large

overlap in gene content among ecologically distinct genera, with subtle but consistent

genus-specific features. The high overlap in gene content across the *Pelagibacteraceae* and small

number genus-specific gene clusters suggests differentiation between genera of the

Pelagibacteraceae may be determined by only a small portion of the genome. The polyphyletic

distribution of habitat preference (**Fig. 1**) supports that genes and functions that are linked to

success in these different habitats likely evolved multiple times through independent processes.

Given this evolutionary trajectory, it is highly plausible that instead of vertical inheritance,

horizontal gene transfer led to the acquisition of most of these habitat-specific genes and

subsequent high rates of recombination and strong selective forces resulted in the increased

prevalence among *Pelagibacteraceae* in a given ecological niche. This process, referred to as

gene-specific sweeps, frequently results in a more subtle display of habitat-specific traits[25] as is

observed here. To further examine the potential importance of these environmentally-distinct

functions in providing advantages to *Pelagibacteraceae* prevalent in coastal or offshore habitats,

we next focused on the subset of the environmentally-distinct functions characterized here that

are involved in nutrient resource utilization and cellular stress. Furthermore, in examining the

functions involved in nutrient resource utilization and cellular stress, we broadened our

investigation beyond the functions that were strictly found in coastal genera or strictly found in

offshore genera (e.g. environmentally-distinct functions), to also include functions that were

enriched in either coastal or offshore genera.

a) Conventional pangenome
8,242 gene clusters

Gene clusters specific to multiple genera n=182

Core gene clusters n=784

Gene clusters found in all 11 genera n=127

Genus-specific gene clusters n=206

Singletons n=4201

Environmentally-distinct gene clusters/ functions specific to **offshore genera** or **coastal genera**

Genera
Proprepelagibacter
Pelagibacter
Atlantikopelagibacter
Superopelagibacter
Xanthipelagibacter
Semelpelagibacter
Lacunapelagibacter
Tantilluspelagibacter
Littoralipelagibacter
Coralipelagibacter
Ampluspelagibacter

b)

Rare genera   Offshore genera   Coastal genera

Genus-specific functions n=14

Functions specific to multiple genera n=32

Functions found in all 11 genera n=136

Core functions n=612

Functional pangenome
1,276 KOfam functions

Fig 2. *Pelagibacteraceae* genomes share high overlap in genes and functions among ecologically-distinct groups. The presence and absence of gene clusters and KOfams are shown across **a)** a conventional pangenome and **b)** a functional pagenome, respectively. Genomes (inner rings) are grouped by genus and further grouped by ecological distribution in the tropical Pacific as coastal, offshore, or rare. The distribution of the gene pool and functions in the second most outer layer (e.g., core, genus-specific) show that the gene pool and functions are shared

extensively across all 11 genera and 92 genomes. The outer layer shows the sparse environmentally-distinct portions of the gene pool and functional pool that are specific to coastal or offshore genera.

Potential determinants of habitat-specificity in *Pelagibacteraceae* include molybdenum utilization and nitrogen and carbon metabolisms

Despite a high amount of water exchange, coastal Kāneʻohe Bay and the adjacent

offshore vary in nutrient availability, osmotic conditions (e.g. salinity, temperature, pH), and

phytoplankton communities that determine the organic carbon pool[26,27], suggesting that this

would enforce distinct environmental pressures on coastal and offshore *Pelagibacteraceae*

genera. To characterize ecologically-relevant genes that may support coastal and offshore

habitat-preferences of *Pelagibacteraceae* across the KByT system, we used an enrichment

analysis to examine metabolic traits related to nutrient acquisition and cellular stress that were

differentially distributed across genomes from either coastal or offshore genera. Genes that were

core across genera of the *Pelagibacteraceae* or did not show easily distinguishable distributions

(e.g. driven by phylogeny or environment) are further discussed in **Supplementary Information**

Note 2. Unless noted otherwise, the following metabolic traits were found outside of

hyper-variable regions of genomes and thus represent genes that are encoded within the

relatively stable genomic backbones of a given genus or set of genera.

Molybdenum enzyme utilization.

Molybdenum enzymes are nearly ubiquitous among organisms from all domains of life,

catalyze important oxidation-reduction reactions involved in carbon, sulfur, and nitrogen

metabolisms, and require a cofactor scaffold to hold the molybdenum in place[28]. While most *Pelagibacteraceae* genomes contained genes involved in molybdenum cofactor biosynthesis as well as molybdenum cofactor-dependent enzymes (**Fig. 3a, Supplementary file 2d**), these genes were missing from two offshore genera, *Lacunapelagibacter* and *Tantilluspelagibacter*. In contrast to the lack of molybdenum enzymes found in the offshore *Lacunapelagibacter* and *Tantilluspelagibacter*, other offshore genera, *Xanthipelagibacter* and *Semelpelagibacter*, appeared to increase reliance on molybdenum-dependent enzymes by uniquely harboring genes encoding molybdenum-dependent enzymes associated with the oxidation of purines (xanthine; *xdhAB*) (**Fig. 3a, Supplementary file 2d, Supplementary Note 3**). The distribution of molybdenum enzymes are polyphyletically distributed among offshore genera, where some offshore groups possess no molybdenum enzymes while others harbor all the molybdenum enzymes detected within *Pelagibacteraceae* (**Fig. 3a**). We hypothesize that the absence of molybdenum enzymes within the offshore *Lacunapelagibacter* and *Tantilluspelagibacter*, which have some of the smallest genomes within *Pelagibacteraceae* (~1.2Mbp; **Supplemental file 1a**), is likely driven by streamlining selection to minimize metabolic requirements and genome size[13,29].

Fig. 3. *Pelagibacteraceae* **genera fill distinct niches in molybdenum enzyme utilization. a)**
The distribution of molybdenum cofactor (MoCo) biosynthesis genes and genes involved in
metabolisms that require a molybdenum cofactor across *Pelagibacteraceae* genera. Metabolisms
within *Pelagibacteraceae* that require a molybdenum cofactor included those involved in the
repair of methionine sulfoxides and the catalytic subunit of periplasmic dimethylsulfoxide

(DMSO; *msrPQ*), the detoxification of 6-N-hydroxylaminopurine (HAP) to adenine (*yiiM*,
ycbX), the oxidation of formate to carbon dioxide (*fdhFD*), the oxidation of carbon monoxide
(*coxSMLG*), and the oxidation of purines (*xdhAB*). Genes that are predicted to represent proteins
with molybdenum cofactor binding sites are noted in bold. **b)** Genes encoding for purine
catabolism and transport are only present in some offshore genera. **c)** Genes encoding for
allantonate metabolism and purine salvage are present only in coastal genera. MoCo:
molybdenum cofactor, DMSO: dimethylsulfoxide, HAP: hydroxylaminopurine, CO: carbon
monoxide, NH_3: ammonia, dR-Pur-5P: purine 2'-deoxyribonucleoside 5'-monophosphate, Pur:
purine base, dRib5P: 2'-deoxyribonucleoside 5'-monophosphate, (d)R-Pur-1P:
beta-(deoxy)ribonucleosides.
262

Nitrogen metabolisms.

Purines are abundant in aquatic environments[30], contain a high number of nitrogen atoms

per molecule[31], and likely contribute to niche-differentiation between coastal and offshore

SAR11[32]. Our metabolic reconstructions show that offshore genera *Xanthipelagibacter* and

Semelpelagibacter harbored genes to catabolize purines through the utilization of a molybdenum

cofactor-dependent xanthine dehydrogenase (**Fig. 3b, Supplementary file 2e**), as well as genes

with functions predicted to support the efficient transport of nucleotides (*Xanthipelagibacter*:

nupNOPQ system; *Xanthipelagibacter* and *Semelpelagibacter*: xanthine/uracil permease *uraA*).

To make hypoxanthine available for degradation, *Semelpelagibacter* can convert adenine to

hypoxanthine using adenine deaminase (*ade*) and both genera *Semelpelagibacter* and

Xanthipelagibacter also uniquely possess a tRNA(adenine34) deaminase (*tadA*), which can

convert adenine to hypoxanthine (**Fig. 3b**). Most *Xanthipelagibacter* genomes (6 of 8) also

harbor a urease system (*ureABCDEFG*) to liberate ammonia in the final steps of the xanthine

degradation pathway.

In contrast to the purine degradation pathways found within the offshore genera, we

found coastal genera specialize on the degradation of allatonate, a bi-product of purine

metabolism, as well as the recycling of nucleotides. Members of the coastal genus

Ampluspelagibacter (and a single *Coralipelagibacter* genome, HIMB5) have the functional

capacity to degrade allantoate to (S)-ureiodoglycine and ammonia via an allantoate deiminase

and ureidoglycine to ureidoglycolate and ammonia, via a nucleophile (Ntn)-hydrolase (**Fig. 3c,**

Supplementary file 2e). It is important to note that the allantoin degradation genes found in

coastal genomes characterized here appear to provide a source of nitrogen, and differ from those

found in the offshore genera. All coastal genera possess a deoxynucleoside 5-monophosphate

N-glycosidase (*rcl*) that putatively breaks the N-glycosidic bond of purine nucleotides (i.e.

purine 2'-deoxyribonucleoside 5'-monophosphates; **Fig. 3c**). This differs from the purine

nucleoside salvage pathway core to both coastal and offshore genera, which requires

orthophosphate to cleave the N-glycosidic bond of beta-(deoxy)ribonucleosides to yield

alpha-(deoxy)ribose 1-phosphate (**Supplementary file 2e**). All *Pelagibacteraceae* are missing

genes to utilize deoxyribose-5-phosphate or deoxyribose-1-phosphate via a

deoxyribose-phosphate aldolase (*deoC*). However, ribose-1-phosphate could be catabolized to

D-fructose-6P and D-glyceraldehyde-3P (**Supplementary file 2e**).

In addition to differences in purine metabolism, offshore genera uniquely harbor a

transaminase (*agxt*) that may help these cells to overcome an auxotrophy for the biosynthesis of

the amino acid glycine (**Fig. 4, Supplementary file 2f**), which is a common feature across

Pelagibacteraceae, but likely restricts important biosynthetic pathways including protein

synthesis and central carbon metabolism[33]. Offshore genera also uniquely contain genes involved

in glutamate deamination, glutathione and carnitine metabolism, and the synthesis of

glucosylglycerate, which is used as an alternative osmolyte under nitrogen-limited conditions in

cyanobacteria[34]. In contrast, the coastal genera contain nitrogen metabolisms generally involved

in the deamination of various amino acids (e.g. hydroxyprolines, proline, ornithine, and

threonine). A subset of coastal *Ampluspelagibacter* genomes also have the potential to metabolize the polyamines putrescine and spermidine via *puuBCD* (**Fig. 4**). The puuD gene was located in hypervariable regions within coastal *Ampluspelagibacter* genomes.

Phosphate & Metals.

Some offshore *Pelagibacteraceae* genera exhibit a reduced genomic capacity to transport phosphate relative to coastal genera. High-affinity phosphate acquisition genes (*pstSCAB-phoU*; *phoB*; *phoR*) were core among coastal genera but, while present in some genomes, were not core among the offshore genera. Offshore genomes lacking this system appear to rely solely on the *pitA* gene, a presumably lower-affinity proton-motive force permease of phosphate or sulfur (**Fig. 4, Supplementary file 2f**). Some offshore genera contain an increased genomic capacity to transport or capture iron relative to coastal genera. While all *Pelagibacteraceae* contain an iron (III) transport system (*afuAB*), the offshore *Semelpelagibacter* genus alone possesses a high-affinity iron permease (*ftr1*) (**Fig. 4, Supplementary file 2f)**. Offshore genera *Superopelagibacter*, *Lacunapelagibacter*, and *Tantilluspelagibacter* also contain a heme oxidase, which could catalyze the oxidative degradation of the heme porphyrin ring to release iron. Unique genes to utilize copper, potassium, and nickel were found among the coastal genera (**Fig. 4**). We observed a copper-containing plastocyanin gene within a small subset of coastal genomes (**Fig. 4; Supplementary file 2f**). Plastocyanins serve as an electron carrier from cytochrome *f* to photosystem I and are generally found within eukaryotic phytoplankton and cyanobacteria[35]. Future studies are needed to understand the functional role of plastocyanins in *Pelagibacteraceae*, although one potential use could be the storage of copper to cope with periods of reduced availability[36].

Cellular stress.

Differences in the capacities to respond to osmotic and oxidative stresses were found among offshore and coastal *Pelagibacteraceae* genera. Most offshore genera shared a catalase-peroxidase (*katG*) that may help to cope with oxidative stress and has also been suggested to play a role in the co-evolution of marine microbial communities[15] (**Fig. 4, Supplementary file 2f**). Coastal genera contained two genes involved in osmotic stress that were not prevalent among offshore genomes: a voltage-gated sodium channel (*vgsc*) and a cation:H+antiporter (*yrbG*) (**Fig. 4, Supplementary file 2f**). Voltage-gated sodium channels utilize a structurally complex gating system to physically open and close the pore in response to changes in fluid shear or membrane stretch from dehydration[37,38], and may be particularly useful in coastal Kāneʻohe Bay where spatiotemporal shifts in salinity are common[39].

Carbon metabolisms.

Carbon metabolisms unique to coastal and offshore genera were remarkably distinct. SAR11's specialization on low-molecular weight, labile carbon sources has likely facilitated the success of this marine oligotroph in open oceans where carbon substrates are limited in availability and under high competition[40–42]. Consistent with these observations, the carbon metabolisms shared by offshore genera did not rely on glycolysis and involved low-molecular-weight compounds (one-carbon to three-carbon; C1-C3) that are common molecules and/or typical waste products in the marine environment. This includes the genetic potential to metabolize organic acids such as D-lactate, oxalate, haloacetate, and formate, as well as diverse aldehydes (**Fig. 4**, **Supplementary file 2f**). In contrast, coastal genera contain gene content to specialize in the degradation of relatively larger carbon compounds (C4-C6), including

the sugars D-xylulose and D-ribulose via *xylB* and sugar alcohols arabinitol and mannitol via an

arabinitol dehydrogenase, both of which subsequently enter glycolysis through the

non-phosphorylative Entner–Doudoroff (np-ED) pathway. Some coastal groups also have the

capacity to metabolize the sugar acids D-tartrate and malate via *tarD* (**Fig. 4; Supplementary**

file 2f). Genomes belonging to the coastal genus *Ampluspelagibacter*, which have the largest

genomes among known *Pelagibacteraceae*, harbored gene content potentially involved in the

transport and metabolism of glycerophospholipids and sn-glycerol-3-phosphate (*glpA*, *upgEBQ*,

gpr), the capacity to metabolize L-fucose, D-arabinose, and L-xylose to pyruvate through a

L-fucono-1,5-lactonase (*fucB*) and the glycolytic Entner-Doudoroff (ED) glycolytic pathway, and

a fructose transport system (*frcABC*) (**Fig.4; Supplementary file 2f**). The ED glycolytic

pathway was core among *Ampluspelagibacter*, *Xanthipelagibacter*, and *Proprepelagibacter*

genomes and variable across genomes in other genera (**Supplementary file 2f**). The increased

prevalence of sugar metabolisms among coastal genera, especially the *Ampluspelagibacter*

genus, suggests broadened metabolic versatility among coastal *Pelagibacteraceae* in response to

a diverse and/or more readily available set of organic carbons found within coastal Kāneʻohe

Bay relative to offshore waters. The distinct distributions of carbon metabolisms resolved here

also suggests fine-scale niche partitioning along the sugar-acid spectrum within

Pelagibacteraceae[43].

Fig. 4. Metabolic determinants of habitat-specificity in *Pelagibacteraceae*. Metabolic genes associated with nutrient acquisition and cellular stress vary among coastal and offshore *Pelagibacteraceae* genera.

Adhesion.

Finally, while investigating the functional differences between coastal and offshore genera, we observed homologs of a large gene (~5,000 to 9,500 bp) annotated as an autotransporter adhesin that was almost always present among coastal genomes and absent in all offshore genomes (**Fig. 4**). Autotransporters provide a simple and relatively minimal mechanism for the delivery of a passenger protein to the surface of Gram-negative bacteria[44], and can vary greatly in size due to changes in the number of repeating sequences in the passenger protein[45]. Because repeating regions cause difficulty for short-read assemblers such as those used in our genome assemblies[22], we further established confidence in the assembly of this gene in our coastal genomes by successfully finding homologs in long-read sequencing libraries from the coastal Kāneʻohe Bay environment and comparing protein structure models of genes annotated from the long-read data to a known autotransporter (**Extended Data Fig. 4; Supplementary Information Note 4**). We also examined the genome context of the gene, which was always positioned next to genes involved in type IV pilus systems and/or type II secretory systems and sometimes, but not always, within hyper-variable regions (**Extended Data Fig. 4, Supplementary Information Note 4, Supplementary file 2f**).

Adhesion to surfaces has not been observed in *Pelagibacteraceae* in culture or through environmental studies, although long filaments believed to be pili were observed within dividing cells of *Pelagibacteraceae*[46] and are associated with multiple processes, including adhesion to surfaces, transformation competence, and DNA uptake[47,48]. While speculative, it is possible that

adhesion by the autotransporter adhesin gene may be involved in the uptake of nutrients also in *Pelagibacteraceae*[49]. The coastal distribution of the autotransporter gene suggests that the advantage of having this large of a gene is worth the trade-off of maintenance in more nutrient-rich coastal environments, but potentially not in nutrient-limited offshore environments.

In summary, our investigation of the distribution of genes related to nutrient acquisition and cellular stress that are unique to coastal and offshore genera reveals numerous subtle differences in metabolic traits driven by the presence or absence of single genes (e.g. high affinity iron transporters, voltage-gated sodium channel genes), but few differences in multi-gene metabolic pathways (e.g. purine metabolism, molybdenum cofactor biosynthesis, fucose metabolism). Differences in genes related to nutrient acquisition were predominantly related to organic carbon and nitrogen metabolisms, and help to further define the unique and diverse roles *Pelagibacteraceae* play in oceanic biogeochemical cycles. The distribution of these metabolic traits generally do not follow phylogenetic distributions, but environmental patterns, suggesting that these metabolisms may be driven by selective pressures from the environment.

Gene determinants of habitat-specificity are under relatively higher selective pressures compared to non-diagnostic genes

Our analyses indicated a relatively small number of functions in the *Pelagibacteraceae* appear to be associated with its strict niche partitioning. Assuming that the requirement to perform optimally may be higher for genes that determine fitness to particular lifestyles, we hypothesized that metabolic traits that have distinct environmental distributions should be under higher selective pressures compared to other genes in the environment. To test this hypothesis we examined the proportion of non-synonymous to synonymous (pN/pS) sites per gene across

Pelagibacteraceae genomes using four deeply sequenced metagenomic samples from the KByT

system.

To define the set of genomes to be included in this analysis, we first evaluated whether

the gene-level selective pressures were uniform across genomes from the same genus. We

examined variation in pN/pS across genes with shared KOfam assignments for genomes

belonging to the genus *Coralipelagibacter,* which has a low minimum gANI (84.5%, n=6) and is

abundant in the tropical Pacific. Gene-to-gene variation explained 82% of the variation, while

sample-to-sample variation and genome-to-genome variation only explained 0.26 and 0.09%,

respectively (ANOVA, **Supplemental file 3a**). The little genome-to-genome variation showed

that genomes within the same genus likely experience similar patterns of selection on shared

genes, and supported using a single genome as a representative per genus. Thus, our downstream

analyses utilized a single genome for each genus in *Pelagibacteraceae*, except

Atlantikopelagibacter, *Pelagibacter*, *Proprepelagibacter*, and *Semelpelagibacter*, which lacked

sufficient coverage in the KByT system (**Extended Data Fig. 1**, **Supplemental file 3b**).

Across the *Pelagibacteraceae* genomes examined, strong purifying selection (pN/pS ≪ 1)

was a general characteristic of most genes as observed previously by Kiefl et al.,[18] where over

75% of genes in each genome representative had a pN/pS value of ≤ 0.1 (**Supplemental file 3c**).

However, not all genes experienced similar magnitudes of purifying selection, with gene-to-gene

variation in pN/pS explaining between 88-99% of the variation in pN/pS across genomes (n=7,

ANOVA, **Supplemental file 3d**). Sample-to-sample variation in pN/pS values varied for most

genomes, however it explained a very small proportion of the overall variation (<0.5%, n=7,

ANOVA). Gene coverage showed no correlation between per gene values of pN/pS (absolute

value of r=0.02-0.28; Pearson correlation coefficient; **Extended Data Fig. 5**). Gene coverage

explained an extremely small portion of the overall pN/pS variation (<0.07%, n=7, ANOVA,
Supplemental file 3d), which indicates that pN/pS values are unlikely to be driven by artifacts
associated with variation in coverage.

We next evaluated the selective pressures on genes that were differentially distributed
across coastal and offshore *Pelagibacteraceae* genera (**Fig. 5; Extended Data Fig. 6**). The
differentially distributed genes with the highest selective pressures in offshore genera included a
malonyl-CoA decarboxylase (*mlycd;* pN/pS= 0.019±0.010, mean ± sd), an aldehyde
dehydrogenase (*aldB*; pN/pS= 0.033±0.018), a formate dehydrogenase
(*fdh;*pN/pS=0.0127±0.001), a D-lactate dehydrogenase (*ldhD*; pN/pS=0.018±0.003) and an
alanine-glyoxylate / serine-glyoxylate / serine-pyruvate transaminase (*agxt*; pN/pS=
0.022±0.008) (**Fig. 5**). These genes are generally involved in the metabolism of small and
common carbon substrates and the capacity to produce glycine, which is an auxotrophy in other
Pelagibacteraceae. Within *Xanthipelagibacter*, genes involved in purine uptake (*uraA*,
pN/pS=0.040±0.001; *nupNOPQ*, pN/pS=0.045±0.010) experienced higher purifying selection
than purine degradation pathways (*xdhAB*, pN/pS=0.084±0.028), which is in line with previous
observations that high-affinity transport is key to *Pelagibacteraceae*'s success as an
oligotroph[50,51]. The low pN/pS values for these distinct carbon and nitrogen related genes support
the hypothesis that these metabolisms provide a fitness advantage in offshore waters.

Purifying selection

Strong   Weak

Genes ranked across genomes by pN/pS

lowest pN/pS   highest pN/pS

differentially distributed metabolic genes

Ampluspelagibacter
HIMB1556
0.001 0.025   0.050   0.075 0.10   0.25

frcA, frcC, vgsc   frcB allC rcl

Xanthipelagibacter
HIMB83
0.001 0.025   0.050   0.075 0.10   0.25

nupP, nupQ, uraA nupN   nupO

Lacunapelagibacter
HIMB1437
0.001 0.025   0.050   0.075 0.10   0.25

fdh, mylcd, aldB, ldhD

Tantilluspelagibacter
RS39
0.001 0.025   0.050   0.075 0.10   0.25

mylcd, fdh, ldhD, agxt   aldB

Coralipelagibacter
HIMB5
0.001 0.025   0.050   0.075 0.10   0.25

vgsc   allC   rcl

Littoralipelagibacter
HIMB4
0.001 0.025   0.050   0.075 0.10   0.25

vgsc   tarD rcl

Superopelagibacter
RS40
0.001 0.025   0.050   0.075 0.10   0.25

mylcd, aldB, agxt   mylcd

Ecologically-relevant genes for **offshore** and **coastal** genera

agxt	alanine-glyoxylate/ serine-glyoxylate/ serine-pyruvate transaminase	allC	allantoate deiminase
aldB	aldehyde dehydrogenase	frcABC	fructose transport system
fdh	formate dehydrogenase	rcl	nucleoside 2-deoxyribosyltransferase
ldhD	d-lactate dehydrogenase	tarD	d-tartrate dehydrogenase
mlycd	malonyl-CoA decarboxylase	vgsc	voltage-gated sodium channel
nupNOPQ	guanosine transporter		
uraA	xanthine permease		

Fig. 5. **Habitat-specific genes are under relatively higher selective pressures compared to non-diagnostic genes.** pN/pS values per gene ranked from lowest to highest across each genome. pN/pS values of 0.01, 0.025, 0.05, 0.075, 0.1, and 0.25 are shown in blue. Genes that are differentially distributed between offshore and coastal *Pelagibacteraceae* genera in **Fig. 4** are colored red (offshore) or black (coastal) across the genomes. For genes that are discussed in the text, the gene names are noted and the description of the gene provided in the key. The full suite of gene names and descriptions can be found in **Extended Data Fig. 6**.

Among the genes unique to coastal genera, the voltage-gated sodium channel was under high selective pressure (*vgsc*, pN/pS=0.016±0.004). Coastal Kāneʻohe Bay harbors a larger range of salinity, temperature, and pH conditions compared to the adjacent offshore[26,27,39], and thus the voltage-gated sodium channels may be critical for coastal genera as they rapidly regulate ionic composition[37]. The allantoin deiminase (*allC*, pN/pS=0.039±0.002) and nucleoside 2-deoxyribosyltransferase (*rcl*, pN/pS=0.054±0.006) were also under moderately high selective pressures in coastal genomes, further supporting the specialization of coastal genera on allantonate and nucleotides and a separation in purine metabolism niche from offshore genera. Three sugar transporter genes (*frcABC*, pN/pS=0.024±0.006) within *Ampluspelagibacter* and a tartrate dehydratase (*tarD*, pN/pS=0.051±0.001) within *Littoralipelagibacter* exhibited low pN/pS values suggesting that sugar metabolisms and transporters are not only a more common genomic feature in the coastal genera, but also likely essential for their fitness.

Finally, to understand which *Pelagibacteraceae* core genes were under the highest selective pressures in the KByT system, we examined the mean pN/pS values for core genes found across the same seven genomes. Not surprisingly, ribosomal proteins were found among the genes with the highest purifying selection (low pN/pS values; **Table 2**). Genes related to baseline functions of the cell including general energy production and conversation, translational, ribosomal structure and biogenesis, transcription, and nucleotide transport and metabolism were also under high selective pressure (**Table 2**). Two genes involved in regulatory systems for

responses to acidity and glycine were among the top 25 core genes with the most purifying

selection. The low pN/pS values for these genes supports that although *Pelagibacteraceae* may

have few regulatory genes generally[29], the ones it does have likely impact the the performance

and survival of *Pelagibacteraceae* cells.

Only two genes related to nutrient acquisition were found among the top 25 core genes

with the highest purifying selection. This included both subunits of the adenylylsulfate reductase

(*aprAB*), which is involved in dissimilatory sulfur metabolism, taking sulfite to adenylyl sulfate

(APS). Incomplete dissimilatory and assimilatory sulfate reduction pathways were core among

all *Pelagibacteraceae* examined in this study (**Extended Data Fig. 7**), in line with past analysis

of *Pelagibacteraceae* genomes[52]. As reported by Smith et al.[53], adenylylsulfate reductase likely

removes sulfite that accumulates during organic sulfur compound degradation. Further

supporting this hypothesis, we found multiple core or near core genes involved in the production

of sulfite from reduced organosulfur compounds: (2R)-sulfolactate sulfo-lyase (*suyAB*), a

sulfoacetaldehyde acetyltransferase (*xsc*), and a sulfur dioxygenase (**Extended Data Fig. 7**). Our

analyses reveal the genetic potential to metabolize a wide diversity of reduced organosulfur

compounds as alternative sources of sulfur and that adenylylsulfate reductases (*aprAB*) likely

plays an important role in accommodating *Pelagibacteraceae*'s unique sulfur metabolisms.

Table 2. **Core *Pelagibacteraceae* genes under high selective pressures in the KByT system**. The lowest 25 mean pN/pS values for core *Pelagibacteraceae* genes with shared KOfam annotations examined from genomes representing the seven *Pelagibacteraceae* genera that are highly abundant in the tropical Pacific. pN/pS: proportion of non-synonymous to synonymous. Enzymes involved in regulatory processes are denoted by a pound sign (#).

Accession	Description	pN/pS (mean±s.d.)
K02946	small subunit ribosomal protein S10	0.012±0.004
K00340	NADH-quinone oxidoreductase subunit K	0.013±0.008
K03560	biopolymer transport protein TolR	0.014±0.007
K03628	transcription termination factor Rho	0.014±0.004
K02078	acyl carrier protein	0.014±0.007
K03544	ATP-dependent Clp protease, subunit ClpX	0.017±0.004
K00053	ketol-acid reductoisomerase	0.017±0.005
K00395	adenylylsulfate reductase, subunit B	0.018±0.008
K00394	adenylylsulfate reductase, subunit A	0.018±0.005
K00412	ubiquinol-cytochrome c reductase cytochrome b subunit	0.018±0.006
K02274	cytochrome c oxidase subunit I	0.018±0.005
K02982	small subunit ribosomal protein S3	0.019±0.007
K03566#	LysR family transcriptional regulator, glycine cleavage system transcriptional activator	0.019±0.011
K01494	dCTP deaminase	0.019±0.006
K02878	large subunit ribosomal protein L16	0.019±0.008
K03798	cell division protease FtsH	0.019±0.005
K03702	excinuclease ABC subunit B	0.020±0.008
K00330	NADH-quinone oxidoreductase subunit A	0.020±0.010
K03569	rod shape-determining protein MreB	0.020±0.011
K02874	large subunit ribosomal protein L14	0.020±0.011
K14981#	two-component system, OmpR family, response regulator ChvI	0.020±0.01
K02988	small subunit ribosomal protein S5	0.021±0.005
K03046	DNA-directed RNA polymerase subunit beta	0.021±0.003
K03701	excinuclease ABC subunit A	0.021±0.003
K00331	NADH-quinone oxidoreductase subunit B	0.021±0.006

Discussion

Through genomic and metagenomic characterizations of a recently expanded collection of SAR11 isolates[22], our study reveals a polyphyletic distribution of coastal ocean versus offshore habitat specialists among closely related SAR11 populations, where the emergence of key genetic features that underlie this strong habitat preferences seem to have occurred through independent evolutionary events rather than transfer from a single common ancestor. Genera that share the same environment (e.g. coastal or offshore) have also accumulated shared and unique gene content that is predominantly involved in the metabolism of organic carbon and the acquisition of nitrogen from organic sources. A subset of these metabolic genes were under high purifying selection, emphasizing the importance of these functions to the fitness of distinct *Pelagibacteraceae* genera and highlighting potential determinants of niche differentiation of *Pelagibacteraceae* in coastal and offshore environments.

The cohesion of both ecological and genetic diversity observed within genera of *Pelagibacteraceae* most closely resembles ecotypes: ecologically homogeneous groups of closely related bacteria whose genetic diversity is guided by cohesive forces of selection, recombination, and genetic drift[54,55]. Ecotypic differentiation has previously been suggested to describe the spatial and temporal variation partitioning genetic and genomic SAR11 diversity[9,12,22,26,56–61], however the underlying evolutionary processes or functional differences contributing to this diversification have rarely been linked to patterns of distribution. Our findings reveal evolutionary drivers behind ecotypic differentiation by explaining distinct ecological distribution patterns with differences in gene content, metabolic potential, and selective pressures across ecotypes. The use of a constrained model system that provided both

metagenomic sequence data as well as isolate genomes from local *Pelagibacteraceae*

populations was paramount to being able to resolve signatures of environmental selection in

sympatric and parapatric SAR11 populations. The power of this approach is further demonstrated

in the fact that most of the *Pelagibacteraceae* ecotypes defined in this constrained system as

having preferences for coastal oceans or the open ocean appear to hold when global read

recruitment data were examined[22].

Our results suggest that *Pelagibacteraceae* has transitioned between offshore and coastal

environments multiple times, with a handful of genes that support these lifestyles likely acquired

through multiple independent processes. The small number of adaptive genes are shared among

ecologically similar, but polyphyletically-distributed genera, where genomes have maintained a

substantial amount of genetic diversity. Given the distribution patterns of the adaptive genes and

the genetic diversity maintained within the populations that carry them, the proliferation of

adaptive genes unlikely occurred via genome-wide selective sweep[62], a process that describes the

clonal expansion of an individual subpopulation that carries an adaptive gene leading to a

genome-wide reduction in genetic diversity. Instead, the maintenance of genetic diversity and the

relatively strong purifying selection to maintain the functional identity of these adaptive genes is

consistent with a gene-specific selective sweep[25,63–66]. Consistent with previous studies that have

identified a handful of genes responsible for habitat-specific adaptation[25,66,67], the gene content

differences attributed to stable ecological speciation here are generally found within the

relatively stable genomic backbone, rather than within hyper-variable genomic islands. This is a

critical observation as it links eco-evolutionary processes to specific portions of the immense

genomic diversity that exists within *Pelagibacteraceae*, and underscores the importance of

analyses informed by genomic architecture. Future studies that examine the contribution of gene

content found within hyper-variable islands to stable ecological speciation as opposed to incipient speciation or intraspecific diversity would provide a better understanding of the role of hyper-variable regions in long-term evolutionary processes.

In closing, our analyses broaden the metabolic diversity known from *Pelagibacteraceae* genomes and suggest that metabolic versatility has contributed to SAR11's success in the global ocean. While metabolic reconstructions are subject to uncertainties that necessitate physiological exploration with controlled experimentation in the laboratory, the metabolic analysis of high-quality genomes from cultivated isolates provides the opportunity to delineate metabolic specialization that could advance cultivation approaches and allow for more targeted media recipes[68–71]. SAR11 is estimated to oxidize between 6% to 37% of gross ocean primary productivity[20]. Future work combining these metabolic predictions with *in situ* measures will greatly improve our understanding of how SAR11 impacts the dissolved organic matter pool and biogeochemical cycles in the ocean.

Methods

Pangenome analyses

Publicly-available *Pelagibacteraceae* isolate genomes downloaded from the National Center for Biotechnology Information (NCBI) or the Joint Genome Institute (JGI) were combined with genomes sequenced from newly isolated strains[22] (**Supplementary file 1a**). Genome completion and contamination were examined with checkM v1.1.2[72] and only isolates above 95% completion with less than 5% redundancy[73] and fewer than 50 contigs were kept (**Supplementary file 1a).** High-quality *Pelagibacteraceae* genomes were used to construct a

pangenome using anvi'o v8.0[74] following previously described pipelines[67]. Briefly, anvi'o

database was created using `anvi-gen-contigs-db` and Prodigal v2.6.3[75] was used to identify

open reading frames (ORFs) from contigs. Single-copy core genes were identified using

HMMER v3.2.1[76]. ORFs with associated functions were annotated from NCBI's Clusters of

Orthologous Groups (COGs)[77] and KOfams[23,24] (a customized HMM database of KEGG

orthologs). The pangenome was created with `anvi-pan-genome`, which uses NCBI's Basic

Local Alignment Search Tool (BLAST)[78] to quantify the similarity between pairs of gene clusters

and the Markov Cluster algorithm (MCL)[79] to define homologous gene clusters with a MCL

inflation parameter of 2. The pangenomes were visualized using the command

`anvi-display-pan` and summary tables exported using the command `anvi-summarize`.

Genome-wide average nucleotide identity (gANI) was estimated using FastANI in

anvi'o[80]. A phylogenomic tree of the *Pelagibacteraceae* was estimated using IQ-Tree v2.12[81]

using 1000 ultrafast bootstraps and model LG+F+R10 from a concatenated alignment of a

custom gene set for SAR11 (SAR11_165[22]). Phylogenies were rooted and branches trimmed in R

v 4.4.1[82] using treeio v1.28.0[83] and visulized in R using phytools v 2.3.0[84] to examine the

phylogenomic relationships between *Pelagibacteraceae* groups detected in our environmental

study.

Gene clusters found in 100% of *Pelagibacteraceae* genomes were considered core while

gene clusters found in only one genome were considered singletons. Gene clusters were also

assessed as genus specific (shared among all representatives of a genus and not found in other

genera) and multi-genus specific (shared among all representatives of two or more genera and

not found among all others). Shared gene content between genera was assessed using

ComplexUpset v 1.3.3[85].

Metagenomic read recruitment and environmental analyses

To examine the distribution of *Pelagibacteraceae* genera within the environments from where the majority were isolated, metagenomic read recruitment was conducted using surface ocean metagenomes from 10 stations within and adjacent to Kānʻeohe Bay, Hawaiʻi (**Fig. 1**) collected as part of the Kāneʻohe Bay Time-series (KByT; PRJNA971314)[26,27]. In addition, metagenomes collected in the surrounding North Pacific Subtropical Gyre at Station ALOHA, a sampling location within the Hawaii Ocean Time-series (HOT)[86,87] (**Fig. 1; Supplementary file 1c**; PRJNA352737[88]) were also used. Metagenomes were competitively mapped with Bowtie2 v 2.3.5[89] to the anviʻo contig database of *Pelagibacteraceae* isolate genomes. The `anvi-profile` function stored coverage and detection statistics of each *Pelagibacteraceae* genome found in the KByT and Station ALOHA metagenomic samples and the `anvi-meta-pan-genome` function[67] was used to bring together the pangenomic information with the read recruitment data.

To evaluate the distribution of individual genomes, a detection metric- the proportion of the nucleotides in a given sequence that are covered by at least one short read- was used to define whether a population was detected in a metagenomic sample. A detection value of at least 0.25 was used as criteria to eliminate false positives that could arise if an isolate genome was falsely found within a sample[90].

The average depth of coverage excluding nucleotide positions with coverages in the 1st and 4th quartiles (mean coverage Q2Q3) was mapped for each genome in each sample. To avoid biased estimates of coverage that can occur due to highly recruiting accessory genes, read recruitment was analyzed using only single-copy genes core to each genus. Using a custom script in R, gene clusters that were found within all genomes of the genus (e.g. core), but found only in a single copy within each genome were identified. Then the mean coverage Q2Q3 read

recruitment data per gene were then subsetted for only single copy core genes. The mean

coverage Q2Q3 of each single-copy core gene was summed per genome per sample and then this

value was normalized per genome per sample, based on whether the genome was detected at

>0.25[90]. Next, when evaluating read recruitment at the genus-level, the read recruitment was

summed for all genomes in a genus per sample. These outputs were then divided across the total

Pelagibacteraceae read recruitment for the sample to yield a relative estimate of each genome or

genus within a sample. For the genus with only a single genome representative

(*Semelpelagibacter*), the single-copy core genes shared with the HIMB1483 genome and nearest

neighbor, genus *Lacunapelagibacter,* were used.

K-means clustering analysis of a scaled matrix of biogeochemical conditions was used to

characterize the environmental background from which the metagenomic data were derived.

Metadata from Station ALOHA were downloaded from

https://hahana.soest.hawaii.edu/hot/hot-dogs/ (accessed 11 Oct 2021). All KByT (n=158) and

Station ALOHA (n=34) surface seawater samples (depth of <30m) with data for flow

cytometrically-determined cellular abundance, chlorophyll *a* concentrations, silicate

concentrations, temperature, pH, and salinity were included in the analyses (**Supplementary file**

1c). The number of clusters were estimated using the `kmeansruns` command in the R fpc v

2.2.13 package[91] using a Calinski Harabasz index. Principal Component Analysis (PCA) was

conducted using the `prcomp` function in R v 4.4.1[82] and visualized using 'ggbiplot' in the

ggplot2 v 3.5.1[92].

The sampling locations of the Kāneʻohe Bay Time-series and Station ALOHA were

mapped using R with `geom_sf` from ggplot2 with geospatial data of the main Hawaiian Islands

(USGS Digital Line Graphs). To visualize *Prochlorococcus* cellular abundance across the

Kāneʻohe Bay Time-series, `mba.surf` from MBA v 0.1.2[93] was used to interpolate data over the

KByT stations.

Functional inferences from *Pelagibacteraceae* genera

Functional enrichment analysis was performed in anviʼo and has been described

previously[94]. Briefly, the `anvi-compute-functional-enrichment` command, as well as the

`anvi-script-gen-function-matrix-across-genomes`, was used to assess the enrichment of

Clusters of Orthologous Groups (COG), KEGG orthologs (KOs), and KEGG Modules across

genomes and genus affiliations. The degree of completeness of individual KEGG modules[95,96] in

the genomes and genera was evaluated using `anvi-estimate-metabolism`[97], prior to

`anvi-compute-functional-enrichment`. The functional pangenome was visualized using

`anvi-display-functions`. Central and alternative carbon metabolisms, amino acid, vitamin, and

cofactor biosynthesis, and genes to cope with nutrient, osmotic, and oxidative stress were

examined across the *Pelagibacteraceae* isolate genomes. The distribution of functional genes

were evaluated for each genus as follows: core genes found in all genomes of a genus,

intermediate genes found in 15-99% of genomes, and rare genes found in less than 15%.

Heatmaps of the metabolic gene distribution across genera were made in the R package

pheatmap v 1.0.12[98].

Selection within *Pelagibacteraceae* genes

To evaluate the selective pressures on individual metabolic genes of interest, the

proportion of non-synonymous to synonymous (pN/pS) sites per gene across genomes was

evaluated. The pipeline followed those developed in Kiefl et al.,[18]. Briefly, contig databases from

the genomes of type specimens from each of the *Pelagibacteraceae* genera[22] were first

functionally annotated in anvi'o (see above) and metagenomic reads from deeply sequenced

metagenomes from either coastal Kāneʻohe Bay or the adjacent offshore were non-competitively

recruited using Bowtie2 v 2.3.5[89] (**Supplementary file 1c**).

Next, the`anvi-profile` function with the flag `--profile-SCVs` was used to characterize

single codon variants across the read recruitment data and then `anvi-gen-variability-profile`

was used to export the single codon variants per gene per sample using the flags --engine CDN

--include-site-pN/pS --kiefl-mode[18]. Finally, pN/pS values were reported per gene with at least

20x coverage in each sample using 'anvi-get-pn-ps-ratio` with the -m 20 flag. Per gene coverage

statistics were exported using `anvi-summarize` with the `--init-gene-coverages` flag. Functional

annotations were exported using `anvi-export-functions`.

To evaluate whether a single genome representative contained similar pN/pS gene values

as other genomes within a given genus, these steps were repeated with all representatives of the

genus sharing the lowest minimum gANI (among the genera that were abundant in the tropical

Pacific): genus *Coralipelagibacter*. The exported data files were brought into R, where variation

in pN/pS per sample, per genome, and across genomes of the same genus was evaluated.

Relationships between pN/pS and coverage were examined and genes per genome representative

were ranked from lowest pN/pS value to highest and visualized in R using ggplot2 v 3.5.1[92].

Figures were made in R v 4.4.1[82] or anvi'o[74] and further edited with Inkscape

(https://inkscape.org/).

Data availability

Accession numbers for genome sequences used to conduct pangenomic analyses and read

recruitment in this study are found in **Supplementary file 1a**. Short-read metagenome accession

IDs and environmental data used in this study are provided in **Supplementary file 1c**. Long-read

metagenomes from coastal Kāneʻohe Bay are available under NCBI Project PRJNA1201851.

Code availability

Code to run analyses and create figures can be found at

https://github.com/tucker4/SAR11_metabolism.

Competing interests

The authors declare no competing interests.

Acknowledgements

We thank Oscar Ramfelt and Evan B. Freel for their insights on computational analyses. This research was funded by the Simons Postdoctoral Fellowships in Marine Microbial Ecology (grant #989028), NOAA Margaret A. Davidson Fellowship, the Colonel Willys E. Lord, DVM and Sadina L. Lord Scholarship, the ARCS Foundation George Orton and Mona Marie Elmore Award in Marine Biology, the Research Corporation of the University of Hawaiʻi Graduate Research Fellowship, the EECB Hampton and Meredith Carson Fellowship, and the Center for Microbial Analysis through Island Knowledge & Investigation Research Grant to SJ Tucker, and NSF grants OCE-1538628 and OCE- 2149128 to MS Rappé. AME acknowledges support from the Center for Chemical Currencies of a Microbial Planet (C-CoMP) (NSF Award OCE-2019589, C-CoMP publication #065), and Simons Foundation (grant #687269)

References

1. Louca, S. *et al.* Integrating biogeochemistry with multiomic sequence information in a model oxygen minimum zone. *Proc. Natl. Acad. Sci. U. S. A.* **113**, E5925–E5933 (2016).

2. Hao, L. *et al.* Novel syntrophic bacteria in full-scale anaerobic digesters revealed by genome-centric metatranscriptomics. *ISME J.* **14**, 906–918 (2020).

3. Wang, C. *et al.* Bacterial genome size and gene functional diversity negatively correlate with taxonomic diversity along a pH gradient. *Nat. Commun.* **14**, 7437 (2023).

4. Cordero, O. X. & Polz, M. F. Explaining microbial genomic diversity in light of evolutionary ecology. *Nat. Rev. Microbiol.* **12**, 263–273 (2014).

5. Sheppard, S. K., Guttman, D. S. & Fitzgerald, J. R. Population genomics of bacterial host adaptation. *Nat. Rev. Genet.* **19**, 549–565 (2018).

6. Whelan, F. J., Hall, R. J. & McInerney, J. O. Evidence for selection in the abundant accessory gene content of a prokaryote pangenome. *Mol. Biol. Evol.* **38**, 3697–3708 (2021).

7. Conrad, R. E. *et al.* Toward quantifying the adaptive role of bacterial pangenomes during environmental perturbations. *ISME J.* **16**, 1222–1234 (2022).

8. Morris, R. M. *et al.* SAR11 clade dominates ocean surface bacterioplankton communities. *Nature* **420**, 806–810 (2002).

9. Brown, M. V. *et al.* Global biogeography of SAR11 marine bacteria. *Mol. Syst. Biol.* **8**, 595 (2012).

10. Grote, J. *et al.* Streamlining and core genome conservation among highly divergent members of the SAR11 clade. *MBio* **3**, (2012).

11. Delmont, T. O. *et al.* Single-amino acid variants reveal evolutionary processes that shape the biogeography of a global SAR11 subclade. *Elife* **8**, (2019).

12. Haro-Moreno, J. M. *et al.* Ecogenomics of the SAR11 clade. *Environ. Microbiol.* **22**, 1748–1763 (2020).

13. Giovannoni, S. J. *et al.* Genome streamlining in a cosmopolitan oceanic bacterium. *Science* **309**,

1242–1245 (2005).

14. Noell, S. E. *et al.* Differences in the regulatory strategies of marine oligotrophs and copiotrophs

reflect differences in motility. *Environ. Microbiol.* **25**, 1265–1280 (2023).

15. Morris, J. J., Lenski, R. E. & Zinser, E. R. The Black Queen Hypothesis: evolution of dependencies

through adaptive gene loss. *MBio* **3**, (2012).

16. Braakman, R., Follows, M. J. & Chisholm, S. W. Metabolic evolution and the self-organization of

ecosystems. *Proc. Natl. Acad. Sci. U. S. A.* **114**, E3091–E3100 (2017).

17. Vergin, K. L. *et al.* High intraspecific recombination rate in a native population of Candidatus

pelagibacter ubique (SAR11). *Environ. Microbiol.* **9**, 2430–2440 (2007).

18. Kiefl, E. *et al.* Structure-informed microbial population genetics elucidate selective pressures that

shape protein evolution. *Sci Adv* **9**, eabq4632 (2023).

19. Grant, S. R., Church, M. J., Ferrón, S., Laws, E. A. & Rappé, M. S. Elemental composition,

phosphorous uptake, and characteristics of growth of a SAR11 strain in batch and continuous culture.

mSystems **4**, (2019).

20. White, A. E., Giovannoni, S. J., Zhao, Y., Vergin, K. & Carlson, C. A. Elemental content and

stoichiometry of SAR11 chemoheterotrophic marine bacteria. *Limnol. Oceanogr. Lett.* **4**, 44–51

(2019).

21. Acker, M. *et al.* Phosphonate production by marine microbes: Exploring new sources and potential

function. *Proc. Natl. Acad. Sci. U. S. A.* **119**, e2113386119 (2022).

22. Freel, K. C., Tucker, S. J., Freel, E. B., Eren, A. M. & Rappé, M. S. New isolate genomes and global

marine metagenomes resolve ecologically relevant units of SAR11. *In prep.*

23. Kanehisa, M. & Goto, S. KEGG: kyoto encyclopedia of genes and genomes. *Nucleic Acids Res.* **28**,

27–30 (2000).

24. Aramaki, T. *et al.* KofamKOALA: KEGG Ortholog assignment based on profile HMM and adaptive

score threshold. *Bioinformatics* **36**, 2251–2252 (2020).

25. Shapiro, B. J. *et al.* Population genomics of early events in the ecological differentiation of bacteria. *Science* **336**, 48–51 (2012).

26. Tucker, S. J. *et al.* Spatial and temporal dynamics of SAR11 marine bacteria across a nearshore to offshore transect in the tropical Pacific Ocean. *PeerJ* **9**, e12274 (2021).

27. Tucker, S. J. *et al.* Sharp transitions in phytoplankton communities across estuarine to open ocean waters of the tropical Pacific. *bioRxiv* (2024) doi:10.1101/2024.05.23.595464.

28. Zhang, Y. & Gladyshev, V. N. Molybdoproteomes and evolution of molybdenum utilization. *J. Mol. Biol.* **379**, 881–899 (2008).

29. Giovannoni, S. J., Cameron Thrash, J. & Temperton, B. Implications of streamlining theory for microbial ecology. *ISME J.* **8**, 1553–1565 (2014).

30. Berman, T. & Bronk, D. A. Dissolved organic nitrogen: a dynamic participant in aquatic ecosystems. *Aquat. Microb. Ecol.* **31**, 279–305 (2003).

31. Kornberg, R. D. Chromatin structure: a repeating unit of histones and DNA. *Science* **184**, 868–871 (1974).

32. Braakman, R. *et al.* Global niche partitioning of purine and pyrimidine cross-feeding among ocean microbes. *bioRxiv* (2024) doi:10.1101/2024.02.09.579562.

33. Tripp, H. J. *et al.* Unique glycine-activated riboswitch linked to glycine-serine auxotrophy in SAR11. *Environ. Microbiol.* **11**, 230–238 (2009).

34. Klähn, S., Steglich, C., Hess, W. R. & Hagemann, M. Glucosylglycerate: a secondary compatible solute common to marine cyanobacteria from nitrogen-poor environments. *Environ. Microbiol.* **12**, 83–94 (2010).

35. Castell, C., Rodríguez-Lumbreras, L. A., Hervás, M., Fernández-Recio, J. & Navarro, J. A. New insights into the evolution of the electron transfer from cytochrome f to photosystem I in the Green and red branches of photosynthetic eukaryotes. *Plant Cell Physiol.* **62**, 1082–1093 (2021).

36. Peers, G. & Price, N. M. Copper-containing plastocyanin used for electron transport by an oceanic diatom. *Nature* **441**, 341–344 (2006).

37. Ren, D. *et al.* A prokaryotic voltage-gated sodium channel. *Science* **294**, 2372–2375 (2001).

38. Strege, P. R. *et al.* Mechanosensitive pore opening of a prokaryotic voltage-gated sodium channel.

Elife **12**, (2023).

39. Yeo, S. K., Huggett, M. J., Eiler, A. & Rappé, M. S. Coastal bacterioplankton community dynamics

in response to a natural disturbance. *PLoS One* **8**, e56207 (2013).

40. Sun, J. *et al.* One carbon metabolism in SAR11 pelagic marine bacteria. *PLoS One* **6**, e23973 (2011).

41. Giovannoni, S. J. SAR11 bacteria: The most abundant plankton in the oceans. *Ann. Rev. Mar. Sci.* **9**,

231–255 (2017).

42. Moore, E. R., Weaver, A. J., Davis, E. W., Giovannoni, S. J. & Halsey, K. H. Metabolism of key

atmospheric volatile organic compounds by the marine heterotrophic bacterium Pelagibacter

HTCC1062 (SAR11). *Environ. Microbiol.* **24**, 212–222 (2022).

43. Gralka, M., Pollak, S. & Cordero, O. X. Genome content predicts the carbon catabolic preferences of

heterotrophic bacteria. *Nat. Microbiol.* **8**, 1799–1808 (2023).

44. Leyton, D. L., Rossiter, A. E. & Henderson, I. R. From self sufficiency to dependence: mechanisms

and factors important for autotransporter biogenesis. *Nat. Rev. Microbiol.* **10**, 213–225 (2012).

45. Doyle, M. T., Tran, E. N. H. & Morona, R. The passenger-associated transport repeat promotes

virulence factor secretion efficiency and delineates a distinct autotransporter subtype. *Mol.*

Microbiol. **97**, 315–329 (2015).

46. Zhao, X. *et al.* Three-Dimensional Structure of the Ultraoligotrophic Marine Bacterium 'Candidatus

Pelagibacter ubique'. *Appl. Environ. Microbiol.* **83**, (2017).

47. Hazes, B. & Frost, L. Towards a systems biology approach to study type II/IV secretion systems.

Biochim. Biophys. Acta **1778**, 1839–1850 (2008).

48. Ellison, C. K. *et al.* Retraction of DNA-bound type IV competence pili initiates DNA uptake during

natural transformation in Vibrio cholerae. *Nat. Microbiol.* **3**, 773–780 (2018).

49. Meuskens, I., Saragliadis, A., Leo, J. C. & Linke, D. Type V secretion systems: An overview of

passenger domain functions. *Front. Microbiol.* **10**, 1163 (2019).

50. Noell, S. E. & Giovannoni, S. J. SAR11 bacteria have a high affinity and multifunctional glycine

betaine transporter. *Environ. Microbiol.* **21**, 2559–2575 (2019).

51. Clifton, B. E., Alcolombri, U., Uechi, G.-I., Jackson, C. J. & Laurino, P. The ultra-high affinity

transport proteins of ubiquitous marine bacteria. *Nature* **634**, 721–728 (2024).

52. Tripp, H. J. *et al.* SAR11 marine bacteria require exogenous reduced sulphur for growth. *Nature* **452**,

741–744 (2008).

53. Smith, D. P. *et al.* Proteome Remodeling in Response to Sulfur Limitation in 'Candidatus

Pelagibacter ubique'. *mSystems* **1**, (2016).

54. Cohan, F. M. Towards a conceptual and operational union of bacterial systematics, ecology, and

evolution. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **361**, 1985–1996 (2006).

55. Koeppel, A. *et al.* Identifying the fundamental units of bacterial diversity: a paradigm shift to

incorporate ecology into bacterial systematics. *Proc. Natl. Acad. Sci. U. S. A.* **105**, 2504–2509

(2008).

56. Field, K. G. *et al.* Diversity and depth-specific distribution of SAR11 cluster rRNA genes from

marine planktonic bacteria. *Appl. Environ. Microbiol.* **63**, 63–70 (1997).

57. Schwalbach, M. S., Tripp, H. J., Steindler, L., Smith, D. P. & Giovannoni, S. J. The presence of the

glycolysis operon in SAR11 genomes is positively correlated with ocean productivity. *Environ.*

Microbiol. **12**, 490–500 (2010).

58. Vergin, K. L. *et al.* High-resolution SAR11 ecotype dynamics at the Bermuda Atlantic Time-series

Study site by phylogenetic placement of pyrosequences. *ISME J.* **7**, 1322–1332 (2013).

59. Tsementzi, D. *et al.* Ecogenomic characterization of widespread, closely-related SAR11 clades of the

freshwater genus 'Candidatus Fonsibacter' and proposal of Ca. Fonsibacter lacus sp. nov. *Syst. Appl.*

Microbiol. **42**, 495–505 (2019).

60. Kraemer, S., Ramachandran, A., Colatriano, D., Lovejoy, C. & Walsh, D. A. Diversity and

biogeography of SAR11 bacteria from the Arctic Ocean. *ISME J.* **14**, 79–90 (2020).

61. Larkin, A. A., Hagstrom, G. I., Brock, M. L., Garcia, N. S. & Martiny, A. C. Basin-scale

biogeography of Prochlorococcus and SAR11 ecotype replication. *ISME J.* **17**, 185–194 (2023).

62. Shapiro, B. J., Leducq, J.-B. & Mallet, J. What is speciation? *PLoS Genet.* **12**, e1005860 (2016).

63. Falush, D. *et al.* Recombination and mutation during long-term gastric colonization by Helicobacter pylori: estimates of clock rates, recombination size, and minimal age. *Proc. Natl. Acad. Sci. U. S. A.* **98**, 15056–15061 (2001).

64. Whitaker, R. J., Grogan, D. W. & Taylor, J. W. Recombination shapes the natural population structure of the hyperthermophilic archaeon Sulfolobus islandicus. *Mol. Biol. Evol.* **22**, 2354–2361 (2005).

65. Rosen, M. J., Davison, M., Bhaya, D. & Fisher, D. S. Microbial diversity. Fine-scale diversity and extensive recombination in a quasisexual bacterial population occupying a broad niche. *Science* **348**, 1019–1023 (2015).

66. Bendall, M. L. *et al.* Genome-wide selective sweeps and gene-specific sweeps in natural bacterial populations. *ISME J.* **10**, 1589–1601 (2016).

67. Delmont, T. O. & Eren, A. M. Linking pangenomes and metagenomes: the Prochlorococcus metapangenome. *PeerJ* **6**, e4320 (2018).

68. Rappé, M. S., Connon, S. A., Vergin, K. L. & Giovannoni, S. J. Cultivation of the ubiquitous SAR11 marine bacterioplankton clade. *Nature* **418**, 630–633 (2002).

69. Stingl, U., Tripp, H. J. & Giovannoni, S. J. Improvements of high-throughput culturing yielded novel SAR11 strains and other abundant marine bacteria from the Oregon coast and the Bermuda Atlantic Time Series study site. *ISME J.* **1**, 361–371 (2007).

70. Song, J., Oh, H.-M. & Cho, J.-C. Improved culturability of SAR11 strains in dilution-to-extinction culturing from the East Sea, West Pacific Ocean. *FEMS Microbiol. Lett.* **295**, 141–147 (2009).

71. Henson, M. W. *et al.* Artificial seawater media facilitate cultivating members of the microbial majority from the Gulf of Mexico. *mSphere* **1**, (2016).

72. Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res.*

25, 1043–1055 (2015).

73. Bowers, R. M. *et al.* Minimum information about a single amplified genome (MISAG) and a

    metagenome-assembled genome (MIMAG) of bacteria and archaea. *Nat. Biotechnol.* **35**, 725–731

    (2017).

74. Eren, A. M. *et al.* Community-led, integrated, reproducible multi-omics with anvi'o. *Nat. Microbiol.*

    6, 3–6 (2021).

75. Hyatt, D. *et al.* Prodigal: prokaryotic gene recognition and translation initiation site identification.

    BMC Bioinformatics **11**, 119 (2010).

76. Eddy, S. R. Accelerated profile HMM searches. *PLoS Comput. Biol.* **7**, e1002195 (2011).

77. Tatusov, R. L. *et al.* The COG database: an updated version includes eukaryotes. *BMC*

    Bioinformatics **4**, 41 (2003).

78. Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment search tool.

    J. Mol. Biol. **215**, 403–410 (1990).

79. Enright, A. J., Van Dongen, S. & Ouzounis, C. A. An efficient algorithm for large-scale detection of

    protein families. *Nucleic Acids Res.* **30**, 1575–1584 (2002).

80. Jain, C., Rodriguez-R, L. M., Phillippy, A. M., Konstantinidis, K. T. & Aluru, S. High throughput

    ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat. Commun.* **9**, 5114

    (2018).

81. Minh, B. Q. *et al.* IQ-TREE 2: New models and efficient methods for phylogenetic inference in the

    genomic era. *Mol. Biol. Evol.* **37**, 1530–1534 (2020).

82. R Core Team. *R: A Language and Environment for Statistical Computing.* (2023).

83. Wang, L.-G. *et al.* Treeio: An R package for phylogenetic tree input and output with richly annotated

    and associated data. *Mol. Biol. Evol.* **37**, 599–603 (2020).

84. Revell, L. J. phytools: an R package for phylogenetic comparative biology (and other things).

    Methods Ecol. Evol. **3**, 217–223 (2012).

85. Lex, A., Gehlenborg, N., Strobelt, H., Vuillemot, R. & Pfister, H. UpSet: Visualization of

intersecting sets. *IEEE Trans. Vis. Comput. Graph.* **20**, 1983–1992 (2014).

86. Karl, D. M. & Lukas, R. The Hawaii Ocean Time-series (HOT) program: Background, rationale and

field implementation. *Deep Sea Res. Part 2 Top. Stud. Oceanogr.* **43**, 129–156 (1996).

87. Karl, D. M. & Church, M. J. Microbial oceanography and the Hawaii Ocean Time-series

programme. *Nat. Rev. Microbiol.* **12**, 699–713 (2014).

88. Mende, D. R. *et al.* Environmental drivers of a microbial genomic transition zone in the ocean's

interior. *Nat. Microbiol.* **2**, 1367–1373 (2017).

89. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**,

357–359 (2012).

90. Utter, D. R., Borisy, G. G., Eren, A. M., Cavanaugh, C. M. & Mark Welch, J. L. Metapangenomics

of the oral microbiome provides insights into habitat adaptation and cultivar diversity. *Genome Biol.*

21, 293 (2020).

91. Hennig, C. *Fpc: Flexible Procedures for Clustering. R Package Version 2.2-13.* (2024).

92. Wickham, H. *ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.* (2016).

93. Finley, A., Banerjee, S. & Hjelle, Ø. *MBA: Multilevel B-Spline Approximation. R Package Version*

0.1-2. (2017).

94. Shaiber, A. *et al.* Functional and genetic markers of niche partitioning among enigmatic members of

the human oral microbiome. *Genome Biol.* **21**, 292 (2020).

95. Kanehisa, M. *et al.* Data, information, knowledge and principle: back to metabolism in KEGG.

Nucleic Acids Res. **42**, D199–205 (2014).

96. Kanehisa, M., Furumichi, M., Tanabe, M., Sato, Y. & Morishima, K. KEGG: new perspectives on

genomes, pathways, diseases and drugs. *Nucleic Acids Res.* **45**, D353–D361 (2017).

97. Veseli, I. *et al.* Microbes with higher metabolic independence are enriched in human gut

microbiomes under stress. (2023) doi:10.7554/elife.89862.

98. Kolde, R. *Pheatmap: Pretty Heatmaps. R Package Version 1.0.12.* (2019).

 # Extended Data Figures

Extended Data Fig. 1. a) Phylogenomic tree showing 11 monophyletic clusters within
Pelagibacteraceae, their genus assignments, and source of isolation. Circles at nodes indicate
ultrafast bootstrap support values ≥90% from 1000 replicates. **b)** Detection of all
Pelagibacteraceae genomes in coastal and offshore metagenomic samples.

Extended Data Fig. 2. K-means analyses of environmental parameters from 192 surface ocean
samples from KByT and Station ALOHA grouped into two clusters, herein referred to as coastal
and offshore.

Extended Data Fig. 3. Relative read recruitment of each isolate genome across the coastal and
offshore environment. Data for genomes that are not detected or have a relative recruitment less
than 0.25% are not shown.

a)

b)

genes without near exact matches in long-read data
near exact matches in long-read data
● bootstrap support >= 90%

2.0

c)

HIMB5- *Coralipelagibacter*
Autotransporter in HVR

HIMB4- *Littoralipelagibacter*
Autotransporter not in HVR

fibronectin-binding
autotransporter adhesin
type IV pilus system
type II secretory system

Extended Data Fig. 4. **a)** A protein structure model of a gene annotated as an autotransporter adhesin from KByT long-read metagenomes and sharing close sequence similarity to gene sequences in *Pelagibacteraceae* genomes shows characteristics expected of autotransporter proteins: 1) α-helical linker, 2) β-barrel translocator domain, and 3) passenger domains containing β helices (Leyton et al., 2012). **b)** Homolog searches to recover autotransporter adhesin genes in long-read metagenomes recovered nine non-redundant gene sequences that were near exact matches to most gene sequences in *Pelagibacteraceae* genomes (33 of 43). **c)**

Patterns of detection of short-read metagenomic read recruitment data across genes in two genomes of *Pelagibacteraceae* show that autotransporter adhesion genes are sometimes located in hypervariable regions and sometimes not. Genes for each *Pelagibacteraceae* genome are ordered by genome synteny with detection values of read recruitment data per gene shown as a bar per metagenomic sample. Areas of low detection are indicative of hypervariable regions (HVRs), and the autotransporter adhesin genes were located in HVRs within the HIMB5 genome, but not within HIMB4 genome. The autotransport adhesion genes were always positioned next to genes involved in type IV pilus systems and/or type II secretory systems.

957

959 **Extended Data Fig. 5**. The coverage of genes from metagenomic read recruitment did not
960 correlate with the pN/pS values. pN/pS: proportion of non-synonymous to synonymous.

958

aidA	fibronectin-binding autotransporter adhesin
agxt	transaminase
aldB	aldehyde dehydrogenase
allC	allantoate deiminase
amyA	glycosidase/amylase
c3lhypd	cis-L-3-hydroxyproline dehydratase
cutA	copper tolerance
dehH	haloacetate dehalogenase
dhaK, dhaL	phosphoenolpyruvate: glycerone phosphotransferase
fdh	formate dehydrogenase
frcABC	fructose transport system
fucB	L-fuconolactonase
gdhA	glutamate dehydrogenase
glpA	glycerol-3-phosphate dehydrogenase
gpgS, yedP	glucosylglycerate synthesis
pucL	uric acid degradation
putA	proline dehydrogenase
puuC	putrescine glutamylation
rcl	nucleoside 2-deoxyribosyltransferase
tadA	tRNA adenosine deaminase
tarD	D-tartrate dehydratase
tdcB	threonine dehydratase
unch	n terminal nucleophile
uraA	xanthine permease
vgsc	voltage-gated sodium channel
xdhAB	xanthine dehydrogenase
xylB	xylulokinase
yedP	mannosyl-3-phosphoglycerate phosphatase
yfdE	CoA:oxalate CoA-transferase
yrbG	cation:H+ antiporter

gpr	L-glyceraldehyde 3-phosphate reductase
hupE	nickel transport
isdG	heme oxygenase
katG	catalase-peroxidase
kce, lcdH	L-carnitine degradation
ldhD	D-lactate dehydrogenase
mcl, mtkA	S-malate metabolism
mdcF	malonate transporter
mmcO	copper tolerance
mylcd	malonyl-CoA decarboxylase
nit2	omega-amidase
nupNOPQ	guanosine uptake
ocd	ornithine cyclodeaminase
phnZ	2-aminoethylphosphonic acid degradation
pitA	phosphate/sulfate permease

Extended Data Fig. 6. All genes in a given genome are ordered by lowest pN/pS value to highest, with genes that are differentially distributed between offshore and coastal Pelagibacteraceae genera in Fig. 4 colored red (offshore) or black (coastal) across the genomes. pN/pS values of 0.01, 0.025, 0.05, 0.075, 0.1, and 0.25 are shown in blue. pN/pS: proportion of non-synonymous to synonymous.

969 **Extended Data Fig. 7**. *Pelagibacteraceae* has incomplete dissimilatory and assimilatory sulfur
970 reduction pathways and the capacity to utilize a wide diversity of organosulfate compounds.
971 APS: adenylyl sulfate, PAPS: 3'-phosphoadenylyl sulfate, DHSP:
972 2,3-dihydroxypropane-1-sulfonate, DMSP: dimethylsulfoniopropionate, DMS: dimethyl sulfide,
973 MMPA: methylmercaptopropionate, CoA: coenzyme A.

Supplementary Information

978 All supplementary infomation text and data files are available on Figshare at:

979 10.6084/m9.figshare.28087925.