TSvelo: Comprehensive RNA velocity by modeling the cascade of gene regulation, transcription and splicing

Jiachen Li[1], Zhe Wang[1], Hong-Bin Shen[2], Ye Yuan[1,2]*

[1]State Key Laboratory of Biopharmaceutical Preparation and Delivery, Institute of Process Engineering, Chinese Academy of Sciences, Beijing 100190, China

[2]Institute of Image Processing and Pattern Recognition, Shanghai Jiao Tong University, and Key Laboratory of System Control and Information Processing, Ministry of Education of China, Shanghai, 200240, China

*Address correspondence to Y. Y. at: yyuan@ipe.ac.cn or yuanye_auto@sjtu.edu.cn

Abstract

RNA velocity approaches fit gene dynamics and infer cell fate by modeling the splicing process using single-cell RNA sequencing (scRNA-seq) data. However, due to short time scale of splicing, high noise and large complexity of data, existing RNA velocity methods often fail to precisely capture the complex velocity dynamics for individual gene and single cell, which makes its downstream analysis less reliable and less robust. We propose **TSvelo**, a comprehensive RNA **velo**city mathematics framework that can model the cascade of gene regulation, **T**ranscription and **S**plicing using highly interpretable neural Ordinary Differential Equations (ODEs). TSvelo can precisely capture the transcription-unspliced-spliced 3D dynamics of all genes simultaneously, infer unified latent time shared by genes within single cell, detect key gene regulatory relations and be applied to multi-lineage datasets. Experiments on six scRNA-seq datasets, including two multi-lineage datasets, demonstrate TSvelo's superiority.

Introduction

Single-cell RNA sequencing (scRNA-seq) enables the detailed exploration of gene expression at the individual cell level. To move beyond static snapshots and capture the dynamic behavior of cells over time, several trajectory inference (TI) methods have been developed, such as PAGA[1], Monocle[2], Slingshot[3] and Palantir[4]. These methods typically estimate pseudotime using diffusion processes and require prior annotation of initial cells. In contrast, RNA velocity[5] offers a more interpretable approach by modeling the time derivative of gene expression, linking unspliced (immature) and spliced (mature) mRNA levels through Ordinary Differential Equations (ODEs).

Several RNA velocity methods have been proposed to capture splicing dynamics. The first approach, Velocyto[5], uses least squares solutions to estimate parameters under the

assumption of steady-state kinetics. ScVelo[6] improves upon this by employing an
Expectation-Maximization (EM) approach for better fitting splicing kinetics.
UniTVelo[7] propose a top-down idea that directly models the spliced RNA levels with
a time-dependent function. More recently, generative models such as VeloVI[8],
veloVAE[9] and Pyrovelocity[10] have been introduced, utilizing Bayesian frameworks to
estimate RNA velocity. To handle multi-lineage datasets, methods like CellDancer[11],
DeepVelo[12] and LatentVelo[13] extend RNA velocity by modeling local dynamics with
neural networks, rather than assuming a globally constant transcriptional rate. Apart
from using unspliced/spliced data, Dynamo[14] enhances RNA velocity further by
incorporating labeled RNA-seq data. The advent of single-cell multi-omics[15,16]
technologies has allowed RNA velocity analysis to extend to protein abundance (e.g.,
protaccel[17]) and single-cell ATAC-seq datasets (e.g., MultiVelo[18]). Additionally,
methods like DeepCycle[19] and VeloCycle[20] have been developed to focus on cell
cycle processes, with specialized modules for capturing periodic signals. STT[21] and
SIRV[22] proposed the idea on extending RNA velocity analysis to spatial
transcriptomics.
57
58
Although RNA velocity theory has significantly advanced the inference of single-cell
trajectories, pseudotime, and gene regulation[23], several challenges persist for current
RNA velocity models. Firstly, RNA velocity models primarily infer cell fate based on
phase portrait fitting of unspliced and spliced dynamics for each gene. However, they
often fail to capture the correct phase portrait for most genes, due to the sparsity and
noise in unspliced and spliced mRNA abundance for individual genes, the short time
scale of the splicing process, and the mixing of cells from different types on the phase
portrait[24-26]. Secondly, The majority of existing RNA velocity models treat each gene
independently and fail to incorporate the underlying regulatory interactions[27].
Although some approaches (e.g. TFvelo[26], PHOENIX[28] and scKINETICS[29]) have
constructed ODE models to capture gene dynamics by integrating regulation
information, these methods overlook the splicing signal so that they cannot jointly
model the transcription and splicing dynamics into one unified form. Thirdly, classical
RNA velocity approaches, such as scVelo, use interpretable parameters in constructing
dynamic model for single gene. By contrast, to model flexible transcriptional rates or
integrate multiple genes, several recent methods employ latent space embeddings or
neural network-based encoders[11]. It makes the model parameters less interpretable in
detailed gene level, which is crucial for understanding the underlying biological
mechanisms. Fourthly, multi-lineage tasks remain a significant challenge for current
RNA velocity models due to the complexity in large scale scRNA-Seq datasets.
79
To address the challenges outlined above, we propose TSvelo, a method that
intergrates gene regulation, transcription and splicing of all genes into a single ODE
model, whose parameters are highly interpretable. TSvelo directly learns the global
latent time without the need to separately learn gene-specific latent times. By
leveraging both unspliced and spliced scRNA-seq data, along with gene regulatory

knowledge from TF-target databases, TSvelo iteratively optimizes the parameters in
the ODE model and the unified latent time using Expectation-Maximization (EM)
algorithm. Experiments on six scRNA-seq datasets, including two multi-lineage
datasets, demonstrate that TSvelo outperforms existing methods in modeling gene
dynamics, inferring cell fates, and is also effective in analyzing multi-lineage datasets.
90

Results

Estimate RNA velocity with TSVelo
We present TSvelo, a method for estimating RNA velocity of multiple target genes
simultaneously using both of transcriptional regulation and splicing information.
Initially, the unspliced and spliced RNA abundances are preprocessed for velocity
gene selection and pseudotime initialization (**Fig. 1a**). See **Methods** for details about
preprocessing. Next to model the velocity gene g's expression dynamics, we suppose
u_g and s_g are the abundance of unspliced and spliced RNA, and $\alpha_g(t), \beta_g, \gamma_g$ are
the transcription rate, splicing rate and degradation rate, respectively. The
dynamics is modeled as:

$$\frac{du_g(t)}{dt} = \alpha_g(t) - \beta_g u_g(t) \#(1)$$

$$\frac{ds_g(t)}{dt} = \beta_g u_g(t) - \gamma_g s_g(t) \#(2)$$

101 Furthermore, we assume that the gene and cell-specific transcriptional rate
102 $\alpha_g(t)$ is influence by the expression of Transcriptional Factors (TFs).
103 Considering the wide usage and interpretability of linear models for gene-relations
104 in previous studies[26,30-33], we model $\alpha_g(t)$ as

$$\alpha_g(t) = ReLU(\sum_{i \in TFs(g)} w_{gi} * s_i(t)) \#(3)$$

$TFs(g)$ refers to the TFs that could regulated the target gene g, which are
selected according to the ChEA and ENCODE TF-target database. In order to
include more TFs in the model, we also reserve those TFs that are not selected as
velocity genes, and directly model the dynamics between transcription and spliced
RNA without using the unspliced abundance. Finally, we combine the dynamic
models on all selected genes into the Ordinary Differentia Equation (ODE) matrix
form (**Fig. 1b**), which enables directly inferring a unified latent time for each cell
using its all genes. See **Methods** for details.

Figure 1. The framework of TSvelo. (**a**) The preprocessing strategy in TSvelo, including velocity genes selection and initial state detection. (**b**) The Neural ODE model and its optimization in TSvelo, where the parameters and latent time is optimized iteratively. (**c-e**) The downstream application tasks of TSvelo, including precise transcription-unspliced-spliced 3D phase portrait fitting (c), cell fate prediction using predicted RNA velocity (d) and gene expression pattern analysis for multi-lineage dataset (e).

TSvelo employs an EM framework to iteratively optimize both latent time   and the parameters in ODE. The global pseudotime   is assigned to each cell through grid search. Due to the difficulty in calculating analytical solutions of   or   , TSvelo adopted a Neural ODE for estimating those parameters of transcription rate, splicing rate and degradation rate. Next, TSvelo can accurately model the gene

dynamics of both transcription, unsplicing and splicing processes (**Fig. 1c**), predict cell states using both pseudotime and velocity stream analysis (**Fig. 1d**), and is also applicable to multi-lineage tasks for analyzing expression patterns across different lineages in large complex scRNA-Seq datasets (**Fig. 1e**).

TSvelo can accurately model 3D gene dynamics and predict cell fate on pancreas dataset

To validate the TSvelo model, we applied it to the pancreas scRNA-seq dataset[5], which is widely used in RNA velocity studies to model cell differentiation from ductal to endocrine cells. Using the predicted RNA velocity, TSvelo can generate velocity stream plot using the functions provided in scVelo as well. Both the final pseudotime t (**Fig. 2a**) and the velocity stream plot (**Fig. 2b**) effectively capture the cell differentiation process. We quantitively compare TSvelo and baseline approaches, including scVelo [6], dynamo[14], UniTVelo[7], cellDancer[11] and TFvelo[26]. TSvelo achieves the highest median velocity consistency (**Fig. 2c**), which demonstrates that the high-dimensional velocity vectors learned by TSvelo are mostly coherent within neighbor cells. TSvelo also achieves the highest median in-cluster coherence and cross-boundary correctness, which validates that TSvelo best fit the differentiation process within these cell types according to the ground truth annotation (**Fig. S2** in **Supplementary Information**).

Figure 2. Results on pancreas dataset. (**a**) The pseudotime learned with TSvelo. (**b**) The Stream-plot for visualization the RNA velocity inferred by TSvelo. (**c**) The quantitative comparison between TSvelo and multiple baseline approaches. The down, central and up hinges correspond to the first quartile, median value and third quartile, respectively. The whiskers extend to $1.5\times$ the interquartile range of the distribution from the hinge. 3,696 samples are included in the boxplot for each method. (**d**) The dynamics fitting on MAML3, ANXA4 and GSTZ1. For each gene, four plots are displayed in a 2x2 layout: the u-t, s-t, u-s, and alpha-u plots. (**e**) The unspliced-spliced phase portrait fitting on MAML3, ANXA4 and GSTZ1 obtained by four baseline RNA velocity approaches, including scVelo, Dynamo, UniTVelo and cellDancer. (**f**) The dynamics fitting in transcription-unspliced-spliced 3D phase portrait for MAML3, ANXA4 and GSTZ1.

Next, we show the dynamics for individual genes and demonstrate how the TSvelo model aids gene dynamics analysis by incorporating both transcriptional and splicing information. Using MAML3, ANXA4 and GSTZ1 as examples, **Fig. 2d** present the results of the TSvelo model on these genes, and **Fig. 2e** present the results of baseline approaches. Many previous approaches assume that the unspliced-spliced phase portrait exhibits an almond-shaped distribution[34]which may not actually hold in the data. For MAML3, while the unspliced-spliced distribution predominantly follows the almond-shaped pattern, some cells, specifically Alpha cells (blue) and Beta cells (light blue), overlap with Ngn3 high EP cells (yellow), indicating that these cell types cannot be distinctly separated using only the unspliced-spliced 2D phase portrait. Nevertheless, TSvelo can infer transcription representation from the expression of multiple transcription factors, which enables the model to distinguish these cell types (**Fig. 2d** and **Fig. 2f**). For ANXA4, the expression pattern exhibits an initial decrease followed by an increase (in red), a dynamic that is not easily captured in the unspliced-spliced phase portrait by previous approaches. By comparison, TSvelo can still accurately predicts the abundance of unspliced spliced RNA. The visualization of dynamics fitting on more genes are provided at **Fig. S3** in **Supplementary Information**.

TSvelo can better predict RNA velocity on gastrulation erythroid dataset
We applied TSvelo to the gastrulation erythroid dataset, which is derived from the transcriptional profile of mouse embryos [35] and has been used in previous RNA velocity studies. This dataset primarily describes the differentiation process from blood progenitors to erythroid cells. First, using Gene Ontology (GO) term enrichment analysis (**Fig. 3a**), we find that the velocity genes selected through TSvelo's preprocessing strategies are mostly enriched in heme biosynthetic process (GO:0006783) and porphyrin-containing compound biosynthetic process (GO:0006779), both of which are related to the erythropoiesis process.

Figure 3. Results on gastrulation erythroid dataset. (**a**) The GO terms which are mostly enriched in the selected velocity genes of TSvelo. (**b**) The pseudotime learned with TSvelo. (**c**) The Stream-plot for visualization the RNA velocity inferred by TSvelo. (**d-f**) The quantitative comparison between TSvelo and multiple baseline approaches in terms of velocity consistency (d), in-cluster coherence (e), and cross boundary direction correctness (f). The down, central and up hinges correspond to the first quartile, median value and third quartile, respectively. The whiskers extend to $1.5\times$ the interquartile range of the distribution from the hinge. 9,815 samples are included in the boxplot for each method. (**g**) The dynamics fitting on HSP90AB1 obtained by TSvelo. Four plots are displayed in a 2x2 layout: the u-t, s-t, u-s, and alpha-u plots, where u, a and alpha mean the abundance of unspliced mRNA, the abundance of spliced mRNA, and the learned transcriptional representation, respectively. (**i**) The phase portrait fitting on HSP90AB1 obtained by baseline approaches. (**i**) The dynamics fitting on RPS26 obtained by TSvelo. (**j**) The TFs with the highest ranked weights as identified by TSvelo. (**k**) Weights for KLF1's targets, with the highest absolute weight. (**l**) Temporal dynamics of KLF1 and its target genes

with the highest weights along pseudotime, which includes HBA-X, ALAS2 and GYPA.

After applying TSvelo, both the inferred pseudotime (**Fig. 3b**) and the velocity stream plot (**Fig. 3c**) accurately reflect the cell differentiation process. We also compare TSvelo with previous approaches. The results, shown in **Fig. 3d-f**, demonstrate that TSvelo achieves the highest velocity consistency, the highest in-cluster coherence, and the highest cross-boundary correctness, showing that the high-dimensional dynamics predicted by TSvelo best capture the underlying biological processes.

We next analyze TSvelo modeling at the gene level in **Fig. 3g-i**. For genes exhibiting higher noise or more complex patterns, TSvelo demonstrates its robustness in dynamic fitting by integrating both transcriptional and splicing signals. For the gene HSP90AB1, TSvelo accurately captures the dynamics (**Fig. 3g**), whose pattern can not be well modeled by previous approaches (**Fig. 3h**). For genes such as RPS26[36], which have critical roles in the development in blood progenitors to erythroid, the unspliced-spliced data is so noisy that cells of different types overlap in phase portrait. TSvelo can still accurately captures the gene dynamics and reveals differences in transcription rates (Alpha) across cell types (**Fig. 3i**). In contrast, methods that rely solely on unspliced-spliced data from individual genes, such as scVelo, fail to accurately fit these dynamics and can not identify them as velocity genes, which further validates TSvelo's performance in complex scenarios. The visualization of dynamics fitting on more genes are provided at **Fig. S5** in **Supplementary Information**.

From the TF-target weight matrices learned by TSvelo, we can extract key regulatory relationships. First, we calculate the mean absolute weight of each TF across its interactions with target genes. The TFs with the highest mean absolute weights are ranked and presented in **Fig. 3j**. Notably, KLF1, a critical TF in blood progenitors[37], is assigned the highest weight. We then examine the target genes of KLF1, highlighting those with the highest weights (**Fig. 3k**), among which HBA-X[38], ALAS2[39], and GYPA[40] have been previously identified as key genes in erythroid differentiation and are known to be regulated by KLF1. A time-delay pattern is also observed between KLF1 and its target genes (**Fig. 3l**), indicating that KLF1 expression increases first, followed by a corresponding upregulation of its target genes.

TSvelo can accurately fit cell fate and gene dynamics on mouse brain data
We apply TSvelo to a 10x multi-omics dataset from the embryonic mouse brain, which includes both Assay for Transposase-Accessible Chromatin with sequencing (ATAC-Seq) [41] and scRNA-seq data. This dataset was previously used in the Multivelo study, which introduced an RNA velocity model designed for multi-omics

datasets to capture the dynamics between chromatin accessibility and unspliced mRNA[19]. In this dataset, Radial Glia (RG) cells located in the outer subventricular zone give rise to neurons, astrocytes, and oligodendrocytes. During cortical development, neurons migrate in an inside-out manner, with younger cells moving toward the upper layers of the cortex, while older cells settle in the deeper layers. Additionally, RG cells are capable of producing intermediate progenitor cells (IPCs), which serve as neural stem cells and generate a variety of mature excitatory neurons within the cortical layers.

Figure 4. Results on mouse brain dataset. (**a**) The velocity stream inferred by Multivelo. (**b**) The Stream-plot for visualization the RNA velocity inferred by TSvelo. (**c**) The pseudotime learned with TSvelo. (**d-f**) The dynamics fitting on gene MEIS2 (d), BASP1 (e) and MSI2 (f) by both Multivelo and TSvelo. In each panel, the

leftmost plot shows the phase portrait fitting of Multivelo, and the next two columns
show TSvelo's results. The rightmost plot shows the learned transcriptional rate (in
green), unspliced abundance (in blue), and spliced abundance (in red) along the
pseudotime. Since the transcriptional rate is calculated for each individual cell, we
apply a Generalized Additive Model (GAM) to transcriptional representation across
cells along the pseudotime and present GAM-fitting results to better visualize its
trends in these plots.

Multivelo is chosen as the baseline on this study since it connects the regulation signal
and RNA velocity in ATAC-unspliced-spliced space., and TSvelo uses the same data
processed be Multivelo for a fair comparison. As shown in **Fig. 4a** and **Fig. 4b**, the
velocity stream obtained by TSvelo is smoother than Multivelo and can correctly
reflect the differentiation from RG (in red) to IPCs (in brown). The pseudotime
inferred by TSvelo is also coherent with the whole cell development process (**Fig. 4c**).

We next further analyze the details in the phase portrait fitting. On MEIS2 (**Fig. 4d**),
both Multivelo and TSvelo successfully models its dynamics, which follows a clear
pattern on the unspliced-spliced phase portrait. However, due to the high noise and
sparsity inherent in ATAC-seq data, Multivelo encounters difficulty on genes that
show significant overlap between different cell types in the unspliced–spliced phase
portrait. For example, in the case of BASP1 (**Fig. 4e**), Multivelo incorrectly models
their expression patterns as monotonically increasing, failing to capture the true
dynamics, which involve initial upregulation followed by downregulation. In contrast,
TSvelo successfully captures the correct gene dynamics and identifies the delay from
transcription to the unspliced mRNA and also the delay from unspliced to spliced
mRNA (The rightmost plots in **Fig. 4e**). MSI2, which has been reported to be highly
expressed in neural stem/progenitor cells[42], is accurately modeled by TSvelo,
exhibiting a decreasing expression pattern during the differentiation process. By
comparison, Multivelo fails to capture the correct trend at the initial stage (**Fig. 4f**).
We also show the learned transcriptional rate α as well as the predicted dynamics of
unspliced and spliced RNA along pseudotime t, which clearly illustrate the time
delays between transcription to unspliced and unspliced to spliced RNAs. The
visualization of dynamics fitting on more genes are provided at **Fig. S6** in
Supplementary Information. Given that transcriptional regulation involves the
binding of TF and chromatin accessibility as measured by ATAC, TSvelo shows that it
is possible to model transcriptional signals using only scRNA-Seq data.

TSvelo can accurately infer cell fate and model lineage-specific gene dynamics
for multi-lineage task
Given that multi-lineage differentiation is a common phenomenon in larger and larger
scRNA-seq datasets, developing the RNA velocity model that can handle such
complexity is essential. Because TSvelo demonstrates robust performance across
various tasks, its applicability can be extended to more complex situations, such as

scRNA-seq datasets where cells differentiate into multiple fates. We apply TSvelo to a multi-lineage dentate gyrus scRNA-seq dataset, which captures the differentiation process from neural blast cells to various cell types[43]. During preprocessing, the velocity genes selected by TSvelo are enriched in GO terms related to neural development, such as axonogenesis (GO:0007409), axon guidance (GO:0007411) and axon development (GO:0061564) (**Fig. 5a**), which aligns with the biological processes described by the dataset.

Figure 5. Results on the multi-lineage dentate gyrus dataset. (**a**) The GO terms enriched in the selected velocity genes of TSvelo. (**b**) The pseudotime learned with TSvelo. (**c**) The Stream-plot for visualization the RNA velocity inferred by TSvelo. Three lineages are detected, which are Granule lineage, CA lineage and glial lineage.

(**d**) The velocity stream inferred by scVelo. (**e**) The velocity stream inferred by cellDancer. (**f-h**) The dynamics modeling of TSvelo on three axonogenesis-related genes, ANK3 (f), MAP1B (g) and SLC1A2 (h). In the leftmost plot of each panel, the lines represent the predicted spliced abundance across all lineages, with the color indicating the cell types most associated with each pseudotime point along the corresponding lineage. Additionally, expression data for each lineage are shown as translucent points. The remaining plots in each panel display the dynamics of the learned transcriptional rate (in green), unspliced abundance (in blue), and spliced abundance (in red) along pseudotime for each lineage. The transcriptional representation in these plots is also processed using GAM fitting.

TSvelo could correctly identify three lineages from this data. Details about the lineage segmentation are provided at **Methods** and **Fig. S7** in **Supplementary Information**. After applying the TSvelo model to each lineage and combine the results, the inferred pseudotime and velocity stream plots (**Fig. 5b** and **Fig. 5c**) align with the ground truth differentiation trajectory well, which begins with neural blast cells. We also compare TSvelo to baseline methods, including scVelo and cellDancer. Notably, cellDancer is designed to handle such multi-lineage data by modeling cell- and gene-specific transcriptional rates, splicing rates, and degradation rates using neural networks. As shown in **Fig. 5d** and **Fig. 5e**, scVelo struggles with this multi-lineage dataset, and cellDancer fails to correctly capture the trajectory of immature granule cells (colored in purple).

Next, we analyze gene expression patterns across different lineages. TSvelo provides inferred dynamics along each lineage, revealing that the expression of ANK3 in the granule and Cornu Ammonis (CA) lineages follows an increasing and then decreasing pattern. In contrast, the expression of ANK3 in the glial lineage decreases to zero (**Fig. 5f**). The similar pattern is also observed for other genes, such as MAP1B (**Fig. 5g**). Both ANK3[44] and MAP1B[45] are associated with GO terms axonogenesis and axon guidance. This observation is consistent with the fact that the granule and Cornu Ammonis[46] predominantly consist of neurons, where axons are a critical component. In contrast, glial cells, such as astrocytes, typically lack axons, providing a potential explanation for these expression patterns. SLC1A2 exhibits a distinct expression pattern, showing a significant increase in the glial lineage and a decrease in the granule and CA lineages. This pattern aligns with the observation that astrocytes are the primary cell type expressing SLC1A2[47]. Details about dynamics fitting for genes on each lineage are provided at **Fig. S8** in **Supplementary Information**.

TSvelo can accurately infer cell fate and model lineage-specific gene dynamics on Larry dataset

The Lineage and RNA Recovery (LARRY) method utilizes barcoded hematopoietic cells to trace both cell lineage and gene expression over time. LARRY has been

372 successfully employed to track the in vitro differentiation of human blood cells,
373 accurately capturing lineage trajectories and cell fates[48].
374

Fig 6. Results on the LARRY dataset. (**a**) The Leiden clustering on LARRY. (**b**) The initial Leiden cluster detection in preprocessing. (**c**) The pseudotime learned with TSvelo. (**d**) The Stream-plot for visualization the RNA velocity inferred by TSvelo. (**e**) The GO terms enriched in the selected velocity genes of TSvelo. (**f**) The dynamics modeling of TSvelo on four genes related to neutrophil development, which are PYGL, MS4A3, CLEC12A and LTA4H. The lines represent the predicted spliced abundance across all lineages, with the color indicating the cell types most strongly associated with each pseudotime point along the corresponding lineage. (**g**) The dynamics modeling of PYGL, MS4A3, CLEC12A and LTA4H on the neutrophil lineage.

On this LARRY dataset which encompass a total of 49,302 cells on multiple lineages, TSvelo effectively detects the initial Leiden clusters based on the unspliced-to-spliced delay (**Fig. 6a** and **Fig. 6b**) and separate its lineages. Furthermore, **Fig. 6c** and **Fig. 6d**

show that the pseudotime and velocity stream inferred by TSvelo can accurately capture the differentiation process, progressing from undifferentiated cells to distinct cell fates. The velocity genes selected by TSvelo are significantly enriched in processes related to neutrophil biology, such as neutrophil degranulation (GO:0043312), neutrophil activation in immune response (GO:0002283), and neutrophil-mediated immunity (GO:0002446).

We further analyzed the expression patterns of genes associated with neutrophil degranulation, focusing specifically on the neutrophil lineage. Using four representative genes as examples (**Figs. 6f, g**), TSvelo can help observe the significant variation in the expression patterns across these genes. PYGL, which has been widely reported as a key gene in neutrophil[49], is almost exclusively expressed in neutrophil cells and exhibits an increasing expression pattern during neutrophil differentiation. In contrast, MS4A3 was examined across all lineages, and using TSvelo, we observed that its expression initially increases and then decreases in both the neutrophil and monocyte lineages. This pattern is consistent with prior studies identifying MS4A3 as a gene specifically expressed by granulocyte-monocyte progenitors[42]. Similarly, CLEC12A, another gene associated with neutrophil degranulation, shows a comparable expression trajectory during neutrophil differentiation. Previous research has indicated that CLEC12A expression is highest in granulocyte-macrophage progenitors[50]. LTA4H exhibits a pattern similar to that of CLEC12A, suggesting it may play a significant role in the early stages of neutrophil differentiation. These findings further confirm the utility of TSvelo for gene-level analysis in multi-lineage differentiation datasets.

Discussion

RNA velocity, utilizing unspliced/spliced data, has become a widely adopted concept for predicting cell fate and modeling gene dynamics. While several RNA velocity models have been proposed, most of they are based on phase portrait fitting in the unspliced-spliced space. However, due to the limited information and high noise inherent in the splicing data of individual genes, the short time delays between unspliced and spliced abundance, and the challenges posed by large-scale datasets with complex processes, most genes cannot be accurately modeled by previous RNA velocity methods. This limitation reduces the reliability and robustness of downstream analyses.

Gene expression is a complex biological process within cells, involving multiple regulatory mechanisms from DNA to the matured RNA. In this process, transcription and splicing are two crucial steps to determine the final gene expression. Due to the mathematical complexity, previous methods cannot jointly model gene regulation, transcription and splicing. TSvelo comprehensively models the cascade of the whole process using an interpretable ODE framework, allowing for learning dynamics of all

velocity genes simultaneously. TSvelo could accurately model gene dynamics, predict cell fate, detect the key regulatory relations, and handle multi-lineage datasets. Results on six scRNA-seq datasets demonstrate that TSvelo is a valuable approach for RNA velocity modeling (results on pons dataset are provided in **Fig. S1** in **Supplementary Information**).

Looking ahead, TSvelo could be further improved by more accurately modeling the splicing and degradation process. Currently, TSvelo assumes a constant gene-specific splicing and degradation rates. Since splicing factors also play significant roles in regulating other genes during the splicing process[51], and there are still complex mechanisms of mRNA degradation[52], further exploration could enhance velocity's modelling and bring new biological insights in future.

Methods

Preprocessing for scRNA-seq data
The unspliced and spliced RNA abundances are preprocessed with multiple steps, which includes highly variable genes (HVGs) selection, normalization, log transformation, k-nearest neighbor (KNN) smoothing, and clustering. Following the data preprocessing procedures outlined in scVelo[6], we first select the top 2,000 highly variable genes (HVGs) and normalize their expression profiles by dividing by the total counts in each cell. A nearest-neighbor graph (with 30 neighbors by default) was constructed based on Euclidean distances in principal component analysis space (with 30 principal components by default) on log-transformed gene expression data. Subsequently, we compute the first- and second-order moments (mean and uncentered variance) for each cell across its nearest neighbors. These steps are performed by using scvelo.pp.filter_and_normalize() and scvelo.pp.moments().

Acquiring Prior Knowledge of Gene Regulatory Relations
To construct a prior gene network for the selected HVGs, we utilize TF-target annotations from the ENCODE[53] and ChEA[54] databases. If a regulatory interaction between a transcription factor (TF) and its target gene is identified in either of these two databases, the TF is included in the set of regulators for modeling the dynamic behavior of the target gene.

Velocity genes selection
Previous studies[6] have demonstrated that only a subset of genes, termed "velocity genes," can be accurately fitted in the unspliced-spliced phase portrait by RNA velocity models. Inspired by the notion that velocity genes provide high-quality data for dynamic modeling in the phase portrait, we propose to select velocity genes during the preprocessing step. The velocity genes are selected based on the premise that they

exhibit clear dynamics in the unspliced-spliced phase portrait, which is necessary for
precise modeling splicing dynamics. The selection is based on the similarity between
cell-cell neighborhood relationships in UMAP space and those observed in
gene-specific unspliced-spliced phase portraits. In detail, we first compute the
cell-cell neighborhood graph, $Graph$, using scvelo.pp.neighbors() on all highly
variable genes (HVGs). Next, we construct an anndata object $adata_g$ for each gene,
which contains only the unspliced and spliced expression data of that gene. We then
apply scvelo.pp.neighbors() to each $adata_g$ to calculate the gene-specific
neighborhood graph, denoted as $Graph_g$. Subsequently, we compute the similarity
between $Graph$ and each $Graph_g$. The top 100 genes with the highest similarity are
selected as velocity genes. These genes are characterized by phase portraits that
exhibit a structure similar to that of the UMAP space, thereby enhancing the
separation of cells from different types. As a result, the splicing dynamics of these
velocity genes are more likely to be captured by RNA velocity models.
491
492

lineages segmentation and pseudotime initialization
494 Based on the normalized gene expression data, we perform Leiden clustering using
495 scanpy.tl.leiden() with a low resolution (default resolution = 0.1). These Leiden
496 clusters are utilized to identify lineages and determine the differentiation direction for
497 each lineage. Subsequently, we apply PAGA (scanpy.tl.paga()) to assess the
498 connectivity between Leiden groups.
499

By setting one Leiden group as the initial state, we infer pseudotime using diffusion
pseudotime (DPT) with scanpy.tl.dpt(). To detect lineages, we start from the selected
initial Leiden cluster and filter the PAGA graph by applying a threshold (default =
0.02). Edges with weights below this threshold are removed, leaving only the shortest
path from the initial state to each group. The remaining paths are considered as the
detected lineages.
506

Given a lineage, TSvelo could detect the orientation of differentiation along it, which
is based on the fact that unspliced RNA precedes spliced RNA in the expression
pattern[55]. For each gene, TSvelo computes the Spearman correlation between
unspliced and spliced abundance across different moving steps along the DPT. The
time number of moving steps along unspliced to spliced expression, which maximizes
the Spearman correlation, is considered the U-to-S delay. **Fig. 1a** illustrates how
U-to-S delay can be used to validate the accuracy of pseudotime, with EML5 serving
as an example. Then the U-to-S delay for that lineage is calculated as the mean U-to-S
delay for all genes on it. And the average U-to-S delay across all lineages provides the
overall U-to-S delay for the initial Leiden cluster. Finally, the initial Leiden cluster
with the highest U-to-S delay is considered the correct initiation. The DPT under this
condition is used to initialize pseudotime for TSvelo. If multiple lineages are detected,
TSvelo will model each lineage independently and subsequently merge them in the
final step. A detailed explanation with example illustrating how lineages are

segmented is provided in the **Fig. S7** in **Supplementary Information**.
522
523
524
525   **The ODE model in TSvelo**
526   Suppose u_g and s_g are the abundance of unspliced and spliced RNA, and
527    $\alpha_g(t), \beta_g, \gamma_g$ are the transcription rate, splicing rate and degradation rate,
528   respectively. The dynamics of a velocity gene g is modeled as:

$$\frac{du_g(t)}{dt} = \alpha_g(t) - \beta_g u_g(t) \#(4)$$

$$\frac{ds_g(t)}{dt} = \beta_g u_g(t) - \gamma_g s_g(t) \#(5)$$

529   We assume that the gene and cell-specific transcriptional rate $\alpha_g(t)$ is influence
530   by the expression of Transcriptional Factors (TFs), while β_g and γ_g are
531   gene-specific constant parameters. Considering the wide usage of linear models for
532   gene-relations in previous studies[26,30-33], we model $\alpha_g(t)$ as

$$\alpha_g(t) = ReLU\left(\sum_{i \in TFs(g)} w_{gi} * s_i(t) \right) \#(6)$$

In order to include more TFs in the modeling, we reserve those TFs that are not
selected as velocity genes. These TFs are excluded from the velocity genes selection
because their unspliced-spliced expression does not provide sufficient information,
primarily due to high noise in the unspliced RNA. Consequently, we incorporate these
TFs, whose spliced abundance is denoted as s', into the TSvelo by directly modeling
the process between transcriptional signal to mature RNA,

$$\frac{ds'_g(t)}{dt} = \alpha_g(t) - \gamma_g s'_g(t) \#(7)$$

539   As a result, we can get the ODE model with parameters matrix A:

$$\begin{bmatrix} dU/dt \\ dS/dt \\ dS'/dt \end{bmatrix} = A \begin{bmatrix} U \\ S \\ S' \end{bmatrix} = \begin{bmatrix} -B & W & W' \\ B & -\Gamma & 0 \\ 0 & W & W'-\Gamma' \end{bmatrix} \begin{bmatrix} U \\ S \\ S' \end{bmatrix} \#(8)$$

$U = [u_1, u_2 \ldots, u_G]^T \in R^G, S = [s_1, s_2 \ldots, s_G]^T \in R^G, \ S' = [s_{G+1}, s_{G+2} \ldots, s_{G+K}]^T \in$
R^K, G is the number of velocity genes and K is the number of additional TFs which
are not selected as velocity genes. $B \in R^{G \times G}$ and $\Gamma \in R^{G \times G}$ are the diagonal
matrices consist of splicing rates $[\beta_1, \beta_2 \ldots, \beta_G]$ and degradation rates
$[\gamma_1, \gamma_2 \ldots, \gamma_G]$ for velocity genes, respectively. $\Gamma' \in R^{K \times K}$ is the diagonal matrices
consist of degradation rates $[\gamma_{G+1}, \gamma_{G+2} \ldots, \gamma_{G+K}]$ for these additional TFs.
$W \in R^{G \times G}$ and $W' \in R^{K \times K}$ denote the matrices representing the TF-target
relationships for the TFs selected as velocity genes and those not selected as
velocity genes, respectively. Using these gene-gene weight matrices, TSvelo can
model the gene- and cell-specific transcriptional rate α_g. The whole parameters
matrix $A \in R^{(2G+K) \times (2G+K)}$
551
552

Optimizing global time and Neural ODE in EM framework

In TSvelo, the dynamics described in **Eq. 7** are implemented using a neural ordinary differential equation (ODE) model. Given an initial state and the parameters in matrix A, the neural ODE model can compute the values of U and S at any time step. This approach eliminates the need for an analytical solution for unspliced and spliced expression, enhancing the flexibility of TSvelo. By default, the number of time steps in the ODE is set to 1,000. The ReLU activation function is applied to the parameters α_g, β_g and γ_g, in order to prevent them from taking negative values. Notably, TSvelo does not incorporate deep neural networks or encoders. Since TSvelo models all genes simultaneously, we normalize the unspliced and spliced abundances of each gene by its standard deviation before inputting the data into the Neural ODE model to ensure that the influence of each gene is balanced.

TSvelo optimize the pseudotime t and parameters matrix A in ODE iteratively using an Expectation-Maximization (EM) approach. The maximum number of iterations is set to 30 by default, with an early stopping criterion applied. Given the parameters in ODE system, we can compute the predicted values of unspliced (for velocity genes) and spliced (for velocity genes and additional TFs) at all time steps, denoted as $\widehat{U}(t\_step; A)$, $\hat{S}(t\_step; A)$ and $\widehat{S'}(t\_step; A)$, where $t_{step} \in \{0,1,2 \dots ,999\}$. Given the time steps assignment t_c for each cell c, we can get the expected values for velocity genes at all cells as $\widehat{U}(t_c; A)$, $\hat{S}(t_c; A)$ and $\widehat{S'}(t_c; A)$. The loss function used in EM algorithm is the mean squared error between the expected value with the observed data.

$$Loss = \frac{1}{c}\sum_c [(\widehat{U}(t_c; A) - U_c)^2 + (\hat{S}(t_c; A) - S_c)^2 + (\widehat{S'}(t_c; A) - S'_c)^2] \# (9)$$

The EM algorithm will iteratively update the time assignment and parameters in neural ODE by minimizing the above loss function.

In the E-step, TSvelo updates the time step assigned to each cell using grid search with time steps range $t_{step} \in \{0,1,2 \dots ,999\}$, which is achieved by finding the minimum distance between the observed unspliced-spliced expression and the model prediction ($\widehat{U}$ and $\hat{S}$) at all time steps.

$$min_{t_c} \frac{1}{c}\sum_c [(\widehat{U}(t_c; A) - U_c)^2 + (\hat{S}(t_c; A) - S_c)^2 + (\widehat{S'}(t_c; A) - S'_c)^2] \# (10)$$

In the M-step, the parameter matrix A is updated using gradient descent within the neural ODE model to minimize the loss function. Here we use the prior gene regulation knowledge to constraint the weight matrix W and W' in A. Only If gene i is annotated as a TF for gene j in the ENCODE or ChEA databases, the corresponding w_{ij} could be learnable. Otherwise w_{ij} is fixed at zero. The usage of prior knowledge could keep the matrix sparse and avoid overfitting. The neural ODE module is implemented using the torchdiffeq package and trained with the Adam

optimizer, a learning rate of 0.05, and a maximum of 500 epochs, incorporating an early stopping criterion to prevent overfitting.

$$min_A \frac{1}{c}\sum_c [(\widehat{U}(t_c; A) - U_c)^2 + (\widehat{S}(t_c; A) - S_c)^2 + (\widehat{S'}(t_c; A) - S'_c)^2] \#(11)$$

By performing the EM optimization, TSvelo could fit the high-dimensional unspliced-spliced data across multiple genes well, and learn the global pseudotime for each cell. After the training procedure, the RNA velocity could be calculated using those parameters in matrix A, which could be further used for cell fate prediction.

Metrics for evaluating
(1) Velocity consistency. We used the scvelo.velocity_confidence() function from scVelo to evaluate velocity consistency, interpreting the results as a measure of how consistent velocities are within neighboring cells.
(2) Cross-boundary direction correctness. Cross-boundary direction correctness assesses the accuracy of transitions from a source cluster to a target cluster by examining the boundary cells, and requires ground truth annotations. We directly run the function unitvelo.evaluate() provided in UniTVelo[7] to obtain the Cross-boundary direction correctness.
(3) Within-cluster velocity coherence. Within-cluster velocity coherence measures the coherence of velocities within a single cluster using a cosine similarity score between cell velocities. We applied the function unitvelo.evaluate() provided by UniTVelo[7] to directly compute the within-cluster velocity coherence.

GO term enrichment analysis
Gene Ontology (GO) terms enrichment analysis is a widely used method for identifying biological processes, molecular functions, or cellular components that are overrepresented in a given set of genes compared to a background set. Here we use the biological processes enrichment analysis in GO terms database to explore the associated biological roles of those selected velocity genes. To perform the GO term enrichment analysis, we utilized the "gseapy.enrichr()" function in Python with using Fisher's exact test by default.

Data Availability Statement

The pancreatic endocrinogenesis dataset[56] comprises the single-cell RNA-seq (10X) data of pancreatic epithelial and Ngn3-Venus fusion cells sampled from mouse embryonic day 15.5, which could be loaded using scVelo's package scvelo.datasets.pancreas().

The gastrulation erythroid dataset, which is selected from the transcriptional profiles of mouse embryos[35], which could be loaded using scVelo's package scvelo.datasets.pancreas().

10x embryonic mouse brain dataset is provided at the 10x website at https://www.10xgenomics.com/resources/datasets/fresh-embryonic-e-18-mouse-brain-5-k-1-standard-1-0-0. The data preprocessed by Multivelo is utilized in this study, (https://multivelo.readthedocs.io/en/latest/MultiVelo_Fig2.html).

The dentate gyrus neurogenesis data is available at http://pklab.med.harvard.edu/velocyto/DentateGyrus/DentateGyrus.loom

The LARRY dataset has been shared by pyrovelocity, which could be accessed at https://figshare.com/articles/dataset/larry_invitro_adata_sub_raw_h5ad/20780344.

The raw data of Hindbrain (pons) of adolescent mice is from https://pklab.med.harvard.edu/ruslan/velocity/oligos/.

The ENCODE TF-target database[53] website: https://maayanlab.cloud/Harmonizome/dataset/ENCODE+Transcription+Factor+Targets.

The ChEA TF-target database[54] website: https://maayanlab.cloud/Harmonizome/dataset/CHEA+Transcription+Factor+Targets.

Code Availability Statement

TSvelo is implemented in Python. The source code can be downloaded from the GitHub repository, https://github.com/lijc0804/TSvelo.

Acknowledgement

This work was supported by National Natural Science Foundation of China (No. 62103262 to Y.Y.), and the National Key R&D Program of China (2023YFF1204500 to Y.Y.).

Competing Interests Statement

The authors declare no competing interests.

1   Wolf, F. A. *et al.* PAGA: graph abstraction reconciles clustering with trajectory inference through a topology preserving map of single cells. *Genome biology* **20**, 1-9 (2019).

2   Qiu, X. *et al.* Reversed graph embedding resolves complex single-cell trajectories.

668   *Nature methods* **14**, 979-982 (2017).

669 3  Street, K. *et al.* Slingshot: cell lineage and pseudotime inference for single-cell

670   transcriptomics. *BMC genomics* **19**, 1-16 (2018).

671 4  Setty, M. *et al.* Characterization of cell fate probabilities in single-cell data with Palantir.

672   *Nature biotechnology* **37**, 451-460 (2019).

673 5  La Manno, G. *et al.* RNA velocity of single cells. *Nature* **560**, 494-498 (2018).

674 6  Bergen, V., Lange, M., Peidli, S., Wolf, F. A. & Theis, F. J. Generalizing RNA velocity

675   to transient cell states through dynamical modeling. *Nature biotechnology* **38**,

676   1408-1414 (2020).

677 7  Gao, M., Qiao, C. & Huang, Y. UniTVelo: temporally unified RNA velocity reinforces

678   single-cell trajectory inference. *Nature Communications* **13**, 6586 (2022).

679 8  Gayoso, A. *et al.* Deep generative modeling of transcriptional dynamics for RNA

680   velocity analysis in single cells. *Nature methods* **21**, 50-59 (2024).

681 9  Gu, Y., Blaauw, D. T. & Welch, J. in *International Conference on Machine Learning.*

682   7887-7901 (PMLR).

683 10  Qin, Q., Bingham, E., La Manno, G., Langenau, D. M. & Pinello, L. Pyro-Velocity:

684   Probabilistic RNA Velocity inference from single-cell data. *bioRxiv* (2022).

685 11  Li, S. *et al.* A relay velocity model infers cell-dependent RNA velocity. *Nature*

686   *biotechnology*, 1-10 (2023).

687 12  Cui, H. *et al.* DeepVelo: deep learning extends RNA velocity to multi-lineage systems

688   with cell-specific kinetics. *Genome Biology* **25**, 27 (2024).

689 13  Farrell, S., Mani, M. & Goyal, S. Inferring single-cell transcriptomic dynamics with

structured latent gene expression dynamics. *Cell Reports Methods* **3** (2023).

14   Qiu, X. *et al.* Mapping transcriptomic vector fields of single cells. *Cell* **185**, 690-711.

e645 (2022).

15   Ma, A., McDermaid, A., Xu, J., Chang, Y. & Ma, Q. Integrative methods and practical

challenges for single-cell multi-omics. *Trends in biotechnology* **38**, 1007-1022 (2020).

16   Subramanian, I., Verma, S., Kumar, S., Jere, A. & Anamika, K. Multi-omics data

integration, interpretation, and its application. *Bioinformatics and biology insights* **14**,

1177932219899051 (2020).

17   Gorin, G., Svensson, V. & Pachter, L. Protein velocity and acceleration from single-cell

multiomics experiments. *Genome biology* **21**, 1-6 (2020).

18   Li, C., Virgilio, M. C., Collins, K. L. & Welch, J. D. Multi-omic single-cell velocity models

epigenome–transcriptome interactions and improves cell fate prediction. *Nature*

Biotechnology **41**, 387-398 (2023).

19   Riba, A. *et al.* Cell cycle gene regulation dynamics revealed by RNA velocity and

deep-learning. *Nature communications* **13**, 1-13 (2022).

20   Lederer, A. R. *et al.* Statistical inference with a manifold-constrained RNA velocity

model uncovers cell cycle speed modulations. *Nature Methods*, 1-16 (2024).

21   Zhou, P., Bocci, F., Li, T. & Nie, Q. Spatial transition tensor of single cells. *Nature*

Methods, 1-10 (2024).

22   Abdelaal, T. *et al.* SIRV: Spatial inference of RNA velocity at the single-cell resolution.

NAR genomics and bioinformatics **6**, lqae100 (2024).

23   Liu, R., Pisco, A. O., Braun, E., Linnarsson, S. & Zou, J. Dynamical Systems Model of

RNA Velocity Improves Inference of Single-cell Trajectory, Pseudo-time and Gene

Regulation. *Journal of Molecular Biology*, 167606 (2022).

24   Gorin, G., Fang, M., Chari, T. & Pachter, L. RNA velocity unraveled. *PLOS*

Computational Biology **18**, e1010492 (2022).

25   Soneson, C., Srivastava, A., Patro, R. & Stadler, M. B. Preprocessing choices affect

RNA velocity results for droplet scRNA-seq data. *PLoS computational biology* **17**,

e1008585 (2021).

26   Li, J., Pan, X., Yuan, Y. & Shen, H.-B. TFvelo: gene regulation inspired RNA velocity

estimation. *Nature Communications* **15**, 1387 (2024).

27   Bergen, V., Soldatov, R. A., Kharchenko, P. V. & Theis, F. J. RNA velocity—current

challenges and future perspectives. *Molecular systems biology* **17**, e10282 (2021).

28   Hossain, I., Fanfani, V., Fischer, J., Quackenbush, J. & Burkholz, R. Biologically

informed NeuralODEs for genome-wide regulatory dynamics. *Genome Biology* **25**,

127 (2024).

29   Burdziak, C. *et al.* scKINETICS: inference of regulatory velocity with single-cell

transcriptomics data. *Bioinformatics* **39**, i394-i403 (2023).

30   Tran, A., Yang, P., Yang, J. Y. & Ormerod, J. T. scREMOTE: Using multimodal single

cell data to predict regulatory gene relationships and to build a computational cell

reprogramming model. *NAR genomics and bioinformatics* **4**, lqac023 (2022).

31   Song, Q., Ruffalo, M. & Bar-Joseph, Z. Using single cell atlas data to reconstruct

regulatory networks. *Nucleic Acids Research* **51**, e38-e38 (2023).

32   Liu, Y.-Y., Slotine, J.-J. & Barabási, A.-L. Controllability of complex networks. *nature*

473, 167-173 (2011).

33 Wang, L. *et al.* Dictys: dynamic gene regulatory network dissects developmental

736 continuum with single-cell multiomics. *Nature Methods* **20**, 1368-1378 (2023).

34 Weiler, P., Van den Berge, K., Street, K. & Tiberi, S. in *Single Cell Transcriptomics:*

738 *Methods and Protocols* 269-292 (Springer, 2022).

35 Pijuan-Sala, B. *et al.* A single-cell molecular map of mouse gastrulation and early

740 organogenesis. *Nature* **566**, 490-495 (2019).

36 Liu, Y. & Karlsson, S. Perspectives of current understanding and therapeutics of

742 Diamond-Blackfan anemia. *Leukemia* **38**, 1-9 (2024).

37 Mukherjee, K. & Bieker, J. J. EKLF/Klf1 regulates erythroid transcription by its

744 pioneering activity and selective control of RNA Pol II pause-release. *Cell reports* **41**

745 (2022).

38 Perseu, L. *et al.* KLF1 gene mutations cause borderline HbA2. *Blood, The Journal of*

747 *the American Society of Hematology* **118**, 4454-4458 (2011).

39 Singleton, B. K., Burton, N. M., Green, C., Brady, R. L. & Anstee, D. J. Mutations in

749 EKLF/KLF1 form the molecular basis of the rare blood group In (Lu) phenotype. *Blood,*

750 *The Journal of the American Society of Hematology* **112**, 2081-2088 (2008).

40 Kuvardina, O. N. *et al.* RUNX1 represses the erythroid gene expression program

752 during megakaryocytic differentiation. *Blood, The Journal of the American Society of*

753 *Hematology* **125**, 3570-3579 (2015).

41 Buenrostro, J. D., Giresi, P. G., Zaba, L. C., Chang, H. Y. & Greenleaf, W. J.

Transposition of native chromatin for fast and sensitive epigenomic profiling of open

chromatin, DNA-binding proteins and nucleosome position. *Nature methods* **10**,

1213-1218 (2013).

42   Liu, Z. *et al.* Fate mapping via Ms4a3-expression history traces monocyte-derived

cells. *Cell* **178**, 1509-1525. e1519 (2019).

43   Hochgerner, H., Zeisel, A., Lönnerberg, P. & Linnarsson, S. Conserved properties of

dentate gyrus neurogenesis across postnatal development revealed by single-cell

RNA sequencing. *Nature neuroscience* **21**, 290-299 (2018).

44   Leussis, M. P., Madison, J. M. & Petryshen, T. L. Ankyrin 3: genetic association with

bipolar disorder and relevance to disease pathophysiology. *Biology of mood & anxiety*

disorders **2**, 1-13 (2012).

45   Meixner, A. *et al.* MAP1B is required for axon guidance and is involved in the

development of the central and peripheral nervous system. *The Journal of cell biology*

151, 1169-1178 (2000).

46   Douglas, R. J. The hippocampus and behavior. *Psychological bulletin* **67**, 416 (1967).

47   Sun, Y. *et al.* Promotion of astrocyte-neuron glutamate-glutamine shuttle by SCFA

contributes to the alleviation of Alzheimer's disease. *Redox Biology* **62**, 102690

(2023).

48   Weinreb, C., Rodriguez-Fraticelli, A., Camargo, F. D. & Klein, A. M. Lineage tracing on

transcriptional landscapes links state to fate during differentiation. *Science* **367**,

eaaw3381 (2020).

49   Borella, R. *et al.* Metabolic reprograming shapes neutrophil functions in severe

COVID-19. *European Journal of Immunology* **52**, 484-502 (2022).

50   Bill, M. *et al.* Mapping the CLEC 12A expression on myeloid progenitors in normal

bone marrow; implications for understanding CLEC 12A-related cancer stem cell

biology. *Journal of Cellular and Molecular Medicine* **22**, 2311-2318 (2018).

51   Rogalska, M. E. *et al.* Transcriptome-wide splicing network reveals specialized

regulatory functions of the core spliceosome. *Science* **386**, 551-560 (2024).

52   Shyu, A. B., Wilkinson, M. F. & Van Hoof, A. Messenger RNA regulation: to translate

or to degrade. *The EMBO journal* **27**, 471-481 (2008).

53   Feingold, E. *et al.* The ENCODE (ENCyclopedia of DNA elements) project. *Science*

306, 636-640 (2004).

54   Lachmann, A. *et al.* ChEA: transcription factor regulation inferred from integrating

genome-wide ChIP-X experiments. *Bioinformatics* **26**, 2438-2444 (2010).

55   Lin, Z. TIVelo: RNA velocity estimation leveraging cluster-level trajectory inference.

(2024).

56   Bastidas-Ponce, A. *et al.* Comprehensive single cell mRNA profiling reveals a detailed

roadmap for pancreatic endocrinogenesis. *Development* **146**, dev173849 (2019).