Revisiting fungal polarized growth: from spitzenkörper to crescent accumulation of secretory vesicles

Adrien Hamandjian[1,2,*], Glen Calvar[1,2], Matthieu Blandenet[1,2], Mélanie Crumière[1,2], Nathalie Poussereau[1,2], Mathias Choquer[1,2#] and Christophe Bruel[1,2#*]

[1] Université Claude Bernard Lyon 1, CNRS, INSA-Lyon, UMR5240 Microbiologie, Adaptation et Pathogénie. 10 Rue Raphaël Dubois, Villeurbanne, F-69622, France.

[2] Bayer SAS, Crop Science Division. Laboratoire Mixte, 14 Impasse Pierre Baizet, Lyon, F-69263, France.

[#] These authors contributed equally

[*] Corresponding authors : Adrien Hamandjian, Christophe Bruel

Abstract

Polarized elongation in filamentous fungi depends on secretory vesicles being supplied and fusing at the apex. In Ascomycota and Basidiomycota, these vesicles have mostly been reported to accumulate into a spheroidal structure known as the spitzenkörper. Using time-lapse microscopy, the spatial and temporal dynamics of fluorescently-labelled vesicles were investigated in 13 apices of the phytopathogenic fungus *Botrytis cinerea*. Over time, the fluorescent signal highlighted the spheroid spitzenkörper in half the sample and a crescent-shaped region of the apical dome in the other half. A linear relationship was found between the roundness of the fluorescent region and the hyphal elongation rate. Temporal Dynamics Clustering and Fourier transform showed periodic pulses of fluorescence intensity in the spitzenkörper that were absent in the crescent-shape region. These results reveal a dual mode of secretory vesicles accumulation at the apex of growing hyphae.

Highlights

- Secretory Vesicles accumulation at the apex form a spheroid spitzenkörper or a crescent

- Vesicles accumulation of at the spitzenkörper is periodic

- Roundness of SV accumulation regions correlates with elongation rates

Introduction

Sustained polar cell elongation is a shared trait in various branches of the tree of life, from filamentous bacteria such as actinomycetes[1], to tip-growing eukaryotic cells found in plants[2–4] (e.g., pollen tubes and root hairs) and algae[5] (e.g., protonemata). In eukaryotic microorganisms, it has been described in filamentous yeasts, including pathogens such as *Candida albicans*[6,7] or *Ashbya gossypii*[8] whose respective virulence to human and cotton relies on the production of filaments. More notably, polar cell elongation is a hallmark of filamentous fungi, supporting continuous hyphal extension.

Polarized cell elongation results from the spatial restriction of exocytosis within the fungal cell. By limiting vesicle fusion to a specific region, lipids are locally supplied to the plasma membrane, resulting in the unidirectional elongation of the cell[9]. The development of the mycelium, formed by extending and branching hyphae, therefore depends on the production, transport, and regulated exocytosis of secretory vesicles (SV) at the hyphal tip. SV are Golgi-derived vesicles transported to the apex by microtubules[10–12] or directly produced on-site by apical Golgi

compartments[13]. Once at the hyphal tip, also called the apex, SV are locally transported along
actin filaments to interact with the exocyst, a protein complex that facilitates vesicle docking and
fusion with the plasma membrane during exocytosis[14–16].

In addition to transporting and releasing lipids, SV also carry transmembrane proteins destined
for the plasma membrane, and soluble proteins delivered to the extracellular space, including the
cell wall. Among the transmembrane protein cargoes, chitin synthases (CHS) have received
particular attention. CHS are type 2 glycosyltransferases that synthesize chitin, a major structural
component of the fungal cell wall that is composed of N-acetylglucosamine residues linked by
beta-(1,4) linkages[17]. Because CHS are transported by SV to the apical plasma membrane,
fluorescently tagged CHS have been widely used to visualize SV trafficking[11,18–20]. The delivery by
SV of enzymes involved in cell wall synthesis and remodeling ensures that cell wall expansion
occurs in coordination with plasma membrane extension during apical elongation.

SV are the final carriers of all proteins that follow the canonical secretory pathway, i.e. proteins
translated in the endoplasmic reticulum and matured in the Golgi apparatus[21]. These proteins
include lytic enzymes involved in fungal nutrition, which play a key role in ecosystems due to their
importance in nutrient cycling[22,23]. Additionally, SV transport proteins that mediate interactions
between fungi and other organisms, notably virulence factors secreted by pathogenic species[24].
Beyond its ecological function, fungal secretion is also crucial to various industrial processes,
ranging from fermentation in the food industry to the secretion of proteins and enzymes of
economic interest[25]. Consequently, a deeper understanding of temporal and spatial dynamics of
exocytosis in fungi is needed to better understand their biology, but also to better control
pathogens or, on the contrary, maximize production yields in industrial settings.

In Ascomycota and Basidiomycota, the apex of elongating filaments is marked by a visible
accumulation of vesicles forming a spheroidal structure known as the spitzenkörper[26]. This
structure has been proposed to act as a vesicle supply center (VSC), where vesicles accumulate
before being delivered to the plasma membrane[27]. The spitzenkörper is located at the interface
between the microtubules transporting SV to the apex[28], the polarisome nucleating actin
filaments at the apex[29], and the exocyst mediating the vesicle tethering and fusion with the
plasma membrane[30]. In *Neurospora crassa*, it was shown to be stratified with SV subpopulations
known as microvesicles and macrovesicles accumulating at the core and in the outer layer of the
spitzenkörper, respectively[31].

The spitzenkörper is therefore an organized accumulation of SV at the apex which can be
described as both dynamic and stable. It is dynamic by the constant entry and exit of vesicles[26,32–
34], but stable by its importance to direct growth in actively growing filaments[35–38]. This stability is
however challenged. In the filament-producing yeast *Ashbya gossypii*, chemical labelling of lipids
showed the presence of a spitzenkörper in fast elongating hyphae, but a crescent-shaped
accumulation of vesicles in slower elongating hyphae[39]. Interestingly, both spitzenkörper and
crescent-shaped accumulation of SV have also been documented in *Aspergillus nidulans*[19,40].
While significant progress has been made in understanding the apical machinery and vesicle
trafficking during polarized cell elongation, the consistency of vesicle accumulation through time
and space across filaments remains unclear. In addition, studies on polarized cell elongation
have mainly focused on cells displaying a spitzenkörper and have been conducted in saprotrophic
model species, potentially underestimating the diversity of vesicle accumulation patterns and
behaviors both between hyphae and across fungal species.

In this study, a fluorescently-labelled class III chitin synthase was used to label SV in the phytopathogenic necrotrophic fungus *Botrytis cinerea*. Time-lapse confocal acquisitions revealed two populations of growing hyphae displaying either a spheroid spitzenkörper or a fluorescent crescent at the apex. Further temporal analysis revealed periodic variations of fluorescence intensity at the apex of hyphae displaying a spitzenkörper, that were not detected in hyphae displaying a crescent. These results reveal the existence of at least two populations of hyphae that display different spatial and temporal dynamics of SV distribution in the apical dome.

Results

1. The fluorescently-labelled class III chitin synthase BcCHSIIIa localizes at the tip of growing hyphae in *B. cinerea*

To visualize SV at the apex of *B. cinerea* hyphae, a reporter system was required. Based on the successful labelling of SV through the fluorescent tagging of class III chitin synthases in different ascomycetes[11,18–20,40], the *B. cinerea* chitin synthase IIIa (BcCHSIIIa - Bcin04g03120) was chosen[41]. The gene encoding an optimized version of the fluorescent protein eGFP[42] was fused at the 3' end of the BcCHSIIIa gene by homologous recombination to produce the chimeric BcCHSIIIa::eGFP protein (Fig.S1). In parallel, a control strain was constructed that produces that eGFP alone under the control of the constitutive pOliC promoter (eGFP[Cyt] strain).

Following growth on a glass surface, confocal microscopy was performed on both strains. As expected, the eGFP[Cyt] control strain displayed a diffuse cytosolic fluorescent signal (Fig.1A). In contrast, mobile fluorescent speckles and an apical accumulation of fluorescence were observed in the BcCHSIIIa::eGFP strain (Fig.1B) that were consistent with previously reported localizations of SV transporting chitin synthases[18–20,40]. These observations therefore validated the creation of the desired reporter strain.

2. Patterns of apical BcCHSIIIa::eGFP accumulation are linked to hyphal elongation rate

Noticeably, in different hyphae of the BcCHSIIIa::eGFP strain, the fluorescence signal at the apex appeared either as a spheroid spot or was more diffused (Fig.1B, right panel top and bottom). To exclude the possibility of an artifact, 10 min time-lapse acquisitions (1 frame every 4.455 sec) were collected on 13 BcCHSIIIa::eGFP hyphae (Fig.1C). For each acquisition, the apex was tracked and cropped to accommodate for hyphal elongation. The overall distribution of fluorescence over the course of the acquisition was visualized by performing time-projections (Fig. 1C, top). The region of maximal fluorescence accumulation (see *Material and methods*) was then extracted for each hypha using a threshold-based approach (Fig.1C middle). In some hyphae, this region showed as a spheroid spot, reflecting a spatially confined accumulation of the fluorescent SV over time. In other hyphae it showed as a crescent. This crescent was not necessarily occupied at once, but resulted from changes over time in the position of the fluorescent signal that shaped a crescent when cumulating all the frames in a projection. The area of both the spheroid and crescent-shaped regions of fluorescence accumulation were measured and followed a normal distribution (Shapiro-Wilk test, *p*-value 0.297, mean area of $0.428 \pm 0.019 \, \mu m^2$). The roundness of these regions were also measured and were normally distributed, ranging between 0.290 and 0.818 (Shapiro-Wilk test, *p*-value 0.446, mean roundness of 0.529 ± 0.154). This last observations indicate that the apical region of fluorescence accumulation spans a continuum of shapes, ranging from crescent-like to spherical.

For each acquisition, kymographs were then produced to estimate the growth rate of each hypha during acquisition (Fig.S2). This rate varied between 104 and 753 nm.min^{-1} (mean value 301 ± 173 nm.min^{-1}) and no statistical difference was observed between the control (eGFP[Cyt]) and

the BcCHIIIa::eGFP strain (Student t-test, *p*-value 0.190, mean growth rates of 450 ± 247 nm.min⁻
^{1} and 301 ± 173 nm.min^{-1} with n = 5 and n = 13, respectively). In contrast, a positive linear
correlation was observed between the roundness of the region of fluorescence accumulation and
the hyphal elongation rate in acquisitions of the BcCHSIIIa::eGFP strain (Fig.1F, Pearson
correlation 0.675, *p*-value 0.011, analysis of residues in Figure S3).

3. Temporal Dynamics Clustering reveals two clusters based on fluorescence variability over
time

In addition to analyzing the spatial organization of SV at the apex, we examined the temporal
dynamics of their accumulation. For each acquisition and at every frame, the mean intracellular
fluorescence intensity was measured in the apical region (4 µm long from the tip, see *Materials*
and Methods) of hyphae from the control eGFP^{Cyt} strain expressing cytoplasmic eGFP and from
the BcCHSIIIa::eGFP strain. This yielded, for each acquisition, a time series representing the
mean fluorescence intensity at the apex over time. In all time series this intensity fluctuated over
time, independently of movements in the z dimension (Fig.S4). Since such fluctuations can result
from intrinsic properties of eGFP (folding, maturation or stability), we compared the dynamics
observed in the BcCHSIIIa::eGFP strain to those of the control strain expressing cytosolic eGFP.

To characterize the temporal dynamics of the fluorescence signals, we applied Temporal
Dynamics Clustering (TDC)[43], a features-based clustering approach that groups time series based
on fluorescence variability over time. As drawn in figure 2A, the signals were detrended and the
first-order differences were computed for each time series and expressed as a cumulative first-
order difference (Fig.2B). Next, TDC parameters *u* and *lq* were calculated to partition each
cumulative distribution into two sections: the *main* section, representing common variations, and
the *tail* section, capturing extreme fluctuations (Fig. 2B). For each distribution, the mean value of
elements in the *main* section (μ_{main}), the associated standard deviation (σ_{main}), and the mean value
of elements from the *tail* section (μ_{tail}) were calculated. These three metrics were then used as
features for *k*-means clustering, which identified two distinct clusters, termed Cluster 1 and
Cluster 2 (Fig.2C).

The clustering quality was supported by a silhouette score of 0.761, indicating strong and well-
defined separation between clusters. Moreover, statistical analysis revealed significant
differences between clusters for all 3 TDC metrics (Fig.2D, Student's t-test, *p*-values of 6.7e-9 for
μ_{main}, 2.7e-9 for σ_{main}, and 7.7e-9 for μ_{tail}). These differences in the 3 metrics suggest that
fluorescence variability between clusters differs not only in extreme events (μ_{tail}) but also in their
overall temporal behavior (μ_{main}). In contrast, the median fluorescence intensity before detrending
was not significantly different between clusters (Mann-Whitney U test, *p*-value 0.073), suggesting
that the distinction arose from differences in temporal variability rather than absolute intensity.
Together, these results demonstrate the presence of two statistically distinct patterns of apical
fluorescence dynamics in our sample of hyphae.

4. Distinct fluorescence dynamics are linked to the roundness of apical vesicle
accumulation

Cluster 1 included all the acquisitions recorded on the 5 hyphae from the control eGFP^{Cyt} strain.
Since clustering was based on fluorescence variability over time, Cluster 1 therefore likely reflects
the intrinsic fluorescence fluctuations of eGFP itself. Interestingly, the 13 acquisitions recorded
on the 13 hyphae from the BcCHSIIIa::eGFP strain distributed across both clusters (6 in Cluster 1
and 7 in Cluster 2). This indicated the presence of two populations of growing hyphae in the
BcCHSIIIa::eGFP strain, differing in apical fluorescence variability though time. In all the hyphae

grouped in Cluster 2, the temporal fluorescence behavior at the apex was distinct from the one
observed in hyphae of the control strain, suggesting a specific dynamic of SV in these hyphae.

Interestingly, a link between clustering and roundness of the apical fluorescent signal was noticed
in hyphae in the BcCHSIIIa::eGFP strain. Cluster 1 grouped the hyphae with roundness values
below 0.5 while Cluster 2 grouped the hyphae with roundness values above 0.5 (Fig.3B, Mann–
Whitney U test, *p*-value 0.001). In comparison, neither hyphal width (Fig.3C) nor apex curvature
(Fig.3D) significantly influenced the clustering outcome (Mann–Whitney U test, *p*-value 0.478,
Student's t-test, *p*-value 0.219, respectively). This indicates that the temporal dynamics which
separate the acquisitions of Cluster 1 from those of Cluster 2 are associated with differences in
the spatial organization of the SV-related apical fluorescence. On one hand, the hyphae belonging
to Cluster 1 display a crescent accumulation of fluorescence at the apex (roundness < 0.5) whose
variations over time could not be distinguished from the intrinsic fluorescence variations
displayed by cytosolic eGFP. On the other hand, the hyphae belonging to Cluster 2 display a
spherical accumulation of fluorescence at the apex (roundness > 0.5) whose variations over time
statistically differed from the intrinsic dynamic of eGFP. This suggests that the measured
variations of fluorescence in these hyphae originates from significant fluctuations in vesicle
accumulation at the apex over time. The spherical accumulation of fluorescence at the apex being
consistent with the presence of a spitzenkörper[18–20,40], we considered that the Cluster 2 hyphae
displayed a spitzenkörper.

5.   Fluorescence variations are periodic in hyphae exhibiting a spitzenkörper

The production of kymographs from the time-lapse acquisitions of the BcCHSIIIa::eGFP strain
revealed occurrences of fluorescence hotspots (Fig.S2), as exemplified by one acquisition (Hypha
12) shown in Figure 4A. These spots reflected spatially and temporally restricted events of
fluorescence accumulation which appear as oscillations when the mean fluorescence intensity
at the apex was plotted against time (Fig.4B). Noticeably, these events appeared regularly spaced,
suggesting the presence of an underlying periodic behavior.

To evaluate the presence of a putative period governing the detected oscillations, the Fast Fourier
Transform (FFT) was used. Pre-processed time series were analyzed (see *Material and methods*)
and the outputs were expressed as amplitudes over periods, producing amplitude spectra
(Fig.4C, Fig.5 left column). In the example shown in Figure 4C, the amplitude spectrum revealed
a prominent oscillatory component with a period of 39.75 seconds. When this period was used to
generate a sinusoidal model (Fig.4D, green line), this model closely aligned with the experimental
data (Fig. 4D, grey line). Comparable results were obtained across all acquisitions grouped in
Cluster 2 (Fig.5), each displaying a dominant oscillator with periods ranging from 31.39 to
64.93 seconds (mean period of 45.55 ± 10.09 seconds). All these acquisitions therefore shared a
temporal dynamic marked by periodic events of fluorescence increases. In contrast, no such
periodic behavior could be detected in the acquisitions grouped in Cluster 1. The existence of a
linear correlation between the oscillation period and the hyphal elongation rate was explored but
not found (Fig.S5, Pearson correlation, *p*-value 0.944). More acquisitions would however be
needed to draw definitive conclusions regarding the existence of such a relationship.

Discussion

Polar cell elongation is a hallmark of vegetative growth in filamentous fungi. While the apical
accumulation of Secretory Vesicles (SV) in a spitzenkörper has been broadly documented in
Ascomycota and Basidiomycota, scarce observations of hyphae showing vesicle accumulation
in a crescent[19,39,40] suggested a diversity of SV accumulation patterns in growing hyphae. Here we

used the fusion of a class III chitin synthase BcCHSIIIa (common SV marker) with the fluorescent
protein eGFP to study the apical accumulation of SV in growing hyphae of the plant
phytopathogen *B. cinerea*. As expected, we observed a concentrated accumulation of
fluorescence at the apex of growing hyphae, but the shape of the region of fluorescence
accumulation varied between hyphae, from spheroid to crescent-like. In line with the observation
of a spitzenkörper in *B. cinerea* hyphae by electron microscopy[44], the spheroid accumulation of
fluorescence in some growing hyphae is consistent with the existence of a spitzenkörper in these
hyphae, reflecting a spatially restricted accumulation of SV at their apex.

The crescent-shaped region of fluorescence accumulation reported here corresponds to an area
of the apex within which the position of the recorded maximum fluorescence moves over time.
Although less documented than accumulation at the spitzenkörper, a crescent organization was
previously reported in *Aspergillus nidulans* expressing a fluorescently labeled class III chitin
synthase[19]. It was also reported in the filamentous yeast *Ashbya gossypii* where SV and
components of the exocyst and polarisome were labeled[39]. Altogether these observations
suggest that a dual mode of apical SV accumulation can exist in filamentous fungi and may be
shared among species. Our results document the existence of these two modes of SV
distributions in growing hyphae of *B. cinerea*, but they also suggest that these two modes could
be endpoints of a continuum rather than discrete categories, with intermediate forms distributed
normally.

We observed a positive correlation between hyphal growth rate and the spatial organization of the
fluorescence accumulating at the apex. Hyphae with fast elongation rates (>350 nm.min^{-1})
consistently exhibited a spitzenkörper. This relationship is reminiscent of findings in *A. gossypii*,
where the organization of the polarisome and exocyst, two complexes known to interact with the
spitzenkörper, were shown to depend on growth rate, with faster-growing hyphae displaying a
spherical organization and slower-growing ones showing a crescent-like arrangement[39]. Similarly,
in *N. crassa*, immature hyphae lacking a spitzenkörper exhibit lower elongation rates than mature
hyphae in which a spitzenkörper is present[33,45]. Altogether, these findings suggest a
interdependence between spitzenkörper formation and hyphal elongation. However, whether
vesicle accumulation facilitates faster growth or, conversely, rapid growth promotes
spitzenkörper assembly remains to be elucidated. In addition, we also observed hyphae growing
at similar rates (233 and 221 nm.min^{-1} for example) but exhibiting dissimilar apical SV
organizations (crescent and spitzenkörper with roundness of 0.290 and 0.597, respectively in this
example). Intermediate elongation rates may therefore be permissive to either the presence or
absence of the spitzenkörper. Such a buffer zone was observed in *A. gossypii*, where some
elongation rates were observed in hyphae with crescent or spitzenkörper-like organization of the
exocyst or of the polarisome[39]. Altogether this suggests a not-strictly linear relationship between
elongation rate and SV organization and these observations across species could support a
model in which SV apical accumulation undergoes growth rate-dependent reorganization,
transitioning from a crescent to a spitzenkörper as the elongation rate increases. With none of our
recordings capturing hyphae undergoing substantial stable change in elongation rate, such a
transition could not be observed. Whether such a reorganization occurs progressively within
individual hyphae or whether crescent and spitzenkörper represent distinct and stable states
associated with different hyphal populations therefore remains an open question.

We next investigated the temporal dynamics of apical fluorescence accumulation. Time-lapse
confocal imaging revealed that fluorescence intensity fluctuated over time. To characterize these
temporal patterns, we applied Temporal Dynamics Clustering[43] (TDC) and this identified two

distinct clusters: Cluster 1 grouping time series whose fluorescence profiles were
indistinguishable from those of the control strain expressing cytosolic eGFP, and Cluster 2
grouping time series with marked fluctuations. Remarkably, all hyphae showing fluorescence
accumulation in a crescent-shaped apical region grouped in Cluster 1, whereas hyphae with a
spitzenkörper grouped in Cluster 2. Differences in the spatial organization of SV accumulation are
therefore also associated with differences in temporal behavior (the fluorescence in hyphae
displaying a spitzenkörper exhibit marked variations in intensity over time).

We further characterized the temporal dynamics present in acquisitions of hyphae displaying a
spitzenkörper (Cluster 2) using the Fourier transform analysis. In all these hyphae, a dominant
oscillatory component was found, indicating a repeated succession of high and low fluorescence
intensity events regularly spaced over time. These variations are consistent with a periodic
succession of SV accumulation and dispersion events at the apex and such periodic apical
dynamics were previously described in *N. crassa* and *A. nidulans*[46] as well as in
Arthrobotrys flagrans[18]. Interestingly, the study by Takeshita *et al.*[46] led to the proposition of an
inverse relationship between the hyphal elongation rate and the oscillation period across species.
In *N. crassa,* elongation is rapid (12 µm.min^{-1}) and the period governing the apical fluorescent
oscillation is short (17 ± 6 seconds)[46]. In *A. nidulans,* elongation is slower (1.8 µm.min^{-1}) and the
period is longer (30 ± 7 seconds)[46]. Our results in *B. cinerea* would support this trend with an even
slower hyphal mean elongation rate (0.34 ± 0.21 µm.min^{-1}) and a longer period governing the
apical fluorescent oscillation (45 ± 10 seconds). While the methods used to identify this
periodicity differ (measurement of the temporal distance between fluorescence peaks in the
study of Takeshita *et al.*[46] and Fourier transform analysis in our study), the observed pattern
suggests that faster elongation rates may be associated with shorter oscillation periods at the
species level. Such a relationship could be explained by the demand for lipids at the membrane
level. Assuming stable size of SV, an increase in elongation rate would require an increase in lipid
supply, made possible by a reduction in the time between exocytosis events. However, no such
inverse relationship was observed when we compared periods and elongation rates between the
7 Cluster 2 hyphae of our study in *B. cinerea*. This absence may stem from insufficient sampling
to detect subtler trends. A similar situation has been reported in pollen tubes, a tip-growing plant
cell type that also exhibits periodic elongation, and in which a correlation between period and
growth rate became apparent only with large sample sizes, due to high variability around the
trend[47].

Altogether, our study reveals the presence of two subpopulations in growing hyphae of *B. cinerea*
that differ in the temporal dynamics of SV accumulation at the apex. One subpopulation shows
relatively constant SV accumulation, while the other displays periodic fluctuations reflecting
cycles of SV accumulation and release. This dichotomy was also observed in space, as the first
subpopulation exhibited a crescent-shaped fluorescence signal, while the second showed a
spherical accumulation reminiscent of the spitzenkörper. However, these two types of
accumulation were not discrete classes but rather both ends of a continuum, revealing a plastic
dimension to SV organization and accumulation in filamentous fungi. Recognizing this plasticity
may reshape current models of fungal growth regulation, with broader implications for fungal
development, adaptability, and pathogenicity. In light of these findings, it becomes interesting to
investigate hyphal elongation in filamentous fungi with greater attention to the diversity of apical
dynamics. Studies focusing solely on hyphae with a spitzenkörper may capture only a subset of
fungal behavior and risk overlooking alternative mechanisms underlying polarized growth.

Limitations of the study

While class III chitin synthases (CHS) are commonly used to visualize secretory vesicles via fusion
with a fluorescent protein, it is important to note that these enzymes may be transported by only
a subpopulation of vesicles. In *N. crassa*, CHS are specifically transported by microvesicles while
the spitzenkörper also contains macrovesicles[31]. In *B. cinerea*, transmission electron microscopy
has revealed the presence of both micro- and macrovesicles, but the distribution of BcCHSIIIa
between these subpopulations remains unknown.

While the TDC approach did not identify fluorescence variations at the apex of the Cluster 1
hyphae (displaying a crescent-shaped region of fluorescence accumulation) when compared to
the control strain hyphae, the existence of subtle variations cannot be ruled out. Acquisitions
using more spatially and temporally resolutive microscopy are needed to further explore these
dynamics.

The experimental design and confocal microscope limited the number of acquisitions. Additional
data would help strengthen the analysis of the linear correlation between roundness of the region
of accumulation and elongation rate. Nonetheless, statistical metrics and residual analysis
provide a strong support for this relationship, even with the current sample size. Additional
acquisitions of hyphae displaying a spitzenkörper will also help to uncover the relationship that
connects period and elongation rate.

The signal pre-processing used in this study was intentionally stringent to minimize false
positives, but it may have filtered out biologically informative signals, notably low-frequency
patterns. Based on this work, we encourage future studies to further explore temporal dynamics
that may have gone undetected in our study.

Author contribution

AH conceived the study, performed the experiments, analyzed the data, developed the software
tools, and prepared the figures. GC, MB, and M. Crumière produced the fungal strains used in this
study. NP, M. Choquer, and CB secured funding and supervised the project. AH wrote the initial
draft of the manuscript, which was reviewed and edited by AH, M. Choquer and CB, with input
from all authors.

Material and methods

1. Fungal strains

The strains BcCHSIIIa::eGFP and eGFP[Cyt] were obtained by *Agrobacterium tumefasciens*-
mediated transformation of the *B. cinerea* strain B05.10[48]. The BcCHSIIIa::eGFP strain was
generated by the insertion, through homologous recombination, of an eGFP-tNiaD-NatR DNA
fragment in frame with the BcCHSIIIa (Bcin04g03120) coding sequence (Fig.S1A). This DNA
fragment, containing eGFP optimized for *B. cinerea*[42], the terminator of the nitrate reductase from
B. cinerea (tNiaD) and a nourseothricin resistance gene (NatR)[49], was amplified from a donor
plasmid. A 1031 bp fragment corresponding to the end of the BcCHSIIIa coding sequence, minus
the STOP codon, was then amplified from B05.10 genomic DNA. A 924 bp fragment was amplified,
corresponding to the 3' UTR and terminator region of the BcCHSIIIa gene. The eGFP-tNiaD-NatR
fragment was then assembled between the BcCHSIIIa coding sequence and the BcCHSIIIa
terminator fragment using double-joint PCR[50]. This DNA cassette was inserted by *In Vivo*
Assembly (IVA) cloning[51] into a *A. tumefasciens* transfer-DNA carried by the p7 vector. Following
verification by restriction enzyme digestion and sequencing, the vector was introduced in
A. tumefasciens (strain LDA 1126) and the transformed bacteria was used to mutagenize
B. cinerea[48].

For the eGFP^{Cyt} strain, eGFP was placed under the control of the constitutive OliC promoter
(pOliC) from *A. nidulans*[52] and the tNiaD terminator. The pOliC-eGFP-tNiaD cassette was then
inserted in the *A. tumefasciens* tDNA carried out by a pBHT2 plasmid containing a hygromycin
resistance cassette[53]. Transformation led to the obtention of *A. tumefasciens* transformants that
were later used to transform conidia of the parental B05.10 strain. *B. cinerea* transformants
displaying hygromycin and nourseothricin resistance where then screened using epifluorescence
microscopy. One transformant with strong signal was then selected for use in this study.

For all strains, conidia were harvested after 10 days of culture at 21°C with near-UV light on a
medium containing 0.5% of glucose, 0.1% of malt extract, 0.1% of tryptone, 0.1% of casamino
acids, 0.1% of yeast extract, 0.02% of ribonucleic acids and 1.6% of agar[54]. Conidia were then
stored at -80°C and defrosted on ice before use in experiments.

2. Image acquisition

Before acquisition, conidia were diluted to 1.10^3 conidia.ml^{-1} in MMII medium (NaNO$_3$ 24 mM,
Glucose 111 mM, KH$_2$PO$_4$ 0.147 mM, MgSO$_4$ 0.039 mM, KCl 0.134 mM, FeSO$_4$/7H$_2$O 0.7 μM) to
inoculate wells of a 96 wells SensoPlate microplate (GREINER) with 200μl per well. Incubation
was performed at 21°C, in the dark. Time-lapse acquisitions (10 min) were conducted using a
laser scanning confocal microscope (ZEISS Observer 7 equipped with an LSM 800) and a Plan-
Apochromat 63x Oil DIC objective (ZEISS). Excitation was achieved using a 493 nm laser, and
fluorescence emissions were detected within the range of 491 to 573 nm. The laser scanning was
performed with an averaging of 16 on a single z-position. To ensure a uniform acquisition speed,
the size of the acquired region was standardized (295x224 pixels) leading to an acquisition rate of
4.456 seconds per frame.

3. Tracking of apexes

To compensate for the elongation of filaments occurring during the acquisition, a custom FIJI[55]
macro was designed to move a region of interest (ROI) of 50x50 pixels (6.78x6.78 μm) through
time. Each frame was then cropped to obtain a 50x50 time-lapse acquisition with the apex (4 first
micrometers of the hyphae) centered at each frame.

4. Morphological measures

The curvature of the apex was measured on non-cropped acquisitions using the Kappa plugin[56] in
FIJI. A 10 μm curved was placed to follow the apical dome, the curvature was then estimated as
the curvature of the most apical point on the first frame of each acquisition. The hyphal width was
calculated as the average of three measures performed in the distal part of the apical region
(3.3 ± 1.2 μm) on the first frame of each non-cropped acquisition.

To measure the geometry and area of maximal fluorescence intensity at the apex, a time-
projection was first performed using the average intensity function in FIJI. Then, images were
thresholded to only keep pixels in the 1% of highest intensity. The shape and area of the obtained
structure was measured directly in FIJI.

Kymographs were produced using the KymographBuilder[57] plugin (v.3.0.0) in FIJI. For growth
speed estimation, kymographs were produced using all time points of the acquisition (135 total
points with a time between frames of 4.456 seconds).

5. From acquisition to time series

Before measuring the intensity of fluorescence, two approaches were used to suppress
background pixels that would bias the measure of fluorescence per frame: one to suppress
extracellular pixels and one to suppress cytosolic background signals.

To limit the impact from extracellular pixels, a 50x50 pixels ROI was placed in the extracellular
region uncropped acquisitions from both the BcCHSIIIa::eGFP and the eGFP^{Cyt} strains. The mean
fluorescence intensity in this ROI was measured for all acquisition times and added to twice the
associated standard deviation such as 95% of extracellular pixels have values bellow the set
threshold[58]. All pixels with intensity values below this threshold were replaced by NaN values. The
threshold was computed for each acquisition independently. Thresholded acquisitions were then
cropped (see *Tracking of apexes*). Once cropped, the mean fluorescence intensity was measured
at all time points of an acquisition to produce a time series (*i.e.* the mean fluorescence intensity
over time). This approach restricts the measure of fluorescence intensity to the inside of the
hyphae (termed *threshold technic 1* in this section).

To limit the impact from cytosolic background fluorescence present in the acquisitions of the
BcCHSIIIa::eGFP strain, and thus limit the measure of pixel intensity to the regions of
fluorescence accumulation, a 16x16 pixels ROI was placed in the sub-apical region (5 ± 2 μm
bellow the hyphal tip) post-acquisition. The mean fluorescence intensity in this ROI was
measured for the first 20 frames and added to twice the associated standard deviation such as
95% of background signals are smaller than the threshold value[58]. This threshold was used as an
estimated of the intensity of the cytosolic background signal. All fluorescence values below this
threshold were replaced by NaN values. Thresholded acquisitions were then cropped (see
Tracking of apexes). Once cropped, the mean fluorescence intensity was measured at all time
points of an acquisition to produce a time series. This approach restricts the measure of
fluorescence intensity to regions of strong fluorescence (termed *threshold technic 2* in this
section).

Following both approaches, the mean pixel intensity was measured at each frame producing one
time series per acquisition corresponding to the mean fluorescence intensity over time. For each
signal (*i.e.* time series) a trend was identified using a Savitzky-Golay filter of window length 10
(polynomial degree of 2). Each signal was then subtracted by its own trend to obtain detrended
signals. No normalization was applied to preserve the relative amplitude of short-term
fluctuations.

6. Temporal Dynamics Clustering

Temporal Dynamics Clustering (TDC) is a featured-based clustering approach for time series[43].
This approach was performed on detrended signals obtained after the threshold technic 1 (see
From acquisition to time series). Following the approach detailed by Shuyang Li *et al.*[43], the first
order difference was computed independently for all signals such as:

$$\Delta y_t = |y_t - y_{t-1}|$$

In our context, Δy_t therefore corresponds to the variation of fluorescence intensity between two
successive frames. The second step of the TDC is the computation of u and of the subsequent I_q,
with:

$$u = \frac{(T-1)^{2/3}}{log(log(T-1))}$$

and :

439
$$I_q = 1 - \frac{u}{T - 1}$$

With T the length of the original time series. In our study, the number of time points is content
between acquisitions leading to a single u and I_q values for all time series. I_q is a quantile index
allowing the spreading of a time series into a *main* section containing all y_t values below the and
at the quantile I_q and a *tail* section containing all y_t values above I_q. Following this partition of first-
order differences, the *main* section contains moderate variations of fluorescence intensities
while the *tail* section contains extreme events of fluorescence variation. These *main* and *tail*
sections allow the production of metrics μ_{main} and μ_{tail} corresponding to the mean fluorescence
intensities of the *main* and *tail* section respectively, and of σ_{main} that is the standard deviation of
the *main* section. The values of μ_{main}, μ_{tail} and σ_{main} are features used for the clustering using k-
means clustering[59]. For a number of clusters from 2 to T-1, silhouette scores were computed to
measure cluster quality and determine the optimal number of clusters, defined as the highest
silhouette score with the minimal number of clusters.

7.   Fast Fourier Transform

To study temporal variations in fluorescence intensity in regions of fluorescence accumulation,
we further analyzed the detrended time series obtained after the threshold technic 2 (see section
From acquisition to time series) using Fast Fourier Transform (FFT). Time series were first
smoothed using a Savitzky-Golay filter (window size of 10, polynomial degree of 2) in combination
with a moving average filter (window size of 3). To improve the frequency resolution, all signals
were zero-padded to reach a total of 16 384 time points.

For each signal S, the FFT of S (FFT(S)) was computed for all discrete frequency bins f using the
NumPy package[60]. The amplitude $A(f)$ associated with each frequency f was computed as :

461
$$A(f) = 2\frac{\sqrt{Re[FFT(S)(f)]^2 + Im[FFT(S)(f)]^2}}{N}$$

Where Re and Im denote the real and imaginary part of the FFT output, respectively, and N the
number of points in the signal before zero padding. The factor 2 is used to correct the symmetrical
distribution of the FFT output over positive and negative frequencies. For easier biological
interpretation, amplitudes were plotted over periods (in seconds) instead of frequencies (in Hz).
To satisfy the Nyquist sampling theorem, periods below 9 seconds were not considered.
Additionally, only periods that could be sampled in at least four complete cycles during the 10-
minute time-lapse acquisitions were considered, excluding periods longer than 150 seconds.

8.   Sinusoidal model

For a period p and its associated amplitude A_p, we created a sinusoidal model using the equation

471
$$M(t) = A_p \cdot sin\left(\frac{2\pi}{p}(t - \phi)\right)$$

472   with ϕ the phase shift defined as

473
$$\phi = \theta\frac{p}{360}$$

The value of ϕ was estimated as the value minimizing the Mean Squared Error between the model
and the experimental data for values of θ between 0 and 360°.

9. Linear modeling and residual analysis

To produce linear models, the normality of the datasets was first evaluated using a Shapiro-Wilk
test with a threshold p-value of 0.05. Peason and Spearman correlations were performed. The
linear regression model was obtained by fitting an ordinal least squares regression using the
statsmodels package[61]. A constant was added to the model to allow for a regression line with a
non-zero intercept. Once the model was obtained, the 95% confidence interval was extracted
using the statsmodels package and the R^2 computed. The influence of individual observations on
the linear model was then assessed using residuals, standardized residuals and Cook's distance.
Observations with a Cook's distance exceeding the 4/N threshold were considered influential
outliers[62].

10. Control for z-axis movement confirms genuine fluorescence dynamics

To rule out the possibility that the observed fluorescence variations resulted from z-axis
movements during the recorded single-plane time-lapse acquisitions, we performed a time-lapse
z-stack imaging on an hypha displaying a spitzenkörper (Fig.S4). The stack consisted of four focal
planes (Z1 to Z4), with Z1 capturing the extracellular space and Z2–Z4 intersecting the fungal cell
(Fig.S4A). First, time projection following apex tracking revealed a spheroid fluorescence
accumulation in the 3 planes intersecting the fungal cell, with roundness values of 0.510, 0.832,
and 0.575, respectively. Then, for each plane and frame, the mean intracellular fluorescence
signals were extracted over time to produce time series (Fig.S4B). These signals were detrended
to center them around zero (Fig.S4D) and finally compared through cross-correlation at lag 0. As
shown in Figure S6E, the extracellular plane (Z1) and the intracellular planes (Z2-Z4) showed low
similarity (correlation ≤ 0.3), whereas intracellular planes exhibited strong to very strong
correlations (correlation = 0.6-0.9) between them. This indicates that the signals recorded from
the 3 intracellular planes correspond to a single, consistent, intracellular fluorescence source.
Artifacts caused by z-axis movements are therefore unlikely to cause the observed periodic
fluorescence variations in the Cluster 2 hyphae.

11. Statistics

Throughout the article, measures are presented as mean ± standard deviation. When statistical
tests were used, the name of the statistical test is indicated in parenthesis along with its
associated p value. Statistic tests were performed in python using the scipy package[63], with a
significance threshold set at 0.05. In plots, '*ns*' indicates 'non-significant' tests ($p > 0.05$). A single
asterisk (*) indicates $p < 0.05$, double asterisks (**) indicate $p < 0.01$ and triple asterisks (***)
indicate $p < 0.001$.

Figure legends

1. Figure 1. Distinct accumulation patterns of BcCHSIIIa::eGFP correlate with hyphal growth
rates

(A–B) Fluorescence intensity measured along the longitudinal axis of growing hyphae from the
eGFP^{Cyt} strain (A) and the BcCHSIIIa::eGFP strain (B). Solid lines represent mean fluorescence
intensity, shaded areas indicate standard deviation (n = 5 for A, n = 12 for B). Confocal images
show representative hyphae (scale bar = 3 µm). (C, top) Time projections after apex tracking,
showing the overall fluorescence distribution over the acquisition time. (C, middle) Region of
fluorescence accumulation, defined by the pixels with the top 1% intensity in (C, top). Dotted
white lines outline hyphae. Regions of interest displayed in (C, top) and (C, middle) are
6.78x6.78 µm. (C, bottom) Scatter plot of roundness values measured from regions of
accumulation in (C, middle). Colors indicate roundness values. (D) The distribution of roundness

measures follows a normal distribution (Shapiro-Wilk test, *p*-value 0.446). (E) The distribution of
growth rate measures also follows a normal distribution (Shapiro-Wilk test, *p*-value 0.063). (F)
Scatter plot of roundness versus growth rate for each acquisition. The solid grey line shows the
linear regression, the grey-shaded area represents the 95% prediction interval for individual
observations. Data points are color-coded according to roundness values as in (C). Numbers in
(F) correspond to Hypha 1 to 13 as defined in (C).

2.   Figure 2. Temporal Dynamics Clustering reveals two distinct clusters differing in their
528   variability over time

(A) Schematic overview of the Temporal Dynamics Clustering (TDC) approach. Original signals
were detrended (first line), and first-order differences were computed and sorted (second line).
The sorted values were partitioned into *main* and *tail* sections using the TDC-defined parameters
u and *lq* (third line). From each acquisition, three parameters were extracted, the mean of
elements in the *main* section (μ_{main}), the associated standard deviation (σ_{main}) and the mean of
elements in the *tail* section (μ_{tail}) (fourth line). These features were used for *k*-means clustering to
group signals based on fluorescence variability (fifth line). (B) Sorted first-order differences for
acquisitions of the control eGFP^{Cyt} strain (left, n = 5), BcCHSIIIa::eGFP strain (middle, n = 13), or
combined data (right). Dashed horizontal lines indicate the threshold *lq* used to separate the *main*
and *tail* sections. (C) Scatter plot of μ_{tail} over μ_{main} for all acquisitions of the eGFP^{Cyt} strain
(diamond) and the BcCHSIIIa::eGFP strain (circles). Dashed circle outlines are used to visualize
Cluster 1 and Cluster 2 identified by *k*-means. BcCHSIIIa::eGFP data points are color-coded
according to the roundness of the fluorescence accumulation region as in Figure 1, and numbered
to match the corresponding hyphae shown in Figure 1C. (D) Comparison of μ_{main}, μ_{tail} and σ_{main}
between Cluster 1 and Cluster 2. Student's t-test *p*-values are 6.7e-9 for μ_{main}, 7.7e-9 for μ_{tail} and
2.7e-9 for σ_{main}.

3.   Figure 3. The roundness of the region of fluorescence accumulation differs between
546   acquisitions belonging to the TDC Cluster 1 or Cluster 2

(A) Schematic representation of a hypha showing hyphal width, the point of maximal curvature
and the region of fluorescence accumulation whose roundness was measured. (B) Roundness of
the region of fluorescence accumulation measured on time projection of BcCHSIIIa::eGFP
acquisitions, grouped by TDC clusters (Mann–Whitney U test, *p*-value 0.001). (C) Hyphal width
measured for acquisitions of both control eGFP^{Cyt} and BcCHSIIIa::eGFP strains, grouped by
cluster (Mann–Whitney U test *p*-value 0.479). (D) Maximum curvature of the apex grouped by
cluster (Student's t-test *p*-value 0.219). (B–D) Data points are color-coded according to the
roundness measure, consistent with Figures 1 and 2. Data points of the eGFP^{Cyt} strain acquisitions
are shown in gray.

4.   Figure 4. The spherical accumulation of fluorescence is marked by periodic variations of
557   fluorescence

(A, left) Frame at t_0 after tracking and cropping of the apex in the acquisition used as an example
in this figure (Hypha 12). The yellow line indicates the position used to generate the kymograph.
Scale bar = 1 µm. (A, right) Kymograph showing fluorescence intensity along the 3.8 µm line
(yellow line in the left panel) over 135 frames (10 minutes total, 4.455 s interval). The absence of
slope results from the dynamic tracking of the apex. White asterisks mark approximate positions
of fluorescence hotspots. (B) Mean fluorescence intensity measured per frame over time. The
blue square highlights a timepoint of high fluorescence and its corresponding frame (top right),
the red square marks a low-intensity timepoint and its frame (bottom right). (C) Amplitude

spectrum from Fast Fourier Transform analysis of the time series in (B), cropped to periods shorter than 150 seconds. The green marker indicates a peak corresponding to a period of 39.75 s.(D, top) Kymograph, (middle) the corresponding time series, (bottom) a sinusoidal model with a 39.75 s period. Grey zones represent predicted fluorescence peaks from the model and are overlaid on both the time series and the kymograph.

5. Figure 5. All hyphae assigned to Cluster 2 display a dominant oscillatory component

(Left column) Amplitude spectra obtained for acquisitions belonging to the TDC Cluster 2. Colored markers indicate the peak of maximal amplitude in each spectrum. For each acquisition, the peak amplitude and corresponding period were used to generate a sinusoidal model (colored bold line), overlaid on the experimental signal (grey line). Acquisitions are sorted in ascending order based on the roundness of the fluorescence accumulation region, such that the topmost acquisition has the lowest roundness value, and the bottommost has the highest. Models are color-coded according to the roundness measure, consistent with previous figures.

6. Figure S1. Construction of the BcCHSIIIa::eGFP strain

(A) Schematic representation of the insertion of the *BcchsIIIa::eGFP-nat^R-3'UTR* cassette by homologous recombination at the *BcchsIIIa* locus. A 1 031 bp fragment corresponding to the end of the *BcchsIIIa* coding sequence was fused with the coding sequence of the fluorescent protein eGFP, the *nat^R* resistance cassette (conferring resistance to nourseothricin), and the 3'UTR of *BcchsIIIa*. (B) PCR amplification using primers AtMT_S-IIIa-GFP-RFP-F and AtMT_S_GFP-R confirms the presence of the genetic construct in the BcCHSIIIa::eGFP strain, and its absence in the parental strain. (C) Western-blot detection of the chimeric BcCHSIIIa::eGFP protein in the BcCHSIIIa::eGFP strain. The three lanes correspond to fractions obtained by differential centrifugation at 1 000g (first lane), 36 000g (second lane), and 100 000g (third lane), used to enrich secretory vesicles.

7. Figure S2. Kymographs obtained for all acquisitions of the BcCHSIIIa::eGFP strain

Kymographs were generated along the elongation axis of individual hyphae to capture tip growth dynamics. For each acquisition, the fluorescence signal was measured over a 10 μm line at each time point, for a total duration of 10 minutes (135 time points with a 4.455-second interval). Each kymograph corresponds to one hypha. Kymographs are color-coded to represent fluorescence intensity. Scale bar = 5 μm. The table contains elongation rates as measures from kymographs for the 13 hyphae of the BcCHSIIIa::eGFP strain (Hypha 1 to Hypha 13) and the 5 hyphae of the eGFP^Cyt strain (Hypha 14 to 18). Measures are in nm.min^{-1}.

1. Figure S3. Validation metrics confirm linearity between roundness and growth rate

(A) Summary of key statistical metrics assessing the relationship between roundness and growth rate. Shapiro-Wilk tests support the normality of both distributions. (B, C) Q-Q plots for growth rate (B) and roundness (C). (D) Distribution of residuals from the linear model. (E) Q-Q plot of residuals. (F) Plot of standardized residuals versus fitted values. Orange and red lines indicate thresholds at ±2 and ±3, commonly used to identify moderate and extreme outliers, respectively. (G) Cook's distance values used to identify influential points. Red line indicates the threshold 0.308 (calculated as 4/N with N=13).

2. Figure S4. Cross-correlation analysis confirms consistent fluorescence oscillations across Z-positions

(A) Mean fluorescence intensity over time at the apex of a growing *Botrytis cinerea* hypha, obtained from the acquisition of time-lapse confocal images at four distinct Z-positions (Z1-Z4).

(B) Schematic representation of the hypha and of focal planes. Z1 corresponds to a position outside the hypha, Z2 to Z4 are cross-sections of the cell. (C) Overview of the signal processing pipeline. (D) Processed fluorescence signals corresponding to the four Z-positions. (E) Cross-correlation matrix at lag 0 between processed signals. Black dashed box surrounds the region where the correlation score at lag0 is above 0.5.

3. Figure S5. Periods of fluorescence variation are not linearly correlated with growth rate or the roundness of the region of fluorescence accumulation

(A) Scatter plot of period values measured in Cluster 2 acquisitions versus the elongation rate of the corresponding hyphae. (B) Scatter plot of period values measured in Cluster 2 acquisitions versus the roundness of the region of fluorescence accumulation. Points are numbered to match the corresponding hyphae shown in Figure 1C.

Acknowledgement

We thank Isabelle Gonçalves for her critical reading of the manuscript. The authors also thank Bayer SAS for access to specific equipment.

Declaration of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Resource availability

For further information, contact Adrien Hamandjian (contact@adrien-h.com). Data and code will be made available upon publication in a peer-reviewed journal.

References

1. Goriely, A., and Tabor, M. (2003). Biomechanical models of hyphal growth in actinomycetes. Journal of Theoretical Biology *222*, 211–218. https://doi.org/10.1016/S0022-5193(03)00029-8.

2. Domozych, D.S., Fujimoto, C., and LaRue, T. (2013). Polar Expansion Dynamics in the Plant Kingdom: A Diverse and Multifunctional Journey on the Path to Pollen Tubes. Plants *2*, 148–173. https://doi.org/10.3390/plants2010148.

3. Rensing, S.A. (2016). Plant Evo–Devo: How Tip Growth Evolved. Current Biology *26*, R1228–R1230. https://doi.org/10.1016/j.cub.2016.09.060.

4. Boutillon, A., Banavar, S.P., and Campàs, O. (2024). Conserved physical mechanisms of cell and tissue elongation. Development *151*, dev202687. https://doi.org/10.1242/dev.202687.

5. Braun, M., and Limbach, C. (2006). Rhizoids and protonemata of characean algae: model cells for research on polarized growth and plant gravity sensing. Protoplasma *229*, 133–142. https://doi.org/10.1007/s00709-006-0208-9.

6. Mitchell, A.P. (1998). Dimorphism and virulence in *Candida albicans*. Current Opinion in Microbiology *1*, 687–692. https://doi.org/10.1016/S1369-5274(98)80116-1.

7. Kadosh, D., and Johnson, A.D. (2005). Induction of the Candida albicans Filamentous Growth Program by Relief of Transcriptional Repression: A Genome-wide Analysis. Mol Biol Cell *16*, 2903–2912. https://doi.org/10.1091/mbc.E05-01-0073.

8. Wendland, J. (2020). Sporulation in Ashbya gossypii. Journal of Fungi *6*, 157.
https://doi.org/10.3390/jof6030157.

9. Riquelme, M. (2013). Tip Growth in Filamentous Fungi: A Road Trip to the Apex. Annu. Rev.
Microbiol. *67*, 587–609. https://doi.org/10.1146/annurev-micro-092412-155652.

10. Peñalva, M.A., Zhang, J., Xiang, X., and Pantazopoulou, A. (2017). Transport of fungal RAB11
secretory vesicles involves myosin-5, dynein/dynactin/p25, and kinesin-1 and is independent
of kinesin-3. Mol Biol Cell *28*, 947–961. https://doi.org/10.1091/mbc.E16-08-0566.

11. Takeshita, N., Wernet, V., Tsuizaki, M., Grün, N., Hoshi, H., Ohta, A., Fischer, R., and Horiuchi,
H. (2015). Transportation of Aspergillus nidulans Class III and V Chitin Synthases to the
Hyphal Tips Depends on Conventional Kinesin. PLOS ONE *10*, e0125937.
https://doi.org/10.1371/journal.pone.0125937.

12. Mouriño-Pérez, R.R., Riquelme, M., Callejas-Negrete, O.A., and Galván-Mendoza, J.I. (2016).
Microtubules and associated molecular motors in Neurospora crassa. Mycologia *108*, 515–
527. https://doi.org/10.3852/15-323.

13. Weiner, A., Orange, F., Lacas-Gervais, S., Rechav, K., Ghugtyal, V., Bassilana, M., and
Arkowitz, R.A. (2019). On-site secretory vesicle delivery drives filamentous growth in the
fungal pathogen Candida albicans. Cellular Microbiology *21*, e12963.
https://doi.org/10.1111/cmi.12963.

14. TerBush, D.R., Maurice, T., Roth, D., and Novick, P. (1996). The Exocyst is a multiprotein
complex required for exocytosis in Saccharomyces cerevisiae. *The EMBO journal*, *15*(23),
6483–6494.

15. Guo, W., Grant, A., and Novick, P. (1999). Exo84p Is an Exocyst Protein Essential for Secretion.
Journal of Biological Chemistry *274*, 23558–23564.
https://doi.org/10.1074/jbc.274.33.23558.

16. Zuriegat, Q., Abubakar, Y.S., Wang, Z., Chen, M., and Zhang, J. (2024). Emerging Roles of
Exocyst Complex in Fungi: A Review. Journal of Fungi *10*, 614.
https://doi.org/10.3390/jof10090614.

17. Gonçalves, I.R., Brouillet, S., Soulié, M.-C., Gribaldo, S., Sirven, C., Charron, N., Boccara, M.,
and Choquer, M. (2016). Genome-wide analyses of chitin synthases identify horizontal gene
transfers towards bacteria and allow a robust and unifying classification into fungi. BMC Evol
Biol *16*, 252. https://doi.org/10.1186/s12862-016-0815-9.

18. Wernet, V., Kriegler, M., Kumpost, V., Mikut, R., Hilbert, L., and Fischer, R. (2023).
Synchronization of oscillatory growth prepares fungal hyphae for fusion. eLife *12*, e83310.
https://doi.org/10.7554/eLife.83310.

19. Fukuda, K., Yamada, K., Deoka, K., Yamashita, S., Ohta, A., and Horiuchi, H. (2009). Class III
Chitin Synthase ChsB of Aspergillus nidulans Localizes at the Sites of Polarized Cell Wall
Synthesis and Is Required for Conidial Development. Eukaryot Cell *8*, 945–956.
https://doi.org/10.1128/EC.00326-08.

20. Sánchez-León, E., Verdín, J., Freitag, M., Roberson, R.W., Bartnicki-Garcia, S., and Riquelme,
M. (2011). Traffic of Chitin Synthase 1 (CHS-1) to the Spitzenkörper and Developing Septa in

Hyphae of Neurospora crassa: Actin Dependence and Evidence of Distinct Microvesicle
Populations. Eukaryotic Cell *10*, 683–695. https://doi.org/10.1128/EC.00280-10.

21. Viotti, C. (2016). ER to Golgi-Dependent Protein Secretion: The Conventional Pathway. In
Unconventional Protein Secretion: Methods and Protocols, A. Pompa and F. De Marchis, eds.
(Springer), pp. 3–29. https://doi.org/10.1007/978-1-4939-3804-9_1.

22. Girard, V., Dieryckx, C., Job, C., and Job, D. (2013). Secretomes: The fungal strike force.
PROTEOMICS *13*, 597–608. https://doi.org/10.1002/pmic.201200282.

23. Bani, A., Pioli, S., Ventura, M., Panzacchi, P., Borruso, L., Tognetti, R., Tonon, G., and Brusetti,
L. (2018). The role of microbial community in the decomposition of leaf litter and deadwood.
Applied Soil Ecology *126*, 75–84. https://doi.org/10.1016/j.apsoil.2018.02.017.

24. de Vallée, A., Bally, P., Bruel, C., Chandat, L., Choquer, M., Dieryckx, C., Dupuy, J.W., Kaiser,
S., Latorse, M.-P., Loisel, E., et al. (2019). A Similar Secretome Disturbance as a Hallmark of
Non-pathogenic Botrytis cinerea ATMT-Mutants? Front Microbiol *10*, 2829.
https://doi.org/10.3389/fmicb.2019.02829.

25. Cairns, T.C., Zheng, X., Zheng, P., Sun, J., and Meyer, V. (2021). Turning Inside Out: Filamentous
Fungal Secretion and Its Applications in Biotechnology, Agriculture, and the Clinic. J Fungi
(Basel) *7*, 535. https://doi.org/10.3390/jof7070535.

26. Riquelme, M., and Sánchez-León, E. (2014). The Spitzenkörper: a choreographer of fungal
growth and morphogenesis. Curr Opin Microbiol *20*, 27–33.
https://doi.org/10.1016/j.mib.2014.04.003.

27. Gierz, G., and Bartnicki-garcia, S. (2001). A Three-Dimensional Model of Fungal
Morphogenesis Based on the Vesicle Supply Center Concept. Journal of Theoretical Biology
208, 151–164. https://doi.org/10.1006/jtbi.2000.2209.

28. Egan, M.J., McClintock, M.A., and Reck-Peterson, S.L. (2012). Microtubule-based transport in
filamentous fungi. Curr Opin Microbiol *15*, 637–645.
https://doi.org/10.1016/j.mib.2012.10.003.

29. Xie, Y., and Miao, Y. (2021). Polarisome assembly mediates actin remodeling during polarized
yeast and fungal growth. Journal of Cell Science *134*, jcs247916.
https://doi.org/10.1242/jcs.247916.

30. Riquelme, M., Bredeweg, E.L., Callejas-Negrete, O., Roberson, R.W., Ludwig, S., Beltrán-
Aguilar, A., Seiler, S., Novick, P., and Freitag, M. (2014). The Neurospora crassa exocyst
complex tethers Spitzenkörper vesicles to the apical plasma membrane during polarized
growth. Mol Biol Cell *25*, 1312–1326. https://doi.org/10.1091/mbc.E13-06-0299.

31. Verdín, J., Bartnicki-Garcia, S., and Riquelme, M. (2009). Functional stratification of the
Spitzenkörper of Neurospora crassa. Molecular Microbiology *74*, 1044–1053.
https://doi.org/10.1111/j.1365-2958.2009.06917.x.

32. Steinberg, G. (2007). Hyphal Growth: a Tale of Motors, Lipids, and the Spitzenkörper.
Eukaryotic Cell *6*, 351. https://doi.org/10.1128/EC.00381-06.

33. Araujo-Palomares, C.L., Castro-Longoria, E., and Riquelme, M. (2007). Ontogeny of the
Spitzenkörper in germlings of Neurospora crassa. Fungal Genetics and Biology *44*, 492–503.
https://doi.org/10.1016/j.fgb.2006.10.004.

34. Schuster, M., Treitschke, S., Kilaru, S., Molloy, J., Harmer, N.J., and Steinberg, G. (2012).
Myosin-5, kinesin-1 and myosin-17 cooperate in secretion of fungal chitin synthase. EMBO J
31, 214–227. https://doi.org/10.1038/emboj.2011.361.

35. Bracker, C.E., Murphy, D.J., and Lopez-Franco, R. (1997). Laser microbeam manipulation of
cell morphogenesis growing in fungal hyphae. In Functional Imaging and Optical
Manipulation of Living Cells (SPIE), pp. 67–80. https://doi.org/10.1117/12.274325.

36. Held, M., Kašpar, O., Edwards, C., and Nicolau, D.V. (2019). Intracellular mechanisms of
fungal space searching in microenvironments. PNAS *116*, 13543–13552.
https://doi.org/10.1073/pnas.1816423116.

37. Thomson, D.D., Wehmeier, S., Byfield, F.J., Janmey, P.A., Caballero-Lima, D., Crossley, A., and
Brand, A.C. (2015). Contact-induced apical asymmetry drives the thigmotropic responses of
Candida albicans hyphae. Cell Microbiol *17*, 342–354. https://doi.org/10.1111/cmi.12369.

38. Riquelme, M., Reynaga-Peña, C.G., Gierz, G., and Bartnicki-García, S. (1998). What
Determines Growth Direction in Fungal Hyphae? Fungal Genetics and Biology *24*, 101–109.
https://doi.org/10.1006/fgbi.1998.1074.

39. Köhli, M., Galati, V., Boudier, K., Roberson, R.W., and Philippsen, P. (2008). Growth-speed-
correlated localization of exocyst and polarisome components in growth zones of Ashbya
gossypii hyphal tips. J Cell Sci *121*, 3878–3889. https://doi.org/10.1242/jcs.033852.

40. Zhou, L., Evangelinos, M., Wernet, V., Eckert, A.F., Ishitsuka, Y., Fischer, R., Nienhaus, G.U.,
and Takeshita, N. (2018). Superresolution and pulse-chase imaging reveal the role of vesicle
transport in polar growth of fungal cells. Sci Adv *4*, e1701798.
https://doi.org/10.1126/sciadv.1701798.

41. Soulié, M.-C., Perino, C., Piffeteau, A., Choquer, M., Malfatti, P., Cimerman, A., Kunz, C.,
Boccara, M., and Vidal-Cros, A. (2006). Botrytis cinerea virulence is drastically reduced after
disruption of chitin synthase class III gene (Bcchs3a). Cellular Microbiology *8*, 1310–1321.
https://doi.org/10.1111/j.1462-5822.2006.00711.x.

42. Leroch, M., Mernke, D., Koppenhoefer, D., Schneider, P., Mosbach, A., Doehlemann, G., and
Hahn, M. (2011). Living Colors in the Gray Mold Pathogen Botrytis cinerea: Codon-Optimized
Genes Encoding Green Fluorescent Protein and mCherry, Which Exhibit Bright Fluorescence.
Applied and Environmental Microbiology *77*, 2887–2897.
https://doi.org/10.1128/AEM.02644-10.

43. Li, S., Francini, G., and Magli, E. (2023). Temporal dynamics clustering for analyzing cell
behavior in mobile networks. Computer Networks *223*, 109578.
https://doi.org/10.1016/j.comnet.2023.109578.

44. Roberson, R.W., Abril, M., Blackwell, M., Letcher, P., McLaughlin, D.J., Mouriño-Pérez, R.,
Riquelme, M., and Uchida, M. (2010). Hyphal structure, in cellular and molecular biology of
filamentous fungi. Cellular and Molecular Biology of Filamentous Fungi, 8–24.

45. Guo, M., Kilaru, S., Schuster, M., Latz, M., and Steinberg, G. (2015). Fluorescent markers for
the Spitzenkörper and exocytosis in *Zymoseptoria tritici*. Fungal Genetics and Biology *79*,
158–165. https://doi.org/10.1016/j.fgb.2015.04.014.

46. Takeshita, N., Evangelinos, M., Zhou, L., Serizawa, T., Somera-Fajardo, R.A., Lu, L., Takaya, N.,
Nienhaus, G.U., and Fischer, R. (2017). Pulses of Ca2+ coordinate actin assembly and
exocytosis for stepwise cell extension. Proceedings of the National Academy of Sciences *114*,
5701–5706. https://doi.org/10.1073/pnas.1700204114.

47. Hemelryck, M.V., Bernal, R., Ispolatov, Y., and Dumais, J. (2018). Lily Pollen Tubes Pulse
According to a Simple Spatial Oscillator. Sci Rep *8*, 12135. https://doi.org/10.1038/s41598-
018-30635-y.

48. Rolland, S., Jobic, C., Fèvre, M., and Bruel, C. (2003). Agrobacterium-mediated
transformation of Botrytis cinerea, simple purification of monokaryotic transformants and
rapid conidia-based identification of the transfer-DNA host genomic DNA flanking
sequences. Curr Genet *44*, 164–171. https://doi.org/10.1007/s00294-003-0438-8.

49. Crumière, M., de Vallée, A., Rascle, C., Gillet, F.-X., Nahar, S., van Kan, J.A.L., Bruel, C.,
Poussereau, N., and Choquer, M. (2025). A LysM Effector Mediates Adhesion and Plant
Immunity Suppression in the Necrotrophic Fungus Botrytis cinerea. Journal of Basic
Microbiology *65*, e2400552. https://doi.org/10.1002/jobm.202400552.

50. Yu, J.-H., Hamari, Z., Han, K.-H., Seo, J.-A., Reyes-Domínguez, Y., and Scazzocchio, C. (2004).
Double-joint PCR: a PCR-based molecular tool for gene manipulations in filamentous fungi.
Fungal Genet Biol *41*, 973–981. https://doi.org/10.1016/j.fgb.2004.08.001.

51. García-Nafría, J., Watson, J.F., and Greger, I.H. (2016). IVA cloning: A single-tube universal
cloning system exploiting bacterial In Vivo Assembly. Scientific Reports 6, 27459.
https://doi.org/10.1038/srep27459.

52. Souibgui, E., Bruel, C., Choquer, M., de Vallée, A., Dieryckx, C., Dupuy, J.W., Latorse, M.-P.,
Rascle, C., and Poussereau, N. (2021). Clathrin Is Important for Virulence Factors Delivery in
the Necrotrophic Fungus Botrytis cinerea. Frontiers in Plant Science *12*.

53. Mullins, E.D., Chen, X., Romaine, P., Raina, R., Geiser, D.M., and Kang, S. (2001).
Agrobacterium-Mediated Transformation of Fusarium oxysporum: An Efficient Tool for
Insertional Mutagenesis and Gene Transfer. Phytopathology *91*, 173–180.
https://doi.org/10.1094/PHYTO.2001.91.2.173.

54. Antal, Z., Rascle, C., Cimerman, A., Viaud, M., Billon-Grand, G., Choquer, M., and Bruel, C.
(2012). The Homeobox BcHOX8 Gene in *Botrytis cinerea* Regulates Vegetative Growth and
Morphology. PLOS ONE *7*, e48134. https://doi.org/10.1371/journal.pone.0048134.

55. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,
Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for biological-
image analysis. Nat Methods *9*, 676–682. https://doi.org/10.1038/nmeth.2019.

56. Mary, H., and Brouhard, G.J. (2019). Kappa (κ): Analysis of Curvature in Biological Image Data
using B-splines. Preprint, https://doi.org/10.1101/852772.

57. Mary, H., Rueden, C., and Ferreira, T. (2016). KymographBuilder: Release 1.2.4. (Zenodo).
https://doi.org/10.5281/zenodo.56702.

58. Ross, S.M. (2009). Chapter 2 - DESCRIPTIVE STATISTICS. In Introduction to Probability and
Statistics for Engineers and Scientists (Fourth Edition), S. M. Ross, ed. (Academic Press), pp.
9–53. https://doi.org/10.1016/B978-0-12-370483-2.00007-2.

59. Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M.,
Prettenhofer, P., Weiss, R., Dubourg, V., et al. (2011). Scikit-learn: Machine Learning in Python.
Journal of Machine Learning Research *12*, 2825–2830.

60. Harris, C.R., Millman, K.J., van der Walt, S.J., Gommers, R., Virtanen, P., Cournapeau, D.,
Wieser, E., Taylor, J., Berg, S., Smith, N.J., et al. (2020). Array programming with NumPy. Nature
585, 357–362. https://doi.org/10.1038/s41586-020-2649-2.

61. Seabold, S., and Perktold, J. (2010). Statsmodels: Econometric and Statistical Modeling with
Python. scipy. https://doi.org/10.25080/Majora-92bf1922-011.

62. Altman, N., and Krzywinski, M. (2016). Analyzing outliers: influential or nuisance? Nature
Methods *13*, 281–282. https://doi.org/10.1038/nmeth.3812.

63. Virtanen, P., Gommers, R., Oliphant, T.E., Haberland, M., Reddy, T., Cournapeau, D., Burovski,
E., Peterson, P., Weckesser, W., Bright, J., et al. (2020). SciPy 1.0: fundamental algorithms for
scientific computing in Python. Nat Methods *17*, 261–272. https://doi.org/10.1038/s41592-
019-0686-2.

Secretory vesicles accumulation

Crescent ←→ Spitzenkörper

Botrytis cinerea

Crescent
Fluorescence variations match those of the control strain, likely reflecting intrinsic fluorophore behavior

Spitzenkörper
Fluorescence variations suggest a periodic succession of vesicle accumulation and release events

Methodology

TIME
- Image acquisition
- Fluorescence signal (mean over time)
- Temporal Dynamics Clustering
- Cluster 1 (control-like behavior) | Cluster 2 (distinct behavior)

SPACE
- Measure roundness of fluorescence accumulation (post clustering)
- Crescent shape | Spitzenkörper shape

TIME
- Fast Fourier Transform analysis

A eGFP^{Cyt}

Fluorescence intensity (AU)

Distance to the tip (µm)

B BcCHSIIIa::eGFP

Fluorescence intensity (AU)

Distance to the tip (µm)

C

Hypha 1 Hypha 2 Hypha 3 Hypha 4 Hypha 5 Hypha 6 Hypha 7 Hypha 8 Hypha 9 Hypha 10 Hypha 11 Hypha 12 Hypha 13

Roundness

Acquisitions

D

Count

Roundness

E

Count

Growth rate (nm.min^{-1})

F

Roundness

R²=0.456

Growth rate (nm.min^{-1})

Amplitude spectra	Model and experimental

Hypha 7 — 31.39 / Experimental signal

Hypha 8 — 64.93 / Experimental signal

Hypha 9 — 40.3 / Experimental signal

Hypha 10 — 41.26 / Experimental signal

Hypha 11 — 52.29 / Experimental signal

Hypha 12 — 39.75 / Experimental signal

Hypha 13 — 48.95 / Experimental signal

Amplitude (AU)

Period (s)

Fluorescence intensity (AU)

Time (s)