**Fast and accurate detection of metal resistance genes using MetHMMDb**

**Karol Ciuchcinski[1,*], Mikolaj Dziurzynski[2]**

[1] Department of Environmental Microbiology and Biotechnology, Institute of Microbiology,

Faculty of Biology, University of Warsaw

[2] Institute of Evolutionary Biology, Faculty of Biology, Biological and Chemical Research

Centre, University of Warsaw

* Corresponding author:

Karol Ciuchcinski; phone: +48 225542205; e-mail: k.ciuchcinski@uw.edu.pl

**ABSTRACT**

Heavy metal pollution, driven by industrialization, is a growing environmental challenge with

severe ecological impacts. One of key factors in fighting this issue is microbial resistance to

heavy metals, and its potential for use in bioremediation and environmental monitoring.

Unfortunately, existing detection methods often fail to identify distant homologs or to provide

insight into the function of microbial metal resistance genes (MMRGs). To overcome these

limitations, we developed MetHMMDB - a novel database containing 254 profile Hidden

Markov Models (HMMs) representing 121 microbial MMRGs and functions. Unlike

traditional sequence similarity-based tools, MetHMMDB leverages HMMs to enhance

detection sensitivity and functional specificity across diverse microbial communities.

MetHMMDB was created through iterative database searches, combined with sequence

clustering and structural prediction, combining both recent advances in bioinformatics and

manal curation of data. In its core, the database prioritises functional annotation, ensuring

accurate detection and classification of genes responsible for metal resistance. Our results

indicate that MetHMMDB outperforms sequence-based approaches, identifying more than

twice as many MMRGs in metagenomic datasets, including extreme environments.

Additionally, application to agricultural soil revealed distinct resistance profiles linked to soil

quality. By improving the identification and functional characterization of metal resistance

genes, MetHMMDB facilitates interdisciplinary research in microbial ecology, environmental

monitoring, and bioremediation. This robust and versatile resource represents a significant

advancement in understanding microbial adaptation to heavy metal contamination and supports

the development of effective environmental management strategies. The database is available

at https://github.com/Haelmorn/MetHMMDB.

Heavy metal pollution, driven by industrialization and human activities, is a growing environmental challenge with severe consequences for ecosystems and biodiversity [1–3]. As heavy metal contamination continues to escalate, there is a need for effective solutions for monitoring and mitigating these pollutants. One promising avenue lies in the study of microbial resistance mechanisms, which have the potential to be harnessed for bioremediation and environmental monitoring [4]. Microbial metal resistance is mediated by a diverse set of genes and pathways that enable microorganisms to tolerate or detoxify metal ions in contaminated environments. Identifying these microbial metal resistance genes (MMRGs) is crucial for understanding microbial adaptation, ecological impacts, and for developing biotechnological applications. Current detection methods, however, largely rely on sequence similarity searches against sequence databases like BacMet [5–7], which, while being a valuable source of manually curated data, has not been updated since 2018 [8]. Additionally, sequence-based methods often fail to detect distant homologs or provide functional context, limiting their applicability for analyzing diverse and complex microbial communities. Here we present MetHMMDB, a comprehensive, manually annotated database of 254 profile Hidden Markov Models (HMMs), uniquely designed to detect and characterize bacterial metal resistance genes with enhanced sensitivity and functional specificity in diverse datasets.

The MetHMMDB database was constructed through a systematic, multi-step process starting with 374 experimentally confirmed metal resistance genes from the MetGeneDB database, previously described in our work [9] (Fig 1A). Those sequences served as queries in DIAMOND BLASTP searches against SwissProt (obtained 07.08.2024), using stringent criteria (>90% identity and coverage), yielding 315 new sequences [10]. The combined set of 681 sequences was then used to search NCBI NR (obtained 04.04.2024), identifying 10,344 additional unique sequences and bringing the final dataset to 11,025 proteins. These sequences were then clustered using MMseqs2 [11] with clustering parameters optimized through a

gradient of identity and coverage values (0.5-0.99, 0.02 increments). The 0.90 threshold was

selected, balancing mean cluster size and singleton count (Fig 1B). This resulted in 341 non-

singleton clusters with a mean size of 27.98 proteins. The largest clusters included sequences

encoding genes *mdtC* (811 sequences, represented by protein HCR6247170.1), *mdtB* (632

sequences; P69340), *cutC* (242 sequences; 1X7I), *fieF* (211 sequences; WP_085449461.1), and

*zupT*                    (195                    sequences;                    Q32BU5).

Figure 1. A) Database creation pipeline. Each box represents a separate step in the pipeline. B)

Results of clustering parameters optimisation process. Coverage and identity value cutoffs are

present on X-axis, while mean cluster size and number of singleton clusters can be seen on Y-

axis.

For each non-singleton cluster, a comprehensive functional annotation using multiple

complementary approaches was performed. The analysis combined domain detection using

Pfam_scan.pl (Pfam v.37.1), protein structure prediction via LocalColabFold, and structural

similarity searches against SwissProt using FoldSeek (E-value threshold ≤ 1e-3) [12–14].

Based on the resulting domain architectures and structural homologies to known SwissProt

proteins, we determined the general function, metal specificity, and appropriate gene

nomenclature for each cluster. This process identified 254 clusters associated with metal

resistance mechanisms. Among these, copper-related clusters were most abundant (47),

followed by clusters involved in arsenic and mercury resistance (32 each). For annotated

clusters, multiple sequence alignments were generated using MAFFT (v7.525) and converted

to profile HMMs using HMMER3.4 (http://hmmer.org/) [15]. Each cluster was then assigned

a descriptive name containing the metal specificity, general function, and gene name when

possible. The database, alongside reference data and individual models is available at

https://github.com/Haelmorn/MetHMMDB.

To evaluate the practical utility of MetHMMDB, we compared its search speed to

DIAMOND BLASTP. Although hmmsearch exhibited slower processing times than

DIAMOND, its performance remained within acceptable limits for practical applications

(Supplementary Table 1). To evaluate detection capabilities, we analyzed a metagenome from

a metal-rich Alpine stream (PRJNA689378) containing 215,000+ protein sequences [16].

Using an E-value cutoff of 1e-5, hmmscan with MetHMMDB identified 3,846 metal resistance

genes compared to 1,786 found by DIAMOND blastp with MetGeneDB. Both methods

revealed similar distributions of resistance types, with *czcA*-like cobalt-zinc-cadmium

resistance proteins and *fbpC*-like iron transporters being predominant. The HMM-based

approach demonstrated enhanced sensitivity, successfully detecting previously reported *Acr3*

and *ArsACHM* arsenic resistance genes, notably revealing the presence of multiple copies of

the aioAB arsenite oxidase—a finding not reported in the original study. The methods showed

substantial agreement, with 1,636 proteins (91.6% of blastp results) detected by both

approaches. The HMM-based method primarily missed genes with low sequence conservation

(mean identity 50.1%, coverage 26.3%), including *acn* (aconitate hydratase), *nrsD* (MFS

superfamily transporter), and *pitA* (inorganic phosphate transporter) genes.

Finally, we checked the performance of the database on agricultural soil metagenomes

from BioProject PRJEB22376 (ERR2486626, ERR2486627) to evaluate its ability to

differentiate healthy and unhealthy soil samples. The analysis revealed distinct metal resistance

profiles between metagenomes, with multidrug *mdtBC* genes and iron transport related genes

enriched in healthy soil, while nickel transporters and *fecE*-like iron transporter were more

abundant in unhealthy soil (Supplementary Figure 1). Notably, two *znuC* zinc transporter

variants displayed opposite patterns: a *Bacillus*-derived variant dominated in healthy soil, while

a *Salmonella*-derived variant was more common in unhealthy soil, which might contribute to

perceived soil quality differences (Fig 2).

Figure 2. Comparison of resistance mechanisms between healthy and unhealthy soil samples.

Top half of the plots shows 5 hits highest difference

In summary, MetHMMDB represents the first profile HMM-based database, specifically

designed for MMRGs. By integrating sequence analysis, structural predictions, and manual

curation, we developed 254 functionally annotated models that prioritize molecular function

over traditional gene nomenclature. This approach improves identification accuracy and

enables the detection of resistance determinants across diverse environmental contexts. The

database's utility was demonstrated through applications to both extreme environments and

agricultural soils. These analyses showed that MetHMMDB offers superior detection

sensitivity and functional specificity compared to traditional sequence-based methods,

identifying a broader range of MMRGs while providing ecological insights. Though

computationally more intensive, the database remains practical for large-scale analyses and is

particularly well-suited for environmental monitoring and bioremediation research. With its

focus on functional annotation rather than sequence similarity, MetHMMDB provides a robust

tool for ecological and environmental studies.

**DATA AVAILABILITY**

The datasets generated during and/or analysed during the current study are available in the

GitHub repository, https://github.com/Haelmorn/MetHMMDB.

**REFERENCES**

1.  Das S et al. Heavy Metal Pollution in the Environment and Its Impact on Health: Exploring Green Technology for Remediation. *Environ Health Insights* 2023; **17**: 11786302231201259. https://doi.org/10.1177/11786302231201259

2.  Darham S et al. Antarctic heavy metal pollution and remediation efforts: state of the art of research and scientific publications. *Braz J Microbiol* 2023; **54**: 2011–2026. https://doi.org/10.1007/s42770-023-00949-9

3.  Briffa J, Sinagra E, Blundell R. Heavy metal pollution in the environment and their toxicological effects on humans. *Heliyon* 2020; **6**. https://doi.org/10.1016/j.heliyon.2020.e04691

4.  Roy R et al. An Overview of Bacteria-Mediated Heavy Metal Bioremediation Strategies. *Appl Biochem Biotechnol* 2024; **196**: 1712–1751. https://doi.org/10.1007/s12010-023-04614-7

5.  Li L et al. Genome-resolved metagenomics provides insights into the ecological roles of the keystone taxa in heavy-metal-contaminated soils. *Front Microbiol* 2023; **14**. https://doi.org/10.3389/fmicb.2023.1203164

6.  Ekhlas D et al. Metagenomic comparison of the faecal and environmental resistome on Irish commercial pig farms with and without zinc oxide and antimicrobial usage. *Animal Microbiome* 2023; **5**: 62. https://doi.org/10.1186/s42523-023-00283-3

7.  Balcha ES et al. Shotgun Metagenomics-Guided Prediction Reveals the Metal Tolerance and Antibiotic Resistance of Microbes in Poly-Extreme Environments in the Danakil Depression, Afar Region. *Antibiotics* 2023; **12**: 1697. https://doi.org/10.3390/antibiotics12121697

8.  Pal C et al. BacMet: antibacterial biocide and metal resistance genes database. *Nucleic Acids Res* 2014; **42**: D737–D743. https://doi.org/10.1093/nar/gkt1252

9.  Dziurzynski M et al. Development of the LCPDb-MET database facilitating selection of PCR primers for the detection of metal metabolism and resistance genes in bacteria. *Ecological Indicators* 2022; **145**: 109606. https://doi.org/10.1016/j.ecolind.2022.109606

10. Buchfink B, Reuter K, Drost H-G. Sensitive protein alignments at tree-of-life scale using DIAMOND. *Nat Methods* 2021; **18**: 366–368. https://doi.org/10.1038/s41592-021-01101-x

163    11.    Steinegger M, Söding J. MMseqs2 enables sensitive protein sequence searching for the analysis

164           of massive data sets. *Nat Biotechnol* 2017; **35**: 1026–1028. https://doi.org/10.1038/nbt.3988

165    12.    Mistry J et al. Pfam: The protein families database in 2021. *Nucleic Acids Research* 2021; **49**:

166           D412–D419. https://doi.org/10.1093/nar/gkaa913

167    13.    Mirdita M et al. ColabFold: making protein folding accessible to all. *Nat Methods* 2022; **19**:

168           679–682. https://doi.org/10.1038/s41592-022-01488-1

169    14.    Van Kempen M et al. Fast and accurate protein structure search with Foldseek. *Nat Biotechnol*

170           2024; **42**: 243–246. https://doi.org/10.1038/s41587-023-01773-0

171    15.    Eddy SR. Accelerated profile HMM searches. *PLoS Computational Biology* 2011; **7**.

172           https://doi.org/10.1371/journal.pcbi.1002195

173    16.    Buetti-Dinh A et al. Geochemical and metagenomics study of a metal-rich, green-turquoise-

174           coloured stream in the southern Swiss Alps. *PLOS ONE* 2021; **16**: e0248877.

175           https://doi.org/10.1371/journal.pone.0248877

**AUTHOR CONTRIBUTIONS**

KC and MD conceptualized the study. KC performed data curation, formal analysis, validated

and visualised the data. KC and MD conducted the investigations. KC wrote the original draft.

KC and MD reviewed and edited the manuscript.

**COMPETING INTERESTS**

The authors declare no competing interests.