Shoot nitrate status regulates Arabidopsis shoot growth and systemic transcriptional

responses via shoot adenylate isopentenyltransferase 3

Kota Monden[a†], Takamasa Suzuki[b], Mikiko Kojima[c], Yumiko Takebayashi[c], Daisuke Sugiura[d],

Tsuyoshi Nakagawa[a], Hitoshi Sakakibara[c,d], and Takushi Hachiya[a†]

[a]Department of Molecular and Functional Genomics, Interdisciplinary Center for Science

Research, Shimane University, 1060 Nishikawatsu-cho, Matsue, Shimane 690-8504, Japan

[b]College of Bioscience and Biotechnology, Chubu University, 1200 Matsumoto-cho, Kasugai,

Aichi 487-8501, Japan

[c]RIKEN Center for Sustainable Resource Science, 1-7-22 Suehiro-cho, Tsurumi-ku, Yokohama,

Kanagawa 230-0045, Japan

[d]Graduate School of Bioagricultural Sciences, Nagoya University, Furo-cho, Chikusa-ku, Nagoya,

Aichi 464-8601, Japan

[†]Corresponding authors

Corresponding authors:

Kota Monden (n23d203@matsu.shimane-u.ac.jp)

Takushi Hachiya (takushi.hachiya@life.shimane-u.ac.jp)

Funding:

This study was supported by JSPS KAKENHI [No. 20K05771 and 23K04978 to TH and No.

JP24KJ1709 to KM], by JST SPRING [No. JPMJSP2155 to KM], and by the Sasakawa Scientific

Research Grant from the Japan Science Society [No. 2022-4039 to KM].

Keywords:

Arabidopsis thaliana; cytokinin; IPT3; isopentenyltransferase; micrografting; NR; nitrate

reductase; nitrate signal; nitrate accumulation; systemic regulation

Abstract

Plants systemically regulate growth and gene expression according to their internal nitrate status.

Our previous study reported that shoot nitrate accumulation increases shoot expression of

adenylate *ISOPENTENYLTRANSFERASE 3* (*IPT3*) and shoot levels of N^6-(Δ^2-isopentenyl)

adenine (iP)-type cytokinins (CKs). *IPT3* expression is localized in the phloem, and iP-type CKs,

which are synthesized by IPT3, are phloem-mobile. As CKs are a class of phytohormones that

control growth and genome-wide gene expression, shoot-expressed IPT3 may mediate the

systemic regulation of growth and transcriptomic responses to shoot nitrate status. To examine

this, we developed a novel system to manipulate nitrate levels and *IPT3* expression in a shoot-

specific manner and performed growth analysis, CK determination, and RNA-sequencing. Our

results demonstrated that shoot nitrate accumulation significantly promoted shoot growth and

elevated shoot concentrations of iP ribotides, iP-7-*N*-glucoside, and iP-9-*N*-glucoside through the

action of shoot IPT3. Transcriptomic responses to shoot nitrate accumulation were largely tuned

by shoot IPT3, with opposite effects in shoots and roots. Shoot IPT3 amplified shoot responses

of nitrate-inducible genes and immune response genes to shoot nitrate accumulation, while

dampened root responses of nitrate transport/assimilation genes. This transcriptomic modulation

via shoot IPT3 was accompanied by coherent transcriptional changes in genes encoding mobile

peptides and transcriptional repressors. Here we present a novel scheme integrating shoot nitrate

status responses with CK signaling.

Introduction

48 Land plants use soil nitrate and ammonium as primary nitrogen sources (Hachiya and Sakakibara

49 2017). Root-absorbed nitrate is reduced to ammonium via the sequential actions of nitrate

50 reductase (NR) and nitrite reductase (NiR). Ammonium, whether taken up by the roots or

51 generated through nitrate reduction, is assimilated into Gln and Glu by the enzymes glutamine

52 synthetase (GS) and glutamate synthase (GOGAT). These amino acids serve as precursors for the

biosynthesis of other amino acids. Nitrate serves as a building block for organic nitrogen biosynthesis and as a signaling molecule (Remans et al. 2006; Wang et al. 2004). The addition of exogenous nitrate to nitrate-depleted plants rapidly changes the expression of nitrate-responsive genes, a phenomenon termed as the primary nitrate response (PNR) (Medici and Krouk 2014). In *Arabidopsis*, the nitrate transceptor NITRATE TRANSPORTER 1.1 (NRT1.1) and the transcription-activating nitrate sensor NIN-LIKE PROTEIN 7 (NLP7) play central roles in PNR (Ho et al. 2009; Bouguyon et al. 2015; Liu et al. 2017; Liu et al. 2022). Nitrate uptake via NRT1.1 transiently induces a Ca^{2+} influx, thereby elevating both cytosolic and nuclear Ca^{2+} levels (Riveras et al. 2015; Liu et al. 2017; Wang et al. 2021). NRT1.1 physically interacts with the cyclic nucleotide-gated channel CNGC15 to form a nitrate-activated Ca^{2+}-permeable channel on the plasma membrane (Wang et al. 2021). Additionally, PHOSPHOLIPASE C also contributes to the NRT1.1-mediated increase in cytoplasmic Ca^{2+} levels (Riveras et al. 2015). The nitrate-binding affinity and uptake activity of NRT1.1 are tuned by specific protein kinases and phosphatase, including CBL-INTERACTING PROTEIN KINASEs 8/23, SUCROSE NON-FERMENTING1-RELATED PROTEIN KINASE (SnRK) 2s, and ABA INSENSITIVE 2 (Ho et al. 2009; Hu et al. 2009; Léran et al. 2015; Su et al. 2020). The subgroup III calcium-dependent protein kinases 10/30/32 (CPK10/30/32) perceive Ca^{2+} and phosphorylate NLP7 at S205, causing its nuclear retention by inhibiting nuclear export (Marchive et al. 2013; Liu et al. 2017). Nitrate-bound NLP7, along with transcription factors NLP2/6 and NITRATE REGULATORY GENE 2—which physically interact with NLP7—promotes the transcription of PNR genes including those encoding secondary transcription factors, thereby further altering the expression of downstream target genes (Konishi and Yanagisawa 2013; Xu et al. 2016; Maeda et al. 2018; Liu et al. 2022; Durand et al. 2023). Carbon deficiency leads to NLP7 depletion via SnRK1-mediated phosphorylation, which promotes its degradation, thereby attenuating nitrate signaling (Wang et al. 2022). This suggests that carbon availability modulates PNR.

Besides its role in PNR, nitrate acts as an internal signal of plant nitrogen status,

modulating nitrogen responses under steady-state conditions. In tobacco plants with different NR

activities grown under varying nitrate availability, the shoot-to-root dry weight ratio was

positively correlated with foliar nitrate levels (Scheible et al. 1997). In *Arabidopsis*, abundant

nitrate availability inhibits primary and lateral root growth (Zhang et al. 1998; Signora et al. 2001;

Patterson et al. 2016). Patterson et al. (2016) reported that nitrate application increases the shoot

expression of glutaredoxin (Grx) genes *GrxS3/4/5/8* via Arabidopsis response regulators (ARRs)

1/10/12 for cytokinin (CK) signaling, thereby inhibiting primary root growth. Kobayashi et al.

(2024) suggested that GrxS1–8 are candidate phloem-mobile signals from shoots to roots, and

they bind to TGACG-binding factor 1 (TGA1) and TGA4 (Alvarez et al. 2014) in roots, forming

a corepressor complex with TOPLESS to limit high-affinity nitrate uptake. Meanwhile, Ota et al.

(2020) discovered that under reduced shoot nitrogen, including nitrate depletion, C-terminally

encoded peptide downstream-like 2 (CEPDL2)—a Grx family protein also known as GrxS9—is

induced in the leaf vasculature and subsequently translocated to the roots. The co-activator

complex composed of CEPDL2, its homologs CEPD1/2, and TGA1/4 promotes root nitrate

uptake (Kobayashi et al. 2024). These findings imply that shoot nitrate status influences root

nitrogen acquisition.

In our previous study, to focus on the action of shoot nitrate status, we manipulated

shoot nitrate levels using wild-type *Arabidopsis* and an NR-null mutant (Okamoto et al. 2019).

Subsequent RNA-sequencing (RNA-seq) suggested that shoot nitrate accumulation alone

represses the expression of nitrogen starvation-inducible genes in shoots and roots. Notably, we

observed that shoot nitrate accumulation was accompanied by increases in the shoot expression

of adenylate *ISOPENTENYLTRANSFERASE 3* (*IPT3*) and the shoot levels of N^6-(Δ^2-isopentenyl)

adenine (iP)-type CKs, including iP ribotides (iPRPs), iP-7-*N*-glucoside (iP7G), and iP-9-*N*-

glucoside (iP9G) (Hachiya et al. 2020). Adenylate IPTs are key enzymes that catalyze the

formation of iPRPs using DMAPP as the isoprenoid donor and adenylates as acceptors

(Sakakibara 2021). iP riboside 5′-monophosphate (iPRMP) is then activated to bioactive free base

iP via either the one-step LONELY GUY (LOG) pathway or a two-step pathway involving 5'-

ribonucleotide phosphohydrolase GY3 and CK/purine riboside nucleosidase CPN1/NSH3

(Kojima et al. 2023; Wu et al. 2023). Nitrate-induced expression of *IPT3* is localized mainly in

phloem cells (Miyawaki et al. 2004; Takei et al. 2004a), and iP-type CKs, which are de novo

synthesized by IPT3, are phloem-mobile (Hirose et al. 2008; Bishopp et al. 2011). CKs are a class

of phytohormones that play central roles in regulating growth and development, such as

promoting cell division in shoots and repressing lateral root initiation (Werner et al. 2003; Laplaze

et al. 2007). Multiple transcriptome analyses have demonstrated that CKs regulate specific gene

sets across several tissues and organs (Bhargava et al. 2013). Hence, the systemic regulation of

growth and transcriptomic responses to shoot nitrate status may be mediated via IPT3-synthesized

iP-type CKs. Supporting this idea, several lines of evidence indicate both overlap and crosstalk

between nitrate and CK signaling. For instance, nitrate- and CK-responsive genes have been

found to partially overlap (Sakakibara et al. 2006). In addition, the nitrogen depletion-dependent

induction of shoot *CEPDL2* is enhanced by CKs translocated from roots (Ota et al. 2020).

Furthermore, *trans*-zeatin (*tZ*)-type CKs mediate preferential root growth in nitrate-rich patches

over nitrate-poor patches (Ruffel et al. 2011; Poitout et al. 2018). Based on this context, the

present study examined the systemic responses to shoot nitrate status and their dependence on

shoot IPT3. We developed a novel experimental system to manipulate nitrate levels and *IPT3*

expression in a shoot-specific manner using grafted plants derived from plants lacking NR and/or

IPT3. Using the shoots and roots of the grafted plants, growth analysis, CK species determination,

and RNA-seq were performed. Our results indicated that shoot nitrate accumulation enhances

shoot growth and alters the transcriptomes of shoots and roots, which is elaborately tuned by

shoot IPT3 in the opposite direction in shoots and roots. We herein present a scheme incorporating

shoot nitrate accumulation responses with shoot IPT3 and iP-type CKs.

Materials and Methods

Plant materials

In this study, we used *A. thaliana* accession Columbia (Col) as the control line; NR-null mutant

(NR-null, Wang et al. 2007), *IPT3*-knockout mutant (*ipt3*, Miyawaki et al. 2006), and

IPT3/IPT5/IPT7-knockout mutant (*ipt3,5,7*, Miyawaki et al. 2006); and a transgenic plant

expressing *GFP* driven by the *IPT3* promoter (*proIPT3:GFP*, Takei et al. 2004a). Homozygous

multiple mutants of NR-null *ipt3* and NR-null *proIPT3:GFP* were produced via crossing.

Culture media

Three different solid media were used for *in vitro* plant culture as follows: (i) 2.5 mM NH_4^+

medium, half-strength Murashige and Skoog (1/2MS) modified basal salt mixture without

nitrogen (M531; PhytoTech Labs, Lenexa, KS, USA) supplemented with 1.25 mM $(NH_4)_2SO_4$,

10.0 g L^{-1} sucrose, 1.0 g L^{-1} MES, and 4.0 g L^{-1} gellan gum (Fujifilm Wako, Osaka, Japan) and

adjusted to pH 6.7 with KOH; (ii) 10 mM NO_3^- and 2.5 mM NH_4^+ medium, 1/2MS modified

basal salt mixture without nitrogen (M531; PhytoTech Labs) supplemented with 10 mM KNO_3,

1.25 mM $(NH_4)_2SO_4$, 10.0 g L^{-1} sucrose, 1.0 g L^{-1} MES, and 4.0 g L^{-1} gellan gum (Fujifilm

Wako) and adjusted to pH 6.7 with KOH; and (iii) no nitrogen medium, 1/2MS modified basal

salt mixture without nitrogen (M531; PhytoTech Labs) supplemented with 2.0 g L^{-1} sucrose, 0.5

g L^{-1} MES, and 5.0 g L^{-1} gellan gum (Fujifilm Wako) and adjusted to pH 5.7 with KOH. It should

be noted that pH 6.7 was employed to mitigate ammonium toxicity under 2.5 mM NH_4^+ medium

(Hachiya et al. 2021b) and to promote better growth of NR-deficient plants under 10 mM NO_3^-

and 2.5 mM NH_4^+ medium (Wang et al. 2004).

Transfer experiments using grafted plants

To manipulate nitrate levels and *IPT3* expression in a shoot-specific manner, micrografting and

transfer experiments were performed using Col, NR-null, *ipt3*, NR-null *ipt3*, and *ipt3,5,7* plants

(Figure 1A, B). Surface-sterilized seeds were placed in the dark at 4°C for 3 days to break seed

dormancy. To ensure uniform growth regardless of the NR activity, approximately 120 seeds per

dish were sown in rectangle dishes ($140 \times 100 \times 20$ mm^3; Eiken Chemical Co. Ltd., Tokyo, Japan)

containing 50 mL of 2.5 mM NH_4^+ medium and grown vertically at 22°C for 4 days under a

photosynthetic photon flux density (PPFD) of 25 µmol m^{-2} s^{-1} with a 16/8-h light/dark cycle

(condition 1). Four-day-old seedlings were micrografted aseptically in rectangle dishes containing

50 mL of 2.5 mM NH_4^+ medium using four combinations of scions and rootstocks corresponding

to Col/*ipt3,5,7*, NR-null/*ipt3,5,7*, *ipt3*/*ipt3,5,7*, and NR-null *ipt3*/*ipt3,5,7* (scion line

name/rootstock line name) (Figure 1B). Notably, *ipt3,5,7* was used as the rootstock to avoid the

complementation of shoot-specific CK depletion in *ipt3* by root-derived CKs. The seedlings were

cut perpendicularly at the upper hypocotyl with an injection needle tip (NN-2613S; TERUMO,

Tokyo, Japan) on a membrane filter (HAWP04700; Merck Millipore, Darmstadt, Germany). The

obtained scion was connected with the partner rootstock through a 0.4-mm silicon microtube (1-

8194-04; AS ONE, Osaka, Japan). The grafted plants were incubated in a growth chamber at

27°C for 5 days under a PPFD of 50 µmol m^{-2} s^{-1} with a constant light (condition 2) and further

cultivated at 22°C for 24 h under a PPFD of 25 µmol m^{-2} s^{-1} with a 16/8-h light/dark cycle

(condition 3). Successfully grafted plants without adventitious root formation were selected for

size uniformity. RT-PCR with primers specific for *NIA1*, *NIA2*, *IPT3*, *IPT5*, *IPT7*, and *ACTIN2*

confirmed the successful generation of the desired grafted plants (Figure S1A, B). The grafted

plants were transferred to round dishes (diameter, 90 mm; depth, 20 mm; AS ONE) containing 30

mL of 10 mM NO_3^- and 2.5 mM NH_4^+ medium and grown for 24 h under a PPFD of 25 µmol

m^{-2} s^{-1} with a constant light to promote nitrate accumulation in the grafted plants (condition 4).

Finally, the plants were transferred to round dishes containing 30 mL of no nitrogen medium and

grown for 5 days under a PPFD of 25 µmol m^{-2} s^{-1} with a constant light (condition 5) to allow

nitrate accumulation in the shoots of NR-null/*ipt3,5,7* and NR-null *ipt3*/*ipt3,5,7* but not in

Col/*ipt3,5,7* and *ipt3*/*ipt3,5,7*. A low light intensity of 25 µmol m^{-2} s^{-1} was employed to mitigate

stress, given that high light intensity exacerbates stress caused by nitrogen depletion (Iwagami et

al. 2022). The surfaces of no nitrogen medium were covered with cellophane sheets to allow

nondestructive sampling of the roots (Hachiya et al. 2021a). It should be noted that all plants were

grown under identical external nitrogen conditions to ensure uniform biosynthesis of systemic

signaling molecules responsive to external nitrogen availability—such as tZ-type CKs (Osugi et

al. 2017) and CEP peptides (Tabata et al. 2014)—thereby isolating the systemic effects of shoot

nitrate status and enabling clearer interpretation of the results. Additionally, external nitrogen

sources such as ammonium could enrich internal organic nitrogen pools and inadvertently activate

various nitrogen signaling, potentially masking the effects of shoot nitrate accumulation. To avoid

this, we manipulated shoot nitrate levels without supplying external nitrogen to the plants

(condition 5).

Growth analysis

Shoots and roots were weighed on a precision balance (HR-202i; A&D, Tokyo, Japan), and shoots

were scanned at a resolution of 600 dpi (GT-X980; EPSON, Tokyo, Japan) to measure the rosette

diameter. Leaf blades were harvested from the shoots and scanned at a resolution of 600 dpi to

measure the blade area of the cotyledon and true leaf. The rosette diameter and leaf blade area

were measured using ImageJ version 1.52p (US National Institutes of Health, Bethesda, MD,

USA).

Determination of the total nitrogen content

The total nitrogen concentration was quantified as previously described (Monden et al. 2022).

Harvested shoots and roots were oven-dried at 80°C for 2 days, and the dried samples were

ground using a spoon and stored in a desiccator before use. The dried powder was weighed and

encapsulated in a tin boat. Nitrogen concentrations were measured using a CN analyzer (Vario

EL III; Elementar Analysensysteme GmbH, Hanau, Germany).

Determination of the nitrate content

Nitrate content was quantified as previously described (Okamoto et al. 2019) with slight

modifications. Frozen samples were ground with a Multi-Beads Shocker (Yasui Kikai, Osaka,

Japan) using zirconia beads (diameter, 5 mm). The powder samples were boiled with 10 volumes

of Milli-Q water at 100°C for 20 min. Five microliters of the supernatant or 5 µL of standard

solution as dilution series of potassium nitrate was mixed with 40 µL of 0.05% (w/v) salicylic

acid in sulfuric acid. The resulting mixture was then incubated at room temperature for 20 min.

A mock treatment of 40 µL of sulfuric acid alone was added to 5 µL of the supernatant. One

microliter of 8% (w/v) NaOH solution was added to the mixture, and the nitrate content was

determined by measuring the absorbance at 410 nm using a plate reader (CORONA SH-9000Lab,

Hitachi High-Tech Corp., Tokyo, Japan).

Observation of fluorescent signals

Fluorescent imaging was performed using an All-in-one fluorescence microscope (BZ-X710,

KEYENCE, Osaka, Japan) with a 4× objective (CFI Plan Fluor DL 4×/0.13, Nikon, Tokyo, Japan)

and a metal halide lamp (No. 91056, KEYENCE). A narrow-band GFP filter (ex. 480/20; em.

510/20, 49020, Chroma, Brattleboro, VT, USA) and Cy5 filter (ex. 620/60; em. 700/75, 49006,

Chroma) were used to detect the fluorescence of GFP and chlorophyll, respectively.

Determination of CK species

The extraction and determination of CKs were performed with slight modifications to the method

described by Kojima et al. (2009). Frozen tissue was ground into powder using a TissueLyser

(Qiagen, Hilden, Germany) with a 5 mm zirconia bead. Samples were extracted in 1 ml of

methanol : formic acid : water (15 : 1 : 4), with stable isotope-labeled internal standards. After

incubation at –30°C for ≥16 h, samples were centrifuged at 10,000 × g, for 15 min. The

supernatant was collected, and the pellet was re-extracted with 0.2 ml of the same solvent.

Combined extracts were purified using Oasis HLB 96-well plates (Waters, Milford, MA, USA)

on a STAR (HAMILTON, Reno, NV, USA), followed by evaporation and reconstitution in 1 ml

of 1 M formic acid. Fractionation was performed on an Oasis MCX 96-well plate (Waters). ABA,

auxins, and gibberellins were eluted with methanol (Elution 1); CK nucleotides with 0.35 M

ammonia (Elution 2); and CK nucleobases, nucleosides, and glucosides with 0.35 M ammonia in

60% methanol (Elution 3). All fractions were dried by evaporation. For CK analysis, Elution 3

was reconstituted in 50 μl of 0.1% acetic acid. Elution 2 was reconstituted in 0.84 ml of 0.1 M

CHES-NaOH (pH 9.8), treated with $1 U \mu l^{-1}$ of alkaline phosphatase (Oriental Yeast, Tokyo,

Japan), incubated at 37°C for 1 h, neutralized, desalted via Oasis HLB plate, evaporated, and

reconstituted in 50 μl of 0.1% acetic acid. CKs were analyzed using an ultra-performance liquid

chromatography–tandem quadrupole mass spectrometry system (AQUITY UPLC

System/XEVO-TQS; Waters) with an octadecylsilyl column (ACQUITY UPLC HSS T3, 1.8 μm,

2.1 mm × 100 mm; Waters). CK species were separated at a flow rate of 0.25 mL min-1 with

linear gradients of solvent A (0.06% acetic acid) and solvent B (0.06% acetic acid in methanol)

set according to the following profile: 0 min, 95% A + 5% B; 7min, 60%A + 40% B; 10 min, 1%

A + 99% B; 10 min, 1% A + 99% B; 13 min,95% A + 5% B.

Determination of the total protein content

The total protein content was determined as previously described (Hachiya et al. 2016). Frozen

samples were ground with a Multi-Beads Shocker using zirconia beads (diameter, 5 mm). Total

proteins were extracted with 10 vol. of Laemmli buffer (2% (w/v) SDS, 62.5 mM Tris-HCl

(pH 6.8), 10% (v/v) glycerol, and 0.0125% (w/v) bromophenol blue) containing Halt™ protease

inhibitor cocktail (Thermo Fisher Scientific, Waltman, MA, USA), followed by incubation at

95°C for 5 min. Extracts were centrifuged at 20,400 × g at room temperature for 10 min, and 10

μL of each aliquot was suspended in 500 μL of deionized water. Next, 100 μL of 0.15% (w/v)

sodium deoxycholate was added, and the mixture was incubated at room temperature for 10 min.

One hundred microliters of 72% (v/v) trichloroacetic acid was added, followed by incubation at

room temperature for 15 min and centrifugation at $20,400 \times g$ for 10 min. The precipitates were

air-dried and suspended in 25 µL of deionized water. The suspension was analyzed with a TaKaRa

BCA Protein Assay Kit (T9300A, TaKaRa, Kusatsu, Japan) according to the manufacturer's

instructions. Protein content was determined by measuring the absorbance at 562 nm using a plate

reader (Hitachi High-Tech Corp.).

RNA extraction

Frozen samples were ground with a Multi-Beads Shocker using zirconia beads (diameter, 5 mm).

Total RNA was extracted using an RNeasy Plant Mini Kit (Qiagen) according to the

manufacturer's instructions. For RNA-seq library preparation, RNA was purified via on-column

DNase digestion (Qiagen).

RNA-seq

RNA quality was evaluated using a Qubit RNA IQ Assay Kit with a Qubit 4 Fluorometer (Thermo

Fisher Scientific). The RNA samples for which the RNA IQs ranged 9.4–10.0 were used for RNA-

seq. cDNA libraries were constructed using a NEBNext Ultra II RNA Library Prep with Sample

Purification Beads (E7775S, New England Biolabs, Ipswich, MA, USA), a NEBNext Poly(A)

mRNA Magnetic Isolation Module (E7490S, New England Biolabs), and a NEBNext Multiplex

Oligos for Illumina (E7710S, New England Biolabs) according to the manufacturer's instructions.

These cDNA libraries were sequenced using NextSeq 500 (Illumina, San Diego, CA, USA), and

the produced bcl files were converted to fastq files using bcl2fastq (Illumina). The reads were

analyzed according to a previously described method (Notaguchi et al. 2014) and mapped to the

Arabidopsis reference (TAIR10) using Bowtie (Langmead et al. 2009) with the following options:

"--all --best --strata." The mapped reads were processed using iDEP version 0.96 and 2.01 (Ge et

al. 2018). For data normalization, the read counts data were transformed with rlog and normalized

by subtracting the average expression for each gene according to the default setting of iDEP. For

DEG extraction, DESeq2 was run with a false-discovery rate cutoff of 0.1. The enrichment

analysis of each gene set was conducted using Metascape (Zhou et al. 2019) according to the

default setting.

RT-qPCR

Reverse transcription (RT) was performed using a ReverTraAce qPCR RT Master Mix with

gDNA Remover (Toyobo, Osaka Japan). RNA solution containing 0.5 μg of template RNA

served as the input. Synthesized cDNA was then diluted 5-fold with water and used for

quantitative PCR (qPCR). RT-qPCR was performed on a QuantStudio 1 (Thermo Fisher

Scientific) with KOD SYBR qPCR Mix (Toyobo). Relative transcript levels were calculated

using the comparative cycle threshold method, with *ACTIN3* as the internal standard (Hachiya et

al. 2021b). Primer sequences are presented in Table S1.

Statistical analysis

Unpaired two-tailed Welch's *t*-test and Tukey–Kramer multiple comparison test were performed

using R software v.2.15.3 (R Foundation for Statistical Computing, Vienna, Austria).

Results and discussion

Manipulation of nitrate levels and *IPT3* expression in a shoot-specific manner

After nitrogen removal (condition 5), nitrate concentrations generally decreased over time in the

shoots and roots of all plants (Figure 1C, D). At 3 and 5 days after nitrogen removal, the shoot

nitrate concentrations were higher in NR-null/*ipt3,5,7* and NR-null *ipt3*/*ipt3,5,7* plants than in the

other plants (Figure 1C), whereas root nitrate was depleted to the same level in all plants (Figure

1D). Meanwhile, at 5 days after nitrogen removal, total protein and total nitrogen levels in shoots

and roots differed little among the plants (Figure 1E–H). RT-qPCR revealed that shoot *IPT3*

expression was significantly higher in NR-null/*ipt3,5,7* than in Col/*ipt3,5,7* (Figure 1I). Only

minor differences in transcripts per million (TPM) of shoot *IPT1/4/5/6/7/8* and root *IPT1/4/6/8*

were observed among the four grafted plants (Figure S1C), suggesting that adenylate *IPT* genes

other than *IPT3* did not compensate for the absence of shoot IPT3. To visualize the tissue-specific

induction of *IPT3* as a result of shoot NR deficiency, scions from *proIPT3:GFP* and NR-null

proIPT3:GFP plants were grafted onto *ipt3,5,7* rootstocks, followed by the same transfer

experiment as before. Shoot nitrate accumulation and *GFP* induction were observed in NR-null

proIPT3:GFP/*ipt3,5,7* plants 5 days after nitrogen removal (Figure 1J, K), with strong GFP

fluorescence localized in the leaf vein (Figure 1L). These findings suggest that *IPT3* induction

due to shoot nitrate accumulation occurs mainly in the vasculature. Given that NLP7 promotes

the *IPT3* transcription in response to exogenous nitrate application (Maeda et al. 2018; Liu et al.

2022), it is plausible that *IPT3* induction in NR-null/*ipt3,5,7* is also mediated by NLP7. Overall,

we successfully developed a novel system that manipulates nitrate levels and *IPT3* expression in

a shoot-specific manner, with minimal impact on the total protein and total nitrogen levels within

the plant, in the absence of external nitrogen sources. This system allows the analysis of systemic

responses to shoot nitrate status and their dependence on shoot IPT3 in the following experiments.

All experiments described below were conducted using samples collected 5 days after nitrogen

removal (condition 5).

Shoot nitrate accumulation elevates shoot levels of iP-type CKs via shoot IPT3

To assess the impact of shoot nitrate status and shoot *IPT3* expression on CK activity, we

measured the concentrations of iP-type and *t*Z-type CKs. The experiments were performed using

the shoots and roots of grafted plants, with three biological replicates collected independently

(Figure 2) or simultaneously (Figure S2). The levels of iPRPs, iP7G, iP9G, *t*Z ribotides (*t*ZRPs),

*t*Z-7-*N*-glucoside (*t*Z7G), *t*Z-9-*N*-glucoside (*t*Z9G), *t*Z-*O*-glucoside (*t*ZOG), and *t*ZR-*O*-

glucoside (*tZROG*) were consistently lower in the shoots of *ipt3*/*ipt3,5,7* and NR-null

ipt3/*ipt3,5,7* plants than in those of Col/*ipt3,5,7* and NR-null/*ipt3,5,7* plants (Figure 2C–E, 2H–L

and Figure S2C–E, S2H–L). Meanwhile, little difference was observed in the root levels of CK

species among the grafted plants. These observations indicated that shoot IPT3 is crucial for

maintaining the shoot levels of *tZ*-type and iP-type CKs, but it has only a minor effect on root CK

levels. Notably, the shoot levels of iPRPs, iP7G, and iP9G were significantly increased by shoot

NR deficiency in the presence of shoot IPT3 but not in its absence (Figure 2C–E and Figure S2C–

E). This finding suggests that shoot nitrate accumulation elevates the shoot levels of these iP-type

CKs via shoot IPT3. However, no significant increase was observed in the levels of iP, the

biologically active form (Figure 2A and Figure S2A). In *Arabidopsis*, seven LOG enzymes with

phosphoribohydrolase activity can directly convert iPRMP into iP (Kuroha et al. 2009; Tokunaga

et al., 2011). Thus, elevated levels of iPRPs in the shoot likely serve as a reservoir of latent CK

activity, with the potential to promote growth depending on the spatial and temporal activity of

LOG enzymes. Given that Arabidopsis *LOG* genes exhibit distinct spatial expression patterns

(Kuroha et al. 2009), the iPRPs accumulated in response to shoot nitrate accumulation may be

locally converted to iP in cells or tissues expressing specific LOG isoforms. While our CK

quantifications were performed using whole shoots and roots, it is possible that localized changes

in iP concentrations—especially in actively growing regions such as the shoot apical meristem or

the basal zone of young leaves—were not captured. Targeted sampling of these regions may

therefore be necessary to reveal the spatial dynamics of iP production in response to shoot nitrate

status.

Shoot nitrate accumulation enhances shoot growth via shoot IPT3

To determine the effects of shoot nitrate status and shoot *IPT3* expression on plant growth, we

analyzed growth parameters using all grafted plants from five independent experiments (Figure

3). Shoot fresh weight, leaf blade area, rosette diameter, and leaf number were significantly lower

in *ipt3*/*ipt3,5,7* and NR-null *ipt3*/*ipt3,5,7* plants than in Col/*ipt3,5,7* and NR-null/*ipt3,5,7* plants

(Figure 3B–E). Meanwhile, no significant difference was observed in the root fresh weight among

the grafted plants (Figure 3F). These observations indicate that shoot IPT3 generally contributes

to shoot growth. Importantly, shoot appearance, shoot fresh weight, and leaf blade area were

significantly increased by shoot NR deficiency in the presence of shoot IPT3 but not in its absence

(Figure 3A–C). A similar trend was observed for the rosette diameter, although the differences

were not statistically significant (Figure 3D). These patterns correspond to the findings for shoot

iPRPs, iP7G, and iP9G concentrations (Figure 2C–E and Figure S2C–E). Actually, shoot fresh

weight and leaf blade area were strongly correlated with levels of these iP-type CKs (Figure S3).

Overall, these findings suggest that shoot nitrate accumulation promotes shoot growth by

increasing IPT3 and iP-type CK biosynthesis in the shoot. It is well established that in Arabidopsis

plants, root-supplied nitrate stimulates the biosynthesis of *t*Z-type CKs in the root and facilitates

their transport to the shoot, thereby promoting shoot growth (Kiba et al. 2013; Osugi et al. 2017;

Sakakibara, 2021). Meanwhile, in the present study, shoot nitrate accumulation did not

significantly increase *t*Z-type CK levels in both shoots and roots (Figure 2F–M and Figure S2F–

M), arguing against their involvement in shoot growth regulation in response to internal nitrate

status. Interestingly, the fresh weight ratios of shoot to root were elevated by shoot NR deficiency

in the presence of shoot IPT3 but not in its absence (Figure 3G). Given that, in general, biomass

ratios of shoot to root are positively correlated with overall nitrogen availability within and

outside the plants (Andrews et al. 2006; Krapp et al. 2011; Hachiya et al. 2014), our findings

support the idea that shoot nitrate status serves as an indicator of nitrogen availability.

Furthermore, since CK action is known to enhance sink activity in shoots (Werner et al. 2008), it

is plausible that shoot nitrate status influences biomass allocation between shoot and root via

shoot IPT3-mediated CK signaling.

389 Transcriptomic responses to shoot nitrate accumulation are tuned via shoot IPT3 with

390 opposite effects in shoots and roots

To investigate the effects of shoot nitrate status and shoot *IPT3* expression on systemic responses,

we performed RNA-seq using the shoots and roots of grafted plants, with three biological

replicates collected independently. Reads per million mapped reads (RPM; Table S2) and

normalized transcript levels (Table S3) were calculated according to the read counts. To identify

genes regulated by shoot nitrate accumulation and further modulated by shoot IPT3, we extracted

differentially expressed genes (DEGs) between NR-null/*ipt3,5,7* and Col/*ipt3,5,7* and between

NR-null *ipt3*/*ipt3,5,7* and *ipt3*/*ipt3,5,7* with a minimum fold change of 1.5 (Figure 4A and Figure

S4 and Table S4). Comparison of these DEGs between shoots and roots revealed minimal overlap

(Figure S4), indicating that shoots and roots express distinct sets of genes regulated by shoot

nitrate accumulation. Hence, we analyzed the shoot and root transcriptomes separately. Notably,

ɫin shoots, the number of DEGs was larger between NR-null/*ipt3,5,7* and Col/*ipt3,5,7* than

between NR-null *ipt3*/*ipt3,5,7* and *ipt3*/*ipt3,5,7* (Figure 4A). By contrast, in roots, the number of

DEGs was lower between NR-null/*ipt3,5,7* and Col/*ipt3,5,7* than between NR-null *ipt3*/*ipt3,5,7*

and *ipt3*/*ipt3,5,7* (Figure 4A). The shoot transcriptional responses of 631 and 663 DEGs to shoot

NR deficiency were decreased by shoot IPT3 deficiency (Figure 4B, C), whereas the root

responses of 189 and 126 DEGs were augmented by shoot IPT3 deficiency (Figure 4D, E). Three

independent RNA-seq results exhibited similar trends; therefore, only the mean transcript levels

are presented in subsequent figures for clarity, unless otherwise stated. k-means clustering

identified 11 out of 16 clusters (clusters 1, 2, 3, 6, 7, 8, 11, 12, 14, 15, and 16) in which gene

expression was significantly altered by shoot NR deficiency, with their responses decreased by

shoot IPT3 deficiency in shoots but increased in roots (Figure S5, S6 and Table S5). In summary,

shoot transcriptional responses to shoot nitrate accumulation are generally enhanced by shoot

IPT3, whereas root responses are dampened by shoot IPT3.

Enrichment analyses of the 11 clusters from Figure S5 and S6 revealed a diverse set

of genes significantly regulated by shoot nitrate and shoot IPT3. In shoots, the cluster 6 and 8

enriched terms related to translation, primary carbon metabolism, and biotic stress response such

as "ath03010: Ribosome", "WP2621: Glycolysis", and "GO:0009617: response to bacterium"

(Figure S5A). Previously, we reported that shoot nitrate accumulation is accompanied by the

upregulation of ribosome-associated genes in shoots (Hachiya et al. 2020). It is widely known

that CK signal upregulates genes encoding ribosomal proteins and elongation factors (Sakakibara

et al. 2006). Hence, translation may be activated by shoot nitrate accumulation, some of which

are mediated by shoot IPT3 (Figure S5B). Meanwhile, terms related to wounding and jasmonic

acids were overrepresented in both cluster 3 and 7 (Figure S5A). Thus, stress responses associated

with jasmonic acids signaling could be downregulated by shoot nitrate accumulation, which is

enhanced by shoot IPT3 (Figure S5B). In roots, the cluster 11, 12, and 14 predominantly featured

terms related to abiotic and biotic stresses, particularly hypoxia (Figure S6A). The cluster 15 and

16 enriched terms associated with translation, biosynthesis, and nitrate transport (Figure S6A).

Notably, the genes encoding nitrate reductases (*NIA1*, *NIA2*) and high-affinity nitrate transporters

(*NRT2.1*, *NRT2.4*, *NRT2.5*, *NRT3.1*) were included in the cluster 16 (Table S5). Thus, shoot

nitrate accumulation suppresses anabolism including nitrogen assimilation in the roots, the

responses of which are dampened by shoot IPT3 (Figure S6B).

Shoot nitrate accumulation induces shoot *ARR2* expression via shoot IPT3

We examined the shoot expression of CK-inducible genes using a shared list across multiple

studies (Bhargava et al. 2013). However, our survey failed to detect their transcriptional changes

associated with shoot NR and/or IPT3 deficiency (Figure S7A and Table S6). Notably, previous

studies analyzed early transcriptomic responses to exogenous application of *t*Z and benzyladenine

(Bhargava et al. 2013); therefore, conventional gene lists might be unsuitable for evaluating

steady-state responses to alterations of endogenous iP-type CKs. Our meticulous investigation

discovered that shoot expression of the type-B response regulator gene *ARR2* was induced by

shoot NR deficiency in the presence of shoot IPT3 but not in its absence (Figure S7B), a finding

corroborated by RT-qPCR (Figure S7C). Choi et al. (2010) demonstrated that CK-activated

ARR2 enhances Arabidopsis immunity through direct binding with the transcription factor TGA3.

In Figure S5A, the cluster 8 derived from clustering was overrepresented in terms such as

"GO:0009617: response to bacterium" and "GO:0009627: systemic acquired resistance" (Figure

S5A). Thus, shoot nitrate accumulation might enhance immunity by upregulating *ARR2* via shoot

IPT3. Furthermore, given that *ARR2*-overexpressing plants exhibit larger rosette leaves and

biomass (Arnaud et al. 2017), it is plausible that shoot nitrate accumulation could promote shoot

growth partly via shoot IPT3–ARR2 cascade.

Transcriptional responses of nitrate–CPKs–NLPs-regulatory genes to shoot nitrate

accumulation are enhanced via shoot IPT3

Previously, we reported that the expression of nitrate-responsive genes is influenced by shoot

nitrate accumulation (Okamoto et al. 2019), prompting our focus on these genes. In shoots, the

transcript levels of genes induced by nitrate itself (Wang et al. 2004) were significantly increased

by shoot NR deficiency, and this induction was attenuated by shoot IPT3 deficiency (Figure 5A

and Table S7). By contrast, there was minimal variation in the root expression of nitrate-inducible

genes among the plants. Given that CPK10/30/32, NLP7, and its homologs (NLPs) govern nitrate-

dependent gene expression (Liu et al. 2017; Liu et al. 2022), we analyzed the expression of genes

induced by nitrate–CPKs–NLPs signaling. In shoots, the genes induced via CPK10/30/32 and

those co-induced via all NLPs exhibited similar expression patterns as the nitrate-inducible genes

(Figure 5A–C and Table S7). Meanwhile, none of the genes induced exclusively by individual

NLPs exhibited different expression patterns among the plants (Figure S8 and Table S8). In roots,

the expression of genes regulated via CPK10/30/32 and NLPs showed little variation among the

plants (Figure 5B, C and Figure S8 and Table S7, S8). These findings suggest that shoot nitrate

accumulation upregulates nitrate-inducible genes via CPK10/30/32 and multiple NLPs

specifically in the shoots, with their responses enhanced by shoot IPT3.

Transcriptional responses of nitrogen starvation-inducible genes to shoot nitrate

accumulation are tuned via shoot IPT3 in the opposite direction in shoots and roots

Previously, we reported that shoot nitrate accumulation alleviates nitrogen starvation responses

(Okamoto et al. 2019). Thus, we analyzed the responses of two independent sets of nitrogen

starvation-inducible gene (Krapp et al. 2011; Kiba et al. 2018) (Table S9). Although these gene

sets are quite distinct (Figure S9A), shoot NR deficiency consistently repressed shoot expression

of nitrogen starvation-inducible genes, with this repression slightly alleviated by shoot IPT3

deficiency (Figure S9B, C). The nitrogen starvation-inducible genes common to both gene sets

exhibited a similar but stronger tendency (Figure 5D). Meanwhile, the root expression of nitrogen

starvation inducible-genes were significantly repressed by shoot NR deficiency only in the

absence of shoot IPT3 (Figure 5D and S9B, C). Overall, the repression of nitrogen starvation-

inducible genes to shoot nitrate accumulation is enhanced by shoot IPT3 in the shoot, but

dampened in the root.

Several genes involved in nitrogen transport and assimilation in the root are often

induced under nitrogen starvation (Krapp et al. 2011; Kiba et al. 2018), and thus, we focused on

their root expression. Most of the genes involved in high-affinity nitrate transport and nitrogen

assimilation were slightly downregulated by shoot NR deficiency, and these responses were

remarkably intensified by shoot IPT3 deficiency (Figure 5E). Thus, the degree of downregulation

in root nitrate utilization due to shoot nitrate accumulation is largely dependent on shoot IPT3.

By contrast, genes involved in dual-affinity/low-affinity nitrate transport (*NRT1.1* and *NRT1.2*)

(Liu et al. 1999) were upregulated by shoot NR deficiency regardless of the presence of shoot

IPT3 (Figure 5E). Because NRT1.1 and NRT1.2 can facilitate the uptake of auxin and abscisic

acid, respectively (Krouk et al. 2010; Kanno et al. 2012), their upregulation may not necessarily

be related to nitrate acquisition.

Systemic regulation of nitrogen starvation-inducible genes regulated by shoot nitrate

accumulation may be mediated via transcriptional repressors and mobile peptides

Nitrate signaling induces nitrate-inducible GARP-type transcriptional repressor 1 clade genes

(*NIGT1.1/1.2/1.3/1.4*) and *LOB DOMAIN-CONTAINING PROTEIN 37/38/39* genes

(*LBD37/38/39*) via NLPs (Rubin et al. 2009; Marchive et al. 2013; Kiba et al. 2018; Maeda et al.

2018; Safi et al 2021). These translational products act as repressors to downregulate nitrogen

starvation-inducible genes. Thus, we examined these repressor genes. In the shoot, the transcript

levels of *NIGT1.1* and *LBD37/38* were increased by shoot NR deficiency, and this induction was

attenuated by shoot IPT3 deficiency (Figure 5F and S10A). The expression of these repressors

exhibited a strong negative correlation with that of nitrogen starvation-inducible genes in the

shoot (Figure S10B). In the root, the transcript levels of *NIGT1.1/1.2* and *LBD37/38* were

increased by shoot NR deficiency, and this induction was enhanced by shoot IPT3 deficiency

(Figure 5G and S10C). Interestingly, root *LBD39* expression was induced by shoot NR deficiency

only in the absence of shoot IPT3. These repressors also showed a strong negative correlation

with nitrogen starvation-inducible genes in the root (Figure S10D). These findings suggest that

shoot nitrate accumulation and shoot IPT3 regulate the systemic regulation of nitrogen starvation-

inducible genes via these repressors. It is valuable to explore whether iP-type CKs synthesized

by shoot IPT3 systemically regulate the root expression of repressor genes.

Next, we focused on candidate shoot-to-root mobile peptides regulating systemic

nitrogen responses, i.e., CEPD1/2, CEPDL2, and GrxS1–8 (Ohkubo et al. 2017; Ota et al. 2020;

Kobayashi et al. 2024) (Figure 5H and S11). The co-activator complexes composed of

CEPD1/2/CEPDL2 and TGA1/4 promote root nitrate uptake via upregulation of *NRT2.1* and

CEPH, whereas the co-repressor complexes of GrxS1–8 and TGA1/4 repress it (Kobayashi et al.

2024). In the shoot, *CEPDL2* transcript levels were markedly reduced by shoot NR deficiency,

and this repression was attenuated by shoot IPT3 deficiency (Figure 5H and S11A). In contrast,

shoot *GrxS3* expression was upregulated by shoot NR deficiency, and this induction was slightly

enhanced in the absence of shoot IPT3. Interestingly, shoot *GrxS5* expression was induced by

shoot NR deficiency only in the absence of shoot IPT3. The root expression of *NRT2.1* and *CEPH*

was positively correlated with shoot expression of *CEPDL2*, whereas negatively correlated with

shoot expression of *GrxS3* and *GrxS5* (Figure 5H and S11). These findings suggest that shoot

nitrate accumulation and shoot IPT3 regulate the systemic regulation of root high-affinity nitrate

uptake via the action of shoot-to-root mobile peptides.

Conclusion: a scheme integrating shoot nitrate accumulation responses with shoot IPT3

Our experimental system elucidated the systemic responses to shoot nitrate status and their

dependence on shoot IPT3. A novel scheme for the systemic responses is summarized in Figure

6: shoot nitrate accumulation enhances shoot growth and the shoot expression of *IPT3*, nitrate–

CPKs–NLPs-inducible genes including nitrate uptake/assimilation genes, and immune response

genes, with their responses enhanced by shoot IPT3. Meanwhile, shoot nitrate accumulation

decreases the root expression of nitrogen starvation-inducible genes and nitrate

uptake/assimilation genes, with their responses dampened by shoot IPT3. Hence, under shoot

nitrate accumulation, shoot IPT3 would promote nitrogen utilization at the whole-plant scale,

providing organic nitrogen crucial for shoot growth enhancement. Notably, plant CK levels are

generally reduced by prolonged abiotic stresses including salinity, drought, chilling, and heat

(Cortleven et al. 2019), during which shoot growth is usually suppressed. Thus, shoot growth and

nitrogen utilization, which are tuned by shoot IPT3 depending on shoot nitrate status, would be

further fine-tuned via modulation of CK levels depending on abiotic stresses. Such crosstalk in

CK signaling would optimize plant growth and nitrogen utilization in fluctuating environments.

Despite these insights, it remains challenging to distinguish between primary and secondary

responses to shoot nitrate accumulation, as well as to evaluate the potential effects of organic

nitrogen compounds such as Gln, a candidate nitrogen status signal (Nacry et al., 2013), which

were not quantified in this study. In addition, it is unclear exactly where and how the phloem-

mobile iP-type CKs produced by shoot IPT3 trigger downstream signaling to orchestrate systemic

responses. Furthermore, the cell types, subcellular compartments, and molecular mechanisms

involved sensing shoot nitrate status are entirely unknown. Future work needs to meticulously

investigate these unresolved questions.

Author contributions

TH: conceptualization; TS: data curation; KM and TH: formal analysis; KM and TH: funding

acquisition; KM, TS, MK, YT, and DS: investigation; KM: methodology; KM and TH: project

administration; KM and TH: resources; TN, HS, and TH: supervision; KM and TH: validation;

KM and TH: visualization; TH: writing – original draft preparation; All authors: writing – review

& editing.

Acknowledgements

The NR-null seeds were kindly provided by Dr. Nigel M. Crawford (University of California,

San Diego). The *ipt3* and *ipt3,5,7* seeds were kindly provided by Dr. Tatsuo Kakimoto (Osaka

University). We thank Ms. Yuki Okamoto and Mr. Masahiro Watanabe for their support in the

preparation of the plant materials. This study was supported by JSPS KAKENHI [No. 20K05771

and 23K04978 to TH and No. JP24KJ1709 to KM], by JST SPRING [No. JPMJSP2155 to KM],

and by the Sasakawa Scientific Research Grant from the Japan Science Society [No. 2022-4039

to KM].

Conflicts of interest

The authors declare no competing interests.

Data Availability Statement

The RNA-seq raw data are available in ArrayExpress under the accession number E-MTAB-

14302 (https://www.ebi.ac.uk/biostudies/arrayexpress).

References

Alvarez, J. M., E. Riveras, E. A. Vidal, et al. 2014. "Systems approach identifies TGA1 and TGA4

transcription factors as important regulatory components of the nitrate response of *Arabidopsis*

thaliana roots." *Plant Journal* 80 : 1–13. https://doi.org/10.1111/tpj.12618.

Andrews, M., J. A. Raven, P. J. Lea, and J. I. Sprent. 2006. "A role for shoot protein in shoot–

root dry matter allocation in higher plants." *Annals of Botany* 97: 3–10.

https://doi.org/10.1093/aob/mcj009.

Arnaud, D., S. Lee, Y. Takebayashi, et al. 2017. "Cytokinin-mediated regulation of reactive

oxygen species homeostasis modulates stomatal immunity in Arabidopsis." *Plant Cell* 29: 543–

559. https://doi.org/10.1105/tpc.16.00583.

Bhargava, A., I. Clabaugh, J. P. To, et al. 2013. "Identification of cytokinin-responsive genes

using microarray meta-analysis and RNA-Seq in Arabidopsis." *Plant Physiology* 162: 272–294.

https://doi.org/10.1104/pp.113.217026.

Bishopp, A., S. Lehesranta, A. Vatén, et al. 2011. "Phloem-transported cytokinin regulates polar

auxin transport and maintains vascular pattern in the root meristem." *Current Biology* 21: 927–

932. https://doi.org/10.1016/j.cub.2011.04.049.

595 Bouguyon, E., F. Brun, D. Meynard, et al. 2015. "Multiple mechanisms of nitrate sensing by

596 *Arabidopsis* nitrate transceptor NRT1.1." *Nature Plants* 1: 15015.

597 https://doi.org/10.1038/nplants.2015.15.

599 Choi, J., S. U. Huh, M. Kojima, H. Sakakibara, K. H. Paek, and I. Hwang. 2010. "The cytokinin-

600 activated transcription factor ARR2 promotes plant immunity via TGA3/NPR1-dependent

601 salicylic acid signaling in *Arabidopsis*." *Developmental Cell* 19: 284–295.

602 https://doi.org/10.1016/j.devcel.2010.07.011.

604 Cortleven, A., J. E. Leuendorf, M. Frank, D. Pezzetta, S. Bolt, and T. Schmülling. 2019.

605 "Cytokinin action in response to abiotic and biotic stresses in plants." *Plant, Cell & Environment*

606 42: 998–1018. https://doi.org/10.1111/pce.13494.

608 Durand, M., V. Brehaut, G. Clement, et al. 2023. "The Arabidopsis transcription factor NLP2

609 regulates early nitrate responses and integrates nitrate assimilation with energy and carbon

610 skeleton supply." *Plant Cell* 35: 1429–1454. https://doi.org/10.1093/plcell/koad025.

612 Ge, S. X., E. W. Son, and R. Yao. 2018. "iDEP: an integrated web application for differential

613 expression and pathway analysis of RNA-Seq data." *BMC Bioinformatics* 19: 534.

614 https://doi.org/10.1186/s12859-018-2486-6.

616 Hachiya, T., D. Sugiura, M. Kojima, et al. 2014. "High CO_2 triggers preferential root growth of

617 *Arabidopsis thaliana* via two distinct systems under low pH and low N stresses." *Plant & Cell*

618 *Physiology* 55: 269–280. https://doi.org/10.1093/pcp/pcu001.

620 Hachiya, T., N. Ueda, M. Kitagawa, et al. 2016. "Arabidopsis root-type ferredoxin:NADP(H)

oxidoreductase 2 is involved in detoxification of nitrite in roots." *Plant & Cell Physiology* 57:

2440–2450. https://doi.org/10.1093/pcp/pcw158.

Hachiya, T., and H. Sakakibara. 2017. "Interactions between nitrate and ammonium in their

uptake, allocation, assimilation, and signaling in plants." *Journal of Experimental Botany* 68:

2501-2512. https://doi.org/10.1093/jxb/erw449.

Hachiya, T., Y. Okamoto, M. Watanabe, et al. 2020. "Genome-wide responses to shoot nitrate

satiety are attenuated by external ammonium in *Arabidopsis thaliana*." *Soil Science and Plant*

Nutrition 66: 317–327. https://doi.org/10.1080/00380768.2020.1717905.

Hachiya, T., T. Oya, K. Monden, A. Nagae, and T. Nakagawa. 2021a. "A cellophane-supported

Arabidopsis culture for seamless transfer between different media is useful for studying various

nitrogen   responses."   *Soil   Science   and   Plant   Nutrition*   67:   277–282.

https://doi.org/10.1080/00380768.2021.1908094.

Hachiya, T., J. Inaba, M. Wakazaki, et al. 2021b. "Excessive ammonium assimilation by plastidic

glutamine   synthetase   causes   ammonium   toxicity   in   *Arabidopsis   thaliana*."   *Nature*

Communications 12: 4944. https://doi.org/10.1038/s41467-021-25238-7.

Hirose, N., K. Takei, T. Kuroha, T. Kamada-Nobusada, H. Hayashi, and H. Sakakibara. 2008.

"Regulation of cytokinin biosynthesis, compartmentalization and translocation." *Journal of*

Experimental Botany 59: 75–83. https://doi.org/10.1093/jxb/erm157.

Ho, C. H., S. H. Lin, H. C. Hu, and Y. F. Tsay. 2009. "CHL1 functions as a nitrate sensor in

plants." *Cell* 138: 1184–1194. https://doi.org/10.1016/j.cell.2009.07.004.

Hu, H. C., Y. Y. Wang, and Y. F. Tsay. 2009. "AtCIPK8, a CBL-interacting protein kinase,

regulates the low-affinity phase of the primary nitrate response." *Plant Journal* 57: 264–278.

https://doi.org/10.1111/j.1365-313X.2008.03685.x.

Kanno, Y., A. Hanada, Y. Chiba, et al. 2012. "Identification of an abscisic acid transporter by

functional screening using the receptor complex as a sensor." *Proceedings of the National*

Academy of Sciences of the United States of America 109: 9653–9658.

https://doi.org/10.1073/pnas.120356710.

Kiba, T., K. Takei, M. Kojima, and H. Sakakibara. 2013. "Side-chain modification of cytokinins

controls shoot growth in *Arabidopsis*." *Developmental Cell* 27: 452–461.

https://doi.org/10.1016/j.devcel.2013.10.004.

Kiba, T., J. Inaba, T. Kudo, et al. 2018. "Repression of nitrogen starvation responses by members

of the Arabidopsis GARP-type transcription factor NIGT1/HRS1 subfamily." *Plant Cell* 30: 925–

945. https://doi.org/10.1105/tpc.17.00810.

Ko, D., J. Kang, T. Kiba, et al. 2014. "*Arabidopsis* ABCG14 is essential for the root-to-shoot

translocation of cytokinin." *Proceedings of the National Academy of Sciences of the United States*

of America 111: 7150–7155. https://doi.org/10.1073/pnas.132151911.

Kobayashi, R., Y. Ohkubo, M. Izumi, et al. 2024. "Integration of shoot-derived polypeptide

signals by root TGA transcription factors is essential for survival under fluctuating nitrogen

environments." *Nature Communications* 15: 6903. https://doi.org/10.1038/s41467-024-51091-5.

Kojima, M., T. Kamada-Nobusada, H. Komatsu, et al. 2009. "Highly sensitive and high-

throughput analysis of plant hormones using MS-probe modification and liquid chromatography–

tandem mass spectrometry: an application for hormone profiling in *Oryza sativa*." *Plant & Cell*

Physiology 50: 1201–1214. https://doi.org/10.1093/pcp/pcp057.

Kojima, M., N. Makita, K. Miyata, et al. 2023. "A cell wall–localized cytokinin/purine riboside

nucleosidase is involved in apoplastic cytokinin metabolism in *Oryza sativa*." *Proceedings of the*

National Academy of Sciences of the United States of America 120: e2217708120.

https://doi.org/10.1073/pnas.221770812.

Konishi, M., and S. Yanagisawa. 2013. "Arabidopsis NIN-like transcription factors have a central

role in nitrate signalling." *Nature communications* 4: 1617. https://doi.org/10.1038/ncomms2621.

Krapp, A., R. Berthomé, M. Orsel, et al. 2011. "Arabidopsis roots and shoots show distinct

temporal adaptation patterns toward nitrogen starvation." *Plant Physiology* 157: 1255–1282.

https://doi.org/10.1104/pp.111.179838.

Krouk, G, B. Lacombe, A. Bielach, et al. 2010. "Nitrate-regulated auxin transport by NRT1.1

defines a mechanism for nutrient sensing in plants." *Developmental Cell* 18: 927–937.

https://doi.org/10.1016/j.devcel.2010.05.008.

Kuroha, T., H. Tokunaga, M. Kojima, et al. 2009. "Functional analyses of *LONELY GUY*

cytokinin-activating enzymes reveal the importance of the direct activation pathway in

Arabidopsis." *Plant Cell* 21: 3152–3169. https://doi.org/10.1105/tpc.109.068676.

Langmead, B., C. Trapnell, M. Pop, and S. L. Salzberg. 2009. "Ultrafast and memory-efficient

alignment of short DNA sequences to the human genome." *Genome Biology* 10: R25.

https://doi.org/10.1186/gb-2009-10-3-r25.

Laplaze, L., E. Benkova, I. Casimiro, et al. 2007. "Cytokinins act directly on lateral root founder

cells to inhibit root initiation." *Plant Cell* 19: 3889–3900. https://doi.org/10.1105/tpc.107.055863.

Léran, S., K. H. Edel, M. Pervent, et al. 2015. "Nitrate sensing and uptake in *Arabidopsis* are

enhanced by ABI2, a phosphatase inactivated by the stress hormone abscisic acid." *Science*

Signaling 8: ra43. DOI: 10.1126/scisignal.aaa4829.

Liu, K. H., C. Y. Huang, and Y. F. Tsay. 1999. "CHL1 is a dual-affinity nitrate transporter of

Arabidopsis involved in multiple phases of nitrate uptake." *Plant Cell* 11: 865–874.

https://doi.org/10.1105/tpc.11.5.865.

Liu, K. H., Y. Niu, M. Konishi, et al. 2017. "Discovery of nitrate–CPK–NLP signalling in central

nutrient–growth networks." *Nature* 545: 311–316. https://doi.org/10.1038/nature22077.

Liu, K. H., M. Liu, Z. Lin, et al. 2022. "NIN-like protein 7 transcription factor is a plant nitrate

sensor." *Science* 377: 1419–1425. https://doi.org/10.1126/science.add1104.

Maeda, Y., M. Konishi, T. Kiba, et al. 2018. "A NIGT1-centred transcriptional cascade regulates

nitrate signalling and incorporates phosphorus starvation signals in *Arabidopsis*." *Nature*

Communications 9: 1376. https://doi.org/10.1038/s41467-018-03832-6.

Marchive, C., F. Roudier, L. Castaings, et al. 2013. "Nuclear retention of the transcription factor

NLP7 orchestrates the early response to nitrate in plants." *Nature communications* 4: 1713.

https://doi.org/10.1038/ncomms2650.

Medici, A., and G. Krouk. 2014. "The primary nitrate response: a multifaceted signalling

pathway." *Journal of Experimental Botany* 65: 5567–5576. https://doi.org/10.1093/jxb/eru245.

Miyawaki, K., M. Matsumoto-Kitano, and T. Kakimoto. 2004. "Expression of cytokinin

biosynthetic isopentenyltransferase genes in *Arabidopsis*: tissue specificity and regulation by

auxin, cytokinin, and nitrate." *Plant Journal* 37: 128–138. https://doi.org/10.1046/j.1365-

313X.2003.01945.x.

Miyawaki, K., P. Tarkowski , M. Matsumoto-Kitano, et al. 2006. "Roles of *Arabidopsis*

ATP/ADP isopentenyltransferases and tRNA isopentenyltransferases in cytokinin biosynthesis."

Proceedings of the National Academy of Sciences of the United States of America 103: 16598–

16603. https://doi.org/10.1073/pnas.0603522103.

Monden, K., T. Kamiya, D. Sugiura, T. Suzuki, T. Nakagawa, and T. Hachiya. 2022. "Root-

specific activation of plasma membrane H^+-ATPase 1 enhances plant growth and shoot

accumulation of nutrient elements under nutrient-poor conditions in *Arabidopsis thaliana*."

Biochemical and Biophysical Research Communications 621: 39–45.

https://doi.org/10.1016/j.bbrc.2022.06.097.

Nacry, P., E. Bouguyon, and A. Gojon. 2013. "Nitrogen acquisition by roots: physiological and

developmental mechanisms ensuring plant adaptation to a fluctuating resource." *Plant and Soil*

370: 1–29. https://doi.org/10.1007/s11104-013-1645-9.

Ohkubo, Y., M. Tanaka, R. Tabata, M. Ogawa-Ohnishi, and Y. Matsubayashi. 2017. "Shoot-to-

root mobile polypeptides involved in systemic regulation of nitrogen acquisition." *Nature Plants*

3: 17029. https://doi.org/10.1038/nplants.2017.29.

Okamoto, Y., T. Suzuki, D. Sugiura, T. Kiba, H. Sakakibara, and T. Hachiya. 2019. "Shoot nitrate

underlies a perception of nitrogen satiety to trigger local and systemic signaling cascades in

Arabidopsis thaliana." *Soil Science and Plant Nutrition* 65: 56–64.

https://doi.org/10.1080/00380768.2018.1537643.

Osugi, A., M. Kojima, Y. Takebayashi, N. Ueda, T. Kiba, and H. Sakakibara. 2017. "Systemic

transport of *trans*-zeatin and its precursor have differing roles in *Arabidopsis* shoots." *Nature*

Plants 3: 17112. https://doi.org/10.1038/nplants.2017.112.

Ota, R., Y. Ohkubo, Y. Yamashita, M. Ogawa-Ohnishi, and Y. Matsubayashi. 2020. "Shoot-to-

root mobile CEPD-like 2 integrates shoot nitrogen status to systemically regulate nitrate uptake

in *Arabidopsis*." *Nature Communications* 11: 641. https://doi.org/10.1038/s41467-020-14440-8.

Patterson, K., L. A. Walters, A. M. Cooper, et al. 2016. "Nitrate-regulated glutaredoxins control

Arabidopsis primary root growth." *Plant Physiology* 170: 989–999.

https://doi.org/10.1104/pp.15.01776.

Poitout, A., A. Crabos, I. Petřík, et al. 2018. "Responses to systemic nitrogen signaling in

Arabidopsis roots involve *trans*-zeatin in shoots." *Plant Cell* 30: 1243–1257.

https://doi.org/10.1105/tpc.18.00011.

Remans, T., P. Nacry, M. Pervent, et al. 2006. "The *Arabidopsis* NRT1.1 transporter participates

in the signaling pathway triggering root colonization of nitrate-rich patches." *Proceedings of the*

National Academy of Sciences of the United States of America 103: 19206–19211.

https://doi.org/10.1073/pnas.060527510.

Riveras, E., J. M. Alvarez, E. A. Vidal, C. Oses, A. Vega, and R. A. Gutiérrez. 2015. "The calcium

ion is a second messenger in the nitrate signaling pathway of Arabidopsis." *Plant Physiology* 169:

1397–1404. https://doi.org/10.1104/pp.15.00961.

Rubin, G., T. Tohge, F. Matsuda, K. Saito, and W. R. Scheible. 2009. "Members of the *LBD*

family of transcription factors repress anthocyanin synthesis and affect additional nitrogen

responses in *Arabidopsis*." *Plant Cell* 21: 3567–3584. https://doi.org/10.1105/tpc.109.067041.

Ruffel, S., G. Krouk, D. Ristova, D. Shasha, K. D. Birnbaum, and G. M. Coruzzi. 2011. "Nitrogen

economics of root foraging: transitive closure of the nitrate–cytokinin relay and distinct systemic

signaling for N supply vs. demand." *Proceedings of the National Academy of Sciences of the*

United States of America 108: 18524–18529. https://doi.org/10.1073/pnas.1108684108.

Sakakibara, H., K. Takei, and N. Hirose. 2006. "Interactions between nitrogen and cytokinin in

the regulation of metabolism and development." *Trends in Plant Science* 11: 440–448.

https://doi.org/10.1016/j.tplants.2006.07.004.

Sakakibara, H. 2021. "Cytokinin biosynthesis and transport for systemic nitrogen signaling."

Plant Journal 105: 421–430. https://doi.org/10.1111/tpj.15011.

Safi, A., A. Medici, W. Szponarski, et al. 2021. "GARP transcription factors repress Arabidopsis

nitrogen starvation response via ROS-dependent and -independent pathways." *Journal of*

Experimental Botany 72: 3881–3901. https://doi.org/10.3390/ijms22168685.

Scheible, W. R., M. Lauerer, E. D. Schulze, M. Caboche, and M. Stitt. 1997. "Accumulation of

nitrate in the shoot acts as a signal to regulate shoot-root allocation in tobacco." *Plant Journal* 11:

671–691. https://doi.org/10.1046/j.1365-313X.1997.11040671.x.

Signora, L., I. De Smet, C. H. Foyer, and H. Zhang. 2001. "ABA plays a central role in mediating

the regulatory effects of nitrate on root branching in *Arabidopsis*." *Plant Journal* 28: 655–662.

https://doi.org/10.1046/j.1365-313x.2001.01185.x.

Su, H., T. Wang, C. Ju, et al. 2021. "Abscisic acid signaling negatively regulates nitrate uptake

via phosphorylation of NRT1. 1 by SnRK2s in *Arabidopsis*." *Journal of Integrative Plant Biology*

63: 597–610. https://doi.org/10.1111/jipb.13057.

Tabata, R., K. Sumida, T. Yoshii, K. Ohyama, H. Shinohara, and Y. Matsubayashi. 2014.

"Perception of root-derived peptides by shoot LRR-RKs mediates systemic N-demand signaling."

Science 346: 343–346. DOI: 10.1126/science.1257800.

Takei, K., N. Ueda, K. Aoki, et al. 2004a. "*AtIPT3* is a key determinant of nitrate-dependent

cytokinin biosynthesis in *Arabidopsis*." *Plant & Cell Physiology* 45: 1053–1062.

https://doi.org/10.1093/pcp/pch119.

Takei, K., T. Yamaya, and H. Sakakibara. 2004b. "Arabidopsis *CYP735A1* and *CYP735A2*

encode cytokinin hydroxylases that catalyze the biosynthesis of *trans*-zeatin." *Journal of*

Biological Chemistry 279: 41866–41872. https://doi.org/10.1074/jbc.M406337200.

Tokunaga, H., M. Kojima, T. Kuroha, et al. 2011. "Arabidopsis lonely guy (LOG) multiple

mutants reveal a central role of the LOG-dependent pathway in cytokinin activation." *Plant*

Journal 69: 355–365. https://doi.org/10.1111/j.1365-313X.2011.04795.x.

Wang, R., R. Tischner, R. A. Gutiérrez, et al. 2004. "Genomic analysis of the nitrate response

using a nitrate reductase-null mutant of Arabidopsis." *Plant Physiology* 136: 2512–2522.

https://doi.org/10.1104/pp.104.044610.

Wang, R, X. Xing, and N. M. Crawford. 2007. "Nitrite acts as a transcriptome signal at

micromolar concentrations in Arabidopsis roots." *Plant Physiology* 145: 1735–1745.

https://doi.org/10.1104/pp.107.108944.

Wang, X., C. Feng, L. Tian, et al. 2021. "A transceptor–channel complex couples nitrate sensing

to calcium signaling in *Arabidopsis*." *Molecular Plant* 14: 774–786.

https://doi.org/10.1016/j.molp.2021.02.005.

Wang, H., C. Han, J. G. Wang, et al. 2022. "Regulatory functions of cellular energy sensor SnRK1

for nitrate signalling through NLP7 repression." *Nature Plants* 8: 1094-1107.

https://doi.org/10.1038/s41477-022-01236-5.

Werner, T., V. Motyka, V. Laucou, R. Smets, H. Van Onckelen, and T. Schmülling. 2003.

"Cytokinin-deficient transgenic Arabidopsis plants show multiple developmental alterations

indicating opposite functions of cytokinins in the regulation of shoot and root meristem activity."

Plant Cell 15: 2532–2550. https://doi.org/10.1105/tpc.014928.

Werner, T., K. Holst, Y. Pörs, et al. 2008. "Cytokinin deficiency causes distinct changes of sink

and source parameters in tobacco shoots and roots." *Journal of Experimental Botany* 59: 2659–

2672. https://doi.org/10.1093/jxb/ern134.

Wu, B., J. Meng, H. Liu, et al. 2023. "Suppressing a phosphohydrolase of cytokinin nucleotide

enhances grain yield in rice." *Nature Genetics* 55: 1381–1389. https://doi.org/10.1038/s41588-

023-01454-3.

Xu, N., R. Wang, L. Zhao, et al. 2016. "The Arabidopsis NRG2 protein mediates nitrate signaling

and interacts with and regulates key nitrate regulators." *Plant Cell* 28: 485–504.

https://doi.org/10.1105/tpc.15.00567.

Zhou, Y., B. Zhou, L. Pache, et al. 2019. "Metascape provides a biologist-oriented resource for

the analysis of systems-level datasets." *Nature Communications* 10: 1523.

https://doi.org/10.1038/s41467-019-09234-6.

Zhang, H., A. Jennings, P. W. Barlow, and B. G. Forde. 1999. "Dual pathways for regulation of

root branching by nitrate." *Proceedings of the National Academy of Sciences of the United States*

of America 96: 6529–6534. https://doi.org/10.1073/pnas.96.11.6529.

Zhang, K., O. Novak, Z. Wei, et al. 2014. "*Arabidopsis* ABCG14 protein controls the acropetal

translocation of root-synthesized cytokinins." *Nature Communications* 5: 3274.

https://doi.org/10.1038/ncomms4274.

Supporting Information

Additional supporting information can be found online in the Supporting Information section.

Figure Legends

Figure 1. Micrografting and transfer experiments for manipulation of nitrate levels and *IPT3* expression in a shoot-specific manner. (A) Summary of the micrografting and transfer experiments. (B) A schematic representation of four types of grafted plants. The dotted line represents the grafted seam in the hypocotyl. (C, D) Nitrate concentrations in shoots (C) and roots (D) 0, 3, or 5 days after nitrogen removal. (E, F) Total protein concentrations in shoots (E) and roots (F) 5 days after nitrogen removal. (C–F) Two grafted plants were pooled as one biological replicate. Data are presented as mean ± SD (n = 3). (G, H) Total nitrogen concentrations in shoots (G) and roots (H) 5 days after nitrogen removal. Sampling was performed three times independently. In this sampling, 7, 6, and 5 grafted plants were pooled as one biological replicate. Data are presented as mean ± SD (n = 3). (C–H) Different lowercase letters indicate significant differences, as determined via Tukey–Kramer test ($P < 0.05$). (I) Transcript levels of *IPT3* in shoots 5 days after nitrogen removal, as determined using RT-qPCR. Sampling was done five times independently. In this sampling, 9, 6, 7, 7, and 8 grafted plants were pooled as one biological replicate. Data are presented as mean ± SD (n = 5). (J) Nitrate concentrations in shoots 5 days after nitrogen removal. (K) Transcript levels of *GFP* in shoots 5 days after nitrogen removal, as determined via RT-qPCR. (J, K) One biological replicate consisted of one grafted plant. Data are presented as mean ± SD (n = 3). (I–K) Statistical significance was assessed using unpaired two-tailed Welch's *t*-test ($*P < 0.05$; $**P < 0.01$; $***P < 0.001$). (L) Fluorescence images of true leaves 5 days after nitrogen removal. Green and red colors correspond to GFP fluorescence and autofluorescence, respectively. Scale bars represent 500 μm. In graph legends, the grafted plants

are indicated as "scion line name/rootstock line name." ns, not significant; FW, fresh weight; DW,

dry weight.

Figure 2. Effects of shoot nitrate status and shoot *IPT3* expression on the shoot and root

concentrations of CK species. (A–M) The shoot and root concentrations of iP (A), iPR (B), iPRPs

(C), iP7G (D), iP9G (E), *t*Z (F), *t*ZR (G), *t*ZPRs (H), *t*Z7G (I), *t*Z9G (J), *t*ZOG (K), *t*ZROG (L),

and *t*ZRPsOG (M) in the grafted plants 5 days after nitrogen removal (condition 5). Sampling

was performed three times independently. In the sampling, 7, 6, and 5 grafted plants were pooled

as one biological replicate. Data are presented as mean ± SD [n = 3, excluding iP concentrations

in Col/*ipt3,5,7* roots and *ipt3*/*ipt3,5,7* roots (n = 2)]. Different lowercase letters indicate

significant differences, as determined via Tukey–Kramer test ($P < 0.05$). Regarding the shoot

concentrations of iP and *t*ZRPsOG, unpaired two-tailed Welch's *t*-test was run at $\alpha = 0.05$. iP, N^6-

(Δ^2-isopentenyl)adenine; iPR, iP riboside; iPRPs, iP ribotides; iP7G, iP-7-*N*-glucoside; iP9G, iP-

9-*N*-glucoside; *t*Z, *trans*-zeatin; *t*ZR, *t*Z riboside; *t*ZRPs, *t*Z ribotides; *t*Z7G, *t*Z-7-*N*-glucoside;

*t*Z9G, *t*Z-9-*N*-glucoside; *t*ZOG, *t*Z-*O*-glucoside; *t*ZROG, *t*ZR-*O*-glucoside; *t*ZRPsOG, *t*ZRPs-*O*-

glucoside; ns, not significant; nd, not detected; FW, fresh weight.

Figure 3. Effects of shoot nitrate status and shoot *IPT3* expression on plant growth. (A)

Representative shoot appearance of grafted plants. Scale bars represent 5 mm. (B–F) Shoot fresh

weight (B), leaf blade area (C), rosette diameter (D), leaf number (E), root fresh weights (F), and

fresh weight ratios of shoot to root (G) of grafted plants 5 days after nitrogen removal (condition

5). All data pooled from five independent grafting experiments are presented as mean ± SD (n =

33). Different lowercase letters indicate significant differences, as determined via Tukey–Kramer

test ($P < 0.05$). ns, not significant; FW, fresh weight.

Figure 4. Effects of shoot nitrate status and shoot *IPT3* expression on the shoot and root

transcriptomes. RNA-seq was performed using the shoots and roots of grafted plants 5 days after

nitrogen removal (condition 5). Sampling was performed three times independently. In the

sampling, 9, 6, and 7 grafted plants were pooled as one biological replicate. (A) Venn diagram

presenting the number of DEGs at a minimum fold change of 1.5 in NR-null/*ipt3,5,7* plants

relative to Col/*ipt3,5,7* plants and in NR-null *ipt3*/*ipt3,5,7* plants relative to *ipt3*/*ipt3,5,7* plants.

(B–E) Plots of normalized transcript levels of 631 DEGs (B) and 663 DEGs (C) in shoots and

189 DEGs (D) and 126 DEGs (E) in roots. 1st, 2nd, and 3rd denote the order of three independent

experiments. Different lowercase letters indicate significant differences, as determined via

Tukey–Kramer test ($P < 0.05$). The numbers on the graph represent the difference in mean values.

Figure 5. Effects of shoot nitrate status and shoot *IPT3* expression on nitrogen-related

transcriptional responses in shoots and roots. (A–D) Plots of mean normalized transcript levels of

nitrate (*per se*)-inducible genes in 2 h (A), genes induced via CPK10/30/32 (B), genes commonly

induced via all NLPs (C), and nitrogen starvation-inducible genes (D) in the shoots and roots of

grafted plants 5 days after nitrogen removal (condition 5). The gene lists were obtained from

Wang et al. (2004) for (A), Liu et al. (2017) for (B), Liu et al. (2022) for (C), and Kiba et al.

(2018) and Krapp et al. (2011) for (D). Different lowercase letters indicate significant differences,

as determined via Tukey–Kramer test ($P < 0.05$). The numbers on the graph represent the

difference in mean values. (E–G) Transcriptional changes in genes involved in nitrogen transport

and assimilation in roots (E), genes encoding transcription factors for nitrogen starvation

responses in shoots (F) and in roots (G), and genes encoding shoot-to-root mobile peptides in

shoots (H) in grafted plants 5 days after nitrogen removal. The normalized transcript levels were

visualized as heatmaps using MORPHEUS (https://software.broadinstitute.org/morpheus). The

relative color scheme uses the minimum (blue) and maximum (yellow) values in each row. 1st,

2nd, and 3rd denote the order of three independent experiments. ns, not significant; TF,

transcription factor.

Figure 6. Summary of systemic responses to shoot nitrate accumulation and their dependence by

shoot IPT3.

Figure 1

(A) Procedure for transfer experiment

	Condition 1	Condition 2	Condition 3	Condition 4	Condition 5
Duration	4 days	5 days	24 hours	24 hours	0, 3, 5 days
N source	2.5 mM NH_4^+	2.5 mM NH_4^+	2.5 mM NH_4^+	10 mM NO_3^- & 2.5 mM NH_4^+	No nitrogen
pH	6.7	6.7	6.7	6.7	5.7
PPFD	25 µmol m^{-2} s^{-1}	50 µmol m^{-2} s^{-1}	25 µmol m^{-2} s^{-1}	25 µmol m^{-2} s^{-1}	25 µmol m^{-2} s^{-1}
Temperature	22°C	27°C	22°C	22°C	22°C

Micrografting   NO$_3^-$ supply   Nitrogen removal

Figure 1. Micrografting and transfer experiments for manipulation of nitrate levels and *IPT3* expression in a shoot-specific manner. (A) Summary of the micrografting and transfer experiments. (B) A schematic representation of four types of grafted plants. The dotted line represents the grafted seam in the hypocotyl. (C, D) Nitrate concentrations in shoots (C) and roots (D) 0, 3, or 5 days after nitrogen removal. (E, F) Total protein concentrations in shoots (E) and roots (F) 5 days after nitrogen removal. (C–F) Two grafted plants were pooled as one biological replicate. Data are presented as mean ± SD (n = 3). (G, H) Total nitrogen concentrations in shoots (G) and roots (H) 5 days after nitrogen removal. Sampling was performed three times independently. In this sampling, 7, 6, and 5 grafted plants were pooled as one biological replicate. Data are presented as mean ± SD (n = 3). (C–H) Different lowercase letters indicate significant differences, as determined via Tukey–Kramer test (*P* < 0.05). (I) Transcript levels of *IPT3* in shoots 5 days after nitrogen removal, as determined using RT-qPCR. Sampling was done five times independently. In this sampling, 9, 6, 7, 7, and 8 grafted plants were pooled as one biological replicate. Data are presented as mean ± SD (n = 5). (J) Nitrate concentrations in shoots 5 days after nitrogen removal. (K) Transcript levels of *GFP* in shoots 5 days after nitrogen removal, as determined via RT-qPCR. (J, K) One biological replicate consisted of one grafted plant. Data are presented as mean ± SD (n = 3). (I–K) Statistical significance was assessed using unpaired two-tailed Welch's *t*-test (*P < 0.05; **P < 0.01; ***P < 0.001). (L) Fluorescence images of true leaves 5 days after nitrogen removal. Green and red colors correspond to GFP fluorescence and autofluorescence, respectively. Scale bars represent 500 µm. In graph legends, the grafted plants are indicated as "scion line name/rootstock line name." ns, not significant; FW, fresh weight; DW, dry weight.

Figure 2

Figure 2. Effects of the shoot nitrate status and shoot *IPT3* expression on the shoot and root concentrations of CK species. (A–M) The shoot and root concentrations of iP (A), iPR (B), iPRPs (C), iP7G (D), iP9G (E), *t*Z (F), *t*ZR (G), *t*ZPRs (H), *t*Z7G (I), *t*Z9G (J), *t*ZOG (K), *t*ZROG (L), and *t*ZRPsOG (M) in the grafted plants 5 days after nitrogen removal (condition 5). Sampling was performed three times independently. In the sampling, 7, 6, and 5 grafted plants were pooled as one biological replicate. Data are presented as mean ± SD [n = 3, excluding iP concentrations in Col/*ipt3,5,7* roots and *ipt3*/*ipt3,5,7* roots (n = 2)]. Different lowercase letters indicate significant differences, as determined via Tukey–Kramer test (*P* < 0.05). Regarding the shoot concentrations of iP and *t*ZRPsOG, unpaired two-tailed Welch's *t*-test was run at *α* = 0.05. iP, N^6-(Δ^2-isopentenyl)adenine; iPR, iP riboside; iPRPs, iP ribotides; iP7G, iP-7-*N*-glucoside; iP9G, iP-9-*N*-glucoside; *t*Z, *trans*-zeatin; *t*ZR, *t*Z riboside; *t*ZRPs, *t*Z ribotides; *t*Z7G, *t*Z-7-*N*-glucoside; *t*Z9G, *t*Z-9-*N*-glucoside; *t*ZOG, *t*Z-*O*-glucoside; *t*ZROG, *t*ZR-*O*-glucoside; *t*ZRPsOG, *t*ZRPs-*O*-glucoside; ns, not significant; nd, not detected; FW, fresh weight.

Figure 3

(A)

Col-0/*ipt3,5,7*   NR-null/*ipt3,5,7*   *ipt3*/*ipt3,5,7*   NR-null *ipt3*/*ipt3,5,7*

(B) Shoot FW

(C) Leaf blade area

(D) Rosette diameter

(E) Leaf number

(F) Root FW

(G) Shoot/Root ratio

Legend:
- Col/*ipt3,5,7*
- NR-null/*ipt3,5,7*
- *ipt3*/*ipt3,5,7*
- NR-null *ipt3*/*ipt3,5,7*

Figure 3. Effects of the shoot nitrate status and shoot *IPT3* expression on plant growth. (A) Representative shoot appearance of grafted plants. Scale bars represent 5 mm. (B–F) Shoot fresh weights (B), leaf blade area (C), rosette diameter (D), leaf number (E), root fresh weights (F), and fresh weight ratios of shoot to root (G) of grafted plants 5 days after nitrogen removal (condition 5). All data pooled from five independent grafting experiments are presented as mean ± SD (n = 33). Different lowercase letters indicate significant differences, as determined via Tukey–Kramer test ($P < 0.05$). ns, not significant; FW, fresh weight.

Figure 4

(A) Number of DEGs

(B) 631 DEGs in shoots

(C) 663 DEGs in shoots

(D) 189 DEGs in roots

(E) 126 DEGs in roots

☐ Col/*ipt3,5,7*
☐ NR-null/*ipt3,5,7*
■ *ipt3*/*ipt3,5,7*
■ NR-null *ipt3*/*ipt3,5,7*

Figure 4. Effects of the shoot nitrate status and shoot *IPT3* expression on the shoot and root transcriptomes. RNA-seq was performed using the shoots and roots of grafted plants 5 days after nitrogen removal (condition 5). Sampling was performed three times independently. In the sampling, 9, 6, and 7 grafted plants were pooled as one biological replicate. (A) Venn diagram presenting the number of DEGs at a minimum fold change of 1.5 in NR-null/*ipt3,5,7* plants relative to Col/*ipt3,5,7* plants and in NR-null *ipt3*/*ipt3,5,7* plants relative to *ipt3*/*ipt3,5,7* plants. (B–E) Plots of normalized transcript levels of 631 DEGs (B) and 663 DEGs (C) in shoots and 189 DEGs (D) and 126 DEGs (E) in roots. 1st, 2nd, and 3rd denote the order of three independent experiments. Different lowercase letters indicate significant differences, as determined via Tukey–Kramer test (*P* < 0.05). The numbers on the graph represent the difference in mean values.

Figure 5

Figure 5. Effects of the shoot nitrate status and shoot *IPT3* expression on nitrogen-related transcriptional responses in shoots and roots. (A–D) Plots of mean normalized transcript levels of nitrate (*per se*)-inducible genes in 2 h (A), genes induced via CPK10/30/32 (B), genes commonly induced via all NLPs (C), and nitrogen starvation-inducible genes (D) in the shoots and roots of grafted plants 5 days after nitrogen removal (condition 5). The gene lists were obtained from Wang et al. (2004) for (A), Liu et al. (2017) for (B), Liu et al. (2022) for (C), and Kiba et al. (2018) and Krapp et al. (2011) for (D). Different lowercase letters indicate significant differences, as determined via Tukey–Kramer test (*P* < 0.05). The numbers on the graph represent the difference in mean values. (E–G) Transcriptional changes in genes involved in nitrogen transport and assimilation in roots (E), genes encoding transcription factors for nitrogen starvation responses in shoots (F) and in roots (G), and genes encoding shoot-to-root mobile peptides in shoots (H) in grafted plants 5 days after nitrogen removal. The normalized transcript levels were visualized as heatmaps using MORPHEUS (https:// software.broadinstitute.org/morpheus). The relative color scheme uses the minimum (blue) and maximum (yellow) values in each row. 1st, 2nd, and 3rd denote the order of three independent experiments. ns, not significant; TF, transcription factor.

Figure 6

Shoot IPT3-regulated responses to shoot NO₃⁻ accumulation

Figure 6. Summary of systemic responses to shoot nitrate accumulation and their modulation by shoot IPT3.