Title: Ectopic gene conversion causing quantitative trait variation

Authors: Marina Pfalz[1], Seïf-Eddine Naàdja[1], Jacqui Anne Shykoff[1], Juergen Kroymann[1*]

Affiliations: [1]Ecologie Société Evolution, CNRS/Université Paris-Saclay/AgroParisTech, Gif-sur-Yvette,
France

*juergen.kroymann@universite-paris-saclay.fr

Abstract: Mutation-selection balance explains the occurrence of genetic variation in populations:
mutations arise continuously, and natural selection removes or fixes them at varying rates. However,
there is more genetic variation segregating in populations than can be explained by mutation-selection
balance alone. Here we elucidate a general mechanism for generating this segregating variation using
the example of a gene family involved in plant defense.

Main Text:

Why is there so much genetic variation segregating in populations (*1*) when we expect natural selection
to purge deleterious mutants rapidly? Clearly, other processes than mutation must be generating this
variation. Here we propose a scenario whereby ectopic gene conversion (EGC) generates fitness
variation in populations, illustrated by dissection of a quantitative trait locus for glucosinolates in
Arabidopsis.

Glucosinolates are amino-acid derived defense metabolites found in Brassicales. Due to their extensive
intra- and interspecific genetic variation, they have become model compounds for understanding the
genomic architecture of adaptive quantitative traits (*2-4*). Indole glucosinolates (IGs) are derived from
tryptophan, are inducible by microbial pathogens or phytophagous insects, and play an important role in
plant defense (*5-9*). We mapped leaf QTL for these IGs in Arabidopsis and cloned the *Indole*
Glucosinolate Modifier 1 (*IGM1*) QTL on chromosome 5 (*9,10*), which controls variation for 4-hydroxy-
indol-3ylmethyl glucosinolate (4OHI3M) and 4-methoxy-indol-3ylmethyl glucosinolate (4MOI3M). A
second QTL on chromosome 1, termed *Indole Glucosinolate Modifier 2* (*IGM2*), which we dissect here,
controls variation for 4OHI3M but not 4MOI3M (*10*).

A tightly linked gene family of indole glucosinolate *O*-methyltransferase genes (*IGMTs*) underlies *IGM2*,
via a complex determination of metabolite levels. While dissecting the QTL, we discovered patterns of
nucleotide polymorphism consistent with ectopic gene conversion (EGC), a process of non-reciprocal
transfer of genetic material between paralogous loci during DNA repair. This was interesting because
sequence transfer between specialized gene copies places sequence information into an inappropriate
expression context, which should produce maladapted variants. Here we demonstrate that the QTL
genes are shaped by EGC, but that selection acts against EGC near codons specifying fixed derived amino
acids, *i.e.*, mutations that differentiate *Arabidopsis thaliana* gene copies from those of *Arabidopsis*
lyrata. This allows us to conclude that gene copies have specialized upon subfunctionalization and that
EGC indeed generates maladapted variants under these conditions.

Results

Mapping and dissection of *IGM2*

We fine-mapped *IGM2* using F_2 progeny from a cross between two near-isogenic lines (NILs) of the
Arabidopsis Da(1)-12 × Ei-2 mapping population, DE096 and DE155 (*10*). In leaves, DE096 had more
4OHI3M than DE155, but the NILs did not differ in their 4MOI3M content. In roots, DE096 had more
4OHI3M and less 4MOI3M than DE155 (**fig. S1, A**). DE096 and DE155 differed only in a ~ 20 Mbp region
containing *IGM2*, while their marker genotypes were identical for the rest of the genome (**fig. S1, B**). We
phenotyped for the fraction of 4OHI3M in the combined amount of 4OHI3M and 4MOI3M. *IGM2*
mapped to a ca. 1.6 Mbp interval, with no evidence of additional QTL nearby (**fig. S1, C and D**). Statistical
support for a QTL centered near a family of four tandemly arranged methyltransferase genes. These
genes were previously identified as coding for indole glucosinolate *O*-methyltransferases (IGMTs) that
convert 4OHI3M to 4MOI3M (*11*), making them plausible candidates for the QTL (**fig. S1, E and F**).

We conducted quantitative real-time PCR to assess whether differences in *IGMT* expression could
explain the QTL. We used primers specific to the target genes, ensuring substantial mismatches with
other gene family members while avoiding polymorphisms at primer binding sites between accessions
(**table S1**). Additionally, we employed a primer pair that could simultaneously amplify all genes. Leaf
IGMT1 and *3* transcript levels were higher in DE155 than in DE096, as was the overall transcript level,
while *IGMT2* and *4* transcript levels did not differ between the two NILs (**Fig. 1**). In roots, *IGMT1*
transcript levels were higher in DE155, but *IGMT3*, *4*, and particularly total *IGMT* transcripts were lower.
Thus, genotypic differences in *IGMT* transcript abundance alone could not explain the observed
quantitative metabolite pattern.

Therefore, we employed CRISPR/Cas9 to generate *igmt* mutants in both NILs. Our approach involved a
construct that produced three single guide RNAs (sgRNAs), each targeting distinct sites within the four
IGMT sequences. However, due to sequence variation, the sgRNAs did not match precisely in every
instance. We Sanger-sequenced all Cas9 target sites within the four *IGMT* genes in our mutant lines.
Typically, we detected small insertions or deletions that caused frameshifts. However, in certain cases,
we observed larger deletions resulting in the fusion of mismatched sequence from neighboring genes
(**fig. S2**).

We obtained quadruple knockouts, *igmt1-4*$_{DE155}$ and *igmt1-4*$_{DE096}$, for both NILs. While the leaves of
these mutants lacked 4MOI3M entirely, the roots still exhibited a background level of 4MOI3M,
suggesting the presence of another enzyme capable of converting 4OHI3M to 4MOI3M in roots. Our
prime candidate was IGMT5, encoded on chromosome 5, which normally converts 1-hydroxy-indol-
3ylmethyl to 1-methoxy-indole-3ylmethyl glucosinolate and has high root but low leaf activity (*5*). We
therefore generated an *igmt1-4* quadruple knockout in the *igmt5* mutant background. As expected,
these plants showed no detectable 4MOI3M in either leaves or roots (**Table 1**).

In addition to the quadruple mutant, referred to as *igmt1-4*$_{DE155}$, we obtained the genotypes *igmt3*$_{DE155}$,
igmt1/3$_{DE155}$, and *igmt1-3*$_{DE155}$ for DE155, and *igmt1/4*$_{DE096}$, *igmt1/3/4*$_{DE096}$, and *igmt1/2/4*$_{DE096}$ for DE096
(**fig. S2)**.

In leaves, we observed the most substantial differences in the levels of 4OHI3M, 4MOI3M and/or the
fraction of 4OHI3M when comparing $igmt1\text{-}3_{DE155}$ with $igmt1\text{-}4_{DE155}$, and DE096 wildtype with
$igmt1/4_{DE096}$ (**Fig. 2**). Thus, *IGMT4* had the largest impact on 4MOI3M generation in the leaves of both
NILs. Additionally, we found noticeable differences between $igmt1/2/4_{DE096}$ and $igmt1/4_{DE096}$, and
between $igmt1/3/4_{DE096}$ and $igmt1\text{-}4_{DE096}$ (**Fig. 2**), which highlighted the role of $IGMT2_{DE096}$. In stark
contrast, we only found minor differences for $IGMT2_{DE155}$ when comparing $igmt1/3_{DE155}$ and $igmt1\text{-}3_{DE155}$,
indicating that genotypic variation in *IGMT2* contributed to the leaf QTL.

However, given that the generation of each 4MOI3M molecule consumes one 4OHI3M molecule, the
contrast between an active IGMT2 in DE096 and a largely inactive IGMT2 in DE155 alone could not
explain the leaf QTL pattern. We therefore inspected the effects of IGMT4 in both NILs more closely,
based on the combined analysis of data for $igmt1\text{-}3_{DE155}$, $igmt1\text{-}4_{DE155}$, $igmt1/4_{DE096}$, and DE096 from four
independent experiments. Indeed, this analysis unveiled an additional quantitative difference between
the NILs (**Fig. 3**). While the increase in 4OHI3M was consistent in the comparisons from $igmt1\text{-}3_{DE155}$ to
$igmt1\text{-}4_{DE155}$ and from DE096 to $igmt1/4_{DE096}$, the reduction in 4MOI3M was significantly more
pronounced when *IGMT4* was knocked out in the DE155 background. This indicated that IGMT4 had a
higher level of activity in DE155 compared to DE096, thereby increasing the flux through the pathway.

We also conducted tests to ascertain if there were differences in the levels of *CYP81F* transcripts
between DE155 and DE096, or if mutations in *IGMT* genes influenced the transcript abundance of
wildtype *IGMT*s. However, we observed no substantial alterations (**fig. S3**).

Based on these results, we could explain the leaf QTL as the cumulative effects of two distinct
quantitative trait genes (QTGs), namely *IGMT2* and *IGMT4*. Genotypic variation in *IGMT4* led to an
excess of 4MOI3M in DE155 without altering 4OHI3M levels. This should result in a QTL for 4MOI3M,
with DE155 having the high allele for *IGMT4*. However, genotypic variation in *IGMT2* resulted in an
elevation of 4MOI3M in DE096, which counterbalanced the excess of 4MOI3M in DE155, and
concurrently caused a reduction in 4OHI3M. Consequently, the collective impact of genotypic variation
in both QTGs manifested as a disparity in 4OHI3M levels, with no noticeable difference in the levels of
4MOI3M.

In the roots of DE155, *IGMT1* was the primary contributor to 4MOI3M biosynthesis, as evidenced by the
contrast between $igmt3_{DE155}$ and $igmt1/3_{DE155}$ (**Fig. 2**). IGMT4 and, notably, IGMT2 also played roles, but
IGMT3 had no discernible impact. In the roots of DE096, we noted the largest contrast between
$igmt1/4_{DE096}$ and DE096 wildtype. However, the $igmt1/4_{DE096}$ double mutant alone did not allow us to
discern whether the observed effects were attributable to $IGMT1_{DE096}$, $IGMT4_{DE096}$, or their combination.
Furthermore, $IGMT2_{DE096}$ and $IGMT3_{DE096}$ were implicated in the conversion of 4OHI3M to 4MOI3M,
primarily observed through the impact of the corresponding mutants on the fraction of 4OHI3M.

The influence of $IGMT1_{DE155}$ was substantially more pronounced than the combined effect of $IGMT1_{DE096}$
and $IGMT4_{DE096}$. Hence, *IGMT1* was a root QTG. *IGMT3* was a second, but cryptic, root QTG, with DE096
possessing the high allele that mitigated the impact of $IGMT1_{DE155}$. In contrast to leaves, genotypic
variation in *IGMT2* had no noticeable quantitative effect on roots.

This root QTL pattern corresponded well with the abundance of *IGMT* transcripts in both NILs (**Fig. 1**),
suggesting that quantitative variation in the roots was mainly governed by differences in the expression

of individual genes. The transcript level of *IGMT1* was higher and that of *IGMT3* lower in DE155 than in
DE096, while the transcript level of *IGMT2* was similar between NILs. Interestingly, *IGMT4*$_{DE096}$ exhibited
a higher transcript abundance than *IGMT4*$_{DE155}$, indicating that *IGMT4* could be another cryptic root
QTG, further counteracting the effect of genotypic variation in *IGMT1*.

Ectopic gene conversion in the *IGMT* cluster

In the comprehensive assessment of QTL effects and transcript level variation among NILs, *IGMT2*$_{DE155}$
stood out. The DE155 allele was obviously functional in roots, contributing to 4MOI3M at a level
comparable to its DE096 counterpart (**Fig. 2**). However, *IGMT2*$_{DE155}$ activity in leaves was relatively
minor, despite having a transcript level that was similar, if not slightly higher, than that of *IGMT2*$_{DE096}$
(**Fig. 1**).

To investigate the cause of this peculiar phenomenon, we built a gene tree with all four genes from both
NILs. We expected alleles of the same gene copies to cluster together, which was indeed the case for
IGMT1 and *IGMT4*. Much to our surprise, however, the relationship between *IGMT2* and *IGMT3* genes
and alleles remained unresolved (**fig. S4**), suggesting that *IGMT2*$_{DE155}$ and *IGMT3*$_{DE155}$ did not evolve
independently. Such non-independence could arise from non-homologous recombination between gene
copies, generating chimeras from two or more copies. Alternatively, it could stem from the non-
reciprocal transfer of genetic material between paralogous loci, known as ectopic gene conversion
(EGC).

To examine this further, we procured and manually aligned sequences of the *IGMT* region from the 1001
Arabidopsis Genomes project (1001genomes.org), concentrating on *de novo*, reference-quality
assembles (*12*), and incorporated additional short-read assemblies with near-complete *IGMT* sequences
(*13*) (**table S2**). Our alignment contained data from 28 accessions. All assemblies of the *IGMT* region,
particularly those from long reads, showed exactly four *IGMT* copies. Because non-homologous
recombination usually generates copy number variation, we rule this out for explaining the apparent
non-independent evolution of *IGMT* genes.

We compared the coding sequences of all four *IGMT* genes and found numerous polymorphisms
(**10.6084/m9.figshare.28093955**). To ensure these were not sequencing errors, we amplified and
sequenced *IGMT4* coding sequence from 16 Arabidopsis accessions, confirming our findings. Strikingly,
many of these polymorphisms were not copy-specific, but shared by two or even more gene copies.

We tested whether point mutations could explain the high number of cases where the same
polymorphism segregated at the same site in two or more copies, which we refer to as 'shared
polymorphisms'. We focused on instances with two independent mutations affecting the same site,
either within a single gene copy or across two distinct copies. The relative probabilities of both events
depended on the number of gene copies (n), and were n^{-1} for two mutations at the same site in the
same gene copy and $1 - n^{-1}$ for two mutations at the same site in two distinct copies.

Assuming a large population and no initial polymorphisms (but still allowing fixed differences between
gene copies), two outcomes can occur when the same gene copy mutates twice at the same site. Either

both mutations lead to the same polymorphism, appearing as a single, copy-specific polymorphism, or
they result in three nucleotides segregating at the same site. When mutations hit the same site in two
different copies, this can lead to a shared polymorphism or to two copy-specific polymorphisms. The
probabilities of these outcomes depend on whether the copies initially had identical or different
nucleotides at the side in question, and in the latter case, it also depends on whether the nucleotides
differed by a transition or a transversion (see **Supplementary text** for details). From our data, we
estimated a transition/transversion ratio of 2.5935, which was in line with other studies (*14,15*).
However, we were uncertain about the initial count of identical *versus* different positions in gene copy
comparisons. Therefore, we examined the entire frequency spectrum of identical *versus* different
positions, ranging from zero to 100%, taking into account their mutual exclusivity.

For n = 4 copies and a transition rate of 2.5935, as in the case of the Arabidopsis *IGMT* genes, the
proportion of shared among visible types of polymorphisms can never exceed 50% (**fig. S5**). At higher
gene copy numbers, this proportion plateaus at 56%. In stark contrast, our entire dataset had only three
cases with three different segregating nucleotides in the same gene copy and ten cases with two
different segregating polymorphisms across multiple gene copies. However, shared polymorphisms
across copies occurred at 61 sites, making up over 80% of the total visible polymorphisms that affected
either the same copy twice or two different copies once each.

Thus, observed polymorphism patterns revealed far too many shared polymorphisms across gene copies
than would be expected from independent point mutations, making EGC the most likely explanation.
Indeed, visual inspection identified numerous linked specific variants at polymorphic sites, sometimes
spanning hundreds of positions, indicative of gene conversion tracts.

Reexamining the 13 cases of three segregating nucleotides in the same copy or with two different
segregating polymorphisms across multiple gene copies revealed that most of these events were better
explained by EGC than by independent point mutations. Hence, shared polymorphisms resulting from
two independent point mutations were even more rare than initially suspected. Thus, shared
polymorphisms overall indicate EGC.

EGC and subfunctionalization

What are the consequences of EGC for the *IGMT* gene cluster? All four gene products performed the
same reaction – converting 4OHI3M to 4MOI3M – but showed organ-specific differences in transcript
abundance (**Fig. 1**). CRISPR/Cas9-induced mutations in one gene were not effectively offset by the
remaining gene copies, which implied that different *IGMT*s may be expressed in distinct cells. Indeed,
IGMT2 and *IGMT3* are expressed in different root tissues (*16*). Together, these findings suggest
subfunctionalization, a process that preserves gene duplicates in a genome by partitioning the functions
of the ancestral gene (*17-20*). Subfunctionalization allows different gene copies to express in different
contexts and to acquire mutations that optimize their function within these contexts, such as specific
cellular environments. However, when EGC transfers these mutations to a gene copy specialized for a
different cellular environment, they may impair the function of this copy, as exemplified by $IGMT2_{DE155}$
in leaves (**Fig. 2**). Consequently, amino acids that enhance the environment-specific function of gene
copies should be subject to purifying selection.

To test this, we focused on fixed derived amino acids identified by comparing *A. thaliana* and
Arabidopsis lyrata (*21*) IGMT sequences. We divided each *A. thaliana IGMT* coding sequence into
intervals delimited by adjacent polymorphisms shared with any of the other three gene copies. We then
compared $\log_{10}$-transformed interval lengths with and without codons specifying fixed derived amino
acids, using two-sided *t* tests. Intervals with one or more fixed derived amino acids were significantly
larger than those without for *IGMT1* ($t_{df=19}$ = -3.12, *p* < 0.01), *IGMT3* ($t_{df=46}$ = -2.77, *p* < 0.01), *IGMT2* ($t_{df=42}$
= -2.88, *p* < 0.01), and *IGMT4* ($t_{df=28}$ = -2.42, *p* < 0.05) (**Fig. 4 and fig. S6**). Across all four genes, results
were highly significant ($t_{df=141}$ = -5.72, *p* < 0.0001), suggesting that EGC was indeed selected against
around those amino acids.

Discussion

We used the CRISPR/Cas9 system to dissect a near-cryptic QTL for defense metabolites. We obtained a
set of different combinations of *IGMT* mutants in three genetic backgrounds but we did not obtain all
possible knockout combinations. Consequently, we cannot entirely rule out that some observed effects
result from interactions between gene products, such as heterodimers. However, the tissue-specific
expression of these genes, coupled with the lack of coordinated expression and compensation for
defective genes at the transcript level, argue against this possibility. Our analyses show that *IGM2*
comprises a cluster of QTGs, with DE155 and DE096 both having high and low alleles, but at different
gene copies. Thus, even small-effect QTL can have a complex genetic architecture.

To demonstrate the existence of EGC among *IGMT* copies, we counted sites with multiple
polymorphisms, either within the same gene copy or across different copies. This approach provided a
convenient means to assess EGC without requiring prior knowledge of mutation, recombination, or gene
conversion rates. Independent point mutations are highly unlikely to generate polymorphisms shared
across two or more gene copies, whereas EGC generates such shared polymorphisms, especially visible
when conversion tracts encompass long sequence stretches with multiple nucleotide substitutions.
Quantitative differences in IGMT performance were partially attributable to EGC, as shown by
comparing IGMT2 between two near-isogenic Arabidopsis lines. Both alleles were equally active in roots,
but IGMT2 from DE155, which had a clear signature of EGC, performed worse in leaves compared to its
DE096 counterpart. Furthermore, EGC events have shaped polymorphism patterns across the entire
IGMT gene cluster (**fig. S7**), evidenced by sequence comparison of 28 Arabidopsis accessions, suggesting
that other *IGMT* gene copies may also have experienced performance alterations due to EGC.

Ectopic gene conversion (EGC) has been documented across a wide range of organisms, from bacteria to
humans (*22-30*), indicating its universal influence on gene family evolution. Conversion affects not only
coding sequences but also promoter regions, UTRs and introns (**tables S3 and S4**), potentially altering
regulatory sequences and expression patterns (*28*). EGC tends to increase variation among alleles while
reducing it between paralogs (*31,32*). EGC can spread advantageous mutations across gene copies
(*28,33,34*), but will also place locally adapted variants in a new context where their performance may be
impaired.

Gene duplication is essential for optimizing related functions within a complex fitness landscape of
similar yet distinct tasks (*17-20*). However, EGC, as an unavoidable consequence of having multiple gene

copies, introduces sequence from a gene copy adapted to a specific context into another copy adapted to a different context, as exemplified here. Thereby, EGC generates inferior variants, which, within a population, manifest as QTL segregating fitness variation (**Fig. 5**). Thus, the adaptive process that enables the evolution of specialized gene copies also generates temporary maladaptations until the gene copies have sufficiently diverged to prevent EGC. This can explain why there is more segregating fitness variation within populations than predicted under mutation selection balance (*1*).

References and Notes

1. B. Charlesworth, Causes of natural variation in fitness: Evidence from studies of *Drosophila* populations. *Proc. Natl. Acad. Sci.* **112**, 1662-1669 (2015).

2. D. J. Kliebenstein, J. Kroymann, P. D. Brown, A. Figuth, D. Pedersen, J. Gershenzon, T. Mitchell-Olds, Genetic control of natural variation in *Arabidopsis thaliana* glucosinolate accumulation. *Plant Physiol.* **126**, 811-825 (2001).

3. D. J. Kliebenstein, J. Kroymann, T. Mitchell-Olds, The glucosinolate-myrosinase system in an ecological and evolutionary context. *Curr. Opin. Plant Biol.* **8**, 264-271 (2005).

4. E. K. F. Chan, H. C. Rowe, J. A. Corwin, B. Joseph, D. J. Kliebenstein, Combining genome-wide association mapping and transcriptional networks to identify novel genes controlling glucosinolates in *Arabidopsis thaliana*. *PLOS Biol.* **9**, e1001125 (2011).

5. M. Pfalz, M. Mukhaimar, F. Perreau, J. Kirk, C. I. C. Hansen, C. E. Olsen, N. Agerbirk, J. Kroymann, Methyl transfer in glucosinolate biosynthesis mediated by indole glucosinolate *O*-methyltransferase 5. *Plant Physiol.* **172**, 2190–2203 (2016).

6. P. Bednarek, M. Piślewska-Bednarek, A. Svatoš, B. Schneider, J. Doubskỳ, M. Mansurova, M. Humphry, C. Consonni, R. Panstruga, A. Sanchez-Vallet, A. Molina, P. Schulze-Lefert, A glucosinolate metabolism pathway in living plant cells mediates broad-spectrum antifungal defense. *Science* **323**, 101-106 (2009).

7. N. K. Clay, A. M. Adio, C. Denoux, G. Jander, F. M. Ausubel, Glucosinolate metabolites required for an Arabidopsis innate immune response. *Science* **323**, 95-101 (2009).

8. J. Kim, G. Jander, *Myzus persicae* (Green Peach Aphid) feeding on Arabidopsis induces the formation of a deterrent indole glucosinolate. *Plant J.* **49**, 1008-1019 (2007).

9. M. Pfalz, H. Vogel, J. Kroymann, The gene controlling the *Indole Glucosinolate Modifier 1* QTL alters indole glucosinolate structures and aphid resistance in Arabidopsis. *Plant Cell* **21**, 985-999 (2009).

10. M. Pfalz, H. Vogel, T. Mitchell-Olds, J. Kroymann, Mapping of QTL for resistance against the crucifer specialist herbivore *Pieris brassicae* in a new Arabidopsis inbred line population, Da(1)-12 × Ei-2. *PLOS ONE* **6**, e578 (2007).

11. M. Pfalz, M. D. Mikkelsen, P. Bednarek, C. E. Olsen, B. A. Halkier, J. Kroymann, Metabolic engineering in *Nicotiana benthamiana* reveals key enzyme functions in Arabidopsis indole glucosinolate modification. *Plant Cell* **23**, 716-729 (2011).

12. W. B. Jiao, K. Schneeberger, Chromosome-level assemblies of multiple *Arabidopsis* genomes reveal hotspots of rearrangements with altered evolutionary dynamics. *Nat. Commun.* **11**, 989 (2020).

13. X. Gan, O. Stegle, J. Behr, J. G. Steffen, P. Drewe, K. L. Hildebrand, R. Lyngsoe, S. J. Schultheiss, E. J. Osborne, V.
T. Sreedharan, A. Kahles, R. Bohnert, G. Jean, P. Derwent, P. Kersey, E. J. Belfield, N. P. Harberd, E. Kemen, C.
Toomajian, P. X. Kover, R. M. Clark, G. Rätsch, R. Mott, Multiple reference genomes and transcriptomes for
Arabidopsis thaliana. *Nature* **477**, 419-423 (2011).

14. S. Ossowski, K. Schneeberger, J. I. Lucas-Lledó, N. Warthmann, R. M. Clark, R. G. Shaw, D. Weigel, M. Lynch, The
rate and molecular spectrum of spontaneous mutations in *Arabidopsis thaliana*. *Science* **327**, 92-94 (2010).

15. M. L. Weng, C. Becker, J. Hildebrandt, M. Neumann, M. T. Rutter, R. G. Shaw, D. Weigel,C. B. Fenster, Fine-
grained analysis of spontaneous mutation spectrum and frequency in *Arabidopsis thaliana*. *Genetics* **211**, 703-714
(2019).

16. H. Cao, X. Zhang, F. Li, Z. Han, X. Guo, Y. Zhang, Y., Glucosinolate *O*-methyltransferase mediated callus
formation and affected ROS homeostasis in *Arabidopsis thaliana*. *Physiol. Mol. Biol. Pla.* **30**, 109-111 (2024).

17. A. Force, M. Lynch, F. B. Pickett, A. Amores, Y. L. Yan,J. Postlethwait, Preservation of duplicate genes by
complementary, degenerative mutations. *Genetics* **151**, 1531–1545 (1999).

18. M. Lynch, A. Force, The probability of duplicate gene preservation by subfunctionalization. *Genetics* **154**, 459–
473 (2000).

19. M. Lynch, J. S. Conery, The evolutionary fate and consequences of duplicate genes. *Science* **290**, 1151-1155
(2000).

20. X. He, J. Zhang, Rapid subfunctionalization accompanied by prolonged and substantial neofunctionalization in
duplicate gene evolution. *Genetics* **169**, 1157–1164 (2005).

21. T. T. Hu, P. Pattyn, E. G. Bakker, J. Cao, J. F. Cheng, R. M. Clark, N. Fahlgren, J. A. Fawcett, J. Grimwood, H.
Gundlach, G. Haberer, J. D. Hollister,S. Ossowski, R. P. Ottilar, A. A. Salamov, K. Schneeberger, M. Spannagl, X.
Wang, L. Yang, M. E. Nasrallah, J. Bergelson, J. C. Carrington, B. S. Gaut, J. Schmutz, K. F. X. Mayer, Y. Van de Peer, I.
V. Grigoriev, M. Nordborg, D. Weigel, Y. L. Guo, The *Arabidopsis lyrata* genome sequence and the basis of rapid
genome size change. *Nat. Genet.* **43**, 476-481 (2011).

22. H. Innan, A two-locus gene conversion model with selection and its application to the human RHCE and RHD
genes. *Proc. Natl. Acad. Sci. USA* **100**, 8793-8798 (2003).

23. J. Kroymann, S. Donnerhacke, D. Schnabelrauch, T. Mitchell-Olds, Evolutionary dynamics of an Arabidopsis
insect resistance quantitative trait locus. *Proc. Natl. Acad. Sci. USA* **100**, 14587-14592 (2003).

24. R. T. Morris, G. Drouin, Ectopic gene conversions in four *Escherichia coli* genomes: increased recombination in
pathogenic strains, *J. Mol. Evol.* **58**, 596–605 (2004).

25. N. Osada, H. Innan, Duplication and gene conversion in the Drosophila melanogaster genome. *PLOS Genet.* **4**,
e1000305 (2008).

26. D. Benovoy, G. Drouin, Ectopic gene conversions in the human genome. *Genomics* **93**, 27-32 (2009).

27. J. R. Arguello, T. Connallon, Gene duplication and ectopic gene conversion in *Drosophila*. *Genes* **2**, 131-151
(2011).

28. M. Hanikenne, J. Kroymann, A. Trampczynska, M. Bernal, P. Motte, S. Clemens, U. Krämer, Hard selective sweep and ectopic gene conversion in a gene cluster affording environmental adaptation. *PLOS Genetics* **9**, e1003707 (2013).

29. A. Harpak, X. Lan, Z. Gao, J. K. Pritchard, Frequent nonallelic gene conversion on the human lineage and its effect on the divergence of gene duplicates. *Proc. Natl. Acad. Sci. USA* **114**, 12779-12784 (2017).

30. E. Lamping, J. Y. Zhu, M. Niimi, R. D. Cannon, Role of ectopic gene conversion in the evolution of a *Candida krusei* pleiotropic drug resistance transporter family. *Genetics* **205**, 1619-1639 (2017).

31. H. Innan, A method for estimating the mutation, gene conversion and recombination parameters in small multigene families. *Genetics* **161**, 865-872 (2002).

32. T. Ohta, Allelic and nonallelic homology of a supergene family. *Proc. Natl. Acad. Sci. USA* **79**, 3251–3254 (1982).

33. J. H. Thomas, Concerted evolution of two novel protein families in *Caenorhabditis* species. *Genetics* **172**, 2269–2281 (2006).

34. S. Mano, H. Innan, The evolutionary rate of duplicated genes under concerted evolution. *Genetics* **180**, 493-505 (2008).

35. R. Buchner, in *Glucosinolates in Rapeseed*, J. P. Wathelet, Ed. (Martinus Nijhoff Publishers, 1987), pp. 50-58.

36. F. Fauser, S. Schiml, H. Puchta, Both CRISPR/Cas-based nucleases and nickases can be used efficiently for genome engineering in *Arabidopsis thaliana*. *Plant J.* **79**, 348-359 (2014).

37. C. Morineau, Y. Bellec, F. Tellier, L. Gissot, Z. Kelemen, F. Nogué, J. D. Faur, Selective gene dosage by CRISPR-Cas9 genome editing in hexaploid *Camelina sativa*. *Plant Biotechnol. J.* **15**, 729-739 (2016).

38. S. J. Clough, A. F. Bent, Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J.* **16**, 735-743 (1998).

39. Z. Yang, PAML 4: a program package for phylogenetic analysis by maximum likelihood. *Mol. Biol. Evol.* **24**, 1586-1591 (2007).

40. D. H. Huson, D. Bryant, Application of phylogenetic networks in evolutionary studies. *Mol. Biol. Evol.* **23**, 254-267 (2006).

Acknowledgements

We thank Marine Paupiére for her contribution to QTL mapping.

Funding: Agence Nationale de la Recherche (ANR-10-GENM-005, ANR-20-CE92-0042)

Author contributions: Conceptualization: JK; Methodology: MP, SEN; Investigation: MP, SEN, JK; Funding acquisition: JK; Supervision: JK, MP; Writing: JK, JAS, MP

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: All data are available in the main text or the supplementary
materials.

Supplementary Materials

Materials and Methods

Supplementary Text

Figs. S1 to S7

Tables S1 to S4

Fig. 1. *IGMT* transcript levels in DE155 and DE096. Shown are boxplots with maximum, median and
minimum of ΔC_T values for leaves (L) and roots (R) in comparison to a control gene, *UBQ10*. Each
genotype/transcript combination had n = 3 biological samples. *IGMT1* to *IGMT4* represent individual
genes, while *IGMT1-4* indicates total *IGMT* transcript levels. Note that higher ΔC_T values correspond to
lower expression levels, and *vice versa*.

Fig. 2. Quantitative differences in 4MOI3M and 4OHI3M in leaves and roots. Shown are estimated marginal means of metabolite concentration in nmol per gram fresh weight (± SEM) for Arabidopsis DE155, DE096 and *igmt* mutants. Different upper and lower case letters indicate statistically significant differences. Each genotype had n = 10 or 11 samples. (A) Concentration of 4MOI3M (blue) and 4OHI3M (red). Lines connect meaningful comparisons. (B) Retransformed data for the fraction of 4OHI3M in the combined amount of 4OHI3M and 4MOI3M. Genotypes at *IGMT 1 − 4* are indicated, with "-" for defective and "+" for functional genes. NILs differ significantly for the fraction of 4OHI3M, indicated by asterisks.

Fig. 3. Quantitative effects of *IGMT4* on leaf IGs in DE155 and DE096. Data pooled from four independent experiments show estimated marginal means of metabolite concentration in nmol per gram fresh weight (± SEM) of 4OHI3M (red) and 4MOI3M (blue) with *IGMT4* intact (left) or defective (right) in DE155 (circles and solid lines) and DE096 (diamonds and dashed lines). Each genotype had n = 35 – 61 samples. The interaction of *IGMT4* by background is highly significant for 4MOI3M ($F_{1,164}$ = 18.7; p < 0.0001) but not for 4OHI3M ($F_{1,164}$ = 0.0; p = 0.994). There were no significant genotype or genotype-by-background effects for I3M, 1MOI3M, and total IG.

Fig. 4. Boxplot comparing the log10-transformed interval sizes without (black) and with (red) fixed derived amino acids. These derived amino acids are nonsynonymous substitutions that distinguish Arabidopsis IGMTs from those in *Arabidopsis lyrata*. Each interval is flanked by neighboring shared polymorphisms, defined as the same nucleotides segregating at the same position in two or more gene copies. Shown are mean (crosses), median (black horizontal line), and range of the data including outliers (open circles). Statistical significance is indicated by asterisks (*: $p < 0.05$; **: $p < 0.01$; ***: $p < 0.001$).

Gene duplication

Subfunctionalization
& optimization

EGC

Maladaptation
&
quantitative variation

Fig. 5. Maladaptation caused by EGC in gene families. After gene duplication, the paralogs acquire specific expression patterns and optimize accordingly. EGC introduces sequence from one copy optimized for one context into another copy optimized for a different context, leading to maladaptation. This results in quantitative genetic variation in the population.

Table 1 Estimated Marginal Means - 4MOI3M/(4OHI3M + 4MOI3M) in Roots

Genotype	Mean	SEM	95% Confidence Interval	
			Lower	Upper
DE155	0.533	0.019	0.495	0.570
$igmt1$-4_{DE155}	0.050	0.015	0.020	0.080
DE096	0.361	0.017	0.326	0.396
$igmt1$-4_{DE096}	0.035	0.014	0.006	0.063
Col-0	0.297	0.017	0.264	0.330
$igmt5_{Col-0}$	0.260	0.019	0.223	0.297
$igmt1$-5_{Col-0}	0.000	0.015	-0.030	0.030

388    $8 \leq n \leq 12$

| 390 | **Supplementary Materials** |

| 391 | |

| 392 | **Materials and Methods** |

393	**Culture conditions**
394	Arabidopsis plants were cultivated in growth chambers with 11.5 hours of light at 22°C and 12.5 hours of
395	darkness at 16°C, maintaining approximately around 70 – 80% relative humidity. Seeds were sown on
396	soil and stratified for three days at 6°C. For glucosinolate analysis, one week after germination, plants
397	were transferred to sand and grown randomized in 96-celled trays, fertilized weekly with Hydrocani C2
398	liquid fertilizer (Hydro Agri), as described by (*5*).

399	**Glucosinolate extraction and analysis**
400	Four weeks after germination, approximately 100 g leaf material and the entire root were harvested,
401	weighed and snap-frozen in liquid nitrogen. Glucosinolate extracts were separated in their desulfo-form
402	on a Vanquish HPLC system (Thermo), utilizing a LiChrospher 100 RP-18e LiChroCART column (250 x 4,
403	100A, 5 µm) as described by (*11*). Glucosinolate identification was based on retention time and UV
404	spectra, and quantification was based on integrated absorption peak area at 229 nm, using sinigrin as an
405	external standard and published response factors (*35*) to correct for different UV absorption capacities
406	of various IGs.

407	**RNA extraction and quantitative real-time quantitative PCR (RT-qPCR)**
408	Frozen leaf or root material was finely ground using a TissueLyser II (Qiagen). Total RNA was extracted
409	with TriSure (Bioline) or the NucleoSpin RNA kit (Macherey & Nagel), treated with Turbo DNase
410	(Ambion), and purified using RNeasy MinElute columns (Qiagen) or RNA Clean & Concentrator 5 (Zymo
411	Research). RNA quality and quantity were assessed by agarose gel electrophoresis and a Nanodrop 2000
412	(Thermo). First strand cDNA synthesis was performed with the Maxima First Strand cDNA Synthesis Kit
413	(Thermo), using 500 ng of total RNA. RT-qPCR was conducted with SYBR Green qPCR Master Mix
414	(Thermo) on a StepOnePlus Real Time PCR System (Applied Biosystems), using Arabidopsis *UBQ10*
415	(At4g05320) as the reference gene. Each genotype had three to six biological replicates. Primer
416	sequences are listed in **table S1**.

417	**Generation of mutant plants**
418	Coding sequences of *IGMT1 – 4* from Arabidopsis accessions Col-0, Da(1)-12 and Ei-2 were used to
419	design three single guide RNA (sgRNA) sequences targeting all four genes simultaneously. The design
420	was performed using the CRISPOR software (crispor.tefor.net), with a preference for sites close to the
421	5'-end of the target genes and low off-target probability. A 1026 bp cassette flanked by attB sites,
422	including all three sgRNAs, each driven by an *At*U6-26 promoter and containing the tracrRNA scaffold as
423	well as the U6 terminator, was synthesized by Genewiz (Leipzig, Germany) and cloned into the
424	pDONR207 vector (Thermo). The final cloning step into pDE-Cas9 DsRed (*36,37*), permitting selection by
425	DsRed marker fluorescence, was achieved using Gateway™ LR recombination. This vector was then used
426	to transform *Agrobacterium tumefaciens* GV3101. The floral dip method (*38*) was employed to generate
427	mutants in Arabidopsis DE096, DE155, Col-0 and *igmt5*. After DNA extraction, mutations were identified
428	by Sanger sequencing, using gene-specific primers (**table S1**) covering the target regions of all three
429	sgRNAs.

Statistical analyses

All measurements of were taken from distinct samples. Fine-mapping of *IGM2* was performed using a general linear model in Systat V9, with markers and growth trays as fixed factors. Glucosinolate quantities were analyzed via ANCOVA in jamovi Version 2.3.28.0 (www.jamovi.org), using the weight of the harvested plant tissue as a covariate and genotype as a fixed factor. When pooling data from multiple experiments, experiment was included as an additional fixed factor, along with an interaction term for genotype × experiment. The leaf effect of *IGMT4* was assessed by ANCOVA, with leaf weight as a covariate and genetic background, genotype at *IGMT4* and experiment as fixed factors. The data for the fraction of 4OHI3M in the combined amount of 4OHI3M and 4MOI3M were arcsine square root-transformed before ANOVA, with genotype as a fixed factor. Statistical differences between genotypes were assessed using post-hoc two-sided *t*-tests. PAML (*39*) was used to estimate the transition/transversion ratio in *IGMT* genes. SplitsTree (*40*) was employed to construct a genealogical network for Arabidopsis *IGMT* sequences.

Supplementary text

Calculations of probabilities for different types of nucleotide polymorphisms

We consider a large population with n copies of a gene. Initially, there is no polymorphism within copies
but fixed differences among copies are allowed. We examine cases where two independent point
mutations affect the same site, either twice in the same copy or once in two different copies. The
probabilities of these events are $P_{same\ copy} = n^{-1}$ and $P_{different\ copies} = (1 - n^{-1})$, respectively.

(2 copies: AA*, AB, BA, BB*

3 copies: AA*, AB, AC, BA, BB*, BC, CA, CB, CC*

4 copies: AA*, AB, AC, AD, BA, BB*, BC, BD, CA, CB, CC*, CD, DA, DB, DC, DD*)

Each nucleotide can change into any of the three other nucleotides. There is one possibility for a
transition, t_i (*e.g.*, A -> G) and two possibilities for a transversion, t_v (*e.g.*, A -> C, A -> T). The probability
for a transition is $P_{ti} = x$, the probability for each transversion is $\frac{1}{2} P_{tv} = \frac{1}{2} (1 - x)$. The
transition/transversion ratio t_i/t_v can be estimated from the data.

We first determine the probability that the same mutation occurs twice at the same site, either in one
copy or in two different copies, with μ_1 being the first and μ_2 being the second mutation:

μ_1	$P(X=\mu_1)$	μ_2	$P(X=\mu_2)$	$P(X=\mu_1)*P(X=\mu_2)$
A->C	$\frac{1}{2}(1-x)$	A->C	$\frac{1}{2}(1-x)$	$\frac{1}{4}(1-x)^2$
A->G	x	A->C	$\frac{1}{2}(1-x)$	$\frac{1}{2}x*(1-x)$
A->T	$\frac{1}{2}(1-x)$	A->C	$\frac{1}{2}(1-x)$	$\frac{1}{4}(1-x)^2$
A->C	$\frac{1}{2}(1-x)$	A->G	x	$\frac{1}{2}x*(1-x)$
A->G	x	A->G	x	x^2
A->T	$\frac{1}{2}(1-x)$	A->G	x	$\frac{1}{2}x*(1-x)$
A->C	$\frac{1}{2}(1-x)$	A->T	$\frac{1}{2}(1-x)$	$\frac{1}{4}(1-x)^2$
A->G	$x/(n*l)$	A->T	$\frac{1}{2}(1-x)$	$\frac{1}{2}x*(1-x)$
A->T	$\frac{1}{2}(1-x)$	A->T	$\frac{1}{2}(1-x)$	$\frac{1}{4}(1-x)^2$

The pattern shown above for possible outcomes starting from 'A' is also applicable to the three other
nucleotides. The probability that the same point mutation occurs twice at the same site (red cases) is:

$x^2 + 2 * \frac{1}{4} (1 - x)^2 = x^2 + \frac{1}{2} (1 - 2x + x^2) = 0.5 - x + 1.5 x^2$

Note that identical mutations occurring independently at the same site in different individuals within a
large population go unnoticed.

The probability of different point mutations arising at the same site (blue cases) is:

$1 - (0.5 - x + 1.5 x^2) = 0.5 + x - 1.5 x^2$

If two distinct point mutations occur at the same site in the same copy in different individuals within a
large population, three different nucleotides will segregate at that site.

The first mutation creates a polymorphism at the affected site, and the mutant nucleotide may increase
in frequency (f) within the population. Does this alter the probabilities of the outcomes following a
second mutation at the same site in the same copy?

P (X = 'invisible') = (x * f * x) + (x * (1 − f) * x) + 2 (½ (1 − x) * f * ½ (1 − x)) + 2 (½ (1 − x) * (1 − f) * ½ (1 −
x))

= f * x² + (1 − f) * x² + 2 (¼ f * (1 − x)²) + 2 (¼ (1 − f) * (1 − x)²)

= (f + 1 − f) * x² + 2 (¼ f * (1 − x)² + ¼ (1 − f) * (1 − x)²) = x² + 2 * ¼ (f + 1 − f) * (1 − x)²

= x² + ½ (1 − x)² = x² + ½ (1 − 2x + x²) = x² + 0.5 − x + 0.5 x² = 0.5 − x + 1.5 x²

Thus, the probabilities for 'invisible' events and those where three different nucleotides segregate at the
same site after the second mutation are independent of the mutant allele frequency.

If two mutations hit the same site in the same copy, the probabilities of the outcomes are:

1. **Invisible**: $(0.5 - x + 1.5\, x^2)$

2. **Three nucleotides segregating at the same site** in one copy: $(0.5 + x - 1.5\, x^2)$

If two mutations occur at the same site in different copies, assuming <u>both copies initially have the same</u>
<u>nucleotide</u>, two outcomes are possible:

1. **Shared polymorphism**: Identical mutations result in the same nucleotides segregating at the same
site in different copies. Probability: $(0.5 - x + 1.5\, x^2)$

2. **Copy-specific polymorphisms**: Different mutations differ result in unique polymorphisms at the same
site in each copy. Probability: $(0.5 + x - 1.5\, x^2)$

If the two copies initially had different nucleotides due to an ancient mutation, the outcomes depend on
whether the differences are transitions or transversions.

a) The copies differ by a transition: A & G or C & T

513	μ_1	$P(X=\mu_1)$	μ_2	$P(X=\mu_2)$	$P(X=\mu_1)*P(X=\mu_2)$
514	A->C	½ (1-x)	G->A	x	½ x*(1-x)
515	A->G	x	G->A	x	x²
516	A->T	½ (1-x)	G->A	x	½ x*(1-x)
517					
518	A->C	½ (1-x)	G->C	½ (1-x)	¼ (1-x)²
519	A->G	x	G->C	½ (1-x)	½ x*(1-x)
520	A->T	½ (1-x)	G->C	½ (1-x)	¼ (1-x)²
521					
522	A->C	½ (1-x)	G->T	½ (1-x)	¼ (1-x)²
523	A->G	x	G->T	½ (1-x)	½ x*(1-x)
524	A->T	½ (1-x)	G->T	½ (1-x)	¼ (1-x)²
525					

526   A 'shared polymorphism' (red) arises with probability $P = x^2$. With probability $P = (1 - x^2)$, two copy-
527   specific polymorphisms occur at the same site in both copies.

529   b) The copies differ by a transversion: A & C, A & T, G & C, or G & T

530	μ_1	$P(X=\mu_1)$	μ_2	$P(X=\mu_2)$	$P(X=\mu_1)*P(X=\mu_2)$
531	A->C	½ (1-x)	C->A	½ (1-x)	¼ (1-x)²
532	A->G	x	C->A	½ (1-x)	½ x*(1-x)
533	A->T	½ (1-x)	C->A	½ (1-x)	¼ (1-x)²

535	A->C	½ (1-x)	C->G	½ (1-x)	¼ (1-x)²
536	A->G	x	C->G	½ (1-x)	½ x*(1-x)
537	A->T	½ (1-x)	C->G	½ (1-x)	¼ (1-x)²

539	A->C	½ (1-x)	C->T	x	½ x*(1-x)
540	A->G	x	C->T	x	x²
541	A->T	½ (1-x)	C->T	x	½ x*(1-x)

A 'shared polymorphism' (red) occurs with a probability of $P = \frac{1}{4}(1-x)^2$. With a probability of $P = 1 -$
$(\frac{1}{4}(1-x)^2)$, we obtain two copy-specific polymorphisms at the same site in both copies.

The relative frequencies of scenarios a) and b) are themselves determined by the probabilities of
transitions and transversions. Specifically, in x instances, genes with a transition difference serve as the
starting point. Conversely, in $(1 - x)$ instances, genes with a transversion difference are the starting
point.

Therefore, the probability of obtaining a shared polymorphism starting with a transition difference is
$P = x^3$, and the probability when starting with a transversion difference is $P = \frac{1}{4}(1-x)^3$. This sums up to a
total probability of $P = \frac{1}{4}(1 - 3x + 3x^2 + 3x^3)$ for copies that initially had different nucleotides at the
relevant position. Conversely, the total probability of ending up with two different copy-specific
polymorphisms at the same position in both copies is $P = \frac{3}{4}(1 + x - x^2 - x^3)$.

The exact number of mutable sites in our dataset is uncertain. An initial approximation can be based on
the number of synonymous sites, though many synonymous sites in *IGMT*s remain unmutated.
Conversely, some non-synonymous sites exhibit polymorphisms or substitutions.

To address this, we have explored scenarios ranging from zero to 100% identical nucleotides,
considering that the events of 'two mutations of identical nucleotides at the same position' and 'two
mutations of distinct nucleotides at the same position' are mutually exclusive.

572 Probabilities of different outcomes for two mutations occurring at the same site X:

x: transition probability
n: number of gene copies
N: percentage of identical nucleotides

Fig. S1. Fine mapping of *Indole Glucosinolate Modifier 2*. (A) Indole glucosinolate phenotypes of near-
isogenic lines DE096 (light gray) and DE155, shown as percentages of total indole glucosinolate. (B)
Genotypes of DE096 and DE155 on chromosome 1. DE096 carries parental alleles from the Arabidopsis
accession Ei-2 (light gray) in the *IGM2* candidate (dashed lines) region, whereas DE155 carries alleles
from Da(1)-12 (dark gray). DE096 and DE155 marker genotypes do not differ on chromosomes 2 − 5. (C)
Fine mapping. We fine-mapped the leaf QTL as the arcsine square root-transformed proportion of
4OHI3M among modified indole glucosinolates [c4OHI3M x c(4OHI3M + 4MOI3M)$^{-1}$] in 184 F_2 progeny of
a cross between near isogenic lines DE096 and DE155 with twelve markers. Marker positions refer to
their physical position in the Col-0 reference genome. Statistical support (green line) for the presence of
a QTL peaked close to *IGMT1-4*. Subsequently, we confirmed the presence of a similar QTL in roots
(orange line) with a subset of markers in an independent experiment with 184 F_2 progeny. (D) Estimated
proportion (± SEM) of 4OHI3M by genotype at marker C7676 in leaves (green) and roots (orange). (E)
Organization of IGMT1 − 4. Exon-intron structures are based on the Col-0 accession and drawn to scale,
with the intervening sequence shortened. The gene order is the same in all investigated accessions. (F)
Chemical structures of indole glucosinolates and reactions catalyzed by CYP81F1 − 3 and IGMT 1 − 4.
CYP81F1 − 3 gene products hydroxylate indol-3ylmethyl glucosinolate (I3M) at the 4-position of the
indole ring, producing 4-hydroxy-I3M (4OHI3M). This compound then serves as the substrate for *IGMT1*
− 4 gene products, which generate 4-methoxy-I3M (4MOI3M).

Fig. S2. Schematic representation of *IGMT* wild types and mutants. Locations of sgRNAs (red, blue and green colors) in *IGMT1 – 4*, corresponding sequences and flanking nucleotides (capital letters) in DE096, *igmt5*_{Col-0}, and DE155. Sequence deviations are shown in black. Mutations rendering genes non-functional are shown in colors corresponding to the different target sites. Insertions are denoted by letters, deletions by a Δ proceeding letters (in case of single-nucleotide deletions) or numbers (in case of multiple nucleotide deletions). Larger deletions, some resulting in fusions between parts of adjacent gene copies, are marked by yellow triangles. Note that some of the target sites were still segregating for polymorphisms when analyzed, leading to heterozygous configurations; however, additional mutations at other target sites ensured non-functionality of the gene products.

Fig. S3. *IGMT* and *CYP81F* transcript levels in wild type and mutant leaves. (A) Boxplots showing *IGMT1* – *4* transcript levels in NILs and selected mutant leaves. (B) *CYP81F1* – *F3* and *IGMT5* transcript levels. Data are presented as median and range of ΔC_T values in comparison to the control gene *UBQ10*. Genotypes at *IGMT1* – *4* are indicated, with "-" for defective and "+" for functional genes.

Fig. S4. Genealogical networks of *IGMT* genes. (A) Network based on DE155 and DE096 *IGMT* genes. **(B)**
Network based on *IGMT1 – 4* genes from 28 Arabidopsis accessions, IGMT5, and corresponding
sequences from *A. lyrata*. Colored dots indicate the positions of DE155 (blue) and DE096 (red). The
IGMT5 branch was shortened to 1/5 of its original length. Note that there are two clusters with *IGMT3*
genes.

Fig. S5. Expected proportion of shared among visible types of polymorphisms for different
transition/transversion (t_i/t_v) ratios and copy numbers (n). % identity refers to cases where two
mutations hit different copies at the same site, which may carry either identical or different nucleotides.
Data for Arabidopsis *IGMT1 – 4* are indicated by a solid red line.

Fig. S6. Position of derived amino acids and shared polymorphisms in *IGMT1 – 4* coding sequences. A polymorphism is considered 'shared' when the same nucleotides segregate at the same site in two or more gene copies. Intervals flanked by two adjacent shared polymorphisms (black) are significantly larger when they contain derived amino acids (red). Positions are shown in base pairs.

Fig. S7. Number of shared polymorphisms in pairwise comparisons of *IGMT* genes. Note that shared polymorphisms are more frequent when genes are in close proximity. Also, the two central genes exhibit the highest number of shared polymorphisms.

Table S1. Primer sequences used in this study.

Fine mapping of *IGM2*			
Marker	Sequence (5' → 3')	Sequence (5' → 3')	Physical position [Mbp] on Chr 1 in Col-0
C1P42	GACATAGTTGGCGATGATTG	CAAACAATGGGAGAAGTTGGTC	1.34
C1P43	CTTTTGGAGACACTCAAAGTTG	TAGAGTAAGGTACAAGACCTC	3.02
c3519	CAGTTTGGTAGTTGGATCTGG	GTATAACACTAATCCGAGAACC	3.52
c4556	CGCATGATTCGTCTTGTAAC	CTTTAGAGAGAGGGATCAATAC	4.56
c5020	AGTTTGCTGGTGGATTCGTG	CATTGCATGGCCAAATTCCTC	5.02
F20D23_2	CCGTCACACCATTCACAATC	CCAACCCCTTATATATCGTTC	5.82
c6061	ACTAAAGGAAGAGATCAATCAC	GCTGATGCTTAGTCCTTTC	6.06
c7676	GTGATTTAGTTAGGCATTAAGTTTC	CTCTATATTCTCAGCTTTCTTTC	7.68
M3	GCTTTTATCAGCTCAAACAT	ACTCTTACGTTTGGAGTTCA	7.99
c8654	CACCTGATCTAACGAGTGTTTC	GTTAGCATTTTTATGCCACTATTAG	8.65
M5	TCTGTATCTCGGTGAATTCTCC	TACCGAACCAAAACACAAAG	9.89
F1K23	GAACCAATAAGGAGGCTCAAC	CCATACGGAGAAACCTTCTTC	10.00

Amplification and sequencing of *IGMT1-4* promoters & genes		
Target	Primer name	Sequence (5' → 3')
IGMT1 (At1g21100)	PromMT 100_1F	CATTTTCCTTTGTTTTTGTTTC
	PromMT 100_F	TTTCCTTTGTTTTTGTTTCTTTCTTC
	PromMT 100_R	CTATAACAATTGAGATGGGAAAC
IGMT3 (At1g21110)	PromMT 110_F	TTCTCTTTCTTTTTTGTTTCTTTCTTC
	PromMT 110_1F	CATTTCTCTTTCTTTTTTGTTTC
	PromMT 110_R	CAACTGTTTCTAAATGGCTATC
	PromMT 110_2R	GTAGCGGTGTAATCATTACTTC
	PromMT 110_3R	GGTATAAGAGATATTGCGTGTG
IGMT2 (At1g21120)	PromMT 120_F	TTCTCTTTCTTTTTTGTTACTTTCTTC
	PromMT 120_1F	CATTTCTCTTTCTTTTTTGTTAC
	PromMT 120_R	GAATCAAACCGACGATGAAC
	PromMT 120_2R	GACTAACGGCCGAAAGGGAC
IGMT4 (At1g21130)	PromMT 130_1F	CATTTTATTTCTTTTTGTTTCTTT
	PromMT 130_R	GAAACTTAACTGGGCCAAGC
	PromMT 130_2R	GCATAACGTAACAGACCATG
	RIK-3F	GGGATACCTTCTAGAAGAAAC
	RIK-3R	CAGAACTCAATAATCCAGCAG

Detection of changes in sgRNA sequences (PCR & Sanger sequencing)		
Target	Primer name	Sequence (5' → 3')
IGMT1 (At1g21100)	gRNA21100F	GAAAGTAGCCGACCGTGGAG
	gRNA21100R	GCACAAGAGGCTCTTACTTTTC
IGMT2 (At1g21120)	gRNA21120F	CAGAACTCAATAATCCAGCAGT
	gRNA21120R	CTCTTACCTTTCTAAACAGAGAC
IGMT3 (At1g21110)	gRNA21110F	CAAATGAAAGCAGCTGGTCTTG
	gRNA21110R	CTACACAGAGACTACTAACTCAG
IGMT4 (At1g21130)	gRNA21130F	CAGAACTCAATAATCCAGCAGT
	gRNA21130R	CTACGGTACATTACATAGAGAC

Real Time quantitative PCR		
Target	Primer name	Sequence (5' → 3')
IGMT1 (At1g21100)	RT-MT100for	CTCTCACCCTATGAAATAGCAA
	RT-MT100rev	GCTCTGTAGACCCTCTCGC
IGMT2 (At1g21120)	120new_f	CATGGTCAAGTGTGAGAAGGTC
	120new_r	GAGGCAAGGGATCCAATATCC
IGMT3 (At1g21110)	110new_f	CATGGTCAAGTGTGAGAAG
	110new_r	GTTTGCGTTGATGTCACC
IGMT4 (At1g21130)	130new_f	GCGAGAGGCTTCAGCATAGAG
	130new_r	GGGATACCTTCTAGAAGAAACC
IGMT5 (At5g76790)	RT-76790F	CAGTCCAGTCATGAAGTATAC
	RT-76790R	GTGCTTTAACACAAGCTCCTTC
CYP81F1 (At4g37430)	RT-CYP430F	GAACCGCTCTTATCTGATTG
	RT-CYP430R	GGAAATAGCCGGAATGTTTC
CYP81F2 (At5g57220)	Cyt2F	GTCACAGGGAGACGCTACTAC
	Cyt2R	CACCACTGTTGTCATTGATGTC
CYP81F3 (At4g37400)	RT-CYP400F	GCCTAATGCTGGGCATGATG
	RT-CYP400R	CTTTAATATCCTCTGTTGGTG
CYP81F4 (At4g37410)	37410seq2	CCCTGAGTACTACACTGAC
	RT-410-(09)R	CAAAACTTTTGGGTGGTTCAAC
IGMT1-4	Universal (IGMT)-for	ACGTATACTTCATGATTGGAC
	Universal (IGMT)-rev	CAATAATCCAGCAGTGATAAG
UBQ10 (At4g05320)	UBQ10f	AAAGAGATAACAGGAACGGAAACATAGT
	UBQ10r	GGCCTTGTATAATCCCTGATGAATAAG

641 **Table S2. Arabidopsis accessions used in this study.**

Accession	Country	Note
Col-0	Poland	reference genome
An-1	Belgium	
C24		
Cvi-0	Cape Verdi Islands	
Eri-1	Sweden	Jiao & Schneeberger, 2020
Kyo	Japan	
Ler	Poland	
Sha	Tadjikistan	
Bur-0	Ireland	
Can-0	Canary Isles	
Ct-1	Italy	
Edi-0	Scotland	
Hi-0	Netherlands	
Kn-0	Lithuania	
Ler-0	Poland	
Mt-0	Libya	
No-0	Germany	
Oy-0	Norway	Gan et al., 2011
Po-0	Germany	
Rsch-4	Russia	
Sf-2	Spain	
Tsu-0	Japan	
Wil-2	Russia	
Ws-0	Russia	
Wu-0	Germany	
Zu-0	Germany	
Da(1)-12	Czech Republic	
Ei-2	Germany	*IGMT1*, *2*, *3*, *4*; this study
No-0	Germany	
Aa-0	Germany	
Ag-0	France	
Bl-1	Italy	
Cal-0	United Kingdom	
Can-0	Canary Islands	
Cnt-1	Canterbury	
Cvi-0	Cape Verdi Islands	
Ema-1	United Kingdom	near-complete *IGMT4*; this study
Hodja	Tadjikistan	
Kas-0	India	
Ler-0	Germany	
Lip-0	Poland	
Ma		
Mr-0	Italy	
Oy-0	Norway	
Tsu-0	Japan	

Table S3. Shared polymorphisms in the 5'UTR of *IGMT1* and *2*.

IGMT1		
An-1	G T T **A T** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T
Bur-0	G T T **T C** T T T C T T C T C **T** T T C A A G G C A C T C G A T C T A C G T G A T C A **C** A C T G A T
C24	G T T **A T** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T
Can-0	G T T **A T** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T
Col-0	G T T **T C** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T
Ct-1	G T T **T C** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T
Cvi	G T T **T C** T T T C T T C T C **T** T T C A A G G C A C T C G A T C T A C G T G A T C A **C** A C T G A T
Da(1)-12	
Edi-0	T T T **T C** T T T C T T C T C **T** T T C A A G G C A C T C G A T C T A C G T G A T C A **C** A C T G A T
Ei-2	G T T **T C** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T
Eri	G T T **T C** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T
Hi-0	G T T **T C** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T
Kn-0	G T T **T C** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T
Kyo	G T T **T C** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T
Ler	G T T **T C** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T
Ler-0	G T T **T C** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A **M** A C T G A T
Mt-0	G T T **T C** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T
No	G T T **T C** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T
Oy-0	T T T T **S** T T T C T T C T C **C** T T C A A G G C A C T C G A T C T A C G T G A T C A **C** A C T G A T
Po-0	G T T **T C** T T T C T T C T C **T** T T C A A G G C A C T C G A T C T A C G T G A T C A A **A** A C T G A T
Rsch-4	G T T **T C** T T T C T T C T C **C** T T C A A G G C A C T C - - - T A C G T G A T C A A **A** A C T G A T
Sf-2	G T T **A T** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T
Sha	G T T **A T** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T
Tsu-0	G T T **A T** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T
Wil-2	G T T **T C** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T
Ws-0	G T T **T C** T T T C T T C T C **T** T T C A A G G C A C T C G A T C T A C G T G A T C A **C** A C T G A T
Wu-0	G T T **T C** T T T C T T C T C **C** T T C A A G G C A C T C - - - T A C G T G A T C A A **A** A C T G A T
Zu-0	G T T **T C** T T T C T T C T C **C** T T C A A G G C A C T C - - - T A C G T G A T C A A **A** A C T G A T

IGMT2		
An-1	G T T **A T** T T T C T T C T C **C** T T C A A G G C A C T C - - - T A C G T G A T C A A **A** A C T G A T
Bur-0	G T T **T C** T T T C T T C T C **T** T T C A A G G C A C T C G A T C T A C G T G A T C A **C** A C T G A T
C24	G T T **A T** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T
Can-0	G T T **A T** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T
Col-0	G T T **A C** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T
Ct-1	G T T **A C** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T
Cvi	G T T **T C** T T T C T T C T C **T** T T C A A G G C A C T C G A T C T A C G T G A T C A **C** A C T G A T
Da(1)-12	G T T **T C** T T T C T T C T C **T** T T C A A G G C A C T C G A T C T A C G T G A T C A **C** A C T G A T
Edi-0	G T T **T C** T T T C T T C T C **T** T T C A A G G C A C T C G A T C T A C G T G A T C A **C** A C T G A T
Ei-2	G T T **A C** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T
Eri	G T T **T C** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T
Hi-0	G T T **A C** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T
Kn-0	G T T **A C** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T
Kyo	G T T **A T** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T
Ler	G T T **T C** T T T C T T C T C **T** T T C A A G G C A C T C G A T C T A C G T G A T C A **C** A C T G A T
Ler-0	G T T **A C** T T T C T T C T C **C** T T C A A G G C A C T C G A T C T A C G T G A T C A A **A** A C T G A T
Mt-0	G T T **A T** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T
No	G T T **A T** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T
Oy-0	G T T **A C** T T T C T T C T C **C** T T C A A G G C A C T C G A T C T A C G T G A T C A A **A** A C T G A T
Po-0	G T T **A T** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T
Rsch-4	G T T **A C** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T
Sf-2	G T T **A T** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T
Sha	G T T **A T** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T
Tsu-0	G T T **A T** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T
Wil-2	G T T **A T** T T T C T T C T C **C** T T C A A G G C A C T C G A T C T A C G T G A T C A A **A** A C T G A T
Ws-0	G T T **A C** T T T C T T C T C **C** T T C A A G G C A C T C G A T C T A C G T G A T C A **C** A C T G A T
Wu-0	G T T **T C** T T T C T T C T C **T** T T C A A G G C A C T C G A T C T A C G T G A T C A **C** A C T G A T
Zu-0	G T T **A C** T T T C T T C T C **C** T T C A A G G C A C T C - - - - T A C G T G A T C A A **A** A C T G A T

646 Shared polymorphisms ↑↑   ↑   ↑↑↑↑   ↑

Table S4. Shared polymorphisms in an *IGMT* intron.