Long-term adaptation to hypoxia provides insight into

mechanisms facilitating the switch of *Pseudomonas*

***aeruginosa* to chronic lung infections.**

Joanna Drabinska[1,2], Lucia O'Connor[1,2], Niamh Duggan[1,2], Ciarán J. Carey[1,2], Siobhán McClean[1,2]*

[1]School of Biomolecular and Biomedical Sciences, University College Dublin, Belfield, Dublin, Ireland.

[2]Conway Institute of Biomolecular and Biomedical Science, University College Dublin, Belfield, Dublin,
Ireland.

*Corresponding author

Email: siobhan.mcclean@ucd.ie

Abstract

Opportunistic bacterial infections emerging from the environment are an increasing threat, especially for immunocompromised individuals such as people with cystic fibrosis (CF), driving morbidity, and mortality. *Pseudomonas aeruginosa* is a key pathogen from this group associated with chronic lung infections that has been extensively studied in this context, but the mechanism driving its adaptation towards chronic colonisation in the CF lung is not fully understood.

This work focuses on the adaptations of *P. aeruginosa* to long-term hypoxia, one of the important environmental pressures present in the CF lung, to investigate whether it drives the development of persistence in CF patients. We examined how an early CF isolate adapted to 6% oxygen over 28 days using an experimental evolution approach and showed that two distinct *P. aeruginosa* small colony variants (SCVs) emerged, one exclusively in hypoxia exposure cultures. Importantly, SCVs were more prevalent in hypoxia-adapted cultures than in normoxia-adapted cultures (98% vs <35%). Proteomic analysis revealed significant changes in the abundance of >200 proteins within 28 days, including proteins involved in antibiotic resistance, stress response, iron acquisition, biofilm formation and those previously associated with chronic infection. Importantly the observed changes in the proteome were distinct from changes reported after short-term exposure to hypoxia. Phenotypic characterisation of adapted cultures supported these findings, showing that hypoxia-adapted cultures developed higher resistance to 8 of 13 antibiotics tested; increased biofilm (1.8-4.1-fold (p<0.0001) and decreased pyocyanin production (2.6-fold, p<0.0001). All hypoxia-adapted cultures showed decreased siderophore production (2.9-3.6-fold, p<0.02). Two distinct patterns of adaptation to hypoxic conditions were observed, with the phenotypes of one culture being markedly different from the other two. Overall, we demonstrate that long-term hypoxia exposure contributes to multiple phenotype and proteome changes that are also observed in *P. aeruginosa* CF lung chronic infection, possibly opening the path to treatment of chronic infections.

Author Summary

Opportunistic antibiotic-resistant bacterial infections emerging from the natural environment are an increasing threat, particularly for immunocompromised individuals such as people with cystic fibrosis (CF). These infections increase disease severity, the number of deaths and healthcare costs. *Pseudomonas aeruginosa* is a major pathogen in this context, causing long-term infections, yet the mechanisms driving its adaptation towards chronic colonisation in the CF lung are not understood.

We investigated the adaptations of *P. aeruginosa* exposed to low oxygen conditions (an important environmental pressure in the CF lung), for 28 days, to examine whether it drives the process of persistence. We found that prolonged exposure to hypoxia induced changes in the pathogen's characteristics, which were distinct from those reported after short term exposure. We observed multiple adaptations in hypoxia adapted cultures, many of which were associated with important signalling networks linked to lifestyle shifts in most bacteria and several of them are also observed in isolates that had adapted to chronic infection. Similar adaptive responses in other opportunistic environmental pathogens suggest that this adaptation could be targeted to create universal therapeutic approaches preventing antibiotic overuse and improving patient outcomes.

Key words

Pseudomonas aeruginosa; hypoxia; adaptation; experimental evolution; chronic infection; cystic fibrosis

Introduction

Antibiotic-resistant bacterial infections are a growing threat to society, were associated with nearly five million deaths in 2019 (1), and are expected to cause more than 10 million deaths annually by 2050 (2). Opportunistic pathogens such as *Pseudomonas aeruginosa* play a crucial role in escalating this crisis. While these bacteria typically do not cause infections in healthy people, immunocompromised individuals and people with underlying conditions such as cystic fibrosis (CF) and chronic obstructive pulmonary disease (COPD) or those on ventilator support are at great risk of developing chronic *P. aeruginosa* lung infections (3,4). Moreover, opportunistic pathogens are responsible for the majority of healthcare-associated infections, contributing to increasing morbidity, mortality and costs (5).

P. aeruginosa is a Gram-negative opportunistic human pathogen, a leading cause of nosocomial infections, constituting 11-26% of ventilator-associated pneumonia globally (6). Overall, it causes more than 7% of all health-associated infections and is a major contributor to morbidity and mortality among people with CF (7–9). It has many intrinsic and acquired mechanisms of antibiotic resistance contributing to its spread and persistence (10), and prolonged antibiotic treatments necessary to manage chronic infections contribute to the spread of antimicrobial resistance (11,12). Furthermore, the World Health Organisation recognised *P. aeruginosa* as a high priority pathogen in 2024 while the ECDC considers it a serious threat (13–16).

A key feature of *P. aeruginosa* is its exquisite metabolic flexibility and adaptability to harsh conditions, allowing it to transition from the colonisation of the rhizosphere to colonising niches within human or animal hosts (4), and establishing persistent infections (17). *P. aeruginosa* isolates from chronic infections generally show different characteristics to those isolated from acute infections and chronic infections are extremely difficult to eradicate. Its adaptability is driven by a complex regulatory

network involving transcriptional regulators, DNA modifications, non-coding RNAs, quorum sensing,

and a range of virulence factors (18–23). Despite this extensive research, the exact mechanism(s)

enabling *P. aeruginosa* to transition from acute to chronic infection remain unclear.

P. aeruginosa, a facultative anaerobe, thrives under low oxygen/anaerobic conditions using

alternative terminal electron acceptors and/or fermentation substrates. Previous studies showed that

P. aeruginosa preferentially colonizes hypoxic niches within mucus layers of the lung over more

oxygenated surfaces (24). *P. aeruginosa* responds to changes in oxygen levels via Anr, which regulates

the expression many of the genes shown to be altered in CF chronic infection (25,26). Studies on

another CF-associated pathogen, *Burkholderia cenocepacia* have shown that 40% of genes within a

low oxygen activated (Lxa) locus were upregulated in long-term infection isolates from CF patients

(27). We subsequently showed that deletion of single genes from the Lxa or the entire locus resulted

in changes in phenotypes crucial for the virulence of this bacterium including host cell attachment,

intracellular survival and stress responses (27,28). Given that hypoxia is an important environmental

pressure present in the lung, we wanted to evaluate the impact of long-term hypoxia exposure on *P.*

aeruginosa.

To examine the effect of long-term exposure to a low oxygen environment, we developed an

experimental evolution approach and demonstrated the significance of long-term exposure to low

oxygen conditions on the adaptation of *P. aeruginosa* to chronic lung infections. Our findings provide

new insights into the mechanisms that underlie the persistence and virulence of opportunistic

bacteria within the host.

Results

Design and Optimisation of Experimental Evolution

To investigate the impact of long-term exposure to low oxygen conditions on the adaptation of *P. aeruginosa*, we designed an experimental evolution approach (Fig 1A) exposing the *P. aeruginosa* early infection strain AMT 0023-30 (29,30) to either 6% oxygen or to normoxia (21%) for 28 days. This is a well-characterized and sequenced cystic fibrosis (CF) paediatric early infection strain that was selected from the international *P. aeruginosa* panel, for which a sequential chronic infection isolate is available (AMT 0023-31) (30–32). After considering ranges of oxygen bacteria are exposed to in soil and water, the range of hypoxic conditions in the CF lung and available literature on physiological tissue hypoxia (24,33,34), we chose 6% oxygen as it allowed us to observe the response to hypoxia, while not extensively stressing the cells and distinguish this response from changes associated with anoxia.

Daily culture transfer volumes were optimised by evaluating a range of volumes (1-10%) of overnight *P. aeruginosa* cultures transferred into fresh media and analysing their growth patterns (S1 Figure). A 5% volume transfer was chosen as optimal, as it allowed the transfer of the maximum number of bacterial cells without disrupting the growth pattern and resulted in a starting OD_{600} of ~0.1. Examination of growth patterns of the early infection strain revealed a significant difference in the generation times observed during hypoxic and normoxic conditions, with the strain growing 1.8-fold slower under hypoxia (g = 107 minutes) compared to normoxia (g = 60 minutes) (S2 Figure). Consequently, during the 28 day adaptation period, the cultures exposed to hypoxia underwent only 377 generations in contrast to 677 generations under normoxic conditions.

To facilitate further investigation, including phenotyping, proteome analysis, and whole genome

sequencing (WGS), three independent flasks were inoculated and incubated under either 6% oxygen

or normoxia (21% O_2), and samples were collected every day over 28 days, at time of culture transfer.

The three 28 day independently adapted populations are referred to as N1, N2, and N3 for replicate

normoxia-adapted control populations (NACs) and H1, H2, and H3 for the three replicate hypoxia-

adapted cultures (HACs) (Table 1). All the phenotypic characterisation of the individual adapted

populations (Both HACs and NACs) was performed under normoxic conditions, to ensure the

observed changes are stable in the absence of continued hypoxia.

Table 1. List of strains and populations used in this study

	Strain/Population Name	Description	Source
	E	Paediatric early infection cystic fibrosis strain AMT 0023-30	(29)
	L	Paediatric early infection cystic fibrosis strain AMT 0023-31	(29)
NACs	N1	28 day normoxia-adapted population, replicate 1	This study
	N2	28 day normoxia-adapted population, replicate 2	This study
	N3	28 day normoxia-adapted population, replicate 3	This study
HACs	H1	28 day hypoxia-adapted population, replicate 1	This study
	H2	28 day hypoxia-adapted population, replicate 2	This study
	H3	28 day hypoxia-adapted population, replicate 3	This study

Changes in colony morphology and pyocyanin production during

adaptation

Changes in colony morphology often serve as key indicators of adaptation to the environment. Plating

of daily samples from the adapting cultures showed the emergence of small colony variants (SCVs)

both in NACs and HACs, however, the SCVs appeared earlier during hypoxia exposure (Fig 1B). SCVs

first appeared on day 2 in the H3 population, and by day 3 in the H1 and H2 populations. The SCV

abundance peaked at 98% on day 22, with 77% remaining on day 28 in H1. Similarly, the highest

proportion of SCVs was reached on day 20 with 82% of the population and 80% on the last day of the

experimental evolution experiment in flask H2. In contrast, in population H3 the SCV abundance was

lower and more variable, reaching only 50% of the population by day 15, and falling to 1% on day 21.

The SCVs appeared much later in normoxic cultures, on days 6, 7 and 8 in flasks N1, N2 and N3

respectively and only reached a maximum of 25-33% of the population by day 14 (Fig 2A).

Two distinct SCV morphologies were observed (Fig 1C and D). The first type, termed SCV 1, was

observed both under normoxia and hypoxia and resembled the morphology most often described as

a rugose small colony variant characterised by small size, irregular edges and a heavily wrinkled, dry

surface (35). The second type, SCV 2, was slightly bigger than SCV 1 with a similar rough, dry surface;

however it was easily distinguished by a more regular edge and shape and importantly, it was

exclusive to HACs.

Throughout the 28 days of adaptation, we also observed dynamic changes in culture colour (Fig 2A),

suggesting changes in pigments, such as pyocyanin. Quantification of pyocyanin production at 28

days, despite almost no visible change of colour being observed on the day 28 of the experiment (Fig

2A), revealed that populations H1, H2 and one NAC, N1 produced comparable pyocyanin levels to the

late infection strain (AMT 0023-31) (Fig 2B and Fig 2C), suggesting that pyocyanin production is

independent of environmental oxygen levels.

Influence of hypoxia on the whole proteome

The proteomes of 28 day adapted cultures were analysed to elucidate the overall influence of long-

term hypoxia on *P. aeruginosa*. The individual flasks were treated as technical replicates, thus showing

the global changes across 3 independent populations. A total of 244 proteins showed significant

differences (p≤0.05) in abundance between 28 day HACs and 28 day NACs, of which 205 were altered

by 1.5-fold or more (Table 2, S3. Table). The majority of these proteins showed decreased abundance

following exposure to hypoxic conditions (n=148; 72.55%), while only 56 proteins (27.45%) showed

increased abundance (Table 2). A constructed heatmap showed divergence between the samples

from cultures adapted to the same condition (Fig 3A).

Table 2. Proteome comparison of 28 day hypoxia-adapted PA populations to normoxia-adapted populations.

Protein groups	Number of proteins*
Total number of identified proteins	1718
Proteins with significantly changed abundance (total)	244
Proteins with significantly changed abundance (by ≥1.5-fold)	205
Proteins unique to hypoxia	5
Proteins unique to normoxia (absent in hypoxia)	16
Proteins with a significantly increased abundance (by ≥1.5-fold) in hypoxia	56
Proteins with a significantly decreased abundance (by ≥1.5-fold) in hypoxia	149

*The proteins with significantly changed abundance were determined using mean LFQ intensities of three independent populations of normoxia and hypoxia-adapted cultures. A fold-change cutoff ≥1.5 was to compare hypoxic and normoxic conditions (T.test $p \leq 0.05$)

Functional enrichment

GO biological process functional enrichment analysis identified 35 enriched categories (S4 Table), including those associated with stimulus response, iron-sulphur cluster assembly, and amino acid metabolism. A separate analysis of proteins with increased or decreased abundance in HACs provided further insights into these important processes (Table 3, S2 Table). Overall, 38 GO biological processes were enriched in the increased abundance in hypoxia group (Fig 3B), while 37 processes were enriched in the decreased abundance group (Figure 3B; S2 Table). As expected, hypoxia led to upregulation of processes related to oxygen transport and nitrogen metabolism. Additionally, eight categories relating to nucleotide, nucleoside, ribonucleotide, and ribose phosphate processes were upregulated in hypoxia, together with those associated with gene expression, translation, peptide biosynthesis, and amide biosynthesis. Glutamate biosynthesis and glutamine family amino acid biosynthesis were the only two processes related to amino acid metabolism associated with increased abundance in HACs. In contrast, 18 categories related to amino acid metabolism, including those involving valine, tyrosine, phenylalanine, glutamate, and glutamine, were in the reduced abundance group, suggesting they may be downregulated. Sulphur and iron-sulphur cluster associated processes, chemotaxis, and motility were also associated with proteins that were reduced in abundance.

Table 3. Comparison of GO term biological function enrichment of proteins increased in abundance, decreased in abundance or both in the 28 day hypoxia adapted cultures

Increased in abundance	Decreased in abundance
Amino acid metabolism *Glutamate biosynthesis*	Amino acid metabolism *Glutamate metabolism* *L-Phenylalanine metabolism* *Tyrosine catabolism* *Valine metabolism*
Oxygen transport	Iron-sulphur cluster assembley
Nucleotide metabolism *Nucleotide biosynthesis* *Ribonucleotide metabolism*	Response to stimulus *Locomotion* *Chemotaxis*
Protein metabolism	
Peptide biosynthesis	
Amide biosynthesis	
Gene expression	

Blue: functional enrichment of proteins increased in abundance in hypoxia adapted cultures; Orange: functional enrichment of proteins decreased in abundance in hypoxia adapted cultures.

A STRING local network cluster analysis confirmed that the processes associated with proteins that were increased in abundance included translation, protein synthesis, and nucleoside synthesis in hypoxia. It also revealed enrichment in proteins associated with glycolysis and pentose phosphate pathways, including those that were increased and decreased in abundance. Additionally, two mixed categories comprising the *P. aeruginosa* biofilm formation pathway appeared to be upregulated, based on changes in protein abundance. Beyond the downregulation of amino acid and chemotaxis-related pathways identified by GO term enrichment analysis, STRING analysis also highlighted reduced abundance of proteins associated with siderophore production, lipopolysaccharide biosynthesis, fatty acid degradation, quinone biosynthesis, and stress response pathways in hypoxia (S4 Table). KEGG pathway enrichment analysis further highlighted the upregulation of pyrimidine metabolism, ribosome biogenesis, and the alanine, aspartate, and glutamate pathways, alongside the downregulation of siderophore biosynthesis, propanoate metabolism, and amino acid metabolism under hypoxia (S4 Table).

Proteins unique to hypoxia or normoxia

Proteins unique to hypoxia or normoxia

Interestingly, 8.61 % of proteins that showed changes in abundance were either unique to HACs or

NACs, as they were present in one condition and were not detected in any of the 3 replicates in the

other group. Their consistent presence or absence across all three HACs suggests that their expression

is either switched on or off, and consequently, they play critical roles in adapting to low oxygen

environments. Only five proteins were unique to HACs (Table 4). These were associated with antibiotic

resistance, exopolysaccharide transport, glutamine and arginine metabolism and cell division. The

protein showing the highest abundance among these was PA4456, the ATP-binding component of an

ABC transporter. Although other ABC transporter components also showed changes in abundance,

they were less dramatic, with fold changes below 6.5-fold. Another protein unique to hypoxia, outer

membrane lipoprotein PelC is essential for Pel exopolysaccharide export (36)**.** Glutamate

dehydrogenase GdhA, an enzyme in the glutamate biosynthesis pathway and arginine catabolism,

was also exclusively detected under hypoxic conditions. Finally, two hypothetical proteins were also

unique to HACs, PA3402, involved in transmembrane transport (37), and PA4278, identified as an *E.*

coli DedD homolog, a crucial component of the bacterial divisome (38).

Proteins that were absent in HACs, and thus potentially redundant in low oxygen conditions, are

involved in essential processes such as stress resistance, virulence, regulation (two-component

systems), polyamine catabolism, tryptophan biosynthesis, QS, bacterial competition, motility,

attachment, biofilm formation and antibiotic resistance (Table 4). Surprisingly, the NarL component

of the nitrate-responsive, Anr-regulated system crucial for anaerobic metabolism (39,40), was

undetectable in hypoxic conditions. Proteins important for *P. aeruginosa* virulence were also

undetectable in the HACs, including the HptA histidine phosphotransporter, a component of an

unorthodox/hybrid two-component system. NqrA NADH dehydrogenase is involved in bacterial

competition and protects *P. aeruginosa* against autotoxicity (41), while AsnB asparagine synthase and

YhfE glutamyl aminopeptidase are both involved in osmotic stress resistance and cell wall stress

response (42–45) and the Sbp substrate binding protein is associated with sulphate starvation

response and virulence (46,47), were all absent in HACs. Examples of proteins associated with biofilm

formation and/or motility that were absent in HACs, include the RidA deaminase (48,49), the PlaB

phospholipase B (50) and the PelC exopolysaccharide export associated protein.

Table 4. Proteins unique to hypoxia-adapted or normoxia-adapted cultures grouped according to

function.

Gene ID[a]	Protein Name	Protein Product Name	H/N[b]	NP[c]	MW[d]	SC[e]
Stress response						
PA3459	AsnB	probable glutamine amidotransferase		8	66.03	17.50
PA3461	YhfE	conserved hypothetical protein		6	43.29	21.60
PA0283	Sbp	sulfate-binding protein precursor		4	36.43	20.20
Biofilm/ exopolysaccharide production						
PA3062	PelC	PelC		6	18.68	58.10
PA0537	PA0537	conserved hypothetical protein		4	22.18	25.20
PA5234	PA5234	probable oxidoreductase		4	34.01	14.50
PA2927	PlaB	hypothetical protein		4	49.48	5.20
PA3123	RidA	RidA subfamily protein		4	34.40	13.52
Amino acid metabolism						
PA4588	GdhA	glutamate dehydrogenase		4	48.85	13.5
PA3713	SpdH	spermidine dehydrogenase SpdH		9	68.86	19.5
PA0036	TrpB	tryptophan synthase beta chain		6	43.68	18.4
Transport						
PA4456	YrbF	probable ATP-binding component of ABC transporter		5	29.62	26.00
PA3402	PA3402	hypothetical protein		5	36.21	21.10
Other						
PA4278	PA4278	hypothetical protein		3	26.08	10.70
PA0991	HptA	Histidine phosphotransfer protein HptA		3	38.40	13.29
PA2999	NqrA	Na+-translocating NADH: ubiquinone oxidoreductase subunit Nrq1		2	48.08	6.10
PA3879	NarL	two-component response regulator NarL		4	24.26	30.10
PA2786	PA2786	hypothetical protein		4	17.54	33.50
PA3808	IscX	conserved hypothetical protein		4	7.80	62.10
Function unknown						
PA2530	PA2530	hypothetical protein		7	48.96	22.00
PA5136	PA5136	hypothetical protein		3	54.32	6.90

[a]Gene ID – *Pseudomonas aeruginosa* PAO1 gene locus ID
[b]H/N – Blue: proteins unique to hypoxia-adapted cultures; Red: Proteins absent in hypoxia-adapted cultures
[c]NP - Unique peptides: The number of peptide sequences unique to a protein
[d]MW - Molecular weight as determined by Q-Exactive LC-MS using *P. aeruginosa* PAO1 reference
[e]SC - % sequence covered by matching peptides for the identified protein

Antibiotic resistance of adapted cultures

A change in the abundance of 48 proteins associated with antibiotic resistance was observed in HACs

(Table 5), with 25 showing increased abundance. Notably, type VI secretion system proteins, and TopA

topoisomerase, all previously associated with resistance to various classes of antibiotics, were increased in abundance; however, in contrast, MexA and OprM efflux pump proteins, DNA turnover proteins (e.g. elongation factor G, MutL) and cell envelope biosynthesis proteins showed decreased abundance.

Table 5. Examples of proteins altered by ≥ 1.5-fold in hypoxia-adapted cultures relative to normoxia-adapted cultures.

Gene ID[a]	Protein Name	Fold Change	Protein Product Name	Function[b]	UP[c]	MW[d]	SC[e]
PA3402	PA3402	U[f]	hypothetical protein	Antibiotic res	5	36.21	21.10
PA3062	PelC	U[f]	PelC	Biofilm	6	18.68	58.10
PA4456	YrbF	U[f]	probable ATP-binding component of ABC transporter	Antibiotic res	5	29.62	26.00
PA2664	Fhp	6.52	flavohemoprotein	Stress res	7	43.66	27.50
PA3163	Cmk	5.50	cytidylate kinase	Antibiotic res	3	24.64	21.80
PA5346	SadB	3.39	SadB	Biofilm, Motility	5	51.43	19.60
PA3559	PA3559	3.20	Probable nucleotide sugar dehydrogenase	Biofilm	3	51.80	10.60
PA0091	VgrG1	2.60	VgrG1	Antibiotic res	5	72.01	22.90
PA2900	PA2900	2.47	probable outer membrane protein precursor	Antibiotic res	5	29.85	30.90
PA4753	YhbY	2.46	conserved hypothetical protein	Antibiotic res	9	11.66	88.50
PA5549	GlmS	2.34	glucosamine-fructose-6-phosphate aminotransferase	Antibiotic res	30	66.29	49.60
PA0762	AlgU	2.19	Sigma factor AlgU	Biofilm, Motility	7	22.20	52.30
PA0083	TssB1	2.10	TssB1	Antibiotic res, Biofilm	13	18.79	73.80
PA0084	TssC1	2.00	TssC1	Antibiotic res, Biofilm	31	55.77	69.70
PA0141	Ppk2	1.97	GDP-polyphosphate phosphotransferase	Biofilm	20	40.79	72.80
PA4131	PA4131	1.90	probable iron-sulfur protein	Fe acq.	15	63.99	24.80
PA5117	TypA	1.88	regulatory protein TypA	Antibiotic res, Motility	22	66.93	35.70
PA0067	PrlC	1.85	oligopeptidase A	Antibiotic res, Motility	35	76.04	40.80
PA1673	Mrh	1.82	Microoxic hemerythrin	Biofilm	5	17.85	37.30
PA0998	PqsC	1.70	PqsC	Antibiotic res	16	38.24	55.50
PA3011	TopA	1.70	DNA topoisomerase I	Antibiotic res, Stress res, Biofilm	37	97.28	51.80
PA3617	RecA	1.60	RecA protein	Antibiotic res	12	36.88	36.40
PA4748	TpiA	1.59	triosephosphate isomerase	Antibiotic res	10	25.62	44.20
PA3810	HscA	1.57	Heat shock protein HscA	Fe acq.	13	66.45	22.60
PA0588	yeaG	-1.56	conserved hypothetical protein	Antibiotic res, Motility	55	73.72	67.70
PA4595	ArtR	-1.56	ATP-activated Repressor of T3SS regulator	Antibiotic res	39	61.32	61.40
PA5078	OpgG	-1.59	OpgG	Stress res, biofilm	14	59.48	34.90
PA2553	PA2553	-1.60	probable acyl-CoA thiolase	Antibiotic res	43	41.40	90.90
PA3529	TsaA	-1.66	alkylhydroperoxide reductase C	Stress res	28	21.82	84.00
PA4406	LpxC	-1.70	UDP-3-O-acyl-N-acetylglucosamine deacetylase	Antibiotic res	17	33.44	54.50
PA3723	YqjM	-1.72	probable FMN oxidoreductase	Antibiotic res, Stress res	24	40.32	80.40
PA0758	PA0758	-1.74	hypothetical protein	Stress res	10	30.88	47.50
PA2552	AcdB	-1.85	probable acyl-CoA dehydrogenase	Antibiotic res, Motility	14	40.91	46.40
PA2532	Tpx	-1.90	thiol peroxidase	Antibiotic res, Stress res, Biofilm	18	17.23	99.40
PA2071	FusA2	-1.96	elongation factor G	Antibiotic res	31	77.57	70.40
PA4946	MutL	-2.08	DNA mismatch repair protein MutL	Antibiotic res	12	69.75	25.60
PA4876	OsmE	-2.11	osmotically inducible lipoprotein OsmE	Stress res	7	12.52	41.20
PA1097	FleQ	-2.14	Transcriptional regulator FleQ	Biofilm, Motility	23	55.28	58.00
PA5285	SutA	-2.17	SutA	Biofilm	12	11.17	93.30
PA0427	OprM	-2.20	Major intrinsic multiple antibiotic resistance efflux outer membrane protein OprM precursor	Antibiotic res, Stress res	14	52.60	32.60
PA3017	PA3017	-2.32	Universal stress protein	Stress res	11	15.84	84.80
PA2015	LiuA	-2.38	putative isovaleryl-CoA dehydrogenase	Antibiotic res	20	42.21	55.80
PA2639	NuoD	-2.49	NADH dehydrogenase I chain C,D	Antibiotic res	27	68.32	50.30

Gene ID	Protein	Fold	Description	Function	UP	MW	SC
PA1041	**PA1041**	-2.55	Probable outer membrane protein precursor	Biofilm	13	21.75	49.50
PA2009	**HmgA**	-2.55	homogentisate 1,2-dioxygenase	Stress res	17	47.84	50.50
PA3922	**PA3922**	-2.55	conserved hypothetical protein	Antibiotic res, Biofilm	17	51.23	54.30
PA0473	**PsfA**	-2.62	probable glutathione S-transferase	Fe acq.	6	28.33	22.90
PA1561	**Aer**	-2.65	Aerotaxis receptor Aer	Motility	18	57.18	45.70
PA5111	**GloA3**	-2.73	lactoylglutathione lyase	Antibiotic res	3	20.21	31.80
PA4290	**PA4290**	-2.92	probable chemotaxis transducer	Antibiotic res, Motility	16	58.40	37.20
PA4515	**PiuC**	-3.04	conserved hypothetical protein	Antibiotic res, Fe acq.	7	25.62	31.00
PA2867	**PA2867**	-3.07	Probable chemotaxis transducer	Motility	8	51.46	34.90
PA1753	**PA1753**	-3.10	conserved hypothetical protein	Antibiotic res, Stress res	4	18.37	33.30
PA4310	**PctB**	-3.13	Chemotactic transducer PctB	Motility	8	68.08	45.80
PA2008	**FahA**	-3.16	fumarylacetoacetase	Antibiotic res	15	46.34	41.20
PA2764	**PA2764**	-3.27	Hypothetical protein	Motility	7	24.52	33.30
PA0887	**AcsA**	-3.28	acetyl-coenzyme A synthetase	Antibiotic res	45	71.76	64.40
PA3795	**PA3795**	-3.50	probable oxidoreductase	Stress res	7	34.80	31.30
PA2920	**PA2920**	-3.57	probable chemotaxis transducer	Stress res, Motility	10	58.32	23.10
PA4221	**FptA**	-3.76	Fe(III)-pyochelin outer membrane receptor precursor	Fe acq.	25	79.99	42.20
PA4738	**YjbJ**	-4.09	conserved hypothetical protein	Stress res	5	7.61	75.40
PA2638	**NuoB**	-4.11	NADH dehydrogenase I chain B	Antibiotic res	9	25.43	36.40
PA0250	**PA0250**	-4.35	conserved hypothetical protein	Stress res	6	15.67	59.00
PA2652	**PA2652**	-4.67	Methyl-accepting chemotaxis protein	Motility	10	60.34	26.60
PA1092	**FliC**	-4.76	Flagellin type B	Bbiofilm, Motility	6	49.24	15.60
PA0425	**MexA**	-4.83	Resistance-Nodulation-Cell Division (RND) multidrug efflux membrane fusion protein MexA precursor	Antibiotic res	19	40.97	48.60
PA4223	**PchH**	-4.88	probable ATP-binding component of ABC transporter	Antibiotic res, Motility	9	61.34	20.70
PA0744	**PA0744**	-4.89	Probable enoyl-CoA hydratase/isomerase	Mot	14	40.59	38.70
PA3938	**TauA**	-5.11	probable periplasmic taurine-binding protein precursor	Antibiotic res, Stress res	9	35.86	50.10
PA3812	**IscA**	-5.34	probable iron-binding protein IscA	Antibiotic res, Fe acq.	4	11.57	51.40
PA3819	**YcfJ**	-5.67	conserved hypothetical protein	Antibiotic res, Stress res	9	18.97	63.20
PA2363	**HsiJ3**	-5.94	HsiJ3	Antibiotic res	5	47.78	19.20
PA4225	**PchF**	-5.98	pyochelin synthetase	Fe acq.	67	197.08	38.50
PA2788	**PA2788**	-6.13	Probable chemotaxis transducer	Motility	14	56.39	36.90
PA4226	**PchE**	-6.46	dihydroaeruginoic acid synthetase	Fe acq.	47	156.42	34.10
PA4224	**PchG**	-7.13	pyochelin biosynthetic protein PchG	Fe acq.	12	37.70	35.50
PA1065	**PA1065**	-7.68	conserved hypothetical protein	Antibiotic res	3	17.99	24.20
PA3441	**SsuF**	-7.71	probable molybdopterin-binding protein	Stress res	3	7.59	47.90
PA4230	**PchB**	-7.77	salicylate biosynthesis protein PchB	Antibiotic res, Fe acq.	12	11.43	62.40
PA2640	**NuoE**	-8.00	NADH dehydrogenase I chain E	Antibiotic res, Mot	6	18.09	60.20
PA2717	**Cpo**	-8.42	chloroperoxidase precursor	Stress res, Biofilm	9	30.44	59.10
PA2033	**ViuB**	-10.88	hypothetical protein	Fe acq.	6	33.65	28.50
PA5131	**Pgm**	-12.57	phosphoglycerate mutase	Antibiotic res	7	55.60	17.70
PA4825	**MgtA**	-17.09	Mg(2+) transport ATPase, P-type 2	Stress res, Fe acq.	41	100.05	32.80
PA4227	**PchR**	-20.62	transcriptional regulator PchR	Fe acq.	3	32.34	14.90
PA0283	**Sbp**	U[f]	sulfate-binding protein precursor	Stress res	4	36.43	20.20
PA0537	**PA0537**	U[f]	Conserved hypothetical protein	Biofilm	4	22.18	25.20
PA2786	**PA2786**	U[f]	hypothetical protein	Antibiotic res	4	17.54	33.50
PA2927	**PlaB**	U[f]	Hypothetical protein	Biofilm	4	49.48	5.20
PA3123	**RidA**	U[f]	RidA subfamily protein	Biofilm, Motility	4	13.52	34.40
PA3459	**AsnB**	U[f]	probable glutamine amidotransferase	Stress res	8	66.03	17.50
PA3461	**YhfE**	U[f]	conserved hypothetical protein	Stress res	6	43.29	21.60
PA3808	**IscX**	U[f]	conserved hypothetical protein	Fe acq.	4	7.80	62.10

| | | | Increased in abundance (fold) | | 1.5-2 | 2-5 | ≥5 |
| | | | Decreased in abundance (fold) | | 1.5-2 | 2-5 | ≥5 |

Blue shading represents proteins increased in abundance or unique to hypoxia. Dark blue: ≥5-fold increase in abundance; medium blue:
2-5-fold increase in abundance, light blue: 1.5-2-fold increase in abundance. Orange shading represents proteins decreased in
abundance in hypoxia or absent in hypoxia. Dark orange: ≥5-fold decrease in abundance; medium orange: 2-5-fold decrease of
abundance; light orange: 1.5-2-fold decrease in abundance.
[a]Gene ID – *Pseudomonas aeruginosa* PAO1 gene locus ID.
[b]Relevant protein function. Antibiotic res: antibiotic resistance; Stress res: stress resistance; biofilm: Biofilm formation and/or
exopolysaccharide production; Fe acq.: iron acquisition and/or homeostasis; Motility: motility.
[c]UP - Unique peptides: The number of peptide sequences unique to a protein
[d]MW - Molecular weight as determined by Q-Exactive LC-MS using *P. aeruginosa* PAO1 reference (kDa)
[e]SC - % sequence covered by matching peptides for the identified protein

fU – Proteins unique to hypoxia or absent in hypoxia

To confirm whether the alterations of proteins associated with antibiotic resistance affect the

phenotype of each of the individually adapted cultures, antibiotic susceptibility was examined.

Changes in sensitivity to a wide range of antibiotic classes were observed in both populations (Fig 4).

However, both H1 and H2 populations clearly showed more increases in antibiotic resistance than any

other cultures, correlating with the changes in protein abundance. Increased resistance to

fluoroquinolones, macrolides, cephalosporins, carbapenems and penicillins was observed in both H1

and H2 cultures, together with decreased resistance to aztreonam. Interestingly, H3 culture only

showed an increased resistance to aztreonam with no change in susceptibility to any other antibiotic

tested. All strains were resistant to tetracycline.

Hypoxia-adapted cultures show changes in stress response

Many proteins related to stress responses showed changes in abundance in response to long-term

exposure to hypoxia (Table 4), including universal stress proteins (PA3017, PA1753); osmotic stress

response associated proteins (OpgG, OsmE, PA2920, PA4738, PA3461, AsnB); oxidative stress

response associated proteins (Ftp, TsaA, PA3723, Tpx, PA0250, TauA, OprM, MexA, TopA, Hmg, Cpo);

heat-shock stress response associated proteins (TopA, PA0758), sulphate starvation (SsuF, Sbp); cell

wall stress/envelope stress response proteins (PA3819, PA3795) and cadmium stress response (MgtA)

(Table 5). To examine the impact of these alterations on phenotype of each of the individual

populations, droplet stress plates were prepared to test a range of stresses described as hallmarks of

the CF lung environment. No clear pattern resistance to high temperature, osmotic or acidic stress

between the HACs and NACs was observed (Fig 4). Interestingly, increased resistance to oxidative

stress was observed in both H1 and H2 cultures and in the N3 culture, which were also the only

populations where SCVs had been detected on day 28, and where increased biofilm production and

exopolysaccharide production were observed (Fig 4).

Hypoxia-adapted cultures show increased biofilm formation and exopolysaccharide production

Typically anaerobic conditions promote biofilm formation, with greater numbers of live bacteria, increased thickness, and more compact structures (51). Changes in the abundance of 20 proteins associated with biofilm and/or exopolysaccharide production were observed (Table 5). Nine showed increased abundance or were unique to hypoxia, including the PelC exopolysaccharide export protein, as already mentioned. Others included proteins involved in surface attachment (SadB), exopolysaccharide production (PA3559, AlgU), and type VI secretion system (TssB1, TssC1). Proteins that showed decreased abundance are involved in the switch from a planktonic to a biofilm lifestyle (RidA), biofilm formation and structure, and cell attachment (PlaB, PA1041, SutA, FleQ, OpgG), alginate production (Cpo), and flagellin (FliC).

To confirm the impact of these proteome changes on biofilm and exopolysaccharide production, these phenotypes were examined in each individual adapted culture (Fig 5A, Fig 5B). Only H1 and H3 populations showed a significantly increased biofilm formation as compared to the early infection patient isolate (6.1- and 4.3-fold increases respectively) after 24 h, although significant increases in biofilm production were also observed in the late strain (1.6-fold), H1 (3.9-fold) and H2 (1.4-fold) over the early strain at 48 h, and this trend persisted at 72h (Fig 5A). The increased biofilm production of the H3 population was transient and reverted to AMT 0023-30 strain levels by 48 hours. Congo Red staining revealed darker colonies for H1 and H2 populations which were more wrinkly than in the other strains indicating a higher production of exopolysaccharides (Fig 5B). Interestingly, a similar phenotype was observed in culture N3, again suggesting that this phenotype is not dependent on oxygen exposure.

Adaptation to hypoxia affected iron acquisition and homeostasis

Proteomic analysis showed changes in the abundance of 14 proteins related to iron homeostasis and acquisition pathways (Table 5). More than half of the proteins in the pyochelin biosynthesis pathway were decreased in abundance by 6-fold or more (PchB, PchE, PchF, PchG) (Fig 5C), and the master regulator of pyochelin production, PchR, was decreased in abundance by over 20-fold. Additionally, the abundance of several iron transport proteins was decreased, including FptA (iron-bound pyochelin receptor), ViuB (siderophore uptake protein), PiuC (Fe(II)-dependent dioxygenase), MgtA (involved in citrate-bound iron transport into the cell), and PsfA (a stress protein involved in iron acquisition). Moreover, the abundance of proteins involved in iron-sulphur cluster biosynthesis was also affected by hypoxia with HscA increased in abundance nearly 1.6-fold, IscA decreased in abundance over 5.3-fold and IscX undetectable in HACs.

Differences in iron acquisition and consequently siderophore levels of all individual populations were examined and results were consistent with the overall proteome analysis, showing a clear decrease in siderophore production in all of the HACs by 2.8- to- 3.6-fold (p<0.05), but no significant changes in the NACs (Fig 5D). Moreover, a similar change was observed between the early infection isolate and late infection isolate, with a 4-fold decrease in siderophores (p<0.01).

Adaptation to hypoxia induces changes in motility

Over 16 proteins associated with motility showed decreased abundance in the HACs, while only three (flagellar reversal regulator SadB, TypA GTPase and the PrlC oligopeptidase), were increased in abundance (Table 5). The proteins showing decreased abundance included those directly associated with motility (FliC, FleQ, PA0588, PchH, NuoE) and a large group of chemotaxis associated proteins (PA4290, PA2867, PctB, PA2920, PA2652, PA2764, AcdB, PA0744, PA2788) and receptor protein Aer,

suggesting that motility was downregulated in HACs. Notably, many of these proteins are also

associated with biofilm formation and antibiotic resistance.

To elucidate the effect of hypoxia on motility further, the motility of the individual cultures was

assessed. The late patient isolate showed a 2-fold reduction in swarming ($p<0.05$) (Fig 6B) and a

moderate reduction in twitching motility (1.3-fold; $p<0.01$) (Fig 6C), while H1 and H2 showed an

increased ability to swim (1.5 and 1.2-fold, $p<0.05$) compared to the early infection strain (Fig 6A). All

forms of motility were decreased in culture H3 by over 2.4-fold ($p<0.01$) (Fig 6A, 6B, 6C). A similar

pattern was observed in culture N3, with a 1.6-fold decrease in swimming motility ($p<0.05$) and a 1.9-

fold decrease in twitching motility ($p<0.01$) (Fig 6A, 6C). The pattern of swarming motility was clearly

different between NACs and HACs. N1 and N2 populations showed a classic swarming pattern with

clearly distinguishable dendrites despite showing no changes in diameter (Fig 6B). N3 showed a

different pattern, forming colonies with smooth edges. The H1 and H2 colonies on swarm plates were

clearly distinguishable from the N1 and N2 counterparts, with no clear dendrites being formed (Fig

6).

359 Effects of hypoxia-adaptation on *in vivo* virulence

360 The proteome and phenotype characterisation of HACs showed that adaptation to hypoxia altered

361 the abundance of proteins involved in virulence, and the appearances of many virulence associated

362 phenotypes, including SCV formation, stress and antibiotic resistance, and motility. Unexpectedly

363 however, no differences in virulence were observed *in vivo* in *Galleria mellonella* between the early

364 infection isolate and the HACs (Fig. 6D).

Discussion

Summary

Understanding the complex network of pathway interactions and the evolutionary pressures *P. aeruginosa* undergoes in different environments is the key to understanding its biology and the mechanisms behind chronic infection, facilitating the development of effective therapeutics. Hypoxic niches are abundant in the CF lung due to increased mucus thickness, obstructive mucus plugging and increased oxygen consumption by epithelial and immune cells (24,52,53). Studies have shown that microaerobic respiration is the predominant mode of *P. aeruginosa* growth in the CF lung (54,55), with two transcription factors, Anr and Dnr, tightly regulating the response to changes in oxygen tension (56,57), and giving *P. aeruginosa* the ability to respond to changes in oxygen concentration along a continuum (25).

While the impact of short-term hypoxia on *P. aeruginosa* has been previously studied (a few hours, until reaching OD_{600} = 1.0), the impact of hypoxia stress has not been examined over extended periods. We demonstrate that prolonged low oxygen conditions alone can induce stable proteomic and phenotypic changes which are typically associated with chronic infection, and thus potentially facilitate the transition from acute to chronic infection.

Diverse patterns of adaptation to long-term hypoxia

We have observed two distinct patterns of adaptation of *P. aeruginosa* to hypoxic conditions, consistent with existing literature which demonstrates evolution of different phenotypic and genetic characteristics with bacterial populations isolated from the same lung niche, or different parts of the biofilm, enabling bacterial competitiveness (22,58–62). Flynn et. al (2006), using a similar experimental evolution approach as we used in this work, albeit over 90 days, compared three biofilm

cultures and three planktonic cultures of *P. aeruginosa* PA14 and determined that different

phenotypes appeared at different intervals with different abilities to persist not only between

replicate cultures, but even between colonies isolated from the same population (59). Furthermore,

the various colony morphologies observed in this study correlated with temporal variations in c-di-

GMP levels (59), which aligns with the changes in the abundance of proteins associated with the

response to c-di-GMP in HACs.

Two of the HACs in our study (H1 and H2) showed consistent alterations across multiple phenotypes,

while the third population displayed only slight alterations relative to the early strain and were

effectively comparable to the NACs (Fig. 7A). This was confirmed by a phylogenetic tree that we

generated based on our sequencing data (data unpublished), which showed that H1 and H2 are

closely related to each other, but distant from all other populations, and the third HAC was most

closely related to the N3 population (Fig 7B). Previous investigation of the short term hypoxia

response (over 24 hours) revealed that the proteome changes were strain-specific (63), suggesting

that different adaptation patterns can occur as the population diversifies over the longer term. A

further, deeper analysis of the WGS data will allow us to elucidate the genetic background of the

observed differences.

Adaptation of *Pseudomonas aeruginosa* to long-term hypoxia is consistent with changes observed during chronic infection

Interestingly, the adaptation of *P. aeruginosa* to long-term hypoxia mirrors the phenotypic and

molecular changes that have been reported for *P. aeruginosa* late infection isolates as a result of

adaptation to chronic infection (8,18,20,23,35,62,64). These include increased SCV prevalence,

increased biofilm and exopolysaccharide production, increased antibiotic resistance and decreased

siderophore production. The stability of these changes was evident, as all the phenotypes were tested

in normoxic conditions.

The first phenotypic change observed during the experimental adaptation of *P. aeruginosa* to long-

term hypoxia was the emergence of two distinct SCV types. They were more prevalent in the HACs,

and one of the morphotypes was exclusive to hypoxia. Surprisingly, the abundance of c-di-GMP

signalling pathways proteins, Wsp and YfiBNR, typically associated with SCVs (35,65), were unchanged

in the HACs, suggesting an alternative mode of regulation. SCVs are strongly associated with

pathogenicity and persistence among several bacterial pathogens, (66–68) and are prevalent among

chronic *P. aeruginosa* isolates (66,67,69). Specifically, SCVs of *P. aeruginosa* correlate with increased

exopolysaccharide and biofilm production, antimicrobial resistance, stress resistance and host

immune responses (35,69). Their emergence has been linked to redox imbalances which can be

caused by low oxygen conditions (70), although other environmental stressors or co-colonisation with

other species also contribute to their increased prevalence (66,69,71). This confirms directly, for the

first time, that prolonged hypoxia exposure is associated with increased SCV occurrence in *P.*

aeruginosa populations. Although SCVs temporarily occurred under normoxic conditions, they were

not selected as a beneficial trait in the longer term. Identification of the factors that contribute to SCV

emergence will be crucial due to the challenges they pose in clinical settings (66) and recognising the

environmental factors that select this phenotype may help us limit their occurrence.

Chronic infection *P. aeruginosa* CF isolates typically exhibit increased biofilm and exopolysaccharide

production, and reduced motility (18,62,72–74), with biofilm formation relying on hypoxia responses

due to the presence of steep oxygen gradients (26,75,76). Moreover, the awareness of the role of

biofilms in antibiotic treatment efficiency is being acknowledged increasingly (77,78).

The two HACs that showed a high SCV prevalence (H1, H2) also showed increased resistance to a

broad spectrum of antimicrobials together with increased biofilm and exopolysaccharide production.

Surprisingly, no trends in motility relating to hypoxia exposure or biofilm formation were observed,

although some changes in patterns of swarming colonies were noted.

Proteins associated with biofilm formation, exopolysaccharide production and motility were changed

in abundance, including an increase of abundance of SadB, which is required for irreversible cell

attachment, exopolysaccharide production and is associated with decreased swarming motility

(79,80), and the alternative sigma factor AlgU (σ22), which is one of the key factors leading to the

development of chronic *P. aeruginosa* infection in the CF lung and regulates the mucoid phenotype,

and contributing to stress and antibiotic resistance (81–83). A interesting cascade of regulation is

observed, as both SadB and AlgU act through the c-di-GMP and Gac/Rsm signalling pathways (84–

86). Further AlgU inhibits the FleQ master regulator of flagellar motility and exopolysaccharide

secretion, indirectly decreasing the abundance of the FliC flagellin type B (87). Noteworthy, another

modulator of the virulence, biofilm and antibiotic resistance of many Gram-negative bacteria, the

TypA GTPase (88,89), was also increased in abundance in HACs.

Antibiotic resistance is a major hallmark of *P. aeruginosa* infections in CF (90), and short-term hypoxic

and anaerobic conditions have been previously shown to increase *P. aeruginosa* antibiotic resistance

(91–93). We demonstrate that long-term exposure to hypoxia can induce stable changes in resistance

to antibiotics, which didn't revert following subsequent culture in normoxic conditions during

phenotyping or other investigations. Additionally, a wider range proteins associated with antibiotic

resistance were altered in abundance, including an increase in abundance of secretion systems

proteins and ABC transporter systems, outer membrane proteins, the TopA topoisomerase,

peptidoglycan/LPS biosynthesis proteins and quinolone signalling proteins. Disc diffusion assays

confirmed that two of the HACs showed increased resistance to a wide range of antibiotics from

different classes, further supporting the evidence of the extent of the influence of hypoxia pre-

adaptation on drug resistance and the wide range of diverse mechanisms affected.

Bacteria that colonise the CF lung undergo constant stress, including increased osmotic, acid and

oxidative stress (53,94,95). An increased resistance to oxidative stress and increased antibiotic

resistance was observed in two HACs, H1 and H2, which aligns with the available literature data

showing that preexposure to oxidative stress increases the resistance to many antibiotics (96–98).

Considering responses to different stress conditions involve overlapping systems (99), prolonged

hypoxia adaptation is likely to drive the organism's stress resistance in general. Understanding the

specific dynamics of this co-dependency may guide therapy choices in the future (Dawan & Ahn,

2022).

The airways of people with CF are reported to show an increased abundance of available iron

(101,102). *P. aeruginosa* expresses many iron acquisition systems including siderophore production,

haem acquisition systems and complexes importing iron-bound citrate or catechol (Schalk & Perraud,

2023). Importantly, a strong positive correlation between increased sputum iron and chronic *P.*

aeruginosa infections has been shown (104). HACs showed a decrease in the abundance of proteins

involved in pyochelin production (Pch cluster proteins), iron/siderophore uptake proteins/receptors

(FptA and ViuB), and citrate-bound iron transporter MgtA and PsfA, which alters iron uptake under

stress which correlate with the reports of low activity of pyochelin in CF sputum (105). The highly

conserved iron-sulphur cluster group, which includes the IscA and IscX (both of which were decreased

in abundance in HACs) is associated with iron storage and homeostasis, but also respiration, DNA and

RNA repair contributing to adaptation to various environments (106–108). Furthermore, iron uptake

and homeostasis are regulated by the c-di-GMP through the Gac/Rsm cascade (109), in line with other

alterations observed and underlines the role of hypoxia leading to changes in iron acquisition on

chronic infection development, which led to recognising them as therapy targets for *P. aeruginosa*

treatment (102,110).

It has been shown that exposure to hypoxia attenuated the virulence of isolates from acute, but not

chronic infections (111) and decreased their interactions with host cells, demonstrating the

importance of this environmental pressure. Moreover, despite the lower bacterial concentrations *in*

vivo in hypoxia, the mortality of mice was accelerated (112,113). Previous reports demonstrated that

decreased pyocyanin production is a characteristic of chronic infection (114). However, no changes in

the abundance of proteins associated with this phenotype were detected in HACs, and its production

was decreased in both HACs and NACs, suggesting that even though hypoxia may contribute to

reduced pyocyanin production, it is not dependent on low oxygen conditions. Additionally, we haven't

observed any changes in the *Galleria mellonella* infection model related to hypoxia exposure.

A recent study defined 224 genes of *P. aeruginosa* as pathoadaptive, and determined that the

expression of certain genes was associated with the host preference towards people with CF (116).

Nine proteins increased in abundance in HACs were encoded by pathoadaptive genes identified by

Weimann et al., including AlgU, FleQ, NarL, FptA, MexA, MutL, OprM, YrbF (previously discussed) and

additionally the RmlB rhamnose/lipopolysaccharide biosynthetic protein (117,118). Moreover, our

proteomics analysis showed changes in the abundance of 21 proteins encoded by genes whose

expression was positively associated with CF host preference and 38 that were associated with non-

CF infections (116), suggesting long-term hypoxia may play a role in *P. aeruginosa* infection both in CF

and non-CF cohorts. Although many of the phenotypes that were altered by extended culturing in

hypoxia can contribute to the pathogenicity of *P. aeruginosa*, the virulence of the HADs was

unchanged in the *G. mellonela* infection model. However, this is an acute infection model, and

consequently is not an optimal model to assess the impact of long-term hypoxia adaptation on

chronic infection.

Differences between long-term adaptation and short-term response to hypoxia

The influence of various environmental pressures on the metabolism of pathogenic bacteria is

typically investigated by exposing bacteria to that factor for 24 h or less, which captures early

responses, but it does not evaluate the prolonged environmental pressures experienced during

infection. No studies on the effects of long-term exposure to hypoxia have been investigated in *P.*

aeruginosa.

Our findings highlight that the bacterial strategies employed to overcome short-term and long-term

stress exposure are quite distinct. Short-term adaptation facilitates quick, immediate responses

including the stringent response and/or metabolic adjustments, whereas prolonged exposure to

specific conditions is more likely to foster adaptation through the acquisition and accumulation of

genetic changes (119–122). Specific examples include the osmotic stress response of *Pseudomonas*

putida, which initially involved upregulation of osmoprotectant production, while long-term exposure

led to adjustments in the outer membrane's composition to reduce permeability (99). Furthermore,

Helicobacter pylori responded to acidic stress by rapidly expressing urease to neutralise the acid, but

over time altered its membrane structure and proton-pumping systems (123).

To date, only one study has specifically investigated the global proteome changes in response to short-

term hypoxia (63). In contrast, to the that study, which examined the *P. aeruginosa* PAO1 reference

strain and two adult cystic fibrosis strains (PASS2, PASS3), we used an early-infection paediatric isolate

(AMT 0023-30) with no prior lung adaptation, providing a unique perspective (29,30,32). Only three

proteins among the 101 proteins that changed in abundance both after long-term and short-term

hypoxia exposure were consistently altered across all four strains examined in both studies,

highlighting key proteins for hypoxia adaptation: the Mhr microaerophilic hemerythrin and the

PA0141 GDP-polyphosphate phosphotransferase were increased in abundance in hypoxia, while the

MaiA maleylacetoacetate isomerase decreased in abundance. Their role(s) in the pathogenicity of *P.*

aeruginosa suggests they may be promising candidates for further studies in relation to chronic

infection and its treatment (124–127). The abundance of 107 proteins was altered only in long-term

hypoxia cultures, but not after short-term exposure. Of the 31 proteins uniquely increased in

abundance during long-term adaptation, notable examples previously connected to chronic infection

adaptation and/or virulence include the AlgU sigma factor, T6SS-associated proteins, transporters,

and LPS associated proteins. Contrary to short-term hypoxia exposure which led to changes in the

abundance of cytochromes and denitrification pathways, no changes in the abundance of these

groups were detected in the long-term HACs. Additionally, no increases in the abundance of proteins

involved in oxidative stress resistance were observed, unlike the reported short-term hypoxia

response (63). This suggests that while oxidative stress responses are crucial for initial response to

low oxygen conditions, they are unlikely to play a significant role in the adaptation to prolonged

hypoxia.

Hypoxia-induced redox imbalance changes c-di-GMP levels

Hypoxia causes a redox imbalance in the bacterial cell which needs to be mitigated for the

maintenance of function and viability, and this process is regulated by c-di-GMP signalling. Previous

studies reported the role of c-di-GMP in regulating low oxygen response and stress resistance in *P.*

aeruginosa and highlight its function in the metabolic switch in response to environmental stimuli

including the switch from a planktonic to a biofilm lifestyle (85,128,129)**.** Moreover, the changes in c-

di-GMP levels were associated with diversification of populations and variability in phenotypes

expressed after adaptation to one condition (59). As we identified several proteins associated with

the GacAS system, it is tempting to speculate that hypoxia modulates the second messenger c-di-

GMP and consequently alters multiple c-di-GMP-regulated phenotypes, including the increased

biofilm formation and exopolysaccharide production, altered motility, and increased antibiotic

resistance we observe. For example, FleQ is a c-di-GMP receptor regulating biofilm formation,

exopolysaccharide production, and type VI secretion system expression (130–132). Considering the

above, we speculate that hypoxia adaptation modulates the swich from acute to chronic infection

through the c-di-GMP second messenger.

Common adaptation mechanisms of opportunistic pathogens to hypoxia

Several oxygen-sensing mechanisms, homologous genes in hypoxia regulons and several phenotypes

altered in chronic infection are shared across other opportunistic bacteria causing pulmonary

infections (Carey et al., *under review).* This suggests that adaptation to prolonged hypoxia could be a

shared mechanism between many low oxygen and/or microaerophilic habitats and are not limited to

human infections. These environments are increasingly abundant in water and soil, where

opportunistic bacteria reservoirs exist and play significant roles in shaping their physiology, with

increasingly contributing to their increased prevalence (133–137). As many opportunistic bacterial

pathogens causing CF infections reside in these niches, common adaptations are not unexpected.

Proteome analysis of HACs showed a decrease in abundance of ten proteins encoded on the Anr

regulon (56) that are involved in energy and amino acid metabolism (Qor, Hpd, GapN, PepN), or

associated with pathogenesis (LpxC – lipopolysaccharide production; ArtR – T3SS activity regulation;

Aer, PA2867 – chemotaxis/motility). Meanwhile, two proteins involved in biofilm production, Ppk2

and Mrh were increased in abundance after long-term hypoxia adaptation. Anr was reported to show

high activity in chronic *P. aeruginosa* infection isolates (25,26). It is noteworthy that its regulon shares

homologous genes with the DosR regulon of *Mycobacterium abscessus* and Lxa locus of *Burkholderia*

cenocepacia (Carey et al., *under review*), highlighting that aspects of the hypoxia response may be

shared across several opportunistic CF lung infections more generally. Targeting shared mechanisms

of adaptation to chronic infection could present a new opportunity for the treatment of multiple

pathogens simultaneously.

Impact

Recognizing the adaptive patterns of microorganisms is critical to devising targeted treatment

approaches. Crucially, this study demonstrates that alterations in phenotype after long-term exposure

to hypoxic conditions shows several similarities with those observed during adaptation to chronic

infection. Thus, further development of our understanding of the specific mechanisms by which *P.*

aeruginosa adapts to hypoxia in chronic infection and the interplay with other factors present in the

lung environment will potentially enable us to prevent or treat the switch from acute to chronic

infection. This could inevitably limit antibiotic use, severity of the disease and ultimately, decrease

the socioeconomic impact of infections caused by *P. aeruginosa*. Furthermore, the first drug to alter

the human response to hypoxia was recently approved for use by the FDA (138), thus it is tempting

to speculate that the response of bacteria to low oxygen pressures could be targeted to prevent a

switch from acute to chronic infection.

To summarise, the changes in proteome and phenotypes (increased biofilm formation, increased

antibiotic resistance, decreased siderophore production, increased SCV prevalence) further deepens

the knowledge on how hypoxia leads to adaptation of *P. aeruginosa*, shedding light on how prolonged

low oxygen exposure can impact on adaptation to chronic infection and providing insights into the

differences between short-term responses and long-term adaptation. Overall, we confirm that

hypoxia contributes to multiple phenotypic changes that are prevalent in chronic infection, opening

a path towards designing therapies targeting these.

Materials and methods

Bacterial strains

Pseudomonas aeruginosa AMT 0023-30 early infection clinical isolate (E) and AMT 0023-31 late infection clinical isolate (L) are two well characterised strains in the International *P. aeruginosa* panel (30). Hypoxia- and normoxia-adapted *P. aeruginosa* strains were obtained in this study as described below and are referred to as H1, H2, H3, N1, N2, and N3 (Table 1). If not stated otherwise, bacteria were routinely grown in LB broth (Sigma-Aldrich, St. Louis, MO, USA) at 37°C with orbital agitation (200 rpm) or plated on LB agar (Sigma-Aldrich, St. Louis, MO, USA) and grown at 37°C.

Experimental evolution

Overnight cultures of *P. aeruginosa* strain AMT 0023-30 (E) were inoculated into 30 ml LB media to OD_{600} ~0.1. Prior to use, the media was equilibrated for 18 h at 37°C in an O_2 Control InVitro Glove Box (PHGB model, Coy Laboratory Products, MI, United States) in 6% O_2, 5% CO_2 or in a Steri-Cycle CO_2 Incubator (model 371, Thermo Scientific, Waltham, MA, USA) at 21% O_2, 5% CO_2. Three flasks were inoculated for each of the conditions being examined. Additionally, aliquots (2 ml) of the overnight *P. aeruginosa* cultures were retained as a controls for proteomic and transcriptomic analysis. All flasks were incubated for 24h with agitation (200 rpm, CO_2 resistant orbital shaker, model 15351105, Thermo Fisher Scientific, Waltham, MA, USA) in hypoxic and normoxic conditions. Daily aliquots (2 ml) were removed for proteomic analysis from each of the flasks and also additional aliquots for OD_{600} measurements and plating. Glycerol stocks (50%, LB (v/v)) were prepared for subsequent phenotype analysis and frozen immediately to ensure minimal oxygen exposure. A volume of 5% of the bacterial culture was also transferred into 30 ml of pre-adapted media each day and this procedure was continued for up to 28 days to determine long-term adaptation patterns. To ensure stability of the hypoxic conditions, all of the samples from H1, H2 and H3 were taken inside

621 the hypoxia chamber. To ensure that there were no other environmental differences between the test

622 conditions other than oxygen levels, the same batch of media was used for both normoxia-adapting

623 and hypoxia-adapting flasks and shaking.

Whole proteome analysis

Samples (2 ml) were collected throughout the experimental evolution for proteome analysis. Bacterial

cells were pelleted at 14000-16000 g for 2 min and stored at -80°C until further analysis. Whole cell

lysates were extracted from each of the hypoxia- and normoxia exposed cultures by resuspending in

1 ml of ice-cold lysis buffer containing 40mM Tris-HCl (pH 7.8) and 1x cOmplete™, EDTA-free protease

inhibitor cocktail (Roche, Basel, Switzerland). Cells were disrupted by a sonication with a sonic

dismembrator and microtip 6mm probe (model 505; Fisherbrand, Thermo Fisher Scientific, Waltham,

MA, USA) (set with 20% output power using eight 10 s bursts on ice). Cell debris and any remaining

intact cells were pelleted using centrifugation (12000 g for 30 min). Supernatants were transferred

into fresh tubes and the proteins were precipitated with four volumes of methanol and incubated in

-80°C overnight. The samples were centrifuged at 12000 g for 30 min at 4°C, supernatants

were removed, and the protein pellets were resuspended in 200 µl of resuspension buffer containing

8 M urea, 50 mM Tris-HCL (pH 8.0), before 1 M 1,4-dithiothreitol (DTT) was added to each supernatant

(10 µl/ml lysate) and incubated at 56°C for 30 min, followed by the addition of 1 M iodoacetamide

(55 µl/ml lysate) which was incubated in darkness at room temperature for 20 min. Lysates were

dialyzed in SnakeSkin™ tubing (Thermo Fisher Scientific, Waltham, MA, USA) with a cut-off of 3.5 kDa

against 100 mM ammonium bicarbonate overnight, with stirring at 4°C, followed by at least 4 more

hours incubation with fresh volumes of ammonium bicarbonate. Trypsin (400 ng/µl) (Sigma-Aldrich,

St. Louis, MO, USA) was added to the dialyzed protein (5 µl /100 µl protein) and tubes were incubated

overnight at 37°C. Aliquots from each sample were placed in fresh tubes and samples were dried in a

vacufuge concentrator system (model 5301; Eppendorf, Hamburg, Germany) at 65°C and

resuspended in 20 µl of ZipTips resuspension buffer containing 0.5% trifluoroacetic acid (TFA) in MilliQ

water. ZipTips (Merc Millipore, Burlington, MA, USA) were used for peptide purification as per the

manufacturers' instructions. Nanodrop spectrophotometer DS-11+ (DeNovix, Wilmington, DE, USA)

was used for peptide concentration measurements. The samples were analysed on a Bruker TimsTOF

Pro mass spectrometer connected to a Evosep One chromatography system. Peptides were separated

on an 8 cm analytical C18 column (Evosep, 3 µm beads, 100 µm ID) using the pre-set 33 samples per

day gradient on the Evosep one. A trapped ion mobility (TIMS) analyser was synchronized with a

quadrupole mass filter to enable the highly efficient PASEF (Parallel Accumulation Serial

Fragmentation acquisition) procedure with acquisition rates of 100 Hz. The accumulation and ramp

times for the TIMS were both set to 100 ms., with an ion mobility (1/k0) range from 0.62 to 1.46

Vs/cm. Spectra were recorded in the mass range from 100 to 1,700 m/z. The precursor (MS) Intensity

Threshold was set to 2,500 and the precursor Target Intensity set to 20,000. Each PASEF cycle

consisted of one MS ramp for precursor detection followed by 10 PASEF MS/MS ramps, with a total

cycle time of 1.16 s.

Protein identification and label-free quantitative (LFQ) analysis was conducted using MaxQuant (v

2.4.2.0, https://maxquant.org/) by searching against the reference strain *P. aeruginosa* PAO1 (ATCC

15692). For protein identification, the following search parameters were used: trypsin was the

digesting enzyme with up to two missed cleavages allowed for; oxidation on methionine and

acetylation on the N terminus were selected as variable modifications with a fixed modification of

carbamidylation on cysteine; Label Free Quantitation (LFQ) and Match Between Runs (MBR) options

were selected. FDR for protein and peptide were set at 1%.

One sample from each 28 day culture was analysed, with one technical replicate for a randomly

chosen sample and LFQ intensities were compared. All missing LFQ intensities were imputed with a

fixed value of 1. The mean was calculated separately for the hypoxia and normoxia samples, followed

by fold change (FC) calculation. T-test statistical analysis was performed to identify proteins with a

significantly changed abundance (p-value ≤ 0.05). Proteins with a -1.5 ≤ FC ≥ 1.5 were considered.

Antibiotic resistance

Overnight bacterial cultures were diluted in 10 ml of fresh LB to OD_{600} equal to 0.1. Bacteria were

spread evenly on Mueller–Hinton agar plates (Neogen, Lansing, MI, USA) using cotton swabs, in three

directions, to give a homogenous lawn. Antibiotic disks (Oxoid, Hampshire, UK) were placed on the

surface of the agar and the plates were incubated overnight in normoxic conditions at 37ºC. Growth

inhibition zone diameters were measured in at least two points, and the mean was used for statistical

analysis.

Motility

Motility assays were performed as describe by Rashid & Kornberg (2000). Swimming motility was

assessed by inoculating the surface of a 0.3 % (w/v) LB agar plate with overnight culture using a sterile

toothpick. Swimming and swarming plates were incubated overnight in normoxic conditions at 37ºC

for ~18 h. For swarming motility 0.8% nutrient broth (Sigma-Aldrich, St. Louis, MO, USA), 0.5%

dextrose, 0.5% agar plates were inoculated identically as for swimming motility assessment. The

diameter of bacterial growth zones was measured in at least two directions, and the mean was used

for statistical analysis. Twitching motility was assessed by inoculating 1.5 % (w/v) LB agar plate by

piercing through the agar. Plates were incubated overnight in normoxic conditions at 37ºC for ~18 h,

and the agar was removed. Bacteria on the surface of the plate were stained with 1% (w/v) crystal

violet for 20 min. The plates were washed with water until all excess dye was removed and dried at

room temperature. The diameter of the growth zone was measured in at least two points, and the

mean was used for statistical analysis.

Biofilm production

Overnight cultures of PA strains were diluted in fresh LB media to OD_{600} of ~0.1, and 100 µl of each diluted culture was transferred into 6 wells of a 96-well plate. The plates were incubated statically in normoxic conditions at 37ºC for 24, 48 or 72 h and OD_{600} was measured with the Synergy H1 microplate reader (Mason Technology, Dublin, Ireland). The medium with planktonic bacteria was removed, and the biofilm fixed by incubation with 200ul of methanol for 10 min. The wells were washed three times with phosphate buffered saline (PBS) (Sigma-Aldrich, St. Louis, MO, USA), followed by the addition of 200 µl of 0.1% crystal violet solution and 30 min incubation at room temperature. The solution was removed, and the wells were washed three times with water and once with PBS. The plates were dried for at least 30 min at room temperature and 150 µl of 96% ethanol was added to dissolve the bound stain. After 10 min of incubation at room temperature, the solution was mixed by pipetting, OD_{590} was determined, and the OD_{590}/OD_{600} ratio was calculated.

Exopolysaccharide production - Congo Red agar assay

Overnight cultures of PA strains were pelleted and washed with 1 ml of PBS. The pellets were resuspended in PBS to OD_{600} of ~0.1 and 1 µl of the suspension was plated on a Congo Red agar plate (10 g/L Tryptone, 10 g/ L Bactoagar, 40mg/L Congo Red, 20mg/L Coomassie Brilliant Blue). The plates were incubated for 48 h in normoxic conditions and colony morphology was assessed.

Stress sensitivity assays

Overnight cultures of PA strains were diluted to a CFU/ml of 10^8 and a range of serial dilutions was prepared. Suspensions (1ul) were plated on one of the following: LB agar control plates, LB agar supplemented with 0.5 M NaCl (osmotic stress), LB agar supplemented with 1mM H_2O_2 (oxidative stress), or LB agar, pH 5 (acidic stress). The plates were incubated in normoxic conditions at 37ºC for

713   24 or 48 h. Temperature stress was determined by incubation of an LB agar plate at 45°C for 24 h.

714   Change of sensitivity was determined by determining the lowest dilution at which bacterial growth

715   was observed.

Galleria mellonella infection model

Virulence was determined as previously described (32). Wax moth larvae (*Galleria mellonella*) (The

KammieShop, Waalwijk, Denmark) were equilibrated in normoxic conditions at 15°C for at least 7 days

after arrival and stored at the same conditions for up to 4 weeks, during which larvae with a weight

range of 0.2-0.4 g were selected for experiments. Overnight bacterial cultures in LB were diluted in

fresh media to OD$_{600}$ of ~0.1 and grown for 2-3 h. Cultures were centrifuged, and bacterial cells were

washed with PBS. The bacteria were diluted to 10^6 CFU/ml and serially diluted to 10^{-6}. Each dilution

was injected (20 µl) into the hindmost left proleg of healthy larvae (10 per group) and the same

volume of PBS was injected into one group as a control. Injected larvae were incubated at 37°C for

72 h and the number of cells causing an LD$_{50}$ after 24 h was determined through bioburden

assessment by plating bacteria onto LB agar and counting colonies after 24 h.

Pyocyanin production quantification

P. aeruginosa strains were inoculated to OD$_{600}$ in fresh PB media (20g/L Bacto-Peptone, 1.4g/L MgCl$_2$,

10g/L K$_2$SO$_4$) and cultured with aeration in normoxic conditions at 37°C to maximize pyocyanin

production. The bacterial cultures were pelleted (10 min at 12300 g) and two 7.5-ml replicates were

withdrawn from each of the 18 h stationary phase cultures. Pyocyanin was extracted with 4.5 ml of

chloroform and a total of 1.5 ml of 0.2 M HCl was added to each extract causing colour change from

blue to pink. The absorbance of this solution was measured at 520 nm, and the obtained values were

converted to pyocyanin content following Essar et al. (1990) (139). The experiment was repeated at

least two times.

Automatic growth curves

To assess the influence of the inoculation volume on the growth patterns of *P. aeruginosa* AMT 0023-30, 24h stationary phase cultures were inoculated into fresh LB media in a way they constituted 1%, 5% or 10% of the new culture. Three 200 µl technical replicates of the cultures were transferred into a 96-well plate and incubated in normoxic conditions at 37°C with agitation.

Growth curves

To assess the differences in the numbers of generations in hypoxic conditions and normoxic conditions of the AMT0063-30 early infection strain overnight cultures were prepared. The following day they were inoculated in triplicate into fresh LB media and incubated with aeration at 37°C in the O_2 Control InVitro Glove Box in 6% O_2, 5% CO_2 or in 21% O_2, 5% CO_2 or in the Steri-Cycle CO_2 Incubator at 21% O_2, 5% CO_2. Every hour a sample was taken from each culture for OD_{600} measurements and plating on LB agar. The agar plates were incubated overnight at 37°C and the colony forming units were calculated and OD_{600} to CFU/ml was plotted, a trend line was determined and R^2 calculated. The doubling time was calculated for the logarithmic phase of growth using the following formula: Doubling time = (Duration (min)*ln(2))/[ln(final cell concentration/initial cell concentration)].

Acknowledgements

JD, CJC are supported by an SFI Frontiers of the Future award (20/FFP-P/8717) to SMcC. ND was supported by the Irish Research Council, GOIPG/2021/1443 IRC Award. The authors are grateful to Dr David Gomez-Matallanas for his advice in proteomic analysis and for the support of the staff and facilities in the UCD Conway proteomics core.

References

1. Antimicrobial resistance collaborators. Antimicrobial Resistance Collaborators: Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. Lancet. 2022 Feb 12;399(10325):629–55.

2. O'Neill J. Tackling drug-resistant infections globally: final report and recommendations [Internet]. 2016 May [cited 2024 Oct 4]. Available from: https://wellcomecollection.org/works/thvwsuba

3. Blanchard AC, Waters VJ. Opportunistic Pathogens in Cystic Fibrosis: Epidemiology and Pathogenesis of Lung Infection. Journal of the Pediatric Infectious Diseases Society. 2022 Sep 1;11(Supplement_2):S3–12.

4. Sanz-García F, Gil-Gil T, Laborda P, Ochoa-Sánchez LE, Martínez JL, Hernando-Amado S. Coming from the Wild: Multidrug Resistant Opportunistic Pathogens Presenting a Primary, Not Human-Linked, Environmental Habitat. International Journal of Molecular Sciences. 2021 Jan;22(15):8080.

5. Serra-Burriel M, Keys M, Campillo-Artero C, Agodi A, Barchitta M, Gikas A, et al. Impact of multi-drug resistant bacteria on economic and clinical outcomes of healthcare-associated infections in adults: Systematic review and meta-analysis. PLoS One. 2020;15(1):e0227139.

6. Reynolds D, Kollef M. The Epidemiology and Pathogenesis and Treatment of Pseudomonas aeruginosa Infections: An Update. Drugs. 2021 Dec 1;81(18):2117–31.

7. Li Y, Roberts JA, Walker MM, Aslan AT, Harris PNA, Sime FB. The global epidemiology of ventilator-associated pneumonia caused by multi-drug resistant *Pseudomonas aeruginosa*: A systematic review and meta-analysis. International Journal of Infectious Diseases. 2024 Feb 1;139:78–85.

8. Rossi E, La Rosa R, Bartell JA, Marvig RL, Haagensen JAJ, Sommer LM, et al. Pseudomonas aeruginosa adaptation and evolution in patients with cystic fibrosis. Nat Rev Microbiol. 2021 May;19(5):331–42.

9. Smith WD, Bardin E, Cameron L, Edmondson CL, Farrant KV, Martin I, et al. Current and future therapies for Pseudomonas aeruginosa infection in patients with cystic fibrosis. FEMS Microbiology Letters. 2017 Aug 1;364(14):fnx121.

10. Botelho J, Grosso F, Peixe L. Antibiotic resistance in *Pseudomonas aeruginosa* – Mechanisms, epidemiology and evolution. Drug Resistance Updates. 2019 May 1;44:100640.

11. Bakkeren E, Diard M, Hardt WD. Evolutionary causes and consequences of bacterial antibiotic persistence. Nat Rev Microbiol. 2020 Sep;18(9):479–90.

12. Sett A, Dubey V, Bhowmik S, Pathania R. Decoding Bacterial Persistence: Mechanisms and Strategies for Effective Eradication. ACS Infect Dis [Internet]. 2024 Jun 28 [cited 2024 Jul 1]; Available from: https://doi.org/10.1021/acsinfecdis.4c00270

793   13.   ECDC. Healthcare-associated infections acquired in intensive care units - Annual Epidemiological
794   Report for 2016 [Internet]. Stockholm; 2018 [cited 2024 May 15]. Available from:
795   https://www.ecdc.europa.eu/en/publications-data/healthcare-associated-infections-intensive-
796   care-units-2019

797   14.   ECDC. Healthcare-associated infections acquired in intensive care units - Annual Epidemiological
798   Report for 2019 [Internet]. Stockholm; 2023 [cited 2024 May 15]. Available from:
799   https://www.ecdc.europa.eu/en/publications-data/healthcare-associated-infections-intensive-
800   care-units-2019

801   15.   Tacconelli E, Carrara E, Savoldi A, Harbarth S, Mendelson M, Monnet DL, et al. Discovery,
802   research, and development of new antibiotics: the WHO priority list of antibiotic-resistant
803   bacteria and tuberculosis. The Lancet Infectious Diseases. 2018 Mar 1;18(3):318–27.

804   16.   WHO. WHO bacterial priority pathogens list, 2024: Bacterial pathogens of public health
805   importance to guide research, development and strategies to prevent and control antimicrobial
806   resistance [Internet]. 2024 [cited 2024 Dec 5]. Available from:
807   https://www.who.int/publications/i/item/9789240093461

808   17.   Cullen L, McClean S. Bacterial Adaptation during Chronic Respiratory Infections. Pathogens. 2015
809   Mar;4(1):66–89.

810   18.   Jurado-Martín I, Sainz-Mejías M, McClean S. Pseudomonas aeruginosa: An Audacious Pathogen
811   with an Adaptable Arsenal of Virulence Factors. Int J Mol Sci. 2021 Mar 18;22(6):3128.

812   19.   Moradali MF, Ghods S, Rehm BHA. Pseudomonas aeruginosa Lifestyle: A Paradigm for
813   Adaptation, Survival, and Persistence. Front Cell Infect Microbiol [Internet]. 2017 Feb 15 [cited
814   2024 Jul 1];7. Available from: https://www.frontiersin.org/journals/cellular-and-infection-
815   microbiology/articles/10.3389/fcimb.2017.00039/full

816   20.   Camus L, Vandenesch F, Moreau K. From genotype to phenotype: adaptations of Pseudomonas
817   aeruginosa to the cystic fibrosis environment. Microbial Genomics. 2021;7(3):000513.

818   21.   Chadha J, Harjai K, Chhibber S. Revisiting the virulence hallmarks of Pseudomonas aeruginosa: a
819   chronicle through the perspective of quorum sensing. Environmental Microbiology.
820   2022;24(6):2630–56.

821   22.   Cramer N, Klockgether J, Tümmler B. Microevolution of Pseudomonas aeruginosa in the airways
822   of people with cystic fibrosis. Current Opinion in Immunology. 2023 Aug 1;83:102328.

823   23.   Rosales-Reyes R, Vargas-Roldán SY, Lezana-Fernández JL, Santos-Preciado JI. *Pseudomonas
824   Aeruginosa*: Genetic Adaptation, A Strategy for its Persistence in Cystic Fibrosis. Archives of
825   Medical Research. 2021 May 1;52(4):357–61.

826   24.   Worlitzsch D, Tarran R, Ulrich M, Schwab U, Cekici A, Meyer KC, et al. Effects of reduced mucus
827   oxygen concentration in airway Pseudomonas infections of cystic fibrosis patients. J Clin Invest.
828   2002 Feb;109(3):317–25.

829   25.   Alvarez-Ortega C, Harwood CS. Responses of Pseudomonas aeruginosa to low oxygen indicate
830   that growth in the cystic fibrosis lung is by aerobic respiration. Molecular Microbiology.
831   2007;65(1):153–65.

832   26.   Hassett DJ, Sutton MD, Schurr MJ, Herr AB, Caldwell CC, Matu JO. </i>Pseudomonas
833   aeruginosa</i> hypoxic or anaerobic biofilm infections within cystic fibrosis airways. Trends in
834   Microbiology. 2009 Mar 1;17(3):130–8.

835   27.   Cullen L, O'Connor A, McCormack S, Owens RA, Holt GS, Collins C, et al. The involvement of the
836   low-oxygen-activated locus of Burkholderia cenocepacia in adaptation during cystic fibrosis
837   infection. Sci Rep. 2018 Sep 6;8(1):13386.

838   28.   O'Connor A, Jurado-Martín I, Mysior MM, Manzira AL, Drabinska J, Simpson JC, et al. A universal
839   stress protein upregulated by hypoxia has a role in Burkholderia cenocepacia intramacrophage
840   survival: Implications for chronic infection in cystic fibrosis. Microbiologyopen. 2022 Dec
841   9;12(1):e1311.

842   29.   Mulcahy LR, Burns JL, Lory S, Lewis K. Emergence of Pseudomonas aeruginosa Strains Producing
843   High Levels of Persister Cells in Patients with Cystic Fibrosis. Journal of Bacteriology. 2010 Oct
844   8;192(23):6191.

845   30.   De Soyza A, Hall AJ, Mahenthiralingam E, Drevinek P, Kaca W, Drulis-Kawa Z, et al. Developing an
846   international Pseudomonas aeruginosa reference panel. Microbiologyopen. 2013
847   Dec;2(6):1010–23.

848   31.   Freschi L, Bertelli C, Jeukens J, Moore MP, Kukavica-Ibrulj I, Emond-Rheault JG, et al. Genomic
849   characterisation of an international Pseudomonas aeruginosa reference panel indicates that the
850   two major groups draw upon distinct mobile gene pools. FEMS Microbiol Lett. 2018 Jul
851   1;365(14).

852   32.   Cullen L, Weiser R, Olszak T, Maldonado RF, Moreira AS, Slachmuylders L, et al. Phenotypic
853   characterization of an international Pseudomonas aeruginosa reference panel: strains of cystic
854   fibrosis (CF) origin show less in vivo virulence than non-CF strains. Microbiology.
855   2015;161(10):1961–77.

856   33.   Aanaes K, Rickelt LF, Johansen HK, von Buchwald C, Pressler T, Høiby N, et al. Decreased mucosal
857   oxygen tension in the maxillary sinuses in patients with cystic fibrosis. J Cyst Fibros. 2011
858   Mar;10(2):114–20.

859   34.   Walkiewicz A, Brzezińska M, Bieganowski A. Methanotrophs are favored under hypoxia in
860   ammonium-fertilized soils. Biol Fertil Soils. 2018 Oct 1;54(7):861–70.

861   35.   Xu A, Zhang X, Wang T, Xin F, Ma LZ, Zhou J, et al. Rugose small colony variant and its hyper-
862   biofilm in Pseudomonas aeruginosa: Adaption, evolution, and biotechnological potential.
863   Biotechnology Advances. 2021 Dec 1;53:107862.

864   36.   Marmont LS, Rich JD, Whitney JC, Whitfield GB, Almblad H, Robinson H, et al. Oligomeric
865   lipoprotein PelC guides Pel polysaccharide export across the outer membrane of Pseudomonas
866   aeruginosa. Proceedings of the National Academy of Sciences. 2017 Mar 14;114(11):2892–7.

867   37.   Poudyal B, Sauer K. The ABC of Biofilm Drug Tolerance: the MerR-Like Regulator BrlR Is an
868   Activator of ABC Transport Systems, with PA1874-77 Contributing to the Tolerance of
869   Pseudomonas aeruginosa Biofilms to Tobramycin. Antimicrobial Agents and Chemotherapy.
870   2018 Jan 25;62(2):10.1128/aac.01981-17.

38. Avila-Cobian LF, De Benedetti S, Kim C, Feltzer R, Champion MM, Fisher JF, et al. In vitro studies
of the protein-interaction network of cell-wall lytic transglycosylase RlpA of Pseudomonas
aeruginosa. Commun Biol. 2022 Nov 30;5(1):1–16.

39. Benkert B, Quäck N, Schreiber K, Jaensch L, Jahn D, Schobert M. Nitrate-responsive NarX-NarL
represses arginine-mediated induction of the Pseudomonas aeruginosa arginine fermentation
arcDABC operon. Microbiology. 2008;154(10):3053–60.

40. Schreiber K, Krieger R, Benkert B, Eschbach M, Arai H, Schobert M, et al. The Anaerobic
Regulatory Network Required for Pseudomonas aeruginosa Nitrate Respiration. Journal of
Bacteriology. 2007 Jun;189(11):4310–4.

41. Raba DA, Yuan M, Fang X, Menzer WM, Xie B, Liang P, et al. Role of Subunit D in Ubiquinone-
Binding Site of Vibrio cholerae NQR: Pocket Flexibility and Inhibitor Resistance. ACS Omega. 2019
Nov 19;4(21):19324–31.

42. Kotecka K, Kawalek A, Kobylecki K, Bartosik AA. The MarR-Type Regulator PA3458 Is Involved in
Osmoadaptation Control in Pseudomonas aeruginosa. International Journal of Molecular
Sciences. 2021 Jan;22(8):3982.

43. Aspedon A, Palmer K, Whiteley M. Microarray Analysis of the Osmotic Stress Response in
Pseudomonas aeruginosa. J Bacteriol. 2006 Apr;188(7):2721–5.

44. Wood LF, Ohman DE. Identification of Genes in the σ22 Regulon of Pseudomonas aeruginosa
Required for Cell Envelope Homeostasis in Either the Planktonic or the Sessile Mode of Growth.
mBio. 2012 May 15;3(3):10.1128/mbio.00094-12.

45. D'Souza-Ault MR, Smith LT, Smith GM. Roles of N-acetylglutaminylglutamine amide and glycine
betaine in adaptation of Pseudomonas aeruginosa to osmotic stress. Appl Environ Microbiol.
1993 Feb;59(2):473–8.

46. Delden C van, Page MGP, Köhler T. Involvement of Fe Uptake Systems and AmpC β-Lactamase in
Susceptibility to the Siderophore Monosulfactam BAL30072 in Pseudomonas aeruginosa.
Antimicrobial Agents and Chemotherapy. 2013 May;57(5):2095.

47. Déziel E, Gopalan S, Tampakaki AP, Lépine F, Padfield KE, Saucier M, et al. The contribution of
MvfR to Pseudomonas aeruginosa pathogenesis and quorum sensing circuitry regulation:
multiple quorum sensing-regulated genes are modulated without affecting lasRI, rhlRI or the
production of N-acyl- l-homoserine lactones. Molecular Microbiology. 2005;55(4):998–1014.

48. Irons JL, Hodge-Hanson K, Downs DM. RidA Proteins Protect against Metabolic Damage by
Reactive Intermediates. Microbiology and Molecular Biology Reviews. 2020 Jul
15;84(3):10.1128/mmbr.00024-20.

49. Irons J, Hodge-Hanson KM, Downs DM. PA5339, a RidA Homolog, Is Required for Full Growth in
Pseudomonas aeruginosa. Journal of Bacteriology. 2018 Oct 23;200(22):10.1128/jb.00434-18.

50. Weiler AJ, Spitz O, Gudzuhn M, Schott-Verdugo SN, Kamel M, Thiele B, et al. A phospholipase B
from *Pseudomonas aeruginosa* with activity towards endogenous phospholipids affects biofilm
assembly. Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids. 2022 Apr
1;1867(4):159101.

51.   Yoon SS, Hennigan RF, Hilliard GM, Ochsner UA, Parvatiyar K, Kamani MC, et al. Pseudomonas
aeruginosa Anaerobic Respiration in Biofilms: Relationships to Cystic Fibrosis Pathogenesis.
Developmental Cell. 2002;

52.   König J, Schreiber R, Voelcker T, Mall M, Kunzelmann K. The cystic fibrosis transmembrane
conductance regulator (CFTR) inhibits ENaC through an increase in the intracellular Cl–
concentration. EMBO Rep. 2001 Nov 15;2(11):1047–51.

53.   Cowley ES, Kopf SH, LaRiviere A, Ziebis W, Newman DK. Pediatric Cystic Fibrosis Sputum Can Be
Chemically Dynamic, Anoxic, and Extremely Reduced Due to Hydrogen Sulfide Formation. mBio.
2015 Jul 28;6(4):10.1128/mbio.00767-15.

54.   Line L, Alhede M, Kolpen M, Kühl M, Ciofu O, Bjarnsholt T, et al. Physiological levels of nitrate
support anoxic growth by denitrification of Pseudomonas aeruginosa at growth rates reported
in cystic fibrosis lungs and sputum. Front Microbiol. 2014 Oct 24;5:554.

55.   Borriello G, Richards L, Ehrlich GD, Stewart PS. Arginine or Nitrate Enhances Antibiotic
Susceptibility of Pseudomonas aeruginosa in Biofilms. Antimicrobial Agents and Chemotherapy.
2006 Jan;50(1):382–4.

56.   Tribelli PM, Lujan AM, Pardo A, Ibarra JG, Fernández Do Porto D, Smania A, et al. Core regulon
of the global anaerobic regulator Anr targets central metabolism functions in Pseudomonas
species. Sci Rep. 2019 Jun 21;9(1):9065.

57.   Trunk K, Benkert B, Quäck N, Münch R, Scheer M, Garbe J, et al. Anaerobic adaptation in
Pseudomonas aeruginosa: definition of the Anr and Dnr regulons. Environmental Microbiology.
2010;12(6):1719–33.

58.   Ashish A, Paterson S, Mowat E, Fothergill JL, Walshaw MJ, Winstanley C. Extensive diversification
is a common feature of Pseudomonas aeruginosa populations during respiratory infections in
cystic fibrosis. J Cyst Fibros. 2013 Dec;12(6):790–3.

59.   Flynn KM, Dowell G, Johnson TM, Koestler BJ, Waters CM, Cooper VS. Evolution of Ecological
Diversity in Biofilms of Pseudomonas aeruginosa by Altered Cyclic Diguanylate Signaling. Journal
of Bacteriology. 2016 Sep 9;198(19):2608–18.

60.   Jorth P, Staudinger BJ, Wu X, Hisert K, Hayden H, Garudathri J, et al. Regional Isolation Drives
Bacterial Diversification within Cystic Fibrosis Lungs. Cell Host Microbe. 2015 Sep 9;18(3):307–
19.

61.   Markussen T, Marvig RL, Gómez-Lozano M, Aanæs K, Burleigh AE, Høiby N, et al. Environmental
heterogeneity drives within-host diversification and evolution of Pseudomonas aeruginosa.
mBio. 2014 Sep 16;5(5):e01592-01514.

62.   Winstanley C, O'Brien S, Brockhurst MA. Pseudomonas aeruginosa Evolutionary Adaptation and
Diversification in Cystic Fibrosis Chronic Lung Infections. Trends Microbiol. 2016 May;24(5):327–
37.

63.   Kamath KS, Krisp C, Chick J, Pascovici D, Gygi SP, Molloy MP. Pseudomonas aeruginosa Proteome
under Hypoxic Stress Conditions Mimicking the Cystic Fibrosis Lung. J Proteome Res. 2017 Oct
6;16(10):3917–28.

64. Sommer LM, Johansen HK, Molin S. Antibiotic resistance in Pseudomonas aeruginosa and
adaptation to complex dynamic environments. Microbial Genomics. 2020;6(5):e000370.

65. Besse A, Groleau MC, Trottier M, Vincent AT, Déziel E. Pseudomonas aeruginosa Strains from
Both Clinical and Environmental Origins Readily Adopt a Stable Small-Colony-Variant Phenotype
Resulting from Single Mutations in c-di-GMP Pathways. J Bacteriol. 2022 Oct
18;204(10):e0018522.

66. Johns BE, Purdy KJ, Tucker NP, Maddocks SE. Phenotypic and Genotypic Characteristics of Small
Colony Variants and Their Role in Chronic Infection. Microbiol Insights. 2015 Jan 1;8:MBI.S25800.

67. Ryan H, Ballard E, Stockwell RE, Duplancic C, Thomson RM, Smith K, et al. A systematic review of
the clinical impact of small colony variants in patients with cystic fibrosis. BMC Pulm Med. 2023
Sep 1;23(1):323.

68. Schneider M, Mühlemann K, Droz S, Couzinet S, Casaulta C, Zimmerli S. Clinical Characteristics
Associated with Isolation of Small-Colony Variants of Staphylococcus aureus and Pseudomonas
aeruginosa from Respiratory Secretions of Patients with Cystic Fibrosis. Journal of Clinical
Microbiology. 2008 Mar 5;46(5):1832.

69. Malone JG. Role of small colony variants in persistence of Pseudomonas aeruginosa infections
in cystic fibrosis lungs. Infection and Drug Resistance. 2015 Jul 29;8(null):237–47.

70. Besse A, Groleau MC, Déziel E. Emergence of Small Colony Variants Is an Adaptive Strategy Used
by Pseudomonas aeruginosa to Mitigate the Effects of Redox Imbalance. mSphere. 2023 Feb
28;8(2):e00057-23.

71. Hoffman LR, Déziel E, D'Argenio DA, Lépine F, Emerson J, McNamara S, et al. Selection for
Staphylococcus aureus small-colony variants due to growth in the presence of Pseudomonas
aeruginosa. Proceedings of the National Academy of Sciences. 2006 Dec 26;103(52):19890–5.

72. Bjarnsholt T, Jensen PØ, Fiandaca MJ, Pedersen J, Hansen CR, Andersen CB, et al. Pseudomonas
aeruginosa biofilms in the respiratory tract of cystic fibrosis patients. Pediatr Pulmonol. 2009
Jun;44(6):547–58.

73. Høiby N, Ciofu O, Bjarnsholt T. Pseudomonas Aeruginosa Biofilms in Cystic fibrosis. Future
Microbiology. 2010 Nov 1;5(11):1663–74.

74. Maurice NM, Bedi B, Sadikot RT. Pseudomonas aeruginosa Biofilms: Host Response and Clinical
Implications in Lung Infections. Am J Respir Cell Mol Biol. 2018 Apr;58(4):428–39.

75. Klementiev AD, Jin Z, Whiteley M. Micron Scale Spatial Measurement of the O2 Gradient
Surrounding a Bacterial Biofilm in Real Time. mBio. 2020 Oct 20;11(5):10.1128/mbio.02536-20.

76. Wu Y, Klapper I, Stewart PS. Hypoxia arising from concerted oxygen consumption by neutrophils
and microorganisms in biofilms. Pathogens and Disease. 2018 Apr 23;76(4):fty043.

77. Bowler P, Murphy C, Wolcott R. Biofilm exacerbates antibiotic resistance: Is this a current
oversight in antimicrobial stewardship? Antimicrob Resist Infect Control. 2020 Oct 20;9(1):162.

78. Liang Z, Nilsson M, Kragh KN, Hedal I, Alcàcer-Almansa J, Kiilerich RO, et al. The role of individual
exopolysaccharides in antibiotic tolerance of Pseudomonas aeruginosa aggregates. Front
Microbiol [Internet]. 2023 Jun 2 [cited 2024 Jul 2];14. Available from:
https://www.frontiersin.org/journals/microbiology/articles/10.3389/fmicb.2023.1187708/full

79. Caiazza NC, O'Toole GA. SadB is required for the transition from reversible to irreversible
attachment during biofilm formation by Pseudomonas aeruginosa PA14. J Bacteriol. 2004
Jul;186(14):4476–85.

80. Caiazza NC, Merritt JH, Brothers KM, O'Toole GA. Inverse Regulation of Biofilm Formation and
Swarming Motility by Pseudomonas aeruginosa PA14. J Bacteriol. 2007 May;189(9):3603–12.

81. Bazire A, Shioya K, Soum-Soutéra E, Bouffartigues E, Ryder C, Guentas-Dombrowsky L, et al. The
sigma factor AlgU plays a key role in formation of robust biofilms by nonmucoid Pseudomonas
aeruginosa. J Bacteriol. 2010 Jun;192(12):3001–10.

82. Candido Caçador N, Paulino da Costa Capizzani C, Gomes Monteiro Marin Torres LA, Galetti R,
Ciofu O, da Costa Darini AL, et al. Adaptation of Pseudomonas aeruginosa to the chronic
phenotype by mutations in the algTmucABD operon in isolates from Brazilian cystic fibrosis
patients. PLoS One. 2018;13(11):e0208013.

83. Hershberger CD, Ye RW, Parsek MR, Xie ZD, Chakrabarty AM. The algT (algU) gene of
Pseudomonas aeruginosa, a key regulator involved in alginate biosynthesis, encodes an
alternative sigma factor (sigma E). Proc Natl Acad Sci U S A. 1995 Aug 15;92(17):7941–5.

84. Moscoso JA, Jaeger T, Valentini M, Hui K, Jenal U, Filloux A. The Diguanylate Cyclase SadC Is a
Central Player in Gac/Rsm-Mediated Biofilm Formation in Pseudomonas aeruginosa. Journal of
Bacteriology. 2014 Oct 29;196(23):4081–8.

85. Schmidt A, Hammerbacher AS, Bastian M, Nieken KJ, Klockgether J, Merighi M, et al. Oxygen-
dependent regulation of c-di-GMP synthesis by SadC controls alginate production in
seudomonas aeruginosa. Environmental Microbiology. 2016;18(10):3390–402.

86. Stacey SD, Pritchett CL. Pseudomonas aeruginosa AlgU Contributes to Posttranscriptional
Activity by Increasing rsmA Expression in a mucA22 Strain. Journal of Bacteriology. 2016 Jun
13;198(13):1812–26.

87. Vijay-Kumar M, Gewirtz AT. Flagellin: key target of mucosal innate immunity. Mucosal
Immunology. 2009 May 1;2(3):197–205.

88. Neidig A, Yeung AT, Rosay T, Tettmann B, Strempel N, Rueger M, et al. TypA is involved in
virulence, antimicrobial resistance and biofilm formation in Pseudomonas aeruginosa. BMC
Microbiol. 2013 Apr 9;13(1):77.

89. Overhage J, Lewenza S, Marr AK, Hancock REW. Identification of genes involved in swarming
motility using a Pseudomonas aeruginosa PAO1 mini-Tn5-lux mutant library. J Bacteriol. 2007
Mar;189(5):2164–9.

90. Sindeldecker D, Stoodley P. The many antibiotic resistance and tolerance strategies of
Pseudomonas aeruginosa. Biofilm. 2021 Dec 1;3:100056.

91. Borriello G, Werner E, Roe F, Kim AM, Ehrlich GD, Stewart PS. Oxygen Limitation Contributes to
Antibiotic Tolerance of Pseudomonas aeruginosa in Biofilms. Antimicrobial Agents and
Chemotherapy. 2004 Jul;48(7):2659–64.

92. Schaible B, Taylor CT, Schaffer K. Hypoxia Increases Antibiotic Resistance in Pseudomonas
aeruginosa through Altering the Composition of Multidrug Efflux Pumps. Antimicrobial Agents
and Chemotherapy. 2012 Mar 16;56(4):2114–8.

93. Zhao X, Drlica K. Reactive oxygen species and the bacterial response to lethal stress. Current
opinion in microbiology. 2014 Jul 30;0:1.

94. Puchelle E, Bajolet O, Abély M. Airway mucus in cystic fibrosis. Paediatric Respiratory Reviews.
2002 Jun 1;3(2):115–9.

95. Bhagirath AY, Li Y, Somayajula D, Dadashi M, Badr S, Duan K. Cystic fibrosis lung environment and
Pseudomonas aeruginosa infection. BMC Pulm Med. 2016 Dec 5;16:174.

96. Lan L, Murray TS, Kazmierczak BI, He C. Pseudomonas aeruginosa OspR is an oxidative stress
sensing regulator that affects pigment production, antibiotic resistance and dissemination
during infection. Molecular Microbiology. 2010;75(1):76–91.

97. Fraud S, Poole K. Oxidative Stress Induction of the MexXY Multidrug Efflux Genes and Promotion
of Aminoglycoside Resistance Development in Pseudomonas aeruginosa. Antimicrobial Agents
and Chemotherapy. 2011 Feb 24;55(3):1068–74.

98. Mohamed FA, Shaker GH, Askoura MM. Oxidative Stress Influences Pseudomonas aeruginosa
Susceptibility to Antibiotics and Reduces Its Pathogenesis in Host. Curr Microbiol. 2020 Mar
1;77(3):479–90.

99. Bojanovič K, D'Arrigo I, Long KS. Global Transcriptional Responses to Osmotic, Oxidative, and
Imipenem Stress Conditions in Pseudomonas putida. Applied and Environmental Microbiology.
2017 Mar 17;83(7):e03236.

100. Dawan J, Ahn J. Bacterial Stress Responses as Potential Targets in Overcoming Antibiotic
Resistance. Microorganisms. 2022 Jul;10(7):1385.

101. Hunter RC, Asfour F, Dingemans J, Osuna BL, Samad T, Malfroot A, et al. Ferrous Iron Is a
Significant Component of Bioavailable Iron in Cystic Fibrosis Airways. mBio. 2013 Aug
20;4(4):10.1128/mbio.00557-13.

102. Moreau-Marquis S, Bomberger JM, Anderson GG, Swiatecka-Urban A, Ye S, O'Toole GA, et al.
The DeltaF508-CFTR mutation results in increased biofilm formation by Pseudomonas
aeruginosa by increasing iron availability. Am J Physiol Lung Cell Mol Physiol. 2008
Jul;295(1):L25-37.

103. Schalk IJ, Perraud Q. and its multiple strategies to access iron. Environmental Microbiology.
2023;25(4):811–31.

104. Reid DW, Carroll V, O'May C, Champion A, Kirov SM. Increased airway iron as a potential factor
in the persistence of Pseudomonas aeruginosa infection in cystic fibrosis. European Respiratory
Journal. 2007 Aug 1;30(2):286–92.

105. Konings AF, Martin LW, Sharples KJ, Roddam LF, Latham R, Reid DW, et al. Pseudomonas
aeruginosa Uses Multiple Pathways To Acquire Iron during Chronic Infection in Cystic Fibrosis
Lungs. Infection and Immunity. 2013 Jul 11;81(8):2697–704.

106. Lo Sciuto A, D'Angelo F, Spinnato MC, Garcia PS, Genah S, Matteo C, et al. A molecular
comparison of [Fe-S] cluster-based homeostasis in Escherichia coli and Pseudomonas
aeruginosa. mBio. 2024 Oct 3;0(0):e01206-24.

107. Py B, Barras F. Building Fe–S proteins: bacterial strategies. Nat Rev Microbiol. 2010 Jun;8(6):436–
46.

108. Romsang A, Duang-Nkern J, Leesukon P, Saninjuk K, Vattanaviboon P, Mongkolsuk S. The Iron-
Sulphur Cluster Biosynthesis Regulator IscR Contributes to Iron Homeostasis and Resistance to
Oxidants in Pseudomonas aeruginosa. PLOS ONE. 2014 sty;9(1):e86763.

109. Frangipani E, Visaggio D, Heeb S, Kaever V, Cámara M, Visca P, et al. The Gac/Rsm and cyclic-di-
GMP signalling networks coordinately regulate iron uptake in seudomonas aeruginosa.
Environmental Microbiology. 2014;16(3):676–88.

110. Banin E, Brady KM, Greenberg EP. Chelator-induced dispersal and killing of Pseudomonas
aeruginosa cells in a biofilm. Appl Environ Microbiol. 2006 Mar;72(3):2064–9.

111. Schaible B, Rodriguez J, Garcia A, von Kriegsheim A, McClean S, Hickey C, et al. Hypoxia Reduces
the Pathogenicity of Pseudomonas aeruginosa by Decreasing the Expression of Multiple
Virulence Factors. J Infect Dis. 2017 May 1;215(9):1459–67.

112. Gil-Marqués ML, Pachón-Ibáñez ME, Pachón J, Smani Y. Effect of Hypoxia on the Pathogenesis of
Acinetobacter baumannii and Pseudomonas aeruginosa In Vitro and in Murine Experimental
Models of Infection. Infection and Immunity. 2018 Sep 21;86(10):10.1128/iai.00543-18.

113. Schaible B, McClean S, Selfridge A, Broquet A, Asehnoune K, Taylor CT, et al. Hypoxia Modulates
Infection of Epithelial Cells by Pseudomonas aeruginosa. PLOS ONE. 2013 Feb 13;8(2):e56491.

114. Vilaplana L, Marco MP. Phenazines as potential biomarkers of Pseudomonas aeruginosa
infections: synthesis regulation, pathogenesis and analytical methods for their detection. Anal
Bioanal Chem. 2020 Sep 1;412(24):5897–912.

115. Jiao H, Li F, Wang T, Yam JKH, Yang L, Liang H. The Pyocin Regulator PrtR Regulates Virulence
Expression of Pseudomonas aeruginosa by Modulation of Gac/Rsm System and c-di-GMP
Signaling Pathway. Infection and Immunity [Internet]. 2021 Jan 19 [cited 2024 Oct 23];89(2).
Available from: https://journals.asm.org/doi/full/10.1128/iai.00602-20

116. Weimann A, Dinan AM, Ruis C, Bernut A, Pont S, Brown K, et al. Evolution and host-specific
adaptation of Pseudomonas aeruginosa. Science. 2024 Jul 5;385(6704):eadi0908.

117. Aguirre-Ramírez M, Medina G, González-Valdez A, Grosso-Becerra V, Soberón-Chávez G. The
Pseudomonas aeruginosa rmlBDAC operon, encoding dTDP-l-rhamnose biosynthetic enzymes,
is regulated by the quorum-sensing transcriptional regulator RhlR and the alternative sigma
factor σS. Microbiology. 2012;158(4):908–16.

118. Rahim R, Burrows LL, Monteiro MA, Perry MB, Lam JS. Involvement of the rml locus in core
oligosaccharide and O polysaccharide assembly in Pseudomonas aeruginosa. Microbiology.
2000;146(11):2803–14.

119. Koska M, Kordes A, Erdmann J, Willger SD, Thöming JG, Bähre H, et al. Distinct Long- and Short-
Term Adaptive Mechanisms in Pseudomonas aeruginosa. Microbiology Spectrum. 2022 Nov
14;10(6):e03043-22.

120. McHugh JP, Rodríguez-Quiñones F, Abdul-Tehrani H, Svistunenko DA, Poole RK, Cooper CE, et al.
Global Iron-dependent Gene Regulation in *Escherichia coli*. Journal of Biological Chemistry. 2003
Aug 8;278(32):29478–86.

121. Touati D. Iron and oxidative stress in bacteria. Arch Biochem Biophys. 2000 Jan 1;373(1):1–6.

122. Zhang Y, Zhang X, Zhang H, He Q, Zhou Q, Su Z, et al. Responses of Soil Bacteria to Long-Term
and Short-Term Cadmium Stress as Revealed by Microbial Community Analysis. Bull Environ
Contam Toxicol. 2009 Mar 1;82(3):367–72.

123. Krulwich TA, Sachs G, Padan E. Molecular aspects of bacterial pH sensing and homeostasis.
Nature reviews Microbiology. 2011 Apr 5;9(5):330.

124. Clay ME, Hammond JH, Zhong F, Chen X, Kowalski CH, Lee AJ, et al. Pseudomonas aeruginosa
lasR mutant fitness in microoxia is supported by an Anr-regulated oxygen-binding hemerythrin.
Proc Natl Acad Sci U S A. 2020 Feb 11;117(6):3167–73.

125. Neville N, Roberge N, Ji X, Stephen P, Lu JL, Jia Z. A Dual-Specificity Inhibitor Targets
Polyphosphate Kinase 1 and 2 Enzymes To Attenuate Virulence of Pseudomonas aeruginosa.
mBio. 2021 Jun 15;12(3):e00592.

126. Ortiz-Severín J, Varas M, Bravo-Toncio C, Guiliani N, Chávez FP. Multiple antibiotic susceptibility
of polyphosphate kinase mutants (ppk1 and ppk2) from Pseudomonas aeruginosa PAO1 as
revealed by global phenotypic analysis. biol res. 2015 Apr 25;48(1):22.

127. Salunkhe P, Smart CHM, Morgan JAW, Panagea S, Walshaw MJ, Hart CA, et al. A Cystic Fibrosis
Epidemic Strain of Pseudomonas aeruginosa Displays Enhanced Virulence and Antimicrobial
Resistance. Journal of Bacteriology. 2005 Jul;187(14):4908–20.

128. An S, Wu J, Zhang LH. Modulation of Pseudomonas aeruginosa Biofilm Dispersal by a Cyclic-Di-
GMP Phosphodiesterase with a Putative Hypoxia-Sensing Domain. Appl Environ Microbiol. 2010
Dec;76(24):8160–73.

129. Feng Q, Zhou J, Zhang L, Fu Y, Yang L. Insights into the molecular basis of c-di-GMP signalling in
Pseudomonas aeruginosa. Critical Reviews in Microbiology. 2024 Jan 2;50(1):20–38.

130. Hickman JW, Harwood CS. Identification of FleQ from Pseudomonas aeruginosa as a c-di-GMP-
responsive transcription factor. Mol Microbiol. 2008 Jul;69(2):376–89.

131. Matsuyama BY, Krasteva PV, Baraquet C, Harwood CS, Sondermann H, Navarro MVAS.
Mechanistic insights into c-di-GMP-dependent control of the biofilm regulator FleQ from
Pseudomonas aeruginosa. Proc Natl Acad Sci U S A. 2016 Jan 12;113(2):E209-218.

132. Zhou T, Huang J, Liu Z, Lin Q, Xu Z, Zhang LH. The Two-Component System FleS/FleR Represses
H1-T6SS via Cyclic di-GMP Signaling in Pseudomonas aeruginosa. Appl Environ Microbiol. 2022
Jan 25;88(2):e0165521.

133. Amoureux L, Bador J, Fardeheb S, Mabille C, Couchot C, Massip C, et al. Detection of
Achromobacter xylosoxidans in Hospital, Domestic, and Outdoor Environmental Samples and
Comparison with Human Clinical Isolates. Appl Environ Microbiol. 2013 Dec;79(23):7142–9.

134. Crone S, Vives-Flórez M, Kvich L, Saunders AM, Malone M, Nicolaisen MH, et al. The
environmental occurrence of Pseudomonas aeruginosa. APMIS. 2020;128(3):220–31.

135. Honda JR, Virdi R, Chan ED. Global Environmental Nontuberculous Mycobacteria and Their
Contemporaneous Man-Made and Natural Niches. Front Microbiol. 2018 Aug 30;9.

136. Honda JR, Virdi R, Chan ED. Global Environmental Nontuberculous Mycobacteria and Their
Contemporaneous Man-Made and Natural Niches. Front Microbiol. 2018;9:2029.

137. Price CTD, Hanford HE, Al-Quadan T, Santic M, Shin CJ, Da'as MSJ, et al. Amoebae as training
grounds for microbial pathogens. mBio. 2024 Jul 8;15(8):e00827-24.

138. Mullard A. 2023 FDA approvals. Nat Rev Drug Discov. 2024 Feb;23(2):88–95.

139. Essar DW, Eberly L, Hadero A, Crawford IP. Identification and characterization of genes for a
second anthranilate synthase in Pseudomonas aeruginosa: interchangeability of the two
anthranilate synthases and evolutionary implications. J Bacteriol. 1990 Feb;172(2):884–900.

140. Ge SX, Jung D, Yao R. ShinyGO: a graphical gene-set enrichment tool for animals and plants.
Bioinformatics. 2020 Apr 15;36(8):2628-9.

141. Letunic I, Bork P. Interactive Tree of Life (iTOL) v6: recent updates to the phylogenetic tree display
and annotation tool. Nucleic Acids Research. 2024 Apr 13:gkae268.

1157

1158

1159

1160

1161

1162

1163

1164

1165

1166

1167

1168

Supporting information captions

S1. Figure. Differenced of growth patterns of the *P. aeruginosa* AMT 0023-30 early infection isolate with different inoculation volumes.

S2. Figure. Differences in doubling times of the *P. aeruginosa* AMT 002330 early infection strain in hypoxic and normoxic conditions. A. OD_{600} measurements over time. **B.** CFU/ml counts over time **C.** OD_{600} to CFU/ml trend lines and linear regression calculations. **D.** Differences in generation times in normoxia and hypoxia.

S3. Table. Proteins differentially expressed after 28 day exposure to hypoxia

S4. Table. Functional enrichment of proteins changed in abundance after 28 days of exposure to hypoxia

Figure 1. Long-term adaptation of *Pseudomonas aeruginosa* to hypoxic conditions. Experimental evolution study design (**A.**) and influence of hypoxia on small colony variants emergence **B.** Percentage of SCVs in individual normoxia- and hypoxia adapting cultures over 28 days. **C.** Hypoxia-adapted colony morphotypes observed throughout the long-term adaptation. **D.** Ratios of SCV 1 and SCV 2 in the hypoxia-adapting populations over 28 days. The percentage of the small colony variants in the whole populations was caried out for each flask individually each day of the experiment.

Figure 2. Influence of hypoxia on pyocyanin production. A. Representative photographs of flasks during the

28 day experimental evolution study showing changes in pyocyanin production. **B.** Quantification of pyocyanin

production by individual populations. **C.** Quantification of pyocyanin production, pooled data for normoxia-

and hypoxia-adapted cultures. The data represents the concentration of pyocyanin in 1ml of bacterial culture,

calculated as described in the methods. The data represents two separate experiments performed in normoxic

conditions, with at least 3 technical replicates for each strain. Statistically significant differences by ordinary

one-way ANOVA test compared to the early infection strain with an uncorrected Fishers LSD test (B.) or

Kruskal-Wallis one-way ANOVA test comparing all groups with an uncorrected Dunn's test (C.) are shown (**p<

0.01; ***p<0.001; ****p< 0.0001). E: early infection clinical isolate; L: late infection clinical isolate; N1, N2,

N3: 28-day normoxia-adapted populations; H1, H2, H3: 28-day hypoxia adapted populations.

Figure 3. Whole proteome comparison of long-term hypoxia-adapted and normoxia-adapted *P. aeruginosa.*

A. Heat map representing expression patterns in adapted populations. The heatmap was generated with the use of Heatmapper (Babicki et al., 2016). **B.** Dot plot representing functional GoTerm Biological Process enrichment of proteins with increased and decreased abundance in hypoxia. The enrichment analysis and dot plot were prepared using ShinyGo 0.80 (140).

Figure 4. Effects of long-term hypoxia adaptation on antibiotic resistance and stress resistance. **A.** Graphs represent the antibiotic susceptibility of early infection control strain AMT 0023-30, late infection strain, AMT 0023-31, and 28-day adapted populations using diffusion discs, tested in normoxic conditions. Bars represent the mean inhibition zone diameter. The data represents three biological replicates and error bars represent the standard deviation. The significance was determined with the nonparametric Kruskal-Wallis test in combination with uncorrected Dunns multiple comparisons test. **B**. Summary of resistance to all tested antibiotics. **C.** Droplet assays. Representative pictures of colonies from serial dilutions of each strain plated on agar with stress factor added, tested in normoxic conditions. The experiment was performed at least 3 times for each stress type with the early infection isolate as a control. **D.** Table summarising changes in stress sensitivity. ↑ - increased resistance; ↓ - decreased resistance; = no significant difference in resistance; R – no inhibition zone observed. E: early infection clinical isolate; L: late infection clinical isolate; N1, N2, N3: 28-day normoxia-adapted populations; H1, H2, H3: 28-day hypoxia adapted populations.

Figure 5. Effects of long-term hypoxia adaptation on biofilm formation, exopolysaccharide production and siderophore production. A. Biofilm formation capability of hypoxia-adapted cultures determined with the crystal violet assay after 24, 48 or 72h of static incubation on a 96-well plate in normoxic conditions. Data represents the mean value of the OD_{590}/OD_{600} ratio from three separate experiments with at least 6 technical replicates for each strain. Statistically significant difference by Kruskal-Wallis one-way ANOVA test compared to the early infection strain (E) with an uncorrected Dunn's test (**$p< 0.01$; ***$p<0.001$; ****$p< 0.0001$). B. Exopolysaccharide production by the adapted cultures. The photographs show colonies of different populations of bacteria. The change of colour to a darker shade and a wrinkly colony surface indicate a higher exopolysaccharide production. C. Pathway of pyochelin synthesis. Pink: proteins decreased in abundance in the proteome of hypoxia-adapted populations. D. Siderophore production represented as an average of siderophore units produced by each colony forming unit, tested in normoxic conditions. Statistically significant differences by ordinary one-way ANOVA test compared to the early infection strain with an uncorrected Fishers LSD test are shown (*$p<0.05$; **$p< 0.01$). E: early infection clinical isolate; L: late infection clinical isolate; N1, N2, N3: 28-day normoxia-adapted populations; H1, H2, H3: 28-day hypoxia adapted populations. E: early infection clinical isolate, AMT 0023-30; L: late infection clinical isolate, AMT 0023-31; N1, N2, N3: 28-day normoxia-adapted populations; H1, H2, H3: 28-day hypoxia adapted populations.

Figure 6. Effects of long-term hypoxia adaptation on *P. aeruginosa* motility and virulence in the *Galleria mellonela* model. A. Changes in swimming motility**. B.** Changes in swarming motility. **C.** Changes in twitching motility. For each type of motility, a graph representing the average of diameter of the colony with bars representing data from at least 3 separate experiments conducted in normoxic conditions, with at least two technical replicates. Statistically significant difference by Kruskal-Wallis one-way ANOVA test compared to the early infection strain (E) with an uncorrected Dunn's test (*p< 0.05; **p< 0.01; ***p<0.001; ****p< 0.0001). **D.** Graph representing the average CFU needed to reach LD50 within 24h of the experiment from at least 3 separate repeats of the experiment**.** A touch to the larvae head was used to determine if they are alive. Statistically significant difference by Kruskal-Wallis one-way ANOVA test compared to the early infection strain (E) with an uncorrected Dunn's test is presented (*p< 0.05). **E.** A representative picture showing the *G. mellonela* larvae melanisation. Healthy larvae with no disease symptoms are a bright cream colour. Larvae showing a slightly beige colour show first symptoms of infection, larvae with dark spots/stipes are clearly affected and the severely melanised larvae are in the late stages of the infection or dead. E: early infection clinical isolate; L: late infection clinical isolate; N1, N2, N3: 28-day normoxia-adapted populations; H1, H2, H3: 28 day hypoxia adapted populations.

A.

Phenotype		L	N1	N2	N3	H1	H2	H3
SCVs (%)		58.0%	0.00	0.00	2.7%	76.9%	79.8%	0.7%
Pyocyanin		↓	↓	=	=	↓	↓	=
Antibiotic resistance		=	=	=	=	↑	↑	=
Stress resistance	Osmotic	↓	↑	=	↑	↑	↑	↑
	Oxidative	=	=	=	↑	↑	↑	=
	Heat	↓	↓	↓	=	↓	↓	=
	Acidic	=	↑	↑	↑	=	=	=
Biofilm 72h		↑	=	↑	=	↑	↑	=
Motility	Swimming	↓	↑	↑	↓	↑	↑	↓
	Swarming	↓	=	↑	↓	↑	↑	↓
	Twitching	↓	=	↑	=	=	↑	↓
Siderophore production		↓	=	=	=	↓	↓	↓
Virulence (*Galleria mellonela*)		↑	=	=	=	=	=	=

B.

Figure 7. Patterns of adaptation to long-term hypoxia. A. Summary of the phenotypic changes of adapted cultures tested in normoxic conditions. B. Phylogenetic tree of adapted *P. aeruginosa* populations generated by the iTOL Interactive tree of life (v6) (141). The branch lengths are presented for each isolate.