Long-term adaptation to hypoxia provides insight into mechanisms facilitating the switch of *Pseudomonas aeruginosa* to chronic lung infections

Joanna Drabinska[1,2]*, Lucia O'Connor[1,2], Niamh Duggan[1,2], Ciarán J. Carey[1,2], Siobhán McClean[1,2]*

[1]School of Biomolecular and Biomedical Sciences, University College Dublin, Belfield, Dublin, Ireland.

[2]Conway Institute of Biomolecular and Biomedical Science, University College Dublin, Belfield, Dublin, Ireland.

*Corresponding authors

Email: siobhan.mcclean@ucd.ie; joanna.drabinska@ucd.ie

Abstract

Opportunistic bacterial infections are an increasing threat, especially for immunocompromised individuals such as people with cystic fibrosis (CF), driving morbidity and mortality. *Pseudomonas aeruginosa* is a key pathogen associated with chronic lung infections, that has been extensively studied in this context, but the mechanism(s) driving its adaptation towards chronic colonisation in the lung are not fully understood. This work focuses on the adaptations of *P. aeruginosa* to long-term hypoxia, one of the important environmental pressures present in the CF lung, to investigate whether it drives the development of persistence in CF patients. We used an experimental evolution approach to investigate how an early CF strain adapted to 6% oxygen over 28 days. We focussed on the impact of long-term hypoxia on the proteome and investigated the emergence of stable changes in phenotype. Changes in the abundance of >140 proteins were observed compared to the ancestral strain, including proteins involved in antibiotic resistance, stress response, iron homeostasis, biofilm formation and those previously associated with chronic infection. Significant changes in the abundance of proteins regulating cellular c-di-GMP levels were also observed. We show that two distinct *P. aeruginosa* small colony variants (SCVs) emerged, one exclusively in hypoxia exposed cultures. Hypoxia-adapted cultures developed resistance to 8 out of 13 antibiotics tested; increased biofilm and exopolysaccharide production; and decreased pyocyanin production, consistent with the changes in the proteome. All hypoxia-adapted cultures showed decreased siderophore production. Overall, we demonstrate that long-term hypoxia exposure contributes to multiple changes in phenotype and proteome that are frequently observed in *P. aeruginosa* CF lung chronic infection isolates. This suggests that hypoxia is driving these adaptations, at least in part, and opens a new path to treatment.

Author Summary

Opportunistic antibiotic-resistant bacterial infections are an increasing threat, particularly for immunocompromised individuals such as people with cystic fibrosis (CF). These infections increase disease severity, the number of deaths and healthcare costs. *Pseudomonas aeruginosa* is a major pathogen in this context, causing persistent infections, yet the mechanisms driving its adaptation towards chronic colonisation in the CF lung are not understood.

We investigated the adaptations that *P. aeruginosa* exposed to low oxygen conditions (an important environmental pressure in the CF lung) for 28 days, to examine whether it drives adaptations that lead to chronic infection. We found that prolonged exposure to hypoxia induced stable changes in *P. aeruginosa*, which are distinct from those reported after short term exposure. We observed multiple adaptations in hypoxia adapted cultures, many of which were associated with important signalling networks linked to lifestyle shifts in bacteria and several were also observed in isolates that had adapted to chronic infection. Similar adaptive responses in other opportunistic environmental pathogens suggest that this process of adaptation could be targeted to create universal therapeutic approaches preventing antibiotic overuse and improving patient outcomes.

Introduction

Antibiotic-resistant bacterial infections are a growing threat to society. They were associated with nearly five million deaths in 2019 (1), and are expected to cause more than 10 million deaths annually by 2050 (2). Opportunistic pathogens such as *Pseudomonas aeruginosa* play a crucial role in escalating this crisis. While these bacteria typically do not cause infections in healthy people, immunocompromised individuals and people with underlying conditions such as cystic fibrosis (CF) and chronic obstructive pulmonary disease (COPD) or those on ventilator support are at great risk of developing chronic *P. aeruginosa* lung infections (3,4). Moreover, opportunistic pathogens are responsible for the majority of healthcare-associated infections, contributing to increasing morbidity, mortality and costs (5).

P. aeruginosa is a Gram-negative opportunistic human pathogen, which is a leading cause of nosocomial infections, constituting 11-26% of ventilator-associated pneumonia globally (6). Overall, it causes more than 7% of all health-associated infections and is a major contributor to morbidity and mortality among people with CF (7–9). It has many intrinsic and acquired mechanisms of antibiotic resistance contributing to its spread and persistence (10), and the prolonged antibiotic treatments necessary to manage *P. aeruginosa* chronic infections contribute to the spread of antimicrobial resistance (11,12). Furthermore, the World Health Organisation recognised *P. aeruginosa* as a high priority pathogen in 2024 while the European Centre for Disease Prevention and Control (ECDC) considers it a serious threat (13–16).

A key feature of *P. aeruginosa* is its exquisite metabolic flexibility and adaptability to harsh conditions, allowing it to transition from the colonisation of the rhizosphere or hydrosphere to colonising niches within human or animal hosts, and establishing persistent infections (4,17). *P. aeruginosa* chronic infection isolates show multiple changes that attenuate virulence, allowing it to

avoid immune recognition, thereby enhancing survival, and consequently, chronic infections are

extremely difficult to eradicate (18). Its adaptability is driven by a complex regulatory network

involving transcriptional regulators, DNA modifications, non-coding RNAs, quorum sensing, and a

range of virulence factors (18–23). These adaptations and associated lifestyle changes are also

driven by the alterations in c-di-GMP second messenger levels (24–27). Despite this extensive

research, the mechanism(s) enabling *P. aeruginosa* to transition from acute to chronic infection

remain unclear.

P. aeruginosa, a facultative anaerobe, thrives under low oxygen and anaerobic conditions

using alternative terminal electron acceptors and fermentation substrates. It preferentially colonizes

hypoxic niches within mucus layers of the lung over more oxygenated surfaces (28). *P. aeruginosa*

responds to changes in oxygen availability via the transcriptional regulator Anr, which regulates

many genes shown to be altered in CF chronic infection (29,30). Studies on another CF-associated

pathogen, *Burkholderia cenocepacia* have shown that 40% of proteins encoded within a low oxygen

activated (Lxa) locus showed increased abundance in chronic infection isolates from CF patients

(31). Subsequent studies showed that deletion of single genes from the Lxa or the entire locus

resulted in changes in phenotypes crucial for the virulence of this bacterium including host cell

attachment, intracellular survival and stress responses (31–33). Additionally, the *sicX* sRNA has been

proposed to be a biomarker of *P. aeruginosa* chronic infection and is also oxygen responsive (34).

Given that hypoxia is an important environmental pressure in the CF lung, we wanted to evaluate

the impact of long-term hypoxia exposure on *P. aeruginosa.*

We developed an experimental evolution approach and demonstrate that the impact of

long-term exposure to low oxygen on the proteome and phenotype of *P. aeruginosa* is consistent

with many adaptations that are associated with chronic lung infection. We specifically evaluated on

the proteome and phenotype to focus on the effective changes in the strains following long-term

hypoxia, which is not widely studied. Our findings provide new insights into the mechanisms that underlie the persistence and pathogenesis of *P. aeruginosa* in chronic infection and possibly other opportunistic pathogens, within the host.

Results

Design and Optimisation of Experimental Adaptation

To investigate the impact of long-term exposure to low oxygen conditions on the adaptation of *P. aeruginosa*, we designed an experimental evolution/adaptation approach (Fig 1A), exposing the *P. aeruginosa* early infection strain AMT 0023-30 to either 6% oxygen or to normoxia (21%) for 28 days. AMT 0023-30 is a well-characterised and sequenced CF paediatric early infection strain that was selected from the international *P. aeruginosa* panel (35–38). This strain was isolated from a 6 month old paediatric patient with no former history of *P. aeruginosa* infection, thus the likelihood of preadaptation to the lung or hypoxic niches is very low. Furthermore, a sequential, 96-month, chronic infection strain is available (AMT 0023-34) which has also been sequenced and well characterised (35–38). This strain is the ancestor of the adapted cultures further described in this manuscript (Table 1). After considering the ranges of oxygen that *P. aeruginosa* is exposed to in soil and water, together with the range of hypoxic conditions reported in the CF lung and available literature on physiological tissue hypoxia (28,39,40), we chose 6% oxygen as a low oxygen concentration. Soil aeration ranges from 10 to 20% but decreases with depth and microoxic and anoxic niches are available (39). Similarly a range of oxic, hypoxic and anoxic conditions have been reported for CF patient lungs (28, 40) ranging from180 mmHg in the lumen to 2.5 mmHg within the mucus. Thus 6% O_2 is in the higher hypoxia range and it allowed us to observe the response to hypoxia, while not excessively stressing the cells, so that we can distinguish this response from changes associated with anoxia.

Figure 1. Long-term adaptation of *Pseudomonas aeruginosa* to hypoxic conditions. Experimental evolution study design (**A.**) and influence of hypoxia on small colony variants emergence (**B.-D.**). **A.** Experimental evolution study design. The overnight culture of the ancestral strain was inoculated into three flasks with hypoxia or normoxia equilibrated medium and incubated with agitation in 6% O_2 or 21% O_2. Every 24 h 5% of the culture was transferred into a fresh portion of equilibrated media. The transfers of the hypoxia-adapting cultures were performed under hypoxic conditions to avoid oxygen level fluctuation. The samples from the cultures adapted for 28 days were taken for further proteomic and phenotypic analysis. The details of the procedure are described in the method section. **B.** Hypoxia-adapted colony morphotypes observed throughout the long-term adaptation. **C.** Percentage of SCVs in individual normoxia- and hypoxia adapting cultures over 28 days. **D.** Ratios of SCV 1 and SCV 2 in the hypoxia-adapting populations over 28 days. The percentage of the small colony variants in the whole populations was carried out for each flask individually each day.

To maintain stable growth over the period of the evolution experiment, daily culture transfer volumes were optimised by evaluating a range of volumes (1-10%) of overnight *P. aeruginosa* cultures transferred into fresh media and analysing their growth patterns (S1 Figure). A 5% volume transfer was chosen as optimal, as it allowed the transfer of the maximum number of bacterial cells without disrupting the growth pattern and resulted in a starting OD_{600} of ~0.1. Examination of growth patterns of the early infection strain revealed small differences in the generation times observed during hypoxic and normoxic conditions, with the strain growing 1.8-fold slower under hypoxia (g = 107 minutes) compared to normoxia (g = 60 minutes), and showing logarithmic phases of 3 h in normoxia and 6 h in hypoxia (S2 Figure). Consequently, we estimate that during the 28 day adaptation period, all the cultures underwent 84-94 generations.

To facilitate further investigation, including phenotyping and proteome analysis, three independent flasks were inoculated and incubated under either 6% oxygen or normoxia (21% O_2), and samples were collected every day over 28 days, at the time of culture transfer. Three replicate cultures were chosen for each condition to facilitate deeper analysis over a broad range of phenotypes. To maintain stable hypoxic conditions and avoid exposure to fluctuating oxygen levels, the transfers of hypoxia-adapting cultures were conducted within the hypoxic chamber. The three

28 day independently adapted populations evolved from the ancestral AMT 0023-30 strain are

referred to as N1, N2, and N3 for replicate normoxia-adapted control populations (NACs) and H1,

H2, and H3 for the three replicate hypoxia-adapted cultures (HACs) (Table 1). All characterisation of

the individual adapted populations (both HACs and NACs) was performed under normoxic

conditions, to ensure the observed changes were stable in the absence of continued hypoxia and

are not a transient, plastic response.

	Strain/Population Name	Description	Source
	E	Paediatric (6-month old) early infection cystic fibrosis strain AMT 0023-30. Ancestral strain.	(38)
	L	Paediatric late infection cystic fibrosis strain AMT 0023-34, sequential (96 months) isolate of AMT 0023-30.	(38)
NACs	N1	28 day normoxia-adapted population, replicate 1. Evolved from AMT 0023-30 (E).	This study
	N2	28 day normoxia-adapted population, replicate 2. Evolved from AMT 0023-30 (E).	This study
	N3	28 day normoxia-adapted population, replicate 3. Evolved from AMT 0023-30 (E).	This study
HACs	H1	28 day hypoxia-adapted population, replicate 1. Evolved from AMT 0023-30 (E).	This study
	H2	28 day hypoxia-adapted population, replicate 2. Evolved from AMT 0023-30 (E).	This study
	H3	28 day hypoxia-adapted population, replicate 3. Evolved from AMT 0023-30 (E).	This study

163 **Table 1. List of strains and populations used in this study**

Changes in colony morphology and pyocyanin production during adaptation

Changes in colony morphology often serve as key indicators of adaptation to the environment.

Plating of daily samples from the adapting cultures showed the emergence of small colony variants

(SCVs) in both NACs and HACs; however, the SCVs appeared earlier in HACs over the timeframe (Fig

1B and 1C). SCVs first appeared on day 2 in the H3 population, and by day 3 in the H1 and H2

populations. The SCV abundance peaked at 98% on day 22, with 77% remaining on day 28 in H1. A

similar profile was shown in H2, with SCVs comprising 82% of the population by day 20. In contrast

in population H3, the SCV abundance was generally lower and more variable, reaching only 50% of

the population by day 15, and falling to 1% on day 21 (Fig 1C). The SCVs appeared much later in

normoxic cultures, on days 6, 7 and 8 in flasks N1, N2 and N3 respectively, and only reached a

maximum of 25-33% of the population by day 14 (Fig 1C).

Two distinct SCV morphologies were observed (Fig 1B and 1D). The first type, termed SCV 1,

was observed both under normoxia and hypoxia, and resembled the morphology most often

described as a rugose small colony variant characterised by small size, irregular edges and a heavily

wrinkled, dry surface (41). The second type, SCV 2, was bigger than SCV 1 with a similar rough, dry

surface; however it was easily distinguished by a more regular edge and shape and importantly, it

was exclusive to HACs. The SCVs were stable after replating in normoxic conditions, although SCV 2

morphology differed, with a smoother surface and edges (S3 Figure).

Throughout the 28 days of adaptation, we also observed dynamic changes in culture colour

(Fig 2A), suggesting changes in pigments, such as pyocyanin. Despite almost no visible change in

colour being observed on day 28 (Fig 2A), quantification of pyocyanin production revealed that

populations H1, H2 and one NAC (N1) produced comparable pyocyanin levels at day 28 to the late

infection strain (AMT 0023-34) (Fig 2B and Fig 2C). Although pyocyanin was higher in the NACs

overall compared with HACs, the differences between biological replicates suggests that pyocyanin

production is independent of environmental oxygen levels or is not stable in normoxic conditions.

Figure 2. Influence of hypoxia on pyocyanin production. A. Representative photographs of flasks during the
28 day experimental evolution study showing changes in pyocyanin production. **B. and C.** Quantification of
pyocyanin production by individual populations **(B.)** and mean values for normoxia- and hypoxia-adapted
cultures **(C.)** at 28 days, relative to early (E) and late (L) infection strains. The data represent the
concentration of pyocyanin in 1ml of bacterial culture, calculated as described in the methods. The data
represent two independent experiments performed in normoxic conditions, with at least three technical
replicates for each strain. Statistically significant differences were determined by ordinary one-way ANOVA
test compared to the early infection strain with an uncorrected Fishers LSD test (**B.**) or Kruskal-Wallis
one-way ANOVA test comparing all groups with an uncorrected Dunn's test (**C.**) are shown (**p< 0.01;
***p<0.001; ****p< 0.0001). E: early infection clinical strain; L: late infection clinical strain; N1, N2, N3: 28-
day normoxia-adapted populations; H1, H2, H3: 28-day hypoxia adapted populations.

Influence of hypoxia on the whole proteome

The proteomes of 28 day adapted cultures were analysed to elucidate the overall influence of long-term hypoxia on *P. aeruginosa*. The individual flasks were treated as biological replicates, thus showing the global changes across three independent populations. Principal component analysis (PCA) showed a clear adaptation to the conditions tested, PC1 and PC2 accounted for 84.0% and 7.3% of the total variance respectively (Fig 3A). A total of 257 proteins showed significant differences (adjusted p-value ≤0.10) in abundance between HACs and the ancestor strain, NACs and the ancestor strain and/or between HACs and NACs (Table 2, S4 Table). The majority of proteins that were altered in abundance between HACs, and the ancestral strain were decreased following exposure to hypoxia (n=106; 72.55%), while only 36 proteins (27.45%) showed increased abundance (Table 2, volcano plot, Fig 3C). An additional 13 proteins were identified as significantly altered in abundance after comparing HACs to NACs. Eighty proteins were changed both in HACs and NACs relative to the ancestral strain and, suggesting they may be affected by long-term culture generally and not hypoxia. A total of 89 proteins were further considered as significantly altered during hypoxia adaptation (Fig 3B).

Table 2. Proteome comparison of 28 day the *P. aeruginosa* HACs and NACs vs the early infection ancestral strain and HACs vs NACs.

Protein groups	Number of proteins*
Total number of identified proteins across the ancestral strain, HACs and NACs	1788
Proteins with significantly changed abundance one-way ANOVA (p-value ≤0.05))	749
Proteins with significantly changed abundance in at least one of the comparisons with an adjusted p-value ≤0.1	257
Proteins significantly changed in abundance in HACs vs ancestral strain	142
Proteins significantly changed in abundance in NACs vs ancestral strain	169
Proteins significantly changed in abundance in HACs vs NACs	25
Proteins absent across all replicates in at least one group	117
Proteins with a significantly increased abundance in HACs vs the ancestral strain	36
Proteins with a significantly decreased abundance in HACs vs the ancestral strain	106

*The proteins with significantly changed abundance were determined using mean LFQ intensities of three independent populations of the ancestral, early infection strain, NACs and HACs. One-way ANOVA (p≤0.05) was combined with a FDR adjustment using the Benjamini-Hochberg method. An adjusted p-value cutoff of 0.1 was used.

Figure 3. Whole proteome comparison of long-term hypoxia-adapted and normoxia-adapted *P.*
***aeruginosa.* A.** Principal component analysis plot of quantified proteins from label-free LC-MS/MS analysis of
whole cell extracts from *P. aeruginosa* adapted to hypoxic conditions (HACs) or normoxic conditions (NACs)
for 28 days and the ancestral strain (D0). The PCA plot was performed using LFQ intensities of 1785 proteins
across three biological replicates per condition. Clustering by condition indicates specific global proteomic
shifts associated with long-term hypoxia adaptation. The PCA plot was generated using the Galaxy server v
3.4.0 (42) with ggplot2 using default settings with Euclidean distance clustering. **B.** A Venn diagram of the
number of proteins significantly changed in abundance between HACs and the ancestral strain; NACs and
the ancestral strain; and HACs vs NACs. The analysis was performed using one-way ANOVA with Benjamini-
Hochberg correction and proteins with an adjusted p-value <0.1 were considered significantly altered in
abundance. The 89 proteins in the coloured sections were considered for further analysis. The white sections
were not considered due to lack of relevance to hypoxia adaptation. Only the numbers in parenthesis in the
cross section of NACs and HACs compared to the ancestral strains were considered due to distinct change
between HACs and NACs (1.5 fold-change). **C.** Volcano plot of differentially expressed proteins between
HACs vs the ancestral strain (D0) representing $\log_{10}$ of adjusted p-values and $\log_2$ transformed fold-change
values. Proteins significantly decreased in abundance are highlighted in red and proteins significantly
increased in abundance are highlighted blue. Examples of proteins of interest are labelled. A value of 1 was
imputed for all the zero LFQ values, thus they are represented on the plot, despite the lack of detection. **D.**
STRING-based KEGG pathway enrichment of 89 proteins significantly altered in abundance in HACs vs the
ancestral strain and HACs vs NACs. Enrichment analysis was conducted using STRING functional enrichment
(FDR <0.05) (v12.0) (43). The dot plot shows the top 10 enriched pathways, where dot size corresponds to
the number of hits per pathway and colour represents statistical significance (FDR-corrected p-value). The
pathways are grouped by similarity (0.8). **E.** and **F.** Dot plots representing functional GoTerm Biological
Process enrichment of proteins with increased (E) and decreased abundance (F). Size corresponds to the
number of hits per pathway and colour represents statistical significance. The enrichment analysis and dot
plot were prepared using ShinyGo v0.82 (44).

Proteins unique to hypoxia or normoxia

Interestingly, 117 (15.62%) of the 749 proteins that showed significant changes in abundance (one-
way ANOVA) were undetectable in at least one of the groups: all three HACs, all three NACs or all
three ancestral strain replicates. Their consistent presence or absence suggests that their expression
was either switched on or off, and consequently, they are likely to play critical roles under the
specific conditions. Among these, only three proteins were unique to HACs and not detected in any

NAC or in the ancestral strain: phenazine biosynthesis protein, PhzE; Pel polysaccharide secretion

protein, PelC; and glutamine amidotransferase, HisH2 (Table 3). Additionally, anthranilate synthase

component I PhnA and iron(II)/α-ketoglutarate-dependent oxygenase AmbC, involved in toxin

production were undetected in the ancestral strain, and expressed in all three HACs, but only one of

the NACs.

Many proteins that were undetectable in HACs but expressed in the ancestral strain and all

NACs and thus potentially redundant in low oxygen conditions, are involved in virulence, including

flagellar hook protein FlgE, involved in motility, virulence, host interactions, biofilm formation and

antibiotic resistance; lipid A-disaccharide synthase LpxB, involved in immune evasion; spermidine

dehydrogenase SpdH, involved in chemotaxis and motility. In addition, GacS and ColR, components

of two-component response regulator systems involved in virulence regulation, and phospholipase

B were also undetectable in all HACs, but were present in two of the replicates of the ancestral

strain, and all three NACs. Moreover, oxidoreductase MexS associated with antibiotic resistance and

virulence factor production, chromosome partitioning protein MksE, and iron-sulphur cluster

assembly protein CyaY were detected in all replicates of the ancestral strain and two of the NACs,

while undetectable in all three HACs (Table 3). This attenuation of virulence proteins mirrors the loss

of virulence during chronic infection.

Proteins which were detected both in the ancestral strain and HACs but not detected in

NACs may also be important in the process of adaptation to hypoxia, as their abundance is

maintained. The transcription termination factor Rho associated with biofilm formation and stress

resistance, and UDP-N-acetylglucosamine acyltransferase LpxA belong to this group. Additionally,

polymer-forming cytoskeletal protein PA0664 involved in biofilm formation, and L-lactate

dehydrogenase LldD were detected in all replicates of the ancestral strain and two of the HACs, but

none of the NACs, suggesting their maintenance is required for fitness under low oxygen. Lastly, the

proteins which were undetected in the ancestral strain replicates and all HACs include the acyl-homoserine lactone synthase LasI which produces a quorum-sensing signal and positively regulates numerous *P. aeruginosa* virulence-associated traits including biofilm formation, oxidative stress resistance and elastase production. Additionally, lipase PA1615 and transcriptional regulator PchR, controlling pyochelin production were not detected in all replicates of the ancestral strain and in two of the HACs while expressed in all three NACs. Interestingly, the Anr transcriptional regulator was undetectable both in normoxic and hypoxic conditions.

Table 3. Examples of proteins not detected in at least one of the groups (Day 0, HACs, NACs).

Gene ID[a]	Protein Name	Protein Product Name	DO[b]	HACs[b]	NACs[b]	Function	NP[c]	MW[d]	SC[e]
PA3062	PelC	PelC				Exopolysaccharide (EPS) production, Biofilm formation (45)	6	18.7	58.1
PA1903	PhzE2	Phenazine biosynthesis protein				Pyocyanin production (46)	4	69.0	7.3
PA3152	HisH2	Glutamine amidotransferase				Amino acid biosynthesis (47)	5	22.7	38.6
PA1001	PhnA	Anthranilate synthase component I				QS, pyocyanin production (48)	6	58.7	14
PA2304	AmbC	Iron(II)/α-ketoglutarate-dependent oxygenase				Toxin production, virulence (49)	7	40.7	27.9
PA5139	PA5139	Hypothetical protein				Biofilm formation, Stress resistance (50)	7	27.6	32.3
PA3644	LpxA	UDP-N-acetylglucosamine acyltransferase				LPS production (51)	8	28.0	34.9
PA0664	PA0664	Polymer-forming cytoskeletal protein				Biofilm formation (52)	2	14.8	23.6
PA4771	LldD	L-lactate dehydrogenase				Stress resistance, persistence (53)	16	41.1	45.7
PA1080	FlgE	Flagellar hook protein FlgE				Host interaction, virulence, biofilm formation, antibiotic resistance, motility (54–56)	2	48.3	7.4
PA1157	PA1157	Probable two-component response regulator				Two component system, regulator (57)	3	26.4	13.1
PA3643	LpxB	Lipid A-disaccharide synthase				LPS production, immune evasion (51)	6	41.3	23
PA3713	SpdH	Spermidine dehydrogenase				Motility, chemotaxis (56,58)	9	68.9	19.5
PA4381	ColR	Two-component response regulator ColR				Virulence (57,59,60)	2	25.2	16.3
PA0928	GacS	Sensor/response regulator hybrid				QS, virulence, SCVs (61,62)	5	101.1	8.2
PA2927	PlaB	Phospholipase B				Biofilm formation (63)	4	49.5	5.2
PA2491	MexS	Oxidoreductase MexS				Antibiotic resistance, virulence, pyocyanin production, rhamnolipid production (64–67)	10	36.8	44.2
PA5275	CyaY	Iron-sulphur cluster assembly protein				Iron homeostasis (68)	4	11.9	41.7
PA1432	LasI	Autoinducer synthesis protein LasI				QS, Elastase production, virulence (69,69,70)	3	22.7	20.4
PA1615	PA1615	Probable lipase				Lipase activity (69)	1	12.5	21.2
PA4227	PchR	Transcriptional regulator PchR				Pyochelin production (70)	3	32.3	14.9
PA1544	Anr	Transcriptional regulator Anr				Anaerobic metabolism regulation (71)	4	27.1	15.6

[a]Gene ID – *P. aeruginosa* PAO1 gene locus ID
[b]H/N – Dark blue: protein detected in all three replicates; light blue: protein detected in two out of three replicates; light orange: proteins not detected in two out of three replicates; Dark orange: Protein not detected in all three replicates.
[c]NP - Unique peptides: The number of peptide sequences unique to a protein
[d]MW - Molecular weight as determined by Q-Exactive LC-MS using *P. aeruginosa* PAO1 reference
[e]SC - % sequence covered by matching peptides for the identified protein
Proteins consistently absent or consistently present in all three replicates the ancestral strain HACs and NACs are marked in bold

Functional enrichment

Functional enrichment analysis of the 89 proteins identified in the previous step (Fig 3B) was performed. GO biological process identified 53 enriched categories (S5 Table), including various processes associated with carbon, nitrogen and amino acid metabolism; fatty acid and lipid metabolism; nucleotide and ribonucleotide metabolism. A separate analysis of proteins with increased or decreased abundance in HACs provided further insights into these processes (S5 Table). Overall, 25 GO biological processes were enriched among proteins increased abundance in hypoxia (Fig 3E), while 27 processes were enriched in the decreased abundance group (Fig 3F). Hypoxia led to upregulation of processes related to ribonucleotide metabolism, but a downregulation of pyrimidine-containing compound metabolism. Additionally, processes associated with lipid and fatty acid metabolism, oxidation-reduction and chromosome segregation were upregulated in HACs. Multiple processes associated with amino acid metabolism were also downregulated including glutamine and arginine. Additionally, nitrogen metabolism associated processes were downregulated as well. KEGG pathway enrichment analysis further highlighted alterations of pyrimidine metabolism, changes of RNA degradation pathways, and various carbon, amino acid, and energy metabolism associated pathways (Fig 3D).

Antibiotic resistance of adapted cultures

A change in the abundance of many proteins associated with the resistance to multiple classes of antibiotics was observed in HACs (Table 4). Six proteins were increased in abundance including UDP-N-acetylmuramate-alanine ligase MurC, outer membrane protein H1 precursor OprH, and the Topoisomerase IV subunit A ParC, while five were decreased in abundance.

Table 4. Examples of proteins altered in abundance in *P. aeruginosa* in three comparisons.

			Fold-change[b]						
Gene ID[a]	Protein Name	Protein Product Name	HACs vs D0	NACs vs D0	HACs vs NACs	Function[c]	NP[d]	MW[e]	SC[f]

Gene ID	Gene	Description				Function	NP	MW	SC
PA1903	phzE2	phenazine biosynthesis protein PhzE	30383.00	---	30383.00	Pyocyanin and phenazine biosynthesis (46,72)	4	69.0	7.3
PA2444	glyA2 / ShrA	serine hydroxymethyltransferase	21.96	---	---	SCVs, biofilm formation, EPS production, motility, redox control (73)	10	45.0	47.6
PA3559	PmrE	probable UDP-glucose dehydrogenase	21.53	---	---	EPS production, antibiotic resistance, LPS modification (74,75)	3	51.8	10.6
PA1327	PA1327	probable protease	11.20	---	---	Protease activity	12	72.5	20.3
PA3723	PA3723	probable FMN oxidoreductase	5.46	---	---	Stress resistance, antibiotic resistance (76)	24	40.3	80.4
PA4547	PilR	two-component response regulator PilR	4.52	---	---	Motility, virulence (58)	4	49.7	13.3
PA4411	MurC	UDP-N-acetylmuramate--alanine ligase	3.05	---	---	Antibiotic resistance, peptidoglycan biosynthesis (77,78)	13	52.0	25.8
PA1772	RraA	probable methyltransferase. Ribonuclease activity regulator protein	2.64	---	---	RNA turnover, global regulator (79)	11	17.4	67.3
PA0083	TssB1	TssB1	2.46	---	---	Antibiotic resistance, T6SS (80)	13	18.8	73.8
PA1178	OprH	PhoP/Q and low Mg^{2+} inducible outer membrane protein H1 precursor	2.17	---	---	Antibiotic resistance (81)	23	21.6	69
PA4964	ParC	Topoisomerase IV subunit A	1.77	---	---	Antibiotic resistance (82,83)	18	83.4	34.4
PA0141	Ppk2	conserved hypothetical protein	---	---	1.88	Antibiotic resistance (84)	20	40.8	72.8
PA3262	PA3262	probable peptidyl-prolyl cis-trans isomerase, FkbP-type	---	---	1.74	Virulence; peptidoglycan biosynthesis (85)	21	26.8	48.6
PA4542	ClpB	ClpB protein	---	---	1.37	Stress resistance, virulence (86)	72	95.0	70.3
PA4163	Tse8	hypothetical protein	-2.29	---	---	Virulence (87)	20	60.5	49.6
PA0428	RhlE2	probable ATP-dependent RNA helicase	-2.38	---	---	RNA turnover, global regulator, virulence (72)	30	70.1	51
PA2991	Sth	soluble pyridine nucleotide transhydrogenase	-2.61	---	---	Redox control (88)	30	51.2	64.9
PA2667	MvaT/ MvaU	MvaU	-2.68	---	---	Virulence, pyocyanin prod., motility, protease prod., biofilm formation, QS (89–91)	19	13.4	87.2
PA3083	PepN	aminopeptidase N	-4.26	-2.38	---	Virulence, competition (92)	45	100.1	41.5
PA5078	OpgG	OpgG	-4.44	-2.79	---	Stress resistance, biofilm formation (93)	14	59.5	34.9
PA3831	PepA	leucine aminopeptidase	-5.07	---	---	Virulence, protein turnover (94)	33	52.3	71.1
PA5507	PA5508	hypothetical protein	-5.56	---	---	QS (95)	12	24.3	68.2
PA0301	SpuE	polyamine transport protein	-6.02	---	---	TSS3 regulation, virulence (96,97)	12	40.1	55.9
PA0423	PasP	lipocalin protein family member	-6.04	---	---	Antibiotic resistance, virulence (98,99)	17	20.8	73.3
PA1847	NfuA	NfuA	-7.77	---	---	Stress resistance, iron homeostasis, virulence (100,101)	8	21.1	47.9
PA3470	PA3470	NUDIX hydrolase	-105969.67	---	---	Antibiotic & stress resistance, pyocyanin prod (102)	3	16.7	13.2
PA3115	FimV	motility protein FimV	---	---	-1.40	Motility, T2SS (103,104)	23	96.9	27.4
PA4427	SspB	stringent starvation protein B	---	---	-5.65	Antibiotic resistance, virulence Iron homeostasis, QS (105)	3	14.5	47.4
PA4525	PilA	type 4 fimbrial precursor PilA	---	---	-13.11	Virulence, biofilm formation, motility (58,106)	1	15.5	19.5
PA2927	PlaB	phospholipase B	---	---	-23843.67	Biofilm formation (63)	4	49.5	5.2

Blue shading represents proteins increased in abundance or unique to hypoxia. Dark blue: ≥5-fold increase in abundance; medium
blue: 2- to 5-fold increase in abundance, light blue: 1.5- to 2-fold increase in abundance. Orange shading represents proteins
decreased in abundance in hypoxia or absent in hypoxia. Dark orange: ≥5-fold decrease in abundance; medium orange: 2- to 5-fold
decrease of abundance; light orange: 1.5- to 2-fold decrease in abundance.

[a]Gene ID – *P. aeruginosa* PAO1 gene locus ID.

[b]Proteins not significantly altered in abundance between given conditions are marked as "---"

[c]Relevant protein function assigned by literature searches, UniProt Database or Pseudomonas Genome Database.

[d]NP - Unique peptides: The number of peptide sequences unique to a protein

[e]MW - Molecular weight as determined by Q-Exactive LC-MS using *P. aeruginosa* PAO1 reference (kDa)

[f]SC - % sequence covered by matching peptides for the identified protein

We examined whether the alterations in protein abundance associated with antibiotic

resistance affect affected antibiotic susceptibility of adapted cultures. Changes in sensitivity to a

wide range of their antibiotic classes were observed in both HACs and NACs compared to the early

infection strain (Fig 4A and 4B). However, both H1 and H2 populations clearly showed greater

antibiotic resistance than NACs or ancestral cultures, correlating with the relative changes in protein

abundance. Increased resistance to fluoroquinolones, macrolides, cephalosporins, carbapenems

and penicillins was observed in both H1 and H2 cultures, together with decreased resistance to

aztreonam. Interestingly, the H3 culture only showed an increased resistance to aztreonam with no

change in susceptibility to any other antibiotic tested. All strains were resistant to tetracycline.

Figure 4. Effects of long-term hypoxia adaptation on antibiotic resistance and stress resistance. **A.** Graphs
represent the antibiotic susceptibility of early infection control strain AMT 0023-30, late infection strain,
AMT 0023-34, and 28-day adapted populations using diffusion discs, tested in normoxic conditions. Bars
represent the mean inhibition zone diameter. The data represent three biological replicates and error bars
represent the standard deviation. The statistical significance was determined with the nonparametric
Kruskal-Wallis test in combination with uncorrected Dunns multiple comparisons test (*p< 0.05; **p< 0.01;
***p<0.001; ****p< 0.0001). **B**. Summary of antibiotic resistance to all tested antibiotics. **C.** Determination
of stress resistance with droplet assays. Representative pictures of colonies from serial dilutions of each
strain plated on agar with the specific stress factor added. The experiments were performed in normoxic
conditions at least three times for each stress type with the early infection strain as a control. **D.** Table
summarising changes in stress sensitivity. ↑ - increased resistance; ↓ - decreased resistance; = no significant
difference in resistance; R – no inhibition zone observed. E: early infection clinical strain; L: late infection
clinical strain; N1, N2, N3: 28-day normoxia-adapted populations; H1, H2, H3: 28-day hypoxia adapted
populations.

Hypoxia-adapted cultures show changes in stress response

Many proteins related to stress responses showed changes in abundance in response to long-term

exposure to hypoxia (Table 4), including the osmotic stress response associated protein, OpgG, and

the oxidative stress response associated protein, PA3723. The hypothetical protein, PA5139

previously associated with stress resistance, and lactate dehydrogenase LldD, associated with

oxidative stress resistance were detected in the ancestral strain and HACs, but not any of the NACs.

To examine the impact of these alterations on the phenotype of each of the individual

populations, droplet stress plates were prepared to test a range of stresses described as hallmarks

of the CF lung environment. No clear pattern of resistance to high temperature, osmotic or acidic

stress between the HACs and NACs was observed (Fig 4C and 4D). Interestingly, increased resistance

to oxidative stress was observed in both H1 and H2 cultures and in the N3 culture, which were also

the only populations where SCVs had been detected on day 28, and where increased biofilm

production and exopolysaccharide production were observed.

360   Hypoxia-adapted cultures show increased biofilm formation and
361   exopolysaccharide production

Typically, anaerobic conditions promote biofilm formation, with greater numbers of live bacteria,

increased thickness, and more compact structures (107). In this work, changes in the abundance of

proteins associated with biofilm and/or exopolysaccharide production were also observed in

response to exposure to 6% oxygen (Table 3 and Table 4). Two proteins associated with biofilm

associated antibiotic resistance were increased in abundance, while three were decreased.

Additionally, some of the proteins associated with biofilm formation or EPS production were not

detected in at least one of the groups: exopolysaccharide export protein, PelC, was identified solely

in HACs but not in NACs or ancestral strain (Table 3); solute-binding protein, PA5139, is present in

the ancestral strain, maintained in HACs, while undetectable in NACs; the polymer-forming

cytoskeletal protein PA0664 is detected in the ancestral strain, and in two HACs, but none of the

NACs; phospholipase PlaB was detected in two replicates of the ancestral strain, detected in all of

the NACs, but none of the HACs.

To confirm the impact of these proteome changes on biofilm and exopolysaccharide

production, these phenotypes were examined in each individual adapted culture (Fig 5A and 5B).

Both H1 and H3 populations showed a significant, dramatic increase in biofilm formation as

compared to the early infection strain after 24 h (6.1- and 4.3-fold increases respectively).

Significant increases in biofilm production were also observed in the late infection strain (1.6-fold),

H1 (3.9-fold) and H2 (1.4-fold) over the ancestral strain at 48 h, and this trend persisted at 72h (Fig

5A). The increased biofilm production of the H3 population was transient and reverted to AMT

0023-30 early infection strain levels by 48 h. Congo Red staining revealed darker colonies for H1 and

H2 populations which were more wrinkly than in the other strains indicating a higher production of

exopolysaccharides (Fig 5B). Interestingly, a similar phenotype was observed in culture N3, again

suggesting that this phenotype is not solely dependent on oxygen availability.

Figure 5. Effects of long-term hypoxia adaptation on biofilm formation, exopolysaccharide production and
siderophore production. A. Biofilm formation capability of hypoxia-adapted cultures determined with the
crystal violet assay after 24, 48 or 72h of static incubation on a 96-well plate in normoxic conditions. Data
represent the mean value of the OD_{590}/OD_{600} ratio from three separate experiments with at least six
technical replicates for each strain. Statistically significant difference by Kruskal-Wallis one-way ANOVA test
compared to the ancestral, early infection strain (E) with an uncorrected Dunn's test (**p< 0.01; ***p<0.001;
****p< 0.0001). **B.** Exopolysaccharide production by the adapted cultures. The photographs show colonies
of different populations of bacteria. The change of colour to a darker shade and a wrinkly colony surface
indicate a higher exopolysaccharide production. **C.** Siderophore production represented as the mean of
siderophore units produced by each colony forming unit, tested in normoxic conditions. Statistically
significant differences by ordinary one-way ANOVA test compared to the ancestral, early infection strain
with an uncorrected Fishers LSD test are shown (*p<0.05; **p< 0.01). E: early infection clinical strain (AMT
0023-30); L: late infection clinical strain (AMT 0023-34); N1, N2, N3: 28-day normoxia-adapted populations;
H1, H2, H3: 28-day hypoxia adapted populations.

Adaptation to hypoxia affected iron acquisition and homeostasis

Proteomic analysis showed changes in the abundance of proteins related to iron homeostasis and

acquisition pathways (Table 4). CyaY protein involved in iron-sulphur cluster assembly was

undetected in HACs but was present in the ancestral strain and maintained in the NACs (Table 3).

NfuA, previously associated with iron homeostasis was decreased in abundance by nearly 8-fold

relative to the ancestral strain (Table 4). In contrast, the stringent starvation protein SspB was

increased in abundance over 5-fold in HACs compared to NACs.

Differences in iron acquisition and siderophore levels of all individual populations showed a

consistent, clear decrease in siderophore production in all HACs by 2.8- to 3.6-fold but no significant

change in NACs (Fig 5D). No significant differences were detected in the pyochelin and pyoverdine

production pathways, but the PchR transcriptional regulator was undetectable in all the replicates of

the ancestral strain, two HACs and present in all NACs. Moreover, a 4-fold decrease in siderophores

was also observed between the early infection strain (ancestral) and late infection strain.

Adaptation to hypoxia induces changes in motility

A common adaptation associated with P. aeruginosa chronic infection isolates is loss of motility.

Multiple proteins associated with motility showed changes in abundance in the HACs. Flagellar hook

protein FlgE, and spermidine dehydrogenase SpdH were undetectable in the HACs while present in

the ancestral strains and NACs (Table 3). The two-component response regulator, PilR, and serine

hydroxymethyltransferase ShrA were increased in abundance in HACs (4.5 and 22.0-fold

respectively), but not in NACs compared to the ancestral strain. In contrast, DNA-binding protein

MvaT was decreased in abundance by 2.7-fold. Additionally, type 4 fimbrial precursor PilA, and FimV

were both decreased in abundance (13- and 1.4 fold respectively) in HACs vs NACs. Notably, many

of these proteins are also associated with biofilm formation and/or antibiotic resistance (ShrA,

MvaT, PilA, FlgE).

To elucidate the effect of hypoxia on motility further, the motility of the individual cultures

was assessed. The late infection strain showed a 2-fold reduction in swarming (Fig 6B) and a

moderate reduction in twitching motility (1.3-fold) (Fig 6C), while H1 and H2 showed an increased

swimming (1.5 and 1.2-fold) compared to the early strain (Fig 6A). All forms of motility were

decreased in H3 by over 2.4-fold (Fig 6A, 6B, 6C). A similar pattern was observed in culture N3, with

a 1.6-fold decrease in swimming motility and a 1.9-fold decrease in twitching motility (Fig 6A, 6C).

The pattern of swarming motility was clearly different between NACs and HACs. N1 and N2

populations showed classic swarming patterns with clearly distinguishable dendrites despite

showing no changes in diameter (Fig 6B). N3 showed a different pattern, forming colonies with

smooth edges. The H1 and H2 colonies on swarm plates were clearly distinguishable from the N1

and N2 counterparts, with no clear dendrites being formed (Fig 6). Overall, exposure to hypoxia led

to clear changes in strain motility.

Figure 6. Effects of long-term hypoxia adaptation on *P. aeruginosa* motility and virulence in the *Galleria*

***mellonella* model. A.** Changes in swimming motility**. B.** Changes in swarming motility. **C.** Changes in

twitching motility. For each type of motility, a graph representing the mean of diameter of the colony with

bars representing data from at least three separate experiments conducted in normoxic conditions, with at

least two technical replicates. Statistically significant difference by Kruskal-Wallis one-way ANOVA test

compared to the ancestral, early infection strain (E) with an uncorrected Dunn's test (*p< 0.05; **p< 0.01;

***p<0.001; ****p< 0.0001). **D.** Graph representing the mean CFU required to reach a lethal dose leading to

50% mortality (LD50) within 24h of the experiment from at least three independent experiments. A touch to

the larvae head was used to determine if they are alive. Statistically significant difference by Kruskal-Wallis

one-way ANOVA test compared to the ancestral, early infection strain (E) with an uncorrected Dunn's test is

presented (*p< 0.05). **E.** A representative image showing the *G. mellonella* larvae melanisation. Healthy

larvae with no disease symptoms are a bright cream colour. Larvae showing a slightly beige colour show first

symptoms of infection, larvae with dark spots/stipes are clearly affected and the severely melanised larvae

are in the late stages of the infection or dead. E: early infection clinical strain AMT 0023-30; L: late infection

clinical strain AMT 0023-34; N1, N2, N3: 28-day normoxia-adapted populations; H1, H2, H3: 28 day hypoxia

adapted populations.

Effects of hypoxia-adaptation on *in vivo* virulence

Previously we showed that three different late infection CF isolates were less virulent in the *G. mellonella* in vivo model than their earlier infection counterparts (37). HACs showed that adaptation to hypoxia altered the abundance of several proteins involved in virulence, and many virulence-associated phenotypes, including SCV formation, stress and antibiotic resistance, toxin and LPS production, and motility. As mentioned, phenazine biosynthesis protein PhzE was unique to HACs; Iron(II)/α-ketoglutarate-dependent oxygenase AmbC involved in toxin production was undetectable in the ancestral strain and two of the NACs while present in all HACs. The earlier mentioned two component system proteins GacS and ColR were undetectable in all HACs while present in two replicates of the ancestral strain and all NACs (Table 3). Additionally, virulence factors, leucine aminopeptidase PepA and lipocalin protein PasP, were decreased in abundance in HACs (Table 4). Unexpectedly, no differences in virulence were observed *in vivo* in *G. mellonella* between the ancestral, early infection strain and the HACs (Fig. 6D and 6E), showing hypoxia is unlikely to drive this adaptation.

Discussion

Understanding the complex network of pathway interactions and the evolutionary pressures *P. aeruginosa* undergoes in different environments is the key to understanding its biology and the mechanisms behind chronic infection, thereby, facilitating the development of effective therapeutics. Hypoxic niches are abundant in the CF lung due to increased mucus thickness, obstructive mucus plugging and increased oxygen consumption by epithelial and immune cells (28,108,109). Studies have shown that microaerobic respiration is the predominant mode of *P. aeruginosa* growth in the CF lung (110,111), with two transcription factors, Anr and Dnr, tightly regulating the response to changes in oxygen tension (29,71,112,113).

While the impact of short-term severe hypoxia and anoxia on *P. aeruginosa* has been

previously described (114,115), the impact of moderately-low hypoxia stress has not been

examined over extended periods. To enable an in-depth phenotype and proteome analysis of

adapted cultures, we restricted the study to three replicates in each condition. This focused

approach allowed us to demonstrate that prolonged low oxygen conditions alone can induce stable

changes to the proteome and phenotype, many of which are typically associated with chronic

infection, and thus may facilitate the transition from acute to chronic infection.

Diverse patterns of adaptation to long-term hypoxia

We observed two distinct patterns of hypoxia adaptation in *P. aeruginosa*, with H1 and H2 adapting

differently from H3 (Fig. 7). This is consistent with reports on evolution of different characteristics

under the same environmental pressures, including bacterial populations isolated from the same

lung niche, or in different regions of a biofilm, and enables bacterial competitiveness (21,116–120).

Flynn et. al (2016), using a similar experimental evolution approach compared three biofilm cultures

and three planktonic cultures of *P. aeruginosa* PA14 over 90 days and determined that diverse

phenotypes appeared over time with different abilities to persist between replicate cultures, and

also between colonies isolated from the same population (117). Furthermore, the various colony

morphologies observed in their study correlated with temporal variations in c-di-GMP levels (117),

which aligns with the changes in the abundance of proteins associated with the response to c-di-

GMP in HACs. We will evaluate the potential genetic changes that may have led to this in a future

study.

Two of the HACs in this study (H1 and H2) showed consistent alterations across multiple

phenotypes, while the third population displayed only slight alterations relative to the ancestral,

early infection strain and was closer in phenotype to the NACs (Fig. 7). This pattern aligns with

earlier observations that proteome changes caused by short term hypoxia response were strain-

specific (115), suggesting that different adaptation patterns can occur as the population diversifies

over the longer term. Given the diversity of *P. aeruginosa* isolates within patients and between

patients, it should be noted that the late infection strain which was examined in parallel in

phenotyping experiments represents one branch within a potentially more complex evolutionary

tree in that patient. Further genetic characterisation of the populations and specific subpopulations

will provide additional information on diversity.

Figure 7. Patterns of adaptation to long-term hypoxia. Summary of the phenotypic changes of adapted
cultures tested in normoxic conditions compared to the ancestral strain AMT 0023-30 (E) and the late
infection strain AMT 0023-34 (L). It should be noted that L is a single clinical strain and not representative of
all chronic infection isolates or strains. ↑ - indicates an increase; ↓ - indicates a decrease; = indicates no
significant difference in resistance; L: late infection clinical isolate; N1, N2, N3: 28-day normoxia-adapted
populations; H1, H2, H3: 28-day hypoxia adapted populations. (121).

Adaptation of *P. aeruginosa* to long-term hypoxia is consistent with changes observed during chronic infection

When oxygen availability decreases, electron transport chain efficiency also decreases, leading to

NADH accumulation and an increased NADH/NAD$^+$ ratio (122), creating a more reducing

intracellular environment (defined as redox stress) which must be mitigated to maintain cellular

function and viability. This is regulated by c-di-GMP signalling, which plays a central role in

mediating low oxygen responses and stress resistance in *P. aeruginosa* (123,124). It facilitates

metabolic switching in response to environmental stimuli, including the transition from a planktonic

to a biofilm lifestyle (24,124,125) and regulates multiple phenotypes such as EPS production,

motility, pyocyanin production, and antibiotic resistance (126–128).

Based on previous studies (113,115,129,71), we expected decreased oxygen availability

would lead to an increased abundance of the Anr regulator. However, in contrast, Anr was not

detected in any of the HACs, although it was present in the ancestral strain. Previous studies

investigating the Anr regulon were performed under either anoxia or severe hypoxia (0.2-2% O_2),

rather than the moderately-low hypoxia used here. There are three possible explanations: 1) the

oxygen level may be too high for Anr activity; 2) there are other levels of regulation of Anr as yet

undiscovered; 3) hypoxia adaptation may have selected for strains that downregulate Anr to reduce

energy costs. This is important in the context of the CF lungs, where oxygen gradients from anoxia

to physiological oxygen are present, suggesting more variable responses and adaptation

mechanisms than assumed in previous studies.

Given the absence of Anr in HACs, alternative regulatory pathways are likely, as exemplified

by >20-fold increase in abundance of the serine hydroxy methyltransferase, ShrA in HACs. Deletion

of *shrA* was previously shown to lead to increased c-di-GMP levels, emergence of SCVs, increased

biofilm, EPS and pyocyanin production, decreased swarming motility and suppression of iron

acquisition genes and denitrification operons (Fig 8A) (73) which align with many of the phenotypes

identified in response to long-term hypoxia. We propose that the redox imbalance caused by

hypoxia increases the abundance of ShrA *via* a yet unknown mechanism.

Figure 8. Suggested mechanism of regulation of c-di-GMP levels by 6% oxygen hypoxia exposure. A.
Mechanism of regulation of c-di-GMP levels by ShrA and the Gac/Rsm system system in normoxic conditions
based on available literature. When NADH/NAD+ ratio is balanced, ShrA activity leads mostly to the
production of (6S)-THF, which leads to limited SAM production and WspA methylation. This leads to a low
production of c-di-GMP by WspR. **B.** Proposed mechanism of regulation of c-di-GMP levels by ShrA and the
Gac/Rsm system in hypoxic conditions based on the proteomic analysis and phenotype studies presented in
this work. An increased NADH/NAD+ ratio may lead to a shift ShrA activity via an yet unknown mechanism,
which leads to an increased (6S)-5,10-CH_4-THF production, increasing SAM production and WspA
methylation. This leads to a increased production of c-di-GMP by WspR. Anr: anaerobic regulator; GacA:
response regulator; GacS: sensor kinase; LasI: acyl-homoserine lactone (AHL) synthase; LasR: Las quorum
sensing system transcriptional regulator; MetF: methylenetetrahydrofolate reductase; MetH: methionine
synthase; MetK: S-adenosylmethionine synthetase; MetY: homocysteine synthase; MvaT: hitone-like nucleoid
structuring global transcriptional regulator; RhlE2: Dead-boxRNA helicase; RsmA: RNA-binding regulator of

secondary metabolites A; RsmY: rsm system small regulatory RNA Y; RsmZ: rsm system small regulatory RNA Z; SadC: suppressor of alginate production diguanylate cyclase; ShrA: serine hydroxymethyltransferase; WspA: methyl-accepting chemotaxis protein; WspC: methyltransferase; WspE: histidine kinase; WspF: methylesterase; WspR: diguanylate cyclase; THF: tetrahydrofolate; (6S)-THF: (6S)-tetrahydrofolate; (6S)-5,10-CH_4-THF: (6S)-5,10-methylenetetrahydrofolate; SAM: S-adenosylmethionine; SAH: S-adenosylhomocysteine. Green arrows indicate upregulation or activation. Red arrows indicate downregulation or inactivation. Solid lines define literature described relationships. Dashed lines define hypothetical relationships. Green boxes indicate proteins increased in abundance in the proteomic analysis performed as a part of this work. Red boxes indicate proteins decreased in abundance in the proteomic analysis performed as a part of this work.

ShrA catalyses the reversible conversion of glycine and (6S)-5,10-methylene-THF ((6S)-5,10-CH_4-THF) to serine and (6S)-THF, with a nearly threefold preference for the latter products (73). Besides its role in c-di-GMP biosynthesis regulation, 5,10-CH_4-THF regulates folate and nucleotide biosynthesis and amino acid metabolism. However, as these findings were identified under normoxic conditions, ShrA's role in hypoxia remains unclear and warrants further investigation. In particular, its reaction flux may be influenced by factors such as the increased $NADH/NAD^+$ ratio observed under hypoxic conditions. Given that ShrA contributes to restoring redox balance by reducing the $NADH/NAD^+$ ratio, its elevated abundance is likely to be beneficial under hypoxic stress. The (6S)-5-CH_3-THF synthesized by ShrA is further metabolized through several steps to produce S-adenosylmethionine (SAM), which serves as methyl group donor for the methylation of WspA (73). Methylated WspA activates WspR and c-di-GMP synthesis (130). Thus, a shift towards synthesis of (6S)-5-CH_4-THF would increase the methylation of WspA, leading to increased levels of c-di-GMP under hypoxic conditions (Fig 8B). We acknowledge that ShrA regulation in hypoxia may be more complex and requires further investigation, as it has been previously shown to be induced by homocysteine, repressed by SAM, and activated via quorum sensing and PpyR (131,132).

Sensory histidine kinase GacS is also involved in c-di-GMP regulation. Under non-stress conditions, GacS levels are naturally low or undetectable (61), however, when GacS is absent, less GacA is phosphorylated, leading to reduced expression of RsmY and RsmZ sRNAs (133). This results

in a larger pool of active RsmA, increased repression of target mRNAs, and ultimately, lower c-di-

GMP levels. Unexpectedly, we detected GacS in the ancestral strain and NACs, but not the HACs

which is possibly due to the activity of regulators of RsmZ/RsmY sRNAs. Histone-like protein MvaT

reduces c-di-GMP levels by binding to the upstream regions of rsmY/rsmZ repressing their

transcription, increasing RsmA levels and repressing the activity of diguanylate cyclases (90).

Additionally, the RhlE2 reduces RsmY and RsmZ stability, thus promoting RsmA sequestering and

allowing the production of c-di-GMP (72). In HACs we observed comparable >2.5-fold decreases in

the abundance of MvaT, and RhlE2 compared to the ancestral strain, which was not observed in

NACs. Therefore, we speculate that the presence of GacS in NACs and ancestral strain is overcome

by the activity of RhlE2 and MvaT coupled with the decreased production of c-di-GMP by WspR.

Taken together, our findings suggested that the observed changes in abundance of ShrA,

GacS, RhlE2 and MvaT lead to an increase in cellular c-di-GMP levels, consistent with the observed

phenotypes including increased biofilm and EPS production, emergence of stable SCV phenotypes,

alterations in motility, antibiotic resistance and pyocyanin production which could also be

associated with this signalling molecule. Thus, we hypothesised that while Anr primarily responds

under anoxic conditions or severe hypoxia, other proteins, including ShrA, may regulate relevant

pathways under moderately-low O_2 levels after long-term exposure to hypoxia. However, we have

not observed any changes in the intracellular concentrations of c-di-GMP levels between the

ancestral early infection *P. aeruginosa* strain and any of the HACs and NACs or the late infection

strain despite the observed changes in related phenotypes (S6. Figure). This was not unexpected as

Yan et al. (2017) previously investigated the correlation between c-di-GMP levels, motility, and

biofilm formation (134) and did not observe any direct correlation, despite using well-established

models describing the mechanisms of c-di-GMP action. This finding suggests a more complex

regulatory network, where the final phenotypic outcome depends on the cumulative effect of

multiple small-scale changes rather than a straightforward 1:1 correlation. Such complexity could

potentially explain the seemingly contradictory observations in c-di-GMP concentrations, regulator

protein abundances, and variation in motility and biofilm formation.

Interestingly, the adaptation of *P. aeruginosa* to long-term hypoxia mirrors the phenotype

and molecular changes previously reported for *P. aeruginosa* chronic infection isolates

(8,18,21,23,36,65,68), including increased SCV prevalence, increased biofilm and exopolysaccharide

production, increased antibiotic resistance and decreased siderophore production. Importantly, all

these changes are stable, because all phenotypes were examined in normoxic conditions; thus, they

are a result of adaptation to hypoxia, rather than a plastic temporary response to hypoxic

conditions.

While a potential limitation of our experiment is that growth in relatively rich laboratory

media does not represent the conditions in the lung, the fact that we can observe changes

resembling adaptation to chronic infection even in the absence of nutrient stress, highlights a clear

role for hypoxia in this process. Choosing a planktonic model for this study allowed us to balance

the feasibility of the experimental design and the relevance of the model in the clinical context but

considering the obtained results we argue that this approach effectively reveals the potential role of

long-term hypoxia in driving adaptation during chronic infection.

Two distinct SCV types emerged in the long term cultures which were more prevalent in the

HACs, and one of the morphotypes was exclusive to hypoxia. We observed a temporary emergence

of SCV 1 in the NACs, which was not positively selected in normoxia, possibly due to the lack of an

adaptive benefit long-term. Under optimal laboratory conditions, long-term cultivation can induce

stress due to factors such as high population density, accumulation of metabolic byproducts,

nutrient gradients leading to transient SCV arising in the population. *P. aeruginosa* also possesses

phase variation systems that allow spontaneous phenotypic switching, generating SCVs even in the

absence of external stressors (135,136). These variants may be maintained at low frequencies,

preparing the population for potential environmental shifts (135). The abundance of c-di-GMP

signalling pathways proteins, Wsp, and YfiBNR, typically associated with SCVs (41,135), was

unchanged in the HACs, but as mentioned earlier, the ShrA protein indirectly regulates the Wsp

system activity. Additionally, the inactivation of *gacS* has been shown to be associated with the

emergence of SCVs under stress conditions and suppression of the reversion of this phenotype to a

normal colony variant (62). The latter may explain why the SCV populations in the NACs were

transient, as GacS is expressed in the NACs but not HACs. SCVs are strongly associated with

pathogenicity and persistence among several bacterial pathogens (137–139), and are prevalent

among chronic *P. aeruginosa* isolates (137,138,140). Specifically, SCVs of *P. aeruginosa* correlate

with increased exopolysaccharide and biofilm production, antimicrobial resistance, stress resistance

and host immune responses (41,140). Their emergence has been linked to redox imbalances which

can be caused by low oxygen conditions (136), although other environmental stressors or co-

colonisation with other species also contribute to their increased prevalence (137,140,141). This

confirms directly, for the first time, that prolonged hypoxia exposure is associated with increased

SCV occurrence in *P. aeruginosa* populations. Identification of the factors that contribute to SCV

emergence is crucial due to the challenges they pose in clinical settings (137) and recognising the

environmental factors that select this phenotype may help us limit their occurrence. Both HACs that

showed a high SCV prevalence (H1, H2) also showed increased resistance to a broad spectrum of

antimicrobials together with increased biofilm and exopolysaccharide production. Surprisingly, no

trends in motility relating to hypoxia exposure or biofilm formation were observed, although some

changes in patterns of swarming colonies were noted. Further investigation of NCVs and SCV 1 and

SCV 2 (genotypes and phenotypes) will provide insight on the contribution of the specific

subpopulations to the observed changes in other adaptations and are ongoing.

Chronic infection *P. aeruginosa* CF isolates typically exhibit increased biofilm and

exopolysaccharide production, and reduced motility (18,120,142–144), with biofilm formation

relying on hypoxia responses due to the presence of steep oxygen gradients (30,145,146).

Moreover, the awareness of the role of biofilms in antibiotic treatment efficiency is being

increasingly acknowledged (147,148). Proteins associated with biofilm formation, exopolysaccharide

production, and motility were changed in abundance. Notably, outer membrane lipoprotein PelC,

which is essential for biofilm formation (45), was unique to HACs while in contrast, the flagellar

hook protein FlgE, typically associated with motility, host interactions and immune evasion (54), was

absent in HACs. Both are regulated by FleQ, a c-di-GMP binding transcriptional regulator, which

activates the *pel* operon and inhibits the flagellar gene expression, coordinating this lifestyle switch

(149). Two component system regulator PilR associated with increased biofilm formation, host cell

attachment, and motility (58) was increased in abundance in HACs. However, the PilR-activated type

IV fimbrial precursor, PilA, was decreased in abundance by >13-fold in HACs compared to NACs

which suggests alternative regulation mechanisms are at play. Further highlighting this complexity,

phospholipase PlaB, showing activity towards exogenous lipids in the biofilm matrix (63), was

absent in HACs while UDP-glucose dehydrogenase PA3559, part of the EPS biosynthesis pathway

(74), was increased in abundance in HACs. These changes collectively indicate an adaptive shift

towards a biofilm lifestyle in HACs, aligning with changes observed in chronic infection adaptation.

Antibiotic resistance is a major hallmark of *P. aeruginosa* infections in CF (150), and short-

term hypoxic and anaerobic conditions have been previously shown to increase *P. aeruginosa*

antibiotic resistance (151–153). We demonstrate that long-term exposure to hypoxia induces stable

changes in antibiotic resistance, which importantly, didn't revert during in normoxic conditions used

for phenotyping. Two of the HACs showed increased resistance to a wide range of antibiotics from

different classes, further supporting the evidence of the extent of the influence of hypoxia pre-

adaptation on drug resistance and the wide range of diverse mechanisms affected. Interestingly,

high c-di-GMP levels have been shown to corelate with increased resistance to antibiotic,

independently form the biofilm/ planktonic lifestyle (128). Several proteins involved in lipid A/LPS

biosynthesis (LpxA, LpxB, WaaG), membrane stability (DsbG, ColR, OprH), and peptidoglycan

biosynthesis (MurC, PA3262), many of which have been previously linked to antibiotic resistance

(154–158) were altered in abundance. These may contribute to the observed broad-spectrum

antibiotic resistance by affecting cell envelope permeability and structure. Moreover, other proteins

linked to antibiotic resistance including phosphate kinase Ppk2, previously linked to multidrug

resistance (84), and DNA topoisomerase IV subunit ParC involved in fluoroquinolone resistance

(83,159) were increased in HACs (106).

The airways of people with CF are reported to show an increased abundance of available

iron (160,161). *P. aeruginosa* expresses many iron acquisition systems including siderophore

production, haem acquisition systems and complexes importing iron-bound citrate or catechol

(162). Importantly, a strong positive correlation between increased sputum iron and chronic *P.*

aeruginosa infections has been shown (163). Furthermore, iron uptake and homeostasis are

regulated by the c-di-GMP through the Gac/Rsm cascade (164), in line with other alterations

observed and underlines the role of hypoxia leading to changes in iron acquisition on chronic

infection development, which led to recognising them as therapy targets for *P. aeruginosa*

treatment (161,165). Surprisingly, we didn't observe any alterations associated with pyochelin or

pyoverdine biosynthesis and transport, although a clear decrease in siderophore production was

observed in HACs. Notably, the PchR regulator was not detected in the ancestral strain, nor in two of

the HACs, while present in all of the NACs, suggesting alternative mechanisms of control of

siderophore production or transport. Iron-sulphur cluster proteins are essential for maintaining iron

homeostasis and regulating diverse cellular functions (166). Notably, two iron-cluster proteins

frataxin-like protein CyaY, and iron-sulphur cluster scaffold protein NfuA were absent or decreased

in abundance in HACs, respectively. Additionally, the RhlE2 RNA helicase mentioned earlier has been

shown to regulate expression of iron acquisition genes. These changes may explain the decreased

siderophore levels observed, but require further investigation; notably, they reflect the

downregulation of siderophore production by chronic infection isolates from CF lungs (167).

Adaptation of *P. aeruginosa* to chronic infection is frequently associated with reduction in

expression of multiple virulence factors including pyocyanin, proteases, lipases, urease, elastase and

secretion systems (18). Moreover, exposure to hypoxia attenuated the virulence of isolates from

acute, but not chronic infections (168) and decreased their interactions with host cells,

demonstrating the importance of this environmental pressure. Likewise, proteomic analysis of

HACs, showed a decrease in abundance of many virulence factors: proteases (PasP serine protease,

aminopeptidase PepA, aminopeptidase PepN), Phospholipase B, and type VI secretion system

(TssB1, Tse8 effector protein) and type II secretion system effector proteins (PasP, PilA). LasI, a

component of the QS system was not detected in HACs or the ancestral strain, but present in NACs.

Since it positively regulates protease and lipase synthesis (and indirectly T6SS), its decreased

abundance is consistent with the other alterations. Interestingly, phenazine biosynthesis protein

PhzE was unique to HACs, which would suggest an increase in pyocyanin production in all HACs,

however, we detected a change in pyocyanin in only two cultures. Thus, we might have expected

decreased virulence in HACs, but did not observe any change in the *G. mellonella* infection model

following hypoxia exposure. However, this acute infection model may not be optimal to assess the

impact of long-term hypoxia adaptation on chronic infection.

Some of the observed changes in phenotypes, including increased biofilm and EPS

production, and antibiotic resistance may result from the increased prevalence of the SCVs in the

population, as all of these are altered in HAC 1 and 2, while limited changes were observed in HAC

3. Investigation of this is ongoing, however, given that many alterations in the proteome were conserved across all three HACs, despite divergent phenotypic outcomes suggests a more complex explanation involving additional, SCV-independent adaptive mechanisms.

A recent study defined *P. aeruginosa* 224 genes as pathoadaptive, and demonstrated that specific gene expression was associated with a host preference towards people with CF (169), many of which encode proteins altered in response to long-term hypoxia. Three of these, ColR, GacS and MexS were not detected in HACs, while another, PA0664 protein was maintained in HACs while not detected in NACs. Additionally, PilR and ParC were increased in abundance in HACs respectively, while OprM was decreased in abundance. Moreover, overall changes in the abundance of 15 proteins encoded by genes that were associated with CF host preference (including *oprM*, *pepA*, *fimV*, and *pilR*) and 11 that were associated with non-CF infections (including *mvaT* and *tssB1*) (169), suggest long-term hypoxia may play a role in *P. aeruginosa* infection both in CF and non-CF cohorts.

Common adaptation mechanisms of opportunistic pathogens to hypoxia

Several oxygen-sensing mechanisms, homologous genes in hypoxia regulons and several phenotypes altered in chronic infection are shared across other opportunistic bacteria causing pulmonary infections. This suggests that adaptation to prolonged hypoxia could be a shared mechanism between many low oxygen and/or microaerophilic habitats and not limited to human infections. These environments are increasingly abundant in water and soil, where opportunistic bacteria reservoirs exist, and play significant roles in shaping their physiology, contributing to their increased prevalence (170–174). As many opportunistic bacterial pathogens causing CF infections reside in these niches, common adaptations are not unexpected.

Proteome analysis of HACs showed an increase in abundance of only two proteins encoded

on the Anr regulon (113), the Ppk2 and the AdhA alcohol dehydrogenase while NorC was not

detected in HACs but present in NACs and the ancestral strain. Only three proteins from the Anr

core regulon being altered is surprising, since hypoxia response has been primarily associated with

it. However, as previously mentioned, Anr regulation is complex, and hypoxia adaptation may have

selected for strains that downregulate Anr to conserve energy.   Additionally, as mentioned the

moderate hypoxic conditions utilised may lead to adaptation via other pathways rather than the Anr

regulon. Notably, the Anr regulon shares homologous genes with the DosR regulon of

Mycobacterium abscessus and Lxa locus of *Burkholderia cenocepacia*, highlighting that aspects of

the hypoxia response may be shared across several opportunistic CF lung infections more generally.

Investigations on the impact of different levels of hypoxia on these three pathogens is ongoing in

our group. Targeting shared mechanisms of adaptation to chronic infection could present a new

opportunity for the treatment of multiple pathogens simultaneously.

Impact

Overall, we demonstrate that alterations in phenotype after long-term exposure to hypoxic

conditions shows several similarities associated with chronic infection. Further understanding of the

mechanisms by which *P. aeruginosa* adapts to hypoxia in chronic infection and the interplay with

other factors present in the lung environment will potentially enable us to prevent or treat the

switch from acute to chronic infection. This could inevitably limit antibiotic use, severity of the

disease and ultimately, decrease the socioeconomic impact of infections caused by *P. aeruginosa*.

Furthermore, the first drug to alter the human response to hypoxia was recently approved for use

by the US Food and Drug Administration (175). Thus, it is tempting to speculate that the response of

bacteria to low oxygen pressures could be targeted to prevent a switch from acute to chronic

infection.

The observed changes in proteome and phenotypes (increased biofilm formation, increased

antibiotic resistance, decreased siderophore production, increased SCV prevalence) further deepens

our knowledge on how hypoxia leads to adaptation of *P. aeruginosa*, shedding light on how

prolonged low oxygen exposure can impact on adaptation to chronic infection. Overall, we confirm

that hypoxia contributes to the development of multiple phenotypic changes that are prevalent in

chronic infection, opening a path towards designing therapies targeting these.

Materials and methods

Bacterial strains

Pseudomonas aeruginosa AMT 0023-30 early infection clinical strain (E) and AMT 0023-34 late

infection clinical strain (L) are two well characterised strains in the International *P. aeruginosa* panel

(35). Hypoxia- and normoxia-adapted *P. aeruginosa* strains were obtained in this study as described

below and are referred to as H1, H2, H3, N1, N2, and N3 (Table 1). If not stated otherwise, bacteria

were routinely grown in LB broth (Sigma-Aldrich, St. Louis, MO, USA) at 37°C with orbital agitation

(200 rpm) or plated on LB agar (Sigma-Aldrich, St. Louis, MO, USA) and grown at 37°C.

Experimental evolution

Overnight cultures of *P. aeruginosa* strain AMT 0023-30 (E) were inoculated into 30 ml LB media to

OD$_{600}$ ~0.1. Prior to use, the media was equilibrated for 18 h at 37°C in an O$_2$ Control InVitro Glove

Box (PHGB model, Coy Laboratory Products, MI, United States) at 6% O$_2$, 5% CO$_2$ or in a Steri-Cycle

CO$_2$ Incubator (model 371, Thermo Scientific, Waltham, MA, USA) at 21% O$_2$, 5% CO$_2$. Three flasks

were inoculated as biological replicates for each of the conditions being examined. Additionally,

aliquots (2 ml) of the overnight *P. aeruginosa* cultures were retained as a control for proteomic and

transcriptomic analysis. All flasks were incubated for 24h with agitation (200 rpm, CO$_2$ resistant

orbital shaker, model 15351105, Thermo Fisher Scientific, Waltham, MA, USA) in hypoxic and

normoxic conditions. Daily aliquots (2 ml) were removed for proteomic analysis from each of the

flasks and also additional aliquots retained for OD$_{600}$ measurements and plating. Glycerol stocks

(50%, LB (v/v)) were prepared for subsequent phenotype analysis and frozen immediately to ensure

minimal oxygen exposure. A volume of 5% of the bacterial culture was also transferred into 30 ml of

pre-equilibrated media each day and this procedure was continued for up to 28 days to determine

long-term adaptation patterns. To ensure stability of the hypoxic conditions, all of the samples from

H1, H2 and H3 were taken inside the hypoxia chamber. To ensure that there were no other

environmental differences between the test conditions other than oxygen levels, the same batch of

media was used for both normoxia-adapting and hypoxia-adapting flasks and shaking conditions

were consistently maintained.

Whole proteome analysis

Samples (2 ml) were collected throughout the experimental evolution for proteome analysis.

Bacterial cells were pelleted at 14000-16000 g for 2 min and stored at -80°C until further analysis.

Whole cell lysates were extracted from each of the hypoxia- and normoxia exposed cultures by

resuspending in 1 ml of ice-cold lysis buffer containing 40mM Tris-HCl (pH 7.8) and 1x cOmplete™,

EDTA-free protease inhibitor cocktail (Roche, Basel, Switzerland). Cells were disrupted by a

sonication with a sonic dismembrator and microtip 6mm probe (model 505; Fisherbrand, Thermo

Fisher Scientific, Waltham, MA, USA) (set with 20% output power using eight 10 s bursts on ice). Cell

debris and any remaining intact cells were pelleted using centrifugation (12000 g for 30 min).

Supernatants were transferred into fresh tubes and the proteins were precipitated with four

volumes of methanol and incubated in -80°C overnight. The samples were centrifuged at 12000 g

for 30 min at 4°C, supernatants were removed, and the protein pellets were resuspended in 200 μl

of resuspension buffer containing 8 M urea, 50 mM Tris-HCL (pH 8.0), before 1 M 1,4-dithiothreitol

(DTT) was added to each supernatant (10 μl/ml lysate) and incubated at 56°C for 30 min, followed

by the addition of 1 M iodoacetamide (55 μl/ml lysate) which was incubated in darkness at room

temperature for 20 min. Lysates were dialyzed in SnakeSkin™ tubing (Thermo Fisher Scientific,

Waltham, MA, USA) with a cut-off of 3.5 kDa against 100 mM ammonium bicarbonate overnight,

with stirring at 4°C, followed by at least 4 more hours incubation with fresh volumes of ammonium

bicarbonate. Trypsin (400 ng/μl) (Sigma-Aldrich, St. Louis, MO, USA) was added to the dialyzed

protein (5 μl /100 μl protein) and tubes were incubated overnight at 37°C. Aliquots from each

sample were placed in fresh tubes and samples were dried in a vacufuge concentrator system

(model 5301; Eppendorf, Hamburg, Germany) at 65°C and resuspended in 20 μl of ZipTips

resuspension buffer containing 0.5% trifluoroacetic acid (TFA) in MilliQ water. ZipTips (Merc

Millipore, Burlington, MA, USA) were used for peptide purification as per the manufacturers'

instructions. Nanodrop spectrophotometer DS-11+ (DeNovix, Wilmington, DE, USA) was used for

peptide concentration measurements. The samples were analysed on a Bruker TimsTOF Pro mass

spectrometer connected to a Evosep One chromatography system. Peptides were separated on an

8 cm analytical C18 column (Evosep, 3 µm beads, 100 µm ID) using the pre-set 33 samples per day

gradient on the Evosep one. A trapped ion mobility (TIMS) analyser was synchronized with a

quadrupole mass filter to enable the highly efficient PASEF (Parallel Accumulation Serial

Fragmentation acquisition) procedure with acquisition rates of 100 Hz. The accumulation and ramp

times for the TIMS were both set to 100 ms., with an ion mobility (1/k0) range from 0.62 to 1.46

Vs/cm. Spectra were recorded in the mass range from 100 to 1,700 m/z. The precursor (MS)

Intensity Threshold was set to 2,500 and the precursor Target Intensity set to 20,000. Each PASEF

cycle consisted of one MS ramp for precursor detection followed by 10 PASEF MS/MS ramps, with a

total cycle time of 1.16 s.

Protein identification and label-free quantitative (LFQ) analysis were conducted using MaxQuant (v

2.4.2.0, https://maxquant.org/) by searching against the reference strain *P. aeruginosa* PAO1 (ATCC

15692). For protein identification, the following search parameters were used: trypsin was the

digesting enzyme with up to two missed cleavages allowed for; oxidation on methionine and

acetylation on the N terminus were selected as variable modifications with a fixed modification of

carbamidylation on cysteine; Label Free Quantitation (LFQ) and Match Between Runs (MBR) options

were selected. FDR for protein and peptide were set at 1%.

The three biological replicates from each 28 day culture condition were analysed, with one technical

replicate for a randomly chosen sample and LFQ intensities were compared. All missing LFQ

intensities were imputed with a fixed value of 1. The mean was calculated separately for the hypoxia

and normoxia samples, followed by fold change (FC) calculation.

One-way ANOVA statistical analysis (p-value ≤0.05) together with Benjamini–Hochberg false-

discovery adjustment were performed to identify proteins with a significantly changed abundance

Proteins with adjusted p-values≤0.1 were considered significantly altered in abundance.

Functional enrichment of proteins significantly changed in abundance was performed. For GO

biological process functional enrichment analysis ShinyGO (v 0.82) (44) was used. Species was set to

P. aeruginosa PAO1, FDR cutoff was set to 0.05 and the minimal pathway size was set as two. For

STRING local network and KEGG pathway enrichment the STRING (v 12.0) database was used with

default settings.

862 Antibiotic resistance

Overnight bacterial cultures were diluted in 10 ml of fresh LB to OD_{600} equal to 0.1. Bacteria were

spread evenly on Mueller–Hinton agar plates (Neogen, Lansing, MI, USA) using cotton swabs, in

three directions, to give a homogenous lawn. Antibiotic disks (Oxoid, Hampshire, UK) were placed

on the surface of the agar and the plates were incubated overnight in normoxic conditions at

37°C. Growth inhibition zone diameters were measured in at least two points, and the mean was

used for statistical analysis. The experiment was repeated three times. The following antibiotics

were tested: norfloxacin (10 µg), ciprofloxacin (10 µg), azithromycin (15 µg), clarithromycin (15 µg),

ceftazidime (30 µg), meropenem (10 µg), imipenem (10 µg), piperacillin (100 µg), aztreonam (30

µg), tobramycin (10 µg), and tetracycline (30 µg).

874 Motility

Motility assays were performed as described previously (176). Swimming motility was assessed by

inoculating the surface of a 0.3 % (w/v) LB agar plate with overnight culture using a sterile

toothpick. Swimming and swarming plates were incubated overnight in normoxic conditions at 37°C

for ~18 h. For swarming motility 0.8% nutrient broth (Sigma-Aldrich, St. Louis, MO, USA), 0.5%

dextrose, 0.5% agar plates were inoculated identically as for swimming motility assessment. The

diameter of bacterial growth zones was measured in at least two directions, and the mean was used

for statistical analysis. Twitching motility was assessed by inoculating 1.5 % (w/v) LB agar plate by

piercing through the agar. Plates were incubated overnight in normoxic conditions at 37°C for ~18 h,

and the agar was removed. Bacteria on the surface of the plate were stained with 1% (w/v) crystal

violet for 20 min. The plates were washed with water until all excess dye was removed and dried at

room temperature. The diameter of the growth zone was measured in at least two points, and the

mean was used for statistical analysis. The experiment was repeated three times.

Biofilm production

Overnight cultures of *P. aeruginosa* strains were diluted in fresh LB media to OD_{600} of ~0.1, and

100 μl of each diluted culture was transferred into 6 wells of a 96-well plate. The plates were

incubated statically in normoxic conditions at 37°C for 24, 48 or 72 h and OD_{600} was measured with

the Synergy H1 microplate reader (Mason Technology, Dublin, Ireland). The medium with planktonic

bacteria was removed, and the biofilm fixed by incubation with 200ul of methanol for 10 min. The

wells were washed three times with phosphate buffered saline (PBS) (Sigma-Aldrich, St. Louis, MO,

USA), followed by the addition of 200 μl of 0.1% crystal violet solution and 30 min incubation at

room temperature. The solution was removed, and the wells were washed three times with water

and once with PBS. The plates were dried for at least 30 min at room temperature and 150 μl of

96% ethanol was added to dissolve the bound stain. After 10 min of incubation at room

temperature, the solution was mixed by pipetting, OD_{590} was determined, and the OD_{590}/OD_{600} ratio

was calculated. The experiment was repeated three times.

Exopolysaccharide production - Congo Red agar assay

Overnight cultures of *P. aeruginosa* strains were pelleted and washed with 1 ml of PBS. The pellets

were resuspended in PBS to OD_{600} of ~0.1 and 1 μl of the suspension was plated on a Congo Red

agar plate (10 g/L Tryptone, 10 g/ L Bactoagar, 40mg/L Congo Red, 20mg/L Coomassie Brilliant

Blue). The plates were incubated for 48 h in normoxic conditions and colony morphology was

assessed. The experiment was repeated three times.

Stress sensitivity assays

Overnight cultures of *P. aeruginosa* strains were diluted to a CFU/ml of 10^8 and a range of serial

dilutions was prepared. Suspensions (1μl) were plated on one of the following: LB agar control

plates, LB agar supplemented with 0.5 M NaCl (osmotic stress), LB agar supplemented with 1mM

H_2O_2 (oxidative stress), or LB agar, pH 5 (acidic stress). The plates were incubated in normoxic

conditions at 37°C for 24 or 48 h. Temperature stress was determined by incubation of an LB agar

plate at 45°C for 24 h. Change of sensitivity was determined by determining the lowest dilution at

which bacterial growth was observed. The experiment was repeated three independent times.

Galleria mellonella infection model

Virulence was determined as previously described (37). Wax moth larvae (*Galleria mellonella*) (The

KammieShop, Waalwijk, Denmark) were equilibrated in normoxic conditions at 15°C for at least 7

days after arrival and stored at the same conditions for up to 4 weeks, during which larvae with a

weight range of 0.2-0.4 g were selected for experiments. Overnight bacterial cultures in LB were

diluted in fresh media to OD_{600} of ~0.1 and grown for 2-3 h. Cultures were centrifuged, and

bacterial cells were washed with PBS. The bacteria were diluted to 10^6 CFU/ml and serially diluted

to 10^{-6}. Each dilution was injected (20 μl) into the hindmost left proleg of healthy larvae (10 per

group) and the same volume of PBS was injected into one group as a control. Injected larvae were

incubated at 37°C for 72 h. To have an accurate count of the bioburden inoculated, aliquots of the

bacteria injected were serially diluted and plated onto LB agar and colonies counted after 24 h. The

experiment was repeated at least three times for each strain tested.

Pyocyanin production quantification

P. aeruginosa strains were inoculated to OD_{600} in fresh PB media (20g/L Bacto-Peptone, 1.4g/L

$MgCl_2$, 10g/L K_2SO_4) and cultured with aeration in normoxic conditions at $37^{o}C$ to maximize

pyocyanin production. The bacterial cultures were pelleted (10 min at 12300 g) and two 7.5-ml

replicates were withdrawn from each of the 18 h stationary phase cultures. Pyocyanin was extracted

with 4.5 ml of chloroform and a total of 1.5 ml of 0.2 M HCl was added to each extract causing

colour change from blue to pink. The absorbance of this solution was measured at 520 nm, and the

obtained values were converted to pyocyanin content following Essar et al. (1990) (48). The

experiment was repeated at least two times.

Siderophore production

The siderophore production was quantified with a modified version of a microplate method

developed previously (177). Prior to siderophore quantification all glassware was rinsed with a 3M

solution of HCl to remove trace iron and rinsed thoroughly with deionised water. The CAS reagent

was prepared as described before (177).

Three independent cultures of each tested population or strain were grown in fresh LB medium for

48 h at $37^{o}C$ with agitation in normoxic conditions. Aliquots of the cultures were taken for CFU

counts, serially diluted and plated on LB agar. The cultures were spun down (10 min at 12300 g), and

50 µl aliquots of the supernatant were transferred into separate wells of a microplate followed by

an immediate addition of 100 µl of CAS reagent. As a control LB broth was used. The plates were

incubated for 20 min at room temperature in the dark and the optical density was measured at 630

nm using the Synergy H1 microplate reader. The Siderophore units were calculated as follows:

siderophore production = $[(A_{LB} - A_s) * 100]/A_{LB}$, where A_{LB} = absorbance of LB control and A_s =

absorbance of sample. The earlier plated bacteria were incubated at 37°C overnight and colony

forming units were counted. Siderophore production per colony forming unit was calculated to

account for any differences in growth. The experiment was repeated two times.

Determination of growth by optical density

952   To assess the influence of the inoculation volume on the growth patterns of *P. aeruginosa* AMT

953   0023-30, 24h stationary phase cultures were inoculated into fresh LB media in a way they

954   constituted 1%, 5% or 10% of the new culture. Three 200 µl technical replicates of the cultures were

955   transferred into a 96-well plate and incubated in the Synergy H1 microplate reader in normoxic

956   conditions at 37°C with agitation with OD_{600} measurements taken every 30 min.

Determination of growth by CFU

To assess the differences in the numbers of generations in hypoxic conditions and normoxic

conditions of the AMT0063-30 early infection strain overnight cultures were prepared. The

following day they were inoculated in triplicate into fresh LB media and incubated with aeration at

37°C in the O_2 Control InVitro Glove Box in 6% O_2, 5% CO_2 or in 21% O_2, 5% CO_2 or in the Steri-Cycle

CO_2 Incubator at 21% O_2, 5% CO_2. Every hour a sample was taken from each culture for OD_{600}

measurements and plating on LB agar. The agar plates were incubated overnight at 37°C and the

colony forming units were calculated and OD_{600} to CFU/ml was plotted, a trend line was determined

and R^2 calculated. The doubling time was calculated for the logarithmic phase of growth using the

following formula: Doubling time = (Duration (min)*ln(2))/[ln(final cell concentration/initial cell

concentration)].

Intracellular c-di-GMP concentration measurements:

C-di-GMP was extracted from *P. aeruginosa* cells as previously described (178). Briefly, the OD_{600} of overnight cultures was measured, and a culture volumes equivalent to 4 ml of OD_{600} = 1.8 were centrifuged and washed twice with PBS. Pellets were resuspended in 400 µl PBS, incubated at 100°C for 5 min before adding ice-cold ethanol (final concentration of 65%), and mixing. The supernatants were collected after centrifugation and the extraction was repeated three times. Pooled supernatants were dried at 30°C, resuspended in 200 µl nanopure water, and c-di-GMP levels were quantified using a cyclic-di-GMP assay kit (Lucerna Technologies, Brooklyn, NY, USA), as per the manufacturer's recommendations. C-di-GMP concentrations were normalized to total protein content, measured using the Bradford reagent (Sigma-Aldrich, St. Louis, MO, USA) on pellets that were resuspended in TE buffer, sonicated on ice for 1 min (10 s bursts, 20% output power), centrifuged and the supernatants used for protein quantification.

Statistical analysis of phenotype data

Initial data preparation was performed using Microsoft Excel and statistical analysis was performed using GraphPad Prism (v 8.0.2, GraphPad Software, Boston, MA, US). Only p-values of 0.05 or lower were considered statistically significant. First, the normality was assessed using the Shapiro-Wilk test. Data with normal distribution was analysed using the Ordinary one-way ANOVA with the Uncorrected Fisher's LSD; data with non-normal distribution was analysed using the Kruskal-Wallis test with an uncorrected Dunn's test. Details of the analysis and significance are provided in figure legends.

Acknowledgements

989 JD, CJC are supported by an SFI Frontiers of the Future award (20/FFP-P/8717) to SMcC. ND was

990 supported by the Irish Research Council, GOIPG/2021/1443 IRC Award. The authors are grateful to

991 Dr David Gomez-Matallanas for his advice in proteomic analysis and for the support of the staff and

992 facilities in the UCD Conway proteomics core.

References

1.　Antimicrobial resistance collaborators. Antimicrobial Resistance Collaborators: Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. Lancet. 2022 Feb 12;399(10325):629–55.

2.　O'Neill J. Tackling drug-resistant infections globally : final report and recommendations. Wellcome Trust; 2016.

3.　Blanchard AC, Waters VJ. Opportunistic Pathogens in Cystic Fibrosis: Epidemiology and Pathogenesis of Lung Infection. Journal of the Pediatric Infectious Diseases Society. 2022 Sep 1;11(Supplement_2):S3–12.

4.　Sanz-García F, Gil-Gil T, Laborda P, Ochoa-Sánchez LE, Martínez JL, Hernando-Amado S. Coming from the Wild: Multidrug Resistant Opportunistic Pathogens Presenting a Primary, Not Human-Linked, Environmental Habitat. International Journal of Molecular Sciences. 2021 Jan;22(15):8080.

5.　Serra-Burriel M, Keys M, Campillo-Artero C, Agodi A, Barchitta M, Gikas A, et al. Impact of multi-drug resistant bacteria on economic and clinical outcomes of healthcare-associated infections in adults: Systematic review and meta-analysis. PLoS One. 2020;15(1):e0227139.

6.　Reynolds D, Kollef M. The Epidemiology and Pathogenesis and Treatment of Pseudomonas aeruginosa Infections: An Update. Drugs. 2021 Dec 1;81(18):2117–31.

7.　Li Y, Roberts JA, Walker MM, Aslan AT, Harris PNA, Sime FB. The global epidemiology of ventilator-associated pneumonia caused by multi-drug resistant *Pseudomonas aeruginosa*: A systematic review and meta-analysis. International Journal of Infectious Diseases. 2024 Feb 1;139:78–85.

8.　Rossi E, La Rosa R, Bartell JA, Marvig RL, Haagensen JAJ, Sommer LM, et al. Pseudomonas aeruginosa adaptation and evolution in patients with cystic fibrosis. Nat Rev Microbiol. 2021 May;19(5):331–42.

9.　Smith WD, Bardin E, Cameron L, Edmondson CL, Farrant KV, Martin I, et al. Current and future therapies for Pseudomonas aeruginosa infection in patients with cystic fibrosis. FEMS Microbiology Letters. 2017 Aug 1;364(14):fnx121.

10.　Botelho J, Grosso F, Peixe L. Antibiotic resistance in *Pseudomonas aeruginosa* – Mechanisms, epidemiology and evolution. Drug Resistance Updates. 2019 May 1;44:100640.

11.　Bakkeren E, Diard M, Hardt WD. Evolutionary causes and consequences of bacterial antibiotic persistence. Nat Rev Microbiol. 2020 Sep;18(9):479–90.

12.　Sett A, Dubey V, Bhowmik S, Pathania R. Decoding Bacterial Persistence: Mechanisms and Strategies for Effective Eradication. ACS Infect Dis. 2024 Aug;

13.　ECDC. Healthcare-associated infections acquired in intensive care units - Annual Epidemiological Report for 2016. Stockholm; 2018.

14. ECDC. Healthcare-associated infections acquired in intensive care units - Annual
Epidemiological Report for 2019. Stockholm; 2023.

15. Tacconelli E, Carrara E, Savoldi A, Harbarth S, Mendelson M, Monnet DL, et al. Discovery,
research, and development of new antibiotics: the WHO priority list of antibiotic-resistant
bacteria and tuberculosis. The Lancet Infectious Diseases. 2018 Mar 1;18(3):318–27.

16. WHO. WHO bacterial priority pathogens list, 2024: Bacterial pathogens of public health
importance to guide research, development and strategies to prevent and control
antimicrobial resistance. 2024.

17. Cullen L, McClean S. Bacterial Adaptation during Chronic Respiratory Infections. Pathogens.
2015 Mar;4(1):66–89.

18. Jurado-Martín I, Sainz-Mejías M, McClean S. Pseudomonas aeruginosa: An Audacious
Pathogen with an Adaptable Arsenal of Virulence Factors. Int J Mol Sci. 2021 Mar
18;22(6):3128.

19. Camus L, Vandenesch F, Moreau K. From genotype to phenotype: adaptations of Pseudomonas
aeruginosa to the cystic fibrosis environment. Microbial Genomics. 2021;7(3):000513.

20. Chadha J, Harjai K, Chhibber S. Revisiting the virulence hallmarks of Pseudomonas aeruginosa:
a chronicle through the perspective of quorum sensing. Environmental Microbiology.
2022;24(6):2630–56.

21. Cramer N, Klockgether J, Tümmler B. Microevolution of Pseudomonas aeruginosa in the
airways of people with cystic fibrosis. Current Opinion in Immunology. 2023 Aug 1;83:102328.

22. Moradali MF, Ghods S, Rehm BHA. Pseudomonas aeruginosa Lifestyle: A Paradigm for
Adaptation, Survival, and Persistence. Front Cell Infect Microbiol. 2017 Feb 15;7(39).

23. Rosales-Reyes R, Vargas-Roldán SY, Lezana-Fernández JL, Santos-Preciado JI. *Pseudomonas
Aeruginosa*: Genetic Adaptation, A Strategy for its Persistence in Cystic Fibrosis. Archives of
Medical Research. 2021 May 1;52(4):357–61.

24. Feng Q, Zhou J, Zhang L, Fu Y, Yang L. Insights into the molecular basis of c-di-GMP signalling in
Pseudomonas aeruginosa. Critical Reviews in Microbiology. 2024 Jan 2;50(1):20–38.

25. Krasteva PV, Giglio KM, Sondermann H. Sensing the messenger: The diverse ways that bacteria
signal through c-di-GMP. Protein Science. 2012;21(7):929–48.

26. Valentini M, Filloux A. Multiple Roles of c-di-GMP Signaling in Bacterial Pathogenesis. Annual
Review of Microbiology. 2019 Sep 8;73(Volume 73, 2019):387–406.

27. Hall CL, Lee VT. Cyclic-di-GMP regulation of virulence in bacterial pathogens. WIREs RNA.
2018;9(1):e1454.

28. Worlitzsch D, Tarran R, Ulrich M, Schwab U, Cekici A, Meyer KC, et al. Effects of reduced mucus
oxygen concentration in airway Pseudomonas infections of cystic fibrosis patients. J Clin Invest.
2002 Feb;109(3):317–25.

29. Alvarez-Ortega C, Harwood CS. Responses of Pseudomonas aeruginosa to low oxygen indicate
that growth in the cystic fibrosis lung is by aerobic respiration. Molecular Microbiology.
2007;65(1):153–65.

30. Hassett DJ, Sutton MD, Schurr MJ, Herr AB, Caldwell CC, Matu JO. Pseudomonas aeruginosa
hypoxic or anaerobic biofilm infections within cystic fibrosis airways. Trends in Microbiology.
2009 Mar 1;17(3):130–8.

31. Cullen L, O'Connor A, McCormack S, Owens RA, Holt GS, Collins C, et al. The involvement of the
low-oxygen-activated locus of Burkholderia cenocepacia in adaptation during cystic fibrosis
infection. Sci Rep. 2018 Sep 6;8(1):13386.

32. O'Connor A, Jurado-Martín I, Mysior MM, Manzira AL, Drabinska J, Simpson JC, et al. A universal stress protein upregulated by hypoxia has a role in Burkholderia cenocepacia intramacrophage survival: Implications for chronic infection in cystic fibrosis. Microbiologyopen. 2022 Dec 9;12(1):e1311.

33. Cullen L, O'Connor A, Drevinek P, Schaffer K, McClean S. Sequential Burkholderia cenocepacia Isolates from Siblings with Cystic Fibrosis Show Increased Lung Cell Attachment. Am J Respir Crit Care Med. 2017 Mar 15;195(6):832–5.

34. Cao P, Fleming D, Moustafa DA, Dolan SK, Szymanik KH, Redman WK, et al. A Pseudomonas aeruginosa small RNA regulates chronic and acute infection. Nature. 2023 Jun;618(7964):358–64.

35. De Soyza A, Hall AJ, Mahenthiralingam E, Drevinek P, Kaca W, Drulis-Kawa Z, et al. Developing an international Pseudomonas aeruginosa reference panel. Microbiologyopen. 2013 Dec;2(6):1010–23.

36. Freschi L, Bertelli C, Jeukens J, Moore MP, Kukavica-Ibrulj I, Emond-Rheault JG, et al. Genomic characterisation of an international Pseudomonas aeruginosa reference panel indicates that the two major groups draw upon distinct mobile gene pools. FEMS Microbiol Lett. 2018 Jul 1;365(14).

37. Cullen L, Weiser R, Olszak T, Maldonado RF, Moreira AS, Slachmuylders L, et al. Phenotypic characterization of an international Pseudomonas aeruginosa reference panel: strains of cystic fibrosis (CF) origin show less in vivo virulence than non-CF strains. Microbiology. 2015;161(10):1961–77.

38. Mulcahy LR, Burns JL, Lory S, Lewis K. Emergence of Pseudomonas aeruginosa Strains Producing High Levels of Persister Cells in Patients with Cystic Fibrosis. Journal of Bacteriology. 2010 Oct 8;192(23):6191.

39. Aanaes K, Rickelt LF, Johansen HK, von Buchwald C, Pressler T, Høiby N, et al. Decreased mucosal oxygen tension in the maxillary sinuses in patients with cystic fibrosis. J Cyst Fibros. 2011 Mar;10(2):114–20.

40. Walkiewicz A, Brzezińska M, Bieganowski A. Methanotrophs are favored under hypoxia in ammonium-fertilized soils. Biol Fertil Soils. 2018 Oct 1;54(7):861–70.

41. Xu A, Zhang X, Wang T, Xin F, Ma LZ, Zhou J, et al. Rugose small colony variant and its hyper-biofilm in *Pseudomonas aeruginosa*: Adaption, evolution, and biotechnological potential. Biotechnology Advances. 2021 Dec 1;53:107862.

42. The Galaxy Community. The Galaxy platform for accessible, reproducible, and collaborative data analyses: 2024 update. Nucleic Acids Research. 2024 Jul 5;52(W1):W83–94.

43. Szklarczyk D, Kirsch R, Koutrouli M, Nastou K, Mehryary F, Hachilif R, et al. The STRING database in 2023: protein-protein association networks and functional enrichment analyses for any sequenced genome of interest. Nucleic Acids Res. 2023 Jan 6;51(D1):D638–46.

44. Ge SX, Jung D, Yao R. ShinyGO: a graphical gene-set enrichment tool for animals and plants. Bioinformatics. 2020 Apr 15;36(8):2628–9.

45. Marmont LS, Rich JD, Whitney JC, Whitfield GB, Almblad H, Robinson H, et al. Oligomeric lipoprotein PelC guides Pel polysaccharide export across the outer membrane of Pseudomonas aeruginosa. Proceedings of the National Academy of Sciences. 2017 Mar 14;114(11):2892–7.

46. Culbertson JE, Toney MD. Expression and characterization of PhzE from *P. aeruginosa* PAO1: aminodeoxyisochorismate synthase involved in pyocyanin and phenazine-1-carboxylate production. Biochimica et Biophysica Acta (BBA) - Proteins and Proteomics. 2013 Jan 1;1834(1):240–6.

47.   Wang Y, Wang L, Zhang J, Duan X, Feng Y, Wang S, et al. PA0335, a Gene Encoding Histidinol
Phosphate Phosphatase, Mediates Histidine Auxotrophy in Pseudomonas aeruginosa. Applied
and Environmental Microbiology. 2020 Feb 18;86(5):e02593-19.

48.   Essar DW, Eberly L, Hadero A, Crawford IP. Identification and characterization of genes for a
second anthranilate synthase in Pseudomonas aeruginosa: interchangeability of the two
anthranilate synthases and evolutionary implications. J Bacteriol. 1990 Feb;172(2):884–900.

49.   Rojas Murcia N, Lee X, Waridel P, Maspoli A, Imker HJ, Chai T, et al. The Pseudomonas
aeruginosa antimetabolite L -2-amino-4-methoxy-trans-3-butenoic acid (AMB) is made from
glutamate and two alanine residues via a thiotemplate-linked tripeptide precursor. Front
Microbiol [Internet]. 2015 Mar 12 [cited 2025 Jun 27];6. Available from:
https://www.frontiersin.org/journals/microbiology/articles/10.3389/fmicb.2015.00170/full

50.   Müsken M, Di Fiore S, Dötsch A, Fischer R, Häussler S. Genetic determinants of Pseudomonas
aeruginosa biofilm establishment. Microbiology (Reading). 2010 Feb;156(Pt 2):431–41.

51.   Smith EW, Zhang X, Behzadi C, Andrews LD, Cohen F, Chen Y. Structures of Pseudomonas
aeruginosa LpxA Reveal the Basis for Its Substrate Selectivity. Biochemistry. 2015 Sep
29;54(38):5937–48.

52.   Hou M, Huang J, Jia T, Guan Y, Yang F, Zhou H, et al. Deep Profiling of the Proteome Dynamics
of Pseudomonas aeruginosa Reference Strain PAO1 under Different Growth Conditions. J
Proteome Res. 2023 Jun 2;22(6):1747–61.

53.   Florek LC, Lin X, Lin YC, Lin MH, Chakraborty A, Price-Whelan A, et al. The L-lactate
dehydrogenases of Pseudomonas aeruginosa are conditionally regulated but both contribute
to survival during macrophage infection. mBio. 2024 Aug 20;15(9):e00852-24.

54.   You Y, Ye F, Mao W, Yang H, Lai J, Deng S. An overview of the structure and function of the
flagellar hook FlgE protein. World J Microbiol Biotechnol. 2023 Mar 21;39(5):126.

55.   Valentin JDP, Straub H, Pietsch F, Lemare M, Ahrens CH, Schreiber F, et al. Role of the flagellar
hook in the structural development and antibiotic tolerance of Pseudomonas aeruginosa
biofilms. The ISME Journal. 2022 Apr 1;16(4):1176–86.

56.   Dasgupta N, Wolfgang MC, Goodman AL, Arora SK, Jyot J, Lory S, et al. A four-tiered
transcriptional regulatory circuit controls flagellar biogenesis in Pseudomonas aeruginosa.
Molecular Microbiology. 2003;50(3):809–24.

57.   Francis VI, Stevenson EC, Porter SL. Two-component systems required for virulence in
Pseudomonas aeruginosa. FEMS Microbiology Letters. 2017 Jun 15;364(11):fnx104.

58.   Kilmury SLN, Burrows LL. The Pseudomonas aeruginosa PilSR Two-Component System
Regulates   Both   Twitching   and   Swimming   Motilities.   mBio.   2018   Jul
24;9(4):10.1128/mbio.01310-18.

59.   Wiesmann CL, Zhang Y, Alford M, Hamilton CD, Dosanjh M, Thoms D, et al. The ColR/S two-
component system is a conserved determinant of host association across Pseudomonas
species. The ISME Journal. 2023 Feb 1;17(2):286–96.

60.   Gellatly SL, Bains M, Breidenstein EBM, Strehmel J, Reffuveille F, Taylor PK, et al. Novel roles for
two-component regulatory systems in cytotoxicity and virulence-related properties in
Pseudomonas aeruginosa. AIMS Microbiol. 2018 Mar 8;4(1):173–91.

61.   Latour X. The Evanescent GacS Signal. Microorganisms. 2020 Nov 6;8(11):1746.

62.   Davies JA, Harrison JJ, Marques LLR, Foglia GR, Stremick CA, Storey DG, et al. The GacS sensor
kinase controls phenotypic reversion of small colony variants isolated from biofilms of
Pseudomonas aeruginosa PA14. FEMS Microbiology Ecology. 2007;59(1):32–46.

63. Weiler AJ, Spitz O, Gudzuhn M, Schott-Verdugo SN, Kamel M, Thiele B, et al. A phospholipase B
from *Pseudomonas aeruginosa* with activity towards endogenous phospholipids affects biofilm
assembly. Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids. 2022 Apr
1;1867(4):159101.

64. Wu W, Huang J, Xu Z. Antibiotic influx and efflux in Pseudomonas aeruginosa: Regulation and
therapeutic implications. Microbial Biotechnology. 2024;17(5):e14487.

65. Alcalde-Rico M, Hernando-Amado S, Blanco P, Martínez JL. Multidrug Efflux Pumps at the
Crossroad between Antibiotic Resistance and Bacterial Virulence. Front Microbiol [Internet].
2016 Sep 21 [cited 2025 Jun 27];7. Available from:
https://www.frontiersin.org/journals/microbiology/articles/10.3389/fmicb.2016.01483/full

66. Jin Y, Yang H, Qiao M, Jin S. MexT Regulates the Type III Secretion System through MexS and
PtrC in Pseudomonas aeruginosa. Journal of Bacteriology. 2011 Jan 3;193(2):399–410.

67. Kristensen R, Andersen JB, Rybtke M, Jansen CU, Fritz BG, Kiilerich RO, et al. Inhibition of
Pseudomonas aeruginosa quorum sensing by chemical induction of the MexEF-oprN efflux
pump. Antimicrobial Agents and Chemotherapy. 2024 Jan 8;68(2):e01387-23.

68. Bandyopadhyay S, Chandramouli K, Johnson MK. Iron–sulfur cluster biosynthesis. Biochemical
Society Transactions. 2008 Nov 19;36(6):1112–9.

69. De Los Santos PE, Parret AHA, De Mot R. Stress-related Pseudomonas genes involved in
production of bacteriocin LlpA. FEMS Microbiology Letters. 2005 Mar 1;244(2):243–50.

70. Michel L, González N, Jagdeep S, Nguyen-Ngoc T, Reimmann C. PchR-box recognition by the
AraC-type regulator PchR of Pseudomonas aeruginosa requires the siderophore pyochelin as
an effector. Molecular Microbiology. 2005;58(2):495–509.

71. Trunk K, Benkert B, Quäck N, Münch R, Scheer M, Garbe J, et al. Anaerobic adaptation in
Pseudomonas aeruginosa: definition of the Anr and Dnr regulons. Environmental Microbiology.
2010;12(6):1719–33.

72. Hausmann S, Gonzalez D, Geiser J, Valentini M. The DEAD-box RNA helicase RhlE2 is a global
regulator of Pseudomonas aeruginosa lifestyle and pathogenesis. Nucleic Acids Research. 2021
Jul 9;49(12):6925–40.

73. Pu M, Sheng L, Song S, Gong T, Wood TK. Serine Hydroxymethyltransferase ShrA (PA2444)
Controls Rugose Small-Colony Variant Formation in Pseudomonas aeruginosa. Front Microbiol
[Internet]. 2018 Feb 27 [cited 2025 May 7];9. Available from:
https://www.frontiersin.orghttps://www.frontiersin.org/journals/microbiology/articles/10.338
9/fmicb.2018.00315/full

74. Hung RJ, Chien HS, Lin RZ, Lin CT, Vatsyayan J, Peng HL, et al. Comparative analysis of two UDP-
glucose dehydrogenases in Pseudomonas aeruginosa PAO1. J Biol Chem. 2007 Jun
15;282(24):17738–48.

75. Moskowitz SM, Ernst RK, Miller SI. PmrAB, a Two-Component Regulatory System of
Pseudomonas aeruginosa That Modulates Resistance to Cationic Antimicrobial Peptides and
Addition of Aminoarabinose to Lipid A. J Bacteriol. 2004 Jan;186(2):575–9.

76. Salunkhe P, Smart CHM, Morgan JAW, Panagea S, Walshaw MJ, Hart CA, et al. A Cystic Fibrosis
Epidemic Strain of Pseudomonas aeruginosa Displays Enhanced Virulence and Antimicrobial
Resistance. Journal of Bacteriology. 2005 Jul;187(14):4908–20.

77. Bugg TD, Walsh CT. Intracellular steps of bacterial cell wall peptidoglycan biosynthesis:
enzymology, antibiotics, and antibiotic resistance. Nat Prod Rep. 1992 Jun;9(3):199–215.

78. El Zoeiby A, Sanschagrin F, Lamoureux J, Darveau A, Levesque RC. Cloning, over-expression and
purification of Pseudomonas aeruginosa murC encoding uridine diphosphate N-
acetylmuramate: l-alanine ligase. FEMS Microbiology Letters. 2000 Feb 1;183(2):281–8.

79. Tang J, Luo M, Niu S, Zhou H, Cai X, Zhang W, et al. The Crystal Structure of Hexamer RraA from
Pseudomonas Aeruginosa Reveals Six Conserved Protein–Protein Interaction Sites. Protein J.
2010 Nov 1;29(8):583–90.

80. Lossi NS, Manoli E, Simpson P, Jones C, Hui K, Dajani R, et al. The archetype Pseudomonas
aeruginosa proteins TssB and TagJ form a novel subcomplex in the bacterial type VI secretion
system. Molecular Microbiology. 2012;86(2):437–56.

81. Chevalier S, Bouffartigues E, Bodilis J, Maillot O, Lesouhaitier O, Feuilloley MGJ, et al. Structure,
function and regulation of Pseudomonas aeruginosa porins. FEMS Microbiology Reviews. 2017
Sep 1;41(5):698–722.

82. Nguyen KV, Nguyen ,Trung Vu, Nguyen ,Hang Thi Thuy, and Le DV. Mutations in the gyrA, parC,
and mexR genes provide functional insights into the fluoroquinolone-resistant Pseudomonas
aeruginosa isolated in Vietnam. Infection and Drug Resistance. 2018 Feb 2;11:275–82.

83. Arabameri N, Heshmatipour Z, Eftekhar Ardebili S, Jafari Bidhendi Z. The role of Gene
Mutations (gyrA, parC) in Resistance to Ciprofloxacin in Clinical Isolates of Pseudomonas
Aeruginosa. Iran J Pathol. 2021;16(4):426–32.

84. Ortiz-Severín J, Varas M, Bravo-Toncio C, Guiliani N, Chávez FP. Multiple antibiotic susceptibility
of polyphosphate kinase mutants (ppk1 and ppk2) from Pseudomonas aeruginosa PAO1 as
revealed by global phenotypic analysis. biol res. 2015 Apr 25;48(1):22.

85. Huang Q, Yang J, Li C, Song Y, Zhu Y, Zhao N, et al. Structural characterization of PaFkbA: A
periplasmic chaperone from Pseudomonas aeruginosa. Computational and Structural
Biotechnology Journal. 2021 Jan 1;19:2460–7.

86. Alam A, Bröms JE, Kumar R, Sjöstedt A. The Role of ClpB in Bacterial Stress Responses and
Virulence. Front Mol Biosci [Internet]. 2021 Apr 22 [cited 2025 Jun 29];8. Available from:
https://www.frontiersin.org/journals/molecular-
biosciences/articles/10.3389/fmolb.2021.668910/full

87. Nolan LM, Cain AK, Clamens T, Furniss RCD, Manoli E, Sainz-Polo MA, et al. Identification of
Tse8 as a Type VI secretion system toxin from Pseudomonas aeruginosa that targets the
bacterial transamidosome to inhibit protein synthesis in prey cells. Nat Microbiol. 2021
Sep;6(9):1199–210.

88. Nikel PI, Pérez-Pantoja D, de Lorenzo V. Pyridine nucleotide transhydrogenases enable redox
balance of Pseudomonas putida during biodegradation of aromatic compounds. Environ
Microbiol. 2016 Oct;18(10):3565–82.

89. Li C, Wally H, Miller SJ, Lu CD. The Multifaceted Proteins MvaT and MvaU, Members of the H-
NS Family, Control Arginine Metabolism, Pyocyanin Synthesis, and Prophage Activation in
Pseudomonas aeruginosa PAO1. Journal of Bacteriology. 2009 Oct 15;191(20):6211–8.

90. Williams McMackin EA, Marsden AE, Yahr TL. H-NS Family Members MvaT and MvaU Regulate
the Pseudomonas aeruginosa Type III Secretion System. Journal of Bacteriology. 2019 Jun
21;201(14):10.1128/jb.00054-19.

91. Castang S, Dove SL. Basis for the Essentiality of H-NS Family Members in Pseudomonas
aeruginosa. Journal of Bacteriology. 2012 Aug 28;194(18):5101–9.

92. Nganje CN, Haynes SA, Qabar CM, Lent RC, Ghanem ENB, Shainheit MG. PepN is a non-
essential, cell wall-localized protein that contributes to neutrophil elastase-mediated killing of
Streptococcus pneumoniae. PLOS ONE. 2019 Feb 1;14(2):e0211632.

93. Lequette Y, Rollet E, Delangle A, Greenberg EP, Bohin JP. Linear osmoregulated periplasmic
glucans are encoded by the opgGH locus of Pseudomonas aeruginosa. Microbiology.
2007;153(10):3255–63.

94. Hauser AR, Kang PJ, Engel JN. PepA, a secreted protein of Pseudomonas aeruginosa, is
necessary for cytotoxicity and virulence. Mol Microbiol. 1998 Feb;27(4):807–18.

95. Tipton KA, Coleman JP, Pesci EC. Post-transcriptional regulation of gene PA5507 controls
Pseudomonas quinolone signal concentration in P. aeruginosa. Molecular Microbiology.
2015;96(3):670–83.

96. Lin Q, Wang H, Huang J, Liu Z, Chen Q, Yu G, et al. Spermidine Is an Intercellular Signal
Modulating T3SS Expression in Pseudomonas aeruginosa. Microbiology Spectrum. 2022 Apr
18;10(3):e00644-22.

97. Wu D, Lim SC, Dong Y, Wu J, Tao F, Zhou L, et al. Structural Basis of Substrate Binding Specificity
Revealed by the Crystal Structures of Polyamine Receptors SpuD and SpuE from *Pseudomonas
aeruginosa*. Journal of Molecular Biology. 2012 Mar 9;416(5):697–712.

98. Marquart ME, Caballero AR, Chomnawang M, Thibodeaux BA, Twining SS, O'Callaghan RJ.
Identification of a Novel Secreted Protease from Pseudomonas aeruginosa that Causes Corneal
Erosions. Investigative Ophthalmology & Visual Science. 2005 Oct 1;46(10):3761–8.

99. Lee C, Kim MI, Park J, Kim J, Oh H, Cho Y, et al. Crystal structure of the *Pseudomonas
aeruginosa* PA0423 protein and its functional implication in antibiotic sequestration.
Biochemical and Biophysical Research Communications. 2020 Jul 12;528(1):85–91.

100. Romsang A, Duang-nkern J, Saninjuk K, Vattanaviboon P, Mongkolsuk S. Pseudomonas
aeruginosa nfuA: Gene regulation and its physiological roles in sustaining growth under stress
and anaerobic conditions and maintaining bacterial virulence. PLOS ONE. 2018 Aug
9;13(8):e0202151.

101. Daung-nkern J, Vattanaviboon P, Mongkolsuk S. Inactivation of nfuA enhances susceptibility of
Pseudomonas aeruginosa to fluoroquinolone antibiotics. Journal of Antimicrobial
Chemotherapy. 2010 Aug 1;65(8):1831–2.

102. Drabinska J, Ziecina M, Modzelan M, Jagura-Burdzy G, Kraszewska E. Individual Nudix
hydrolases affect diverse features of Pseudomonas aeruginosa. Microbiologyopen. 2020 May
17;9(8):e1052.

103. Semmler ABT, Whitchurch CB, Leech AJ, Mattick JS. Identification of a novel gene, fimV,
involved in twitching motility in Pseudomonas aeruginosaThe GenBank accession number for
the sequence determined in this work is U93274. Microbiology. 2000;146(6):1321–32.

104. Michel GPF, Aguzzi A, Ball G, Soscia C, Bleves S, Voulhoux R. Role of fimV in type II secretion
system-dependent protein secretion of Pseudomonas aeruginosa on solid medium.
Microbiology. 2011;157(7):1945–54.

105. Shen L, Gao L, Yang M, Zhang J, Wang Y, Feng Y, et al. Deletion of the PA4427-PA4431 Operon
of Pseudomonas aeruginosa PAO1 Increased Antibiotics Resistance and Reduced Virulence and
Pathogenicity by Affecting Quorum Sensing and Iron Uptake. Microorganisms. 2021 May
14;9(5):1065.

106. Duque E, de la Torre J, Bernal P, Molina-Henares MA, Alaminos M, Espinosa-Urgel M, et al.
Identification of reciprocal adhesion genes in pathogenic and non-pathogenic Pseudomonas.
Environmental Microbiology. 2013;15(1):36–48.

107. Yoon SS, Hennigan RF, Hilliard GM, Ochsner UA, Parvatiyar K, Kamani MC, et al. Pseudomonas
aeruginosa Anaerobic Respiration in Biofilms: Relationships to Cystic Fibrosis Pathogenesis.
Developmental Cell. 2002;

108. König J, Schreiber R, Voelcker T, Mall M, Kunzelmann K. The cystic fibrosis transmembrane
conductance regulator (CFTR) inhibits ENaC through an increase in the intracellular Cl−
concentration. EMBO Rep. 2001 Nov 15;2(11):1047–51.

109. Cowley ES, Kopf SH, LaRiviere A, Ziebis W, Newman DK. Pediatric Cystic Fibrosis Sputum Can Be
Chemically Dynamic, Anoxic, and Extremely Reduced Due to Hydrogen Sulfide Formation.
mBio. 2015 Jul 28;6(4):10.1128/mbio.00767-15.

110. Line L, Alhede M, Kolpen M, Kühl M, Ciofu O, Bjarnsholt T, et al. Physiological levels of nitrate
support anoxic growth by denitrification of Pseudomonas aeruginosa at growth rates reported
in cystic fibrosis lungs and sputum. Front Microbiol. 2014 Oct 24;5:554.

111. Borriello G, Richards L, Ehrlich GD, Stewart PS. Arginine or Nitrate Enhances Antibiotic
Susceptibility of Pseudomonas aeruginosa in Biofilms. Antimicrobial Agents and
Chemotherapy. 2006 Jan;50(1):382–4.

112. Carey CJ, Duggan N, Drabinska J, McClean S. Harnessing hypoxia: bacterial adaptation and
chronic infection in cystic fibrosis. FEMS Microbiology Reviews. 2025 Jan 1;49:fuaf018.

113. Tribelli PM, Lujan AM, Pardo A, Ibarra JG, Fernández Do Porto D, Smania A, et al. Core regulon
of the global anaerobic regulator Anr targets central metabolism functions in Pseudomonas
species. Sci Rep. 2019 Jun 21;9(1):9065.

114. Papa R, Imperlini E, Trecca M, Paris I, Vrenna G, Artini M, et al. Virulence of Pseudomonas
aeruginosa in Cystic Fibrosis: Relationships between Normoxia and Anoxia Lifestyle.
Antibiotics. 2024 Jan;13(1):1.

115. Kamath KS, Krisp C, Chick J, Pascovici D, Gygi SP, Molloy MP. Pseudomonas aeruginosa
Proteome under Hypoxic Stress Conditions Mimicking the Cystic Fibrosis Lung. J Proteome Res.
2017 Oct 6;16(10):3917–28.

116. Ashish A, Paterson S, Mowat E, Fothergill JL, Walshaw MJ, Winstanley C. Extensive
diversification is a common feature of Pseudomonas aeruginosa populations during respiratory
infections in cystic fibrosis. J Cyst Fibros. 2013 Dec;12(6):790–3.

117. Flynn KM, Dowell G, Johnson TM, Koestler BJ, Waters CM, Cooper VS. Evolution of Ecological
Diversity in Biofilms of Pseudomonas aeruginosa by Altered Cyclic Diguanylate Signaling.
Journal of Bacteriology. 2016 Sep 9;198(19):2608–18.

118. Jorth P, Staudinger BJ, Wu X, Hisert K, Hayden H, Garudathri J, et al. Regional Isolation Drives
Bacterial Diversification within Cystic Fibrosis Lungs. Cell Host Microbe. 2015 Sep 9;18(3):307–
19.

119. Markussen T, Marvig RL, Gómez-Lozano M, Aanæs K, Burleigh AE, Høiby N, et al.
Environmental heterogeneity drives within-host diversification and evolution of Pseudomonas
aeruginosa. mBio. 2014 Sep 16;5(5):e01592-01514.

120. Winstanley C, O'Brien S, Brockhurst MA. Pseudomonas aeruginosa Evolutionary Adaptation
and Diversification in Cystic Fibrosis Chronic Lung Infections. Trends Microbiol. 2016
May;24(5):327–37.

121. Letunic I, Bork P. Interactive Tree of Life (iTOL) v6: recent updates to the phylogenetic tree
display and annotation tool. Nucleic Acids Research. 2024 Jul 5;52(W1):W78–82.

122. Bueno E, Mesa S, Bedmar EJ, Richardson DJ, Delgado MJ. Bacterial Adaptation of Respiration
from Oxic to Microoxic and Anoxic Conditions: Redox Control. Antioxidants & Redox Signaling.
2012 Apr 15;16(8):819–52.

123. Scribani Rossi C, Eckartt K, Scarchilli E, Angeli S, Price-Whelan A, Di Matteo A, et al. Molecular
insights into RmcA-mediated c-di-GMP consumption: Linking redox potential to biofilm
morphogenesis in Pseudomonas aeruginosa. Microbiol Res. 2023 Dec;277:127498.

124. Schmidt A, Hammerbacher AS, Bastian M, Nieken KJ, Klockgether J, Merighi M, et al. Oxygen-
dependent regulation of c-di-GMP synthesis by SadC controls alginate production in
seudomonas aeruginosa. Environmental Microbiology. 2016;18(10):3390–402.

125. An S, Wu J, Zhang LH. Modulation of Pseudomonas aeruginosa Biofilm Dispersal by a Cyclic-Di-GMP Phosphodiesterase with a Putative Hypoxia-Sensing Domain. Appl Environ Microbiol. 2010 Dec;76(24):8160–73.

126. Wang H, de Carvalho LPS. Metabolomic profiling reveals bacterial metabolic adaptation strategies and new metabolites. Current Opinion in Chemical Biology. 2023 Jun 1;74:102287.

127. Banerjee P, Sahoo PK, Sheenu null, Adhikary A, Ruhal R, Jain D. Molecular and structural facets of c-di-GMP signalling associated with biofilm formation in Pseudomonas aeruginosa. Mol Aspects Med. 2021 Oct;81:101001.

128. Gupta K, Liao J, Petrova OE, Cherny KE, Sauer K. Elevated levels of the second messenger c-di-GMP contribute to antimicrobial resistance of Pseudomonas aeruginosa. Mol Microbiol. 2014 May;92(3):488–506.

129. Arai H. Regulation and Function of Versatile Aerobic and Anaerobic Respiratory Metabolism in Pseudomonas aeruginosa. Front Microbiol [Internet]. 2011 May 5 [cited 2025 May 20];2. Available from: https://www.frontiersin.org/journals/microbiology/articles/10.3389/fmicb.2011.00103/full

130. O'Connor JR, Kuwada NJ, Huangyutitham V, Wiggins PA, Harwood CS. Surface sensing and lateral subcellular localization of WspA, the receptor in a chemosensory-like system leading to c-di-GMP production. Mol Microbiol. 2012 Nov;86(3):720–9.

131. Dev IK, Harvey RJ. Regulation of synthesis of serine hydroxymethyltransferase in chemostat cultures of Escherichia coli. Journal of Biological Chemistry. 1984 Jul 10;259(13):8394–401.

132. Attila C, Ueda A, Wood TK. PA2663 (PpyR) increases biofilm formation in Pseudomonas aeruginosa PAO1 through the psl operon and stimulates virulence and quorum-sensing phenotypes. Appl Microbiol Biotechnol. 2008 Feb;78(2):293–307.

133. Brencic A, McFarland KA, McManus HR, Castang S, Mogno I, Dove SL, et al. The GacS/GacA signal transduction system of Pseudomonas aeruginosa acts exclusively through its control over the transcription of the RsmY and RsmZ regulatory small RNAs. Molecular Microbiology. 2009;73(3):434–45.

134. Yan J, Deforet M, Boyle KE, Rahman R, Liang R, Okegbe C, et al. Bow-tie signaling in c-di-GMP: Machine learning in a simple biochemical network. PLOS Computational Biology. 2017 Aug 2;13(8):e1005677.

135. Besse A, Groleau MC, Trottier M, Vincent AT, Déziel E. Pseudomonas aeruginosa Strains from Both Clinical and Environmental Origins Readily Adopt a Stable Small-Colony-Variant Phenotype Resulting from Single Mutations in c-di-GMP Pathways. J Bacteriol. 2022 Oct 18;204(10):e0018522.

136. Besse A, Groleau MC, Déziel E. Emergence of Small Colony Variants Is an Adaptive Strategy Used by Pseudomonas aeruginosa to Mitigate the Effects of Redox Imbalance. mSphere. 2023 Feb 28;8(2):e00057-23.

137. Johns BE, Purdy KJ, Tucker NP, Maddocks SE. Phenotypic and Genotypic Characteristics of Small Colony Variants and Their Role in Chronic Infection. Microbiol Insights. 2015 Jan 1;8:MBI.S25800.

138. Ryan H, Ballard E, Stockwell RE, Duplancic C, Thomson RM, Smith K, et al. A systematic review of the clinical impact of small colony variants in patients with cystic fibrosis. BMC Pulm Med. 2023 Sep 1;23(1):323.

139. Schneider M, Mühlemann K, Droz S, Couzinet S, Casaulta C, Zimmerli S. Clinical Characteristics Associated with Isolation of Small-Colony Variants of Staphylococcus aureus and Pseudomonas aeruginosa from Respiratory Secretions of Patients with Cystic Fibrosis. Journal of Clinical Microbiology. 2008 Mar 5;46(5):1832.

140. Malone JG. Role of small colony variants in persistence of Pseudomonas aeruginosa infections
in cystic fibrosis lungs. Infection and Drug Resistance. 2015 Jul 29;8(null):237–47.

141. Hoffman LR, Déziel E, D'Argenio DA, Lépine F, Emerson J, McNamara S, et al. Selection for
Staphylococcus aureus small-colony variants due to growth in the presence of Pseudomonas
aeruginosa. Proceedings of the National Academy of Sciences. 2006 Dec 26;103(52):19890–5.

142. Bjarnsholt T, Jensen PØ, Fiandaca MJ, Pedersen J, Hansen CR, Andersen CB, et al. Pseudomonas
aeruginosa biofilms in the respiratory tract of cystic fibrosis patients. Pediatr Pulmonol. 2009
Jun;44(6):547–58.

143. Høiby N, Ciofu O, Bjarnsholt T. Pseudomonas Aeruginosa Biofilms in Cystic fibrosis. Future
Microbiology. 2010 Nov 1;5(11):1663–74.

144. Maurice NM, Bedi B, Sadikot RT. Pseudomonas aeruginosa Biofilms: Host Response and Clinical
Implications in Lung Infections. Am J Respir Cell Mol Biol. 2018 Apr;58(4):428–39.

145. Klementiev AD, Jin Z, Whiteley M. Micron Scale Spatial Measurement of the O2 Gradient
Surrounding a Bacterial Biofilm in Real Time. mBio. 2020 Oct 20;11(5):10.1128/mbio.02536-20.

146. Wu Y, Klapper I, Stewart PS. Hypoxia arising from concerted oxygen consumption by
neutrophils and microorganisms in biofilms. Pathogens and Disease. 2018 Apr 23;76(4):fty043.

147. Bowler P, Murphy C, Wolcott R. Biofilm exacerbates antibiotic resistance: Is this a current
oversight in antimicrobial stewardship? Antimicrob Resist Infect Control. 2020 Oct 20;9(1):162.

148. Liang Z, Nilsson M, Kragh KN, Hedal I, Alcàcer-Almansa J, Kiilerich RO, et al. The role of
individual exopolysaccharides in antibiotic tolerance of Pseudomonas aeruginosa aggregates.
Front Microbiol. 2023 Jul 2;14(1187708).

149. Oladosu VI, Park S, Sauer K. Flip the switch: the role of FleQ in modulating the transition
between the free-living and sessile mode of growth in Pseudomonas aeruginosa. Journal of
Bacteriology. 2024 Mar 4;206(3):e00365-23.

150. Sindeldecker D, Stoodley P. The many antibiotic resistance and tolerance strategies of
Pseudomonas aeruginosa. Biofilm. 2021 Dec 1;3:100056.

151. Borriello G, Werner E, Roe F, Kim AM, Ehrlich GD, Stewart PS. Oxygen Limitation Contributes to
Antibiotic Tolerance of Pseudomonas aeruginosa in Biofilms. Antimicrobial Agents and
Chemotherapy. 2004 Jul;48(7):2659–64.

152. Schaible B, Taylor CT, Schaffer K. Hypoxia Increases Antibiotic Resistance in Pseudomonas
aeruginosa through Altering the Composition of Multidrug Efflux Pumps. Antimicrobial Agents
and Chemotherapy. 2012 Mar 16;56(4):2114–8.

153. Zhao X, Drlica K. Reactive oxygen species and the bacterial response to lethal stress. Current
opinion in microbiology. 2014 Jul 30;0:1.

154. Macfarlane ELA, Kwasnicka A, Ochs MM, Hancock REW. PhoP–PhoQ homologues in
Pseudomonas aeruginosa regulate expression of the outer-membrane protein OprH and
polymyxin B resistance. Molecular Microbiology. 1999;34(2):305–16.

155. Edrington TC, Kintz E, Goldberg JB, Tamm LK. Structural Basis for the Interaction of
Lipopolysaccharide with Outer Membrane Protein H (OprH) from Pseudomonas aeruginosa*.
Journal of Biological Chemistry. 2011 Nov 11;286(45):39211–23.

156. Hameed P S, Manjrekar P, Chinnapattu M, Humnabadkar V, Shanbhag G, Kedari C, et al.
Pyrazolopyrimidines Establish MurC as a Vulnerable Target in Pseudomonas aeruginosa and
Escherichia coli. ACS Chem Biol. 2014 Oct 17;9(10):2274–82.

157. Nalca Y, Jänsch L, Bredenbruch F, Geffers R, Buer J, Häussler S. Quorum-Sensing Antagonistic
Activities of Azithromycin in Pseudomonas aeruginosa PAO1: a Global Approach. Antimicrob
Agents Chemother. 2006 May;50(5):1680–8.

158. Rausch M, Deisinger JP, Ulm H, Müller A, Li W, Hardt P, et al. Coordination of capsule assembly
and cell wall biosynthesis in Staphylococcus aureus. Nat Commun. 2019 Mar 29;10(1):1404.

159. Higgins PG, Fluit AC, Milatovic D, Verhoef J, Schmitz FJ. Mutations in GyrA, ParC, MexR and
NfxB in clinical isolates of *Pseudomonas aeruginosa*. International Journal of Antimicrobial
Agents. 2003 May 1;21(5):409–13.

160. Hunter RC, Asfour F, Dingemans J, Osuna BL, Samad T, Malfroot A, et al. Ferrous Iron Is a
Significant Component of Bioavailable Iron in Cystic Fibrosis Airways. mBio. 2013 Aug
20;4(4):10.1128/mbio.00557-13.

161. Moreau-Marquis S, Bomberger JM, Anderson GG, Swiatecka-Urban A, Ye S, O'Toole GA, et al.
The DeltaF508-CFTR mutation results in increased biofilm formation by Pseudomonas
aeruginosa by increasing iron availability. Am J Physiol Lung Cell Mol Physiol. 2008
Jul;295(1):L25-37.

162. Schalk IJ, Perraud Q. and its multiple strategies to access iron. Environmental Microbiology.
2023;25(4):811–31.

163. Reid DW, Carroll V, O'May C, Champion A, Kirov SM. Increased airway iron as a potential factor
in the persistence of Pseudomonas aeruginosa infection in cystic fibrosis. European Respiratory
Journal. 2007 Aug 1;30(2):286–92.

164. Frangipani E, Visaggio D, Heeb S, Kaever V, Cámara M, Visca P, et al. The Gac/Rsm and cyclic-di-
GMP signalling networks coordinately regulate iron uptake in seudomonas aeruginosa.
Environmental Microbiology. 2014;16(3):676–88.

165. Banin E, Brady KM, Greenberg EP. Chelator-induced dispersal and killing of Pseudomonas
aeruginosa cells in a biofilm. Appl Environ Microbiol. 2006 Mar;72(3):2064–9.

166. Miller HK, Auerbuch V. Bacterial iron-sulfur cluster sensors in mammalian pathogens.
Metallomics. 2015 Jun 10;7(6):943–56.

167. Konings AF, Martin LW, Sharples KJ, Roddam LF, Latham R, Reid DW, et al. Pseudomonas
aeruginosa Uses Multiple Pathways To Acquire Iron during Chronic Infection in Cystic Fibrosis
Lungs. Infection and Immunity. 2013 Jul 11;81(8):2697–704.

168. Schaible B, Rodriguez J, Garcia A, von Kriegsheim A, McClean S, Hickey C, et al. Hypoxia
Reduces the Pathogenicity of Pseudomonas aeruginosa by Decreasing the Expression of
Multiple Virulence Factors. J Infect Dis. 2017 May 1;215(9):1459–67.

169. Weimann A, Dinan AM, Ruis C, Bernut A, Pont S, Brown K, et al. Evolution and host-specific
adaptation of Pseudomonas aeruginosa. Science. 2024 Jul 5;385(6704):eadi0908.

170. Amoureux L, Bador J, Fardeheb S, Mabille C, Couchot C, Massip C, et al. Detection of
Achromobacter xylosoxidans in Hospital, Domestic, and Outdoor Environmental Samples and
Comparison with Human Clinical Isolates. Appl Environ Microbiol. 2013 Dec;79(23):7142–9.

171. Crone S, Vives-Flórez M, Kvich L, Saunders AM, Malone M, Nicolaisen MH, et al. The
environmental occurrence of Pseudomonas aeruginosa. APMIS. 2020;128(3):220–31.

172. Honda JR, Virdi R, Chan ED. Global Environmental Nontuberculous Mycobacteria and Their
Contemporaneous Man-Made and Natural Niches. Front Microbiol. 2018 Aug 30;9.

173. Honda JR, Virdi R, Chan ED. Global Environmental Nontuberculous Mycobacteria and Their
Contemporaneous Man-Made and Natural Niches. Front Microbiol. 2018;9:2029.

174. Price CTD, Hanford HE, Al-Quadan T, Santic M, Shin CJ, Da'as MSJ, et al. Amoebae as training
grounds for microbial pathogens. mBio. 2024 Jul 8;15(8):e00827-24.

175. Mullard A. 2023 FDA approvals. Nat Rev Drug Discov. 2024 Feb;23(2):88–95.

176. Rashid MH, Kornberg A. Inorganic polyphosphate is needed for swimming, swarming, and
twitching motilities of Pseudomonas aeruginosa. Proc Natl Acad Sci U S A. 2000 Apr
25;97(9):4885–90.
177. Arora NK, Verma M. Modified microplate method for rapid and efficient estimation of
siderophore produced by bacteria. 3 Biotech. 2017 Dec;7(6):381.
178. Roy AB, Petrova OE, Sauer K. Extraction and Quantification of Cyclic Di-GMP from P.
aeruginosa. Bio Protoc. 2013;3(14):e828.

Supporting information captions

S1. Figure. Differences in growth patterns of the *P. aeruginosa* AMT 0023-30 early infection strain with different inoculation volumes.

S2. Figure. Differences in doubling times of the *P. aeruginosa* AMT 002330 early infection strain in hypoxic and normoxic conditions. A. OD_{600} measurements over time. **B.** CFU/ml counts over time **C.** OD_{600} to CFU/ml trend lines and linear regression calculations. **D.** Differences in generation times in normoxia and hypoxia.

S3. Figure. Stability of SCVs in normoxic conditions. A. Photograph of different *P. aeruginosa* colony variants streaked on LB agar and grown overnight in normoxic conditions. **B.** Enlarged fragment of photograph in section A.

S4. Table. Proteomic analysis of long-term hypoxia-adapted *P. aeruginosa* isolates. A. All proteins differentially expressed between HACs and the ancestral strain (D0), NACs and the ancestral strain (D0) or HACs and NACs. **B.** Proteins not detected in at least two replicates in at least one of the groups (Day 0, HACs, NACs). C. All proteins detected in the proteomics analysis of the ancestral strain, HACs and NACs

S5. Table. Functional enrichment of the 89 proteins of interest significantly altered in abundance after long-term adaptation of *P. aeruginosa* to hypoxia

S6. Figure. Effects of long-term hypoxia adaptation on intracellular c-di-GMP concentrations. The graph represents the concentration of c-di-GMP in the cell extract from an equivalent of 4 ml of culture of OD_{600} = 1.8. No statistically significant differences between cultures were shown with the Brown-Forsythe ANOVA test. E: early infection clinical strain (AMT 0023-30); L: late infection clinical strain (AMT 0023-34); N1, N2, N3: 28-day normoxia-adapted populations; H1, H2, H3: 28-day hypoxia adapted populations.

A.

Norfloxacin — Inhibition zone diameter (cm): E, L(****), N1, N2, N3, H1(**), H2(**), H3

Ciprofoxacin — E, L(***), N1, N2, N3, H1(**), H2(*), H3

Azithromycin — E, L, N1(**), N2(*), N3, H1(*), H2, H3

Clarithromycin — E, L, N1, N2, N3, H1(**), H2(**), H3

Ceftazidime — E, L, N1, N2, N3, H1(***), H2(**), H3

Meropenem — Inhibition zone diameter (cm): E, L, N1, N2, N3, H1(***), H2(**), H3

Imipenem — E, L, N1, N2, N3(**), H1, H2, H3

Piperacillin — E, L, N1, N2, N3, H1(**), H2(**), H3

Aztreonam — E, L, N1, N2, N3, H1(**), H2(**), H3(*)

Tobramycin — E, L(****), N1(***), N2(**), N3(***), H1, H2, H3

B.

Antibiotic	L	N1	N2	N3	H1	H2	H3	Antibiotic Class
Norfloxacin	↑	=	=	=	↑	↑	=	Fluoroquinolones
Ciprofoxacin	↑	=	=	=	↑	↑	=	
Azithromycin	=	↑	↑	=	↑	=	=	Macrolides
Clarithromycin	=	=	=	=	↑	↑	=	
Ceftazidime	=	=	=	=	↑	↑	=	Cephalosporins
Cefotaxime	=	=	=	=	=	=	=	
Meropenem	=	=	=	=	↑	↑	=	Carbapenems
Imipenem	=	=	=	↑	=	=	=	
Tobramycin	↑R	↑	↑	↑	=	=	=	Aminoglycosides
Piperacillin	↓	=	=	=	↑	↑	=	Penicillins
Aztreonam	=	=	=	=	↓	↓	↑	Monobactams
Tetracycline	=R	=R	=R	=R	=R	=R	=R	Tetracyclines

C.

Control	Temperature	Oxidative	Osmotic	Acidic
LB agar, 37°C	45°C	1 mM H₂O₂	0.5 M NaCl	pH 5

E, L, N1, N2, N3 (top rows); E, H1, H2, H3 (bottom rows)

Dilution series: 0, -1, -2, -3, -4, -5

24h — 48h — 24h

D.

Stress type	L	N1	N2	N3	H1	H2	H3
Temperature	↓	↓	↓	=	↓	↓	=
Osmotic	↓	↑	=	↑	↑	↑	↑
Oxidative	=	=	=	↑	↑	↑	=
Acidic	=	↑	↑	↑	=	=	=

A.

24 h

OD_{590}/OD_{600}

E L N1 N2 N3 H1 H2 H3

**** (H1) **** (H3)

48 h

E L N1 N2 N3 H1 H2 H3

** (L) ** (N2) **** (H1) ** (H2)

72 h

E L N1 N2 N3 H1 H2 H3

*** (L) ** (N2) **** (H1) **** (H2) ** (H3)

B.

E   L   N1   N2   N3   H1   H2   H3

C.

Siderophore Units / CFU

E L N1 N2 N3 H1 H2 H3

** (L) * (H1) ** (H2) * (H3)

Phenotype		L	N1	N2	N3	H1	H2	H3
SCVs (%)		58.0%	0.00	0.00	2.7%	76.9%	79.8%	0.7%
Pyocyanin		↓	↓	=	=	↓	↓	=
Antibiotic resistance		=	=	=	=	↑	↑	=
Stress resistance	Osmotic	↓	↑	=	↑	↑	↑	↑
	Oxidative	=	=	=	↑	↑	↑	=
	Heat	↓	↓	↓	=	↓	↓	=
	Acidic	=	↑	↑	↑	=	=	=
Biofilm 72h		↑	=	=	↑	↑	↑	=
Exopolysaccharide production		=	=	=	↑	↑	↑	=
Motility	Swimming	↓	↑	↑	↓	↑	↑	↓
	Swarming	↓	=	↑	↓	↑	↑	↓
	Twitching	↓	=	↑	=	=	↑	↓
Siderophore production		↓	=	=	=	↓	↓	↓
Virulence (*G. mellonella*)		↑	=	=	=	=	=	=