Tertiary and quaternary structure remodeling by occupancy of the substrate binding

pocket in a large glutamate dehydrogenase

Melisa Lázaro [1,#], Nicolás Chamorro [2,#], Jorge P. López-Alonso [3,4], Diego Charro [1],

Rodolfo M. Rasia [2,5], Gonzalo Jiménez-Osés [1,6], Mikel Valle [1,*], María-Natalia Lisa [2,4,*]

[1] Center for Cooperative Research in Biosciences (CIC bioGUNE), Basque Research and

Technology Alliance (BRTA), Bizkaia Technology Park, Building 801A, 48160 Derio,

Spain.

[2] Instituto de Biología Molecular y Celular de Rosario (IBR CONICET-UNR), Ocampo y

Esmeralda, Rosario S2002LRK, Argentina.

[3] Instituto Biofisika (UPV/EHU, CSIC), 48940 Leioa, Spain.

[4] Basque Resource for Electron Microscopy, 48940 Leioa, Spain.

[5] Plataforma de Biología Estructural y Metabolómica (PLABEM), Ocampo y Esmeralda,

Rosario S2002LRK, Argentina.

[6] Ikerbasque, Basque Foundation for Science, 48013 Bilbao, Spain.

# Equal contribution

* Correspondence: MNL: lisa@ibr-conicet.gov.ar; MV: mvalle@cicbiogune.es

ABSTRACT

Glutamate dehydrogenases (GDHs) catalyze the oxidative deamination of L-glutamate to 2-oxoglutarate using $NAD(P)^+$ as a cofactor. The large type of GDHs (L-GDHs) displays a dynamic homotetrameric architecture that alternates between open and closed states. The catalytic mechanism and the role of the large conformational changes of L-GDHs in enzymatic function are unknown. Here, we explore by cryoEM the structure and the conformational space of the mycobacterial L-GDH composed of 180 kDa subunits (mL-GDH_{180}) when incubated with L-glutamate and NAD^+. Classification of the heterogeneous population of tetramers into different subsets reveals opening-closing motions and sorting of individual subunits resolves the occupancy of the cofactor and substrate binding pockets. Maps show that ligand binding to the glutamate binding pocket is accompanied by structural changes in a distant region that mediates a novel interaction between the catalytic domains of neighboring subunits in closed tetrameric states. Our findings indicate that the occupancy of the orthosteric site of mL-GDH_{180} is linked to a remodeling of the enzyme tertiary and quaternary structure.

INTRODUCTION

Amino acid dehydrogenases have been a focus of interest for decades due to their physiological roles and potential industrial applications. Among them, glutamate dehydrogenases (GDHs; EC 1.4.1.2) catalyze the reaction L-glutamate + NAD(P)$^+$ + H$_2$O <=> 2-oxoglutarate + NAD(P)H + H$^+$ + NH$_3$, at the crossroads of carbon and nitrogen metabolism. These enzymes are oligomeric and are classified based on the molecular weight of their subunits. Small GDHs (S-GDHs), present in prokaryotes and eukaryotes, are composed of 50 kDa subunits [1]. The conserved catalytic domain spans nearly the entire length of the primary structure of S-GDHs and provides the oligomerization surfaces. Biochemical and structural studies of diverse S-GDHs have allowed elucidating their catalytic and regulatory mechanisms. However, knowledge about such aspects for large GDHs (L-GDHs) present in prokaryotes and lower eukaryotes is comparatively scarce [2]. Among these enzymes are L-GDHs composed of 115 kDa subunits (L-GDHs$_{115}$) and those with 180 kDa subunits (L-GDHs$_{180}$), such as the L-GDH$_{180}$ found in mycobacteria (mL-GDH$_{180}$). Distinctive long N- and C-terminal regions flank the catalytic domain of L-GDHs [3].

In a previous work we discovered that the L-GDH$_{180}$ from *Mycobacterium smegmatis* (mL-GDH$_{180}$; 76% sequence identity with the L-GDH$_{180}$ from *Mycobacterium tuberculosis*) assembles into a homo-tetramer, adopting a quaternary structure that is radically different from that of S-GDHs [3]. Such architecture depends on interactions provided by an N-terminal region containing ACT- and PAS-type domains [4,5] with regulatory potential as well as a C-terminal domain of unknown function, both flanking the catalytic domain in mL-GDH$_{180}$ subunits.

Here we report cryoEM evidence of up to 3 Å resolution of alternative conformers of mL-GDH$_{180}$ detected in the presence of NAD$^+$ and L-glutamate. We found that ligand binding to the glutamate binding pocket is accompanied by changes in the tertiary structure in a

region distant from the active site, which in turn promotes interactions between the catalytic domains of neighboring monomers and modifications of the quaternary structure. Despite decades of research into the molecular basis of GDHs activity and regulation, experimental structural information on enzyme-cofactor-L-glutamate ternary complexes is still scarce, especially for NAD^+-dependent enzymes. Here, we present cryoEM evidence for enzyme species present upon incubation with L-glutamate and NAD^+ of a key L-GDH_{180} for the virulence of a long-attacked pathogen [6].

METHODS

Protein production

Plasmid pLIC-His-mL-GDH$_{180}$ [3] was used to produce N-terminally His6-tagged mL-GDH180 (MSMEG_4699, Uniprot A0R1C2) in *E. coli* cells. Transformed *E. coli* BL21(DE3) cells were grown at 37 °C in LB broth supplemented with 100 µg/ml ampicillin until reaching 0.8 units of optical density at 600 nm. Protein expression was then induced by adding isopropyl β-D-1-thiogalactopyranoside (IPTG) to a final concentration of 0.5 mM and incubation was continued for 18 hours at 14 °C. The culture was centrifuged and the pellet was resuspended in 25 mM Hepes, 500 mM NaCl, 10 % V/V glycerol, 10 mM imidazole, pH 8.0. The suspension was supplemented with 0.01 µg/mL DNAse, 4 µM MgSO$_4$ and the protease inhibitor cOmplete™ EDTA-free Protease Inhibitor Cocktail (Roche). Cells were then sonicated and the mixture was clarified by centrifugation. The supernatant was loaded onto a HisTrap HP column (Cytiva) equilibrated with buffer 25 mM Hepes, 500 mM NaCl, 10% v/v glycerol, 20 mM imidazole, pH 8.0, and His6-tagged mL-GDH$_{180}$ was purified by applying a linear imidazole gradient (20-500 mM). The protein was then further purified by size-exclusion chromatography, using a Superose 6 10/300 GL column (Cytiva) equilibrated in 20 mM Mes, 300 mM NaCl, pH 6.5. mL-GDH$_{180}$ containing fractions, as confirmed by SDS-PAGE, were pooled and used immediately or stored at -80 °C for use in subsequent studies. The protein was quantified by electronic absorption using the molar absorption coefficient of 171,090 M^{-1} cm^{-1}, predicted from the amino acid sequence by the ProtParam tool (http://web.expasy.org/protparam/).

Steady state kinetics

Reactions were performed at 25 °C, using 11.2 nM mL-GDH$_{180}$. Oxidative deamination of L-glutamate catalyzed by mL-GDH$_{180}$ was monitored by following the increase in absorbance at 340 nm resulting from NADH production. To calculate reaction rates, the

molar extinction coefficient 6220 M^{-1} cm^{-1} for NADH at 340 nm was used. The kinetic

parameters k_{cat} and K_M were derived by fitting the Michaelis-Menten equation to initial

rates measured in 100 mM NaH_2PO_4/Na_2HPO_4, pH 7.0, recorded on a Shimadzu UV-120-

02 spectrophotometer. k_{cat} and K_M values and their errors reported correspond to the

mean and maximum dispersion observed, respectively, between at least two independent

measurements performed with different protein preparations. The catalytic efficiency

(k_{cat}/K_M) was calculated as the quotient between k_{cat} and K_M, and its error was obtained by

propagating the error of such parameters.

Cryo-electron microscopy

4 µl of a mixture containing 42 mM L-glutamate, 4 mM NAD^+ and 0.125 mg/ml mL-GDH$_{180}$

was prepared in 20 mM Mes, 300 mM NaCl, pH 6.5, at 10 ºC, while monitoring NADH

production. Samples were applied to Quantifoil R2/2 holey carbon grids and vitrified using

a Leica EM GP2 within the first minute after mixing, ensuring the presence of catalytically

active enzyme molecules. In addition, a similar protein solution was applied onto the same

type of grids bearing an additional thin layer of carbon. Vitrification on these two types of

grids was required to obtain top and side views of mL-GDH$_{180}$ tetramers. Data collection

was carried out in a ThermoFisher Titan Krios G4 transmission electron microscope paired

with a Gatan BioContinuum Imaging Filter and a K3 direct electron detector device, at the

BREM (Basque Resource for Electron Microscopy). 23,747 and 19,038 movie frames were

acquired for the sample on Q2/2 grids and grids supplemented with an additional thin layer

of carbon, respectively. Both collections were performed at a nominal magnification of

×130,000, resulting in a sampling of 0.6462 Å/pixel. Each movie contained 50 frames with

an accumulated dose of 49 e^-/$Å^2$. Movie frames were aligned using MotionCor2 [7] and the

final average included all dose-weighted frames.

The contrast transfer function (CTF) of the micrographs was estimated using

CTFFIND4 [8]. Particles were automatically selected from the micrographs using

autopicking from RELION-4 [9]. An initial set of 667,866 particles was subjected to 2D and

3D classification procedures. The 3D-classification with imposed D2 symmetry resulted in

eight different classes, but only three achieved good definition and high resolution. Three

different conformations were obtained, called the Open (37,641 particles), Closed1

(45,998 particles) and Closed2 (32,441 particles) conformations, with estimated

resolutions of 3.88 Å, 3.54 Å and 3.57 Å, respectively. Each class was symmetry-

expanded, the star files were joined (116,080x4, *i.e.* 464,320 particles) and an average 3D

map of a subunit was refined using a mask and polished to a final resolution of 2.63 Å. A

3D classification within a sphere mask focused on the active site separated three different

states: empty-monomer, with 131,216 particles and 3.07 Å of resolution, cofactor-

monomer, with 168,668 particles and 2.95 Å of resolution, and cofactor/ligand-monomer,

with 112,701 particles and 3.12 Å of resolution. These three groups were crossed with the

three sets of particles obtained in the initial classification from the conformational changes

of the tetramer, resulting into nine groups of particles that were refined independently. The

three maps with signal in the cofactor and substrate binding pockets were subjected to

polishing and ctfrefine to improve the resolution, each one coming from a conformation of

the tetramer: Open-cofactor/ligand-monomer (3.23 Å), Closed1-cofactor/ligand-monomer

(3.11 Å) and Closed2-cofactor/ligand-monomer (3.18 Å). The cryoEM density maps were

postprocessed by local sharpening using EMReady [10], which improved the structural

details and facilitated the atomic model building.

Model fitting into cryo-EM maps was performed using the programs UCSF Chimera [11],

Coot [12] and phenix.real_space_refine [13]. From the 2.63 Å resolution cryoEM map of a mL-

GDH_{180} monomer and an initial model of the protein built by AlphaFold 3 [14], a model of the

polypeptide chain was refined which was then fitted into the other cryo-EM maps. After

removing parts of the model not defined in the maps, performing manual model building

when necessary and placing ligands manually, using Coot [12], where appropriate, atomic

coordinates and B-factors were refined using phenix.real_space_refine [13] with NCS and

secondary structure restraints.

Figures were generated and rendered with UCSF Chimera [11] or PyMOL version 1.8.x

(Schrödinger, LLC).

Cryo-EM maps obtained for mL-GDH$_{180}$ were deposited in the Electron Microscopy Data

Bank under the accession codes EMD-52419 (Open), EMD-52420 (Closed1), EMD-52421

(Closed2), EMD-52422 (empty-monomer), EMD-52423 (cofactor-monomer), EMD-52424

(cofactor/ligand-monomer), EMD-52425 (Open-cofactor/ligand-monomer), EMD-52426

(Closed1-cofactor/ligand-monomer), EMD-52427 (Closed2-cofactor/ligand-monomer),

EMD-52428 (total-monomer) and EMD-52429 (Closed2-cofactor/ligand-tetramer). Atomic

coordinates were deposited in the Protein Data Bank under the accession codes 9HUX

(Open), 9HUY (Closed1), 9HUZ (Closed2), 9HV0 (empty-monomer), 9HV4 (cofactor-

monomer), 9HV5 (cofactor/ligand-monomer) and 9HV6 (total-monomer).

RESULTS

Three conformers of mL-GDH$_{180}$ detected upon incubation with L-glutamate and NAD^{+} reveal modulations of the enzyme quaternary structure

To investigate the molecular basis of mL-GDH$_{180}$ activity, we first determined the kinetic parameters for the oxidative deamination of L-glutamate catalyzed by the enzyme. The estimated k_{cat}, K_M and k_{cat}/K_M values under steady state conditions, in 100 mM NaH$_2$PO$_4$/Na$_2$HPO$_4$ at pH 7.0, were (20 ± 5) s^{-1}, (34 ± 7) mM and (0.6 ± 0.3) s^{-1} mM^{-1} for L-glutamate, and (21 ± 5) s^{-1}, (0.26 ± 0.07) mM and (80 ± 40) s^{-1} mM^{-1} for NAD^{+}, respectively (Figure S1). Then, single-particle cryoEM experiments were performed with mL-GDH$_{180}$ in the presence of NAD^{+} and L-glutamate, during the course of NADH production (see Methods). As previously reported [3], 2D averages for side views of mL-GDH$_{180}$ tetramers revealed a high degree of flexibility at distal ends, where ACT*1-2 and PAS domains reside, and their corresponding densities vanish in cryoEM maps. A 3D-classification of mL-GDH$_{180}$ particles was performed to distinguish alternate conformers of the enzyme. Three mL-GDH$_{180}$ conformers were found, called the Open, Closed1 and Closed2 conformations (Figure 1, Table 1, Figure S2), for which the ACT*3 module, the helical motif 3 (HM3), the catalytic domain and the C-terminal region were defined in each monomer, achieving estimated resolutions of 3.88 Å, 3.54 Å and 3.57 Å, respectively. The three conformers differ in the relative positions of the subunits (Figure 1A, Figure S3A and Movie S1). The catalytic domains in mL-GDH$_{180}$ monomers in contact through their N-terminal segments are closer to each other in the Closed conformations compared to the Open form (Figure 1B).

A comparison of the atomic models derived from the obtained 3D cryoEM maps shed initial insights on the molecular determinants involved in the modulation of the architecture of mL-GDH$_{180}$. An average RMSD of 0.4 Å (1075 aligned residues) was estimated after pairwise comparisons of monomers in the three mL-GDH$_{180}$ conformers, suggesting that

the observed conformational changes were largely related to modifications of the

quaternary structure. Consistent with this, the contact area at the AB-CD interface (Figure

S3B) is 550 Å^2, 750 Å^2 and 800 Å^2, while at the AD-BC interface it is 1200 Å^2, 1000 Å^2 and

1200 Å^2, as measured for the Open, Closed1 and Closed2 forms of mL-GDH$_{180}$,

respectively. At the AB-CD interface of mL-GDH$_{180}$ conformers, intersubunit contacts are

established by hydrogen bonds between residues of the ACT*3 domains in neighboring

monomers (Figure S3B and Table S1). Remarkably, a previously undetected interfacial

interaction was found between the side chains of Ser650 in the HM3 and Glu790 in the

catalytic domain, exclusively in the Closed2 mL-GDH$_{180}$ tetramer (Figure 1B and Figure

S3B). On the other hand, at the AD-BC interface, mL-GDH$_{180}$ monomers connect via

Asp1516:Arg1582 or Arg1504:Glu1562 salt bridges in the Open and Closed1 forms,

respectively, while both interactions were observed in the Closed2 conformer (Figure S3B

and Table S1). Additionally, in all three conformations extensive series of hydrogen bonds

are established that alternately link a conserved set of secondary structure elements in the

C-terminal domain. Most of the interfacial amino acids in mL-GDH$_{180}$ tetramers found upon

incubation with of NAD$^+$ and L-glutamate were previously identified as determinants of the

oligomeric state of the enzyme [3] and are conserved in diverse L-GDHs (Figure S4),

pointing out their functional relevance.

As HM3 in one monomer and the region including residue Glu790 in the catalytic domain

of another subunit approach, so do other portions of the catalytic domains in mL-GDH$_{180}$

monomers linked by the N-terminal segment (Figure 1B). Intriguingly, the density of the

zone 825-838 was poorly defined in the maps of the Open, Closed1 and Closed2 mL-

GDH$_{180}$ tetramers distinguished upon incubation with NAD$^+$ and L-glutamate. At the same

time, the nicotinamide portion of the cofactor molecule placed within the active site

displayed high mobility in the three enzyme conformers, although it appeared to decrease

in the Closed1 and Closed2 forms. When analyzing the active sites in mL-GDH$_{180}$

tetramers, no evidence was found for the presence of a chemical entity other than the
cofactor.

Three conformers of mL-GDH$_{180}$ subunits reveal modulations in tertiary structure
associated with differential active site occupancies

In order to clarify the lack of signal for a ligand other than the cofactor in the cryoEM maps
for mL-GDH$_{180}$ tetramers, we performed a hierarchical classification of all mL-GDH$_{180}$
subunits, whether they were part of the Open, Closed1 or Closed2 conformers, applying a
mask that covered the catalytic domain (Figure S2). In this classification we found three
forms for mL-GDH$_{180}$ monomers, called empty-monomer, cofactor-monomer and
cofactor/ligand-monomer, achieving estimated resolutions of 3.07 Å, 2.95 Å and 3.12 Å,
respectively (Figure 2 and Table 1). The empty-monomer map shows some density
beyond that attributable to the polypeptide chain in the cofactor binding pocket, but does
not indicate, however, strong binding to that site nor does it reveal any signal in the
substrate binding pocket. The cofactor-monomer map, on the other hand, clearly reveals a
cofactor molecule bound to the catalytic cavity of mL-GDH$_{180}$ and the cofactor/ligand-
monomer map discloses both the cofactor as well as an additional compound within the
active site of the enzyme. An average of all these variants resulted in a 2.63 Å resolution
map of a mL-GDH$_{180}$ monomer (Figure S5), which was used to unambiguously model the
polypeptide chain prior to subsequent refinement of atomic coordinates linked to species
with differential occupancy in the active site.

The atomic model derived from the empty-monomer map exposed the arrangement of
residues that, according to previous analyses [2,3], constitute the dinucleotide binding region
(DNBR) and the glutamate binding region (GluBR) of mL-GDH$_{180}$ (Figure 2). The cryoEM
maps of the cofactor- and cofactor/ligand- monomers of mL-GDH$_{180}$ allowed to
unambiguously position a NAD$^+$ molecule in extended conformation in the DNBR of mL-

GDH$_{180}$ (Figure 2), with the pyrophosphate moiety located in the crevice formed between

strands β1 and β5 (residues 953-958 and 1070-1073, respectively) of the catalytic domain

subdomain II (SDII) and stabilized by a favorable interaction with the dipole of the helix

(residues 964-969) connecting strands β1 and β2 (residues 976-982), also called the

dinucleotide binding helix (Figure 3). Also, it was possible to unequivocally place the

adenine ring of NAD$^+$, to assign the conformation of the glycosidic bond to the

nicotinamide ring as *syn*, and to assign the conformation of the adenosine and

nicotinamide riboses to C2' *endo*, in agreement with previous reports for S-GDHs [15].

Differences were observed in the pattern of NAD$^+$-protein interactions established in

cofactor- and cofactor/ligand- mL-GDH$_{180}$ subunits (Figure 2). In the cofactor-monomer,

the adenine ring of the dinucleotide interacts with the side chain (sc) of residue Tyr1079,

the 2' and 3' hydroxyl groups of the adenosine ribose point towards the surface of the

enzyme and hydrogen bond Asp983 (sc) and the main chain (mc) of Asp960, respectively,

phosphates interact with the amide nitrogen of Asp964 and Val965, the nicotinamide

ribose hydrogen bonds Gly1075 (mc) and Asn1115 (mc and sc), and the nicotinamide

amide contacts His923 (sc), Thr929 (sc) and Asp964 (sc). In the cofactor/ligand-monomer,

instead, the adenine of NAD$^+$ interacts with Tyr1079 (sc) as well as His984 (sc), both

hydroxyl groups on the adenosine ribose hydrogen bond Asp960 (sc and mc), the

nicotinamide ribose exhibits stabilizing hydrogen bonds only with Asn1115 (mc and sc)

and the nicotinamide amide contacts only Thr929 (sc) and Asp964 (sc). Thereby, together

with subtle but detectable changes in segments 958-966 and 981-987 contributing to the

DNBR, in the cofactor/ligand-monomer a shift of the cofactor towards the GluBR is

observed, where a novel density element becomes evident (Movie S2). It is possible to

locate a substrate molecule within it so that the α-carbon of the amino acid is 3.3 Å from

the C4 atom of the nicotinamide ring of the cofactor, thus explaining the density bridge

observed between them. Hence, the γ-carboxylate of the ligand establishes salt bridges

with Arg780 (sc).

Remarkably, both segment 914-924 delimiting the active site cleft and portion 886-901

sitting on the glutamate binding pocket adopt different conformations in the empty-,

cofactor- and cofactor/ligand- mL-GDH$_{180}$ monomers (Figure 2). This was in turn

associated with alternate conformations of the region 825-838, found better defined in the

electronic maps of mL-GDH$_{180}$ monomers, which allowed modeling residues 825-827 and

830-838 for the empty- and cofactor- monomers and residues 825 and 829-838 for the

cofactor/ligand-monomer. Thus, in the empty-monomer of mL-GDH$_{180}$ Arg835 points

toward Asp964 and Asp960, and Asp834 points toward His984. When the cofactor is

anchored within the DNBR and interacts with Asp964, Asp960 and His984, Arg835 and

Asp834 are displaced from such positions and the segment 830-838 is less tightly packed

against the catalytic core. When the GluBR is also occupied, the region 831-837 is packed

against the segment 889-894. The amino acid string 825-838 of mL-GDH$_{180}$ is found in L-

GDHs$_{180}$ but not in related low molecular weight enzymes (Figure 3 and Figure S4) [16].

GluBR occupancy is linked to the coupling of the catalytic domains of two adjacent

monomers in the Closed mL-GDH$_{180}$ conformations

To elucidate structural changes in the context of mL-GDH$_{180}$ tetramers, we next crossed

the empty-, cofactor- and cofactor/ligand-monomer classes with the tetrameric classes

Open, Closed1 and Closed2, thus obtaining evidence for the Open-empty-monomer,

Closed1-empty-monomer, Closed2-empty-monomer (3.5 Å, 3.35 Å and 3.48 Å resolution,

respectively), Open-cofactor-monomer, Closed1-cofactor-monomer, Closed2-cofactor-

monomer (3.3 Å, 3.23 Å and 3.28 Å resolution, respectively), Open-cofactor/ligand-

monomer, Closed1-cofactor/ligand-monomer and Closed2-cofactor/ligand-monomer (3.23

Å, 3.11 Å and 3.18 Å resolution, respectively) species (Figure S2). While no differences

were noticeable for the empty- or cofactor- monomers depending on whether they came from Open, Closed1 and Closed2 mL-GDH$_{180}$ tetramers, cryoEM maps disclosed heterogeneous density within the glutamate binding pocket of cofactor/ligand-monomers. The cryoEM map of the Open-cofactor/ligand-monomer accounts for the density within the GluBR that is linked to that of the cofactor nicotinamide ring (Figure 4, see Figure 2 as well), while the former vanishes in the area where the γ-carboxylate of the substrate amino acid would be stabilized. Conversely, in the maps of the Closed1- and Closed2- cofactor/ligand monomers density is found for the ligand in the GluBR mainly towards the area closer to Arg780, such element being smaller for the Closed2 conformer.

As previously noted, changes in the active site of mL-GDH$_{180}$ upon incubation with NAD$^+$ and L-glutamate are associated with structural changes in the segment 825-838 (Figure 2). Remarkably, this remodeled segment establishes an interaction with the region 704-706 in the catalytic domain of a neighboring N-terminally-linked subunit, building a molecular bridge in the Closed tetramers (Figure 4), with a consolidation of this in the Closed2 form (Table 1). Such interaction is only observed when for one monomer there is density simultaneously in both the cofactor and substrate binding pockets (Movie S3). Overall, our results point out that the occupancy of the GluBR of mL-GDH$_{180}$ is linked to the coupling of the catalytic domains of two adjacent subunits, which may be the basis for a regulatory mechanism of turnover.

DISCUSSION

GDHs constitute a large family of enzymes that function in all kingdoms of life at the crossroad between the Krebs cycle and ammonium assimilation. While S-GDHs have been studied for more than half a century, it was not until 2021 that the first experimental evidence for the 3D architecture of an L-GDH was obtained [3]. In that work we determined that the quaternary structure of mL-GDH$_{180}$ is radically different from that of S-GDHs and that such assembly depends on the distinctive N- and C-terminal regions found in L-GDHs [3]. Here we present cryoEM evidence of up to 2.63 Å resolution for mL-GDH$_{180}$ (Table1) obtained in the presence of NAD$^+$ during oxidative deamination of L-glutamate.

We have solved the 3D structure of mL-GDH$_{180}$ species with the DNBR and the GluBR simultaneously occupied as well as NAD$^+$-only charged subunits (Figure 1, Figure 2 and Figure 4). The binding mode of NAD$^+$ to mL-GDH$_{180}$ (Figure 2 and Figure 4) is analogous to that described for *Cs*GDH [15]. The adenine ring of NAD$^+$ is stabilized in mL-GDH$_{180}$ by hydrogen bonds with residues His984 (Pro262 in *Cs*GDH) and Tyr1079 (Asp325), the NH and CO groups of the nicotinamide amide contact Asp964 (Asn240) and Thr929 (Thr209), respectively, the phosphates interact with Asp964 and Val965 (Val241), while the nicotinamide ribose hydrogen bonds Gly1075 (Ala321) and Asn1115 (Asn347). On the other hand, the 2'-OH group of the adenosine ribose, which shares with the 3'-OH group the Asp960 (Phe238) ligand and is additionally surrounded by Asp983 (Gly261) and His984, is stabilized in mL-GDH$_{180}$ in a relatively wider pocket than the equivalent in *Cs*GDH [15]. Considering that His984 and Arg985 (Asp263) could potentially provide stabilizing interactions for the distinctive phosphate attached to the 2'-OH in NADP$^+$, a possible explanation for mL-GDH$_{180}$ NAD$^+$ specificity [17,18] is that this enzyme, as well as related L-GDHs [2,19–24], would have evolved the ability to positively select the cofactor through the interactions established by the 2'-OH group with Asp960 and Asp983, which would, in addition, exert a repulsive effect on the 2'-OP in NADP$^+$. This is consistent with

previous proposals that NAD$^+$ specificity may be dictated by interactions of the 2'-OH group of the adenosine ribose with the Gly-rich loop (GxGxxA/S, as defined in L-GDHs[180] [2]) and with an acidic residue at the so-called position 7 (Asp983 in mL-GDH[180]) of the fingerprint region of the nucleotide binding fold [25,26].

In a pioneering work it was determined that amino acid specificity is determined in *Cs*GDH by the interactions between the side chains of Lys89 and Ser380 with the γ-carboxylate of the substrate [27]. Similarly, the position of Arg780 (Lys89) allows it to form salt bridges with the γ-carboxylate of L-glutamate modeled in the GluBR of mL-GDH[180] (Figure 2). Besides, although the distance between the hydroxyl of Ser1147 (Ser380) of mL-GDH[180] and the γ-carboxylate of L-glutamate (4.3 Å) exceeds that of a hydrogen bond, the position of such residue is equivalent to that of Ser360 in *Cs*GDH. Also, residues Ala886 and Val1144 (Ala163 and Val377), Gly814 and Ala815 (Gly122 and Gly123), Gly781 and Gly782 (Gly90 and Gly91) and Gly1143 (Gly376), previously highlighted for their roles in establishing hydrophobic interactions with the γ- and δ-carbons of L-glutamate, shaping the L-glutamate binding pocket, orienting functional groups in the catalytic center and being located near the binding site of the nicotinamide ring of the dinucleotide, respectively [27], are arranged in cofactor/ligand- mL-GDH[180] monomers in a manner similar to that observed for *Cs*GDH. Overall, these observations highlight the preservation of the determinants involved in amino acid specificity in mL-GDH[180] (and related L-GDHs; see Figure S4 and [2]) as compared to *Cs*GDH (and related S-GDHs).

Given the conservation of active site residues, catalysis would proceed by the same mechanism in S-GDH and L-GDH [3]. Our atomic model of the mL-GDH[180] cofactor/ligand-monomer was built in an attempt to explain the observed density within the GluBR (Figure 2). According to this, the α-carboxylate and amino groups of L-glutamate would adopt in the GluBR of mL-GDH[180] a somewhat different orientation to that described for *Cs*GDH [27]. On the one hand, even thought the α-carboxylate of L-glutamate points towards Lys806

(Lys113), the distance between them (4.1 Å) is relatively long. Besides, the α-carboxylate of the substrate is 6.4 Å away from the side chain of Gln803 (Gln110). On the other hand, the amino group of L-glutamate is 4.9 Å away from the carbonyl of Ala887 (Gly164) and 6.7 Å away from the side chain of the catalytic Asp888 (Asp165). It should be noted, however, that although packing of the region 825-838 against the segment 889-894 would stabilize the position of Asp888, the carboxylate of this residue is not completely defined in the density maps of the mL-GDH$_{180}$ cofactor/ligand-monomer. Nevertheless, experimental data for the cofactor/ligand-monomer (Figure 2 and Figure 4) reveal density consistent with positioning of a group, modeled as the α-carbon of L-glutamate, 3.2 Å from the C4 atom of the cofactor nicotinamide, and density 3-3.5 Å from the Ala887 carbonyl, where the leaving NH$_3$ group would be found upon catalysis.

The accumulated cryoEM evidence for mL-GDH$_{180}$ ([3] and this work) indicates that this enzyme has a very dynamic behavior. In the absence of any ligand, two tetrameric forms of mL-GDH$_{180}$ were found [3], named Open and Closed (4.19 and 6.6 Å resolution, respectively). In the present work we used a saturating (4 mM) concentration of NAD$^+$ and a sub-saturating (42 mM) concentration of L-glutamate (see also Figure S1). Consistent with expectations, evidence of DNBR loading was found in all detected tetrameric forms of the enzyme (Figure 1). Thus, under the conditions tested mL-GDH$_{180}$ samples the conformational space in such a way that the stabilization of the Closed tetramers is promoted. The segment 825-838 adopts alternative conformations depending on the anchoring of the cofactor to the DNBR as well as the presence of L-glutamate (or derivatives thereof) in the GluBR (Figure 2). The geometry of the segment 825-838 in mL-GDH$_{180}$ monomers with the GluBR and the DNBR simultaneously loaded (cofactor/ligand-monomers) is compatible with the consolidation of a contact between such a region and the zone 704-706 of a neighboring subunit linked by the N-terminal region (Figure 4).

Unlike for S-GDH [27,28], we have not found evidence of changes in the relative positions of

the SDI and SDII in the catalytic domain of mL-GDH$_{180}$. Instead, we have detected

changes in inter-monomeric couplings involving residues 650 (in the HM3) and 790 (in the

catalytic domain) as well as regions 825-838 and 704-706 (both in the catalytic domain) of

adjacent mL-GDH$_{180}$ subunits. These, in turn, are accompanied by adjustments in the

interfaces provided by the ACT*3 and C-terminal domains of the enzyme. Transitions of

the tertiary and quaternary structure at long distances from the active site must be much

slower than catalysis and are, therefore, detectable in our experiment. Although given the

nature of our data we cannot draw conclusions about positional and/or compositional

heterogeneity within the GluBR and the DNBR of mL-GDH$_{180}$, and therefore not about

singular events during the enzyme-catalyzed deamination, it is reasonable to expect

catalysis to occur when there is evidence of simultaneous loading of the two regions

(cofactor/ligand-monomer, Figure 4), with the cofactor approaching the substrate binding

pocket (Figure 2). Certainly, for catalysis to take place, a relatively rapid and local

conformational change must occur that allows the side chain of the catalytic Asp888 to

approach the amino group of the substrate for proton abstraction in the first stage of the

reaction [27]. Notably, in the absence of effectors, we have not found evidence of

cooperativity for mL-GDH$_{180}$-catalyzed L-glutamate oxidation, neither at optimal pH (6.5-

7.0) nor at other pHs, as tested in buffers of diverse chemical nature. However, it is

intriguing whether [3] and how [17] this might be different in the presence of modulators of mL-

GDH$_{180}$ activity.

We have discovered that ligand binding to the orthosteric site of mL-GDH$_{180}$ is linked to

changes in the enzyme tertiary structure that affect the arrangement of oligomeric

interfaces nanometers away from the active site [29]. Allosteric regulation usually involves

the inverse pathway, with the binding of molecules to an allosteric site modulating the

catalytic center. It is interesting to consider whether and how the binding of natural or

designed molecules to the 825-838 and/or 704-706 segments of mL-GDH$_{180}$ could impact

on catalysis and how the occupancy of the enzyme active site could make such regulation

more or less permissive [30]. Our work opens new perspectives for atomic resolution studies

on catalysis and regulation of GDHs of medical and industrial relevance.

REFERENCES

1.   Li, M., Li, C., Allen, A., Stanley, C. A. & Smith, T. J. Glutamate dehydrogenase: Structure, allosteric regulation, and role in insulin homeostasis. *Neurochemical Research* **39**, 433–445 (2014).

2.   Miñambres, B., Olivera, E. R., Jensen, R. A. & Luengo, J. M. A new class of glutamate dehydrogenases (GDH): Biochemical and genetic characterization of the first member, the AMP-requiring NAD-specific GDH of Streptomyces clavuligerus. *J. Biol. Chem.* **275**, 39529–39542 (2000).

3.   Lázaro, M. *et al.* 3D architecture and structural flexibility revealed in the subfamily of large glutamate dehydrogenases by a mycobacterial enzyme. *Commun. Biol.* **4**, 684 (2021).

4.   Lang, E. J. M., Cross, P. J., Mittelstädt, G., Jameson, G. B. & Parker, E. J. Allosteric ACTion: The varied ACT domains regulating enzymes of amino-acid metabolism. *Curr. Opin. Struct. Biol.* **29**, 102–111 (2014).

5.   Möglich, A., Ayers, R. A. & Moffat, K. Structure and Signaling Mechanism of Per-ARNT-Sim Domains. *Structure* **17**, 1282–1294 (2009).

6.   Gallant, J. L., Viljoen, A. J., Van Helden, P. D. & Wiid, I. J. F. Glutamate dehydrogenase is required by mycobacterium bovis BCG for resistance to cellular stress. *PLoS One* **11**, 1–14 (2016).

7.   Zheng, S. Q. *et al.* MotionCor2: anisotropic correction of beam-induced motion for improved cryo-electron microscopy. *Nat. Methods* **14**, 331–332 (2017).

8.   Rohou, A. & Grigorieff, N. CTFFIND4: Fast and accurate defocus estimation from electron micrographs. *J. Struct. Biol.* **192**, 216–221 (2015).

9.   Zivanov, J. *et al.* New tools for automated high-resolution cryo-EM structure determination in RELION-3. *Elife* **7**, e42166 (2018).

10. He, J., Li, T. & Huang, S. Y. Improvement of cryo-EM maps by simultaneous local

and non-local deep learning. *Nat. Commun.* **14**, 1–16 (2023).

11.   Pettersen, E. F. *et al.* UCSF Chimera - A visualization system for exploratory

research and analysis. *J. Comput. Chem.* **25**, 1605–1612 (2004).

12.   Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of

Coot. *Acta Crystallogr. Sect. D Biol. Crystallogr.* **66**, 486–501 (2010).

13.   Afonine, P. V. *et al.* Real-space refinement in PHENIX for cryo-EM and

crystallography. *Acta Crystallogr. Sect. D Struct. Biol.* **74**, 531–544 (2018).

14.   Abramson, J. *et al.* Accurate structure prediction of biomolecular interactions with

AlphaFold 3. *Nature* **630**, 493–500 (2024).

15.   Baker, P. J. *et al.* Subunit assembly and active site location in the structure of

glutamate dehydrogenase. *Proteins Struct. Funct. Bioinforma.* **12**, 75–86 (1992).

16.   Britton, K. L., Baker, P. J., Rice, D. W. & Stilllman, T. J. Structural relationship

between the hexameric and tetrameric family of glutamate dehydrogenases. *Eur. J.*

Biochem. **209**, 851–859 (1992).

17.   O'Hare, H. M. *et al.* Regulation of glutamate metabolism by protein kinases in

mycobacteria. *Mol. Microbiol.* **70**, 1408–1423 (2008).

18.   Nott, T. J. *et al.* An intramolecular switch regulates phosphoindependent FHA

domain interactions in Mycobacterium tuberculosis. *Sci. Signal.* **2**, ra12 (2009).

19.   Veronese, F. M., Nyc, J. F., Degani, Y., Brown, D. M. & Smith, E. L. Nicotinamide

adenine dinucleotide specific glutamate dehydrogenase of Neurospora. I.

Purification and molecular properties. *J. Biol. Chem.* **249**, 7922–7928 (1974).

20.   Lu, C. D. & Abdelal, A. T. The gdhB gene of Pseudomonas aeruginosa encodes an

arginine-inducible NAD+-dependent glutamate dehydrogenase which is subject to

allosteric regulation. *J. Bacteriol.* **183**, 490–499 (2001).

21.   Camardella, L. *et al.* The Antarctic Psychrobacter sp. TAD1 has two cold-active

glutamate dehydrogenases with different cofactor specificities. Characterisation of

the NAD+-dependent enzyme. *Comp. Biochem. Physiol. - A Mol. Integr. Physiol.*

131, 559–567 (2002).

22. Kawakami, R., Sakuraba, H. & Ohshima, T. Gene cloning and characterization of the

very large NAD-dependent L-glutamate dehydrogenase from the psychrophile

Janthinobacterium lividum, isolated from cold soil. *J. Bacteriol.* **189**, 5626–5633

(2007).

23. Kawakami, R., Oyama, M., Sakubara, H. & Ohshima, T. The Unique Kinetic

Behavior of the Very Large NAD-Dependent Glutamate Dehydrogenase from

Janthinobacterium lividum. *Biosci. Biotechnol. Biochem.* **74**, 884–887 (2010).

24. Beaufay, F. *et al.* A NAD-dependent glutamate dehydrogenase coordinates

metabolism with cell division in Caulobacter crescentus. *EMBO J.* **34**, 1786–1800

(2015).

25. Baker, P. J., Britton, K. L., Rice, D. W., Rob, A. & Stillman, T. J. Structural

consequences of sequence patterns in the fingerprint region of the nucleotide

binding fold. Implications for nucleotide specificity. *J. Mol. Biol.* **228**, 662–671 (1992).

26. Engel, P. C. Glutamate dehydrogenases: The why and how of coenzyme specificity.

Neurochem. Res. **39**, 426–432 (2014).

27. Stillman, T. J., Baker, P. J., Britton, K. L. & Rice, D. W. Conformational flexibility in

glutamate dehydrogenase: Role of water in substrate recognition and catalysis.

Journal of Molecular Biology **234**, 1131–1139 (1993).

28. Borgnia, M. J. *et al.* Using cryo-EM to map small ligands on dynamic metabolic

enzymes: Studies with glutamate dehydrogenase. *Mol. Pharmacol.* **89**, 645–651

(2016).

29. Perica, T., Chothia, C. & Teichmann, S. A. Evolution of oligomeric state through

geometric coupling of protein interfaces. *Proc. Natl. Acad. Sci.* **109**, 8127–8132

(2012).

30.   Schulze, J. O. *et al.* Bidirectional Allosteric Communication between the ATP-

501   Binding Site and the Regulatory PIF Pocket in PDK1 Protein Kinase. *Cell Chem.*

502   *Biol.* **23**, 1193–1205 (2016).

ACKNOWLEDGMENTS

This work was supported by grants PICT 2019-00989 and PICT 2021-00253 from the Agencia Nacional de Promoción de la Investigación, el Desarrollo Tecnológico y la Innovación (Agencia I+D+i, Argentina) to MNL as well as grants PID2021-124074NB-I00 and PID2021-125946OB-I00 from Agencia Estatal de Investigación of Spain to MV and GJO, respectively. We also thank the Severo Ochoa Center of Excellence Accreditation CEX2021-001136-S. MNL is a career investigator from Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET, Argentina). NC holds a CONICET PhD fellowship. Some of this work was performed at the Basque Resource for Electron Microscopy located at Instituto Biofisika (UPV/EHU, CSIC), supported by the Department of Science, Universities and Innovation and the Innovation Fund of the Basque Government, with additional support from MCIN (Recovery, Transformation and Resilience Plan) and the Basque Government "Biotechnology Complementary Plan Applied to Health" with funding from European Union NextGenerationEU (PRTR-C17.I1; PRTR-C17.I01.P01.S13) (AAAA_ACG_AY_2539/22_05).

AUTHOR CONTRIBUTIONS

ML produced protein, prepared EM samples, performed EM experiments, processed and analyzed EM data and performed structural analyses; NC produced protein, performed kinetic analyses, analyzed EM data and performed structural analyses; JPLA collected and analyzed EM data; RMR performed structural analyses; GJO performed structural analyses; MV designed experiments, supervised EM data collection and processing and analyzed results; MNL designed experiments, supervised biochemical experiments, refined atomic models, analyzed data and wrote the paper. All authors read and corrected the paper.

530 **COMPETING INTERESTS**

531 The authors declare no competing interests.

Figure 1

Three conformers of mL-GDH$_{180}$ detected upon incubation with L-glutamate and NAD$^+$ reveal modulations of the enzyme quaternary structure. (A) CryoEM mL-GDH$_{180}$ maps obtained after incubating the enzyme with L-glutamate and NAD$^+$. The dimensions of each conformer are provided in Å. Maps cover, for each monomer, the region indicated by a dotted rectangle in the diagram shown below (residues 500-1588). ACT*, Aspartate kinase-Chorismate mutase-TyrA-like; PAS, Per-Arnt-Sim; HM, helical motif; CD, catalytic

domain; CTD, C-terminal domain. (B) Approach of the catalytic domains of mL-GDH$_{180}$

subunits linked by the N-terminal region. Segment 825-838, poorly defined in maps of the

Open, Closed1 and Closed2 enzyme forms, approaches the catalytic domain in a

neighboring subunit. In each case, a zoomed in view of the dinucleotide binding region

(DNBR) is shown below. The protein is displayed in ribbon representation and cryoEM

maps are depicted in grey. The nicotinamide portion of the NAD$^+$ molecule (in sticks)

exhibits high mobility in all conformations. Subdomains I and II (SDI and SDII) of the

catalytic domain are colored in yellow and orange, respectively, whereas ACT*3 and HM3

motifs are shown in cyan and green, respectively, for one monomer (the one on the left),

with the neighboring subunit (the one on the right) shown in grey. GluBR, glutamate

binding region.

Figure 2

Three conformers of mL-GDH₁₈₀ subunits reveal modulations in tertiary structure associated with differential active site occupancies. Alternative conformations of mL-GDH₁₈₀ subunits are shown, with the active site either empty or containing NAD⁺ or NAD⁺ and L-glutamate. The average RMSD for 1122 aligned residues in pairwise comparisons

of the atomic models (show in ribbon representation) is 0.7 Å. NAD^+ and L-glutamate are

depicted as sticks as well as highlighted residues. For each species, the catalytic domain

subdomains I and II (SDI, residues 702-926 and 1188-1220; SDII, residues 927-1187) are

colored in yellow and orange, respectively. Interactions established by the segment 825-

838 with other portions of the polypeptide chain are shown in the panels on the left. A

comparison of the atomic models of the empty- (green), cofactor- (blue) and

cofactor/ligand-monomer (gray) is presented in the lower left corner. CryoEM maps are

depicted in grey. H-bonds and salt bridges are displayed as dashed lines. GluBR,

glutamate binding region; DNBR, dinucleotide binding region.

Figure 3

Comparison of the catalytic domain of mL-GDH₁₈₀ with that of S-GDHs. A monomer of

mL-GDH₁₈₀ is shown on the left. Segments 824-838 and 858-882 (in dark green), both

located in the catalytic domain subdomain I (SDI), are absent in S-GDHs [2]. For the 824-

838 portion coordinates were taken from a model calculated with AlphaFold 3 [14]. A

monomer of *Cs*GDH (*Cs*, *Clostridium symbiosum*; PDB 1BGV [27]), considered a

representative S-GDH, is shown on the right. The SDI of S-GDHs presents secondary

structure elements (in dark blue, named according to the consensus nomenclature for S-

GDHs) with conserved roles in oligomerization in the enzyme subfamily, which are not

present in mL-GDH₁₈₀ or related L-GDHs. For both mL-GDH₁₈₀ and *Cs*GDH, the six-

stranded 432156 β-sheet that shapes the DNBR in the catalytic domain subdomain II

(SDII) is shown in grey; the linker regions β4-β5 (in light green) and β5-β6 (in cyan)

encompass a larger number of residues in mL-GDH₁₈₀. For the two proteins, the SDI and

SDII are shown, except for otherwise highlighted portions, in yellow and orange,

respectively.

Figure 4

GluBR occupancy is linked to the coupling of the catalytic domain of two adjacent

monomers in the Closed conformations. The atomic model of the cofactor/ligand-

monomer of mL-GDH$_{180}$ is shown together with maps of this species obtained from Open,

Closed1 or Closed2 tetramers (top panel). NAD$^+$ and L-glutamate as well as highlighted

residues are depicted as sticks. The catalytic domain subdomains I and II (SDI and SDII)

are colored in yellow and orange, respectively. In the context of tetramers containing a

cofactor/ligand-monomer (cofactor/ligand-tetramers; bottom panel), in a progression

towards the Closed1 and Closed2 conformers, the stabilization of an interaction between

segment 825-838 of one subunit (colored, on the left) and segment 704-706 of the

neighboring one (in grey, on the right) is observed. The atomic coordinates of the subunit

on the left correspond in each case to those of a cofactor/ligand-monomer, replacing the

equivalent ones presented in Figure 1. H-bonds and salt bridges are displayed as dashed

lines. CryoEM maps are displayed in grey.

    TABLES

604   **Table 1. CryoEM data collection and processing.** LS, local sharpening.

	Open (EMD-52419)	Closed1 (EMD-52420)	Closed2 (EMD-52421)
Data collection and processing			
Magnification	130,000	130,000	130,000
Voltage (kV)	300	300	300
Electron dose ($e^-/Å^2$)	0.979	0.979	0.979
Defocus range (μm)	1.0-1.6	1.0-1.6	1.0-1.6
Pixel size (Å)	1.29	1.29	1.29
Symmetry imposed	D2	D2	D2
Initial particles images (no.)	667,866	667,866	667,866
Final particles images (no.)	37,641	45,998	32,441
Map resolution (Å)	3.88	3.54	3.57
FSC threshold	0.143	0.143	0.143
Map sharpening B factor ($Å^2$)	-50	-50	-50
Refinement			
Model resolution (Å)	3.88	3.54	3.57
Model composition			
Non-hydrogen atoms	33576	33576	33576
Protein residues	4300	4300	4300
Ligands	4 NAD^+	4 NAD^+	4 NAD^+
B factors ($Å^2$)			
Protein	224.33	189.71	159.33
Ligands	273.06	208.84	186.14
R.m.s. deviations			
Bond lengths (Å)	0.001	0.002	0.002
Bond angles (°)	0.371	0.409	0.395
Validation			
MolProbity score	1.69	1.62	1.50
Clashscore	10.54	11.11	9.42
Poor rotamers (%)	0.11	0	0.11
Ramachandran plot			
Favored (%)	97.20	97.76	98.55
Allowed (%)	2.80	2.24	1.45
Disallowed (%)	0	0	0
CC (mask)	0.83	0.77	0.83
CC (box)	0.91	0.87	0.89
CC (peaks)	0.77	0.70	0.77
CC (volume)	0.82	0.76	0.82
Mean CC for ligands	0.85	0.82	0.80

	empty-monomer (EMD-52422)	cofactor-monomer (EMD-52423)	cofactor/ligand-monomer (EMD-52424)
Data collection and processing			
Magnification	130,000	130,000	130,000
Voltage (kV)	300	300	300
Electron dose (e$^-$/Å^2)	0.979	0.979	0.979
Defocus range (μm)	1.0-1.6	1.0-1.6	1.0-1.6
Pixel size (Å)	1.29	1.29	1.29
Symmetry imposed	C1	C1	C1
Initial particles images (no.)	464,320	464,320	464,320
Final particles images (no.)	131,216	168,668	112,701
Map resolution (Å)	3.07	2.95	3.12
FSC threshold	0.143	0.143	0.143
Map sharpening B factor	LS	LS	LS
Refinement			
Model resolution (Å)	3.07	2.95	3.12
Model composition			
Non-hydrogen atoms	8717	8761	8760
Protein residues	1124	1124	1123
Ligands	-	1 NAD$^+$	1 NAD$^+$/ 1 L-Glu
B factors (Å^2)			
Protein	64.32	62.58	72.09
Ligands	-	56.65	81.37
R.m.s. deviations			
Bond lengths (Å)	0.004	0.004	0.004
Bond angles (°)	0.691	0.721	0.693
Validation			
MolProbity score	1.37	1.27	1.37
Clashscore	5.53	5.16	5.80
Poor rotamers (%)	0.55	0.55	0.11
Ramachandran plot			
Favored (%)	97.68	98.48	97.77
Allowed (%)	2.32	1.52	2.23
Disallowed (%)	0	0	0
CC (mask)	0.87	0.88	0.87
CC (box)	0.89	0.89	0.89
CC (peaks)	0.81	0.81	0.79
CC (volume)	0.88	0.88	0.87
Mean CC for ligands	-	0.91	0.84

	Open-cofactor/ligand-monomer (EMD-52425)	Closed1-cofactor/ligand-monomer (EMD-52426)	Closed2-cofactor/ligand-monomer (EMD-52427)
Data collection and processing			
Magnification	130,000	130,000	130,000
Voltage (kV)	300	300	300
Electron dose (e⁻/Å²)	0.979	0.979	0.979
Defocus range (μm)	1.0-1.6	1.0-1.6	1.0-1.6
Pixel size (Å)	0.646	0.646	0.646
Symmetry imposed	C1	C1	C1
Initial particles images (no.)	112,701	112,701	112,701
Final particles images (no.)	33,330	46,380	32,991
Map resolution (Å)	3.23	3.11	3.18
FSC threshold	0.143	0.143	0.143
Map sharpening B factor	LS	LS	LS
Refinement			
Model resolution (Å)			
Model composition			
Non-hydrogen atoms			
Protein residues			
Ligands			
B factors (Å²)			
Protein			
Ligands			
R.m.s. deviations			
Bond lengths (Å)			
Bond angles (°)			
Validation			
MolProbity score			
Clashscore			
Poor rotamers (%)			
Ramachandran plot			
Favored (%)			
Allowed (%)			
Disallowed (%)			
CC (mask)			
CC (box)			
CC (peaks)			
CC (volume)			
Mean CC for ligands			

	total-monomer (EMD-52428)	Closed2-cofactor/ligand-tetramer (EMD-52429)
Data collection and processing		
Magnification	130,000	130,000
Voltage (kV)	300	300
Electron dose (e⁻/Å^2)	0.979	0.979
Defocus range (μm)	1.0-1.6	1.0-1.6
Pixel size (Å)	0.646	1.29
Symmetry imposed	C1	C1
Initial particles images (no.)	464,320	112,701
Final particles images (no.)	464,320	32,991
Map resolution (Å)	2.63	4.1
FSC threshold	0.143	0.143
Map sharpening B factor	LS	-50
Refinement		
Model resolution (Å)	2.63	
Model composition		
Non-hydrogen atoms	8683	
Protein residues	1113	
Ligands	1 NAD$^+$	
B factors (Å^2)		
Protein	56.19	
Ligands	48.51	
R.m.s. deviations		
Bond lengths (Å)	0.002	
Bond angles (°)	0.467	
Validation		
MolProbity score	1.39	
Clashscore	7.12	
Poor rotamers (%)	0.55	
Ramachandran plot		
Favored (%)	99.10	
Allowed (%)	0.90	
Disallowed (%)	0	
CC (mask)	0.80	
CC (box)	0.68	
CC (peaks)	0.65	
CC (volume)	0.78	
Mean CC for ligands	0.81	