Tertiary and quaternary structure remodeling by occupancy of the substrate

binding pocket in a large glutamate dehydrogenase

Melisa Lázaro [1,#], Nicolás Chamorro [2,#], Jorge P. López-Alonso [3,4], Diego Charro [1],

Rodolfo M. Rasia [2,5], Gonzalo Jiménez-Osés [1,6], Mikel Valle [1,*], María-Natalia Lisa

[2,5,*]

[1] Center for Cooperative Research in Biosciences (CIC bioGUNE), Basque Research

and Technology Alliance (BRTA), Bizkaia Technology Park, Building 801A, 48160

Derio, Spain.

[2] Instituto de Biología Molecular y Celular de Rosario (IBR CONICET-UNR), Ocampo

y Esmeralda, Rosario S2002LRK, Argentina.

[3] Instituto Biofisika (UPV/EHU, CSIC), 48940 Leioa, Spain.

[4] Basque Resource for Electron Microscopy, 48940 Leioa, Spain.

[5] Plataforma de Biología Estructural y Metabolómica (PLABEM), Ocampo y

Esmeralda, Rosario S2002LRK, Argentina.

[6] Ikerbasque, Basque Foundation for Science, 48013 Bilbao, Spain.

# Equal contribution

* Correspondence: MNL: lisa@ibr-conicet.gov.ar (+54 341 423 7070; MV:

mvalle@cicbiogune.es (+34 946 572 503)

Running title: Ligand-driven remodeling of mL-GDH$_{180}$

Total number of manuscript pages, 42; supplementary material pages, 16; tables, 1;

figures, 5

Description of supplementary material including filenames:

Supplementary material.doc: Supplementary figures and tables.

Movie S1.mp4: Changes in the quaternary structure of mL-GDH$_{180}$ detected by cryo-

EM in the presence of L-glutamate and NAD$^+$.

Movie S2.mp4: Changes in the active site of mL-GDH$_{180}$ detected by cryo-EM in the

presence of L-glutamate and NAD$^+$.

Movie S3.mp4: In the presence of L-glutamate and NAD$^+$, a contact is established

between the catalytic domains of neighboring mL-GDH$_{180}$ subunits connected

through their N-terminal regions.

ABSTRACT

Glutamate dehydrogenases (GDHs) catalyze the oxidative deamination of L-

glutamate to 2-oxoglutarate using $NAD(P)^+$ as a cofactor. The large type of GDHs (L-

GDHs) displays a dynamic homotetrameric architecture that alternates between open

and closed states. However, the catalytic mechanism and the functional relevance of

the large conformational changes in L-GDHs remain poorly understood. Here, we

use cryo-EM to investigate the structure and the conformational landscape of the

mycobacterial L-GDH composed of 180 kDa subunits ($mL-GDH_{180}$) when incubated

with L-glutamate and NAD^+. Classification of the heterogeneous population of

tetramers reveals opening-closing motions and sorting of individual subunits resolves

the occupancy of the cofactor and substrate binding pockets. Cryo-EM maps show

that ligand binding to the glutamate binding pocket is accompanied by structural

changes in a region approximately two nanometers away from the active site, leading

to the formation of a previously undetected interaction between the catalytic domains

of neighboring subunits in $mL-GDH_{180}$ closed tetrameric states. Our findings indicate

that the occupancy of the substrate binding site of $mL-GDH_{180}$ is linked to a

remodeling of both the tertiary and quaternary structure of the enzyme.

KEYWORDS

glutamate dehydrogenase, large GDH ($mL-GDH_{180}$), cryo-electron microscopy,

conformational dynamics, quaternary structure remodeling, substrate binding pocket,

Mycobacterium metabolism

STATEMENT FOR A BROADER AUDIENCE

This work reveals how the binding of L-glutamate and NAD$^+$ reshapes the

architecture of a large glutamate dehydrogenase, linking active site occupancy to

long-range structural remodeling. By capturing previously unseen conformational

transitions with cryo-electron microscopy, we provide insights into the molecular logic

of enzyme function in mycobacteria. These findings establish a framework to

understand how structural plasticity supports metabolic control.

INTRODUCTION

Amino acid dehydrogenases have long attracted interest due to their physiological functions and potential industrial applications. Among them, glutamate dehydrogenases (GDHs; EC 1.4.1.2) catalyze the reaction L-glutamate + NAD(P)$^+$ + H$_2$O <=> 2-oxoglutarate + NAD(P)H + H$^+$ + NH$_3$, at the crossroads of carbon and nitrogen metabolism. These enzymes are oligomeric and are classified based on the molecular weight of their subunits. Small GDHs (S-GDHs), present in prokaryotes and eukaryotes, consist of 50 kDa subunits [1]. The conserved catalytic domain spans nearly the full length of the polypeptide chain of S-GDHs and mediates oligomerization. Biochemical and structural studies on S-GDHs have enabled detailed characterization of their catalytic and regulatory mechanisms. In contrast, much less is known about large GDHs (L-GDHs), which are found in prokaryotes and lower eukaryotes [2]. This group includes enzymes with 115 kDa subunits (L-GDHs$_{115}$) and those with 180 kDa subunits (L-GDHs$_{180}$), such as the L-GDH$_{180}$ from mycobacteria (mL-GDH$_{180}$). L-GDHs feature distinctive and extended N- and C-terminal regions flanking the catalytic core, which may confer regulatory or structural roles [3].

In a previous work, we demonstrated that the L-GDH$_{180}$ from *Mycobacterium smegmatis* (mL-GDH$_{180}$; 76% sequence identity with the L-GDH$_{180}$ from *Mycobacterium tuberculosis*) assembles into a homotetramer, with a quaternary structure markedly different from that of S-GDHs [3]. Such architecture is stabilized by interactions involving the N-terminal region, which contains ACT- and PAS-type domains typically associated with regulatory functions [4,5], as well as a C-terminal domain of unknown function. Despite such advances, the conformational landscape explored by L-GDHs during catalysis remained poorly unknown.

Here we report cryo-EM structures of mL-GDH$_{180}$ at resolutions up to 3 Å, obtained in

the presence of NAD$^+$ and L-glutamate. We reveal that ligand binding to the

glutamate binding pocket is accompanied by changes in the tertiary structure in a

region approximately 2 nm distant from the active site, promoting interactions

between the catalytic domains of neighboring monomers and modulating the

quaternary structure.

Despite extensive efforts to understand the molecular basis of GDHs activity and

regulation, structural data on ternary complexes containing enzyme, cofactor, and

substrate are scarce, particularly for NAD$^+$-dependent L-GDHs. Here, we present

cryo-EM evidence for enzyme species present upon incubation with L-glutamate and

NAD$^+$ of a key L-GDH$_{180}$ for the virulence of a long-attacked pathogen [6], offering new

insights into the conformational plasticity of L-GDHs and setting the stage for a

deeper understanding of their regulation.

RESULTS

Three conformers of mL-GDH$_{180}$ detected upon incubation with L-glutamate and NAD$^+$ reveal modulations of the enzyme quaternary structure

To investigate the molecular basis of mL-GDH$_{180}$ activity, we first determined its kinetic parameters for the oxidative deamination of L-glutamate. Under steady state conditions, the reaction displayed a k_{cat} of (15 ± 2) s^{-1}; the estimated K_M values were (40 ± 10) mM for L-glutamate and (0.19 ± 0.06) mM for NAD$^+$, yielding catalytic efficiencies (k_{cat}/K_M) of (0.4 ± 0.1) mM^{-1} s^{-1} and (80 ± 30) mM^{-1} s^{-1}, respectively (Figure S1A), as determined in 50 mM Hepes, 50 mM NaCl, at pH 6.8, near the apparent pH optimum of the reaction (Figure S1B). Single-particle cryo-EM experiments were then conducted on mL-GDH$_{180}$ incubated with NAD$^+$ and L-glutamate under NADH-producing conditions, during early turnover (Figure S1C; see also Materials and methods). As previously reported [3], 2D class averages revealed high flexibility at the distal ends of the tetramers, where the ACT*1-2 and PAS domains are located (Figure S2). This flexibility resulted in vanishing density for residues 1-499 in the cryo-EM maps. Subsequently, 3D-classification of mL-GDH$_{180}$ particles revealed three conformational states of the enzyme, named Open (32.4%), Closed1 (39.6%) and Closed2 (28%) (Figure 1, Table I, Figure S3). In these conformers, the ACT*3 module, the helical motif 3 (HM3), the catalytic domain and the C-terminal region were defined in each monomer, achieving estimated global resolutions of 3.88 Å, 3.54 Å and 3.57 Å, respectively. The conformers differ primarily in the relative positions of the subunits (Figure 1A, Figure S4A and Movie S1). In the Closed forms, the catalytic domains of adjacent subunits connected through their N-terminal segments are found in closer proximity compared to the Open state (Figure 1B).

Comparison of the atomic models derived from the 3D cryo-EM maps provided initial

insights on the molecular determinants underlying the modulation of mL-GDH$_{180}$

architecture. Pairwise structural alignments of monomers from the three conformers,

based on all modeled α-carbons, yielded an average RMSD (root-mean-square

deviation) of 0.4 Å (calculated over 1075 aligned residues), indicating that the

observed conformational variability primarily reflects quaternary rearrangements.

Consistent with this, the intersubunit contact area at the AB-CD interface (Figure

S4B) was measured [7] as 550 Å^2, 750 Å^2 and 800 Å^2, and at the AD-BC interface as

1200 Å^2, 1000 Å^2 and 1200 Å^2, for the Open, Closed1 and Closed2 forms,

respectively. At the AB-CD interface, hydrogen bonds between residues in the ACT*3

domains of neighboring monomers mediate the intersubunit contacts in all three

conformers (Figure S4B and Table SI). Remarkably, a previously undetected

interfacial interaction was found in the Closed2 tetramer, involving a hydrogen bond

between the side chains of Ser650 (in the HM3) and Glu790 (in the catalytic domain)

(Figure 1B and Figure S4B). At the AD-BC interface, distinct salt bridge combinations

are found in the different conformers, with Asp1516:Arg1582 present in the Open

form, Arg1504:Glu1562 in the Closed1 and both simultaneously observed in the

Closed2 (Figure S4B and Table SI). Additionally, in all three conformations, extensive

hydrogen bonds networks link a conserved set of secondary structure elements in

the C-terminal domain, with alternative arrangements in each state. Most of the

interfacial residues observed in mL-GDH$_{180}$ tetramers found in the presence of NAD$^+$

and L-glutamate had been previously identified as contributors to oligomer

stabilization [3] and are conserved across diverse L-GDHs (Figure S5), suggesting a

shared structural basis for quaternary modulation within this enzyme family.

As HM3 in one monomer and the region encompassing Glu790 in the catalytic

domain of an adjacent subunit move closer, other portions of the catalytic domains in

mL-GDH$_{180}$ monomers linked via their N-terminal segments also shift toward each

other (Figure 1B). Intriguingly, the density corresponding to residues 825-838 was

poorly defined in the maps of the Open, Closed1 and Closed2 conformers

distinguished in the presence of NAD$^+$ and L-glutamate. At the same time, the

nicotinamide moiety of the cofactor molecule placed within the active site appeared

highly mobile in all three species, although this mobility seemed reduced in the

Closed1 and Closed2 forms. No additional density was detected within the active

sites beyond that attributable to the cofactor.

Three conformers of mL-GDH$_{180}$ subunits reveal modulations in tertiary

structure associated with differential active site occupancies

To clarify the absence of signal for any ligand other than the cofactor in the cryo-EM

maps of mL-GDH$_{180}$ tetramers, we performed a hierarchical classification of all mL-

GDH$_{180}$ subunits, regardless of whether they belonged to the Open, Closed1 or

Closed2 conformers, using a mask that covered the catalytic domain (Figure S3).

This classification yielded three distinct forms for mL-GDH$_{180}$ monomers, termed

empty-monomer, cofactor-monomer and cofactor/ligand-monomer, with estimated

resolutions of 3.07 Å, 2.95 Å and 3.12 Å, respectively (Figure 2 and Table I). The

empty-monomer map shows faint density within the cofactor binding pocket, but does

not reveal strong binding to that site, nor any signal in the substrate binding pocket.

In contrast, the cofactor-monomer map clearly displays a bound cofactor molecule in

the catalytic cavity and the cofactor/ligand-monomer map discloses both the cofactor

and an additional density element within the active site (Figure 2 and Figure S6). An

average of all these variants resulted in a 2.63 Å resolution map of the mL-GDH$_{180}$

monomer (Figure S6 and Figure S7), which was used to model the polypeptide

segment comprising residues 463–1588 before refining the atomic coordinates for

each occupancy state.

The atomic model derived from the empty-monomer map revealed the arrangement

of residues that, according to previous analyses [2,3], define the dinucleotide binding

region (DNBR) and the glutamate binding region (GluBR) of mL-GDH$_{180}$ (Figure 2).

The cryo-EM maps of the cofactor- and cofactor/ligand- monomers of mL-GDH$_{180}$

allowed to place a NAD$^+$ molecule in extended conformation within the DNBR (Figure

2). According to our models, the pyrophosphate moiety is located in the crevice

formed between strands β1 and β5 (residues 953-958 and 1070-1073, respectively)

of subdomain II of the catalytic domain (SDII) and is stabilized by the dipole of the

helix comprising residues 964-969, which connects strands β1 and β2 (residues 976-

982) and is known as the dinucleotide binding helix (Figure 3). The adenine ring of

NAD$^+$ fits well in the density, the glycosidic bond to the nicotinamide ring adopts a

syn conformation and both riboses display C2' *endo* puckering, in agreement with

previous reports for S-GDHs [8].

Notably, the pattern of NAD$^+$-protein interactions differs between cofactor- and

cofactor/ligand- mL-GDH$_{180}$ monomers (Figure 2). In the cofactor-monomer, the

adenine ring interacts with the side chain (sc) of Tyr1079, the 2' and 3' hydroxyl

groups of the adenosine ribose point towards the surface of the protein and hydrogen

bond Asp983 (sc) and the main chain (mc) of Asp960, respectively, phosphates

interact with the amide nitrogen of Asp964 and Val965, the nicotinamide ribose

hydrogen bonds Gly1075 (mc) and Asn1115 (mc and sc), and the nicotinamide

amide contacts His923 (sc), Thr929 (sc) and Asp964 (sc). In contrast, in the

cofactor/ligand-monomer, the adenine of NAD^+ interacts with Tyr1079 (sc) as well as

His984 (sc), both adenosine hydroxyls hydrogen bond Asp960 (sc and mc), the

nicotinamide ribose interacts only with Asn1115 (mc and sc) and the nicotinamide

amide contacts only Thr929 (sc) and Asp964 (sc). These differences, along with

subtle but detectable conformational changes in segments 958-966 and 981-987 that

contribute to the DNBR, correlate with a shift of the cofactor toward the GluBR in the

cofactor/ligand-monomer. In this state, a novel density element becomes evident

(Movie S2), within which it is possible to position a substrate molecule (CC = 0.79)

such that the α-carbon of the amino acid lies 3.3 Å from the C4 atom of the

nicotinamide ring of the cofactor, explaining the continuous density observed

between them. Hence, the γ-carboxylate of the ligand forms salt bridges with Arg780

(sc).

Strikingly, segment 914-924, which lines the active site cleft, and region 886-901,

which borders the glutamate binding pocket, adopt distinct conformations in the

empty-, cofactor- and cofactor/ligand- mL-GDH$_{180}$ monomers (Figure 2). These

variations are associated with alternate conformations of segment 825-838, which is

better defined in the maps of the monomers than in the tetrameric reconstructions.

Thus, residues 825-827 and 830-838 could be built for the empty- and cofactor-

monomers, while residues 825 and 829-838 were modeled for the cofactor/ligand-

monomer. In the empty-monomer, Arg835 is directed toward Asp964 and Asp960,

while Asp834 faces His984. When the cofactor binds within the DNBR, engaging

residues Asp964, Asp960 and His984, Arg835 and Asp834 shift from their positions

and the segment 830-838 loosens its packing against the catalytic core. When the

GluBR is also occupied, segment 831-837 packs against residues 889-894. The

amino acid string 825-838 of mL-GDH$_{180}$ is found in L-GDHs$_{180}$ but not in related low

molecular weight enzymes (Figure 3 and Figure S5) [9].

GluBR occupancy is linked to the coupling of the catalytic domains of two

adjacent monomers in the Closed mL-GDH$_{180}$ conformations

To elucidate structural changes in the context of mL-GDH$_{180}$ tetramers, we next

discriminated the monomeric conformers according to whether they originated from

Open, Closed1 and Closed2. This analysis revealed nine distinct species: Open-

empty-monomer, Closed1-empty-monomer, Closed2-empty-monomer (3.5 Å, 3.35 Å

and 3.48 Å resolution, respectively), Open-cofactor-monomer, Closed1-cofactor-

monomer, Closed2-cofactor-monomer (3.3 Å, 3.23 Å and 3.28 Å), Open-

cofactor/ligand-monomer, Closed1-cofactor/ligand-monomer and Closed2-

cofactor/ligand-monomer (3.23 Å, 3.11 Å and 3.18 Å) (Figure S3).

No discernable structural differences were observed among the empty- or cofactor-

monomers across the tetrameric states. In contrast, cryo-EM maps of the

cofactor/ligand-monomers displayed heterogeneity within the glutamate binding

pocket. In the Open-cofactor/ligand-monomer map, a continuous density extends

from the cofactor nicotinamide ring into the GluBR (Figure 4, see also Figure 2),

although it fades in the region where the γ-carboxylate of the substrate would be

stabilized. Conversely, the maps of the Closed1- and Closed2-cofactor/ligand

monomers display ligand density within the GluBR primarily near Arg780, with a

weaker signal in the Closed2 state.

As noted earlier, incubation of mL-GDH$_{180}$ with NAD$^+$ and L-glutamate induces

remodeling of the region 825-838 (Figure 2). Strikingly, this segment interacts with

residues 704-706 of the catalytic domain of a neighboring N-terminally-linked subunit,

giving rise to an additional molecular bridge in the Closed tetramers (Figure 4). This

intersubunit interaction appears strengthened in the Closed2 state (Table I).

Importantly, it is only observed when the monomer contributing the 825-838 region

has both its cofactor and substrate binding pockets simultaneously occupied. The

interaction occurs approximately 2 nm from the active site of the ligand-loaded

monomer (Movie S3), linking GluBR occupancy to the physical contact between

adjacent catalytic domains and suggesting a structural basis for turnover regulation.

DISCUSSION

GDHs constitute a large family of enzymes that function in all kingdoms of life at the

crossroad between the Krebs cycle and ammonium assimilation. While S-GDHs have

been studied for over half a century, it was not until 2021 that the first experimental

evidence of the 3D architecture of an L-GDH was obtained [3]. In that work, we

determined that the quaternary structure of mL-GDH$_{180}$ differs radically from that of

S-GDHs, and that such assembly is shaped by the distinctive N- and C-terminal

extensions characteristic of L-GDHs [3]. Here we present cryo-EM data reaching

resolutions up to 2.63 Å for mL-GDH$_{180}$ (Table I), captured in the presence of NAD$^+$

during the oxidative deamination of L-glutamate.

Our 3D classification of subunits successfully separated mL-GDH$_{180}$ monomers with

both the DNBR and GluBR simultaneously occupied, as well as monomers bound to

NAD$^+$ alone (Figure 1, Figure 2 and Figure 4). The NAD$^+$ binding mode in mL-GDH$_{180}$

(Figure 2 and Figure 4) mirrors that described for *Cs*GDH (S-GDH from *Clostridium*

symbiosum) [8]. The adenine ring is stabilized by hydrogen bonds with His984 (Pro262

in *Cs*GDH) and Tyr1079 (Asp325), the NH and CO groups of the nicotinamide amide

contact Asp964 (Asn240) and Thr929 (Thr209), respectively, the phosphates interact

with Asp964 and Val965 (Val241), while the nicotinamide ribose hydrogen bonds

Gly1075 (Ala321) and Asn1115 (Asn347). On the other hand, the 2'-OH of the

adenosine ribose, along with the 3'-OH, interacts with Asp960 (Phe238) and is

additionally surrounded by Asp983 (Gly261) and His984, occupying a relatively more

spacious pocket than in *Cs*GDH [8]. Considering that His984 and Arg985 (Asp263)

might stabilize the 2'-phosphate of NADP$^+$, one possible explanation for mL-GDH$_{180}$

specificity for NAD$^+$ [10,11] is that this enzyme, as well as related L-GDHs [2,12–17],

evolved to favor NAD$^+$ via the interactions of the 2'-OH group with Asp960 and

Asp983. These residues might also exert a repulsive effect on the 2'-phosphate of

NADP$^+$. This interpretation aligns with previous suggestions that NAD$^+$ specificity is

determined by interactions between the 2'-OH of the adenosine ribose and both the

Gly-rich loop (GxGxxA/S, as defined in L-GDHs$_{180}$ [2]) and an acidic residue at the so-

called position 7 (Asp983 in mL-GDH$_{180}$) of the fingerprint region of the nucleotide

binding fold [18,19].

A seminal study on *Cs*GDH established that amino acid specificity is conferred by

Lys89 and Ser380, which interact with the γ-carboxylate of the substrate [20]. In mL-

GDH$_{180}$, Arg780 (Lys89) is suitably positioned to form salt bridges with the γ-

carboxylate of L-glutamate (Figure 2). Though the hydroxyl of Ser1147 (Ser380) lies

4.3 Å from the γ-carboxylate of L-glutamate, slightly beyond hydrogen bond range, its

position corresponds to that of Ser360 in *Cs*GDH. Also, residues Ala886 and

Val1144 (Ala163 and Val377), Gly814 and Ala815 (Gly122 and Gly123), Gly781 and

Gly782 (Gly90 and Gly91) and Gly1143 (Gly376), previously highlighted for their

roles in establishing hydrophobic interactions with the γ- and δ-carbons of L-

glutamate, shaping the L-glutamate binding pocket, orienting functional groups in the

catalytic center and being located near the binding site of the nicotinamide ring of the

dinucleotide, respectively [20], are similarly arranged in the mL-GDH$_{180}$ cofactor/ligand-

bound monomers. This structural similitudes underscore the conservation of key

amino acid specificity determinants in mL-GDH$_{180}$ (and related L-GDHs; see Figure

S5 and [2]) and S-GDHs.

Our steady-state data support an ordered sequential mechanism for the oxidative

deamination of L-glutamate catalyzed by mL-GDH$_{180}$, in which NAD$^+$ binds first to the

active site, followed by L-glutamate; chemistry then ensues, 2-oxoglutarate is

released first, and NADH leaves last (Figure S1D and S1E). In line with this, our

cryo-EM 3D classifications recovered empty, cofactor and cofactor/ligand monomer species but not a distinct ligand-only class of subunits. Moreover, given the conservation of residues that mediate catalysis, L-GDHs are expected to function similarly to S-GDHs [3,20]. Our atomic model of the mL-GDH$_{180}$ cofactor/ligand-monomer aimed to account for the density observed in the GluBR (Figure 2). In this model, the α-carboxylate and amino groups of L-glutamate adopt a slightly different orientation than in *Cs*GDH [20]. Although the α-carboxylate of L-glutamate faces Lys806 (Lys113), their separation (4.1 Å) is longer than ideal for direct interaction. Similarly, the α-carboxylate of the substrate lies 6.4 Å from the side chain of Gln803 (Gln110), and the amino group is 4.9 Å from the carbonyl of Ala887 (Gly164) and 6.7 Å from the side chain of the catalytic Asp888 (Asp165). It is worth noting that, at our working resolution, we cannot detect water molecules that might contribute to substrate stabilization within the active site. Besides, the packing of residues 825-838 against segment 889-894 may help stabilize the position of Asp888, even though its side chain is only partially defined in cryo-EM maps. Certainly, catalysis is expected to involve a localized conformational adjustment that enables the side chain of Asp888 to approach the substrate's amino group for proton abstraction in the first step of the reaction [20]. Nevertheless, the cofactor/ligand-monomer maps (Figure 2 and Figure 4) reveal density corresponding to a group modeled as the α-carbon of L-glutamate, located 3.2 Å from the C4 atom of the cofactor nicotinamide, and density 3-3.5 Å from the Ala887 carbonyl, consistent with the position of the leaving NH_3 group during catalysis [20].

Taken together with previous findings [3], our data confirm that mL-GDH$_{180}$ exhibits pronounced conformational flexibility. In the absence of ligands, the enzyme populates two tetrameric forms, Open and Closed, detected at 4.19 and 6.6 Å

resolution, respectively [3]. In the present study, under conditions of saturating NAD$^+$

(4 mM) and sub-saturating L-glutamate (42 mM) (see also Figure S1), three

tetrameric states, Open, Closed1 and Closed2, were observed, with evidence of

DNBR occupancy across the full ensemble (Figure 1). Thus, compared to the ligand-

free state, these conditions appear to shift the conformational equilibrium toward the

Closed forms. The segment 825-838 adopts distinct geometries depending on

whether the cofactor is anchored to the DNBR and whether L-glutamate (or

derivatives thereof) is present in the GluBR (Figure 2). In monomers where both

regions are simultaneously occupied (cofactor/ligand-monomers), the conformation

of this segment is compatible with an interaction with residues 704-706 of the N-

terminally linked neighboring subunit in Closed tetramers (Figure 4).

Unlike in S-GDH [20,21], we did not observe changes in the relative positions of SDI and

SDII within the catalytic domain of mL-GDH$_{180}$. Instead, conformational transitions

among oligomeric states were accompanied by adjustments at the ACT*3 and C-

terminal domain interfaces, changes in the intersubunit contact between residues

650 (in the HM3) and 790 (in the catalytic domain), and the linkage between

segments 825-838 and 704-706 (both in the catalytic domain), the latter forming only

in Closed tetramers when both substrate and cofactor occupy the active site (Figure

5). These long-range transitions are presumably slower than catalysis and are thus

observable in our experiment. While our data do not permit conclusions about

positional and/or compositional heterogeneity within the GluBR or DNBR during

catalysis, the most reasonable scenario is that the reaction occurs when both regions

are loaded (Figure 4), with the cofactor approaching the substrate binding pocket

(Figure 2). Overall, our results indicate that the 825-838/704-706 bridge acts as a

closure-dependent modulator that biases the co-occupied Open-Closed equilibrium

and tunes the dwell time of the Closed ternary complex; the net effect on turnover is

expected to reflect the kinetic balance between chemistry and product release under

the conditions considered.

Notably, in the absence of added effectors, we found no evidence of cooperativity for

mL-GDH$_{180}$ catalyzed L-glutamate oxidation across pH 6.0–7.1, an interval that

spans the reaction's apparent pH optimum (Figure S1B) and overlaps the

physiological pH range [22,23]. At pH 6.8, the initial-rate dependences on L-glutamate

and on NAD$^+$ were Michaelis–Menten both when the co-substrate was fixed at a

saturating concentration and when it was held at several subsaturating levels (Figure

S1D). Consistent with this trend, datasets acquired at pH 6.5 (the condition used for

cryo-EM specimen preparation) were likewise well described by Michaelis–Menten

kinetics. Thus, within this near-optimal pH window, steady-state behavior is

consistent with independent active sites. The structural origin of departures from this

pattern at basic pH (this work and [10]) and under activity modulators [3,10] remains to be

determined, particularly given the regulatory potential of the N- and C-terminal

regions of the enzyme.

We found that ligand binding to the active (orthosteric) site of mL-GDH$_{180}$ is linked to

tertiary structure changes that lead to the consolidation of an oligomeric interface two

nanometers away from the catalytic center [24]. It is worth considering whether and

how natural or designed molecules binding to the 825-838 and/or 704-706 segments

could influence catalysis and whether occupancy of the active site modulates such

regulation [25]. Our findings open new avenues for high-resolution studies on catalysis

and regulation of L-GDHs of medical and industrial relevance.

MATERIALS AND METHODS

Protein production

Plasmid pLIC-His-mL-GDH$_{180}$ [3] was used to produce N-terminally His6-tagged mL-GDH180 (MSMEG_4699, Uniprot A0R1C2) in *E. coli* cells. Transformed *E. coli* BL21(DE3) cells were grown at 37 °C in LB broth supplemented with 100 µg/ml ampicillin until reaching 0.8 units of optical density at 600 nm. Protein expression was then induced by adding isopropyl β-D-1-thiogalactopyranoside (IPTG) to a final concentration of 0.5 mM and incubation was continued for 18 hours at 14 °C. The culture was centrifuged and the pellet was resuspended in 25 mM Hepes, 500 mM NaCl, 10 % V/V glycerol, 10 mM imidazole, pH 8.0. The suspension was supplemented with 0.01 µg/ml DNAse, 4 µM MgSO$_4$ and the protease inhibitor cOmplete™ EDTA-free Protease Inhibitor Cocktail (Roche). Cells were then sonicated and the mixture was clarified by centrifugation. The supernatant was loaded onto a HisTrap HP column (Cytiva) equilibrated with buffer 25 mM Hepes, 500 mM NaCl, 10% v/v glycerol, 20 mM imidazole, pH 8.0, and His6-tagged mL-GDH$_{180}$ was purified by applying a linear imidazole gradient (20-500 mM). The protein was then further purified by size-exclusion chromatography, using a Superose 6 10/300 GL column (Cytiva) equilibrated in 20 mM Mes, 300 mM NaCl, pH 6.5. mL-GDH$_{180}$ containing fractions, as confirmed by SDS-PAGE, were pooled and used immediately or stored at -80 °C for use in subsequent studies. The protein was quantified by electronic absorption using the molar absorption coefficient of 171,090 M^{-1} cm^{-1}, predicted from the amino acid sequence by the ProtParam tool (http://web.expasy.org/protparam/).

Steady state kinetics

Reactions were performed at 25 °C, using 10 nM mL-GDH$_{180}$. Oxidative deamination

of L-glutamate catalyzed by mL-GDH$_{180}$ was monitored by the increase in

absorbance at 340 nm due to NADH production (ε_{340} = 6220 M^{-1} cm^{-1}) on a

Shimadzu UV-120-02 spectrophotometer. Initial rate measurements were carried out

in the following buffers: 50 mM Mes, 50 mM NaCl, pH 6.0; 50 mM Mes, 50 mM NaCl,

pH 6.5; 50 mM Hepes, 50 mM NaCl, pH 6.8; 50 mM Hepes, 50 mM NaCl, pH 7.5; 50

mM Hepes, 50 mM NaCl, pH 8.2; 30 mM sodium phosphate buffer, pH 7.1; 30 mM

sodium phosphate buffer, pH 8.0. Within pH 6.0–7.1, the kinetic parameters k_{cat} and

K_M were obtained by fitting the dependence of the initial rate on substrate

concentration to the Michaelis-Menten equation (Figure S1). At pH 7.5, the

dependence of the initial rate on substrate concentration was sigmoidal; accordingly,

data were fitted to the Hill equation. Parameter estimation for Michaelis-Menten and

Hill models was performed by nonlinear regression. At pH 8.0 and 8.2, where activity

was low and the rate-substrate relationship remained essentially linear across the full

substrate range, catalytic efficiency (k_{cat}/K_M) was obtained by linear regression of rate

versus substrate concentration. Each experiment was performed at least twice, in

some cases using independent protein preparations. When error bars are shown on

kinetic parameters, they represent the standard errors from curve fitting. Mean k_{cat}

and K_M values reported in the text include errors obtained by propagating the

standard errors from individual fits across replicates; catalytic efficiency was

calculated as k_{cat}/K_M with error propagated from those parameter errors.

Cryo-electron microscopy

4 µl of a mixture containing 42 mM L-glutamate, 4 mM NAD$^+$ and 0.125 mg/ml mL-

GDH$_{180}$ was prepared in 20 mM Mes, 300 mM NaCl, pH 6.5, at 10 ºC, using a buffer

closely aligned with conditions previously optimized for single-particle analysis [3].

Samples were applied to Quantifoil R2/2 holey carbon grids and vitrified using a

Leica EM GP2 within the first minute after mixing, during which enzymatic activity

was confirmed by an increase in absorbance at 340 nm (Figure S1C), indicative of

NADH production. In addition, a similar protein solution was applied onto the same

type of grids bearing an additional thin layer of carbon. Vitrification on these two

types of grids was required to obtain top and side views of mL-GDH$_{180}$ tetramers

(Figure S2). Data collection was carried out in a ThermoFisher Titan Krios G4

transmission electron microscope paired with a Gatan BioContinuum Imaging Filter

and a K3 direct electron detector, at the BREM (Basque Resource for Electron

Microscopy). 23,747 and 19,038 movie frames were acquired for the sample on R2/2

grids and grids supplemented with an additional thin layer of carbon, respectively

(Figure S2). Both collections were performed at a nominal magnification of ×130,000,

resulting in a sampling of 0.6462 Å/pixel. Each movie contained 50 frames with an

accumulated dose of 49 e$^-$/Å^2. Movie frames were aligned using MotionCor2 [26] and

the final average included all dose-weighted frames.

The contrast transfer function (CTF) of the micrographs was estimated using

CTFFIND4 [27]. Particles were automatically selected from the micrographs using

autopicking from RELION-4 [28]. To speed up calculations, particles were extracted

with a 2x binning, resulting in a pixel size of 1.29 Å/pixel. In datasets with potential for

high resolution, particles were re-extracted at original sampling. An initial set of

667,866 particles was subjected to 2D and 3D classification procedures (Figure S3).

The 3D-classification with imposed D2 symmetry resulted in eight different classes,

but only three achieved good definition and high resolution. Three different

conformations were obtained, called the Open (37,641 particles), Closed1 (45,998

particles) and Closed2 (32,441 particles) conformations, with estimated resolutions of

3.88 Å, 3.54 Å and 3.57 Å, respectively. Each class was symmetry-expanded, the

star files were joined (116,080x4, *i.e.* 464,320 particles) and an average 3D map of a

subunit was refined using a mask and polished to a final resolution of 2.63 Å. A 3D

classification within a sphere mask focused on the active site separated three

different states: empty-monomer, with 131,216 particles and 3.07 Å of resolution,

cofactor-monomer, with 168,668 particles and 2.95 Å of resolution, and

cofactor/ligand-monomer, with 112,701 particles and 3.12 Å of resolution. These

three groups were crossed (common particles between selection files) with the three

sets of particles obtained in the initial classification from the conformational changes

of the tetramer, resulting in nine groups of particles that were refined independently

(Figure S3). The three maps with signal in the cofactor and substrate binding pockets

were subjected to polishing and ctfrefine to improve the resolution, each one coming

from a conformation of the tetramer: Open-cofactor/ligand-monomer (3.23 Å),

Closed1-cofactor/ligand-monomer (3.11 Å) and Closed2-cofactor/ligand-monomer

(3.18 Å). The resolution of the cryo-EM density maps was estimated using a

threshold of 0.5 in the Fourier Shell Correlation (FSC) between independently

processed halfmaps, *i.e.*, gold standard. The cryo-EM density maps were

postprocessed by local sharpening using EMReady [29], which improved the structural

details and facilitated the atomic model building.

Model fitting into cryo-EM maps was performed using the programs UCSF Chimera

[30], Coot [31] and phenix.real_space_refine [32]. From the 2.63 Å resolution cryo-EM map

of a mL-GDH$_{180}$ monomer and the atomic coordinates from the Protein Data Bank

entry 7A1D, a model of the polypeptide chain was refined which was then fitted into

the other cryo-EM maps. After removing parts of the model not defined in the maps,

performing manual model building when necessary and placing ligands manually,

using Coot [31], where appropriate, atomic coordinates and B-factors were refined

using phenix.real_space_refine [32] with NCS and secondary structure restraints.

RMSDs between models were calculated using Coot [31]. Figures were generated and

rendered with UCSF Chimera [30] or PyMOL version 1.8.x (Schrödinger, LLC).

Cryo-EM maps obtained for mL-GDH$_{180}$ were deposited in the Electron Microscopy

Data Bank under the accession codes EMD-52419 (Open), EMD-52420 (Closed1),

EMD-52421 (Closed2), EMD-52422 (empty-monomer), EMD-52423 (cofactor-

monomer), EMD-52424 (cofactor/ligand-monomer), EMD-52425 (Open-

cofactor/ligand-monomer), EMD-52426 (Closed1-cofactor/ligand-monomer), EMD-

52427 (Closed2-cofactor/ligand-monomer), EMD-52428 (total-monomer) and EMD-

52429 (Closed2-cofactor/ligand-tetramer). Atomic coordinates were deposited in the

Protein Data Bank under the accession codes 9HUX (Open), 9HUY (Closed1), 9HUZ

(Closed2), 9HV0 (empty-monomer), 9HV4 (cofactor-monomer), 9HV5

(cofactor/ligand-monomer) and 9HV6 (total-monomer).

SUPPLEMENTARY MATERIAL

Supplementary material.doc: Supplementary figures and tables.

Movie S1.mp4: Changes in the quaternary structure of mL-GDH$_{180}$ detected by cryo-EM in the presence of L-glutamate and NAD$^+$.

Movie S2.mp4: Changes in the active site of mL-GDH$_{180}$ detected by cryo-EM in the presence of L-glutamate and NAD$^+$.

Movie S3.mp4: In the presence of L-glutamate and NAD$^+$, a contact is established between the catalytic domains of neighboring mL-GDH$_{180}$ subunits connected through their N-terminal regions.

ACKNOWLEDGMENTS

This work was supported by grants PICT 2019-00989 and PICT 2021-00253 from the Agencia Nacional de Promoción de la Investigación, el Desarrollo Tecnológico y la Innovación (Agencia I+D+i, Argentina) to MNL as well as grants PID2021-124074NB-I00 and PID2021-125946OB-I00 from Agencia Estatal de Investigación of Spain to MV and GJO, respectively. We also thank the Severo Ochoa Center of Excellence Accreditation CEX2021-001136-S. MNL is a career investigator from Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET, Argentina). NC holds a CONICET PhD fellowship. NC and MNL are grateful to Liliana Rojas for excellent technical assistance. Some of this work was performed at the Basque Resource for Electron Microscopy located at Instituto Biofisika (UPV/EHU, CSIC), supported by the Department of Science, Universities and Innovation and the Innovation Fund of the Basque Government, with additional support from MCIN (Recovery, Transformation and Resilience Plan) and the Basque Government "Biotechnology Complementary

522 Plan Applied to Health" with funding from European Union NextGenerationEU

523 (PRTR-C17.I1; PRTR-C17.I01.P01.S13) (AAAA_ACG_AY_2539/22_05).

AUTHOR CONTRIBUTIONS

ML produced protein, prepared EM samples, performed EM experiments, processed

and analyzed EM data and performed structural analyses; NC produced protein,

performed kinetic analyses, analyzed EM data and performed structural analyses;

JPLA collected and analyzed EM data; DC assisted during processing of EM data;

RMR performed structural analyses; GJO performed structural analyses; MV

designed experiments, supervised EM data collection and processing and analyzed

results; MNL designed experiments, supervised biochemical experiments, refined

atomic models, analyzed data and wrote the paper. All authors read and corrected

the paper.

COMPETING INTERESTS

537 The authors declare no competing interests.

REFERENCES

1. Li M et al. Glutamate dehydrogenase: Structure, allosteric regulation, and role in insulin homeostasis. Neurochemical Research. 2014;39(3):433–445. https://doi.org/10.1007/s11064-013-1173-2

2. Miñambres B, Olivera ER, Jensen RA, Luengo JM. A new class of glutamate dehydrogenases (GDH): Biochemical and genetic characterization of the first member, the AMP-requiring NAD-specific GDH of Streptomyces clavuligerus. Journal of Biological Chemistry. 2000;275(50):39529–39542. https://doi.org/10.1074/jbc.M005136200

3. Lázaro M et al. 3D architecture and structural flexibility revealed in the subfamily of large glutamate dehydrogenases by a mycobacterial enzyme. Communications Biology. 2021;4(1):684. http://dx.doi.org/10.1038/s42003-021-02222-x. https://doi.org/10.1038/s42003-021-02222-x

4. Lang EJM et al. Allosteric ACTion: The varied ACT domains regulating enzymes of amino-acid metabolism. Current Opinion in Structural Biology. 2014;29:102–111. https://doi.org/10.1016/j.sbi.2014.10.007

5. Möglich A, Ayers RA, Moffat K. Structure and Signaling Mechanism of Per-ARNT-Sim Domains. Structure. 2009;17(10):1282–1294. https://doi.org/10.1016/j.str.2009.08.011

6. Gallant JL, Viljoen AJ, Van Helden PD, Wiid IJF. Glutamate dehydrogenase is required by mycobacterium bovis BCG for resistance to cellular stress. PLoS ONE. 2016;11(1):1–14. https://doi.org/10.1371/journal.pone.0147706

7. Krissinel E, Henrick K. Protein interfaces, surfaces and assemblies' service PISA at the european bioinformatics institute. Journal of Molecular Biology. 2007;372(3):774–797. http://www.ebi.ac.uk/pdbe/prot_int/pistart.html

8. Baker PJ et al. Subunit assembly and active site location in the structure of

glutamate dehydrogenase. Proteins: Structure, Function, and Bioinformatics.

1992;12(1):75–86. https://doi.org/10.1002/prot.340120109

9. Britton KL, Baker PJ, Rice DW, Stilllman TJ. Structural relationship between the

hexameric and tetrameric family of glutamate dehydrogenases. European Journal of

Biochemistry. 1992;209(3):851–859. https://doi.org/10.1111/j.1432-

1033.1992.tb17357.x

10. O'Hare HM et al. Regulation of glutamate metabolism by protein kinases in

mycobacteria. Molecular Microbiology. 2008;70(6):1408–1423.

https://doi.org/10.1111/j.1365-2958.2008.06489.x

11. Nott TJ et al. An intramolecular switch regulates phosphoindependent FHA

domain interactions in Mycobacterium tuberculosis. Science Signaling.

2009;2(63):ra12. https://doi.org/10.1126/scisignal.2000212

12. Veronese FM et al. Nicotinamide adenine dinucleotide specific glutamate

dehydrogenase of Neurospora. I. Purification and molecular properties. Journal of

Biological Chemistry. 1974;249(24):7922–7928. https://doi.org/10.1016/S0021-

9258(19)42053-X

13. Lu CD, Abdelal AT. The gdhB gene of Pseudomonas aeruginosa encodes an

arginine-inducible NAD+-dependent glutamate dehydrogenase which is subject to

allosteric regulation. Journal of Bacteriology. 2001;183(2):490–499.

https://doi.org/10.1128/JB.183.2.490-499.2001

14. Camardella L et al. The Antarctic Psychrobacter sp. TAD1 has two cold-active

glutamate dehydrogenases with different cofactor specificities. Characterisation of

the NAD+-dependent enzyme. Comparative Biochemistry and Physiology - A

Molecular and Integrative Physiology. 2002;131(3):559–567.

https://doi.org/10.1016/S1095-6433(01)00507-4

15. Kawakami R, Sakuraba H, Ohshima T. Gene cloning and characterization of the

very large NAD-dependent L-glutamate dehydrogenase from the psychrophile

Janthinobacterium lividum, isolated from cold soil. Journal of Bacteriology.

2007;189(15):5626–5633. https://doi.org/10.1128/JB.00496-07

16. Kawakami R, Oyama M, Sakubara H, Ohshima T. The Unique Kinetic Behavior

of the Very Large NAD-Dependent Glutamate Dehydrogenase from

Janthinobacterium lividum. Bioscience, Biotechnology, and Biochemistry.

2010;74(4):884–887. http://www.tandfonline.com/doi/full/10.1271/bbb.90925.

https://doi.org/10.1271/bbb.90925

17. Beaufay F et al. A NAD-dependent glutamate dehydrogenase coordinates

metabolism with cell division in Caulobacter crescentus. The EMBO Journal.

2015;34(13):1786–1800.

http://emboj.embopress.org/cgi/doi/10.15252/embj.201490730.

https://doi.org/10.15252/embj.201490730

18. Baker PJ et al. Structural consequences of sequence patterns in the fingerprint

region of the nucleotide binding fold. Implications for nucleotide specificity. Journal of

Molecular Biology. 1992;228(2):662–671. https://doi.org/10.1016/0022-

2836(92)90848-E

19. Engel PC. Glutamate dehydrogenases: The why and how of coenzyme

specificity. Neurochemical Research. 2014;39(3):426–432.

https://doi.org/10.1007/s11064-013-1089-x

20. Stillman TJ, Baker PJ, Britton KL, Rice DW. Conformational flexibility in

glutamate dehydrogenase: Role of water in substrate recognition and catalysis.

Journal of Molecular Biology. 1993;234(4):1131–1139.

https://doi.org/10.1006/jmbi.1993.1665

21. Borgnia MJ et al. Using cryo-EM to map small ligands on dynamic metabolic

enzymes: Studies with glutamate dehydrogenase. Molecular Pharmacology.

2016;89(6):645–651. https://doi.org/10.1124/mol.116.103382

22. Rao M, Streur TL, Aldwell FE, Cook GM. Intracellular pH regulation by

Mycobacterium smegmatis and Mycobacterium bovis BCG. Microbiology.

2001;147(4):1017–1024. https://doi.org/10.1099/00221287-147-4-1017

23. Vandal OH et al. A membrane protein preserves intrabacterial pH in

intraphagosomal Mycobacterium tuberculosis. Nature Medicine. 2008;14(8):849–854.

https://doi.org/10.1038/nm.1795

24. Perica T, Chothia C, Teichmann SA. Evolution of oligomeric state through

geometric coupling of protein interfaces. Proceedings of the National Academy of

Sciences. 2012;109(21):8127–8132.

http://www.pnas.org/cgi/doi/10.1073/pnas.1120028109.

https://doi.org/10.1073/pnas.1120028109

25. Schulze JO et al. Bidirectional Allosteric Communication between the ATP-

Binding Site and the Regulatory PIF Pocket in PDK1 Protein Kinase. Cell Chemical

Biology. 2016;23(10):1193–1205. http://europepmc.org/abstract/med/27693059.

https://doi.org/10.1016/j.chembiol.2016.06.017

26. Zheng SQ et al. MotionCor2: anisotropic correction of beam-induced motion for

improved cryo-electron microscopy. Nature Methods. 2017;14(4):331–332.

https://doi.org/10.1038/nmeth.4193. https://doi.org/10.1038/nmeth.4193

27. Rohou A, Grigorieff N. CTFFIND4: Fast and accurate defocus estimation from

electron micrographs. Journal of Structural Biology. 2015;192(2):216–221.

https://doi.org/10.1016/j.jsb.2015.08.008

28. Zivanov J et al. New tools for automated high-resolution cryo-EM structure

determination in RELION-3 Egelman EH, Kuriyan J, editors. eLife. 2018;7:e42166.

https://doi.org/10.7554/eLife.42166. https://doi.org/10.7554/eLife.42166

29. He J, Li T, Huang SY. Improvement of cryo-EM maps by simultaneous local and

non-local deep learning. Nature Communications. 2023;14(1):1–16.

https://doi.org/10.1038/s41467-023-39031-1

30. Pettersen EF et al. UCSF Chimera - A visualization system for exploratory

research and analysis. Journal of Computational Chemistry. 2004;25(13):1605–1612.

https://doi.org/10.1002/jcc.20084

31. Emsley P, Lohkamp B, Scott WG, Cowtan K. Features and development of Coot.

Acta Crystallographica Section D: Biological Crystallography. 2010;66(4):486–501.

https://doi.org/10.1107/S0907444910007493

32. Afonine P V. et al. Real-space refinement in PHENIX for cryo-EM and

crystallography. Acta Crystallographica Section D: Structural Biology.

2018;74(6):531–544. https://doi.org/10.1107/S2059798318006551.

https://doi.org/10.1107/S2059798318006551

33. Abramson J et al. Accurate structure prediction of biomolecular interactions with

AlphaFold 3. Nature. 2024;630(8016):493–500. http://dx.doi.org/10.1038/s41586-

024-07487-w. https://doi.org/10.1038/s41586-024-07487-w

TABLES

Table I. Cryo-EM data collection and processing. LS, local sharpening.

	Open	Closed1	Closed2
	(EMD-52419)	(EMD-52420)	(EMD-52421)
Data collection and			
Magnification	130,000	130,000	130,000
Voltage (kV)	300	300	300
Electron dose ($e^-/Å^2$)	0.979	0.979	0.979
Defocus range (μm)	1.0-1.6	1.0-1.6	1.0-1.6
Pixel size (Å)	1.29	1.29	1.29
Symmetry imposed	D2	D2	D2
Initial particles images (no.)	667,866	667,866	667,866
Final particles images (no.)	37,641	45,998	32,441
Map resolution (Å)	3.88	3.54	3.57
FSC threshold	0.143	0.143	0.143
Map sharpening B factor ($Å^2$)	-50	-50	-50
Refinement			
Model resolution (Å)	3.88	3.54	3.57
Model composition			
Non-hydrogen atoms	33576	33576	33576
Protein residues	4300	4300	4300
Ligands	4 NAD^+	4 NAD^+	4 NAD^+
B factors ($Å^2$)			
Protein	224.33	189.71	159.33
Ligands	273.06	208.84	186.14
R.m.s. deviations			
Bond lengths (Å)	0.001	0.002	0.002
Bond angles (°)	0.371	0.409	0.395
Validation			
MolProbity score	1.69	1.62	1.50
Clashscore	10.54	11.11	9.42
Poor rotamers (%)	0.11	0	0.11
Ramachandran plot			
Favored (%)	97.20	97.76	98.55
Allowed (%)	2.80	2.24	1.45
Disallowed (%)	0	0	0
CC (mask)	0.83	0.77	0.83
CC (box)	0.91	0.87	0.89
CC (peaks)	0.77	0.70	0.77
CC (volume)	0.82	0.76	0.82
Mean CC for ligands	0.85	0.82	0.80

	empty- monomer (EMD-52422)	cofactor- monomer (EMD-52423)	cofactor/ligand- monomer (EMD-52424)
Data collection and			
Magnification	130,000	130,000	130,000
Voltage (kV)	300	300	300
Electron dose ($e^-/Å^2$)	0.979	0.979	0.979
Defocus range (μm)	1.0-1.6	1.0-1.6	1.0-1.6
Pixel size (Å)	1.29	1.29	1.29
Symmetry imposed	C1	C1	C1
Initial particles images (no.)	464,320	464,320	464,320
Final particles images (no.)	131,216	168,668	112,701
Map resolution (Å)	3.07	2.95	3.12
FSC threshold	0.143	0.143	0.143
Map sharpening B factor	LS	LS	LS
Refinement			
Model resolution (Å)	3.07	2.95	3.12
Model composition			
Non-hydrogen atoms	8717	8761	8760
Protein residues	1124	1124	1123
Ligands	-	1 NAD^+	1 NAD^+/ 1 L-Glu
B factors ($Å^2$)			
Protein	64.32	62.58	72.09
Ligands	-	56.65	81.37
R.m.s. deviations			
Bond lengths (Å)	0.004	0.004	0.004
Bond angles (°)	0.691	0.721	0.693
Validation			
MolProbity score	1.37	1.27	1.37
Clashscore	5.53	5.16	5.80
Poor rotamers (%)	0.55	0.55	0.11
Ramachandran plot			
Favored (%)	97.68	98.48	97.77
Allowed (%)	2.32	1.52	2.23
Disallowed (%)	0	0	0
CC (mask)	0.87	0.88	0.87
CC (box)	0.89	0.89	0.89
CC (peaks)	0.81	0.81	0.79
CC (volume)	0.88	0.88	0.87
Mean CC for ligands	-	0.91	0.84

	Open-cofactor/ligand-monomer (EMD-52425)	Closed1-cofactor/ligand-monomer (EMD-52426)	Closed2-cofactor/ligand-monomer (EMD-52427)
Data collection and			
Magnification	130,000	130,000	130,000
Voltage (kV)	300	300	300
Electron dose ($e^-/Å^2$)	0.979	0.979	0.979
Defocus range (μm)	1.0-1.6	1.0-1.6	1.0-1.6
Pixel size (Å)	0.646	0.646	0.646
Symmetry imposed	C1	C1	C1
Initial particles images (no.)	112,701	112,701	112,701
Final particles images (no.)	33,330	46,380	32,991
Map resolution (Å)	3.23	3.11	3.18
FSC threshold	0.143	0.143	0.143
Map sharpening B factor	LS	LS	LS
Refinement			
Model resolution (Å)			
Model composition			
Non-hydrogen atoms			
Protein residues			
Ligands			
B factors ($Å^2$)			
Protein			
Ligands			
R.m.s. deviations			
Bond lengths (Å)			
Bond angles (°)			
Validation			
MolProbity score			
Clashscore			
Poor rotamers (%)			
Ramachandran plot			
Favored (%)			
Allowed (%)			
Disallowed (%)			
CC (mask)			
CC (box)			
CC (peaks)			
CC (volume)			
Mean CC for ligands			

	total-monomer (EMD-52428)	Closed2-cofactor/ligand-tetramer (EMD-52429)
Data collection and		
Magnification	130,000	130,000
Voltage (kV)	300	300
Electron dose ($e^-/Å^2$)	0.979	0.979
Defocus range (μm)	1.0-1.6	1.0-1.6
Pixel size (Å)	0.646	1.29
Symmetry imposed	C1	C1
Initial particles images (no.)	464,320	112,701
Final particles images (no.)	464,320	32,991
Map resolution (Å)	2.63	4.10
FSC threshold	0.143	0.143
Map sharpening B factor	LS	-50
Refinement		
Model resolution (Å)	2.63	
Model composition		
Non-hydrogen atoms	8683	
Protein residues	1113	
Ligands	1 NAD$^+$	
B factors ($Å^2$)		
Protein	56.19	
Ligands	48.51	
R.m.s. deviations		
Bond lengths (Å)	0.002	
Bond angles (°)	0.467	
Validation		
MolProbity score	1.39	
Clashscore	7.12	
Poor rotamers (%)	0.55	
Ramachandran plot		
Favored (%)	99.10	
Allowed (%)	0.90	
Disallowed (%)	0	
CC (mask)	0.80	
CC (box)	0.68	
CC (peaks)	0.65	
CC (volume)	0.78	
Mean CC for ligands	0.81	

FIGURES

Figure 1

Three conformers of mL-GDH$_{180}$ detected upon incubation with L-glutamate

and NAD$^+$ reveal modulations of the enzyme quaternary structure. (A) Cryo-EM

mL-GDH$_{180}$ maps obtained after incubating the enzyme with L-glutamate and NAD$^+$.

The dimensions of each conformer are provided in Å. Maps cover, for each

monomer, the region indicated by a dotted rectangle in the diagram shown below

(residues 500-1588). ACT*, Aspartate kinase-Chorismate mutase-TyrA-like; PAS,

Per-Arnt-Sim; HM, helical motif; CD, catalytic domain; CTD, C-terminal domain. (B)

Approach of the catalytic domains of mL-GDH$_{180}$ subunits linked by the N-terminal

region. Segment 825-838, poorly defined in maps of the Open, Closed1 and Closed2

enzyme forms, approaches the catalytic domain in a neighboring subunit. In each

case, a zoomed in view of the dinucleotide binding region (DNBR) is shown below.

The protein is displayed in ribbon representation and cryo-EM maps are depicted in

grey. The nicotinamide portion of the NAD^{+} molecule (in sticks) exhibits high mobility

in all conformations. Subdomains I and II (SDI and SDII) of the catalytic domain are

colored in yellow and orange, respectively, whereas ACT*3 and HM3 motifs are

shown in cyan and green, respectively, for one monomer (the one on the left), with

the neighboring subunit (the one on the right) shown in grey. All maps are shown with

a contour level of 0.02. GluBR, glutamate binding region.

Figure 2

Three conformers of mL-GDH$_{180}$ subunits reveal modulations in tertiary

structure associated with differential active site occupancies. Alternative

conformations of mL-GDH$_{180}$ subunits are shown, with the active site either empty or

containing NAD$^+$ or NAD$^+$ and L-glutamate. The average RMSD for 1122 aligned

residues in pairwise comparisons of the atomic models (show in ribbon

representation) is 0.7 Å. NAD^+ and L-glutamate are depicted as sticks as well as

highlighted residues. For each species, subdomains I and II of the catalytic domain

(SDI, residues 702-926 and 1188-1220; SDII, residues 927-1187) are colored in

yellow and orange, respectively. A small portion of the C-terminal domain is

displayed in magenta. Interactions established by the segment 825-838 with other

portions of the polypeptide chain are shown in the panels on the left. Note that within

the region 825–838, residues 825-827 and 830-838 could be built for the empty- and

cofactor- monomers, while residues 825 and 829-838 were modeled for the

cofactor/ligand-monomer. A comparison of the atomic models of the empty- (green),

cofactor- (blue) and cofactor/ligand-monomer (gray) is presented in the lower left

corner. Cryo-EM maps are depicted in grey, with contour levels of 5. H-bonds and

salt bridges are displayed as dashed lines. GluBR, glutamate binding region; DNBR,

dinucleotide binding region.

Figure 3

Comparison of the catalytic domain of mL-GDH$_{180}$ with that of S-GDHs. A

monomer of mL-GDH$_{180}$ is shown on the left. Segments 824-838 and 858-882 (in

dark green), both located in the catalytic domain subdomain I (SDI), are absent in S-

GDHs [2]. For the 824-838 portion coordinates were taken from a model calculated

with AlphaFold 3 [33]. A monomer of *Cs*GDH (*Cs*, *Clostridium symbiosum*; PDB 1BGV

[20]), considered a representative S-GDH, is shown on the right. The SDI of S-GDHs

presents secondary structure elements (in dark blue, named according to the

consensus nomenclature for S-GDHs) with conserved roles in oligomerization in the

enzyme subfamily, which are not present in mL-GDH$_{180}$ or related L-GDHs. For both

mL-GDH$_{180}$ and *Cs*GDH, the six-stranded 432156 β-sheet that shapes the DNBR in

the catalytic domain subdomain II (SDII) is shown in grey; the linker regions β4-β5 (in

light green) and β5-β6 (in cyan) encompass a larger number of residues in mL-

GDH$_{180}$. For the two proteins, the SDI and SDII are shown, except for otherwise

highlighted portions, in yellow and orange, respectively.

Figure 4

GluBR occupancy is linked to the coupling of the catalytic domain of two

adjacent monomers in the Closed conformations. Top panel: the atomic model of

the cofactor/ligand-monomer of mL-GDH$_{180}$ is shown together with maps of this

species obtained from Open, Closed1 or Closed2 tetramers (Open-, Closed1- and

Closed2- cofactor/ligand-monomers; contour levels are 5). NAD$^+$ and L-glutamate as

well as highlighted residues are depicted as sticks. The subdomains I and II of the

catalytic domain (SDI and SDII) are colored in yellow and orange, respectively.

Bottom panel: in the context of tetramers containing a cofactor/ligand-monomer

(cofactor/ligand-tetramers; contour levels are 0.015), in a progression towards the

Closed1 and Closed2 conformers, the stabilization of an interaction between

segment 825-838 of one subunit (colored, on the left) and segment 704-706 of the

neighboring one (in grey, on the right) is observed. The atomic coordinates of the

subunit on the left correspond in each case to those of a cofactor/ligand-monomer,

replacing the equivalent ones presented in Figure 1. H-bonds and salt bridges are

displayed as dashed lines. Cryo-EM maps are displayed in grey.

mL-GDH₁₈₀ + L-glu + NAD⁺

Figure 5

Ligand binding triggers tertiary structure remodeling that links catalysis with

intersubunit communication in mL-GDH₁₈₀. Cryo-EM maps of mL-GDH₁₈₀

incubated with L-glutamate and NAD⁺ illustrate the conformational ensemble

comprising one open and two closed tetrameric states. In the open form, active sites

can be empty or occupied by the cofactor or by both cofactor and ligand. In the

closed conformations, when both the cofactor and the substrate simultaneously

occupy the active site, the catalytic domains of neighboring subunits connected

through their N-terminal regions approach each other, forming an intersubunit contact

approximately 2 nm away from the catalytic center bearing the ligand and cofactor.

This structural transition provides a physical link between active-site occupancy and

long-range communication within the tetramer, coupling catalysis to quaternary-

structure remodeling.