Tertiary and quaternary structure remodeling by occupancy of the substrate binding pocket in a large glutamate dehydrogenase

Melisa Lázaro [a,#], Nicolás Chamorro [b,#], Jorge P. López-Alonso [c,d], Diego Charro [a], Rodolfo M. Rasia [b,e], Gonzalo Jiménez-Osés [a,f], Mikel Valle [a,*], María-Natalia Lisa [b,e,*]

[a] Center for Cooperative Research in Biosciences (CIC bioGUNE), Basque Research and Technology Alliance (BRTA), Bizkaia Technology Park, Building 801A, 48160 Derio, Spain.

[b] Instituto de Biología Molecular y Celular de Rosario (IBR CONICET-UNR), Ocampo y Esmeralda, Rosario S2002LRK, Argentina.

[c] Instituto Biofisika (UPV/EHU, CSIC), 48940 Leioa, Spain.

[d] Basque Resource for Electron Microscopy, 48940 Leioa, Spain.

[e] Plataforma de Biología Estructural y Metabolómica (PLABEM), Ocampo y Esmeralda, Rosario S2002LRK, Argentina.

[f] Ikerbasque, Basque Foundation for Science, 48013 Bilbao, Spain.

# Equal contribution

* Correspondence: MNL: lisa@ibr-conicet.gov.ar; MV: mvalle@cicbiogune.es

ABSTRACT

Glutamate dehydrogenases (GDHs) catalyze the oxidative deamination of L-glutamate to 2-oxoglutarate using $NAD(P)^+$ as a cofactor. The large type of GDHs (L-GDHs) displays a dynamic homotetrameric architecture that alternates between open and closed states. However, the catalytic mechanism and the functional relevance of the large conformational changes in L-GDHs remain poorly understood. Here, we use cryoEM to investigate the structure and the conformational landscape of the mycobacterial L-GDH composed of 180 kDa subunits ($mL\text{-}GDH_{180}$) when incubated with L-glutamate and NAD^+. Classification of the heterogeneous population of tetramers reveals opening-closing motions and sorting of individual subunits resolves the occupancy of the cofactor and substrate binding pockets. CryoEM maps show that ligand binding to the glutamate binding pocket is accompanied by structural changes in a region approximately two nanometers away from the active site, leading to the formation of a previously undetected interaction between the catalytic domains of neighboring subunits in $mL\text{-}GDH_{180}$ closed tetrameric states. Our findings indicate that the occupancy of the substrate binding site of $mL\text{-}GDH_{180}$ is linked to a remodeling of both the tertiary and quaternary structure of the enzyme.

INTRODUCTION

Amino acid dehydrogenases have long attracted interest due to their physiological functions and potential industrial applications. Among them, glutamate dehydrogenases (GDHs; EC 1.4.1.2) catalyze the reaction L-glutamate + $NAD(P)^+$ + H_2O <=> 2-oxoglutarate + NAD(P)H + H^+ + NH_3, at the crossroads of carbon and nitrogen metabolism. These enzymes are oligomeric and are classified based on the molecular weight of their subunits. Small GDHs (S-GDHs), present in prokaryotes and eukaryotes, consist of 50 kDa subunits [1]. The conserved catalytic domain spans nearly the full length of the polypeptide chain of S-GDHs and mediates oligomerization. Biochemical and structural studies on S-GDHs have enabled detailed characterization of their catalytic and regulatory mechanisms. In contrast, much less is known about large GDHs (L-GDHs), which are found in prokaryotes and lower eukaryotes [2]. This group includes enzymes with 115 kDa subunits (L-GDHs$_{115}$) and those with 180 kDa subunits (L-GDHs$_{180}$), such as the L-GDH$_{180}$ from mycobacteria (mL-GDH$_{180}$). L-GDHs feature distinctive and extended N- and C-terminal regions flanking the catalytic core, which may confer regulatory or structural roles [3].

In a previous work, we demonstrated that the L-GDH$_{180}$ from *Mycobacterium smegmatis* (mL-GDH$_{180}$; 76% sequence identity with the L-GDH$_{180}$ from *Mycobacterium tuberculosis*) assembles into a homotetramer, with a quaternary structure markedly different from that of S-GDHs [3]. Such architecture is stabilized by interactions involving the N-terminal region, which contains ACT- and PAS-type domains typically associated with regulatory functions [4,5], as well as a C-terminal domain of unknown function.

Here we report cryoEM structures of mL-GDH$_{180}$ at resolutions up to 3 Å, obtained in the presence of NAD^+ and L-glutamate. We reveal that ligand binding to the glutamate binding pocket is accompanied by changes in the tertiary structure in a region approximately 2 nm distant from the active site, promoting interactions between the catalytic domains of

neighboring monomers and modulating the quaternary structure.

Despite extensive efforts to understand the molecular basis of GDHs activity and

regulation, structural data on ternary complexes containing enzyme, cofactor, and

substrate are scarce, particularly for NAD^+-dependent L-GDHs. Here, we present cryoEM

evidence for enzyme species present upon incubation with L-glutamate and NAD^+ of a key

L-GDH_{180} for the virulence of a long-attacked pathogen [6].

METHODS

Protein production

Plasmid pLIC-His-mL-GDH$_{180}$ [3] was used to produce N-terminally His6-tagged mL-GDH180 (MSMEG_4699, Uniprot A0R1C2) in *E. coli* cells. Transformed *E. coli* BL21(DE3) cells were grown at 37 °C in LB broth supplemented with 100 µg/ml ampicillin until reaching 0.8 units of optical density at 600 nm. Protein expression was then induced by adding isopropyl β-D-1-thiogalactopyranoside (IPTG) to a final concentration of 0.5 mM and incubation was continued for 18 hours at 14 °C. The culture was centrifuged and the pellet was resuspended in 25 mM Hepes, 500 mM NaCl, 10 % V/V glycerol, 10 mM imidazole, pH 8.0. The suspension was supplemented with 0.01 µg/mL DNAse, 4 µM MgSO$_4$ and the protease inhibitor cOmplete™ EDTA-free Protease Inhibitor Cocktail (Roche). Cells were then sonicated and the mixture was clarified by centrifugation. The supernatant was loaded onto a HisTrap HP column (Cytiva) equilibrated with buffer 25 mM Hepes, 500 mM NaCl, 10% v/v glycerol, 20 mM imidazole, pH 8.0, and His6-tagged mL-GDH$_{180}$ was purified by applying a linear imidazole gradient (20-500 mM). The protein was then further purified by size-exclusion chromatography, using a Superose 6 10/300 GL column (Cytiva) equilibrated in 20 mM Mes, 300 mM NaCl, pH 6.5. mL-GDH$_{180}$ containing fractions, as confirmed by SDS-PAGE, were pooled and used immediately or stored at -80 °C for use in subsequent studies. The protein was quantified by electronic absorption using the molar absorption coefficient of 171,090 M^{-1} cm^{-1}, predicted from the amino acid sequence by the ProtParam tool (http://web.expasy.org/protparam/).

Steady state kinetics

Reactions were performed at 25 °C, using 11.2 nM mL-GDH$_{180}$. Oxidative deamination of L-glutamate catalyzed by mL-GDH$_{180}$ was monitored by following the increase in absorbance at 340 nm resulting from NADH production. To calculate reaction rates, the

molar extinction coefficient 6220 M^{-1} cm^{-1} for NADH at 340 nm was used. The kinetic

parameters k_{cat} and K_M were derived by fitting the Michaelis-Menten equation to initial

rates measured in 100 mM NaH_2PO_4/Na_2HPO_4, pH 7.0, recorded on a Shimadzu UV-120-

02 spectrophotometer (Figure S1). k_{cat} and K_M values and their errors correspond to the

mean and maximum dispersion observed, respectively, between at least two independent

measurements performed with different protein preparations. The catalytic efficiency

(k_{cat}/K_M) was calculated as the quotient between k_{cat} and K_M, and its error was obtained by

propagating the error of such parameters.

Cryo-electron microscopy

4 µl of a mixture containing 42 mM L-glutamate, 4 mM NAD^+ and 0.125 mg/ml mL-GDH_{180}

was prepared in 20 mM Mes, 300 mM NaCl, pH 6.5, at 10 ºC. Samples were applied to

Quantifoil R2/2 holey carbon grids and vitrified using a Leica EM GP2 within the first

minute after mixing, during which enzymatic activity was confirmed by an increase in

absorbance at 340 nm, indicative of NADH production. In addition, a similar protein

solution was applied onto the same type of grids bearing an additional thin layer of carbon.

Vitrification on these two types of grids was required to obtain top and side views of mL-

GDH_{180} tetramers (Figure S2). Data collection was carried out in a ThermoFisher Titan

Krios G4 transmission electron microscope paired with a Gatan BioContinuum Imaging

Filter and a K3 direct electron detector device, at the BREM (Basque Resource for

Electron Microscopy). 23,747 and 19,038 movie frames were acquired for the sample on

Q2/2 grids and grids supplemented with an additional thin layer of carbon, respectively

(Figure S2). Both collections were performed at a nominal magnification of ×130,000,

resulting in a sampling of 0.6462 Å/pixel. Each movie contained 50 frames with an

accumulated dose of 49 e^-/$Å^2$. Movie frames were aligned using MotionCor2 [7] and the

final average included all dose-weighted frames.

The contrast transfer function (CTF) of the micrographs was estimated using CTFFIND4 [8]. Particles were automatically selected from the micrographs using autopicking from RELION-4 [9]. To speed up calculations, particles were extracted with a 2x binning, resulting in a pixel size of 1.29 Å/pixel. In datasets with potential for high resolution, particles were re-extracted at original sampling. An initial set of 667,866 particles was subjected to 2D and 3D classification procedures (Figure S3). The 3D-classification with imposed D2 symmetry resulted in eight different classes, but only three achieved good definition and high resolution. Three different conformations were obtained, called the Open (37,641 particles), Closed1 (45,998 particles) and Closed2 (32,441 particles) conformations, with estimated resolutions of 3.88 Å, 3.54 Å and 3.57 Å, respectively. Each class was symmetry-expanded, the star files were joined (116,080x4, *i.e.* 464,320 particles) and an average 3D map of a subunit was refined using a mask and polished to a final resolution of 2.63 Å. A 3D classification within a sphere mask focused on the active site separated three different states: empty-monomer, with 131,216 particles and 3.07 Å of resolution, cofactor-monomer, with 168,668 particles and 2.95 Å of resolution, and cofactor/ligand-monomer, with 112,701 particles and 3.12 Å of resolution. These three groups were crossed (common particles between selection files) with the three sets of particles obtained in the initial classification from the conformational changes of the tetramer, resulting in nine groups of particles that were refined independently (Figure S3). The three maps with signal in the cofactor and substrate binding pockets were subjected to polishing and ctfrefine to improve the resolution, each one coming from a conformation of the tetramer: Open-cofactor/ligand-monomer (3.23 Å), Closed1-cofactor/ligand-monomer (3.11 Å) and Closed2-cofactor/ligand-monomer (3.18 Å). The resolution of the cryoEM density maps was estimated using a threshold of 0.5 in the Fourier Shell Correlation (FSC) between independently processed halfmaps, *i.e.*, gold standard. The cryoEM density maps were postprocessed by local sharpening using

EMReady [10], which improved the structural details and facilitated the atomic model

building.

Model fitting into cryo-EM maps was performed using the programs UCSF Chimera [11],

Coot [12] and phenix.real_space_refine [13]. From the 2.63 Å resolution cryoEM map of a

mL-GDH$_{180}$ monomer and the atomic coordinates from the Protein Data Bank entry 7A1D,

a model of the polypeptide chain was refined which was then fitted into the other cryo-EM

maps. After removing parts of the model not defined in the maps, performing manual

model building when necessary and placing ligands manually, using Coot [12], where

appropriate, atomic coordinates and B-factors were refined using

phenix.real_space_refine [13] with NCS and secondary structure restraints.

Figures were generated and rendered with UCSF Chimera [11] or PyMOL version 1.8.x

(Schrödinger, LLC).

Cryo-EM maps obtained for mL-GDH$_{180}$ were deposited in the Electron Microscopy Data

Bank under the accession codes EMD-52419 (Open), EMD-52420 (Closed1), EMD-52421

(Closed2), EMD-52422 (empty-monomer), EMD-52423 (cofactor-monomer), EMD-52424

(cofactor/ligand-monomer), EMD-52425 (Open-cofactor/ligand-monomer), EMD-52426

(Closed1-cofactor/ligand-monomer), EMD-52427 (Closed2-cofactor/ligand-monomer),

EMD-52428 (total-monomer) and EMD-52429 (Closed2-cofactor/ligand-tetramer). Atomic

coordinates were deposited in the Protein Data Bank under the accession codes 9HUX

(Open), 9HUY (Closed1), 9HUZ (Closed2), 9HV0 (empty-monomer), 9HV4 (cofactor-

monomer), 9HV5 (cofactor/ligand-monomer) and 9HV6 (total-monomer).

RESULTS

Three conformers of mL-GDH$_{180}$ detected upon incubation with L-glutamate and NAD$^+$ reveal modulations of the enzyme quaternary structure

To investigate the molecular basis of mL-GDH$_{180}$ activity, we first determined its kinetic parameters for the oxidative deamination of L-glutamate. Under steady state conditions, in 100 mM NaH$_2$PO$_4$/Na$_2$HPO$_4$ at pH 7.0, the estimated k_{cat}, K_M and k_{cat}/K_M values were (20 ± 5) s^{-1}, (34 ± 7) mM and (0.6 ± 0.3) s^{-1} mM^{-1} for L-glutamate, and (21 ± 5) s^{-1}, (0.26 ± 0.07) mM and (80 ± 40) s^{-1} mM^{-1} for NAD$^+$, respectively (Figure S1). Single-particle cryoEM experiments were then conducted on mL-GDH$_{180}$ incubated with of NAD$^+$ and L-glutamate during NADH production (see Methods). As previously reported [3], 2D class averages revealed high flexibility at the distal ends of the tetramers, where the ACT*1-2 and PAS domains are located (Figure S2). This flexibility resulted in vanishing density for such regions in the cryoEM maps. Subsequently, 3D-classification of mL-GDH$_{180}$ particles revealed three conformational states of the enzyme, named Open, Closed1 and Closed2 (Figure 1, Table 1, Figure S3). In these conformers, the ACT*3 module, the helical motif 3 (HM3), the catalytic domain and the C-terminal region were defined in each monomer, achieving estimated global resolutions of 3.88 Å, 3.54 Å and 3.57 Å, respectively. The conformers differ primarily in the relative positions of the subunits (Figure 1A, Figure S4A and Movie S1). In the Closed forms, the catalytic domains of adjacent subunits connected through their N-terminal segments are found in closer proximity compared to the Open state (Figure 1B).

Comparison of the atomic models derived from the 3D cryoEM maps provided initial insights on the molecular determinants underlying the modulation of mL-GDH$_{180}$ architecture. Pairwise structural alignments of monomers from the three conformers, based on all modeled α-carbons, yielded an average RMSD of 0.4 Å (calculated over 1075 aligned residues), indicating that the observed conformational variability primarily reflects

quaternary rearrangements. Consistent with this, the intersubunit contact area at the AB-

CD interface (Figure S4B) was measured [14] as 550 $Å^2$, 750 $Å^2$ and 800 $Å^2$, and at the

AD-BC interface as 1200 $Å^2$, 1000 $Å^2$ and 1200 $Å^2$, for the Open, Closed1 and Closed2

forms, respectively. At the AB-CD interface, hydrogen bonds between residues in the

ACT*3 domains of neighboring monomers mediate the intersubunit contacts in all three

conformers (Figure S4B and Table S1). Remarkably, a previously undetected interfacial

interaction was found in the Closed2 tetramer, involving a hydrogen bond between the

side chains of Ser650 (in the HM3) and Glu790 (in the catalytic domain) (Figure 1B and

Figure S4B). At the AD-BC interface, distinct salt bridge combinations are found in the

different conformers, with Asp1516:Arg1582 present in the Open form, Arg1504:Glu1562

in the Closed1 and both simultaneously observed in the Closed2 (Figure S4B and Table

S1). Additionally, in all three conformations, extensive hydrogen bonds networks link a

conserved set of secondary structure elements in the C-terminal domain, with alternative

arrangements in each state. Most of the interfacial residues observed in mL-GDH$_{180}$

tetramers found in the presence of NAD$^+$ and L-glutamate had been previously identified

as contributors to oligomer stabilization [3] and are conserved across diverse L-GDHs

(Figure S5), suggesting a shared structural basis for quaternary modulation within this

enzyme family.

As HM3 in one monomer and the region encompassing Glu790 in the catalytic domain of

an adjacent subunit move closer, other portions of the catalytic domains in mL-GDH$_{180}$

monomers linked via their N-terminal segments also shift toward each other (Figure 1B).

Intriguingly, the density corresponding to residues 825-838 was poorly defined in the maps

of the Open, Closed1 and Closed2 conformers distinguished in the presence of NAD$^+$ and

L-glutamate. At the same time, the nicotinamide moiety of the cofactor molecule placed

within the active site appeared highly mobile in all three species, although this mobility

seemed reduced in the Closed1 and Closed2 forms. No additional density was detected

within the active sites beyond that attributable to the cofactor.

Three conformers of mL-GDH$_{180}$ subunits reveal modulations in tertiary structure

associated with differential active site occupancies

To clarify the absence of signal for any ligand other than the cofactor in the cryoEM maps

of mL-GDH$_{180}$ tetramers, we performed a hierarchical classification of all mL-GDH$_{180}$

subunits, regardless of whether they belonged to the Open, Closed1 or Closed2

conformers, using a mask that covered the catalytic domain (Figure S3). This classification

yielded three distinct forms for mL-GDH$_{180}$ monomers, termed empty-monomer, cofactor-

monomer and cofactor/ligand-monomer, with estimated resolutions of 3.07 Å, 2.95 Å and

3.12 Å, respectively (Figure 2 and Table 1). The empty-monomer map shows faint density

within the cofactor binding pocket, but does not reveal strong binding to that site, nor any

signal in the substrate binding pocket. In contrast, the cofactor-monomer map clearly

displays a bound cofactor molecule in the catalytic cavity and the cofactor/ligand-monomer

map discloses both the cofactor and an additional density element within the active site

(Figure 2 and Figure S6). An average of all these variants resulted in a 2.63 Å resolution

map of the mL-GDH$_{180}$ monomer (Figure S6 and Figure S7), which was used to model the

polypeptide segment comprising residues 463–1588 before refining the atomic

coordinates for each occupancy state.

The atomic model derived from the empty-monomer map revealed the arrangement of

residues that, according to previous analyses [2,3], define the dinucleotide binding region

(DNBR) and the glutamate binding region (GluBR) of mL-GDH$_{180}$ (Figure 2). The cryoEM

maps of the cofactor- and cofactor/ligand- monomers of mL-GDH$_{180}$ allowed to place a

NAD$^+$ molecule in extended conformation within the DNBR (Figure 2). According to our

models, the pyrophosphate moiety is located in the crevice formed between strands β1

and β5 (residues 953-958 and 1070-1073, respectively) of subdomain II of the catalytic

domain (SDII) and is stabilized by the dipole of the helix comprising residues 964-969,

which connects strands β1 and β2 (residues 976-982) and is known as the dinucleotide

binding helix (Figure 3). The adenine ring of NAD$^+$ fits well in the density, the glycosidic

bond to the nicotinamide ring adopts a *syn* conformation and both riboses display C2'

endo puckering, in agreement with previous reports for S-GDHs [15].

Notably, the pattern of NAD$^+$-protein interactions differs between cofactor- and

cofactor/ligand- mL-GDH$_{180}$ monomers (Figure 2). In the cofactor-monomer, the adenine

ring interacts with the side chain (sc) of Tyr1079, the 2' and 3' hydroxyl groups of the

adenosine ribose point towards the surface of the protein and hydrogen bond Asp983 (sc)

and the main chain (mc) of Asp960, respectively, phosphates interact with the amide

nitrogen of Asp964 and Val965, the nicotinamide ribose hydrogen bonds Gly1075 (mc)

and Asn1115 (mc and sc), and the nicotinamide amide contacts His923 (sc), Thr929 (sc)

and Asp964 (sc). In contrast, in the cofactor/ligand-monomer, the adenine of NAD$^+$

interacts with Tyr1079 (sc) as well as His984 (sc), both adenosine hydroxyls hydrogen

bond Asp960 (sc and mc), the nicotinamide ribose interacts only with Asn1115 (mc and

sc) and the nicotinamide amide contacts only Thr929 (sc) and Asp964 (sc). These

differences, along with subtle but detectable conformational changes in segments 958-966

and 981-987 that contribute to the DNBR, correlate with a shift of the cofactor toward the

GluBR in the cofactor/ligand-monomer. In this state, a novel density element becomes

evident (Movie S2), within which it is possible to position a substrate molecule (CC = 0.79)

such that the α-carbon of the amino acid lies 3.3 Å from the C4 atom of the nicotinamide

ring of the cofactor, explaining the continuous density observed between them. Hence, the

γ-carboxylate of the ligand forms salt bridges with Arg780 (sc).

Strikingly, segment 914-924, which lines the active site cleft, and region 886-901, which

borders the glutamate binding pocket, adopt distinct conformations in the empty-, cofactor-

and cofactor/ligand- mL-GDH$_{180}$ monomers (Figure 2). These variations are associated with alternate conformations of segment 825-838, which is better defined in the maps of the monomers than in the tetrameric reconstructions. Thus, residues 825-827 and 830-838 could be built for the empty- and cofactor- monomers, while residues 825 and 829-838 were modeled for the cofactor/ligand-monomer. In the empty-monomer, Arg835 is directed toward Asp964 and Asp960, while Asp834 faces His984. When the cofactor binds within the DNBR, engaging residues Asp964, Asp960 and His984, Arg835 and Asp834 shift from their positions and the segment 830-838 loosens its packing against the catalytic core. When the GluBR is also occupied, segment 831-837 packs against residues 889-894. The amino acid string 825-838 of mL-GDH$_{180}$ is found in L-GDHs$_{180}$ but not in related low molecular weight enzymes (Figure 3 and Figure S5) [16].

GluBR occupancy is linked to the coupling of the catalytic domains of two adjacent monomers in the Closed mL-GDH$_{180}$ conformations

To elucidate structural changes in the context of mL-GDH$_{180}$ tetramers, we next discriminated the monomeric conformers according to whether they originated from Open, Closed1 and Closed2. This analysis revealed nine distinct species: Open-empty-monomer, Closed1-empty-monomer, Closed2-empty-monomer (3.5 Å, 3.35 Å and 3.48 Å resolution, respectively), Open-cofactor-monomer, Closed1-cofactor-monomer, Closed2-cofactor-monomer (3.3 Å, 3.23 Å and 3.28 Å), Open-cofactor/ligand-monomer, Closed1-cofactor/ligand-monomer and Closed2-cofactor/ligand-monomer (3.23 Å, 3.11 Å and 3.18 Å) (Figure S3).

No discernable structural differences were observed among the empty- or cofactor-monomers across the tetrameric states. In contrast, cryoEM maps of the cofactor/ligand-monomers displayed heterogeneity within the glutamate binding pocket. In the Open-cofactor/ligand-monomer map, a continuous density extends from the cofactor

nicotinamide ring into the GluBR (Figure 4, see also Figure 2), although it fades in the

region where the γ-carboxylate of the substrate would be stabilized. Conversely, the maps

of the Closed1- and Closed2-cofactor/ligand monomers display ligand density within the

GluBR primarily near Arg780, with a weaker signal in the Closed2 state.

As noted earlier, incubation of mL-GDH$_{180}$ with NAD$^+$ and L-glutamate induces remodeling

of the region 825-838 (Figure 2). Strikingly, this segment interacts with residues 704-706

of the catalytic domain of a neighboring N-terminally-linked subunit, giving rise to an

additional molecular bridge in the Closed tetramers (Figure 4). This intersubunit interaction

appears strengthened in the Closed2 state (Table 1). Importantly, it is only observed when

the monomer contributing the 825-838 region has both its cofactor and substrate binding

pockets simultaneously occupied. The interaction occurs approximately 2 nm from the

active site of the ligand-loaded monomer (Movie S3), linking GluBR occupancy to the

physical contact between adjacent catalytic domains and suggesting a structural basis for

turnover regulation.

DISCUSSION

GDHs constitute a large family of enzymes that function in all kingdoms of life at the crossroad between the Krebs cycle and ammonium assimilation. While S-GDHs have been studied for over half a century, it was not until 2021 that the first experimental evidence of the 3D architecture of an L-GDH was obtained [3]. In that work, we determined that the quaternary structure of mL-GDH$_{180}$ differs radically from that of S-GDHs, and that such assembly is shaped by the distinctive N- and C-terminal extensions characteristic of L-GDHs [3]. Here we present cryoEM data reaching resolutions up to 2.63 Å for mL-GDH$_{180}$ (Table1), captured in the presence of NAD$^+$ during the oxidative deamination of L-glutamate.

Our 3D classification of subunits successfully separated mL-GDH$_{180}$ monomers with both the DNBR and GluBR simultaneously occupied, as well as monomers bound to NAD$^+$ alone (Figure 1, Figure 2 and Figure 4). The NAD$^+$ binding mode in mL-GDH$_{180}$ (Figure 2 and Figure 4) mirrors that described for CsGDH [15]. The adenine ring is stabilized by hydrogen bonds with His984 (Pro262 in CsGDH) and Tyr1079 (Asp325), the NH and CO groups of the nicotinamide amide contact Asp964 (Asn240) and Thr929 (Thr209), respectively, the phosphates interact with Asp964 and Val965 (Val241), while the nicotinamide ribose hydrogen bonds Gly1075 (Ala321) and Asn1115 (Asn347). On the other hand, the 2'-OH of the adenosine ribose, along with the 3'-OH, interacts with Asp960 (Phe238) and is additionally surrounded by Asp983 (Gly261) and His984, occupying a relatively more spacious pocket than in CsGDH [15]. Considering that His984 and Arg985 (Asp263) might stabilize the 2'-phosphate of NADP$^+$, one possible explanation for mL-GDH$_{180}$ specificity for NAD$^+$ [17,18] is that this enzyme, as well as related L-GDHs [2,19–24], evolved to favor NAD$^+$ via the interactions of the 2'-OH group with Asp960 and Asp983. These residues might also exert a repulsive effect on the 2'-phosphate of NADP$^+$. This interpretation aligns with previous suggestions that NAD$^+$ specificity is determined by

interactions between the 2'-OH of the adenosine ribose and both the Gly-rich loop (GxGxxA/S, as defined in L-GDHs$_{180}$ [2]) and an acidic residue at the so-called position 7 (Asp983 in mL-GDH$_{180}$) of the fingerprint region of the nucleotide binding fold [25,26].

A seminal study on *Cs*GDH established that amino acid specificity is conferred by Lys89 and Ser380, which interact with the γ-carboxylate of the substrate [27]. In mL-GDH$_{180}$, Arg780 (Lys89) is suitably positioned to form salt bridges with the γ-carboxylate of L-glutamate (Figure 2). Though the hydroxyl of Ser1147 (Ser380) lies 4.3 Å from the γ-carboxylate of L-glutamate, slightly beyond hydrogen bond range, its position corresponds to that of Ser360 in *Cs*GDH. Also, residues Ala886 and Val1144 (Ala163 and Val377), Gly814 and Ala815 (Gly122 and Gly123), Gly781 and Gly782 (Gly90 and Gly91) and Gly1143 (Gly376), previously highlighted for their roles in establishing hydrophobic interactions with the γ- and δ-carbons of L-glutamate, shaping the L-glutamate binding pocket, orienting functional groups in the catalytic center and being located near the binding site of the nicotinamide ring of the dinucleotide, respectively [27], are similarly arranged in the mL-GDH$_{180}$ cofactor/ligand-bound monomers. This structural similitudes underscore the conservation of key amino acid specificity determinants in mL-GDH$_{180}$ (and related L-GDHs; see Figure S5 and [2]) and S-GDHs.

The conservation of active site residues indicates that L-GDHs operate through a catalytic mechanism essentially identical to that of S-GDHs [3,27]. Our atomic model of the mL-GDH$_{180}$ cofactor/ligand-monomer aimed to account for the density observed in the GluBR (Figure 2). In this model, the α-carboxylate and amino groups of L-glutamate adopt a slightly different orientation than in *Cs*GDH [27]. Although the α-carboxylate of L-glutamate faces Lys806 (Lys113), their separation (4.1 Å) is longer than ideal for direct interaction. Similarly, the α-carboxylate of the substrate lays 6.4 Å from the side chain of Gln803 (Gln110), and the amino group is 4.9 Å from the carbonyl of Ala887 (Gly164) and 6.7 Å from the side chain of the catalytic Asp888 (Asp165). It is worth noting that the packing of

residues 825-838 against segment 889-894 may help stabilize the position of Asp888,

even though its side chain is only partially defined in cryoEM maps. Nevertheless, the

cofactor/ligand-monomer maps (Figure 2 and Figure 4) reveal density corresponding to a

group modeled as the α-carbon of L-glutamate, located 3.2 Å from the C4 atom of the

cofactor nicotinamide, and density 3-3.5 Å from the Ala887 carbonyl, consistent with the

position of the leaving NH_3 group during catalysis [27].

Taken together with previous findings [3], our data confirm that mL-GDH$_{180}$ exhibits

pronounced conformational flexibility. In the absence of ligands, the enzyme populates two

tetrameric forms, Open and Closed, detected at 4.19 and 6.6 Å resolution, respectively [3].

In the present study, under conditions of saturating NAD$^+$ (4 mM) and sub-saturating L-

glutamate (42 mM) (see also Figure S1), three tetrameric states, Open, Closed1 and

Closed2, were observed, with evidence of DNBR occupancy across the full ensemble

(Figure 1). Thus, compared to the ligand-free state, these conditions appear to shift the

conformational equilibrium toward the Closed forms. The segment 825-838 adopts distinct

geometries depending on whether the cofactor is anchored to the DNBR and whether L-

glutamate (or derivatives thereof) is present in the GluBR (Figure 2). In monomers where

both regions are simultaneously occupied (cofactor/ligand-monomers), the conformation of

this segment is compatible with the establishment of an interaction with residues 704-706

of a neighboring subunit connected by the N-terminal region (Figure 4).

Unlike in S-GDH [27,28], we did not observe changes in the relative positions of SDI and

SDII within the catalytic domain of mL-GDH$_{180}$. Instead, conformational transitions among

oligomeric states were accompanied by adjustments at the ACT*3 and C-terminal domain

interfaces, changes in the intermonomeric contact between residues 650 (in the HM3) and

790 (in the catalytic domain), and the linkage between segments 825-838 and 704-706

(both in the catalytic domain), the latter forming only when both substrate and cofactor

occupy the active site. These long-range transitions are presumably slower than catalysis

and are thus observable in our experiment. While our data do no permit conclusions about

positional and/or compositional heterogeneity within the GluBR or DNBR during catalysis,

the most reasonable scenario is that the reaction occurs when both regions are loaded

(cofactor/ligand-monomer, Figure 4), with the cofactor approaching the substrate binding

pocket (Figure 2). Certainly, catalysis is expected to involve a localized conformational

adjustment that enables the side chain of Asp888 to approach the substrate's amino group

for proton abstraction in the first step of the reaction [27]. Notably, in the absence of

effectors, we observed no evidence of cooperativity for mL-GDH$_{180}$-catalyzed L-glutamate

oxidation, either at optimal pH (6.5-7.0) or across a range of pH conditions tested using

diverse buffers. However, it is intriguing whether [3] and how [17] this behavior might

change in the presence of activity modulators.

We found that ligand binding to the active (orthosteric) site of mL-GDH$_{180}$ is linked to

tertiary structure changes that lead to the consolidation of an oligomeric interface two

nanometers away from the catalytic center [29]. It is worth considering whether and how

natural or designed molecules binding to the 825-838 and/or 704-706 segments could

influence catalysis and whether occupancy of the active site modulates such regulation

[30]. Our findings open new avenues for high-resolution studies on catalysis and

regulation of L-GDHs of medical and industrial relevance.

REFERENCES

[1]   M. Li, C. Li, A. Allen, C.A. Stanley, T.J. Smith, Glutamate dehydrogenase: Structure, allosteric regulation, and role in insulin homeostasis, Neurochem. Res. 39 (2014) 433–445. https://doi.org/10.1007/s11064-013-1173-2.

[2]   B. Miñambres, E.R. Olivera, R.A. Jensen, J.M. Luengo, A new class of glutamate dehydrogenases (GDH): Biochemical and genetic characterization of the first member, the AMP-requiring NAD-specific GDH of Streptomyces clavuligerus, J. Biol. Chem. 275 (2000) 39529–39542. https://doi.org/10.1074/jbc.M005136200.

[3]   M. Lázaro, R. Melero, C. Huet, J.P. López-Alonso, S. Delgado, A. Dodu, E.M. Bruch, L.A. Abriata, P.M. Alzari, M. Valle, M.-N. Lisa, 3D architecture and structural flexibility revealed in the subfamily of large glutamate dehydrogenases by a mycobacterial enzyme, Commun. Biol. 4 (2021) 684. https://doi.org/10.1038/s42003-021-02222-x.

[4]   E.J.M. Lang, P.J. Cross, G. Mittelstädt, G.B. Jameson, E.J. Parker, Allosteric ACTion: The varied ACT domains regulating enzymes of amino-acid metabolism, Curr. Opin. Struct. Biol. 29 (2014) 102–111. https://doi.org/10.1016/j.sbi.2014.10.007.

[5]   A. Möglich, R.A. Ayers, K. Moffat, Structure and Signaling Mechanism of Per-ARNT-Sim Domains, Structure 17 (2009) 1282–1294. https://doi.org/10.1016/j.str.2009.08.011.

[6]   J.L. Gallant, A.J. Viljoen, P.D. Van Helden, I.J.F. Wiid, Glutamate dehydrogenase is required by mycobacterium bovis BCG for resistance to cellular stress, PLoS One 11 (2016) 1–14. https://doi.org/10.1371/journal.pone.0147706.

[7]   S.Q. Zheng, E. Palovcak, J.-P. Armache, K.A. Verba, Y. Cheng, D.A. Agard, MotionCor2: anisotropic correction of beam-induced motion for improved cryo-electron microscopy, Nat. Methods 14 (2017) 331–332.

https://doi.org/10.1038/nmeth.4193.

[8]   A. Rohou, N. Grigorieff, CTFFIND4: Fast and accurate defocus estimation from

electron micrographs, J. Struct. Biol. 192 (2015) 216–221.

https://doi.org/10.1016/j.jsb.2015.08.008.

[9]   J. Zivanov, T. Nakane, B.O. Forsberg, D. Kimanius, W.J.H. Hagen, E. Lindahl,

S.H.W. Scheres, New tools for automated high-resolution cryo-EM structure

determination in RELION-3, Elife 7 (2018) e42166.

https://doi.org/10.7554/eLife.42166.

[10] J. He, T. Li, S.Y. Huang, Improvement of cryo-EM maps by simultaneous local and

non-local deep learning, Nat. Commun. 14 (2023) 1–16.

https://doi.org/10.1038/s41467-023-39031-1.

[11] E.F. Pettersen, T.D. Goddard, C.C. Huang, G.S. Couch, D.M. Greenblatt, E.C.

Meng, T.E. Ferrin, UCSF Chimera - A visualization system for exploratory research

and analysis, J. Comput. Chem. 25 (2004) 1605–1612.

https://doi.org/10.1002/jcc.20084.

[12] P. Emsley, B. Lohkamp, W.G. Scott, K. Cowtan, Features and development of Coot,

Acta Crystallogr. Sect. D Biol. Crystallogr. 66 (2010) 486–501.

https://doi.org/10.1107/S0907444910007493.

[13] P. V. Afonine, B.K. Poon, R.J. Read, O. V. Sobolev, T.C. Terwilliger, A. Urzhumtsev,

P.D. Adams, Real-space refinement in PHENIX for cryo-EM and crystallography,

Acta Crystallogr. Sect. D Struct. Biol. 74 (2018) 531–544.

https://doi.org/10.1107/S2059798318006551.

[14] E. Krissinel, K. Henrick, Protein interfaces, surfaces and assemblies' service PISA at

the european bioinformatics institute., J. Mol. Biol. 372 (2007) 774–797.

http://www.ebi.ac.uk/pdbe/prot_int/pistart.html.

[15] P.J. Baker, K.L. Britton, P.C. Engel, G.W. Farrants, K.S. Lilley, D.W. Rice, T.J.

Stillman, Subunit assembly and active site location in the structure of glutamate

dehydrogenase, Proteins Struct. Funct. Bioinforma. 12 (1992) 75–86.

https://doi.org/10.1002/prot.340120109.

[16] K.L. Britton, P.J. Baker, D.W. Rice, T.J. Stilllman, Structural relationship between the

hexameric and tetrameric family of glutamate dehydrogenases, Eur. J. Biochem. 209

(1992) 851–859. https://doi.org/10.1111/j.1432-1033.1992.tb17357.x.

[17] H.M. O'Hare, R. Durán, C. Cerveñansky, M. Bellinzoni, A.M. Wehenkel, O. Pritsch,

G. Obal, J. Baumgartner, J. Vialaret, K. Johnsson, P.M. Alzari, Regulation of

glutamate metabolism by protein kinases in mycobacteria, Mol. Microbiol. 70 (2008)

1408–1423. https://doi.org/10.1111/j.1365-2958.2008.06489.x.

[18] T.J. Nott, G. Kelly, L. Stach, J. Li, S. Westcott, D. Patel, D.M. Hunt, S. Howell, R.S.

Buxton, H.M. O'Hare, S.J. Smerdon, An intramolecular switch regulates

phosphoindependent FHA domain interactions in Mycobacterium tuberculosis, Sci.

Signal. 2 (2009) ra12. https://doi.org/10.1126/scisignal.2000212.

[19] F.M. Veronese, J.F. Nyc, Y. Degani, D.M. Brown, E.L. Smith, Nicotinamide adenine

dinucleotide specific glutamate dehydrogenase of Neurospora. I. Purification and

molecular properties, J. Biol. Chem. 249 (1974) 7922–7928.

https://doi.org/10.1016/S0021-9258(19)42053-X.

[20] C.D. Lu, A.T. Abdelal, The gdhB gene of Pseudomonas aeruginosa encodes an

arginine-inducible NAD+-dependent glutamate dehydrogenase which is subject to

allosteric regulation, J. Bacteriol. 183 (2001) 490–499.

https://doi.org/10.1128/JB.183.2.490-499.2001.

[21] L. Camardella, R. Di Fraia, A. Antignani, M.A. Ciardiello, G. Di Prisco, J.K. Coleman,

L. Buchon, J. Guespin, N.J. Russell, The Antarctic Psychrobacter sp. TAD1 has two

cold-active glutamate dehydrogenases with different cofactor specificities.

Characterisation of the NAD+-dependent enzyme, Comp. Biochem. Physiol. - A Mol.

Integr. Physiol. 131 (2002) 559–567. https://doi.org/10.1016/S1095-6433(01)00507-

4.

[22] R. Kawakami, H. Sakuraba, T. Ohshima, Gene cloning and characterization of the

very large NAD-dependent L-glutamate dehydrogenase from the psychrophile

Janthinobacterium lividum, isolated from cold soil, J. Bacteriol. 189 (2007) 5626–

5633. https://doi.org/10.1128/JB.00496-07.

[23] R. Kawakami, M. Oyama, H. Sakubara, T. Ohshima, The Unique Kinetic Behavior of

the Very Large NAD-Dependent Glutamate Dehydrogenase from Janthinobacterium

lividum, Biosci. Biotechnol. Biochem. 74 (2010) 884–887.

https://doi.org/10.1271/bbb.90925.

[24] F. Beaufay, J. Coppine, A. Mayard, G. Laloux, X. De Bolle, R. Hallez, A NAD-

dependent glutamate dehydrogenase coordinates metabolism with cell division in

Caulobacter crescentus, EMBO J. 34 (2015) 1786–1800.

https://doi.org/10.15252/embj.201490730.

[25] P.J. Baker, K.L. Britton, D.W. Rice, A. Rob, T.J. Stillman, Structural consequences

of sequence patterns in the fingerprint region of the nucleotide binding fold.

Implications for nucleotide specificity, J. Mol. Biol. 228 (1992) 662–671.

https://doi.org/10.1016/0022-2836(92)90848-E.

[26] P.C. Engel, Glutamate dehydrogenases: The why and how of coenzyme specificity,

Neurochem. Res. 39 (2014) 426–432. https://doi.org/10.1007/s11064-013-1089-x.

[27] T.J. Stillman, P.J. Baker, K.L. Britton, D.W. Rice, Conformational flexibility in

glutamate dehydrogenase: Role of water in substrate recognition and catalysis, J.

Mol. Biol. 234 (1993) 1131–1139. https://doi.org/10.1006/jmbi.1993.1665.

[28] M.J. Borgnia, S. Banerjee, A. Merk, D. Matthies, A. Bartesaghi, P. Rao, J. Pierson,

L.A. Earl, V. Falconieri, S. Subramaniam, J.L.S. Milne, Using cryo-EM to map small

ligands on dynamic metabolic enzymes: Studies with glutamate dehydrogenase,

517 Mol. Pharmacol. 89 (2016) 645–651. https://doi.org/10.1124/mol.116.103382.

518 [29] T. Perica, C. Chothia, S.A. Teichmann, Evolution of oligomeric state through

519 geometric coupling of protein interfaces, Proc. Natl. Acad. Sci. 109 (2012) 8127–

520 8132. https://doi.org/10.1073/pnas.1120028109.

521 [30] J.O. Schulze, G. Saladino, K. Busschots, S. Neimanis, E. Süß, D. Odadzic, S.

522 Zeuzem, V. Hindie, A.K. Herbrand, M.-N. Lisa, P.M. Alzari, F.L. Gervasio, R.M.

523 Biondi, Bidirectional Allosteric Communication between the ATP-Binding Site and

524 the Regulatory PIF Pocket in PDK1 Protein Kinase, Cell Chem. Biol. 23 (2016)

525 1193–1205. https://doi.org/10.1016/j.chembiol.2016.06.017.

526 [31] J. Abramson, J. Adler, J. Dunger, R. Evans, T. Green, A. Pritzel, O. Ronneberger, L.

527 Willmore, A.J. Ballard, J. Bambrick, S.W. Bodenstein, D.A. Evans, C.C. Hung, M.

528 O'Neill, D. Reiman, K. Tunyasuvunakool, Z. Wu, A. Žemgulytė, E. Arvaniti, C.

529 Beattie, O. Bertolli, A. Bridgland, A. Cherepanov, M. Congreve, A.I. Cowen-Rivers,

530 A. Cowie, M. Figurnov, F.B. Fuchs, H. Gladman, R. Jain, Y.A. Khan, C.M.R. Low, K.

531 Perlin, A. Potapenko, P. Savy, S. Singh, A. Stecula, A. Thillaisundaram, C. Tong, S.

532 Yakneen, E.D. Zhong, M. Zielinski, A. Žídek, V. Bapst, P. Kohli, M. Jaderberg, D.

533 Hassabis, J.M. Jumper, Accurate structure prediction of biomolecular interactions

534 with AlphaFold 3, Nature 630 (2024) 493–500. https://doi.org/10.1038/s41586-024-

535 07487-w.

ACKNOWLEDGMENTS

This work was supported by grants PICT 2019-00989 and PICT 2021-00253 from the Agencia Nacional de Promoción de la Investigación, el Desarrollo Tecnológico y la Innovación (Agencia I+D+i, Argentina) to MNL as well as grants PID2021-124074NB-I00 and PID2021-125946OB-I00 from Agencia Estatal de Investigación of Spain to MV and GJO, respectively. We also thank the Severo Ochoa Center of Excellence Accreditation CEX2021-001136-S. MNL is a career investigator from Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET, Argentina). NC holds a CONICET PhD fellowship. NC and MNL are grateful to Liliana Rojas for excellent technical assistance. Some of this work was performed at the Basque Resource for Electron Microscopy located at Instituto Biofisika (UPV/EHU, CSIC), supported by the Department of Science, Universities and Innovation and the Innovation Fund of the Basque Government, with additional support from MCIN (Recovery, Transformation and Resilience Plan) and the Basque Government "Biotechnology Complementary Plan Applied to Health" with funding from European Union NextGenerationEU (PRTR-C17.I1; PRTR-C17.I01.P01.S13) (AAAA_ACG_AY_2539/22_05).

AUTHOR CONTRIBUTIONS

ML produced protein, prepared EM samples, performed EM experiments, processed and analyzed EM data and performed structural analyses; NC produced protein, performed kinetic analyses, analyzed EM data and performed structural analyses; JPLA collected and analyzed EM data; DC assisted during processing of EM data; RMR performed structural analyses; GJO performed structural analyses; MV designed experiments, supervised EM data collection and processing and analyzed results; MNL designed experiments, supervised biochemical experiments, refined atomic models, analyzed data and wrote the paper. All authors read and corrected the paper.

COMPETING INTERESTS

565   The authors declare no competing interests.

FIGURES

Figure 1

Three conformers of mL-GDH$_{180}$ detected upon incubation with L-glutamate and

NAD$^+$ reveal modulations of the enzyme quaternary structure. (A) CryoEM mL-GDH$_{180}$

maps obtained after incubating the enzyme with L-glutamate and NAD$^+$. The dimensions

of each conformer are provided in Å. Maps cover, for each monomer, the region indicated

by a dotted rectangle in the diagram shown below (residues 500-1588). ACT*, Aspartate

kinase-Chorismate mutase-TyrA-like; PAS, Per-Arnt-Sim; HM, helical motif; CD, catalytic

domain; CTD, C-terminal domain. (B) Approach of the catalytic domains of mL-GDH$_{180}$

subunits linked by the N-terminal region. Segment 825-838, poorly defined in maps of the

Open, Closed1 and Closed2 enzyme forms, approaches the catalytic domain in a

neighboring subunit. In each case, a zoomed in view of the dinucleotide binding region

(DNBR) is shown below. The protein is displayed in ribbon representation and cryoEM

maps are depicted in grey. The nicotinamide portion of the NAD$^+$ molecule (in sticks)

exhibits high mobility in all conformations. Subdomains I and II (SDI and SDII) of the

catalytic domain are colored in yellow and orange, respectively, whereas ACT*3 and HM3

motifs are shown in cyan and green, respectively, for one monomer (the one on the left),

with the neighboring subunit (the one on the right) shown in grey. All maps are shown with

a contour level of 0.02. GluBR, glutamate binding region.

Figure 2

Three conformers of mL-GDH$_{180}$ subunits reveal modulations in tertiary structure associated with differential active site occupancies. Alternative conformations of mL-GDH$_{180}$ subunits are shown, with the active site either empty or containing NAD$^+$ or NAD$^+$ and L-glutamate. The average RMSD for 1122 aligned residues in pairwise comparisons

of the atomic models (show in ribbon representation) is 0.7 Å. NAD$^+$ and L-glutamate are

depicted as sticks as well as highlighted residues. For each species, subdomains I and II

of the catalytic domain (SDI, residues 702-926 and 1188-1220; SDII, residues 927-1187)

are colored in yellow and orange, respectively. A small portion of the C-terminal domain is

displayed in magenta. Interactions established by the segment 825-838 with other portions

of the polypeptide chain are shown in the panels on the left. Note that within the region

825–838, residues 825-827 and 830-838 could be built for the empty- and cofactor-

monomers, while residues 825 and 829-838 were modeled for the cofactor/ligand-

monomer. A comparison of the atomic models of the empty- (green), cofactor- (blue) and

cofactor/ligand-monomer (gray) is presented in the lower left corner. CryoEM maps are

depicted in grey, with contour levels of 5. H-bonds and salt bridges are displayed as

dashed lines. GluBR, glutamate binding region; DNBR, dinucleotide binding region.

Figure 3

Comparison of the catalytic domain of mL-GDH$_{180}$ with that of S-GDHs. A monomer of

mL-GDH$_{180}$ is shown on the left. Segments 824-838 and 858-882 (in dark green), both

located in the catalytic domain subdomain I (SDI), are absent in S-GDHs [2]. For the 824-

838 portion coordinates were taken from a model calculated with AlphaFold 3 [31]. A

monomer of *Cs*GDH (*Cs*, *Clostridium symbiosum*; PDB 1BGV [27]), considered a

representative S-GDH, is shown on the right. The SDI of S-GDHs presents secondary

structure elements (in dark blue, named according to the consensus nomenclature for S-

GDHs) with conserved roles in oligomerization in the enzyme subfamily, which are not

present in mL-GDH$_{180}$ or related L-GDHs. For both mL-GDH$_{180}$ and *Cs*GDH, the six-

stranded 432156 β-sheet that shapes the DNBR in the catalytic domain subdomain II

(SDII) is shown in grey; the linker regions β4-β5 (in light green) and β5-β6 (in cyan)

encompass a larger number of residues in mL-GDH$_{180}$. For the two proteins, the SDI and

SDII are shown, except for otherwise highlighted portions, in yellow and orange,

respectively.

Figure 4

GluBR occupancy is linked to the coupling of the catalytic domain of two adjacent

monomers in the Closed conformations. Top panel: the atomic model of the

cofactor/ligand-monomer of mL-GDH$_{180}$ is shown together with maps of this species

obtained from Open, Closed1 or Closed2 tetramers (Open-, Closed1- and Closed2-

cofactor/ligand-monomers; contour levels are 5). NAD$^+$ and L-glutamate as well as

highlighted residues are depicted as sticks. The subdomains I and II of the catalytic

domain (SDI and SDII) are colored in yellow and orange, respectively. Bottom panel: in the

context of tetramers containing a cofactor/ligand-monomer (cofactor/ligand-tetramers;

contour levels are 0.015), in a progression towards the Closed1 and Closed2 conformers,

the stabilization of an interaction between segment 825-838 of one subunit (colored, on

the left) and segment 704-706 of the neighboring one (in grey, on the right) is observed.

The atomic coordinates of the subunit on the left correspond in each case to those of a

cofactor/ligand-monomer, replacing the equivalent ones presented in Figure 1. H-bonds

and salt bridges are displayed as dashed lines. CryoEM maps are displayed in grey.

TABLES

Table 1. CryoEM data collection and processing.

	Open (EMD-52419)	Closed1 (EMD-52420)	Closed2 (EMD-52421)
Data collection and processing			
Magnification	130,000	130,000	130,000
Voltage (kV)	300	300	300
Electron dose ($e^-/Å^2$)	0.979	0.979	0.979
Defocus range (μm)	1.0-1.6	1.0-1.6	1.0-1.6
Pixel size (Å)	1.29	1.29	1.29
Symmetry imposed	D2	D2	D2
Initial particles images (no.)	667,866	667,866	667,866
Final particles images (no.)	37,641	45,998	32,441
Map resolution (Å)	3.88	3.54	3.57
FSC threshold	0.143	0.143	0.143
Map sharpening B factor ($Å^2$)	-50	-50	-50
Refinement			
Model resolution (Å)	3.88	3.54	3.57
Model composition			
Non-hydrogen atoms	33576	33576	33576
Protein residues	4300	4300	4300
Ligands	4 NAD$^+$	4 NAD$^+$	4 NAD$^+$
B factors ($Å^2$)			
Protein	224.33	189.71	159.33
Ligands	273.06	208.84	186.14
R.m.s. deviations			
Bond lengths (Å)	0.001	0.002	0.002
Bond angles (°)	0.371	0.409	0.395
Validation			
MolProbity score	1.69	1.62	1.50
Clashscore	10.54	11.11	9.42
Poor rotamers (%)	0.11	0	0.11
Ramachandran plot			
Favored (%)	97.20	97.76	98.55
Allowed (%)	2.80	2.24	1.45
Disallowed (%)	0	0	0
CC (mask)	0.83	0.77	0.83
CC (box)	0.91	0.87	0.89
CC (peaks)	0.77	0.70	0.77
CC (volume)	0.82	0.76	0.82
Mean CC for ligands	0.85	0.82	0.80

	empty-monomer (EMD-52422)	cofactor-monomer (EMD-52423)	cofactor/ligand-monomer (EMD-52424)
Data collection and processing			
Magnification	130,000	130,000	130,000
Voltage (kV)	300	300	300
Electron dose ($e^-/Å^2$)	0.979	0.979	0.979
Defocus range (μm)	1.0-1.6	1.0-1.6	1.0-1.6
Pixel size (Å)	1.29	1.29	1.29
Symmetry imposed	C1	C1	C1
Initial particles images (no.)	464,320	464,320	464,320
Final particles images (no.)	131,216	168,668	112,701
Map resolution (Å)	3.07	2.95	3.12
FSC threshold	0.143	0.143	0.143
Map sharpening B factor	LS	LS	LS
Refinement			
Model resolution (Å)	3.07	2.95	3.12
Model composition			
Non-hydrogen atoms	8717	8761	8760
Protein residues	1124	1124	1123
Ligands	-	1 NAD^+	1 NAD^+/ 1 L-Glu
B factors ($Å^2$)			
Protein	64.32	62.58	72.09
Ligands	-	56.65	81.37
R.m.s. deviations			
Bond lengths (Å)	0.004	0.004	0.004
Bond angles (°)	0.691	0.721	0.693
Validation			
MolProbity score	1.37	1.27	1.37
Clashscore	5.53	5.16	5.80
Poor rotamers (%)	0.55	0.55	0.11
Ramachandran plot			
Favored (%)	97.68	98.48	97.77
Allowed (%)	2.32	1.52	2.23
Disallowed (%)	0	0	0
CC (mask)	0.87	0.88	0.87
CC (box)	0.89	0.89	0.89
CC (peaks)	0.81	0.81	0.79
CC (volume)	0.88	0.88	0.87
Mean CC for ligands	-	0.91	0.84

	Open-cofactor/ligand-monomer (EMD-52425)	Closed1-cofactor/ligand-monomer (EMD-52426)	Closed2-cofactor/ligand-monomer (EMD-52427)
Data collection and processing			
Magnification	130,000	130,000	130,000
Voltage (kV)	300	300	300
Electron dose (e⁻/Å²)	0.979	0.979	0.979
Defocus range (μm)	1.0-1.6	1.0-1.6	1.0-1.6
Pixel size (Å)	0.646	0.646	0.646
Symmetry imposed	C1	C1	C1
Initial particles images (no.)	112,701	112,701	112,701
Final particles images (no.)	33,330	46,380	32,991
Map resolution (Å)	3.23	3.11	3.18
FSC threshold	0.143	0.143	0.143
Map sharpening B factor	LS	LS	LS
Refinement			
Model resolution (Å)			
Model composition			
Non-hydrogen atoms			
Protein residues			
Ligands			
B factors (Å²)			
Protein			
Ligands			
R.m.s. deviations			
Bond lengths (Å)			
Bond angles (°)			
Validation			
MolProbity score			
Clashscore			
Poor rotamers (%)			
Ramachandran plot			
Favored (%)			
Allowed (%)			
Disallowed (%)			
CC (mask)			
CC (box)			
CC (peaks)			
CC (volume)			
Mean CC for ligands			

	total-monomer (EMD-52428)	Closed2-cofactor/ligand-tetramer (EMD-52429)
Data collection and processing		
Magnification	130,000	130,000
Voltage (kV)	300	300
Electron dose (e^-/Å^2)	0.979	0.979
Defocus range (μm)	1.0-1.6	1.0-1.6
Pixel size (Å)	0.646	1.29
Symmetry imposed	C1	C1
Initial particles images (no.)	464,320	112,701
Final particles images (no.)	464,320	32,991
Map resolution (Å)	2.63	4.10
FSC threshold	0.143	0.143
Map sharpening B factor	LS	-50
Refinement		
Model resolution (Å)	2.63	
Model composition		
Non-hydrogen atoms	8683	
Protein residues	1113	
Ligands	1 NAD$^+$	
B factors (Å^2)		
Protein	56.19	
Ligands	48.51	
R.m.s. deviations		
Bond lengths (Å)	0.002	
Bond angles (°)	0.467	
Validation		
MolProbity score	1.39	
Clashscore	7.12	
Poor rotamers (%)	0.55	
Ramachandran plot		
Favored (%)	99.10	
Allowed (%)	0.90	
Disallowed (%)	0	
CC (mask)	0.80	
CC (box)	0.68	
CC (peaks)	0.65	
CC (volume)	0.78	
Mean CC for ligands	0.81	

646
647   LS, local sharpening.