Quantitative analyses of single mitochondrial structure-function heterogeneity

uncovers stemness specification by redox-tuned small-mitochondrial-networks

Mayank Saini[1#], Swati Agarwala[1#], Biratal Wagle[1], Bharti Golchha[1], Brian Spurlock[2,3], Pranit

Sinha[1], Partha Pratim Chakrabarti[5,] Diya Prasad[1], Danitra Parker[2,4], Akshara Kulkarni[1] and

Kasturi Mitra[1,2]*

[1] Department of Biology, Trivedi School of Biological Sciences, Ashoka University, New Delhi

(NCR), India

[2] Department of Genetics, University of Alabama at Birmingham, Alabama, USA

[3] McAllister Heart Institute, Department of Pathology, University of North Carolina, Chapel Hill,

USA

[4] Dept of Integrative Biology and Pharmacology, Institute of Aging, University of Texas Health

Science Center at Houston, USA

[5]Department of Computer Science and Engineering, Indian Institute of Technology Kharagpur

*Corresponding author: Kasturi Mitra

#Equal Contribution

Email: kasturi.mitra@ashoka.edu.in (primary)

Author Contributions: M.S. performed all mito-SinComp analyses, S.A. performed all

transcriptomics analyses, B.W. wrote codes for mito-SinComp analyses, B.S. and D.P. collected

data, Di. P. performed mito-SinComp analysis, A.K. performed validation for redox

measurements, B.G. performed immunoblot analyses and mito-SinComp analyses of mt-DNA,

PS performed RF regression task with consultation from PPC, K.M. supervised all aspects and

wrote the manuscript with help from M.S., S.A., B.W, BG and PS.

Running title: SMNs specify stemness through mt-DNA-stemness transcriptomic network

Abstract

Lack of quantitative understanding of the marked heterogeneity of multifaceted mitochondria poses challenges in unraveling their translatable role. Here, we hierarchically untangled multilevel heterogeneity of mitochondrial networks and non-networks by simultaneously analysing their structure-function within single components and their units, in cells and tissues. Such mito-SinComp quantitative analyses revealed that redox levels of mitochondrial networks and their dramatic intra-network heterogeneity can be predicted by structural features through a translatable machine-learning approach. Mito-SinComp identified and quantitatively characterized a redox-tuned subpopulation of Small-Mitochondrial-Networks(SMNs) that specifies stemness. These SMNs are generated by severing of nodes of oxidized Hyperfused-Mitochondrial-Networks(HMNs), are ten-folds smaller with specific network complexity and elevated mt-DNA nucleoid abundance. Thus, HMN to SMN conversion supports elevated expression of mtDNA genes in establishing MT-ND1(redox)-KRT15(stemness) transcriptomic interaction, as revealed by coupled scRNA-seq. This stemness priming of non-transformed cells is sustained after neoplastic transformation and is detected in patient transcriptome, demonstrating translational relevance of SMNs.

Keywords: Single mitochondrial analyses / Mitochondrial network heterogeneity / Stemness / Microscopy / scRNA-seq

Introduction

Mitochondria, as dynamic organelles, are gatekeepers of multiple cellular homeostatic processes and form signalling hubs[1,2]. Mitochondria can be heterogenous between cells, tissues, and also individuals, while a cell can have few to hundreds of mitochondria of heterogenous structures including their elaborate physical networks. Reorganization of mitochondrial structure by cellular processes is linked to various disease etiologies[3–7]. Mitochondrial structure-function is primarily modulated by the counteracting processes of mitochondrial fission and fusion, and extensive molecular characterization of such processes is an active research area [6,8]. Network forming ability of mitochondria regulates fundamental mitochondrial processes and may have imparted evolutionary advantage [9–13]. However, various fundamental questions remain about how non-networked and networked mitochondrial entities interact to crosstalk with mitochondrial function and thus impact cell physiology.

Lack of quantitative knowledge of mitochondrial network heterogeneity poses challenges in developing generalized understanding of the function of mitochondrial networks, shrouding the field with controversies. Additionally, the natural heterogeneity of mitochondrial structure-function along a wide spectrum is not reflected accurately by genetic alteration of the molecular regulators. Therefore, genetic approaches sometimes lead to confusing findings preventing building knowledge consensus of the exact role of mitochondria. Moreover, genetic experiments may obscure the distinction between primary and secondary impacts due to remodelling ability of mitochondria. Therefore, there is a serious necessity of obtaining direct quantitative understanding of the heterogeneity of mitochondrial structure-function across their natural spectrum towards elucidating the interplay of mitochondria with cellular process. Various laboratories, including our own, have attempted to quantitatively define mitochondrial structure using microscopy[14–22]. We reported that studying mitochondrial structure-function in single cells characterizes mitochondria primed stemness states, similar to single cell transcriptomics[21,23,24].

Contradictory reports about the mitochondrial structure and number in stem cells clouds our understanding of the impact of mitochondria on stem cell physiology [3,25–27]. Lineage specific adult stem cells undergo neoplastic transformation into tumor initiating cells, and some are particularly vulnerable to mitochondrial inhibition [28–30] . Stemness state (characterized by self-renewal) can be sustained by repression of mitochondrial fission [3,21,23,24,31,32], while stem cells are generally considered to have rudimentary fragmented mitochondrial structures [33,34]. However, our quantitative approach revealed repression of mitochondrial fission as well as its regulated activation are required to prime a stem cell state to initiate and sustain neoplasticity [23]. Activation of mitochondrial fission is essential for oncogene induced neoplastic transformation

[35,36]. A recent report proposed that the mitochondrial fission and fusion cycles modulate

oxidative and reductive biosynthesis in same mitochondrial populations [5], highlighting

mitochondria fission and fusion act in concert in regulating distinct aspects of a given cellular

process.

We and others previously identified the hyperfused mitochondria in certain cellular

processes, which is primarily regulated by repression of mitochondrial fission [22,23,37,38]. Here, we

hierarchically untangled inter- and intra-cellular as well as intra-network mitochondrial

heterogeneity by directly quantifying mitochondrial structure-function of single mitochondrial

components. This mito-SinComp analyses revealed novel concepts in mitochondrial redox

physiology and mtDNA functional organization and their mechanistic role in stem cell

physiology. We identified and quantitatively characterized a subpopulation of small

mitochondrial networks (SMNs), which are derived from the hyperfused mitochondrial networks

(HMNs) to specify stemness. Using coupled single cell transcriptomics we identified and

validated key molecules underlying the SMN driven stemness specification, which is also

detectable in patients. Our data demonstrate that unprecedented resolution of mitochondrial

structure-function heterogeneity, enabled by mito-SinComp, allows fundamental discovery and

its translational application.

Results

Modular design of mito-SinComp analyses of single mitochondrial components for

hierarchical untangling of mitochondrial structure(S)-function(F) heterogeneity

Our single cell microscopy based approach of mitochondrial heterogeneity analyses has

revealed novel findings about quantitatively defined mitochondrial structural states in guiding

cell states [23,39]. Here, we unrestricted such analyses to study mitochondrial structure-function

heterogeneity in a 'hierarchical manner' for 'any' given mitochondrial function in 'any' given cell

physiology using a singular quantitative approach. This is a fluorescence pixel level

simultaneous quantitative analyses of structure-function of single mitochondrial components and

its units for live or fixed cells / tissues, namely mito-SinComp. Such analyses can untangle

mitochondrial structure-function heterogeneity between and within individual subjects→

tissues→ cells→ mitochondrial networks or non-networks→ their types and subtypes (based on

their features) →elements forming the networks→zones forming the elements **(Fig. 1).**

The approach involves the following steps on the total mitochondrial population in live or

fixed cells or tissues, within or beyond cellular boundaries (details in Methods): **1)** combinatorial

labelling of mitochondria with a structure probe and one or more compatible function probes

(chemical, immuno- or genetic); **2)** 3D fluorescence confocal microscopy to image the total

population of mitochondria within a cell or tissue allowing wider applicability; **3)** application of

MitoGraph software to obtain 3D 'structure' coordinates of individual mitochondrial components;

4) automated mapping of any mitochondrial 'function' values on structure coordinates of all

resolvable units for hierarchical computation of function; **5)** automated computation of length

and diameter from pixel coordinates to compute the structure (S) and function (F) features for

individual mitochondrial components (**Suppl Table 1**); **6)** studying bivariate or multivariate S-F

or S-S or F-F relationships in associative, predictive or causal manner.

Modularity of mito-SinComp design allows use of any validated single or palate of

mitochondrial functional probes for any mitochondrial sub-compartment, which are compatible

with the structural probe of choice. Its scalability allows exploration with high-content,

multiplexed, time-resolved or super-resolution microscopy, with appropriate modifications. Mito-

SinComp performed in correlative or causative manner with other experimental modalities

allows elucidation of significance of mitochondrial heterogeneity in any cell physiology. Mito-

SinComp allows creation of library or map of mitochondrial S-F quantitative features across cells

/ tissues / organisms for extensive and intensive exploratory analyses, theoretical modelling with

the mitochondrial S-F features and their relationships, and also deep mechanistic understanding

of the mitochondrial properties and their heterogeneity in any given cell physiology. Although the

primary goal of mito-SinComp does not include investigation of molecular mechanism, it can

also be used in elucidation of the impact of genetic mutations or direct studies of mitochondrial

regulators. The simplicity of the mito-SinComp approach allows fundamental and translational

studies on drug, nutrient, macromolecule or genetic screening.

In the subsequent sections, we use mito-SinComp for hierarchical untangling of

mitochondrial structure-function heterogeneity towards unraveling mechanistic role of

mitochondria in specification of stemness state. We used 3 previously demonstrated stemness

supporting conditions and tested in the situation of acute carcinogen exposure. We elucidate

mitochondrial redox function in live cells and functional mtDNA organization within mitochondrial

networks in fixed cells, and use quantitative mito-SinComp analyses in conjunction with

quantitative single cell transcriptomics. The mitochondrial components (CC) having >1

mitochondrial edges (i.e. elements) connected by >2 nodes and with >2 terminal nodes are

classified as 'networked', whereas mitochondrial components (CC) with 1 edge and 2 terminal

nodes are classified as 'non-networked'; the rare looped networked structures with one node

and one edge are exceptions. The features with non-normal distributions are log-normalized

and appropriate outlier filtering is included, as necessary. The numerical data from the

automated analyses is collected in a comprehensive data file for downstream analyses,

including statistical tests with multiple hypothesis testing adjustments. The physical properties of

the networked mitochondria is studied using 10 structure features, while the non-networked

mitochondria present fewer features, while the number of functional features may vary **(Fig. 1)**.

Heterogeneity in the feature distribution is assessed by median and coefficient of variation while

other statistical parameters of choice can be used.

Extraction of quantitative S-F features of single mitochondrial components to study

mitochondrial redox function in live cells

We used the HaCaT normal human keratinocyte model to study specification of

stemness, where we demonstrated mitochondria priming of stemness supports neoplasticity

driven by an oxidative carcinogen [23]. After validating the mito-SinComp computed total

component length of a cell with that of the MitoGraph output **(Fig. S1A)**, we tested the

comprehensive mito-SinComp analytical script against our previous benchmark[23]**.** Mito-

SinComp successfully identified the HaCaT populations harbouring the 'intermediately-fused'

(**INT**) or the 'hyperfused' (**HF**) mitochondria in a pooled population where mitochondrial structure

spectrum is widened with different levels of knockdown of the mitochondrial fission protein,

Drp1, (**Fig. S1B**). Moreover, the 3D Graphs of the mitochondrial structural coordinates and the

computed structural features for the identified networks and non-networks recapitulate the inter-

and intra-cellular mitochondrial heterogeneity of the representative Cell 1-INT and Cell 2-HF

(**Fig. 2A**, **Suppl. Table 2)**. Final validation of the mito-SinComp structural analyses comes from

the successful detection of higher abundance of longer networked mitochondria (i.e.

hyperfused) in the HF-Cells group **(Fig. 2B, left, arrow)**. The validated mito-SinComp structural

analyses revealed the following quantitative comparisons between the INT-Cells and HF-Cells

groups (**Fig. S1C-F**): **a)** comparable abundance of networks and non-networks per cell. **b)**

networks are ~10-100 times larger than the non-networks; **c)** average length of networks in

individual cells is at least 2 folds less in the INT-Cells group in comparison to the HF-Cells

group.

Given stem cells are particularly dependent on redox tuning [2,40], we chose to study

mitochondrial redox function that is bidirectionally linked with mitochondrial structure [41]. We

extracted the mito-SinComp structure and redox features from cells expressing our validated

combination of genetically encoded mitochondrial structural probe mito-PSmO2 and general

redox probe for mitochondrial matrix, mito-roGFP [21] **(Fig. 2C, right)**. First, mito-SinComp

computes the [Oxidation]$_{Px}$ values in the individual pixels and maps them to the mitochondrial
coordinates identified in the binary image (**Fig. 1,** details in Methods). Thereafter, it compute the
[Oxidation] values at different levels of structural hierarchically as follow: average of
[Oxidation]$_{Px}$ within an element yields [Oxidation]$_{EL}$---average of [Oxidation]$_{EL}$ of a component
(network or non-network) yields [Oxidation]$_{CC}$---average of [Oxidation]$_{CC}$ within a cell yields the
[Oxidation]$_{CELL}$. Visually distinct mitochondrial components were chosen to validate the
automated mito-SinComp intensity measurements by their strong linear correlation with values
manually obtained with standard image analyses software **(Fig. 2C, left).** Weaker correlation of
[Oxidation]$_{CELL}$ values further indicates heterogeneity of intracellular mitochondrial signal (**Fig.**
S1G). Nonetheless, the refined mito-SinComp [Oxidation]$_{CELL}$ values reveal statistically
significant differences that are not the evident in manual analyses **(Fig. S1H).**
We demonstrated proof of principle of mito-SinComp S(network)-F(oxidation) analyses
on a physiologically relevant heterogenous data set generated using multiple doses of an
oxidative carcinogen, TCDD, to create a wider spectrum of [Oxidation] within non-transformed
and transformed cells (control and carcinogen exposed). Mito-SinComp analyses of this pooled
cell population of 106 cells from 6 experimental and 4 control groups, revealed that the average
length of the networks in each cell ~20 fold higher than that of the non-networks, and the
average number of networks in each cell is ~4 folds less than those non-networks **(Fig. 2D)**.
Comparison of mito-SinComp S-F features shows that the networks are heterogenous in their
[Size]$_{CC}$ and non-networks are heterogenous in [Thickness]$_{CC}$ and [Oxidation]$_{CC}$ **(Fig. 2E)**.
Individual group analyses reveal bimodal [Oxidation]$_{CC}$ distribution in some and significant
differences in [Oxidation] $_{CC}$ between networks and non-networks in others **(Fig. S1I)**.
Thus, we developed and validated mito-SinComp analyses and extracted the mito-
SinComp quantitative S(network) and F(redox) features from networks and non-networks of live
cells. Using mito-SinComp on genetically or physiologically manipulated cell populations, we
demonstrated the proof of principal of the approach in untangling of mitochondrial structure-
function between networks and non-networks.
228
Mitochondrial redox function and its heterogeneity can be predicted from mitochondrial
network structure using mito-SinComp S-F features in machine learning algorithm
Of various possible analyses, we investigated the quantitative relationships between the
single mitochondrial S-F features in steady state in the heterogenous pooled HaCaT
populations as well as within individual cells. No obvious linear relationship is detected in the
bivariate S-F analyses between [Oxidation]$_{CC}$ and individual structural features of networked

and non-networked mitochondrial components in the pooled data set (**Fig. 2F**). Notably, a

complete range of [Oxidation]$_{CC}$ and its intra-network heterogeneity is observed in the

mitochondrial components in the median range of distribution of each structural feature **(Fig. 2F,**

Fig. S2A). For analyses of intra-cellular heterogeneity, we identified a common cell that

harbours mitochondrial components with S-F features in their respective median bins **(Fig. 2G)**.

These mitochondrial components of this cell exhibit a variety of complex organization of

mitochondrial elements within connected networks **(Fig. S2B).** Notably, the 3D S(structure)-

F(redox) map of the cell reveals dramatic intra-network heterogeneity of mitochondrial redox

levels between their constituent elements, quantified as INH-[Oxidation]$_{CC}$ **(**noted in networks

outlined in **Fig. 2G,** and quantified in inset, formula in Suppl Table 1**)**. Interestingly, an S-F plot

of [Size]$_{CC}$ and [Oxidation]$_{CC}$ and [Oxidation]$_{EL}$ of their constituent elements reveal that the range

of [Oxidation]$_{EL}$ values are similar between networks and non-networks; average of these

values yield [Oxidation]$_{CC}$ of the respective network, which when averaged with the non-network

yields [Oxidation]$_{CELL}$ **(Fig. 2H).** Such intra-network redox heterogeneity may arise from

maintenance of mitochondrial matrix discontinuity at the nodes, or by other undiscovered

mechanisms.

Next, we investigated if the [Oxidation]$_{CC}$ and INH-[Oxidation]$_{CC}$ can be predicted from

multiparametric analyses of their structural features. We used the Machine Learning (ML)

method of Random Forest, to perform regression task towards predicting [Oxidation] levels

considering the mitochondrial elements as the lowest structural unit; the high [Oxidation]$_{CC}$

values consisting of 0.016 % of the total data set were filtered out as outliers. The training was

performed with hyperparameter tuning in 70% of the data and tested in the remaining 30% of

the data, with 5-fold cross validation (see Methods). The prediction was evaluated using R^2

metric and prediction error was measured using Mean Absolute Error (MAE) metric.

Interestingly, [Oxidation]$_{CC}$ levels can be robustly predicted for the mitochondrial networks

(cross validated R^2 of test scores = 0.957) but not for the mitochondrial non-networks, both using

the pooled or individual experimental groups **(Fig. 2I,J)**. The INH-[Oxidation]$_{CC}$ for the networks

is also predictable to the same degree **(**cross validated R^2 of test scores = 0.937**)**. Comparison

of MAE and [Oxidation]$_{CC}$ distribution in the test set revealed that the prediction error increases

with reduction in data abundance, with overall higher error for the non-networks **(Fig.2K)**. Given,

the non-network data set is ~6 fold smaller than that of the networks (due to a singular element),

we tested the performance of the model after randomized reduction of the network data set size

to that of the non-networks. This reduced the model performance for prediction of [Oxidation]$_{CC}$

levels (R^2 = 0.73), but not to the non-predictable level of the non-networks ($R^2 \leq 0.1$), **(Fig. 2L).**

This demonstrates that the combinatorial structural information in the networked mitochondrial

components leads to higher predictability of their redox and its intra-network heterogeneity.

Thus, hierarchical mito-SinComp analyses reveal remarkable intra-cellular and intra-

network heterogeneity of mitochondrial redox, which can be predicted only for networks by ML

based modelling of combinatorial network structural features. Such predictability can be tested

for any mitochondrial function and can be potentially utilized in translational settings.

Use of mito-SinComp structural feature, CRL-mitoNet, reveals conversion of Hyperfused-

Mitochondrial-Networks to Small-Mitochondrial-Networks upon stem cell activation

We tested if untangling mitochondrial heterogeneity can reconcile controversies on

fragmented mitochondrial entities in stem cells[21,24,25]. We used mito-SinComp to identify the

characteristic mitochondrial form of stem cells in benchmarked HaCaT keratinocytes

populations where genetic manipulation of Drp1 enriched or depleted Krt15[hi] stem cells [23], and

in flow sorted Aldh[hi] ovarian cancer tumor initiating (stem) cells where Drp1 modulates Aldh

expression[21,42].

To capture both inter- and intracellular mitochondrial heterogeneity in a single feature,

we computed the 'Cell-Relative Length of mitochondrial network or non-network', [CRL mito-

Net] or [CRL mito-nonNet] (**Suppl Table 1**). Comparison of distribution of [CRL mito-Net] in the

INT-cells and the HF-cells groups of the Drp1 manipulated keratinocytes, indicate that the INT-

cells may be enriched with 'Small Mitochondrial Networks' (**SMNs**), whereas the HF-cells with

'Hyperfused Mitochondrial Networks' (**HMNs**) and 'very Small Mitochondrial Networks'

(**vSMNs**)(the smallest network unit with 4 nodes) (**Fig. 3A, Suppl Table 2**). Indeed,

classification based on [CRL mito-Net] thresholds (see Methods) confirmed remarkable SMN

enrichment in the INT-cells and the enrichment of the HMNs and vSMNs in the HF-cells (**Fig.**

3B); examples of networks and non-networks (which may include the mitochondria derived

vesicles [43]) are shown (**Fig. 3C**). Mito-SinComp (S) features reveal that in comparison to the

HMNs, the SMNs are (**Fig. 3D, Fig. S3A**): **a)** ~10 fold smaller in size and in abundance of nodes

and edges. **b)** comparable in element size within a network; **c)** remarkably heterogenous in

thickness, network complexity (nodes or edge density) and maximum connectivity.

KRT15[hi] stem cells are enriched by the weak (W) Drp1 knockdown that sustains INT-

cells, and depleted due to the strong (S) Drp1 knockdown (kd) that sustains HF-cells [23].

Therefore, we used this benchmark data set and its corresponding scRNA-seq data to

investigate any link of the SMN enrichment to the transcriptomic network of the lineage specific

stemness gene KRT15 [44]. The hdWGCNA **(Fig. S3B-C)** of the combined differentially

expressed gene sets from pseudobulk analyses of NT vs Drp1-kd(W) and NT vs Drp1-kd (S)

identified the 10 genes that directly interact with KRT15 in its transcriptomic network **(Fig. S3D-**

F). HdWGCNA of the marker genes of the KRT15^{hi} stem cell cluster, confirmed 4 out of the

above 10 genes as direct KRT15 interacting partners **(Fig. 3E** with kME bar plot and edge

distance table as insets, **Fig. S3E-F)**. Importantly, these KRT15 interacting genes undergo

marked upregulation across all cell clusters in the KRT15^{hi} stem cell enriched Drp1-kd (W)

population with abundant SMNs in the INT subpopulation **(Fig. 3F, top, Fig. S3G)**. On the other

hand, two of those genes undergo marked downregulation only in the KRT15^{hi} cells that are

depleted in the Drp1-kd (S) population with depletion of SMNs in the HF-cells **(Fig. 3F,**

bottom). Thus, parallel mito-SinComp and hdWGCNA analyses indicate that the SMNs are

linked to KRT15-stemness transcriptomic network in KRT15^{hi} stem cells.

To address if activating stemness can enrich SMNs we performed mito-SinComp

structural analyses in sorted tumor initiating (stem) cells (TICs) and bulk tumor cells (BTCs)

from A2780-CP lines. Self-renewal was activated in the sorted Aldh^{hi} TICs and Aldh^{lo} BTCs with

TIC media (24 hrs), using RPMI control media [21]. The mito-SinComp 3D maps and structural

analyses reveal that the TIC media (with growth factors) reduces the average length of

networked mitochondria while increasing their abundance in both Aldh^{hi} TICs and Aldh^{lo} BTCs

(Fig. S4A-C). TIC media causes a pronounced increase in [CRL mito-Net] in both Aldh^{hi} TICs

and Aldh^{lo} BTCs **(Fig. S4D)**, and induces remarkable enrichment of SMNs and a concomitant

depletion of the HMNs in them **(Fig. 3G).** More importantly, SMN / HMN ratio indicates that

enrichment of SMNs by TIC media is 2 folds higher in the Aldh^{hi} TICs than in the Aldh^{lo} BTCs

(Fig. 3H). These SMNs remain ~10X smaller that of the HMNs and remarkably heterogenous

(Fig. 3I, Fig. S4E, similar to **Fig. 3D)**, Notably, the TIC media dramatically reduces the network

size and abundance of nodes and edges of the HMNs almost to the level of the SMNs, with no

impact on individual element size **(Fig. 3I)**. TIC medium increases thickness in both SMNs and

HMNs, while reducing maximum nodal connectivity in the SMNs **(Fig. S4E-F)**. Together, these

data demonstrate that TIC media converts all the HMNs to SMNs by severing specific nodal

connections of the HMNs and not by reducing the element size.

Next, we examined the heterogenous SMNs and the HMNs in the pooled data set

where mito-SinComp S-F analyses revealed profound intra-network heterogeneity of

[Oxidation]$_{CC}$ (**Fig.2**). Trivariate S-F plot of INH- [Oxidation]$_{CC}$, [Size]$_{CC}$ and [Edge Density] reveal

that the SMNs maintain similar intra-network heterogeneity as the HMNs while higher network

complexity is exhibited in some SMNs with lower intra-network heterogeneity (**Fig. 3J**).

Nonetheless, the [CRL mito-Net], which distinguishes SMNs from the HMNs, accounts for ~30%

of the predictability for redox in the ML modelling, while combination of various structural

features predicts intra-network heterogeneity (**Fig. 3K**, **Fig. 2**); expectedly, the invariant

[Element Size] accounts for only ~1% of both predictability**.**

In summary, we demonstrate that the quantitative feature, [CRL mito-Net], untangles the

SMNs from the HMNs and contributes substantially in prediction of network redox status.

Mechanistic analyses with mito-SinComp demonstrate that activation of stemness converts the

HMNs to ~10 folds smaller heterogenous SMNs by severing specific HMN nodal connections

(**Fig. 3L**). Such precise regulation happens more pronouncedly in the TIC enriched population in

the early during self-renewal induction, linking SMN formation to stemness specification.

Mito-SinComp S-F analyses reveals prompt conversion of oxidized HMNs to redox-tuned

SMNs of restricted complexity in non-transformed cells upon exposure to stem cell

enriching carcinogen

We used mito-SinComp S-F analyses to directly address if any functionally relevant

SMN enrichment is triggered in non-neoplastic HaCaT keratinocytes at the initial stage of stem

cell driven transformation. Thus, we studied the acute impact of reported stem cell enriching

TCDD-1nM, with control TCDD-10nM carcinogenic dose (with reduced stem cell enrichment)

and the Toluene vehicle control [23]. Interestingly, TCDD-1nM promptly depolarizes mitochondria

within one hour (T1-1h) when TCDD-10nM needs 4 hours to cause the same effect **(Fig. 4A,**

Fig. S5A). Mitochondrial depolarization alters their shape [45] and redox function [46]. Therefore, we

used mito-SinComp S(network)-F(redox) analyses to test if stem cell enriching T1-1h induced

mitochondrial depolarization is linked to formation of SMNs with specific redox functions using

the Toluene, T1-1h and T10-1h groups from the experimental pool of **Fig. 2**.

The 3D S(structure)-F(redox) maps of representative cells indicate lack of large

mitochondrial networks particularly after T1-1h exposure, similar to the TIC media, and elevated

[Oxidation]$_{CELL}$ **(Fig. 4B, S5A-B)**. This is quantitatively reflected in reduced average length of

both networked and non-networked mitochondria likely due to the overall increased contribution

of mitochondrial fission, which is modest with T10-1h exposure (**Fig. 4C, S5C-F)**. [CRL mito-

Net] based classification and analyses of various structural features confirms that the stem cell

enriching T1-1h dose promptly converts all the HMNs to 10X smaller and heterogenous SMNs

by severing HMN nodal connections, while T10-1h exposure exhibits less specificity of SMN

enrichment **(Fig. 4D-F, Fig. S5G)**. Although this is similar to stem cell activation with TIC

medium **(Fig. 3G-I)**, a notable difference is that T1-1h exposure elevates network complexity

(node or edge density) **(Fig. 4F)**.

The pronounced specific impact of T1-1h on mitochondrial potential **(Fig. 4A)** and on the
HMN structure **(Fig. 4F)** associates with significant increase in [Oxidation]$_{CC}$ only in the HMNs
that are converting to SMNs, which remain undetected in the uncategorized networked
mitochondria **(Fig. 4G-H)**. On the other hand, the [Oxidation]$_{CC}$ is bimodally distributed in the
SMNs of the Tol-1h and T1-1h groups and reduced to a unimodal distribution in the T10-1h
group (**Fig. 4I**). To tease out such heterogeneity of SMN oxidation, we studied the structural
parameters in [Ox]$_{CC}$ -Hi / Int / Lo zones **(Fig. 4I,** zone demarcations are defined in the legend**).**
T1-1h exposure reduces SMN size and elevates their network complexity prominently in the
[Ox]$_{CC}$ -Lo zones (**Fig. S5H**). To probe further, we performed bivariate S-F analyses between
[Oxidation] and individual structural parameters of the SMNs and quantified their abundance in
the defined [Oxidation]$_{CC}$ zones. The [Ox]$_{CC}$ -Int zone is overall enriched (~1.7 X) with T1-1h
induced SMNs and [Ox]$_{CC}$-Lo zone is enriched (~2X) with T10-1h induced SMNs, with respect to
the Tol-1h group **(Fig. 4J)**. Interestingly, the bivariate S-F plot between [Node or Edge density]
and [Oxidation]$_{CC}$ appears to be unique from other structural parameters **(Fig. S5I)**. Particularly,
the SMNs are primarily restricted within a lower level of [Node or Edge density] and maximal
nodal connectivity only in the [Ox]$_{CC}$ -Int zone **(Fig. 4J, Fig. S5I)**.
Thus, mito-SinComp S-F analyses demonstrate that exposure of non-transformed cells
to the stem cell enriching carcinogen promptly oxidizes their HMNs and severs nodal
connections to convert them to the heterogeneous SMNs that are prevented from conversion to
the vSMNs. Importantly, a subpopulation of SMNs remain tuned at an intermediate redox level
and with restricted network complexity. Such unprecedented untangling of mitochondrial
structure-function heterogeneity can be achieved by mito-SinComp in any cell physiology.
393
Carcinogen induced enrichment of redox-tuned SMNs specifies stemness by
establishing mtDNA-stemness transcriptomic network, revealed by coupled scRNA-seq
We employed scRNA-seq to investigate the immediate cellular impact of T1-1h induced
enrichment of redox-tuned SMNs elucidated by mito-SinComp. Therefore, we performed
scRNA-seq of HaCaT cells following SMN enrichment i.e. after 2.5 hrs exposure to Tol (vehicle
control), T1 (stem cell enriching dose) and T10 (control dose)**.**
Pairwise pseudo-bulk scRNA-seq analyses confirmed upregulation of the canonical
TCDD target gene, *CYP1B1* [47] in T10-2.5h but not in T1-2.5h (not shown). Surprisingly, various
protein coding genes encoded by the mtDNA heavy strand are upregulated presumably in the
SMNs that are the predominant mitochondrial form maintained by T1-2.5h exposure **(Fig. 5A)**;
the *MT-ND6* gene product coded by the mtDNA light strand is downregulated only by T10-2.5h.

This distinction between the mt-DNA heavy and light stand suggests a distinct transcriptional

regulation from the distinct promoters at each stand[48]. In contrast, the nuclear genes encoding

mitochondrial proteins are overall downregulated by both T1-2.5h and T10-2.5h **(Fig. 5A)**.

Immunoblot analyses validated the contrasting effects of T1-2.5h and T10-2.5h on mt-DNA

coded MT-CYB levels, with no impact on nuclear DNA coded NDUFB3 **(Fig. 5B)**. GSEA based

functional annotation of genome wide pairwise pseudo-bulk analyses demonstrate that both T1-

2.5h or T10-2.5h downregulate genes involved in mitochondrial energetics and cell cycle **(Fig.**

S6A). Furthermore, hdWGCNA demonstrates that the modules obtained with (T1 vs Tol)-2.5h

gene sets have higher number of mito-Energetics and CellCycle genes, and their transcriptomic

connectivity is strikingly complex compared to the similar module identified with (T10 vs Tol)-

2.5h gene sets (**Fig. S6B-D)**.

Next, we identified the cell clusters on a UMAP plot, quantified their abundance **(Fig.5C,**

S6E), and assigned cluster functionality by GSEA on their respective gene markers **(Fig. S6F)**.

Both T1-2.5h or T10-2.5h reduces abundance of 'Proliferating' cells (upregulated cell cycle

makers) and increases abundance of the 'CC-Exiting' cells (downregulated cell cycle makers)

(**Fig.5C)**. We re-designated the latter as 'CC-Exiting-K15^{hi}' due to highest contribution of the

stem cell gene KRT15 (among 18 KRT genes) in defining this cluster **(Fig. 5D, Fig.S7A)**.

Noteworthily, the specific T1-2.5h driven upregulation of mtDNA heavy strand genes happens

across all cell clusters and most prominently for the components of the major site of

mitochondrial ROS production namely Complex I [46] **(**marked by * in **Fig. 5E**, outlined in **Fig.**

S7B). This suggests a direct impact of T1-2.5h on mtDNA gene expression in the SMNs**,**

although their lower expression in the 'CC-Exiting-K15^{hi}' cluster is consistent with their

reciprocal relationship with KRT15 in the transformed population [23]. Interestingly, expression of

Drp1(coded by DNM1L gene), among key genes modulating mitochondrial fission and fusion, is

differentially modulated between T1-2.5h and T10-2.5h **(**arrows **in Fig. 5E** inset**),** consistent

with Drp1 knockdown modulating stem cell status in this lineage [23]. Next, we used hdWGCNA to

identify any mtDNA-KRT15 (stemness) interacting modules within each cell cluster and further

assigned functionality to the listed modules using GSEA. Importantly, the mtDNA genes are

found to interact with the stem-cell gene KRT15 only in the Proliferating cluster of the T1-2.5h

group (kME bar plot as inset in **Fig. 5F, Fig. S7C-D**). Hub-gene network analyses of the

mtDNA-KRT15(stemness) module reveal a distinct network of interacting mtDNA genes, out of

which redox related MT-ND1 of Complex I is directly connected to KRT15 that is further

connected to the major redox regulator GPX2 **(Fig. 5F)**.

To confirm the above findings in neoplastic stem cells, we performed parallel mito-

SinComp and hdWGCNA on TCDD-1nM transformed experimental group with pronounced stem

cell enrichment (TF1), TCDD-10 nM transformed control group with substantially weaker stem

cell enrichment (TF10) and non-transformed parental control group (P) [23].   [CRL mito-Net]

based network classification demonstrates that SMN and HMN abundance is distinctly different

between the TF1 and the control groups **(Fig. S7E-F)**. The percentage of cells above or below a

given threshold of SMN / HMN ratio in each cell signifies SMN or HMN enrichment, respectively.

Indeed, SMN enrichment remains consistently higher in the TF1 population between a threshold

of 4 and 6 (**Fig. 5G, Fig. S7G**), while SMNs and HMNs maintain their distinctive properties (**Fig.**

S7H). Moreover, trivariate mito-SinComp S-F analyses demonstrated that the SMNs that exhibit

maximum heterogeneity of $[Edge Density]_{CC}$ and remarkably low $INH-[Oxidation]_{CC}$ belong to the

stem cell enriched TF1 population (outlined in **Fig. 5H**).

Probing into the mt-DNA gene expression confirmed specific upregulation of redox

related MT-ND1 gene for Complex I and MT-CYB gene for Complex III particularly in TF1 (**Fig.**

5I, Fig. S7I). Also, specifically TF1 group exhibits prominent downregulation of the mt-DNA

regulating molecule Prohibitin particularly in the previously labelled KRT15^{hi} stem cell cluster

(**Fig. S7I**, empty triangle, **Fig. S7J**)[49]. Next, we confirmed the SMN related distinction of mtDNA

genes in the available scRNA-seq data from the relevant cutaneous squamous cell carcinoma

(cSCC) patient samples [50]. Interestingly, 1 out of 9 patients exhibits prominent increase in

mtDNA genes (including MT-ND1 and MT-CYB) in the cSCC tumor from the adjacent normal

tissues, similar to the SMN and stem cell enriched TF1 cells **(Fig. 5J, outlined)**. Also, this

patient exhibits no alteration in keratin gene expression in contrast to the other patients showing

decrease in mtDNA genes (similar to TF10) and keratin genes in the tumor. Expectedly, the

nuclear encoded gene for the mitochondrial protein NDUFB3 shows no change across all the 10

patients. Finally, hdWGCNA confirmed the direct interaction of KRT15 with MT-ND1 (and MT-

CB5) in a module detected particularly in the S-Late cluster of the TF1 (**Fig. 5K** with kME plots

as inset**, Fig. S7K)**; although the Cytokine clusters of the TF1 and TF10 populations harbor a

similar module, no direct MT-ND1-KRT15 connection was detected.

In summary, coupled hdWCGNA demonstrated that enrichment of the redox-tuned

SMNs by the stem cell enriching carcinogen is promptly followed by upregulation of their mtDNA

heavy strand genes and establishment of the MT-ND1(redox)-KRT15(stemness) transcriptomic

network in the proliferating non-transformed cells. Given the same features are detected only in

stem cell enriched transformed population, we conclude that SMN induced transcriptomic

modulation prime stemness driven neoplasticity in non-transformed cells. Identification of cSCC

patient with SMN linked mtDNA changes indicate the clinical relevance of our finding while

highlighting patient heterogeneity.

Mito-SinComp analyses of mtDNA nucleoid organization in fixed samples demonstrates

higher nucleoid abundance in the characteristic stemness specifying SMNs.

Efforts are underway to quantitatively resolve the remarkable heterogeneity of the

polyploid mtDNA that is organized in nucleoids of variable size and number within mitochondrial

networks [51]. Using mito-SinComp modular design **(Fig. 1),** we investigated if the mechanism of

increase in mtDNA gene expression with conversion of HMNs to SMNs involves functional

organization of their mtDNA. We performed in-depth mito-SinComp S-F analyses in HaCaT

keratinocyte and beyond cell boundaries in a *Drosophila* tissue amenable to confocal

microscopy, using nucleoid number and size as functional surrogates.

The individual mtDNA nucleoids are identified from the high-resolution 3D confocal

micrographs as pixel clusters using DBSCAN algorithm after appropriate optimization of its

hyperparameters (see Methods) (**Fig. 6A**). The identified clusters are mapped to the

corresponding mitochondrial structural coordinates and mitochondrial components, where each

component can harbor multiple mtDNA nucleoids of varying size **(**arrows in **Fig. 6A)**. The

visually confirmed unmapped clusters **(**arrowheads in **Fig. 6A)** may represent released mtDNA

nucleoids [52] or other unwanted signal. Fixed mito-SinComp analyses confirm the higher

heterogeneity in [Thickness]$_{CC}$ of the non-networked components **(Fig. 6B)**, as in live cells (**Fig.**

2E). However, the absolute [Size]$_{CC}$ and [Overall Nodal Connectivity] are reduced in fixed cells

(**Fig. S8A,B**). We computed the following mtDNA nucleoid functional features for individual

mitochondrial components: [Nucleoid Number]$_{CC}$, [Nucleoid Density]$_{CC}$, [Nucleoid Size]$_{EL}$, [Avg-

Nucleoid Size]$_{CC}$ and INH-[Nucleoid Size]$_{CC}$ (**Suppl. Table 1**). The networked components have

higher [mtDNA nucleoid Number]$_{CC}$ but reduced [Nucleoid Density]$_{CC}$ in comparison to the non-

networked components, while [Nucleoid Size]$_{EL}$ remains comparable **(Fig 6B)**.

Interestingly, S-F plot of [Size]$_{CC}$ vs [Nucleoid Number]$_{CC}$ reveal that while longer non-

networked mitochondria have upto 5 mtDNA nucleoid in this cell, the [Nucleoid Number]$_{CC}$

exponentially increases after formation of mitochondrial network of a critical window of [Size]$_{CC}$

(Fig. 6C, dashed lines**)**. For any given [Nucleoid Number]$_{CC}$, the [Nucleoid Density]$_{CC}$ is

expected to reduce with [Size]$_{CC}$. Deviation from this expectation indicates influence of other

factors, as is evident in certain networked components in the critical window **(**highlighted in **Fig.**

6C). These networks are identified as an SMN subpopulation by quantitatively comparing similar

S-F plots of [Node Density]$_{CC}$ vs [Size]$_{CC}$ in the fixed cell with that of live cells where SMNs were

defined (**Fig. S8B**). Other S-F plots confirm that the higher nucleoid density in these SMN

subpopulation is due to their higher nucleoid number. Importantly, their restricted network

complexity indicate they may represent the redox-tuned SMNs subpopulation with restricted

network complexity (**Fig. 4J**). Like other networks, these SMNs maintain intra-network

heterogeneity of nucleoid number and size between their elements (arrowheads in **Fig. 6C** for

the example SMN in **Fig. 6D**). Visual examinations reveal the positioning of nucleoids at the

nodes of these SMNs (arrows in **Fig. 6D, top**). A trivariate plot of [Nucleoid Density]$_{CC}$, [Node

Density]$_{CC}$ and [Size]$_{CC}$ distinguishes these SMN from the VSMs which exhibit maximized

[Nucleoid Density] and [Node Density] due to their small size (**Fig. 6D, bottom**). Mito-SinComp

analyses in *Drosophila* fixed tissue beyond cell boundaries recapitulates the key distinction of

mtDNA organization between networks and non-networks (**Fig. 6E-G**), highlighting fundamental

importance of our findings. Notably, the S-F plot of [Size]$_{CC}$ and [Nucleoid Density] identifies the

potential SMN subpopulation which when enriched by neoplastic insults may specify stemness

in the cells carrying these SMNs, thus priming neoplasticity (outlined in **Fig.6G**).

In summary, mito-SinComp analyses of mtDNA-nucleoids revealed their distinctive

organization in the networks and non-networks, and also their positioning at the internal and

free nodes. Comparative mito-SinComp analyses between fixed cells and in live cells, in

conjunction with scRNA-seq, reveals a novel mechanistic model of stem cell specification (**Fig.

7**). Stemness specification involves enrichment of a subpopulation of redox-tuned SMNs with

restricted network complexity and greater mtDNA abundance, generated by precise breakdown

of oxidized HMNs. This supports the elevated mtDNA gene expression in these SMNs, which in

turn establishes the transcriptomic link between mtDNA coded mt-ND1 (redox relevant) gene

and nuclear DNA coded Krt15 (stemness gene).

Discussion

Quantitative understanding of mitochondrial heterogeneity is required to elucidate the

complex role of mitochondria in normal and pathophysiological conditions[1,5,53–55]. Understanding

heterogeneity requires quantitative analyses of single entities, which is achieved by Mito-

SinComp structure(S)-function(F) analyses (**Fig. 1**). Mito-SinComp S and F analyses focused

on redox function revealed redox levels and its remarkable intra-network redox heterogeneity

can be quantitively predicted by a combination of the structural features through a machine-

learning approach (**Fig. 2,3**). Mito-SinComp analyses suggest that a characteristic mitochondrial

form, SMNs with specific redox and mtDNA properties, specifies stemness. Our conclusions are

based on mito-SinComp data from 17 experimental groups, involving 772 cells, 12181 networks,

24556 non-networks, 848 HMNs and 5872 SMNs, across 7 cell lines (parental and derived by

genetic manipulation or carcinogen exposure). Moreover, using publicly available data [50], we

demonstrate inter-cancer patient heterogeneity for SMN linked mtDNA modulation, (**Fig. 5J**).

Furthermore, using *Drosophila* we demonstrate that the characteristic SMNs can be identified

using mito-SinComp in normal fixed tissues that are amenable for confocal microscopy (**Fig. 6E-**

G). Molecular surrogates for SMNs remain to be discovered, which can be used in tissues that

are not-amenable for microscopy. Nonetheless, our findings highlight that stem cell functionality

of mitochondria may be exhibited not by fully fragmented mitochondria but rather by

characteristic SMNs which are presumably generated by precisely regulated fission events and

may be maintained by precisely regulated fusion events. This explains need of both fission and

fusion processes in stemness, thus reconciling various contradictory results [3,21,23–25,31–34] .

The unprecedented abilities of mito-SinComp analyses identified and characterized the

subpopulation of stemness specifying SMNs, as well as revealed mechanism of their formation

and function in stemness specification **(Fig. 7)**. The stemness specifying SMNs are 10 times

smaller than HMNs, are redox tuned, have restricted network complexity and elevated mtDNA

nucleoid density. This supports elevated mtDNA gene expression and the resulting transcript of

the redox relevant mt-ND1 gene maintains direct transcriptomic connection with lineage specific

stemness gene, as revealed by coupled hdWGCNA of scRNA-seq data. The redox-tuned

SMNs are formed by severing of the nodes of the oxidized HMNs, likely mitochondrial fission

(**Fig. S5F**). We speculate that inhibition of HMNs to SMNs conversion by obliteration of

mitochondrial fission underlies its marked inhibition of oncogene induced neoplastic

transformation [23,35,36], which can also be tested in other models of neoplastic transformation.

The SMNs need to be sustained further by keeping mitochondrial fission in check, which

explains the fine-tuned regulation of Drp1 in promoting neoplasticity [23]. Mitochondria act as a

hub of redox signaling, while stem cells are particularly dependent on redox tuning [2,40]. SMN

formation could be a potential spatio-temporally controlled mechanism of fine-tuning redox

signaling through mitohormesis [56]. This conception adds a novel direction in elucidation of the

bidirectional crosstalk between mitochondrial structure and redox properties [41,57,58]. hdWGNCA

detected mt-ND1(redox)-KRT15(stemness) transcriptomic network in stemness primed

proliferating cells and in late S phase after transformation (**Fig. 5**). Therefore, the stemness

specification by redox driven SMNs might involve cell cycle regulation, which plays a

deterministic role in stemness [40,59].

We demonstrated the following capabilities of the quantitative mito-SinComp approach,

that can be used for any function of mitochondria for any cell physiology in live and fixed cells

and tissues: **A)** distinguishing mitochondrial networks from non-networks, and their properties;

B) distinguishing component types using quantitative [CRL mito-Net / non-net] feature, and

subpopulation of types based on their function; **C)** understanding query driven quantitative S-S,

S-F or F-F relationships (bivariate or multivariate) within identified types and subtypes; **D)**

quantitative elucidation of intra-network heterogeneity of function; **E)** comparison of quantitative

features between live and fixed samples; **F)** usefulness of S-F features in machine-learning

based predictive modeling for translational research; **G)** distinguishing impact of agents on

specific kinds of networks or non-networks; **H)** informing about mechanism of interconversion

between network forms**; I)** elucidating impact of identified components on cell physiology using

correlatively or causatively with other quantitative modalities. To further elucidate the molecular

mechanism of our novel findings, mito-SinComp can be used in a time-lapse mode to study

activation of Drp1 driven fission sites at the HMN nodes, or to identify players involved in

maintaining the marked intra-network redox heterogeneity, or to link functional mtDNA

organization to their replication, transcription, inheritance etc. Using mito-SinComp on primary

cells from patients will enable studying inter-patient and intra-tumor heterogeneity of SMNs

towards personalized treatment [29,60]. Mito-SinComp can be used to study the impact of genetic

mutations, recruitment and release of mitochondrial regulators and moonlighting proteins.

In conclusion, we untangled multilevel mitochondrial heterogeneity by directly studying

structure-function of single mitochondrial components in the context of stemness using a first of

its kind quantitative approach, mito-SinComp. The unprecedented capabilities of mito-SinComp

can be used in a variety of ways to make fundamental discoveries and support their

translational applications.

Materials and Methods

Cell lines:

We have used the following parental and derived cell lines across 2 cell lineages: **a)** the

parental HaCaT line (immortalized normal skin keratinocytes) and their derived lines

transformed with 1nM or 10nM TCDD (**Fig. 5**); **b)** parental HaCaT lines stably expressing weak

or strong Drp1 shRNA or non-targeted (NT) shRNA (**Fig. 2,3**); **c)** parental double stable HaCAT

line expressing mito-PSmO2 and mito-roGFP and their derived lines transformed with 1nM or

10nM TCDD or 10nM Toluene (**Fig. 2, 4); d)** A2780-CP line (carboplatin resistant ovarian

cancer line derived from parental A2780 line) (**Fig. 3**). All HaCaT cells were maintained in

Dulbecco's Modified Eagle's Medium (DMEM)- high glucose supplemented with 10% FBS and

Penicillin-Streptomycin cocktail (100 µg/mL), in a 5% CO_2 at 37deg C using standard

techniques. A2780-CP cells were similarly maintained in RPMI medium supplemented with 1

mM sodium pyruvate.

The double stable HaCaT lines expressing mito-PSmO2 and mito-roGFP were

generated using the pCDH lentiviral expression system with puromycin and hygromycin

selection, respectively, using standard methods. Transformed HaCaT lines were derived from

the Parental HaCaT cells by selecting multiple clones following treatment with 1nM or 10nM

dose of TCDD for 16 to 20 days, with media replenishment every 2–3 days. The HaCaT cells

with stable knockdown of Drp1 (weak or strong) were generated using Lipofectamine mediated

transfection of human DNM1L shRNA1 and shRNA2 with non-targeted control, and were

selected using puromycin. All stable genetically modified or transformed cell lines, along with

their respective parental controls, were subjected to two consecutive rounds of BM-Cyclin

treatment to ensure the removal of any potential mycoplasma contamination. For experiments

involving acute exposure of TCDD, the parental or double stable HaCaT cells were exposed to

the 1 nM or 10nM TCDD for 1 or 4 hours.

Cell Sorting using flow cytometry

A2780-CP cells were sorted using flow cytometry based on their ALDH (aldehyde

dehydrogenase) activity detected by ALDEFLUOR staining carried out as per the

manufacturer's instructions (Stem Cell Technologies), with minor modifications as required.

Stained cells were analyzed using a 488 nm excitation laser with a 525/50 band-pass filter on

the LSR II flow cytometer, and a 530/30 band-pass filter on the FACS Aria II, both in

combination with a 505 nm long-pass filter. The Aldh activity based cell sorting was performed

using BD FACS AriaII. The sorted cells were collected in complete RPMI medium or TIC

medium composed of RPMI supplemented with 1× N1 supplement, 500 mg/ml insulin, 20 ng/ml

human epidermal growth factor (EGF), and 10 ng/ml basic fibroblast growth factor

(bFGF). Following sorting, cells were plated in Geltrex coated LabTek chambers and stained

with Mitotracker 633 after 24 hours.

MitoTracker and TMRE Staining

Cells were incubated in appropriate medium containing MitoTracker Green FM (50–100 nM) or

MitoTracker Red FM 633 (250 nM) for 15 minutes at 37 °C. For assessment of mitochondrial

membrane potential, TMRE was used at a final concentration of 50 nM under the same

incubation conditions. To assess mitochondrial membrane potential relative to mitochondrial

mass, cells were first stained with TMRE and then mitotracker green, while TCDD/Toluene

doses were maintained in the in the staining and washing media. Stained cells were promptly

imaged live on confocal microscope using appropriate acquisition settings, as described below.

Immunoblotting and Immunofluorescence

For immunoblotting, whole-cell lysates were prepared using Laemmli buffer with reducing agent.

Proteins (cell equivalent) were resolved on 10% SDS–polyacrylamide gels and transferred onto

PVDF membranes. After blocking with 5% bovine serum albumin (BSA), membranes were

incubated with specific primary and then compatible HRP-conjugated secondary antibodies,

followed by detection using chemiluminescence, using standard protocols. Densitometric

analysis of protein bands was performed using ImageJ, with normalization by loading and

experimental controls as appropriate.

For immunofluorescence, cells were cultured on Nunc Lab-Tek chamber slides and

processed using standard protocols. Cells were fixed in freshly prepared 4% paraformaldehyde

containing 4% (w/v) sucrose, then permeabilized using 0.1% Triton X-100. After

permeabilization, cells were blocked in 1% BSA, then stained with specific primary and

secondary antibodies. Finally, samples were mounted using Fluoromount-G containing Hoechst

33342 (10 µg/mL) for nuclear counterstaining. Slides were imaged using a confocal microscope

as described below. Drosophila fat body tissues were dissected in a glass well plate and

immediately fixed in freshly prepared 4% paraformaldehyde. Following fixation, tissues were

permeabilized using 0.1% PBS-Triton X-100 (PBS-TX) and subsequently blocked with an

appropriate concentration of 10% BSA dissolved in 0.1% PBS-TX. The samples were then

incubated with specific primary and compatible secondary antibodies for immunostaining. After

antibody incubation, tissues were washed three times with PBS-TX. During the final wash,

Hoechst 33342 (1:1,000 dilution) was added to stain nuclear DNA. Finally, the stained fat body

tissues were carefully transferred onto a drop of mounting medium on a glass slide. A coverslip

was gently placed over the tissues and lightly pressed to achieve an even and uniform spread.

Confocal Microscopy

Microscopy was performed using either Zeiss LSM 700 laser scanning confocal microscope or

Zeiss LSM 900 laser scanning microscope. LSM 700 was equipped with a 40X Plan-

Apochromat 1.4 NA oil immersion objective, housed within a temperature- and CO_2-controlled

incubation chamber to maintain physiological conditions (for live cell imaging). Appropriate

excitation laser lines were used for each fluorophore (405 nm and 488 nm for mito-roGFP, 555

nm for basal mito-PSmO2, 488 nm for MitoTracker Green FM, 633 nm for MitoTracker Red FM

633 and 555 nm for TMRE). Emission was collected using appropriate detectors optimized for

each fluorophore. Multichannel acquisition protocols were configured to eliminate spectral

crosstalk and cross-excitation. Three-dimensional image stacks were acquired using optical

zoom (3X), at 1 Airy unit pinhole, and 0.5 µm Z-interval. LSM 900 was equipped with a 63X

Plan-Apochromat 1.4 NA oil immersion objective. Appropriate excitation laser lines were used

for each fluorophore (as mentioned above for LSM 700). Three-dimensional image stacks were

acquired using optical zoom (2X), at 1 Airy unit pinhole, and 0.5 µm Z-interval with 2X

averaging. For mitochondrial potential assessment, image analysis was performed on maximum

intensity projections within manually drawn Regions of interest (ROIs) around individual cells,

and with appropriate background correction. The details of the newly developed mito-SinComp

image analyses is described below.

Development of mito-SinComp approach for studying structure-function relationship of

single mitochondrial components

A. Staining / expression of probes: Modularity of the approach allows use of any

appropriate probes alone or in combination. Here, we used: **a)** Structure: Mitotracker staining

(as described above) for structural analyses; TMRE stain for mitochondrial potential was used

while maintaining TCDD in the staining solution; **b)** Structure-function: Previously generated

stable cell lines expressing validated combination of mito-PSmO2 and mito-roGFP was used for

demonstrating use of mito-SinComp in live [Oxidation] function. For analyses of functional

organization of mtDNA, cells or *Drosophila* tissue are co-immunostained samples Tom20 or

Phb2 marking the mitochondrial structure and for DNA antibody marking mtDNA nucleoids.

B. Microscopy: 3D Confocal microscopy for mito-SinComp is performed (as described

above) After the appropriate quality control steps, the resulting confocal micrographs were

passed through mito-SinComp analytical pipeline described below.

C. Mito-SinComp analyses: This uses 3D confocal micrographs with appropriate

combinations of compatible probes reporting mitochondrial structure and function separately. To

develop mito-SinComp structure-function analysis for single mitochondrial components, we

modified the open source MitoGraph code (https://github.com/vianamp/MitoGraph) to identify

single mitochondrial components and include functional analyses for the identified components

within single cells. Mito-SinComp analysis requires a Linux or MacOS system to run natively. It

can be run on Windows through the Windows Subsystem for Linux (WSL) or another Virtual

Machine application like VirtualBox. We developed mito-SinComp analysis using Ubuntu 24.04

LTS with VTK 9.0 after installing all essential libraries and packages (build-essential, cmake, git,

r-base, wget, os, sys, libvtk9-dev, libgl1-mesa-dev, freeglut3-dev, qtbase5-dev, qtdeclarative5-

dev, libqt5opengl5-dev, pandas, networkx, igraph, PIL, numpy (v2.1.1 for mito-SinComp and

v1.1.1 for Foci Detection), tifffile, matplotlib, scikit-learn). Our customized modular script for

mito-SinComp structure-function approach includes the following steps:

1.   Identifying the 'structure' and 'function' channels: The 3D confocal micrographs

for our analysis have 3 channels, excited by 405 nm (function-1), 488 nm (function-2) and 555

nm lasers (structure). MitoGraph exclusively uses only the designated structure channel to

obtain the binary image through an adaptive thresholding process[61] .Therefore, we aimed to

obtain the intensity values of the 'function' channels in the pixels of the MitoGraph binary image.

Here, we used the "splitter.py" function to separate the channels using the AICSImageIO and

tifffile libraries. Each channel is then saved as an individual Z-stack in TIFF format as 'structure'

and 'function' channels.

2.   Running MitoGraph: The open-source MitoGraph (v3.0) script is included to obtain

the MitoGraph output files as described before.

3.   Identifying individual mitochondrial components: Mitograph produces the

following three outputs containing distinct types of information- **a**) connected Components ("cc")

on interconnected elements; **b**) mitochondrial elements ("line id") on individual mitochondrial

segments; **c**) points on the mitochondrial skeleton ("point id") on individual data points. Each

"point id" belongs to a "line id", and each "line id" belongs to a cc. This information is distributed

across the following output files; 1) mitograph.txt contains information about "point id" and the

corresponding "line id" they belong to; 2) mitograph.cc file links "node id" to the "cc" they belong

to; **3**) mitograph.coo file provides geometric positions (x, y, z coordinates) for each "node id". To

streamline our analysis, we consolidated this data into a single, tidy data frame. To incorporate

"cc" information into the mitograph.txt file for network-level analysis, we identified which "point

id" entries are also "node id" entries, where nodes are the terminating points of a "line id".

Towards this, we first computed the Euclidean distance between each pair of "point id" (from

mitograph.txt) and "node id" (from mitograph.coo). Thereafter, we identified the "point id" - "node

id" pair with the smallest distance to identify the "point id" that corresponds to each "node id".

Once the "nodes" were identified, we used the mitograph.cc file to link the cc information to

each "node id". This enables us to append "cc" information to the mitograph.txt file. Thus, the

resulting final data frame integrates "cc", "line id", "point id", x, y, z coordinates and "nodes" in a

single *.csv file where the structure-function features will be appended after computation as

described below. Thus, a comprehensive data file is created for further downstream analyses.

4. Computing structural properties for networks and non-networks: We first

identified the individual networks and non-networks from the connected components ("cc"),

which include both networked and non-networked components. Here, we employed a filtering

approach based on the number of edges and nodes. Specifically, any connected component

with one edge and two nodes, along with their corresponding data, was classified as a non-

networked component. All remaining connected components, along with their corresponding

data, were categorized as networked components. This classification enabled separate analysis

of networked and non-networked components, facilitating a more focused examination of their

distinct properties. The primary measured structural features are length and diameter, from

which other features are derived. Supplementary Table-1 includes the definition and formulae of

all the structural parameters studied.

5. Computing redox function for networks and non-networks in live cells: The

redox function was computed by obtaining the average pixel intensity ratio of individual cc by

mapping the 'function' information to the 'structure' information obtained as above. Thereafter,

the redox functional property of individual mitochondrial component was computed and finally

validated (**Fig. 2C; Fig. S1G-H**). The details of these steps are below:

i. Converting MitoGraph output to pixel coordinates: MitoGraph was run using Z-stack for the

structure channel using runMitograph.py function in a batch processing mode for all cells within

a single directory. MitoGraph outputs the geometric coordinates (x, y, z) of the mitochondrial

skeleton as per the provided scale when running MitoGraph. The pixel coordinates (x_pixel,

y_pixel, z_pixel) can be retrieved from the geometric coordinates through the scale using the

formula, where D represents either x, y, or z. This is then rounded to the nearest integer, as

pixel coordinates are a discrete quantity.

768
$$D_{pixel} = \frac{D}{D_{scale}}$$

Fig. S9A shows an example of the results of this process, where the scale for x and y is 0.104

and is 0.5 for z. The conversion is confirmed by the overall overlap of the computed pixel

coordinates (in red) with the MIP of the structural channel (**Fig. S9B**).

ii. Computing the intensity values of the pixel coordinates from individual 'function'

channels: The Z-stacks Tiff images are converted into a three-dimensional array using tifffile

library. The intensity value of each pixel coordinate generated in step-i is computed from the

individual functional channels. The average intensity of the 9 pixels in a 1X3X3 voxel is

computed and assigned to the central pixel. This is repeated for every x, y, z pixel computed in
step-i. In the following example, in the slice of an image at z = 2, the intensity of a_{222} is the
average of all the colored pixels.

$$z = 2, \begin{pmatrix} b_{11} & b_{12} & b_{13} & b_{14} & b_{15} \\ b_{21} & a_{211} & a_{212} & a_{213} & b_{25} \\ b_{31} & a_{221} & a_{222} & a_{223} & b_{35} \\ b_{41} & a_{231} & a_{232} & a_{233} & b_{45} \\ b_{51} & b_{52} & b_{53} & b_{54} & b_{55} \end{pmatrix}$$

$$Pixel\ intensity\ of\ a_{222} = \frac{a_{211} + a_{212} + a_{213} + a_{221} + a_{222} + a_{223} + a_{231} + a_{232} + a_{333}}{9}$$

<u>iii. Computing pixel intensity ratio to reflect redox function:</u> Once the intensity for
each pixel is obtained (step ii), it is put into the "pixel_intensity_channel_N" column, where "N"
reflects the selected channel. For computing pixel intensity for individual element, the intensities
for each channel are grouped according to their "line_id" and averaged. These average intensity
values were used to compute pixel intensity ratio at the element level. For compute the pixel
intensity ratio at the individual cc level (networks or non-networks), the pixel intensity ratio of the
elements comprising the cc are grouped according to their "cc_id" and averaged to find the pixel
intensity ratio of the "cc" , as exemplified for an individual non-networked "cc" (see **Fig.S9C** and
equations below). These values are validated (see main text) using appropriate modules in
python (scikit-learn).

$$cc\ pixel\ intensity\ channel\ 1 = \frac{256.5 + 243.5 + 374 + 483.5 + 463.5 + 342.75}{6}$$

$$cc\ pixel\ intensity\ channel\ 2 = \frac{12.75 + 35 + 18 + 34.25 + 17 + 0}{6}$$

$$cc\_pixel\_intensity\_ratio = \frac{19.5}{360.625} = 0.05407$$

<u>iv. Visualizing single networks and non-networks within single cells in 3D graphs:</u>
Visualization of the data from the comprehensive data file is performed using the
3D_generate_graphs.py script. The data is plotted by looping through each "cc" and each "line
id". The individual points in each "line id" are then plotted to ensure that the order of the points is
preserved in the visualization. To distinguish between shapes, each "cc" is assigned a random
and unique color. These shapes are then rendered in a 3D space. The same process is
repeated for generating the LUT of the 'function' values, where the color code represents the
[Oxidation] function of individual elements.

6.   Computing mtDNA related functional features for networks and non-
networks in fixed samples: False positive signal from the nucleus was excluded by manually
drawing a region of interest (ROI) around the nucleus using ImageJ. Such processed 3D
confocal micrographs were analysed using our Python-based pipeline created including
DBSCAN (Density-Based Spatial Clustering of Applications with Noise) algorithm for detecting
the mtDNA signal mapped or unmapped to the mitochondrial marker. The following steps are
executed after importing all necessary packages (mentioned in Step C of Methods):
i. Intensity based thresholding and Density-based clustering:
The structural and functional channels are loaded as separate tiff files. Image J is used to define
the threshold from the MIP of the functional channel to include both bright and dim mtDNA
nucleoids. Next, the image is converted to binary where mtDNA signal appeared as foci. The 3D
coordinates of the foci are obtained using DBSCAN algorithm. For DBSCAN clustering, two
defined parameters are considered: epsilon (ε) - defines the maximum distance between two
points to be in one cluster and minimum sample - minimum number of pixels above the
threshold in a cluster. In our study, ε was set to 1 and the minimum sample to 4 by visual
comparison of the DBSCAN output coverage with the mtDNA distribution in the raw image.
ii. Defining quality check of clusters: A unique list of clusters that contain one point is
created and their coordinates are obtained. Thereafter, a 3x3 square is drawn on gray scale
image using each coordinate as the center, and the mean pixel intensity of that square is
computed. If the mean pixel intensity of that square is below 90% of the pixel intensity of the
center, the coordinate is removed. The coordinates in each cluster are obtained and stored in
the form of a pandas data frame.
iii. Extracting cluster information: The clusters are renamed with Z slice number as prefix in
*.csv file with filtered coordinates. The cluster size is then computed as the number of pixels in
one cluster and saved the updated data frame to a new *.csv file. To obtain cluster coordinate,
size and number the data of the *.csv is loaded, ensuring that the necessary columns are in the
data frame. To remove the pixels labelled as noise (-1) during the quality check were filtered out
from the slice of interest. This data is then saved in a separate *.csv file and turned into the final
data frame to be considered for mapping.
iv. Mapping of functional coordinates to the structural coordinates: This involves loading and
extracting coordinates from the function channel data frame created in the previous step and the
mito-SinComp structure channel data frame, separately. Next, the pairwise Euclidean distances
of each coordinate of the above data frames are computed and the minimum distance is
considered as 'mapped'. In this study, we optimized this minimum distance as ≤10 by visually

comparing DBSCAN output coverage with the mtDNA distribution in the raw image. Therefore, if
any DBSCAN cluster is beyond 6 pixels of the structural component was considered unmapped,
and the rest mapped. The final functional channel data frame for both mapped and unmapped
clusters was saved as *.csv separately. Finally, the mapped and unmapped clusters obtained
from individual z-slice are visualized by matching the spatial coordinate information of the
clusters (function) separately onto the structural channel MIP.

v. Computing and storing cluster level information from a single slice with structural level
information: The structure channel data frame and the last updated functional channel data
frame are merged on index followed by computing functional metrics like cluster number, cluster
size, and average cluster size per cc_x. This merged data frame is saved as final
*.csv file for further downstream analysis.

7. Data visualization and representation: The output of mito-SinComp analyses
is a comprehensive data file (*.csv) holding the relevant details for individual mitochondria in
individual cells; rows with point_ids under line_ids under cc_ids while columns with relevant
values for each id. This file is used to obtain and visualize various structural-function
parameters. For inter- and intracellular comparisons of SMNs and HMNs, the *.vtk files
generated by MitoGraph for individual cells were analyzed and visualized using the open-source
software ParaView-v5.13.0. All plots were generated in Python using appropriate modules.
854

Identification of novel structure metric-[CRL mito-Net] to distinguish vSMNs, **SMNs and**
HMNs: We employed a thresholding approach based on the [CRL mito-Net] metric in
combination with the previously defined cellular [Fusion1] metric, which represents the
percentage of the longest component length within a cell. The cell with the minimum Fusion1
value was identified in the HF-cells group. Within this cell, the network with the highest [CRL
mito-Net] value was used as a threshold(-1.4) to demarcate the boundary between the SMN
and HMN zones. To define the SMN region, we visually assessed the peak of this metric within
our positive control INT-Cell group, which showed a significant enrichment of SMNs. This
assessment identified a threshold of - 4.1, with any values equal to or below this point
categorized as vSMNs. This approach effectively divides the entire distribution into three distinct
zones based on the [CRL mito-Net] metric. The same thresholding criteria were consistently
applied across different cell lines and experimental conditions to study mitochondrial networks,
with a primary focus on SMNs and HMNs. Given the versatility and robustness of this
dimensionless metric, it can be adapted for thresholding in future studies, depending on the

specific research question or the mitochondrial components of interest, including both

networked and non-networked mitochondrial structures.

Radom Forest based prediction of mitochondrial function from structure: The random

forest machine learning algorithm is an ensemble learning method that performs regression by

constructing multiple decision trees during training. Each tree is trained on a bootstrapped

subset of the data, and node splits are determined using a random subset of the features in the

dataset. The final prediction is obtained by averaging the predictions of all individual trees.

We implemented the random forest model using the `scikit-learn` package (version

1.5.2). The key parameters optimized during training included: `n_estimators`: The number of

trees in the forest; `max_depth`: The maximum depth of each tree; `ccp_alpha`: The complexity

parameter used for minimal cost-complexity pruning. The algorithm selects features and split

points based on the reduction in variance for regression tasks. While training the model, 70% of

the data was used as the training set and 30% was used as the testing set. A random seed was

set to ensure all experiments were reproducible. Default values were used for parameters not

explicitly mentioned unless otherwise stated.

We performed hyperparameter tuning using the `BayesSearchCV` function from the

`scikit-optimize` (`skopt`) library. This method implements Bayesian optimization to efficiently

explore the hyperparameter space, sampling a fixed number of parameter combinations rather

than exhaustively evaluating all possibilities as in grid search. The search was configured with

the following parameters: Search space: Defined by specifying distributions for each

hyperparameter (`n_estimators`, `max_depth`, `ccp_alpha`); n_iter: The number of parameter

settings sampled from the distributions. We used a value of 100. Default values were used for

unspecified parameters unless otherwise noted.

We assessed model performance using the coefficient of determination (R^2) on the held-

out test set. R^2 is computed as :

$$R^2 = 1 - \frac{RSS}{TSS}$$

Where RSS is the sum of squares of residuals and TSS is the total sum of squares. R^2

quantifies the proportion of variance in the target variable explained by the model, with values

closer to 1 indicating better fit.

We evaluated model performance using 5-fold cross-validation, implemented via the

`cross_val_score` function from `scikit-learn`. The training set was partitioned into five equal-

sized folds, each preserving the overall distribution of the target variable. The model was trained

on four folds and validated on the remaining fold, with this process repeated five times such that
each fold served as the validation set once. The R^2 score were computed for each validation
fold and aggregated to produce an average score with standard deviation. This approach
provided an estimate of generalization error while mitigating bias from any single train-test split.
The same cross-validation folds were used consistently during hyperparameter optimization and
final evaluation to ensure comparability.
To interpret the model, we analyzed feature importance scores from the trained random
forest using the `feature_importances_` attribute of the Random Forest Regressor class. This
attribute computes scores as the mean decrease in impurity (Gini importance) across all trees,
reflecting each feature's relative contribution to predictive accuracy. The importance values are
normalized to sum to 1 for comparative analysis. Performance metrics and feature importance
were computed using native `scikit-learn` implementations to ensure consistency with the
training process. The test set was not used during hyperparameter optimization or feature
selection to prevent data leakage.
916
High Dimensional weighted gene correlation network analyses (hdWGCNA):
<u>Single-Cell RNA Seq:</u> TCDD exposed cells were processed for scRNA-seq using the
10X Genomics 3' v3.1 NextGem platform and analyses were performed using Seurat
(v4.3.3). Notably, the cells were tested for > 90% viability before scRNA-seq. Libraries were
constructed through cDNA fragmentation, end repair, adaptor ligation, and dual-indexed PCR
amplification. Sequencing was performed on Illumina Nextseq500 (28 bp R1, 90 bp R2, 10 bp
i7/i5 indices) with ≥20,000 reads/cell. Only cells expressing a minimum of 2000 detectable
genes were included using the 'nFeature_RNA' filter and variable feature sets were identified by
FindVariableFeatures function in Seurat. This filtering also excluded dead cells with >10%
mtDNA gene expression in all our datasets analyzed. This allowed us to exclude the percent.mt
filter that uses mtDNA-based exclusion of dead cells. *nFeature_RNA* >2000 was applied prior to
normalization of data using the *NormalizeData* function in Seurat. Approximately ~3000 genes
in an average of 5000 cells were covered in our data in each population. The same filters were
used for processing publicly available raw data from cSCC patients[50]. Specific sets of samples
were combined using the *IntegrateData* function for the outlined comparison sets - transformed,
exposed, DRP1 knockdown and cSCC resulting in four sets of scRNASeq analyses. These
integrated datasets were scaled and clustering of cells was performed with up to 20 dimensions
(dim = 1:20) with 0.2 resolution quantified for stem cell high cluster identification, visualized by
UMAP. Marker genes for each cluster were identifed with *FindAllMarkers* function with statistical

significance calculation performed using by Wilcoxon Rank Sum test. While, the FindMarkers

function (pseudobulk analyses) of Seurat was used to compute the differential expression of the

cluster markers (normalized data before integration (assay 'RNA')) using the Wilcoxon rank

sum test. Average Expression was computed for select genes of interest using the *Average*

Expression function. GSEA analyses were performed for scRNASeq clusters with ranked log

fold change (LFC) of markers (LFC > 0.1, p < 0.01, FDR q value p < 0.01) using MsigDB and

Reactome databases. To sum up, Seurat objects from the resultant count matrices were

processed, integrated, visualized, and analyzed for relevant downstream analyses including

pairwise differential gene expression analyses (using FindMarkers function), cell cluster

analyses (using FindAllMarkers function), and Gene Set Enrichment Analysis (GSEA) for cell

cluster annotation. Immunoblotting was performed to validate key scRNAseq finding using

standard methods using commercial antibodies (Key Resource Table).

hdWGCNA design: We applied method on scRNA-seq data to identify transcriptomic

networks[62,63] following the workflow outlined in **Fig. S3B**. We used integrated Seurat objects for

both pseudobulk and cell cluster-based analyses, considered the top 20 / 25 genes (ranked by

kME) to visualize transcriptomic interactions through hub-gene network plots. Functional

annotation of the hdWGCNA modules was performed using GSEA where the modules with ≥ 5

mtDNA genes as the leading-edge genes were reassigned as 'mtDNA' in our analyses. Edge

distance and the number of edges were quantified using TOM, providing insights into gene

connectivity and interaction strength.

Detailed workflow of hdWGCNA: hdWGCNA[62](smorabit.github.io/hdWGCNA/) was

used to identify gene interaction modules amongst differentially expressed genes (> 0.25

avg_log2FC, <10^{-5} p_val_adj, pct ≥ 0.25 in pseudobulk analyses using FindMarkers function)

and amongst the characteristic marker genes for the identified cell clusters (> 0.25 avg_log2FC,

<10^{-5} p_val_adj). QC plots of nCounts_RNA distributions were used to establish lower

thresholds for outlier removal (**Fig. S9D).** This processed integrated Seurat object is used as an

input for respective hdWGCNA, both for pseudobulk and cell cluster levels. The hdWGCNA

object was initialized, and metacells were constructed using K-nearest neighbors (KNN) (**Figs.**

S9E-F). Gene expression matrices were constructed, and pairwise Pearson's correlations were

computed, transformed into adjacency matrices using a soft-thresholding power approach (**Fig.**

S9G), and used to construct networks using Topological Overlap Measures (TOM) values. The

hdWGCNA algorithm ensured a scale-free topology during signed network construction for each

hdWGCNA object for both pseudobulk and cell cluster. Transcriptomic modules were detected

using hierarchical clustering and TOM, with module eigengenes (MEs) computed for further

analysis. Module connectivity (kME) values were computed to identify hub genes characterized

by high kME values. Selection of specific parameter cut-offs for hdWGCNA are elaborated

below: **a.** K-nearest neighbors (KNN)-based dimensionality reduction: Metacells for each

experimental/control group (pseudobulk) and specific clusters were constructed using KNN,

leveraging biological similarity to define cellular subpopulations in each hdWGCNA object.

Pearson correlation-based density plots guided the selection of K value of 20 with a maximum

of 10 shared neighbors for an average of 5000 cells in each population, optimizing the balance

between capturing local structure and minimizing noise in sparse data (**Figs. S9E-F**); **b.** Soft

Thresholding Power (β): The optimal β value was selected by evaluating the scale-free topology

fit index, to enhance strong correlations and reduce weaker ones, aiming for the lowest power

where the network closely approximated a scale-free structure. The β values between 6 and 10

were determined as optimal, balancing network sparsity with scale-free topology at both

pseudobulk and cell cluster-based hdWGCNA objects (solid black circles on plots in (**Fig. S9G**).

This transformation enabled robust, biologically meaningful signed networks across cellular

subpopulations; **c.** Module Connectivity (kME): kME measures the correlation between a gene's

expression profile and its module eigengene, indicating the gene's alignment with the module's

expression pattern. High kME values suggest that a gene has a central role within the module,

while low values indicate a less integral role. Genes in each module were ranked by kME,

identifying the top 20 in pseudobulk (**Fig. 3E, S3D, S6B-C**) and top 25 (**Fig. 5F, 5K, S7D**) in cell

cluster-based hdWGCNA for functional significance. This kME analysis highlighted closely

interacting clusters, enabling us to identify top-ranking hub genes within a network.

Statistical Analyses: Individual experiments were performed at least 3 times, and appropriate

statistics were applied. The N detected for individual mitochondrial component per cell ranges

from 30 to >100. N for each analysis is mentioned in the figure and can be scaled as necessary.

Appropriate statistical analyses for single cell microscopy were conducted using either Excel or

Python (SciPy); The Kolmogorov-Smirnov test was applied to histogram and ridge plots; the

Mann-Whitney U test to violin and box plots; the Student's t-Test to bar plots; and the Chi-

square test to bivariate scatter plots. Corresponding p-values were computed and adjusted for

multiple hypotheses testing using the Holm-Bonferroni correction method. Bootstrapping was

used for resampling, as needed. The hdWGCNA and Random Forest used inbuilt statistical

analyses as required.

1002

1003   **Data Availability:** The raw, analyzed and meta data of scRNA-seq experiment have been

1004   deposited in the Gene expression Omnibus (accession number GSE285728). All data matrices,

1005   scripts, and hdWGCNA results will be deposited in public repositories upon publication, to

1006   ensure transparency and reproducibility for future research.

1007

Acknowledgements: The research, D.P. and B.S were supported by NIH (R33

ES025662/ES/NIEHS) and DBT-Wellcome Trust India Alliance (IA/S/20/2/505198/WTDBT);

S.A. was supported by DBTRA/2023-24/NE/Ashoka/19; M.S. was supported by DBT-JRF-

DBT/2023-24/Ashoka Uni/2196; B.W. was supported by (IA/S/20/2/505198/WTDBT); B.G. was

supported by DBT/2022-23/ASHOKA/2156; PS was supported by Mphasis F1 Foundation. We

acknowledge UAB Flow Cytometry and Single Cell Services Core (supported by P30

AR048311, P30 AI027667) and Dr. Mike Crowley from the Heflin Genomics Core for scRNA-

seq, we sincerely thank Dr. LS Shashidhara, Dr. Alok Bhattacharya and Dr. Gautam Menon for

critically reading the manuscript, and Dr. KM Poonam for crosschecking RF code set.

References

1. Suomalainen, A., and Nunnari, J. (2024). Mitochondria at the crossroads of health and disease. Cell *187*, 2601–2627. https://doi.org/10.1016/J.CELL.2024.04.037.

2. Reczek, C.R., and Chandel, N.S. (2015). ROS-dependent signal transduction. Curr Opin Cell Biol *33*, 8. https://doi.org/10.1016/J.CEB.2014.09.010.

3. Spurlock, B., Tullet, J.M.A., Hartman, J.L., and Mitra, K. (2020). Interplay of mitochondrial fission-fusion with cell cycle regulation: Possible impacts on stem cell and organismal aging. Exp Gerontol *135*, 110919. https://doi.org/10.1016/J.EXGER.2020.110919.

4. Glancy, B., Kim, Y., Katti, P., and Willingham, T.B. (2020). The Functional Impact of Mitochondrial Structure Across Subcellular Scales. Preprint at Frontiers Media S.A., https://doi.org/10.3389/fphys.2020.541040 https://doi.org/10.3389/fphys.2020.541040.

5. Ryu, K.W., Fung, T.S., Baker, D.C., Saoi, M., Park, J., Febres-Aldana, C.A., Aly, R.G., Cui, R., Sharma, A., Fu, Y., et al. (2024). Cellular ATP demand creates metabolically distinct subpopulations of mitochondria. Nature 2024, 1–9. https://doi.org/10.1038/s41586-024-08146-w.

6. Quintana-Cabrera, R., and Scorrano, L. (2023). Determinants and outcomes of mitochondrial dynamics. Preprint at Cell Press, https://doi.org/10.1016/j.molcel.2023.02.012

7. Chan, D.C. (2020). Mitochondrial Dynamics and Its Involvement in Disease. Annual Review of Pathology: Mechanisms of Disease *15*, 235–259. https://doi.org/10.1146/annurev-pathmechdis-012419-032711.

8. Tábara, L.C., Segawa, M., and Prudent, J. (2024). Molecular mechanisms of mitochondrial dynamics. Nature Reviews Molecular Cell Biology 2024, 1–24. https://doi.org/10.1038/s41580-024-00785-1.

9. Viana, M.P., Brown, A.I., Mueller, I.A., Goul, C., Koslover, E.F., and Rafelski, S.M. (2020). Mitochondrial Fission and Fusion Dynamics Generate Efficient, Robust, and Evenly Distributed Network Topologies in Budding Yeast Cells. Cell Syst *10*, 287-297.e5. https://doi.org/10.1016/j.cels.2020.02.002.

10. Glancy, B., Hartnell, L.M., Malide, D., Yu, Z.X., Combs, C.A., Connelly, P.S., Subramaniam, S., and Balaban, R.S. (2015). Mitochondrial reticulum for cellular energy distribution in muscle. Nature *523*, 617–620. https://doi.org/10.1038/nature14614.

11. Nunnari, J., Marshall, W.F., Straight, A., Murray, A., Sedat, J.W., and Walter, P. (1997). Mitochondrial transmission during mating in Saccharomyces cerevisiae is determined by

mitochondrial fusion and fission and the intramitochondrial segregation of mitochondrial
DNA. Mol Biol Cell *8*, 1233–1242. https://doi.org/10.1091/MBC.8.7.1233.

12. Kowald, A., and Kirkwood, T.B.L. (2011). Evolution of the mitochondrial fusion - Fission
cycle and its role in aging. Proc Natl Acad Sci U S A *108*, 10237–10242.
https://doi.org/10.1073/PNAS.1101604108/ASSET/9161FB65-F251-4490-839F-
E80F24BE925C/ASSETS/GRAPHIC/PNAS.1101604108FIG02.JPEG.

13. Shirihai, O.S., Song, M., and Dorn, G.W. (2015). How mitochondrial dynamism
orchestrates mitophagy. Circ Res *116*, 1835.
https://doi.org/10.1161/CIRCRESAHA.116.306374.

14. Rafelski, S.M., Viana, M.P., Zhang, Y., Chan, Y.H.M., Thorn, K.S., Yam, P., Fung, J.C.,
Li, H., Costa, L.D.F., and Marshall, W.F. (2012). Mitochondrial Network Size Scaling in
Budding Yeast. Science *338*, 822. https://doi.org/10.1126/SCIENCE.1225720.

15. Ouellet, M., Guillebaud, G., Gervais, V., Lupien St-Pierre, D., and Germain, M. (2017). A
novel algorithm identifies stress-induced alterations in mitochondrial connectivity and
inner membrane structure from confocal images. PLoS Comput Biol *13*, 1–23.
https://doi.org/10.1371/journal.pcbi.1005612.

16. Ilamathi, H.S., Ouellet, M., Sabouny, R., Desrochers-Goyette, J., Lines, M.A., Pfeffer, G.,
Shutt, T.E., and Germain, M. (2021). A new automated tool to quantify nucleoid
distribution within mitochondrial networks. Scientific Reports 2021 11:1 *11*, 1–14.
https://doi.org/10.1038/s41598-021-01987-9.

17. Rohani, A., Kashatus, J.A., Sessions, D.T., Sharmin, S., and Kashatus, D.F. (2020). Mito
Hacker: a set of tools to enable high-throughput analysis of mitochondrial network
morphology. Sci Rep *10*. https://doi.org/10.1038/S41598-020-75899-5.

18. Karbowski, M., Arnoult, D., Chen, H., Chan, D.C., Smith, C.L., and Youle, R.J. (2004).
Quantitation of mitochondrial dynamics by photolabeling of individual organelles shows
that mitochondrial fusion is blocked during the Bax activation phase of apoptosis. J Cell
Biol *164*, 493–499. https://doi.org/10.1083/JCB.200309082.

19. Iannetti, E.F., Smeitink, J.A.M., Beyrath, J., Willems, P.H.G.M., and Koopman, W.J.H.
(2016). Multiplexed high-content analysis of mitochondrial morphofunction using live-cell
microscopy. Nature Protocols 2016 11:9 *11*, 1693–1710.
https://doi.org/10.1038/nprot.2016.094.

20. Lefebvre, A.E.Y.T., Ma, D., Kessenbrock, K., Lawson, D.A., and Digman, M.A. (2021).
Automated segmentation and tracking of mitochondria in live-cell time-lapse images. Nat
Methods *18*, 1091–1102. https://doi.org/10.1038/s41592-021-01234-z.

21.   Spurlock, B., Gupta, P., Basu, M.K., Mukherjee, A., Hjelmeland, A.B., Darley-Usmar, V.,
Parker, D., Foxall, M.E., and Mitra, K. (2019). New quantitative approach reveals
heterogeneity in mitochondrial structure–function relations in tumor-initiating cells. J Cell
Sci *132*. https://doi.org/10.1242/jcs.230755.

22.   Mitra, K., Wunder, C., Roysam, B., Lin, G., and Lippincott-Schwartz, J. (2009). A
hyperfused mitochondrial state achieved at G1-S regulates cyclin E buildup and entry into
S phase. Proc Natl Acad Sci U S A *106*, 11960–11965.
https://doi.org/10.1073/PNAS.0904875106/SUPPL_FILE/SV5.MOV.

23.   Spurlock, B., Parker, D., Basu, M.K., Sajina, G., Liu, S., Siegal, G.P., Gunter, A., Moran,
A., and Mitra, K. (2021). Fine-tuned repression of Drp1-driven mitochondrial fission
primes a "stem/progenitor-like state" to support neoplastic transformation. Elife *10*.
https://doi.org/10.7554/ELIFE.68394.

24.   Liang, R., Arif, T., Kalmykova, S., Kasianov, A., Lin, M., Menon, V., Qiu, J., Bernitz, J.M.,
Moore, K., Lin, F., et al. (2020). Restraining Lysosomal Activity Preserves Hematopoietic
Stem Cell Quiescence and Potency. Cell Stem Cell *26*, 359-376.e7.
https://doi.org/10.1016/j.stem.2020.01.013.

25.   Folmes, C.D.L., Dzeja, P.P., Nelson, T.J., and Terzic, A. (2012). Metabolic plasticity in
stem cell homeostasis and differentiation. Cell Stem Cell *11*, 596–606.
https://doi.org/10.1016/J.STEM.2012.10.002.

26.   Khacho, M., and Slack, R.S. (2017). Mitochondrial activity in the regulation of stem cell
self-renewal and differentiation. Curr Opin Cell Biol *49*, 1–8.
https://doi.org/10.1016/J.CEB.2017.11.003.

27.   Chakrabarty, R.P., and Chandel, N.S. (2021). Mitochondria as Signaling Organelles
Control Mammalian Stem Cell Fate. Preprint at Cell Press,
https://doi.org/10.1016/j.stem.2021.02.011 https://doi.org/10.1016/j.stem.2021.02.011.

28.   Loh, J.J., and Ma, S. (2024). Hallmarks of cancer stemness. Cell Stem Cell *31*, 617–639.
https://doi.org/10.1016/J.STEM.2024.04.004.

29.   Meacham, C.E., and Morrison, S.J. (2013). Tumor heterogeneity and cancer cell
plasticity. Nature *501*, 328. https://doi.org/10.1038/NATURE12624.

30.   Vasan, K., Werner, M., and Chandel, N.S. (2020). Mitochondrial Metabolism as a Target
for Cancer Therapy. Cell Metab *32*, 341–352.
https://doi.org/10.1016/J.CMET.2020.06.019/ASSET/57DF9986-127A-47AA-BE81-
81EDC3C54529/MAIN.ASSETS/GR5.JPG.

31. Bonnay, F., Veloso, A., Steinmann, V., Köcher, T., Abdusselamoglu, M.D., Bajaj, S.,
Rivelles, E., Landskron, L., Esterbauer, H., Zinzen, R.P., et al. (2020). Oxidative
Metabolism Drives Immortalization of Neural Stem Cells during Tumorigenesis. Cell *182*,
1490-1507.e19. https://doi.org/10.1016/j.cell.2020.07.039.
32. Wu, M.J., Chen, Y.S., Kim, M.R., Chang, C.C., Gampala, S., Zhang, Y., Wang, Y.,
Chang, C.Y., Yang, J.Y., and Chang, C.J. (2019). Epithelial-Mesenchymal Transition
Directs Stem Cell Polarity via Regulation of Mitofusin. Cell Metab *29*, 993-1002.e6.
https://doi.org/10.1016/J.CMET.2018.11.004.
33. Folmes, C.D.L., Ma, H., Mitalipov, S., and Terzic, A. (2016). Mitochondria in pluripotent
stem cells: stemness regulators and disease targets. Curr Opin Genet Dev *38*, 1–7.
https://doi.org/10.1016/J.GDE.2016.02.001.
34. Lisowski, P., Kannan, P., Mlody, B., and Prigione, A. (2018). Mitochondria and the
dynamic control of stem cell homeostasis. EMBO Rep *19*.
https://doi.org/10.15252/EMBR.201745432/ASSET/D6499ABD-4FAC-4033-ACD0-
4F8523E03919/ASSETS/GRAPHIC/EMBR201745432-FIG-0004-M.PNG.
35. Kashatus, J.A., Nascimento, A., Myers, L.J., Sher, A., Byrne, F.L., Hoehn, K.L., Counter,
C.M., and Kashatus, D.F. (2015). Erk2 phosphorylation of Drp1 promotes mitochondrial
fission and MAPK-driven tumor growth. Mol Cell *57*, 537–551.
https://doi.org/10.1016/J.MOLCEL.2015.01.002.
36. Serasinghe, M.N., Wieder, S.Y., Renault, T.T., Elkholi, R., Asciolla, J.J., Yao, J.L.,
Jabado, O., Hoehn, K., Kageyama, Y., Sesaki, H., et al. (2015). Mitochondrial division is
requisite to RAS-induced transformation and targeted by oncogenic MAPK pathway
inhibitors. Mol Cell *57*, 521. https://doi.org/10.1016/J.MOLCEL.2015.01.003.
37. Tondera, D., Grandemange, S., Jourdain, A., Karbowski, M., Mattenberger, Y., Herzig,
S., Da Cruz, S., Clerc, P., Raschke, I., Merkwirth, C., et al. (2009). SLP-2 is required for
stress-induced mitochondrial hyperfusion. EMBO J *28*, 1589–1600.
https://doi.org/10.1038/EMBOJ.2009.89.
38. Gomes, L.C., Benedetto, G. Di, and Scorrano, L. (2011). During autophagy mitochondria
elongate, are spared from degradation and sustain cell viability. Nat Cell Biol *13*, 589–
598. https://doi.org/10.1038/NCB2220.
39. Spurlock, B.M., Xie, Y., Song, Y., Ricketts, S.N., Hua, J.R., Chi, H.R., Nishtala, M.,
Salmenov, R., Liu, J., and Qian, L. (2025). Mitochondrial fusion and cristae reorganization
facilitate acquisition of cardiomyocyte identity during reprogramming of murine fibroblasts.
Cell Rep *44*, 115377. https://doi.org/10.1016/J.CELREP.2025.115377.

40.   Holmström, K.M., and Finkel, T. (2014). Cellular mechanisms and physiological

1153   consequences of redox-dependent signalling. Nature Reviews Molecular Cell Biology

1154   2014 15:6 *15*, 411–421. https://doi.org/10.1038/nrm3801.

1155   41.   Willems, P.H.G.M., Rossignol, R., Dieteren, C.E.J., Murphy, M.P., and Koopman, W.J.H.

1156   (2015). Redox Homeostasis and Mitochondrial Dynamics. Cell Metab *22*, 207–218.

1157   https://doi.org/10.1016/J.CMET.2015.06.006.

1158   42.   Parker, D.J., Iyer, A., Shah, S., Moran, A., Hjelmeland, A.B., Basu, M.K., Liu, R., and

1159   Mitra, K. (2015). A new mitochondrial pool of cyclin E, regulated by Drp1, is linked to cell-

1160   density-dependent cell proliferation. J Cell Sci *128*, 4171–4182.

1161   https://doi.org/10.1242/JCS.172429.

1162   43.   König, T., and McBride, H.M. (2024). Mitochondrial-derived vesicles in metabolism,

1163   disease, and aging. Cell Metab *36*, 21–35. https://doi.org/10.1016/j.cmet.2023.11.014.

1164   44.   Purba, T.S., Haslam, I.S., Poblet, E., Jiménez, F., Gandarillas, A., Izeta, A., and Paus, R.

1165   (2014). Human epithelial hair follicle stem cells and their progeny: Current state of

1166   knowledge, the widening gap in translational research and future challenges. BioEssays

1167   *36*, 513–525. https://doi.org/10.1002/BIES.201300166.

1168   45.   Fenton, A.R., Jongens, T.A., and Holzbaur, E.L.F. (2021). Mitochondrial dynamics:

1169   Shaping and remodeling an organelle network. Curr Opin Cell Biol *68*, 28–36.

1170   https://doi.org/10.1016/J.CEB.2020.08.014.

1171   46.   Murphy, M.P. (2009). How mitochondria produce reactive oxygen species. Biochem J

1172   *417*, 1–13. https://doi.org/10.1042/BJ20081386.

1173   47.   DiMeglio, P., Duarte, J.H., Ahlfors, H., Owens, N.D.L., Li, Y., Villanova, F., Tosi, I., Hirota,

1174   K., Nestle, F.O., Mrowietz, U., et al. (2014). Activation of the Aryl Hydrocarbon Receptor

1175   Dampens the Severity of Inflammatory Skin Conditions. Immunity *40*, 989.

1176   https://doi.org/10.1016/J.IMMUNI.2014.04.019.

1177   48.   Stewart, J.B., and Chinnery, P.F. (2021). Extreme heterogeneity of human mitochondrial

1178   DNA from organelles to populations. Nat Rev Genet *22*, 106–118.

1179   https://doi.org/10.1038/s41576-020-00284-x.

1180   49.   Tatsuta, T., and Langer, T. (2017). Prohibitins. Current Biology *27*, R629–R631.

1181   https://doi.org/10.1016/j.cub.2017.04.030.

1182   50.   Ji, A.L., Rubin, A.J., Thrane, K., Jiang, S., Reynolds, D.L., Meyers, R.M., Guo, M.G.,

1183   George, B.M., Mollbrink, A., Bergenstråhle, J., et al. (2020). Multimodal Analysis of

1184   Composition and Spatial Architecture in Human Squamous Cell Carcinoma. Cell *182*,

1185   497-514.e22. https://doi.org/10.1016/j.cell.2020.05.039.

51.   Rey, T., Tábara, L.C., Prudent, J., and Minczuk, M. (2023). mtFociCounter for automated
single-cell mitochondrial nucleoid quantification and reproducible foci analysis. Nucleic
Acids Res *51*, e107. https://doi.org/10.1093/NAR/GKAD864.

52.   Newman, L.E., and Shadel, G.S. (2023). Annual Review of Biochemistry Mitochondrial
DNA Release in Innate Immune Signaling. https://doi.org/10.1146/annurev-biochem-
032620.

53.   Vafai, S.B., and Mootha, V.K. (2012). Mitochondrial disorders as windows into an ancient
organelle. Nature *491*, 374–383. https://doi.org/10.1038/nature11707.

54.   Chen, W., Zhao, H., and Li, Y. (2023). Mitochondrial dynamics in health and disease:
mechanisms and potential targets. Signal Transduction and Targeted Therapy 2023 8:1
8, 1–25. https://doi.org/10.1038/s41392-023-01547-9.

55.   Chan, D.C. (2020). Mitochondrial Dynamics and Its Involvement in Disease. Annu Rev
Pathol *15*, 235–259. https://doi.org/10.1146/ANNUREV-PATHMECHDIS-012419-032711.

56.   Shadel, G.S., and Horvath, T.L. (2015). Mitochondrial ROS Signaling in Organismal
Homeostasis. Cell *163*, 560. https://doi.org/10.1016/J.CELL.2015.10.001.

57.   Wang, W., Fang, H., Groom, L., Cheng, A., Zhang, W., Liu, J., Wang, X., Li, K., Han, P.,
Zheng, M., et al. (2008). Superoxide flashes in single mitochondria. Cell *134*, 279–290.
https://doi.org/10.1016/J.CELL.2008.06.017.

58.   Ying, Z., Chen, K., Zheng, L., Wu, Y., Li, L., Wang, R., Long, Q., Yang, L., Guo, J., Yao,
D., et al. (2016). Transient Activation of Mitoflashes Modulates Nanog at the Early Phase
of Somatic Cell Reprogramming. Cell Metab *23*, 220–226.
https://doi.org/10.1016/J.CMET.2015.10.002.

59.   van Velthoven, C.T.J., and Rando, T.A. (2019). Stem Cell Quiescence: Dynamism,
Restraint, and Cellular Idling. Cell Stem Cell *24*, 213–225.
https://doi.org/10.1016/J.STEM.2019.01.001.

60.   Kashyap, A., Rapsomaniki, M.A., Barros, V., Fomitcheva-Khartchenko, A., Martinelli,
A.L., Rodriguez, A.F., Gabrani, M., Rosen-Zvi, M., and Kaigala, G. (2022). Quantification
of tumor heterogeneity: from data acquisition to metric generation. Trends Biotechnol *40*,
647–676. https://doi.org/10.1016/J.TIBTECH.2021.11.006/ASSET/89F89478-295E-4185-
88AE-780FF65BA8E0/MAIN.ASSETS/GR4.JPG.

61.   Viana, M.P., Lim, S., and Rafelski, S.M. (2015). Quantifying mitochondrial content in
living cells. Methods Cell Biol *125*, 77–93. https://doi.org/10.1016/BS.MCB.2014.10.003.

62.   Morabito, S., Reese, F., Rahimzadeh, N., Miyoshi, E., and Swarup, V. High dimensional
1219   co-expression networks enable discovery of transcriptomic drivers in complex biological
1220   systems. https://doi.org/10.1101/2022.09.22.509094.
1221   63.   Horvath, S., and Dong, J. (2008). Geometric interpretation of gene coexpression network
1222   analysis. PLoS Comput Biol *4*. https://doi.org/10.1371/journal.pcbi.1000117.

1223

1224

1225

1226

1227

1228

1229

1230

1231

1232

1233

1234

1235

1236

1237

1238

1239

1240

1241

1242

1243

1244

1245

1246

1247

1248

1249

1250

Main Figure Legends

Figure. 1: Modular design of quantitative mito-SinComp approach. Mitochondrial heterogeneity can be untangled at hierarchical levels following 6 steps in live and fixed cells or tissues. Multiple structural features (S) are extracted and the function (F) is computed and mapped on the function. S-F analyses of single mitochondrial components can be performed with various statistical methods. Standalone modular use of mito-SinComp to elucidate mitochondrial physiology is demonstrated through examples of redox function and functional mtDNA organization. Combinatorial use of mito-SinComp to study mitochondrial heterogeneity in the cell physiology is demonstrated using scRNA-seq modality to study stemness.

Figure. 2:

(A). Labelled views of representative MTG stained single HaCaT cells.

(**B**). Histogram distribution of lengths of individual mitochondrial Networks and non-Networks (ignoring cell identity) for the colour-coded cell groups; arrow marks the hyperfused networks.

(C). Bivariate scatter plot of signal intensity values obtained with automated mito-SinComp and manually from ImageJ (left) from 13 mitochondrial components outlined in the representative cell expressing mito-PSMO2 and mito-roGFP (right).

(D). Box plot of average mitochondrial length and total number of mitochondrial Networks and non-Networks within individual cells in a pooled population of 10 experimental groups.

(E). Ridgeline plots comparing distributions of named (S) and (F) features between Networked and non-Networked mitochondria from (D); Median and Coefficient of Variation (in bold) are indicated for each feature.

(**F**). Bivariate S-F scatter plots of $[Oxidation]_{CC}$ and individual structural features with their frequency distributions (dotted line) for Networked and non-Networked mitochondria from (D).

(G). Representative 3D S-F Graph of an HaCaT cell harboring mitochondrial components (outlined) in the median range of features; inset depicts bivariate S-F scatter plot of network $[Size]_{CC}$ and INH-$[Oxidation]_{CC}$ plot of each component.

(H). Bivariate S-F scatter plot of $[Oxidation]_{CC \text{ or } EL}$ and $[Size]_{CC}$ for Networks and non-Networks for the cell in (G).

(I). Bar plot of R^2 values from ML based prediction of $[Oxidation]_{CC}$ on the test set of the Networked and non-Networks from (D).

(J). Scatter plot of predicted $[Oxidation]_{CC}$ values and True values from (I).

1284 **(K).** Plot of [Oxidation]$_{CC}$ prediction error (MAE) on test set (bar) with distribution of [Oxidation]$_{CC}$

1285 (line) for Networks and non-Networks.

1286 **(L).** Plot of R^2 values from ML based prediction of [Oxidation]$_{CC}$ with reduction of total data set

1287 size for the Networks to the level of the non-Networks.

1288

1289 **signifies Holmes-Bonferroni corrected p-value < 0.05; Scale bar: 5 µm; N = sample size.

1290

Figure. 3:

(A). Histogram distribution of [CRL mito-Net] or [CRL mito-nonNet], with demarcation of

thresholds (dashed lines) for vSMNs, SMNs and HMNs. (threshold calculation in Methods).

(B). Violin plot of vSMNs, SMNs, HMNs abundance in individual cells in color-coded cell groups.

(C). Magnified views of the extracted Networks and non-Networks from representative cells

(**Fig. 2A**) with their corresponding micrographs.

(D). Ridgeline plots comparing the distribution of the named (S) features between SMNs and

HMNs; Median and Coefficient of Variation (in bold) are indicated for each feature.

(E). Hub-gene network plot displaying the top 20 genes from the outlined module in **Fig.S3C**;

with kME module plot and table (insets) highlighting KRT15 direct interacting genes.

(**F**). Bar plots of percentage change in median expression of KRT15 and its direct interacting

partners (from E) in the DRP1 manipulated cells from NT control, across identified cell clusters.

(G). Violin plot of SMN and HMN abundance within individual cells of color-coded cell groups.

(H). Table showing number of SMNs, HMNs, and SMN / HMN ratio from (G).

(I). Ridgeline plots comparing the distribution of the named (S) features of SMNs and HMNs

from (G); Median and Coefficient of Variation (in bold) are indicated for each feature.

(J). Trivariate S-F scatter plot of named SMN or HMN S-F features identified from **Fig. 2D**.

(K). Bar plot of feature importance values for individual network features from ML based

predictive model described in Fig. 2I-L.

(L). Schematic describing connection of [CRL mito-Net] classified SMNs and HMNs to KRT15

marked stemness transcriptomic network in Drp1 manipulated cells.

1312

1313 **signifies Holmes-Bonferroni corrected p-value < 0.05; Scale bar: 5 µm for Networks and 1µm

1314 for non-Networks; N = sample size.

1315

1316

1317

Figure. 4:

(A). Mitochondrial potential of individual cells measured from confocal micrographs of individual

HaCaT cells exposed to Toluene, TCDD 1nM or 10nM for 1 or 4 hours.

(**B**). Representative 3D S-F Graphs of HaCaT cells stably expressing mito-PSMO2 and mito-

roGFP from the labelled groups showing [Oxidation]$_{EL}$ with colour-coded scale.

(C). Box plot of average length mitochondrial Networks and non-Networks within individual cells

in labelled groups.

(D). Histogram distribution of [CRL mito-Net] or [CRL mito-nonNet] in labelled groups with

demarcation of thresholds (dashed lines) for SMNs and HMNs (as in **Fig. 3A**).

(E). Violin plot of SMN and HMN abundance within individual cells in labelled cell groups.

(F). Ridgeline plots of distribution of the named (S) features of SMNs and HMNs between

labelled cell groups; Median and Coefficient of Variation (in bold) are indicated for each feature.

(G). Bar plots of mean of [Oxidation]$_{CC}$ of Networked and non-Networked mitochondria of

labelled groups; errors : SD.

(H). Bar plots of [Oxidation]$_{CC}$ in SMNs and HMNs in the color coded groups; errors : SD.

(I). Ridgeline plot of the distribution of [Oxidation]$_{CC}$ in SMNs in the color coded groups. The

solid and dashed lines distinguish the [Ox.]-Hi, [Ox.]-Int and Ox-Lo, where the solid line

demarcates minimum abundance between the two peaks of Tol-1h control and dashed lines on

both sides demarcates the distance to the upper quartile level of the T10-1h control.

(J). Bivariate S-F scatter plots of [Oxidation]$_{CC}$ and [Edge or Node Density] for SMNs in the color

coded groups; vertical and horizontal lines demarcate the S and F features, respectively.

1339

1340 **signifies Holmes-Bonferroni corrected p-value < 0.05; Scale bar: 5 µm; N = sample size

1341

Figure. 5:

(A). Pairwise pseudo-bulk analyses of genes encoding mitochondrial proteins (mtDNA or

Nuclear DNA encoded) from scRNA-seq data of HaCAT cells in labelled groups.

(B). Immunoblot analysis of MT-CYB and NDUFB3, with TOM20 and β-actin as loading controls

in labelled groups; quantified levels are normalized by loading control.

(C). UMAP plot of scRNA-seq derived clusters within cell populations of the labelled groups; the

numbers indicate % abundance of cells in each cluster based on Fig. S5C (see Methods for

functional annotation).

(D). Bar plots showing average logFC of all Keratin genes in the named cell clusters, with stem

cell-specific KRT15 boxed; (Krt15 raw data in Fig. **S8A**).

(E). Bar plots showing the percentage change in median expression of mtDNA-encoded genes relative to the control in labeled groups across cell clusters (raw data in **Fig. S8B**); * mark MT-ND1, MT-ND4, MT-ND5; inset shows bar plots with for the core fission–fusion genes, with arrows on DNM1L gene for Drp1.

(F). Hub-gene network plot displaying the top 25 genes from the kME plot (inset) highlighting KRT15 interacting mtDNA genes.

(G). Line plot showing percentage enrichment of SMNs and HMNs along the threshold values of SMN / HMN ratio within individual cells in the labelled groups.

(H). Trivariate S-F scatter plot of named (S) and (F) features for networked mitochondrial components in color coded groups.

(I). Bar plot showing differential expression of mtDNA genes in the labelled groups.

(J). Absolute average expression of mtDNA genes and major keratins in cSCC patients (top) in normal (N) and tumor (T) tissue samples (top), while change in average expression in tumor from normal (bottom); original data from reference 50.

(K). Hub-Gene Network plot displaying the top 25 genes from the kME plot (inset) highlighting KRT15 interacting mtDNA genes.

Figure. 6: Quantitative mito-SinComp S-F analyses of functional organization of mtDNA within networks and non-networks in fixed cells and tissues.

(A) Representative micrograph of a co-immunostained HaCaT cell with its 3D Structure Graph and Mapped / unmapped mtDNA nucleoids projected on MitoGraph binary image; outlined are identified SMNs and magnified non-network (*) with its feature dimensions mentioned.

(B) Ridgeline plots comparing distribution of the S-F features of mitochondrial Networks and non-Networks; Median and Coefficient of Variation (in bold) are indicated for each feature.

(C) Bivariate S-F scatter plots for and Networks and non-Networks identified SMNs highlighted.

(D) Trivariate S-F 3D scatter plot of S-F features for Networks in color coded groups.

(E). Representative micrograph of co-immunostained *Drosophila* fat body tissue with 3D Structure and the Mapped / unmapped mtDNA nucleoids projected on MitoGraph binary image; representative Networks and non-Network outlined and magnified.

(F). Ridgeline plots comparing distribution of S-F features of mitochondrial Networks and non-Networks; Median and Coefficient of Variation (in bold) are indicated for each feature.

(G). Relevant bivariate S-F scatter plots for mitochondrial non-networked and networks; predicted SMN subpopulation outlined.

1386   ** signifies p-value < 0.0005; Scale bar: 5 µm; N = sample size.

1387

Figure 7: Schematic illustration of stem cell specification by SMNs. The illustration

summarizes the mechanism of stemness specification as revealed by mito-SinComp S-F

analyses and molecular players identification by coupled scRNA-seq analyses. Stemness

specification induces severing of specific nodes of oxidized HMN (*) to convert them to

characteristic redox-tuned SMNs (outlined) where elevated mt-DNA nucleoid density supports

increased mtDNA gene expression. This establishes the MT-ND1(redox)-KRT15(stemness)

transcriptomic network.

Key Resources Table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rabbit-anti-MT-CYB	Invitrogen	Cat#PA5113926
Mouse-anti-NDUFB3	Santa Cruz Biotechnology	Cat#sc-398196
Rabbit anti-TOMM20	Cell Signaling Technology	Cat#42406S
Mouse-anti-B Actin	Santa Cruz Biotechnology	Cat#sc-47778
Rabbit anti-IgG- HRP Linked	Cell Signaling Technology	Cat#7074P2
Mouse anti-IgG- HRP Linked	Cell Signaling Technology	Cat#7076P2
Rabbit-anti-Phb2	Generated in-house with Boster Bio	N/A
Mouse anti-DNA	Millipore	Cat#CBL186
Cy3 conjugated affinypure goat anti-mouse IgG (H+L)	Jackson Immuno research	115-165-003
Cy™5 AffiniPure® Donkey Anti-Rabbit IgG (H+L)	Jackson Immuno research	711-175-152
Mouse anti-DNA	Abcam	Cat#ab27156
Chemicals, peptides, and recombinant proteins		
2,3,7,8-tetrachlorodibenzo-p-dioxin solution (TCDD)	Merck-Supelco	Cat#48599
MitoTracker Green FM	Invitrogen	Cat#M7514
MitoTracker Deep Red 633	Invitrogen	Cat#M22426
ALDEFLOUR	Stem Cell Technologies	Cat#1700
Tetra methyl rhodamine methyl ester perchlorate (TMRE)	Sigma Aldrich	Cat#T5428
Dulbecco's Modified Eagle Medium (DMEM), High glucose with 4.5 g Glucose per liter, L-Glutamine, Sodium pyruvate and Sodium bicarbonate without Calcium chloride	HI-MEDIA	Cat#AL219A
Lipofectamine 3000	Thermo Fisher	Cat#L3000015

Fetal bovine serum	HI-MEDIA	Cat#RM10681
Pen/Strep	HI-MEDIA	Cat#A001A
Trypsin-EDTA (0.25%)	HI-MEDIA	Cat#TCL049
Ethanol	AR China	Amazon
Toluene	Sigma-Aldrich	Cat#179418
Sodium chloride	Sigma-Aldrich	Cat#S9888
Potassium chloride	Sigma-Aldrich	Cat#P3911
Sodium phosphate dibasic	Sigma-Aldrich	Cat#S9763
Potassium phosphate monobasic	Sigma-Aldrich	Cat#P0662
Fluoromount G	Southern Biotech	Cat#0100–01
Paraformaldehyde (EM grade)	Electron Microscopy Solutions	Cat#15711
Nail polish	MAC	Amazon
Human Insulin	Sigma Aldrich	Cat#I2643
N1 supplement	Sigma Aldrich	Cat#N6530
Human EGF	Sigma Aldrich	Cat#E9644
Human bFGF	Thermo Fisher	Cat#100-18B
L-glutamine	Gibco	Cat#25030149
Penicillin-Streptomycin	Gibco	Cat#15140122
RPMI media	Gibco	Cat#11875093
Sodium Pyruvate	Gibco	Cat#11360070
Grace's Media (Insect Dissecting Medium)	Fisher Scientific	30611031-2
Triton X-100	SRL	64518 (2024271)
BSA	Sigma-Aldrich	A9647
Hoechst-33342	Invitrogen	H3570
2X Laemmli Buffer	BIORAD	#1610737
Deposited data		
Raw, analyzed and meta data of scRNA-seq experiment	Gene expression Omnibus	GEO: GSE285728
Code for mitSinComp Oxidation Function, mtDNA Function, hdWGCNA and ML prediction using Random Forest	This paper	https://github.com/mitralaboratory/CellReports_Scripts/tree/main
Experimental models: Cell lines		

Non-transformed parental HaCaT line (immortalised-normal)	Srivastava et al.[1]	Gift
KRT15[hi] stem cell enriched transformed HaCaT line, derived using 1nM TCDD	Spurlock et al.[2]	-
Control transformed HaCaT line, derived using 10nM TCDD	Spurlock et al.[2]	-
KRT15[hi] stem cell enriched double-stable transformed HaCaT line, derived using 1nM TCDD	Spurlock et al.[2]	-
Control double-stable transformed HaCaT line, derived using 10nM TCDD	Spurlock et al.[2]	-
Control double-stable non-transformed HaCaT line	Spurlock et al.[2]	-
Toluene double stable transformed HaCaT line	Spurlock et al.[2]	-
KRT15[hi] stem cell enriched DRP1 weak knockdown HaCaT line	Spurlock et al.[3]	-
Control DRP1 strong knockdown HaCaT line	Spurlock et al.[3]	-
Non Targeted shRNA HaCaT line	Spurlock et al.[3]	-
A2780-CP ovarian cancer line enriched with ALDH[hi] tumor initiating (stem) cells	Spurlock et al.[2]	Gift
Experimental models: Organisms/strains		
Drosophila melanogaster w[1118]	Das et al.[4]	Gift
Recombinant DNA		
Human DNM1L shRNA1 (plasmid)	Parker et al[5]	Dharmacon Clone: TRCN0000001099
Human DNM1L shRNA2 (plasmid)	Parker et al[5]	Dharmacon Clone: TRCN0000001097
Non-targeted shRNA control (plasmid)	Parker et al[5]	Cat#SHC016
mito-PSmOrange (plasmid)	Spurlock et al.[2]	Gift
mito-roGFP (plasmid)	Addgene	Cat#49437
Software and algorithms		

Python 3.9.12	Python Software Foundation	https://www.python.org
ImageJ 1.53t (FIJI)	Schindelin et al.[6]	https://imagej.nih.gov/ij/index.html
MitoGraph v2.1	Spurlock et al.[2] Viana et al.[7]	https://github.com/vianamp/MitoGraph
Paraview 5.13.0	Ahrens et al.[8]	https://www.paraview.org/
R Version 4.3.3	R Core Team (2024)	https://cran.rstudio.com/
Cell Ranger	10X Genomics	RRID:SCR_017344
Seurat	Hao et al.[9]	RRID:SCR_016341
DBSCAN	Ester et al.[10]	https://scikit-learn.org/stable/modules/generated/sklearn.cluster.DBSCAN.html
Random Forest	Breiman et al.[11]	https://scikit-learn.org/stable/modules/generated/sklearn.ensemble.RandomForestClassifier.html
hdWGCNA	Ji et al.[12]	https://github.com/smorabit/hdWGCNA
Other		
LSM-700 Axio-Observer Confocal Microscope	Carl Zeiss	
LSM 900-Confocal Microscope	Carl Zeiss	
Bright field Microscope	Olympus	
Dissecting Microscope	Carl Zeiss	Stemi 2000
FACS Aria II Flow cytometer	BD Biosciences	
60 mm Cell Culture Dishes laboratory	NEST	705001
TC-treated 6-well cell culture plates	NEST	703001
T-25 flask	NEST	707001
Serological pipettes	Corning	4487

Tissue roll	-	Amazon
Auto-pipette tips	Fisher Scientific	-
Disposable filter units	Thermo Scientific	564–0020
Falcons, 15 mL and 50 mL	Corning	430055, 430829
Coverslip	Fisher Scientific	22X60-1; #062915-9
Syringe	DispoVan sterilized 50 mL	Amazon
MF-Millipore membrane filter, 0.22 μm pore size	Merck	GSWP04700
Cell counting slides for TC20 cell counter, dual chamber	Bio-Rad	1450015
Nunc Lab-Tek chamber slides (convenient but expensive, simple glass slides with coverslips can also be used after optimization)	LabTek-II	154534
Nunc Lab-Tek chambered cover glass	LabTek-II	Z734853
Autoclave	Tomy-SX-700	-
Shaker rocker	Rocking shaker II, BOEKEL	-
Class-II bio-safety hood with laminar flow	Microfilt	-
Auto-pipettes	Eppendorf	-
Incubator	Eppendorf	-
Cell counter TC20	Bio-Rad	-
Dry bath	Lab Armor	-
Pipette gun	Drummond Scientific	-
Fluid aspiration system	Vacuubrand-BVC professional	-
Centrifuge 5425/5425 R - microcentrifuge	Eppendorf	-
Geltrex	Gibco	A14133-02
Dumont #5 Forceps	Fine Science Technologies	11251-20
MultiDish 8 Well Square, 24x33mm wells	Electron Microscopy Sciences	70443
Fly Vials	Fisher Scientific	AS-515
Fly Conicals	Fisher Scientific	AS-355
Fly Flugs Vials	Fisher Scientific	AS-273
Fly Flugs Conical	Fisher Scientific	AS-277

References

1. Srivastava, R.K., Li, C., Weng, Z., Agarwal, A., Elmets, C.A., Afaq, F., and Athar, M. (2016). Defining cutaneous molecular pathobiology of arsenicals using phenylarsine oxide as a prototype. Sci Rep *6*. https://doi.org/10.1038/SREP34865,.

2. Spurlock, B., Gupta, P., Basu, M.K., Mukherjee, A., Hjelmeland, A.B., Darley-Usmar, V., Parker, D., Foxall, M.E., and Mitra, K. (2019). New quantitative approach reveals heterogeneity in mitochondrial structure–function relations in tumor-initiating cells. J Cell Sci *132*. https://doi.org/10.1242/jcs.230755.

3. Spurlock, B., Parker, D., Basu, M.K., Sajina, G., Liu, S., Siegal, G.P., Gunter, A., Moran, A., and Mitra, K. (2021). Fine-tuned repression of Drp1-driven mitochondrial fission primes a "stem/progenitor-like state" to support neoplastic transformation. Elife *10*. https://doi.org/10.7554/ELIFE.68394.

4. Das, S., Hegde, S., Wagh, N., Sudhakaran, J., Roy, A.E., Deshpande, G., and Ratnaparkhi, G.S. (2024). Caspar specifies primordial germ cell count and identity in Drosophila melanogaster. Elife *13*. https://doi.org/10.7554/ELIFE.98584,.

5. Parker, D.J., Iyer, A., Shah, S., Moran, A., Hjelmeland, A.B., Basu, M.K., Liu, R., and Mitra, K. (2015). A new mitochondrial pool of cyclin E, regulated by Drp1, is linked to cell-density-dependent cell proliferation. J Cell Sci *128*, 4171–4182. https://doi.org/10.1242/JCS.172429/260130/AM/A-NOVEL-MITOCHONDRIAL-POOL-OF-CYCLIN-E-REGULATED.

6. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: An open-source platform for biological-image analysis. Nat Methods *9*, 676–682. https://doi.org/10.1038/NMETH.2019,.

7. Viana, M.P., Lim, S., and Rafelski, S.M. (2015). Quantifying mitochondrial content in living cells (Elsevier Ltd) https://doi.org/10.1016/bs.mcb.2014.10.003.

8. Ahrens, J., Geveci, B., and Law, C. (2005). ParaView: An End-User Tool for Large-Data Visualization. Visualization Handbook, 717–731. https://doi.org/10.1016/B978-012387582-2/50038-1.

9. Hao, Y., Stuart, T., Kowalski, M.H., Choudhary, S., Hoffman, P., Hartman, A., Srivastava, A., Molla, G., Madad, S., Fernandez-Granda, C., et al. (2024). Dictionary learning for integrative, multimodal and scalable single-cell analysis. Nat Biotechnol *42*, 293–304. https://doi.org/10.1038/S41587-023-01767-Y,.

10. Ester, M., Kriegel, H.-P., Sander, J., and Xu, X. (1996). A Density-Based Algorithm for Discovering Clusters in Large Spatial Databases with Noise.

11. Breiman, L. (2001). Random forests. Mach Learn *45*, 5–32.

12. Ji, A.L., Rubin, A.J., Thrane, K., Jiang, S., Reynolds, D.L., Meyers, R.M., Guo, M.G., George, B.M., Mollbrink, A., Bergenstråhle, J., et al. (2020). Multimodal Analysis of Composition and Spatial Architecture in Human Squamous Cell Carcinoma. Cell *182*, 497-514.e22. https://doi.org/10.1016/j.cell.2020.05.039.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig.5

Fig. 6

Fig. 7