# Population genomics reveals an ancient origin of heartworms in canids

Rosemonde I. Power[a], Swaid Abdullah[b], Heather S. Walden[c], Guilherme G. Verocai[d], Tiana L.

Sanders[d], Joe L. Luksovsky[d], Andrew R. Moorhead[e], Michael T. Dzimianski[f], Jeremy M.

Foster[g], Michelle L. Michalski[h], Alicia Rojas[i,j], Samuel C. Chacón[k], Georgiana Deak[l], Andrei

D. Mihalca[l], Patrizia Danesi[m], Elias Papadopoulos[n], Piyanan Taweethavonsawat[o,p], Dung Thi

Bui[q,r], Anh Do Ngoc[s], Reuben S.K. Sharma[t], Simon Y.W. Ho[u], Stephen R. Doyle[v,*+] and Jan

Šlapeta[a,w,*+]

[a] Sydney School of Veterinary Science, Faculty of Science, The University of Sydney, New

South Wales, Australia

[b] University of Queensland, School of Veterinary Science, Gatton, Queensland, Australia

[c] Department of Comparative, Diagnostic, and Population Medicine, College of Veterinary

Medicine, University of Florida, Florida, USA

[d] Department of Veterinary Pathobiology, College of Veterinary Medicine & Biomedical

Sciences, Texas A&M University, Texas, USA

[e] Department of Clinical Sciences, College of Veterinary Medicine, North Carolina State

University, Raleigh, North Carolina

[f] Department of Infectious Diseases, College of Veterinary Medicine, University of Georgia,

Athens, Georgia, USA

[g] New England Biolabs Inc., Ipswich, Massachusetts, USA

[h] University of Wisconsin Oshkosh, Oshkosh, Wisconsin, USA

[i] Laboratory of Helminthology, Faculty of Microbiology, University of Costa Rica, San José,

Costa Rica

[j] Centro de Investigación en Enfermedades Tropicales, University of Costa Rica, San José,

Costa Rica

[k] Department of Veterinary Parasitology, Veterinary Sciences, Healthy Pet Veterinary Hospital

SC, David, Chiriquí, Panamá

[l] Department of Parasitology and Parasitic Diseases, University of Agricultural Sciences and

Veterinary Medicine of Cluj-Napoca, Calea Mănăștur 3-5, Cluj-Napoca, Romania

[m] SCS3 - Parasitology and Mycology Unit, Istituto Zooprofilattico Sperimentale delle Venezie,

Viale dell'Università 10, Legnaro, Italy

[n] Laboratory of Parasitology and Parasitic Diseases, Faculty of Veterinary Medicine, Aristotle

University of Thessaloniki, Thessaloniki, Greece

[o] Parasitology Unit, Department of Veterinary Pathology, Faculty of Veterinary Science,

Chulalongkorn University, Bangkok, Thailand

[p] Biomarkers in Animals Parasitology Research Unit, Chulalongkorn University, Bangkok,

Thailand

[q] Department of Parasitology, Institute of Ecology and Biological Resources, Vietnam

Academy of Science and Technology, Hanoi, Vietnam

[r] Faculty of Ecology and Biological Resources, Graduate University of Science and

Technology, Vietnam Academy of Science and Technology, Hanoi, Vietnam

[s] Department of Medical Parasitology, Vietnam Military Medical University, Hanoi, Vietnam

[t] Faculty of Veterinary Medicine, Universiti Putra Malaysia, Serdang, Selangor, Malaysia

[u] School of Life and Environmental Sciences, Faculty of Science, The University of Sydney,

New South Wales, Australia

[v] Wellcome Sanger Institute, Hinxton, Cambridgeshire, UK

[w] Sydney Infectious Diseases Institute, The University of Sydney, New South Wales, Australia

+ These authors contributed equally to this work.

* Corresponding authors.

E-mail address: jan.slapeta@sydney.edu.au (J. Šlapeta); stephen.doyle@sanger.ac.uk (S.

Doyle)

## Abstract

Heartworms (*Dirofilaria immitis*) are parasitic nematodes that cause significant cardiopulmonary-associated morbidity and mortality in canids worldwide. The global dissemination of heartworms is believed to have occurred alongside the dispersal of domesticated dogs. To test this theory, we performed the largest population genetics study of heartworms to date, based on whole-genome sequencing of 127 modern adult individuals collected from mammalian carnivore hosts across four continents. Population structure and demographic analyses of the nuclear genome reveal distinct genetic differences between heartworms from different continents, indicating a deeper ancient origin and dispersal in canid hosts than previously recognised. Using admixture analyses, we find an Asian origin for Australian heartworms consistent with the arrival of dingoes thousands of years ago. Finally, the genetic relatedness between European and Central American heartworms suggests that modern dispersal, likely associated with human colonisation of the Americas by Europeans, occurred with domesticated dogs. Our findings shed light on the global population dynamics and evolutionary history of heartworms, which can aid future surveillance and control efforts for this important veterinary parasite.

## Introduction

The heartworm *Dirofilaria immitis* is a parasitic nematode that infects a broad range of mammalian carnivores (order Carnivora) across tropical, subtropical and temperate regions worldwide [1]. Mosquitoes transmit larval stages of heartworms, which develop after infection and reside in their host's heart and pulmonary arteries as adults [2]. These adult worms can cause lesions, vascular damage, pulmonary hypertension, and right-sided congestive heart failure, potentially leading to the death of their host [2]. This is of particular concern for domestic dogs, whereby potentially life-threatening cardiopulmonary disease is globally managed by lifelong and rigorous application of parasiticides [3]. Such treatment regimes have led to the emergence of drug resistance [4,5] that, while currently only well described in the USA, threatens heartworm control globally if it were to spread. Heartworms are predicted to increase in prevalence and expand into new regions due to climate change, pet travel, and habitat alterations [6]. Therefore, defining the origins, distribution, and capacity for transmission will have important implications for implementing effective heartworm surveillance and control.

Current views on the evolutionary origins and subsequent radiation of heartworms worldwide are inherently anthropocentric. Heartworms are believed to have originated in Europe [7] or Asia [2] and spread globally over the past 200–400 years via human-mediated dog movements. As a result of these recent dispersals, heartworms are presumed to form a homogeneous global population. However, these views lack robust empirical support and overlook the deep co-evolutionary history that some parasites have shared with their hosts and environments. The broad host range of heartworm potentially indicates a long co-evolutionary history with mammalian carnivores, especially canids [8,9], a group that dates back approximately 40 Mya [10], predating humans (~2.8 Mya) [11] and the domestication of modern dogs (14–40 kya) [12–14]. Given also that mosquito vectors have existed since the early Triassic (217 Mya) [15], it is possible that heartworms originated and spread long before human involvement. We propose

that heartworms offer a compelling case for assessing the degree to which anthropogenic

activities have influenced parasite distribution, with a potentially much deeper evolutionary

history than previously recognised.

Analyses of genomic data have significant potential to identify the historical origins of

heartworms and infer their global dispersal routes, as demonstrated for other helminths [16–18].

Sequencing adult heartworms is recommended for obtaining high-quality data [19]; however,

obtaining adults from the heart requires invasive surgery or post-mortem, which are often not

feasible. Such sampling is especially difficult in parts of Asia and Africa, where cultural

perceptions of dogs as unclean or carriers of disease can restrict sampling opportunities. In

light of these challenges, here we present the most extensive population genomics study of

heartworms to date. Using whole genome sequencing of 127 adult worms obtained from nine

countries on four continents, we analyse the diversity, genetic structure, and population

demographics of modern heartworms to infer their historical origins and dispersal. Our

analyses suggest that heartworm diversity has been shaped by the distribution of ancestral canid

hosts throughout time and provide evidence of recent human-driven migration.

## Results and Discussion

### Genomic diversity of heartworms from multiple continents, countries, and hosts

We generated whole-genome sequencing data for 127 adult heartworms (*Dirofilaria immitis*) sampled from nine countries in five regions: Australia, North America (USA), Central America (Costa Rica, Panama), Europe (Greece, Italy, Romania), and Asia (Malaysia, Thailand) (Fig. 1a). Samples were primarily collected from domestic dogs ($n = 115$), but we also sampled cats ($n = 4$), foxes ($n = 4$), a ferret ($n = 1$), a golden jackal ($n = 1$), a leopard ($n = 1$), and a wildcat ($n = 1$). Overall, we obtained an average coverage per sample of 75.47× for the nuclear genome, 13,119.80× for the mitochondrial genome, and 1,047.09× for the *Wolbachia* endosymbiont genome (Supplementary Table 1, Supplementary Fig. 1–4). Variant calling, joint genotyping, and filtering of heartworm samples revealed 264,558 nuclear single nucleotide polymorphisms (SNPs) (3.3 SNPs/kb), 32 mitochondrial SNPs (2.3 SNPs/kb), and 280 *Wolbachia* SNPs (0.3 SNPs/kb). The SNP frequencies in the nuclear and mitochondrial genomes are comparable, whereas the frequency in the *Wolbachia* genome is considerably lower. These frequency data are comparable to those of other *Wolbachia*-infected filarial worms, such as *Onchocerca volvulus* [20] and *Wuchereria bancrofti* [18].

### Nuclear variants reveal distinct continental partitioning of heartworms

Analyses of the heartworm cohort's genetic diversity revealed distinct groups of samples based on broad geographical regions. Principal component analysis (PCA) using 196,550 high-quality autosomal SNPs revealed four clusters of samples corresponding to the continents from which the samples were collected (Fig. 1b and Extended Data Fig. 1; PC1 variance: 12.6%, PC2 variance: 10.69%). The third and fourth PCs showed a similar pattern of geographical clustering (Extended Data Fig. 2a; PC3 variance: 5.76%, PC4 variance: 4.39%);

however, although the signal was comparatively weaker, the European and Central American

samples formed separate clusters. Similar findings were observed in a previous study of 31

heartworm samples from Australia, Italy, and the USA, where samples from each country

formed distinct clusters [21]. Independent PCAs of SNPs from each chromosome found the same

continental clustering pattern across all chromosomes, including the sex-linked X chromosome

(Extended Data Fig. 2b–f), suggesting that this is a genome-wide rather than region-specific

observation. The distinct geographical partitioning in our nuclear data was not supported by

the mitochondrial (Extended Data Fig. 3a) or *Wolbachia* data (Extended Data Fig. 3b), likely

due to the few variant sites within each dataset.

Analyses of genome-wide nucleotide diversity (Pi) further demonstrated differences

between heartworm populations across continents. Pi differed significantly between all

populations (Wilcoxon rank sum test, $p < 0.001$), with the exception of Europe and Central

America (Fig. 1c). The highest Pi was observed in the USA (Pi[mean] = $7.39\times10^{-4}$), while the

lowest was in Asia (Pi[mean] = $2.89\times10^{-4}$). The genetic diversity of our USA samples aligned

closely with that of a previous *D. immitis* study (Pi[mean] = $7.36\times10^{-4}$) [21]. However, our

Australian samples (Pi[mean] = $5.61\times10^{-4}$) exhibited higher genetic diversity than those

reported previously (Pi[mean] = $3.9\times10^{-5}$) [21], likely due to the larger and geographically

broader sample here. Overall, the nucleotide diversity of our *D. immitis* samples is comparable

to that of the filarial worm *W. bancrofti* (Pi[mean] = $2.4\times10^{-4}$) [18] but lower than that of *O.*

*volvulus* (Pi[mean] = $4.0\times10^{-3}$) [20]. Each *D. immitis* population exhibited different rates of

linkage disequilibrium (LD) decay and baseline LD (Fig. 1d), with Asia and the USA showing

the highest and lowest LD, respectively. Estimates of admixture provided further support for

continental partitioning in heartworms. To infer the ancestry proportions of samples, we used

NGSadmix with a range of *K* clusters (*K* = 2–10; Extended Data Fig. 4). At *K* = 6, Australian

samples showed the most diverse admixture patterns with strong evidence of shared ancestry

with Asia, whereas USA, Europe, and CENAM were largely distinct with some evidence of

shared ancestry in a minority of European and CENAM samples (Fig. 1e). The strong

population structure in the nuclear genome of heartworms suggests that spatial spread has been

restricted during the species' history, resulting in distinct genetic profiles on each continent.

**Inference of transmission between diverse heartworm hosts**

Although most heartworm samples were obtained from dogs, the analysis of heartworm

genomes from foxes, cats, a golden jackal, a ferret, and a leopard allowed us to evaluate genetic

variation among different host species. We found that geography rather than the host species

explained variation in the global heartworm population, evidenced by samples from cat and

ferret nested within samples from dogs in the USA, fox and dogs in both Australian and Italian

groups, and dogs and jackal, wildcat and leopard from Europe (Fig. 2a). Analyses of nucleotide

diversity (Pi), absolute nucleotide divergence ($D_{XY}$), and genetic differentiation ($F_{ST}$) provided

additional insights on the variation within and between different host species. The ferret

differed from other hosts as its heartworm had the highest diversity (Pi[mean] = $1.21 \times 10^{-3}$)

(Fig. 2b) and the greatest divergence from other hosts (Extended Data Fig. 5a). In contrast, the

samples from the golden jackal, leopard, and wildcat had the lowest diversity (Pi[mean] 1.10–

$1.89 \times 10^{-4}$) and were the least divergent from each other compared with all other host pairs.

These hosts, however, were each represented by only a single worm, and the variation observed

in the jackal, leopard, and wildcat samples may be confounded as they all originated from

Romania. There was little genetic differentiation between heartworms from dogs and ferrets,

and between dogs and foxes, suggesting that heartworms from canids and mustelids may be

more genetically similar to each other than to those from felids (Extended Data Fig. 5b).

Collectively, these data suggest an absence of obvious host-specificity and that transmission is

likely between domestic animals and wildlife. However, further geographic sampling from

non-dog hosts is needed to exclude the possibility of host-specific adaptation.

**Distinguishing modern from ancient origins and dispersal**

The current hypothesis for heartworm distribution around the world is that it has been

largely influenced by human movement in contemporary times. Alternative hypotheses are that

heartworms spread with humans during or shortly after dog domestication ~14–40 kya, or that

heartworms spread with wild canids prior to domestication. To further explore the demographic

history of heartworms, we inferred their effective population size histories and population split

times using SMC++. Effective population sizes were smaller than estimates of past population

sizes for all continents (Fig. 3a). Earlier population splits in the demographic analysis broadly

supported the $D_{XY}$ and $F_{ST}$ results (Extended Data Fig. 6). Notably, the earliest estimated

population splits occurred in Asia, which split from the populations from Europe, the USA,

and Central America between ~46 and 63 kya. Importantly, these data suggest that the

divergence between populations is older than the proposed dates for dog domestication [12–14].

Considering a pre-domestication dispersal hypothesis, we next questioned which host

may have harboured and dispersed heartworms before the emergence of modern dogs. We

hypothesised that longer co-evolution between parasites and hosts would likely result in the

latter being able to carry higher parasite loads. Felids (e.g. domestic cats, wild cats, snow

leopards), procyonids (e.g. raccoon dogs), mustelids (e.g. European badger, ferrets), pinnipeds

(e.g. harbour seal, brown fur seal), and ursids (e.g. brown bear, black bear) have been described

as being infected with heartworms (Supplementary Table 2); however, they typically carry

fewer adult worms and fail to support the parasite's entire life cycle, and so they are not

considered primary reservoirs [22–25] (Fig. 3b). In contrast, wild canids are commonly

microfilaremic and have high worm burdens. Wolves and coyotes have worm burdens similar to dogs, while jackals and foxes have significantly lower burdens than dogs (Kruskal-Wallis test with Dunn's multiple comparisons; Dog vs Jackal: $P \leq 0.05$, Dog vs Fox: $P \leq 0.001$) (Fig. 3b). These findings imply that canids, particularly ancestral wolves or coyotes, were most likely the primary hosts for heartworm during ancient times [26].

**Coevolution of canids and dogs from ancient to contemporary times**

Previous work had proposed that canids originated in North America ~40 Mya when the continent was completely isolated [10], whereas mosquitoes originated much earlier in the Triassic (~217 Mya) [15]. Phylogenetic analyses of mitochondrial DNA suggest that heartworms diverged from their closest relatives in the genus *Onchocerca* ~25 Mya during the Oligocene [27], i.e., when canids and mosquito vectors were already well-established. Together, this timeline supports a long co-evolutionary history between heartworms and ancient canids.

The formation of the Bering Land Bridge and the Isthmus of Panama during the Miocene and Pliocene facilitated the first migrations of canids into Eurasia (~7 Mya) [10] and South America (3.5–3.9 Mya) [28], respectively. Throughout history, fluctuations in sea level have caused the Bering Land Bridge to be repeatedly exposed and submerged. Fossil records and genomic data provide robust evidence that these periods of exposure allowed ancient canids to form continuous populations across the Northern Hemisphere. Examples include Pleistocene wolves, which were highly connected across North America and Eurasia [12,29,30], *Xenocyon lycaonoides,* a large canid distributed across North America and Eurasia in the mid Pleistocene [31] and *Eucyon,* a jackal-sized canid distributed across North America, Eurasia, and Africa in the late Miocene [32]. The existence of these interconnected ancient canid populations

suggests that heartworms may have formed a highly connected population with their hosts at
that time.

If heartworms were once one population, when did they diverge, and why? We
estimated the timing of this contiguous *D. immitis* population to predate ~63 kya before the
species diverged on each continent (Fig. 3a). This roughly aligns with the last interglacial
period (116 – 130 kya), when Earth experienced a climate as warm or even warmer than today
[33]. Such an environment could have been suitable for the development and transmission of
heartworms via mosquito vectors. Following the warm climates of the last interglacial period,
the Earth began to cool. By Marine Isotope Stage 4 (58–72 kya), large continental ice sheets
had formed across North America, Beringia, and northern Eurasia [34]. These ice sheets separated
ancient canid populations [29,35], potentially fragmenting the associated ancient heartworm
population. This fragmentation would have reduced gene flow for extended periods, explaining
our study's continental grouping of samples. Further insights might be obtained by exploring
the presence of heartworm DNA in the remains of ancient canids; if successfully extracted,
these ancient worm DNA sequences may fill in the temporal and geographic gaps needed to
understand heartworm divergence in more detail. Similarly, further sampling in unstudied
regions including South America and Africa, and finer sampling within Asia, would enhance
our understanding of heartworms' evolutionary history.

Considering global warming and cooling cycles, the fragmentation of canid and
heartworm populations would likely have occurred more than once. We found evidence of
secondary admixture between heartworms from Asia and the USA after the initial population
divergence, evident from both the admixture analyses and gene flow between populations (Fig.
4a; delta m = 3.4; all tree migration edges are presented in Extended Data Fig. 7). *D*-statistics
(Fig. 4b,c) provided further support for excessive allele sharing between the USA and Asia
relative to that between the USA and other regions (Fig. 4b). Given that heartworms from the

USA and Asia diverged >45 kya (Fig. 3a), this admixture may reflect multiple waves of

migration with canids across the Bering Land Bridge in the Late Pleistocene, during periods of

ice sheet contraction [12].

**The origin and spread of heartworms in Australia**

Heartworms are endemic in Australia, but the mechanisms and timing of their arrival

in the country have remained an enigma. Consistent with the modern human dispersal

hypothesis, one means of introduction could have been after European colonisation that began

~235 years ago. However, considering our genetic evidence of older, canid-mediated dispersal,

a more likely scenario is that heartworms were brought to Australia via dingoes. Dingoes are

an ancient dog breed from Asia and were the only canids present on the Australian continent

until European settlement. Recent morphometric analysis indicates an East Asian origin of

dingoes via Melanesia [36], whereas genetic studies suggest that dingoes likely came from Island

Southeast Asia and were brought to Australia by humans between 5 and 12 kya [37].

Multiple lines of evidence support the introduction of heartworms into Australia via

Asia. Both admixture and TreeMix analyses highlighted the close relationship, evidenced by

partial or complete admixture profiles (Fig. 1e) and mutual monophyly (Fig. 4a) between Asian

samples and samples collected along the East Coast of Australia (Fig. 5a). Calculations of $D_{XY}$

showed that AUS-ASIA ($D_{XY}$[mean] = 5.13e–4) was the second least divergent pair (after

EUR-CENAM, which indicates much more recent dispersal pattern [see below]), suggesting

that Australian worms are more closely related to those in Asia than elsewhere (Extended Data

Fig. 6a). $D$-statistics solidified this relationship by showing excessive allele sharing between

the two continents relative to that between Australia and other regions (Fig. 5b). These findings

point towards an Asian origin of Australian heartworms, likely transported with a dingo host,

rather than by an introduction through European colonisation.

We estimated the divergence of Asian and Australian heartworms to be ~29 kya (Fig.

3a), which is older than the previous estimates of 5 and 12 kya for the dingo arrival. This

discrepancy likely arises from the limited geographic scope of our Asian samples, which are

only from Thailand and Peninsular Malaysia. Sampling across a broader geographic region,

particularly throughout the Indonesian archipelago and Melanesia, could bridge the gap

between our Australian and Asian samples. Such data may provide further granularity of the

Asia–Australia divergence and potentially coincide more closely with the timeline of dingo

arrival.

Considering the extensive sampling along the north–south transect of Eastern Australia,

we sought to identify factors that might explain current genetic diversity. Australian

heartworms were the second most genetically diverse population globally (Fig. 1c). Within

Australia, diversity was highest in Lockhart River (Pi[mean] = 8.12e-4) and lowest in

Rockhampton (Pi[mean] = 2.15e-4) (Fig. 5c). Heartworms from the northern state of

Queensland were highly admixed, with two unique sources of ancestry not found on any other

continent (Fig. 1e). Interestingly, ten of 16 heartworms from Sydney, the southernmost site,

had a single fixed ancestry that was detected in Queensland samples. Another four Sydney

heartworms had a different fixed ancestry identical to that found in Asia, potentially reflecting

modern movement, though the exact cause remains unclear. Previously, it was believed that

heartworm cases in temperate Sydney and broader New South Wales were primarily caused by

infected pets or mosquitoes travelling down from tropical and subtropical Queensland [38], where

climates are more favourable to heartworm transmission. However, our nuclear PCA showed

a large cluster of samples primarily from Queensland, alongside two distinct clusters from

Sydney (Fig. 5d). This suggests that Sydney heartworms may not necessarily originate from

Queensland, as previously believed. Further investigations are required to determine whether

there are local endemic populations of heartworms in Sydney. Differences in $D_{XY}$ and $F_{ST}$

between Australian heartworm subpopulations were subtle, showing no clear relationship

between genetic and geographical distance as might be expected for a vector-borne parasite

(Fig. 5e, f).

**Modern heartworm migration between Europe and Central America**

Our genomic analysis revealed a close genetic relationship between worms in Europe

and Central America. This finding was most evident in our PCA of nuclear SNPs, where

samples from Greece, Italy, and Romania clustered with those from Costa Rica and Panama

(Fig. 1b). In addition, estimates of $D_{XY}$ revealed that EUR-CENAM ($D_{XY}$[mean] = 4.95e–4)

was the least divergent compared with other populations. Admixture analysis using NGSadmix

further supported this trend as the European and Central American samples shared high levels

of ancestry at lower $K$ values ($K = 2$, 3, and 5; Extended Data Fig. 4). Although increasing $K$

helped to differentiate worms from these two continents, low levels of admixture were still

present (Fig. 1e). These findings were corroborated using $D$-statistics which provided strong

evidence of excessive allele sharing between heartworms from Central America and Europe

relative to that between Central America and other regions (Fig. 4c). These findings support a

modern migration event between European and Central American heartworms which took

place after dog domestication.

One plausible modern migration event between Europe and Central America was the

transatlantic transportation of domestic animals, including dogs, during the European

colonisation of the Americas beginning in the late 15th century. Animals introduced to the

Americas during the Age of Discovery were primarily livestock species, including cattle,

sheep, pigs, goats, horses, and mules, but dogs such as greyhounds and mastiffs were also

included for hunting, guarding, protection, shepherding, and battle purposes [39–41]. The

introduction of European dogs into the Americas is perhaps best documented from Columbus'

voyages, with historical records indicating the presence of at least 20 dogs on board during the

second voyage to Hispaniola [39]. Therefore, infected dogs from Europe may have transported

local strains of this parasite to the Americas. Alternatively, the shared ancestry between

European and Central American heartworms could result from the more recent migration of

humans and their pets. Given the population split times proposed in our demographic analyses

and the lack of evidence for recent migration and transmission between continents elsewhere,

we posit that the former scenario is more biologically plausible and likely occurred over the

past few centuries.

**Concluding remarks**

Our transcontinental genome-wide analysis of heartworm parasites has revealed a much

deeper origin and dispersal history than previously understood, allowing us to propose new

hypotheses on the evolutionary history of heartworms (Fig. 6). Rather than being solely driven

by human-mediated dispersal of dogs over the past few hundred years, we found that ancestral

canid hosts played a pivotal role in the evolution and dissemination of heartworms around the

world. Heartworms have evolved to exploit a unique niche in the heart and bloodstream of

canids, an environment that has likely remained unchanged over millions of years between

hosts. In the absence of niche competition, heartworms could be a relic of the past, which, until

recently, had little need for further adaptation. Anthropogenic influence is, however, likely to

shape heartworm diversity more rapidly than ever before. For example, extensive drug

treatment is already associated with resistance, particularly in the USA; climate change is

expected to shift populations of parasites and their vectors into higher latitudes; and increasing

global connectivity of people and their pets will all influence this distribution and adaptation

of heartworms. Therefore, understanding the global genomic landscape of these parasites, as

demonstrated here, is vital for developing effective surveillance and control strategies in the

future. Our data indicate that such surveillance and control strategies should account for

geographical variation between parasite populations and be aware of transmission within and

between populations to maximise sustainable parasite control.

## Methods

### Sample collection

A total of 120 individual adult *Dirofilaria immitis* were obtained from Australia ($n = 43$), the USA ($n = 37$), Panama ($n = 14$), Greece ($n = 12$), Thailand ($n = 6$), Italy ($n = 3$), Romania ($n = 3$), Costa Rica ($n = 1$), and Malaysia ($n = 1$). The sampling regime aimed to collect adult *D. immitis* from a wide geographical range. Most of these samples were obtained from dogs (*Canis lupus familiaris*; $n = 115$) but also included some from cats (*Felis catus*; $n = 4$), foxes (*Vulpes vulpes*; $n = 4$), a ferret (*Mustela putorius furo*; $n = 1$), a golden jackal (*Canis aureus*; $n = 1$), a leopard (*Panthera pardus*; $n = 1$), and a wildcat (*Felis silvestris*; $n = 1$) (metadata for all samples is presented in Supplementary Table 3). We included five technical replicates (Brisbane: $n = 1$; Sydney: $n = 2$; Malaysia: $n = 1$; Panama: $n = 1$). To provide outgroup samples for our analyses, we collected adult *D. repens* ($n = 2$) from dogs in Italy, *Dirofilaria* sp. 'Thailand II' ($n = 2$) from humans (*Homo sapiens*) in Vietnam and *D. ursi* ($n = 1$) from an American black bear (*Ursus americanus*) in the USA.

### DNA extraction and whole genome sequencing

*Dirofilaria* specimens were frozen at −20°C or stored in ethanol upon collection and sent to the University of Sydney, Australia. A 1–2 cm segment between the first and second third of the anterior portion of the worms was excised using a sterile scalpel blade and placed in a 1.5 mL Eppendorf tube.

***Ethanol-preserved samples.*** Worm segments were centrifuged at maximum speed for 30 s using a benchtop centrifuge (Eppendorf, Australia) and excess ethanol was removed. The segments were washed in 500 mL of phosphate-buffered saline (PBS) and placed on a heat

block (Heat Block Eppendorf Thermomixer Comfort or Heat Block Major Science Dry Bath

Incubator) at 60°C for 20–40 min, depending on their thickness, to evaporate any remaining

ethanol.

***Frozen samples.*** Worm segments were defrosted and washed three times in PBS.

Genomic DNA was extracted from the tissue segments using a Monarch® Genomic DNA

Purification Kit (New England Biolabs, Australia) as per the manufacturer's protocol. Extracted

DNA was shipped to Novogene (HK) Co., Ltd for indexing, library preparation, and whole

genome sequencing using the NovaSeq 6000 platform. Overall, 128/130 and 2/130 DNA

samples were sequenced at expected depths of ~10 G and ~1 G, respectively.

**Bioinformatic analyses**

The bioinformatics workflow of this study is summarised in Extended Data Fig. 8. In

addition to the data generated in this study, we included publicly available datasets of

individual *D. immitis*. These data were derived from (i) five adults from Sydney, Australia

(SRA NCBI: SRR13154013-7) [42]; (ii) one adult from Georgia, USA (ENA Accession:

ERR034940) and one adult from Pavia, Italy (ENA Accession: ERR034941) [43].

**Raw data processing and mapping**

Multiple FastQ files were generated for some samples and merged before data analysis.

The quality of all raw paired-end sequencing reads was inspected using FastQC (version 0.11.8;

https://www.bioinformatics.babraham.ac.uk/projects/fastqc/), and the output was summarised

using MultiQC (version 1.17 [44]). The raw paired-end reads were scanned with a 10-base sliding

window and trimmed to have a minimum average quality score of 20 per base and a minimum

read length of 50 bp using Trimmomatic [45] (version 0.39). We used a Nextflow [46] mapping

pipeline (mapping-helminth/v1.0.9 [47]) to map the trimmed paired-end reads to a combined

reference genome containing *D. immitis* (dimmitis_WSI_2.2 [21]) and the domestic dog

(GenBank accession: GCA_014441545). The dimmitis_WSI_2.2 genome has been previously

supplemented with the *D. immitis* mitochondrial and *D. immitis*-associated *Wolbachia*

endosymbiont genomes [21]. The mapping pipeline was designed for the standard mapping of

helminth genomes using Minimap2 (version 2.17 [48]) as a mapper and GATK (version 4.1.4.1

[49]), Samtools (version 1.14 [50]), and Sambamba (version 1.0 [51]) for further data processing.

Reads aligning to only the *D. immitis* genome were extracted for downstream analysis using

Samtools.

**Sequencing Coverage**

Sequencing coverage was examined using BamTools (version 2.5.1 [52]), BEDTools

makewindows (version 2.31.0 [53]), and SAMtools bedcov. Coverage statistics were obtained for

the nuclear, mitochondrial, and *Wolbachia* genomes of *D. immitis*. To visualise nuclear *D.*

*immitis* sequencing coverage for each sample, the scaffold, mitochondrial and *Wolbachia*

genomes of *D. immitis* were removed, along with any chromosomes of the domestic dog

genome. The ratio of X chromosome to autosome coverage was visualised to determine the sex

of the *D. immitis* samples, identified by a drop in coverage along the X chromosome (ratio ~

0.5) indicative of a male parasite [21].

**Variant calling**

Variants were identified using GATK (version 4.1.4.1; [49]) HaplotypeCaller,

CombineGVCFs, and GenotypeGVCFs. GVCF files were first produced for each sample and

then combined into a cohort-level GVCF file, which was then used for joint genotyping.

Outgroup samples were removed to obtain a *D. immitis* cohort VCF file. Variants in the

nuclear, mitochondrial, and *Wolbachia* datasets were filtered separately using GATK

SelectVariants, GATK VariantFiltration, and VCFtools (version 1.16 [54]). The distribution of

eight quality metrics was established (QUAL, DP, MQ, SOR, QD, FS, MQRankSum, and

ReadPosRankSum), and the relevant upper and/or lower 1% distribution tails were removed.

The single nucleotide variants (SNVs) were further filtered based on the following parameters:

bi-allelic, depth $\geq$3, Hardy-Weinberg Equilibrium p-value $\geq 10^{-6}$ (nuclear), and minor allele

frequency $\geq$0.02. Indels were not included in subsequent analyses. Samples with <50% of the

total variants were removed, and an optimal threshold for missing data of 0.9 was established

(nuclear, mitochondrial, and *Wolbachia*). Additional VCF files were generated from the

filtered nuclear data for autosomes (chromosomes 1–4), the sex-linked X chromosome, and

Australian samples. The same filtering parameters were applied to the VCF file containing both

*D. immitis* and outgroup samples. SNP discordance in the filtered nuclear autosome data for

the technical replicate samples was measured using VCFtools --diff-indv-discordance. All

technical replicates had < 0.5% SNP discordance with their respective original samples.

**Population structure**

To explore the broad-scale genetic relatedness between *D. immitis* individuals and

populations, we performed principal component analysis (PCA) using the R package

SNPRelate (version 1.38.0; [55]). PCA was conducted for the nuclear (autosomes together,

autosomes separate, and chromosome X separate), mitochondrial, and *Wolbachia* data.

Nucleotide diversity (Pi) within populations, as well as absolute nucleotide divergence

($D_{XY}$) and genetic differentiation ($F_{ST}$) between populations, were calculated using pixy

(version 1.2.10.beta2; [56]). First, the GVCF files were genotyped using GATK GenotypeGVCFs

with the inclusion of invariant sites (--all-sites). The nuclear invariants were selected using

GATK SelectVariants (--select-type-to-include NO_VARIATION) and merged with the

previously filtered nuclear SNPs. The VCF file was filtered to only include invariants and SNPs

in the autosomes and sex-linked X chromosome. We conducted the pixy analysis with a

window size of 100 kb. Separate runs were performed, grouping the samples in various ways:

by region (using dog host samples only), by host species (using all samples), and by city/state

(using Australian dog host samples only).

Linkage disequilibrium (LD) statistics were obtained from the nuclear data using

VCFtools. First, we filtered and thinned the data (--maf 0.02 --thin 1000 --max-missing 1). We

then calculated the squared correlation coefficient between genotypes (--ld-window-bp

1000000 --max-alleles 2 --min-alleles 2 --min-r2 0.1 --geno-r2). To visualise patterns of LD

decay, the output was plotted per 0.1 kb for each population.

**Admixture**

The global ancestry and admixture proportions of *D. immitis* were inferred using

NGSadmix from the ANGSD package [57]. First, genotype likelihoods were extracted from the

filtered autosomal *D. immitis* VCF file using VCFtools (--max-missing 1 --BEAGLE-PL).

Genotype likelihoods were extracted separately for each chromosome and merged into a single

dataset. NGSadmix (-minMaf 0.02 -misTol 0.9) was performed using multiple combinations

of different clusters ($K$ = 2 to 10) and seeds (1 to 5). CLUMPAK was used to determine the

optimal $K$ value [58].

We used TreeMix (version 1.12; [59]) to produce a maximum-likelihood tree and infer

population splits and admixture events in the *D. immitis* dataset. To generate the input data,

VCFtools was used to filter the cohort SNV VCF file (--max-missing 1), retaining only

autosomal data and the same samples as in the previous variant-calling filtration step. The

variants were pruned for linkage disequilibrium (10 kb windows, shifting by 10 kb, LD

threshold 0.1), and the VCF file was converted to the appropriate format using custom scripts

('ldPruning.sh'              and              'vcf2treemix.sh')              obtained              from

https://github.com/speciationgenomics/scripts. TreeMix was run across a range of migration

edges (0 to 5) and seeds (0 to 10), with *D. ursi* included as an outgroup. The optimal number

of migration edges was estimated using the R package OptM (version 1.8;

https://github.com/cran/OptM).

To further investigate admixture in the cohort, *D*-statistics were inferred using

AdmixTools (version 5.1 [60]). First, we converted our filtered nuclear VCF file into the required

eigenstrat format using a custom script ('convertVCFtoEigenstrat.sh') obtained from

https://github.com/speciationgenomics/scripts/. Additional tools required for file conversion

were VCFtools and PLINK (version 1.90b6.18 [61]). qpDstat was used to perform *D*-statistic

testing for all population pairs relative to the other populations (PopX), with *D. ursi* as the

outgroup taxa. *Z*-scores with an absolute value >3 were considered significant.

**Population demography**

We estimated the population size history of *D. immitis* and divergence times between

subpopulations using SMC++[71] (version 1.15.3 [62]). Samples from dog hosts in the filtered

autosomal VCF were grouped by continent (*i.e.*, Asia, Australia, Central America, Europe, and

USA) and analysed separately before being plotted together. First, the VCF file was converted

to the SMC++ input format using *vcf2smc*. The model was fitted using *estimate* with a

timepoint range of 1 to 5,000,000 and the nematode *Caenorhabditis elegans* mutation rate of

504 $2.7 \times 10^{-9}$ per site per generation. The *plot* command (-g -c) was used to visualise the output,

505 with the -g parameter specifying the generation time of the target population. *D. immitis* takes

506 ~1 year to produce offspring and can live for up to 7 years in the host's cardiopulmonary

507 system. The development and transmission of this parasite are also highly dependent on local

508 climate and vector availability. To account for potential variability in the generation time of *D.*

509 *immitis*, we tested and compared multiple generation times (-g 1, 2, 4, or 6) (Extended Data

510 Fig. 9). Given the reliance on viable mosquito vectors for transmission, and the presence of

511 both male and female adult worms for sexual reproduction, we determined that a 4-year interval

512 was the most appropriate estimate. In addition, we used *split* to estimate divergence times

513 between all pairs of subpopulations.

## Data availability

516  Raw FastQ sequences generated in this study are available at SRA NCBI BioProject

517 PRJNA1104412.

## Code availability

520  The code used to analyse the data in this study is available from GitHub:

521 https://github.com/rosemondepower/heartworm_genome.git.

## References

1. Noack, S., Harrington, J., Carithers, D. S., Kaminsky, R. & Selzer, P. M. Heartworm disease - Overview, intervention, and industry perspective. *Int. J. Parasitol. Drugs Drug Resist.* **16**, 65–89 (2021).

2. Bowman, D. D. & Atkins, C. E. Heartworm biology, treatment, and control. *Vet. Clin. North Am. Small Anim. Pract.* **39**, 1127–58, vii (2009).

3. Nelson, T., McCall, J. W., Moorhead, A., Starkey, L. & Ames, M. *American Heartworm Society Canine Guidelines for the Prevention, Diagnosis, and Management of Heartworm (Dirofilaria Immitis) Infection in Dogs.* https://d3ft8sckhnqim2.cloudfront.net/images/AHS_Canine_Guidelinesweb22NOV2024.pdf?1732318144 (2024).

4. Hampshire, V. A. Evaluation of efficacy of heartworm preventive products at the FDA. *Vet Parasitol* **133**, 191–195 (2005).

5. Pulaski, C. N. *et al.* Establishment of macrocyclic lactone resistant *Dirofilaria immitis* isolates in experimentally infected laboratory dogs. *Parasit. Vectors* **7**, 494 (2014).

6. Simón, F. *et al.* Human and animal dirofilariasis: the emergence of a zoonotic mosaic. *Clin. Microbiol. Rev.* **25**, 507–544 (2012).

7. Roncalli, R. Tracing the history of heartworms: A 400 year perspective. in *Recent Advances in Heartworm Disease: AHS Symposium* (1998). doi:10.5555/20053201369.

8. Moroni, B. *et al.* Dirofilaria immitis in wolves recolonizing northern Italy: are wolves competent hosts? *Parasit. Vectors* **13**, 482 (2020).

9. Aher, A. M. *et al.* Prevalence, genetic analyses, and risk factors associated with heartworm (*Dirofilaria immitis*) in wild coyotes (*Canis latrans*) from Florida, USA. *J. Wildl. Dis.* **52**, 785–792 (2016).

10. Wang, X. & Tedford, R. H. *Dogs: Their Fossil Relatives and Evolutionary History.*

547   (Columbia University Press, New York, NY, 2010).

548 11. Villmoare, B. *et al.* Paleoanthropology. Early Homo at 2.8 Ma from Ledi-Geraru, afar,

549    Ethiopia. *Science* **347**, 1352–1355 (2015).

550 12. Bergström, A. *et al.* Grey wolf genomic history reveals a dual ancestry of dogs. *Nature*

551    **607**, 313–320 (2022).

552 13. Freedman, A. H. *et al.* Genome sequencing highlights the dynamic early history of dogs.

553    *PLoS Genet.* **10**, e1004016 (2014).

554 14. Botigué, L. R. *et al.* Ancient European dog genomes reveal continuity since the Early

555    Neolithic. *Nat. Commun.* **8**, 16082 (2017).

556 15. Soghigian, J. *et al.* Phylogenomics reveals the history of host use in mosquitoes. *Nat.*

557    *Commun.* **14**, 6252 (2023).

558 16. Sallé, G. *et al.* The global diversity of *Haemonchus contortus* is shaped by human

559    intervention and climate. *Nat. Commun.* **10**, 4811 (2019).

560 17. Doyle, S. R. *et al.* Population genomics of ancient and modern *Trichuris trichiura*. *Nat.*

561    *Commun.* **13**, 3888 (2022).

562 18. Small, S. T. *et al.* Human Migration and the Spread of the Nematode Parasite

563    *Wuchereria bancrofti*. *Mol. Biol. Evol.* **36**, 1931–1941 (2019).

564 19. Power, R. I., Doyle, S. R. & Šlapeta, J. Whole genome amplification and sequencing of

565    individual *Dirofilaria immitis* microfilariae. *Exp. Parasitol.* **263-264**, 108806 (2024).

566 20. Choi, Y.-J. *et al.* Genomic diversity in *Onchocerca volvulus* and its *Wolbachia*

567    endosymbiont. *Nat Microbiol* **2**, 16207 (2016).

568 21. Gandasegui, J. *et al.* Genome structure and population genomics of the canine

569    heartworm *Dirofilaria immitis*. *Int. J. Parasitol.* (2023)

570    doi:10.1016/j.ijpara.2023.07.006.

571 22. Penezić, A., Selaković, S., Pavlović, I. & Ćirović, D. First findings and prevalence of

adult heartworms (*Dirofilaria immitis*) in wild carnivores from Serbia. *Parasitol. Res.*

**113**, 3281–3285 (2014).

23. Ionică, A. M., Deak, G., Boncea, R., Gherman, C. M. & Mihalca, A. D. The European

badger as a new host for *Dirofilaria immitis* and an update on the distribution of the

heartworm in wild carnivores from Romania. *Pathogens* **11**, 420 (2022).

24. McCall, J. W., Genchi, C., Kramer, L. H., Guerrero, J. & Venco, L. Heartworm disease

in animals and humans. *Adv. Parasitol.* **66**, 193–285 (2008).

25. Molnár, V. *et al.* Autochthonous *Dirofilaria immitis* infection in a ferret with aberrant

larval migration in Europe. *J. Small Anim. Pract.* **51**, 393–396 (2010).

26. vonHoldt, B. M. *et al.* Whole-genome sequence analysis shows that two endemic

species of North American wolf are admixtures of the coyote and gray wolf. *Sci. Adv.* **2**,

e1501714 (2016).

27. Small, S. T., Tisch, D. J. & Zimmerman, P. A. Molecular epidemiology, phylogeny and

evolution of the filarial nematode *Wuchereria bancrofti*. *Infect. Genet. Evol.* **28**, 33–43

(2014).

28. Chavez, D. E. *et al.* Comparative genomics uncovers the evolutionary history,

demography, and molecular adaptations of South American canids. *Proc. Natl. Acad.*

*Sci. U. S. A.* **119**, e2205986119 (2022).

29. Leonard, J. A. *et al.* Megafaunal extinctions and the disappearance of a specialized wolf

ecomorph. *Curr. Biol.* **17**, 1146–1150 (2007).

30. Hofreiter, M. Pleistocene extinctions: haunting the survivors. *Curr. Biol.* **17**, R609–11

(2007).

31. Jiangzuo, Q. *et al.* Middle Pleistocene *Xenocyon lycaonoides* Kretzoi, 1938 in

northeastern China and the evolution of Xenocyon-Lycaon lineage. *Hist. Biol.* 1–13

(2022).

32. Rook, L. The wide ranging genus Eucyon Tedford & Qiu, 1996 (Mammalia, Carnivora, Canidae, Canini) in the Mio-Pliocene of the Old World. **31**, 723–741 (2009).

33. Kukla, G. J. *et al.* Last interglacial climates. *Quat. Res.* **58**, 2–13 (2002).

34. Batchelor, C. L. *et al.* The configuration of Northern Hemisphere ice sheets through the Quaternary. *Nat. Commun.* **10**, 3713 (2019).

35. Koblmüller, S. *et al.* Whole mitochondrial genomes illuminate ancient intercontinental dispersals of grey wolves (*Canis lupus*). *J. Biogeogr.* **43**, 1728–1738 (2016).

36. Koungoulos, L. G., Hulme-Beaman, A., Fillios, M. & Willandra Lakes Region World Heritage Aboriginal Advisory Group. Phenotypic diversity in early Australian dingoes revealed by traditional and 3D geometric morphometric analysis. *Sci. Rep.* **14**, 21228 (2024).

37. Fillios, M. A. & Taçon, P. S. C. Who let the dogs in? A review of the recent genetic evidence for the introduction of the dingo to Australia and implications for the movement of people. *J. Archaeol. Sci. Rep.* **7**, 782–792 (2016).

38. McKeever, B., Podadera, J. M., Beijerink, N. J. & Šlapeta, J. Suspect 'baggage canine heartworm' case: canine heartworm disease in a dog from Sydney, New South Wales. *Aust. Vet. J.* **99**, 359–362 (2021).

39. Ensminger, J. J. From hunters to hell hounds: the dogs of Columbus and transformations of the human-canine relationship in the early Spanish Caribbean. *Colon. Latin Am. Rev.* **31**, 354–380 (2022).

40. Symcox, G. & Sullivan, B. Introduction: Columbus—the man, the voyages, the legacy. in *Christopher Columbus and the Enterprise of the Indies* 1–40 (Palgrave Macmillan US, New York, 2005).

41. Welker, M. H. & Dunham, R. Exploring the introduction of European dogs to North America through shoulder height. *Intl J of Osteoarchaeology* **29**, 325–334 (2019).

42.  Lau, D. C.-W. *et al.* Whole-genome reference of from Australia to determine single
nucleotide polymorphisms associated with macrocyclic lactone resistance in the USA.
*Curr Res Parasitol Vector Borne Dis* **1**, 100007 (2021).

43.  Godel, C. *et al.* The genome of the heartworm, *Dirofilaria immitis*, reveals drug and
vaccine targets. *FASEB J.* **26**, 4650–4661 (2012).

44.  Ewels, P., Magnusson, M., Lundin, S. & Käller, M. MultiQC: summarize analysis
results for multiple tools and samples in a single report. *Bioinformatics* **32**, 3047–3048
(2016).

45.  Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

46.  Di Tommaso, P. *et al.* Nextflow enables reproducible computational workflows. *Nat.*
*Biotechnol.* **35**, 316–319 (2017).

47.  McIntyre, J. *et al.* Chromosomal genome assembly resolves drug resistance loci in the
parasitic nematode *Teladorsagia circumcincta. bioRxiv* 2024.07.22.603261 (2024)
doi:10.1101/2024.07.22.603261.

48.  Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**,
3094–3100 (2018).

49.  Van der Auwera, G. A. & O'Connor, B. D. *Genomics in the Cloud: Using Docker,*
*GATK, and WDL in Terra*. (O'Reilly Media, 2020).

50.  Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**, giab008
(2021).

51.  Tarasov, A., Vilella, A. J., Cuppen, E., Nijman, I. J. & Prins, P. Sambamba: fast
processing of NGS alignment formats. *Bioinformatics* **31**, 2032–2034 (2015).

52.  Barnett, D. W., Garrison, E. K., Quinlan, A. R., Strömberg, M. P. & Marth, G. T.
BamTools: a C++ API and toolkit for analyzing and managing BAM files.

*Bioinformatics* **27**, 1691–1692 (2011).

53.    Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing

649           genomic features. *Bioinformatics* **26**, 841–842 (2010).

650    54.    Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158

651           (2011).

652    55.    Zheng, X. *et al.* A high-performance computing toolset for relatedness and principal

653           component analysis of SNP data. *Bioinformatics* **28**, 3326–3328 (2012).

654    56.    Korunes, K. L. & Samuk, K. pixy: Unbiased estimation of nucleotide diversity and

655           divergence in the presence of missing data. *Mol. Ecol. Resour.* **21**, 1359–1368 (2021).

656    57.    Skotte, L., Korneliussen, T. S. & Albrechtsen, A. Estimating individual admixture

657           proportions from next generation sequencing data. *Genetics* **195**, 693–702 (2013).

658    58.    Kopelman, N. M., Mayzel, J., Jakobsson, M., Rosenberg, N. A. & Mayrose, I. Clumpak:

659           a program for identifying clustering modes and packaging population structure

660           inferences across K. *Mol. Ecol. Resour.* **15**, 1179–1191 (2015).

661    59.    Pickrell, J. K. & Pritchard, J. K. Inference of population splits and mixtures from

662           genome-wide allele frequency data. *PLoS Genet.* **8**, e1002967 (2012).

663    60.    Patterson, N. *et al.* Ancient admixture in human history. *Genetics* **192**, 1065–1093

664           (2012).

665    61.    Chang, C. C. *et al.* Second-generation PLINK: rising to the challenge of larger and

666           richer datasets. *Gigascience* **4**, 7 (2015).

667    62.    Terhorst, J., Kamm, J. A. & Song, Y. S. Robust and scalable inference of population

668           history from hundreds of unphased whole genomes. *Nat. Genet.* **49**, 303–309 (2017).

## Acknowledgements

The authors gratefully acknowledge Julie Austin, Iris Cheadle, Constantin Constantinoiu, Carol Esson, Eloise Fenton, Linda Hayes, Tessa Mackie, Yissu Martinez, Sandi McClintock, Ilze Nel, Tony Phillis, Duncan Smith, and Jim Taylor for their generous provision of heartworm specimens for this study. Sampling from the University of Queensland was supported by the University of Queensland, School of Veterinary Science Bequest and Donation Funds for Research (Stone Memorial Fund). We thank Javier Gandasegui for their foundational bioinformatics work that facilitated this work. We acknowledge the Pathogen Informatics group at the Wellcome Sanger Institute and the Sydney Informatics Hub for bioinformatics support. RIP is supported by an Australian Government Research Training Program Stipend and The Jean Walker Trust Fellowship. RIP's collaborative research exchange was supported by the University of Sydney's Eric Horatio Scholarship, the University of Sydney's Postgraduate Research Support Scheme, and the Australian Society for Parasitology's Researcher Exchange, Training and Travel Award. SRD is supported by a UKRI Future Leaders Fellowship [MR/T020733/1] and the Wellcome Trust (UK) through core funding to the Wellcome Sanger Institute (UK) [206194]. This work was funded by the Canine Research Foundation, Dogs Victoria, Australia and the Australian Companion Animal Health Foundation Research Grant.

**Figures**

**Fig. 1: Global sampling and population structure of heartworms. a,** World map showing

the location of adult heartworm specimens used in the study. Adult *Dirofilaria ursi*, *D. repens*,

and *D*. sp. 'Thailand II' were included as outgroup samples. Abbreviations: USA = United

States of America (Georgia: GEO, Illinois: ILL, Louisiana: LOU, Missouri: MIS, Texas: TEX,

Florida: FLO); CRI = Costa Rica; PAN = Panama; ITA = Italy; ROU = Romania; GRC =

Greece; VNM = Vietnam; THA = Thailand; MYS = Malaysia; AUS = Australia (Sydney: SYD,

Brisbane: BNE, Rockhampton: ROK, Townsville: TSV, Cairns: CNS, Lockhart River: LHR).

**b,** Principal component analysis (PCA) of 196,550 nuclear single nucleotide polymorphisms

(SNPs) from 124 heartworms. **c,** Box plots showing the nucleotide diversity (Pi) distribution

within broad geographical regions, with each point representing a 100 kb sliding window. **d,**

Linkage disequilibrium (LD) decay plot for the heartworm cohort, grouped by broad

geographic region. Vertical lines show the distance between SNPs where the LD is half the

maximum value. **e,** Admixture inferred using NGSadmix with six clusters (K). Each vertical

bar represents an individual sample, with colours representing the clusters. Labels at the top of

the bar charts indicate the broad geographic origins of samples, while those at the bottom show

their more specific locations. An asterisk (*) denotes a replicate of the preceding sample.

**Fig. 2: Heartworm variation between carnivoran hosts. a,** Principal component analysis

(PCA) of heartworm nuclear diversity from Fig. 1b, with non-dog hosts labelled. The

geographic origin of samples is indicated on the right (i.e. COUNTRY: city or region). Country

abbreviations: USA = United States of America; CRI = Costa Rica; PAN = Panama; ITA =

Italy; ROU = Romania; GRC = Greece; VNM = Vietnam; THA = Thailand; MYS = Malaysia;

AUS = Australia. **b,** Box plots show the distribution of nucleotide diversity (Pi) within host

populations. Each data point represents a sliding window size of 100 kb.

**Fig. 3: Demographic history and host competency for heartworms. a,** Effective population

size history of heartworm populations inferred using SMC++ based on a four-year generation

time. Each line represents the effective population size of heartworms on each continent over

time. The grey box spans ~14–40 kya, highlighting the previously suggested period of dog

domestication. Vertical dotted lines and associated numbers represent divergence times

between populations. Details of these population splits and their timing are provided on the

right. Abbreviations: AUS = Australia; CENAM = Central America; EUR = Europe; USA =

United States of America. **b,** Support for the heartworm life cycle and adult worm burdens in

carnivoran hosts. The top panel highlights hosts that are frequently microfilaremic and are

hence considered competent reservoirs for heartworms. A black star indicates cases where

microfilaremia has been reported in the host, but they are generally considered to pose a low

risk of being competent reservoirs. The bottom panel is a plot showing adult heartworm

burdens across various hosts. Each point represents data from a study, sourced from the

literature (Supplementary Table 2). Significant differences in worm burdens among competent

reservoirs were assessed using a Kruskal-Wallis test with Dunn's multiple comparisons. P-

values $\leq 0.05$ were considered statistically significant (*: $\leq 0.05$, **: $\leq 0.01$, ***: $\leq 0.001$).

**Fig. 4: Transcontinental admixture in heartworms. a,** Maximum-likelihood tree of heartworm generated using TreeMix, with *Dirofilaria ursi* (DURSI) as an outgroup and one migration edge. **b,c,** *D*-statistics for hypothetical allele-sharing scenarios between heartworm populations using Admixtools qpDstat with 21,088 single nucleotide polymorphisms (SNPs) and *D. ursi* as the outgroup (OUT). *D*-statistics test for admixture between four populations (W, X, Y, Z). A Z-score (*D*-statistic / standard error) is calculated, which provides information about the direction of gene flow (red dotted lines) (Z-score > 0 = gene flow between either [W and Y], or [X and Z]; Z-score < 0 = gene flow between either [W and Z] or [X and Y]). For all data presented, the Z score was significant ($|Z| > 3$). Error bars show the standard error.

**Fig. 5: Gene flow and population structure of Australian heartworms. a,** Distribution of

adult heartworm samples ($n = 48$) collected along the east coast of Australia. **b,** $D$-statistics

showing excessive allele sharing between Australia (AUS) and ASIA relative to that between

AUS and other regions (PopX). Admixtools qpDstat was used to analyse 21,088 single

nucleotide polymorphisms (SNPs), with *Dirofilaria ursi* as the outgroup (OUT). $D$-statistics

test for admixture between four populations (W, X, Y, Z). A Z-score ($D$-statistic / standard

error) is calculated, which provides information about the direction of gene flow (red dotted

lines) (Z-score > 0 = gene flow between [W and Y], or [X and Z]; Z-score < 0 = gene flow

between [W and Z] or [X and Y]). For all data presented, the Z score was significant ($|Z| > 3$).

Error bars show the standard error. **c,** Box plots showing the nucleotide diversity (Pi)

distribution from dog hosts per city. Each data point represents a 100 kb sliding window. **d,**

Principal component analysis (PCA) of 116,492 nuclear SNPs in the Australian cohort. **e,**

Absolute nucleotide divergence ($D_{XY}$) and **f,** genetic differentiation ($F_{ST}$) between Australian

cities using a 100 kb sliding window. Only heartworms from dog hosts were included in the

analysis.

**Fig. 6: A new evolutionary paradigm of heartworms throughout history. a,** Hypothetical heartworm dispersal scenarios supported in the current study. Pink regions represent a panmictic population of ancient canids harbouring heartworms across North America and Eurasia. The Bering Land Bridge is shaded in green, and continental ice sheets are shown in blue. Events are in chronological order (A–E), with arrows illustrating ancient (dashed), old (dotted), and modern (solid) migration events. Small black shapes with adjacent numbers indicate the geographic origin of samples collected in this study and their sample size. The legend specifies the parasite species, including the heartworm *Dirofilaria immitis* (*n* = 127)

771    and outgroups *D. repens* ($n = 2$), D. sp. "Thailand II" ($n = 2$), and *D. ursi* ($n = 1$). **b,** Timeline

772    of key canid evolution and climatic events. Horizontal bars represent Epochs: HOLO =

773    Holocene; PLEI = Pleistocene; PLIO = Pliocene; MIO = Miocene; OLI = Oligocene; EOC =

774    Eocene. The time scale was log-transformed to enhance the visualisation of events.

**Extended Data Figures**

**Extended Data Fig. 1: Principal component analysis (PCA) of 196,550 nuclear single nucleotide polymorphisms (SNPs) from 124 heartworms with labelled metadata.** Samples were collected from Australia (AUS), the USA, Panama (PAN), Greece (GRC), Thailand (THA), Italy (ITA), Romania (ROU), Costa Rica (CRI), and Malaysia (MYS). Replicate samples were included in the analyses. The legend on the right indicates the geographic origin of samples (i.e. COUNTRY: city or region). Principal component 1 (PC1) and principal component 2 (PC2) are shown on the x and y axes, explaining 12.6% and 10.69% of the variation, respectively. These figures are identical to Fig. 1b but include sample labels.

**Extended Data Fig. 2: Principal component analysis (PCA) of 124 heartworms from around the world based on the nuclear genome.** Samples were collected from Australia (AUS), the USA, Panama (PAN), Greece (GRC), Thailand (THA), Italy (ITA), Romania (ROU), Costa Rica (CRI), and Malaysia (MYS). Replicate samples were included in the

analyses. The legend on the right indicates the geographic origin of samples (i.e. COUNTRY:

city or region). The principal components (PCs) are shown on the x and y axes. **a,** PCA of

196,550 nuclear single nucleotide polymorphisms (SNPs) showing PC3 and PC4, which

explained 5.76% and 4.39% of the variation, respectively. **b–f,** PCAs of 44,937 – 67,561 SNPs

in chromosome 1, chromosome 2, chromosome 3, chromosome 4, and the sex-linked X-

chromosome of canine heartworm. PC1 and PC2 explained 11.67 – 14.01% and 10.09 –

11.37% of the variation, respectively.

**Extended Data Fig. 3: Genetic relatedness of the mitochondria and *Wolbachia* endosymbionts of heartworms.** Abbreviations: USA = United States of America; CRI = Costa Rica; PAN = Panama; ITA = Italy; ROU = Romania; GRC = Greece; VNM = Vietnam; THA = Thailand; MYS = Malaysia; AUS = Australia. **a,** Principal component analysis (PCA) of 32 mitochondrial single nucleotide polymorphisms (SNPs) from 127 heartworms. Principal component 1 (PC1) and principal component 2 (PC2) explained 30.32% and 17.98% of the variation, respectively. **b,** PCA of 280 *Wolbachia* endosymbiont SNPs from 121 samples. PC1 and PC2 explained 39.88% and 18.67% of the variation, respectively. Replicate samples were included in all PCAs. The legend on the right indicates the geographic origin of samples (i.e. COUNTRY: city or region).

**Extended Data Fig. 4: Admixture analysis of adult heartworms.** Analysis was performed

in NGSadmix across a range of K values (2 – 10) (colours) for 124 heartworms (vertical bars).

Geographic origins are indicated at the bottom of each chart. Abbreviations: Australia (Sydney:

SYD, Brisbane: BNE, Rockhampton: ROK, Townsville: TSV, Cairns: CNS, Lockhart River:

LHR); Thailand (Bangkok: BKK); Malaysia (Selangor: SEL); USA (Georgia: GEO, Illinois:

ILL, Louisiana: LOU, Missouri: MIS, Texas: TEX, Florida: FLO); Italy (Northeast: NEA,

Pavia: PAV); Romania (Bucharest: BUC, Giurgiu: GIU, Comana: COM); Greece

(Thessaloniki/Xanthi: XAN); Costa Rica (San José: SJO); Panama (Puerto Armuelles: PUE,

San Lorenzo: SLO, Boca Chica: BOC).

**Extended Data Fig. 5: Population genetic summary statistics of heartworms collected from diverse host populations.** Heatmaps show **a,** Absolute nucleotide divergence ($D_{XY}$) and **b,** Genetic differentiation ($F_{ST}$) using a 100 kb sliding window.

**Extended Data Fig. 6: Population genetic summary statistics of heartworms between broad geographical regions.** Heatmaps show **a,** Absolute nucleotide divergence ($D_{XY}$) and **b,** Genetic differentiation ($F_{ST}$) using a 100 kb sliding window. Region abbreviations: ASIA = Asia; AUS = Australia; CENAM = Central America; EUR = Europe; USA = United States of America.

**Extended Data Fig. 7: TreeMix analysis of nuclear heartworm data.** Maximum likelihood trees are shown with migration edges ranging from 0 to 5, using *Dirofilaria ursi* as an outgroup. Coloured lines represent inferred migration events, with the colour indicating support levels. Abbreviations: Australia: AUS (Sydney: SYD, Brisbane: BNE, Rockhampton: ROK, Townsville: TSV, Cairns: CNS, Lockhart River: LHR); USA (Georgia: GEO, Illinois: ILL, Louisiana: LOU, Missouri: MIS, Texas: TEX, Florida: FLO); Thailand: THA; Malaysia: MYS; Italy: ITA; Romania: ROU; Greece: GRC; Costa Rica: CRI; Panama: PAN.

**Extended Data Fig. 8: Summary diagram of the population genomics workflow in this study.** Abbreviations of population structure analyses: PCA = Principal Component Analysis; Pi = nucleotide diversity; $D_{XY}$ = absolute nucleotide divergence; $F_{ST}$ = genetic differentiation.

**Extended Data Fig. 9: Demographic history analysis of heartworms from dogs using a range of generation times.** Size histories were inferred using SMC++, with each line representing a heartworm subpopulation. A range of generation times were tested for *Dirofilaria immitis* (i.e. 1, 2, 4, and 6 years). The grey box indicates previously suggested date ranges for dog domestication (~14–40 kya). Abbreviations: AUS = Australia; CENAM =

848    Central America; EUR = Europe; USA = United States of America; MIS 4 = Marine Isotype

849    Stage 4.

# Supplementary Figures

**Supplementary Fig. 1: Raw nuclear sequencing coverage per sample (Part 1/2).** Coverage

was calculated using 100 kb sliding windows for all mapped samples. Each chromosome is

shown in a different colour, with a legend on the right. The remaining samples are shown in

Supplementary Fig. 2.

**Supplementary Fig. 2: Raw nuclear sequencing coverage per sample (Part 2/2).** Coverage

was calculated using 100 kb sliding windows for all mapped samples. Each chromosome is

shown in a different colour, with a legend on the right.

**Supplementary Fig. 3: Sex determination of samples based on relative coverage of sex-linked vs autosomal chromosomes (Part 1/2).** Sequencing coverage was assessed using a 100kb sliding window and compared to the overall median coverage. Coverage of the sex-

865    linked X chromosome (blue) was compared to an autosome (chromosome 1, red) to determine

866    the parasite's sex. Male canine heartworms show reduced coverage on chromosome X (ratio

867    ~0.5), while females show similar coverage across sex-linked and autosomal chromosomes

868    (ratio ~1). Ambiguous results were observed in samples with low coverage. The remaining

869    samples are shown in Supplementary Fig. 4.

**Supplementary Fig. 4: Sex determination of samples based on relative coverage of sex-linked vs autosomal chromosomes (Part 2/2).** Sequencing coverage was assessed using a 100kb sliding window and compared to the overall median coverage. Coverage of the sex-

875    linked X chromosome (blue) was compared to an autosome (chromosome 1, red) to determine

876    the parasite's sex. Male canine heartworms show reduced coverage on chromosome X (ratio

877    ~0.5), while females show similar coverage across sex-linked and autosomal chromosomes

878    (ratio ~1). Ambiguous results were observed in samples with low coverage.