Title of article:

Mixed DAMP/MAMP oligosaccharides promote both growth and defense against fungal

pathogens of cucumber

Authors:

Sreynich Pring[1], Hiroaki Kato[2], Keiko Taniuchi[1], Maurizio Camagna[1], Makoto Saito[3], Aiko

Tanaka[1], Bryn A. Merritt[4], Orlando Argüello-Miranda[4], Ikuo Sato[1], Sotaro Chiba[1] and Daigo

Takemoto[1]*

Affiliation:

[1] Graduate School of Bioagricultural Sciences, Nagoya University, Chikusa, Nagoya, 464-

8601, Japan

[2] Graduate School of Agriculture, Kyoto University, Muko, Kyoto, 617-0001, Japan.

[3] Resonac Corporation (Showa Denko K.K.), Tokyo, Japan

[4] Department of Plant and Microbial Biology, North Carolina State University, Raleigh, NC

27695-7612 USA

*Corresponding Author:

Daigo Takemoto

E-mail: dtakemo@agr.nagoya-u.ac.jp

Telephone: +81 90 4264 4192

Fax: +81 52 789 5525

Number of figures and tables:

8 Figures and 2 Supplementary Tables

ABSTRACT

Plants recognize a variety of environmental molecules, thereby triggering appropriate responses to biotic or abiotic stresses. Substances containing microbes-associated molecular patterns (MAMPs) and damage-associated molecular patterns (DAMPs) are representative inducers of pathogen resistance and damage repair, thus treatment of healthy plants with such substances can pre-activate plant immunity and cell repair functions. In this study, the effects of DAMP/MAMP oligosaccharides mixture (Oligo-Mix) derived from plant cell wall (cello-oligosaccharide and xylo-oligosaccharide), and fungal cell wall (chitin-oligosaccharide) were examined in cucumber. Treatment of cucumber with Oligo-Mix promoted root germination and plant growth, along with increased chlorophyll contents in the leaves. Oligo-Mix treatment also induced typical defense responses such as MAP kinase activation and callose deposition in leaves. Pretreatment of Oligo-Mix enhanced disease resistance of cucumber leaves against pathogenic fungi *Podosphaera xanthii* (powdery mildew) and *Colletotrichum orbiculare* (anthracnose). Oligo-Mix treatment increased the induction of hypersensitive cell death around the infection site of pathogens, which inhibited further infection and the conidial formation of pathogens on the cucumber leaves. RNA-seq analysis revealed that Oligo-Mix treatment upregulated genes associated with plant structural reinforcement, responses to abiotic stresses and plant defense. These results suggested that Oligo-Mix has beneficial effects on growth and disease resistance in cucumber, making it a promising biostimulant for agricultural application.

Keywords: Anthracnose, Biostimulant, Cucumber, Damage-associated molecular patterns, Elicitors, Microbe-associated molecular patterns, Oligosaccharides, Powdery mildew.

1. Introduction

Plants have an array of mechanisms for sensing and adapting to environmental stresses. Some breakdown molecules of plant cells are released by physical damage, pathogen infection, or insect attack and are recognized as signals by plants to promote cellular repair or defense responses. Such molecules are commonly called DAMPs (damaged-associated molecular patterns), including cello-oligosaccharides (COS), which are constituent molecules of cellulose derived from plant cell walls, and xylo-oligosaccharides (XOS) contained in hemicellulose, which cross-link cellulose and lignin in the cell wall (Souza et al., 2017; Claverie et al., 2018; Pring et al., 2023). Volatiles emitted by damaged plants are also involved in stress responses, thus considered as a type of DAMPs (Yamauchi et al., 2018). It is also known that specific peptides released externally by cell disruption are recognized as DAMPs by receptors on the surface of neighboring cells (Pearce et al., 1991; Yamaguchi et al., 2010; Wang et al., 2018; Hander et al., 2019).

Molecules specific to microorganisms are recognized by plant cells as alert signals of pathogen attack, and the epitope structures of such substances are called MAMPs (microbes-associated molecular patterns). Some MAMPs are highly conserved in large groups (i.e. bacteria, fungi or oomycetes), such as flg22 (a conserved peptide of the bacterial flagellar protein flagellin, Gómez-Gómez et al., 1999), peptidoglycan (the structural polymer of the bacterial cell wall, Gust et al., 2007), chitin and β-glucan (major cell wall components of fungi, Sharp and Albersheim, 1984; Felix et al., 1993) and 9-methyl-4,8-sphingadienine (9Me-Spd, a substructure of ceramides found in the cell membrane of fungi and oomycetes, Kato et al., 2022; Monjil et al., 2024). Other MAMPs are found in certain groups of pathogenic microbes, such as elicitins produced by oomycete pathogens (Derevnina et al., 2016), and fungal ethylene-inducing xylanases (EIXs, Dean et al., 1991). On the other hand, the recognition mechanisms of MAMPs by plant species are also diverse; for example, Solanaceae plants and rice can recognize different parts of bacterial flagellin other than flg22 (Fliegmann et al., 2016). Elicitin and EIX have also been reported to be recognized as MAMPs by a limited range of plant species (Kamoun et al., 1993; Ron and Avni, 2004; Takemoto et al., 2005).

While the understanding of the recognition mechanisms of DAMPs and MAMPs by plants is a subject of basic research, these substances are also anticipated to be utilized as materials to activate plant growth and immunity in agricultural production. Substances that activate beneficial plant responses are collectively referred to as biostimulants. The DAMPs and MAMPs derived from fundamental structures of plants and microbes are expected to be versatile as agricultural materials because they can activate the responses of diverse plant

species. The cell walls of plants and fungi are essentially composed of common materials (Cosgrove 2005; Gow and Lenardon 2023), thus substances in their cell walls are representative candidates for biostimulant.

Previously, we have shown that DAMP and MAMP oligosaccharides, COS prepared from cotton linters, XOS prepared from corn cobs as well as chitin-oligosaccharide (CHOS) from crustacean shells can trigger typical defense responses of Arabidopsis such as production of reactive oxygen species (ROS), MAP kinases phosphorylation and callose depositions (Pring et al., 2023). Gene ontology enrichment analysis of RNA-seq data revealed that simultaneous treatment of COS, XOS and CHOS (Oligo-Mix) effectively activate genes for disease resistance. Generally, it is considered that activating disease resistance in plants suppresses their growth, referred to as "trade-off effect" (Karasov et al., 2017). However, the treatment of Oligo-Mix had no significant defects on the expression of photosynthesis genes. In practice, treatment of the Oligo-Mix enhanced resistance of tomato against powdery mildew pathogen, and moreover, tomato growth was rather promoted by Oligo-Mix treatment (Pring et al., 2023).

In this study, the effects of Oligo-Mix treatment on cucumber were investigated. Cucumber (*Cucumis sativus* L.) is one of the most important vegetable crops cultivated worldwide. Besides the varieties with higher resistance to pathogens, in general, cucumbers are susceptible to a variety of destructive pathogens. Anthracnose, caused by a fungal pathogen *Colletotrichum orbiculare*, is a major foliage disease that reduces the yield and quality of cucumber and other Cucurbitaceae plants such as watermelon and melon. *C. orbiculare* is a hemi-biotrophic pathogen that employs both biotrophic and necrotrophic phases to infect its hosts (Matsuo et al., 2022). Powdery mildew of cucumber is widely distributed, rapidly spreading to cause destructive disease by obligate parasites *Golovinomyces cichoracearum* or *Podosphaera xanthii* (Lebeda et al., 2011). While there are some fungicides accessible for managing these pathogens, the emergence of drug-resistant fungal strains has become a major problem (Vielba-Fernández et al., 2020). Here, we evaluated the influence of the Oligo-Mix treatment on cucumber growth, development, and resistance against anthracnose and powdery mildew pathogens.

2.   Material and methods

2.1. Plant material and growth condition

Cucumber (cv. Shimoshirazu Jibai, Takii seed, Kyoto, Japan) were grown in autoclaved commercial soil (Sakata Super Mix A, Sakata seed, Yokohama, Japan) in a plant growth room at 23°C with 16 h of light per day. For the greenhouse test, two-week-old seedlings grown in a

controlled growth chamber were transferred to pots (28 cm diameter x 22 cm height, filled with

Sakata Super Mix A) and grown in a greenhouse in Togo field, Nagoya University (University

farm, Togo, Aichi, Japan).

2.2. Oligosaccharide mixture (Oligo-Mix)

Oligo-Mix (undiluted solution) used in this study consists of 20 mg/ml cello-oligosaccharide

(COS, prepared from cotton linter), 40 mg/ml xylo-oligosaccharide (XOS, from corn cob), and

20 mg/ml chitin-oligosaccharide (CHOS, from shrimp shell) (Sreynich et al., 2023). Both

Oligo-Mix and the control solutions contain water-soluble P (P_2O_5, 4.7% w/v) and K (K_2O,

3.4% w/v). Oligo-Mix and control solution were diluted to 1/1000 and used for the treatment

of plants by spray or infiltration using a needleless syringe. Oligo-Mix and control solutions

were provided by Resonac Corporation (Tokyo, Japan).

2.3. Determination of root number and length

Cucumber seeds were placed on laboratory tissue soaked with distilled water in Petri dishes

and incubated at 26 ˚C with a light/dark cycle set at 16 h/8 h. The cucumber seeds germinated

at same day were collected and sprayed with Oligo-Mix or control solution (250 µl per petri

dish), and the number of roots per seed and the length of longest root were measured 2 days

after the treatment.

2.4.Determination of cucumber growth

The individual seedlings, treated with control solution or Oligo-Mix as described above,

were transferred into 9-cm-diameter plastic pot filled with sterilized soil mix and grown in

growth room at 23 ˚C with 16 h of light per day. Cucumber plants were sprayed with Oligo-

Mix or control solution once every week (2 ml per plant, 4 times altogether) until sampling.

After one week of the last treatment, the fresh weight of the root and above-ground tissues are

measured.

2.5. Measurement of chlorophyll contents

Chlorophyll was extracted according to the method described in Shibata et al., (2016). One

g of second true leaves of cucumber was soaked in 50 ml methanol overnight. The absorbance

intensity of the solution was quantified by spectrophotometer (Multiskan GO, Thermo Fisher

Scientific, Waltham, MA, USA), and the concentration of chlorophyll a and b was calculated

according to the method of Porra and Scheer (1989) as follows; Chlorophyll a (μg/ml) = 16.29

x A^{665} − 8.54 x A^{652}; Chlorophyll b (μg/ml) = 30.66 x A^{652} − 13.58 x A^{665}.

2.6. MAP kinase activation assay

Activation of MAP kinases in Arabidopsis was detected as previously reported (Kato et al.,

2022). Cucumber seedlings were treated with Oligo-Mix or control solution (by infiltration

with a needleless syringe) for 15 or 30 min and frozen in liquid nitrogen. Proteins were

extracted in extraction buffer (50 mM HEPES-KOH pH 7.4, 5 mM EDTA, 0.5 mM EGTA, 50

mM ß-glycerophosphate, 10 mM NaF, 10 mM Na_3VO_4, and 2 mM DTT). Phosphorylated

MAP kinases were detected by western blot using Phospho-p44/42 MAPK (Erk1/2; Thr-

202/Tyr-204) rabbit monoclonal antibodies #9101 (Cell Signaling Technology). Blots were

stained with PageBlue Protein Staining Solution (Thermo Fisher Scientific) to verify equal

loading.

2.7. Staining of callose deposition

Staining of callose deposition was performed as previously reported (Shibata et al., 2016).

Cucumber seedlings were treated with Oligo-Mix or control solution by infiltration with

needleless syringe, and seedlings were fixed 24 h after treatment in fixation solution (1% [v/v]

glutaraldehyde, 5 mM citric acid, and 90 mM Na_2HPO_4, pH 7.4) overnight. Fixed seedlings

were heated in boiled water for 5 minutes, decolorized in ethanol, and stained in aniline blue

stain (0.1% [w/v] aniline blue in 67 mM phosphate buffer, pH 12.0) to detect callose deposition.

The fluorescence spots of deposited callose were detected by a fluorescence microscope (BX51,

Olympus, Tokyo, Japan) using an excitation wavelength of 365 nm. The number of callose

spots per area was quantified using ImageJ software (Schneider et al., 2012).

2.8. Pathogens and inoculation

Powdery mildew of cucumber, *Podosphaera xanthii* strain SP23*,* was isolated from cucumber

grown in a greenhouse in Togo field, Nagoya University, and maintained by inoculating

healthy cucumbers every 5-10 days. The species of powdery mildew was determined based on

the sequencing of ITS (Internal transcribed spacer) region using the method previously

described (Ashida et al., 2023). Conidia newly formed on cucumber leaves were suspended in

water and adjusted to a concentration of 1×10^5 spores/ml. The 5-leaf-stage cucumber plants

were inoculated by spraying 0.5 ml spore suspension per plant at 24 h after the treatment with

control solution or Oligo-Mix. One week later, spot symptoms were observed on the leaf

surface, and number and type (spored, yellowish, or cell death) of colony were counted. To

measure conidial formation, cut leaves with colonies (1.5 × 1.5 cm) were scrubbed with 2 ml

of distilled water using a cotton bud, then filtered with a stainless filter disc (mesh size 100

μm). Subsequently, the number of conidia was quantified with a hemocytometer (Erma, Tokyo,

Japan).

Anthracnose pathogen, *Colletotrichum orbiculare* strain T-104 (MAFF240422, which was

isolated by Dr. Yasumori, Kyoto University in 1951) was provided by Prof. Yoshitaka Takano

(Kyoto University, Japan). *C. orbiculare* was grown on potato dextrose agar (PDA) in 23°C

for 7 days, and conidia formed on the colony were suspended in water and adjusted to a

concentration of 1×10^5 spores/ml for the inoculation. Cucumber was spray-inoculated with 0.5

ml of spore suspension per plant. The symptoms observed on the leave were photographed 7

days after the inoculation.

2.9. Detection of anthracnose lesions on cucumber leaves

Briefly, the area affected by anthracnose on cucumber leaves was calculated as the

percentage of the affected area divided by the total leaf area. The calculation of the total leave

area was done using the custom-made MATLAB script (OAM_240209_leave_seg) to isolate

all pixels in an image that belonged to a leaf according to a median intensity threshold. To

identify specific infected areas on individual leaves, we trained a modified pixel flow-driven

CNN architecture (Stringer et al., 2021) to identify each pixel on a leaf that could be classified

as affected with more than 0.9 probability. The CNN model was trained using the preliminary

segmentations from the MATLAB script based on median thresholding, which generated

single-leaf masks depicting the affected areas. The preliminary masks were manually corrected

using custom-made software to remove incorrectly classified pixels, and the final corrected

images were split into training (70%), and testing (30%) data sets for deep learning model

training. The CNN architecture was modified to obtain a binary lesion/no-lesion single-pixel

classification (semantic segmentation) as direct output. The CNN model "Takemoto_1" and

the accessory code for whole leaf segmentation, and to obtain the final semantic segmentation

from the CNN, as well as the original image dataset with labels and a toy dataset to test the

segmentation code, are freely available from the GitHub repository of the Miranda Lab:

https://github.com/MirandaLab/Deep_Leaf_Segmentation. Leaf segmentations were

performed on a Dual Intel Xeon Silver 4216 (2.1GHz,3.2GHz Turbo, 16C,9.6GT/s 2UPI,

22MB Cache, HT (100W) equipped with an Nvidia Quadro RTX5000, 16GB, 4DP,
VirtualLink (XX20T) outfitted with 64GB 4x16GB DDR4 2933MHz RDIMM ECC Memory
and a 2TB primary NVMe SSD. Differences between control and treatments were evaluated
using the Kolmogorov-Smirnov tests *kstest2* MATLAB function, with significance set at $p <$
0.05.

2.10.   Trypan blue staining and microscopy

To visualize the pathogen hyphae and conidiophores, or HR-like cell death of plant cells,
the infected leaves were stained with trypan blue as described previously (Takemoto et al.,
2003). Small pieces of leaves were cleared in methanol in 1.5 ml tube for more than 24 h
(methanol was changed once) until the chlorophyll (green color) was removed. Then methanol
was replaced with lactophenol trypan blue stain (10 ml lactic acid, 10 ml glycerol, 10 g phenol,
10 mg trypan blue, 10 ml H_2O) and heated in boiled water for 3 mins. After cooling down to
room temperature, the stain was replaced with 1 g/ml chloral hydrate, and gently shaken
overnight. Pathogen hyphae and plant cells were monitored under a light microscope (BX51,
Olympus, Japan).

2.11.   RNA-seq and data analysis

The four-weeks old cucumber plants were sprayed with Oligo-Mix or control solution and
total RNA was isolated at 24 h after treatment. RNA extraction was performed using the
RNeasy Plant Mini Kit (QIAGEN, Hilden, Germany). Evaluation of RNA quality, construction
of library and sequencing were performed principally as previously described (Kuroyanagi et
al., 2022, Bulasag et al., 2023). Libraries were constructed using KAPA mRNA Capture Kit
(Roche Diagnostics, Tokyo, Japan) and MGIEasy RNA Directional Library Prep Set (MGI,
Shenzhen, China), and sequenced on DNBSEQ-G400RS (MGI) with 150 bp paired-end
protocol. The RNA-seq reads were filtered using trim-galore v.0.6.6 (Martin, 2011,
bioinformatics.babraham.ac.uk) and mapped to the cucumber genome (genome assembly
Cucumber_9930_V3, NCBI RefSeq sequence GCA_000004075.3) using HISAT2 v.2.2.1
(Kim et al., 2019) and abundance inferred via StringTie v.2.1.7 (Kovaka et al., 2019).
Significant differential expression was determined using DESeq2 v.1.32.0 (Love et al., 2014).
All software used during RNA-seq analysis was run with default settings. RNA-seq data for
cucumber reported in this work are available in GenBank under the accession numbers
DRA019928.
SRplot (https://www.bioinformatics.com.cn/en), a free online data analysis and

visualization platform, was used to draw the graphs.

3.   Results and discussion

3.1. Effect of Oligo-Mix treatment on the growth of cucumber

To elucidate the effect of the Oligo-Mix (mixture of cello-oligosaccharide COS, xylo-oligosaccharide XOS, and chitin-oligosaccharide CHOS) on cucumber growth, newly germinated cucumber seeds were spray-treated with Oligo-Mix or control solution, and root number and length were measured 2 days after the treatment. The number of roots per seed and length of longest root were significantly increased (almost 2 times), indicating that Oligo-Mix can promote the growth of cucumber roots (Fig. 1 A). Those seedlings were transplanted to the pot and sprayed with Oligo-Mix weekly (once a week). One week after the last spray (sprayed 4 times and 5 weeks after transplant), the growth of cucumber plants was evaluated. The fresh weight of roots and above-ground tissue was significantly increased by Oligo-Mix treatment (Fig. 1 B). Previously, we have reported the positive effects of Oligo-Mix on root and plant growth of tomato (Sreynich et al., 2023). Additionally, leaves of cucumber treated with Oligo-Mix exhibited a visibly darker green compared to those in the control group. To evaluate the effect of Oligo-Mix on chlorophyll contents, cucumber leaves were immersed with methanol to extract and calculated chlorophylls according to the method reported by Porra and Scheer (1989). Chlorophyll a in leaves treated with Oligo-Mix was approx. 405 µg/ml, whereas control leaves contained 256 µg/ml. Similarly, chlorophyll b contents was 143 µg/ml in leaves treated with Oligo-Mix, while at 89 µg/ml for control leaves. These results suggest that Oligo-Mix can effectively enhance photosynthetic capacity, probably leading to improved plant growth.

3.2 Induction of MAP kinase phosphorylation and callose deposition of cucumber by Oligo-Mix treatment

Activation of MAP kinase cascade is a common initial response of plants to induce disease resistance (Asai et al., 2002). Cucumber cotyledons or Arabidopsis seedlings were treated with Oligo-Mix and phosphorylated MAP kinases were detected using antibody against phospho-MAP kinases (Fig. 2A). Within 15 min after Oligo-Mix treatment, rapid activation of cucumber and Arabidopsis MAP kinases was detected. In Arabidopsis, AtMPK6 (approx. 44 kDa) is the major MAP kinase activated by Oligo-Mix, and relatively minor activation of AtMPK3 and AtMPK4 (approx. 39 kDa and 37 kDa, respectively) were also detected. Similarly, in cucumber, a major activated MAP kinase (approx. 46 kDa) and weak activation of two MAP kinases were

also detected within 15 min after Oligo-Mix treatment, suggesting that analogous activation of

similar MAP kinases is induced by Oligo-Mix treatment in Arabidopsis and cucumber.

Callose deposition is another typical plant defense response to reinforce plant cell walls

during pathogen attack (Ellinger and Voigt, 2014). After treatment of cucumber cotyledons

with Oligo-Mix for 24 h, a significant increase in callose depositions was detected (Fig. 2B).

These results suggested that Oligo-Mix treatment can promote the defense responses of

cucumber.

3.3 Effect of Oligo-Mix treatment on disease resistance of cucumber against powdery mildew

and anthracnose pathogens

Two types of plant pathogens were employed to examine the effect of Oligo-Mix on plant

disease resistance. Cucumber powdery mildew fungus, *Podosphaera xanthii*, is an obligate

parasitic pathogen of Cucurbitaceae. Cucumber plants were sprayed with control solution or

Oligo-Mix, and inoculated with spore suspension of *P. xanthii* 24 h after the treatment.

Although the number of lesions developed on leaves was not affected by Oligo-Mix treatment,

the ratio of yellowish or necrotic lesions was significantly increased for leaves treated with

Oligo-Mix (Fig. 3). Microscopic observation revealed the successful penetration, formation of

haustoria, and massive production of conidiophores and conidia on control leaves, while HR

(hypersensitive response)-like cell death was often found for the leaves treated with Oligo-Mix

(Fig. 4). Consistently, Oligo-Mix treatment significantly reduced the spore formation on the

leaves (Fig. 4). These results indicate that the Oligo-Mix treatment is effective in reducing the

amount of spore formation of powdery mildew, thus presumed to be effective in suppressing

the secondary spread of the disease.

Colletotrichum orbiculare, a cucurbit anthracnose fungus, is a hemibiotrophic pathogen that

infects cucumber, melon, watermelon, and other cucurbitaceous plants. As the experiment for

the powdery mildew, cucumber leaves treated with control solution or Oligo-Mix were

inoculated with spore suspension of *C. orbiculare* 24 h after the treatment. Yellow primary

lesions appeared on inoculated leaves about 4 days after inoculation and were obvious by 7

days. Image analysis to quantify the percentage of infected and yellowed areas (see details in

Material and methods section) indicated that Oligo-Mix treatment reduced the severity of the

anthracnose disease (Fig. 5). In control leaves, necrotic lesions and outgrowth of intracellular

biotrophic hyphae from necrotic area were observed (Fig. 6). Formation of acervuli and setae

(asexual fruiting body), and extensive growth of hyphae on leaf surface were also observed in

control leaves, indicating the successful hemibiotrophic infection of anthracnose pathogen. For

leaves treated with Oligo-Mix, in contrast, only immature acervuli and setae were found and
necrotic lesions with no further extension of pathogen hyphae were observed (Fig. 6). These
observations indicated that pre-treatment of Oligo-Mix can reduce the infection of anthracnose
pathogen.
322

3.4 Expression profile of cucumber genes in response to Oligo-Mix treatment

Four-weeks old cucumber leaves were treated with the control solution or Oligo-Mix, and
expression of cucumber genes at 24 h after treatment was investigated by RNA-seq analysis.
The list of significantly up-regulated genes (based on the average TPM score) in cucumber leaf
is shown as Table S1. Oligo-Mix treatment upregulated 123 genes (FC > 2, $p<0.05$, average
TPM>1) and downregulated 46 genes (FC < 0.5, $p<0.05$, average TPM>1). The upregulated
genes include some genes predicted to be involved in cell reinforcement and expansion (Fig.
7A, Table S1). Exordium-like 1 (CsaV3_3G047580) encodes extracellular protein EXO
mediates leaf cell expansion (Schröder et al., 2009). Xyloglucan
endotransglucosylases/hydrolases (CsaV3_5G031640 and CsaV3_6G038030) are known as
enzymes for the modification of cell wall structure by cleaving and re-joining xyloglucan in
primary cell walls (Eklöf et al., 2010). Arabinogalactans (CsaV3_7G003940,
CsaV3_4G009610 and CsaV3_7G028250) are a class of glycoproteins anchored to the plasma
membrane, presumed to be involved in cellulose synthesis and deposition during plant cell wall
biogenesis (Lin et al., 2022). Expression of genes for stress responses were also upregulated
such as ERF/AP2 family transcription factors (e.g. CsaV3_3G016760, CsaV3_3G018600 and
CsaV3_5G005890, Imano et al., 2022) and chaperon HSP70 (CsaV3_5G026520) (Fig. 7B,
Table S1). Oligo-Mix treatment enhanced genes for plant disease resistance, including
Avr9/Cf-9 rapidly elicited protein-like (CsaV3_4G025220, Rowland et al., 2005), F-box
family protein (CsaV3_6G048900), CCR4-associated factor 1 homolog (CsaV3_6G038490,
Liang et al., 2009), PAR1 protein (CsaV3_3G017150, Herbers et al., 1995, Takemoto et al.,
2003) and thaumatin-like pathogenesis-related protein 5 (PR-5, CsaV3_3G045940, de Jesús-
Pires et al., 2020).

The activation of genes associated with stress tolerance and disease resistance often comes
at a trade-off with the plant growth, such as photosynthesis (Karasov et al., 2017). We assessed
the effect of Oligo-Mix treatment on highly expressed genes, however, Oligo-Mix treatment
did not decrease the expression of genes involved in photosynthesis, but rather increased the
expression of chlorophyll a-b binding protein, although this effect was not statistically
significant (Table S2). These results indicate that Oligo-Mix treatment promotes the activation

of cell growth and disease stress tolerance while maintaining the expression levels of genes
required for photosynthesis.

3.5 Effect of Oligo-Mix treatment on powdery mildew resistance in greenhouse

The effect of Oligo-Mix treatment on disease incidence in cucumbers grown in greenhouses
was investigated. Two-week-old cucumber seedlings grown in a controlled growth chamber
were transferred to pots and grown in a greenhouse. Cucumber plants were sprayed with
control solution or Oligo-Mix once every week (6 times altogether) and the number of naturally
occurring colonies of powdery mildew on leaves was counted. Although no statistically
significant differences were found because some plants did not show natural disease symptoms
in both treatments, but the Oligo-Mix treatment tended to reduce the incidence of powdery
mildew on cucumbers grown in the greenhouse (Fig. 8), suggesting the potential effectiveness
of the practical use of Oligo-Mix in cucumber cultivation.

3.6 Conclusion remarks

The study suggests that treatment with Oligo-Mix, a mixture of the DAMPs COS and XOS
and the MAMP CHOS, promotes both growth and disease resistance in cucumbers. Against
the rational trade-off theory, why Oligo-Mix did not suppress the growth of plants while
improving disease resistance? First, Oligo-Mix promoted root growth, which is expected to
have the effect of allowing plants to grow basically robust and underpin the energy that is spent
in the immune response. Second, the RNA-seq data indicated that the induced genes included
some transcriptional regulators (Table S1). Increased amounts of various transcriptional
regulators are expected to allow plants to respond rapidly and robustly to the next
environmental stresses. Such effect is called priming, and the disease resistance induced by the
Oligo-Mix may be at the level of priming (Conrath et al., 2002). Induction of priming is
presumed to be less energetic loss than actual resistance to a pathogen that exhibits a resistance
effect (e.g., accumulation of antimicrobial substances). In fact, hypersensitive cell death was
not induced by Oligo-Mix treatment alone but was induced after inoculation with the powdery
mildew (Fig. 4).

A mixture of substances promoting growth and weakly activating disease resistance would
be ideal combination as agricultural materials. Biostimulant materials, however, require further
research and study to accumulate experiences for their effective use in agriculture, since
different crops (and different cultivars) often have variable sensitivities to such substances.

Funding

This work was supported partially by a Grant-in-Aid for Scientific Research (B) (23H02212)

to DT from the Japan Society for the Promotion of Science, and donation from Resonac

Corporation (former Showa Denko K.K., Tokyo, Japan) and grant R00GM135487 to OAM

from the National Institute of General Medical Sciences, NIGMS, USA.

Acknowledgment

We are grateful to Ms. Yuki Iwamoto (Nagoya University) for her technical support. We also

express our gratitude to Prof. Yoshitaka Takano (Kyoto University) for providing

Colletotrichum orbiculare. We would like to acknowledge the Japanese Ministry of Education,

Culture, Sports, and Technology (MEXT) for supporting Pring Sreynich in her pursuits. study

in Japan on scholarship.

Figure legends

Fig. 1. Oligo-Mix treatment can promote the growth of cucumber.

(A) Effect of Oligo-Mix treatment on seed germination of cucumber. Cucumber seeds were

placed on laboratory tissue soaked with distilled water in Petri dishes, and germinated seeds

were sprayed with control solution or Oligo-Mix (containing 20 µg/ml cello-oligosaccharide

[COS], 40 µgl/m1 xylo-oligosaccharide [XOS], and 20 µg/ml chitin-oligosaccharide [CHOS]).

The number of roots and root length (longest root) were measured 2 days after spray. **(B)** Effect

of Oligo-Mix treatment on growth of cucumber. Cucumber plants (28 days old) were sprayed

with control solution or Oligo-Mix every week (4 times), and fresh weight of above-ground

tissue (left) and root (right) were measured 7 days after the last treatment. (middle) Chlorophyll

contents in cucumber leaves were quantified according to the method described in Porra and

Scheer (1989). Asterisks indicate a significant difference from control as assessed by a two-

tailed Student's t-test. * $p < 0.05$, ** $p < 0.01$.

Fig. 2. Oligo-Mix treatment induces activation of MAP kinase and callose deposition of

cucumber.

(A) Cucumber or Arabidopsis seedlings were treated with control solution (Cont.) or Oligo-

Mix (containing 20 µg/ml cello-oligosaccharide, 40 µg/m1 xylo-oligosaccharide and 20 µg/ml

chitin-oligosaccharide). The phosphorylation of MAP kinases was determined by western blot

using phospho-p44/42 MAPK (Erk1/2; Thr-202/Tyr-204) antibody. **(B)** Cucumber seedlings

were treated with control solution (Cont.) or Oligo-Mix and callose deposition was visualized
by aniline blue staining. The fluorescence spots were counted 24 h after treatment under a
fluorescence microscope (n = 12). Bar = 200 μm.

Fig. 3. Oligo-Mix induces disease resistance against powdery mildew (*Podosphaera xanthii*)
on cucumber. Cucumber plants (4-weeks-old) were treated with control solution (Cont.) or
Oligo-Mix (containing 20 μg/ml cello-oligosaccharide, 40 μg/m1 xylo-oligosaccharide and 20
μg/ml chitin-oligosaccharide) and spray-inoculated with powdery mildew spore suspension
(1×10^5 spores/ml, 0.5 ml/plant) at 24 h after treatment. The number of powdery mildew colony
types (spored, yellowish , or cell death) per plant was counted one week after the inoculation
(n = 7). Asterisks indicate a significant difference from the control as assessed by a two-tailed
Student's *t*-test. * $p < 0.05$. n.s., not significant.

Fig. 4. Pretreatment of Oligo-Mix can promote the resistance of cucumber against powdery
mildew pathogen *Podosphaera xanthii*. Cucumber plants (4-weeks-old) were treated with
control solution (Cont.) or Oligo-Mix (containing 20 μg/ml cello-oligosaccharide, 40 μg/m1
xylo-oligosaccharide and 20 μg/ml chitin-oligosaccharide) and spray-inoculated with powdery
mildew spore suspension (1×10^5 spores/ml, 0.5 ml/plant) at 24 h after treatment. Inoculated
leaves were stained with lacto-phenol trypan blue 8 days after inoculation. **(A)** Extensive
hyphal growth and conidiophores formation were observed on cucumber leaves 8 days after
the inoculation, while induction of cell death was detected for the leaves pretreated with Oligo-
Mix. Bars = 200 μm. **(B)** Fully developed conidiophores (c) and haustoria (h) were observed
on the leaves of control plants. Local cell death (arrowheads) at the sites of
penetration/haustoria formation of *P. xanthii* was detected on cucumber leaves pretreated with
Oligo-Mix. Bars = 50 μm. **(C)** Number of conidia formed on leaves (with symptoms) was
counted 7 days after the inoculation (n = 16). Asterisks indicate a significant difference from
the control as assessed by a two-tailed Student's *t*-test. ** $p < 0.01$.

Fig. 5. Oligo-Mix induces disease resistance of cucumber against anthracnose disease by
Colletotrichum orbiculare. **(A)** Cucumber plants (4-weeks-old) were treated with control
solution (Cont.) or Oligo-Mix (containing 20 μg/ml cello-oligosaccharide, 40 μg/m1 xylo-
oligosaccharide and 20 μg/ml chitin-oligosaccharide) and spray-inoculated with spore
suspension of *C. orbiculare* (1×10^5 spores/ml, 0.5 ml/plant) at 24 h after treatment.
Photographed 7 days after inoculation. **(B)** The diseased area of leaves was detected using an

image analysis algorithm (See the method section) (n = 10). Asterisks indicate a significant

difference from the control as assessed by a two-tailed Student's *t*-test. * $p < 0.05$.

Fig. 6. Microscopic observation of anthracnose infection on cucumber with or without Oligo-

Mix treatment. Cucumber plants were treated with control solution (Cont.) or Oligo-Mix (20

µg/ml COS, 40 µg/ml XOS, 20 µg/ml CHOS) and conidial suspension of *Colletotrichum*

orbiculare (1x10^5 spores/ml, 0.5 ml/plant) were spray-inoculated. Inoculated leaves were

stained with lacto-phenol trypan blue 8 days after the inoculation. In control cucumber leaves

(left), intracellular biotrophic hyphae (white arrowheads) growing out from necrotic lesions

(black arrowheads) and developing acervulus (a) with multiple setae (s) were often observed.

In Oligo-Mix treated leaves, biotrophic hyphae were hardly detected and fewer and less-

developed acervulus were observed. Bars = 50 µm.

Fig. 7. Expression profiles of representative cucumber genes upregulated by treatment with

Oligo-Mix. Gene expression (TPM value) was determined by RNA-seq analysis of cucumber

leaves treated with control solution (Cont.) or Oligo-Mix (20 µg/ml COS, 40 µg/ml XOS, 20

µg/ml CHOS) for 24 h. **(A)** Genes related to cell wall reinforcement. **(B)** Genes related to the

responses to abiotic stresses. **(C)** Genes related to plant defense. Data marked with asterisks

are significantly upregulated from control as determined by two-tailed Student's *t*-test ** p

<0.01, * p <0.05. See Table S1 for the list of upregulated genes by Oligo-Mix treatment.

Fig. 8. Effect of Oligo-Mix treatment on the appearance of powdery mildew on cucumber

grown in greenhouse. Cucmber plant was sprayed control solution (Cont.) or Oligo-Mix (20

µg/ml COS, 40 µg/ml XOS, 20 µg/ml CHOS) once every week (6 times altogether) and

naturally allowed to get infected by powdery mildew. The number of colonies per plant was

counted at 6 weeks after first treatment. Data are shown as mean ± SD (n = 14).

References

A. Ashida, M. Camagna, I. Sato, S. Chiba, A. Tanaka, D. Takemoto, Draft genome sequences
of *Neofusicoccum parvum* strains isolated from mango and rice-paper plant (*Tetrapanax*
papyrifer), Microbiol. Resour. Announc. 12 (2023) e0025923,
https://doi.org/10.1128/mra.00259-23.

A.S. Bulasag, M. Camagna, T. Kuroyanagi, A. Ashida, K. Ito, A. Tanaka, I. Sato, S. Chiba,
M. Ojika, D. Takemoto, *Botrytis cinerea* tolerates phytoalexins produced by Solanaceae and

Fabaceae plants through an efflux transporter BcatrB and metabolizing enzymes. Front. Plant Sci. 14 (2023) 1177060, https://doi.org/10.3389/fpls.2023.1177060.

J. Claverie, S. Balacey, C. Lemaître-Guillier, D. Brulé, A. Chiltz, L. Granet, E. Noirot, X. Daire, B. Darblade, M.C. Héloir, B. Poinssot, The cell wall-derived xyloglucan is a new DAMP triggering plant immunity in *Vitis vinifera* and *Arabidopsis thaliana*, Front. Plant Sci. 9 (2018) 1725, https://doi.org/10.3389/fpls.2018.01725.

U. Conrath, C.M. Pieterse, B. Mauch-Mani, Priming in plant-pathogen interactions, Trends Plant Sci. 7 (2002) 210–216, https://doi.org/10.1016/s1360-1385(02)02244-6.

D.J. Cosgrove, Growth of the plant cell wall, Nat. Rev. Mol. Cell Biol. 6 (2005) 850–861, https://doi.org/10.1038/nrm1746.

C. de Jesús-Pires, J.R.C. Ferreira-Neto, J. Pacifico Bezerra-Neto, E.A. Kido, R.L. de Oliveira Silva, V. Pandolfi, A.C. Wanderley-Nogueira, E. Binneck, A.F. da Costa, G. Pio-Ribeiro, G. Pereira-Andrade, I.M. Sittolin, F. Freire-Filho, A.M. Benko-Iseppon, Plant thaumatin-like proteins: Function, evolution and biotechnological applications, Curr. Protein Pept. Sci. 21 (2020) 36–51, https://doi.org/10.2174/1389203720666190318164905.

J.F. Dean, K.C. Gross, J.D. Anderson, Ethylene biosynthesis-inducing xylanase : III. Product characterization, Plant Physiol. 96 (1991) 571–576, https://doi.org/10.1104/pp.96.2.571.

L. Derevnina, Y.F. Dagdas, J.C. de la Concepcion, A. Bialas, R. Kellner, B. Petre, E. Domazakis, J. Du, C.H. Wu, X. Lin, C. Aguilera-Galvez, N. Cruz-Mireles, V.G. Vleeshouwers, S. Kamoun, Nine things to know about elicitins, New Phytol. 212 (2016) 888–895, https://doi.org/10.1111/nph.14137.

J.M. Eklöf, H. Brumer, The *XTH* gene family: an update on enzyme structure, function, and phylogeny in xyloglucan remodeling, Plant Physiol. 153 (2010) 456–466, https://doi.org/10.1104/pp.110.156844.

D. Ellinger, C.A. Voigt, Callose biosynthesis in Arabidopsis with a focus on pathogen response: what we have learned within the last decade, Ann. Bot. 114 (2014) 1349–1358, https://doi.org/10.1093/aob/mcu120.

G. Felix, M. Regenass, T. Boller, Specific perception of subnanomolar concentrations of chitin fragments by tomato cells: Induction of extracellular alkalinization, changes in protein phosphorylation, and establishment of a refractory state, Plant J. 4 (1993) 307–316, https://doi.org/10.1046/j.1365-313X.1993.04020307.x.

J. Fliegmann, G. Felix, Immunity: Flagellin seen from all sides, Nat. Plants 2 (2016) 16136, https://doi.org/10.1038/nplants.2016.136.

L. Gómez-Gómez, G. Felix, T. Boller, A single locus determines sensitivity to bacterial flagellin in *Arabidopsis thaliana*, Plant J. 18 (1999) 277–284, https://doi.org/10.1046/j.1365-313x.1999.00451.x.

N.A.R. Gow, M.D. Lenardon, Architecture of the dynamic fungal cell wall, Nat. Rev. Microbiol. 21 (2023) 248–259, https://doi.org/10.1038/s41579-022-00796-9.

A.A. Gust, R. Biswas, H.D. Lenz, T. Rauhut, S. Ranf, B. Kemmerling, F. Götz, E. Glawischnig, J. Lee, G. Felix, T. Nürnberger, Bacteria-derived peptidoglycans constitute pathogen-associated molecular patterns triggering innate immunity in Arabidopsis, J. Biol. Chem. 282 (2007) 32338–32348, https://doi.org/10.1074/jbc.M704886200.

T. Hander, Á.D. Fernández-Fernández, R.P. Kumpf, P. Willems, H. Schatowitz, D. Rombaut, A. Staes, J. Nolf, R. Pottie, P. Yao, A. Gonçalves, B. Pavie, T. Boller, K. Gevaert, F Van Breusegem, S. Bartels, S. Stael, Damage on plants activates Ca_2^+-dependent metacaspases for release of immunomodulatory peptides, Science 363 (2019) eaar7486, https://doi.org/10.1126/science.aar7486.

K. Herbert, G. Mönke, R. Badur, U. Sonnewald, A simplified procedure for the subtractive cDNA cloning of photoassimilate-responding genes: isolation of cDNAs encoding a new class of pathogenesis-related proteins, Plant Mol. Biol. 29 (1995) 1027–1038, https://doi.org/10.1007/BF00014975.

S. Imano, M. Fushimi, M. Camagna, A. Tsuyama-Koike, H. Mori, A. Ashida, A. Tanaka, I. Sato, S. Chiba, K. Kawakita, M. Ojika, D. Takemoto, AP2/ERF transcription factor NbERF-IX-33 is involved in the regulation of phytoalexin production for the resistance of *Nicotiana benthamiana* to *Phytophthora infestans*, Front. Plant Sci. 12 (2022) 821574, https://doi.org/10.3389/fpls.2021.821574.

T.L. Karasov, E. Chae, J.J. Herman, J. Bergelson, Mechanisms to mitigate the trade-off between growth and defense, Plant Cell 29 (2017) 666–680, https://doi.org/10.1105/tpc.16.00931.

S. Kamoun, M. Young, C.B. Glascock, B.M. Tyler, Extracellular protein elicitors from *Phytophthora*: host-specificity and induction of resistance to bacterial and fungal phytopathogens, Mol. Plant Microbe Interact. 6 (1993) 15–25, https://doi.org/10.1094/MPMI-6-015.

H. Kato, K. Nemoto, M. Shimizu, A. Abe, S. Asai, N. Ishihama, S. Matsuoka, T. Daimon, M. Ojika, K. Kawakita, K. Onai, K. Shirasu, M. Yoshida, M. Ishiura, D. Takemoto, Y. Takano, R. Terauchi, Recognition of pathogen-derived sphingolipids in Arabidopsis, Science 376 (2022) 857–860, https://doi.org/10.1126/science.abn0650.

D. Kim, J.M. Paggi, C. Park, C. Bennett, S.L. Salzberg, Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype, Nat. Biotechnol. 37 (2019) 907–915, https://doi.org/10.1038/s41587-019-0201-4.

S. Kovaka, A.V. Zimin, G.M. Pertea, R. Razaghi, S.L. Salzberg, M. Pertea, Transcriptome assembly from long-read RNA-seq alignments with StringTie2, Genome Biol. 20 (2019) 278, https://doi.org/10.1186/s13059-019-1910-1.

T. Kuroyanagi, A.S. Bulasag, K. Fukushima, A. Ashida, T. Suzuki, A. Tanaka, M. Camagna, I. Sato, S. Chiba, M. Ojika, D. Takemoto, *Botrytis cinerea* identifies host plants via the

recognition of antifungal capsidiol to induce expression of a specific detoxification gene, PNAS Nexus 1 (2022) pgac274, https://doi.org/10.1093/pnasnexus/pgac274.

A. Lebeda, E. Křístková, B. Sedláková, M.D. Coffey, J.D. McCreight, Gaps and perspectives of pathotype and race determination in *Golovinomyces cichoracearum* and *Podosphaera xanthii*, Mycoscience 52 (2011) 159–164, https://doi.org/10.1007/S10267-010-0098-8.

W. Liang, C. Li, F. Liu, H. Jiang, S. Li, J. Sun, X. Wu, C. Li, The Arabidopsis homologs of CCR4-associated factor 1 show mRNA deadenylation activity and play a role in plant defence responses, Cell Res. 19 (2009) 307–316, https://doi.org/10.1038/cr.2008.317.

S. Lin, Y. Miao, H. Huang, Y. Zhang, L. Huang, J. Cao, Arabinogalactan proteins: Focus on the role in cellulose synthesis and deposition during plant cell wall biogenesis, Int. J. Mol. Sci. 23 (2022) 6578, https://doi.org/10.3390/ijms23126578.

M.I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2, Genome Biol. 15 (2014) 550, https://doi.org/10.1186/s13059-014-0550-8.

M. Martin, Cutadapt removes adapter sequences from high-throughput sequencing reads, EMBnet J. 17 (2011) 10–12, https://doi.org/10.14806/ej.17.1.200.

H. Matsuo, Y. Ishiga, Y. Kubo, Y. Yoshioka, *Colletotrichum orbiculare* strains distributed in Japan: race identification and evaluation of virulence to cucurbits, Breed Sci. 72 (2022) 306–315, https://doi.org/10.1270/jsbbs.22011.

G. Pearce, D. Strydom, S. Johnson, C.A. Ryan, A polypeptide from tomato leaves induces wound-inducible proteinase inhibitor proteins, Science 253 (1991) 895–897, https://doi.org/10.1126/science.253.5022.895.

R.J. Porra, H. Scheer, Towards a more accurate future for chlorophyll a and b determinations: the inaccuracies of Daniel Arnon's assay, Photosynth Res. 140 (1989) 215–219, https://doi.org/10.1007/s11120-018-0579-8.

S. Pring, H. Kato, S. Imano, M. Camagna, A. Tanaka, H. Kimoto, P. Chen, A. Shrotri, H. Kobayashi, A. Fukuoka, M. Saito, T. Suzuki, R. Terauchi, I. Sato, S. Chiba, D. Takemoto, Induction of plant disease resistance by mixed oligosaccharide elicitors prepared from plant cell wall and crustacean shells, Physiol. Plant. 175 (2023) e14052, https://doi.org/10.1111/ppl.14052.

M. Ron, A. Avni, The receptor for the fungal elicitor ethylene-inducing xylanase is a member of a resistance-like gene family in tomato, Plant Cell 16 (2004) 1604–1615, https://doi.org/10.1105/tpc.022475.

O. Rowland, A.A. Ludwig, C.J. Merrick, F. Baillieul, F.E. Tracy, W.E. Durrant, L. Fritz-Laylin, V. Nekrasov, K. Sjölander, H. Yoshioka, J.D.G. Jones, Functional analysis of Avr9/Cf-9 rapidly elicited genes identifies a protein kinase, ACIK1, that is essential for full Cf-9-dependent disease resistance in tomato, Plant Cell 17 (2005) 295–310, https://doi.org/10.1105/tpc.104.026013.

C.A. Schneider, W.S. Rasband, K.W. Eliceiri, NIH Image to ImageJ: 25 years of image
analysis, Nat. Methods 9 (2012) 671–675, https://doi.org/10.1038/nmeth.2089.

F. Schröder, J. Lisso, P. Lange, C. Müssig, The extracellular EXO protein mediates cell
expansion in Arabidopsis leaves, BMC Plant Biol. 9 (2009) 20, https://doi.org/10.1186/1471-
2229-9-20.

J.K. Sharp, B. Valent, P. Albersheim, Purification and partial characterization of a beta-glucan
fragment that elicits phytoalexin accumulation in soybean, J. Biol. Chem. 259 (1984) 11312–
11320, https://doi.org/10.1016/S0021-9258(18)90864-1.

Y. Shibata, M. Ojika, A. Sugiyama, K Yazaki, D.A. Jones, K. Kawakita, D. Takemoto, The
full-size ABCG transporters Nb-ABCG1 and Nb-ABCG2 function in pre- and postinvasion
defense against *Phytophthora infestans* in *Nicotiana benthamiana*, Plant Cell 28 (2016) 1163–
1181, https://doi.org/10.1105/tpc.15.00721.

C.A. Souza, S. Li, A.Z. Lin, F. Boutrot, G. Grossmann, C. Zipfel, S.C. Somerville, Cellulose-
derived oligomers act as damage-associated molecular patterns and trigger defense-like
responses, Plant Physiol. 173 (2017) 2383–2398, https://doi.org/10.1104/pp.16.01680.

C. Stringer, T. Wang, M. Michaelos, M. Pachitariu, Cellpose: a generalist algorithm for cellular
segmentation, Nat. Methods 18(2021) 100–106, https://doi.org/10.1038/s41592-020-01018-x.

D. Takemoto, A.R. Hardham, D.A. Jones, Differences in cell death induction by *Phytophthora*
Elicitins are determined by signal components downstream of MAP kinase kinase in different
species of *Nicotiana* and cultivars of *Brassica rapa* and *Raphanus sativus*, Plant Physiol. 138
(2005) 1491–1504, https://doi.org/10.1104/pp.104.058388.

D. Takemoto, D.A. Jones, A.R. Hardham, GFP-tagging of cell components reveals the
dynamics of subcellular re-organization in response to infection of Arabidopsis by oomycete
pathogens, Plant J. 33 (2003) 775–792, https://doi.org/10.1046/j.1365-313x.2003.01673.x.

D. Takemoto, H. Yoshioka, N. Doke, K. Kawakita, Disease stress-inducible genes of tobacco:
expression profile of elicitor-responsive genes isolated by subtractive hybridization, Physiol.
Plant. 118 (2003) 545–553, https://doi.org/10.1034/j.1399-3054.2003.00145.x.

A. Vielba-Fernández, Á. Polonio, L. Ruiz-Jiménez, A. de Vicente, A. Pérez-García, D.
Fernández-Ortuño, Fungicide resistance in powdery mildew fungi, Microorganisms 8 (2020)
1431, https://doi.org/10.3390/microorganisms8091431.

L. Wang, E. Einig, M. Almeida-Trapp, M. Albert, J. Fliegmann, A. Mithöfer, H. Kalbacher, G.
Felix, The systemin receptor SYR1 enhances resistance of tomato against herbivorous insects,
Nat. Plants 4 (2018) 152–156, https://doi.org/10.1038/s41477-018-0106-0.

Y. Yamaguchi, A. Huffaker, A.C. Bryan, F.E. Tax, C.A. Ryan, PEPR2 is a second receptor for
the Pep1 and Pep2 peptides and contributes to defense responses in Arabidopsis, Plant Cell 22
(2010) 508–522, https://doi.org/10.1105/tpc.109.068874.

Y. Yamauchi, A. Matsuda, N. Matsuura, M. Mizutani, Y. Sugimoto, Transcriptome analysis
of *Arabidopsis thaliana* treated with green leaf volatiles: possible role of green leaf volatiles

as self-made damage-associated molecular patterns, J. Pestic. Sci. 43 (2018) 207–213,

https://doi.org/10.1584/jpestics.D18-020.

Fig. 1. Oligo-Mix treatment can promote the growth of cucumber.
(A) Effect of Oligo-Mix treatment on seed germination of cucumber. Cucumber seeds were placed on laboratory tissue soaked with distilled water in Petri dishes, and germinated seeds were sprayed with control solution or Oligo-Mix (containing 20 µg/ml cello-oligosaccharide [COS], 40 µg/ml xylo-oligosaccharide [XOS], and 20 µg/ml chitin-oligosaccharide [CHOS]). The number of roots and root length (longest root) were measured 2 days after spray. **(B)** Effect of Oligo-Mix treatment on growth of cucumber. Cucumber plants (28 days old) were sprayed with control solution or Oligo-Mix every week (4 times), and fresh weight of above-ground tissue (left) and root (right) were measured 7 days after the last treatment. (middle) Chlorophyll contents in cucumber leaves were quantified according to the method described in Porra and Scheer (1989). Asterisks indicate a significant difference from control as assessed by a two-tailed Student's t-test. * $p < 0.05$, ** $p < 0.01$.

Fig. 2. Oligo-Mix treatment induces activation of MAP kinase and callose deposition of cucumber.

(A) Cucumber or Arabidopsis seedlings were treated with control solution (Cont.) or Oligo-Mix (containing 20 μg/ml cello-oligosaccharide, 40 μg/m1 xylo-oligosaccharide and 20 μg/ml chitin-oligosaccharide). The phosphorylation of MAP kinases was determined by western blot using phospho-p44/42 MAPK (Erk1/2; Thr-202/Tyr-204) antibody. **(B)** Cucumber seedlings were treated with control solution (Cont.) or Oligo-Mix and callose deposition was visualized by aniline blue staining. The fluorescence spots were counted 24 h after treatment under a fluorescence microscope (n = 12). Bar = 200 μm.

Fig. 3. Oligo-Mix induces disease resistance against powdery mildew (*Podosphaera xanthii*) on cucumber. Cucumber plants (4-weeks-old) were treated with control solution (Cont.) or Oligo-Mix (containing 20 µg/ml cello-oligosaccharide, 40 µg/ml xylo-oligosaccharide and 20 µg/ml chitin-oligosaccharide) and spray-inoculated with powdery mildew spore suspension (1×10^5 spores/ml, 0.5 ml/plant) at 24 h after treatment. The number of powdery mildew colony types (spored, yellowish , or cell death) per plant was counted one week after the inoculation (n = 7). Asterisks indicate a significant difference from the control as assessed by a two-tailed Student's *t*-test. * $p < 0.05$. n.s., not significant.

Fig. 4. Pretreatment of Oligo-Mix can promote the resistance of cucumber against powdery mildew pathogen *Podosphaera xanthii*. Cucumber plants (4-weeks-old) were treated with control solution (Cont.) or Oligo-Mix (containing 20 μg/ml cello-oligosaccharide, 40 μg/ml xylo-oligosaccharide and 20 μg/ml chitin-oligosaccharide) and spray-inoculated with powdery mildew spore suspension (1×10^5 spores/ml, 0.5 ml/plant) at 24 h after treatment. Inoculated leaves were stained with lacto-phenol trypan blue 8 days after inoculation. **(A)** Extensive hyphal growth and conidiophores formation were observed on cucumber leaves 8 days after the inoculation, while induction of cell death was detected for the leaves pretreated with Oligo-Mix. Bars = 200 μm. **(B)** Fully developed conidiophores (c) and haustoria (h) were observed on the leaves of control plants. Local cell death (arrowheads) at the sites of penetration/haustoria formation of *P. xanthii* was detected on cucumber leaves pretreated with Oligo-Mix. Bars = 50 μm. **(C)** Number of conidia formed on leaves (with symptoms) was counted 7 days after the inoculation (n = 16). Asterisks indicate a significant difference from the control as assessed by a two-tailed Student's *t*-test. ** $p < 0.01$.

Fig. 5. Oligo-Mix induces disease resistance of cucumber against anthracnose disease by *Colletotrichum orbiculare*. **(A)** Cucumber plants (4-weeks-old) were treated with control solution (Cont.) or Oligo-Mix (containing 20 µg/ml cello-oligosaccharide, 40 µg/m1 xylo-oligosaccharide and 20 µg/ml chitin-oligosaccharide) and spray-inoculated with spore suspension of *C. orbiculare* (1×10^5 spores/ml, 0.5 ml/plant) at 24 h after treatment. Photographed 7 days after inoculation. **(B)** The diseased area of leaves was detected using an image analysis algorithm (See the method section) (n = 10). Asterisks indicate a significant difference from the control as assessed by a two-tailed Student's *t*-test. * $p <$ 0.05.

Fig. 6. Microscopic observation of anthracnose infection on cucumber with or without Oligo-Mix treatment. Cucumber plants were treated with control solution (Cont.) or Oligo-Mix (20 µg/ml COS, 40 µg/ml XOS, 20 µg/ml CHOS) and conidial suspension of *Colletotrichum orbiculare* ($1x10^5$ spores/ml, 0.5 ml/plant) were spray-inoculated. Inoculated leaves were stained with lacto-phenol trypan blue 8 days after the inoculation. In control cucumber leaves (left), intracellular biotrophic hyphae (white arrowheads) growing out from necrotic lesions (black arrowheads) and developing acervulus (a) with multiple setae (s) were often observed. In Oligo-Mix treated leaves, biotrophic hyphae were hardly detected and fewer and less-developed acervulus were observed. Bars = 50 µm.

Fig. 7. Expression profiles of representative cucumber genes upregulated by treatment with Oligo-Mix. Gene expression (TPM value) was determined by RNA-seq analysis of cucumber leaves treated with control solution (Cont.) or Oligo-Mix (20 µg/ml COS, 40 µg/ml XOS, 20 µg/ml CHOS) for 24 h. **(A)** Genes related to cell wall reinforcement. **(B)** Genes related to the responses to abiotic stresses. **(C)** Genes related to plant defense. Data marked with asterisks are significantly upregulated from control as determined by two-tailed Student's *t*-test ** *p* <0.01, * *p* <0.05. See Table S1 for the list of upregulated genes by Oligo-Mix treatment.

Fig. 8. Effect of Oligo-Mix treatment on the appearance of powdery mildew on cucumber grown in greenhouse. Cucmber plant was sprayed control solution (Cont.) or Oligo-Mix (20 µg/ml COS, 40 µg/ml XOS, 20 µg/ml CHOS) once every week (6 times altogether) and naturally allowed to get infected by powdery mildew. The number of colonies per plant was counted at 6 weeks after first treatment. Data are shown as mean ± SD (n = 14).