A Blueprint of Sex-Specific Neuronal Regulation in *the C. elegans* Nervous System at Single-Cell Resolution

Rizwanul Haque[1,2,*], Hagar Setty[1,2,*], Ramiro Lorenzo[3,*], Gil Stelzer[4], Ron Rotkopf[4], Yehuda

Salzberg[1,2], Gal Goldman[1], Sandeep Kumar[5], Shiraz Nir Halber[1,2], Andrew M. Leifer[5,6], Elad

Schneidman[1], Patrick Laurent[3] and Meital Oren-Suissa[1,2]

[1]Department of Brain Sciences, Weizmann Institute of Science, Rehovot 76100, Israel

[2]Department of Molecular Neuroscience, Weizmann Institute of Science, Rehovot, Israel

[3]Laboratory of Neurophysiology, ULB Neuroscience Institute (UNI), Université Libre de Bruxelles

(ULB), Brussels, Belgium

[4]Bioinformatics Unit- Life Science Core Facilities, Weizmann Institute of Science, Israel

[5]Princeton Neuroscience Institute, Princeton University, Princeton, NJ 08544, USA

[6]Department of Physics, Princeton University, Princeton, NJ 08544, USA

*Equal contribution

ABSTRACT

Sex-specific behaviors within a species are often attributed to variances in neuronal wiring and molecular signatures. However, how the genetic sex shapes the molecular architecture of the nervous system at a single neuron level is still unknown. To address this gap, we used single-cell RNA-sequencing (scRNA-seq) to profile the transcriptome of the sex-shared nervous system of adult male and hermaphrodite *Caenorhabditis elegans*. By ranking neurons based on the degree of molecular dimorphism, we uncover novel sexually dimorphic neurons such as PLM neurons, where functional dimorphism was corroborated by diminished posterior touch responses in hermaphrodites. We identify sex-specific regulators of mechanosensory behavior and neuronal function by combining our dataset with reverse genetic screens. While most sex-shared neurons retain their neurotransmitter identity, the male neuropeptide connectome undergoes substantial remodeling, with most of the neuropeptides showing a male-bias expression. This reinforces the notion that neuropeptides are crucial for diversifying connectome outputs. Furthermore, by correlating gene expression with outgoing synaptic connectivity, we identified regulatory candidates for synaptic wiring, including both shared and sex-specific genes. This comprehensive resource provides foundational insights into the molecular drivers of sexual dimorphism, facilitating future exploration of regulatory mechanisms and their impact on sex-specific behaviors in higher organisms.

INTRODUCTION

The molecular mechanisms that underlie the development of sex differences in the nervous system have been the focus of many studies in the past decade, especially due to the sex bias demonstrated in the prevalence and progression of several neurological diseases such as anxiety and Parkinson's disease[1–3]. Recent studies have made an effort to dissect sexually dimorphic neuronal circuits, uncovering unique circuits across the animal kingdom that operate differently between the sexes to drive behavior (termed "sexually dimorphic"); such as the circuit for empathic behaviors in mice, aggressive behaviors in fruit flies and nociceptive and mechanosensory behaviors in worms[4–7]. While these studies have managed to provide insights into how the nervous system is differentially shaped at the level of specific circuits, a complete picture of the molecular repertoire of the entire nervous system in both sexes is still lacking. With the advancement of RNA sequencing, molecular profiles of both sexes in mice were generated, discovering differentially expressed genes between the sexes in specific neuronal populations such as the medial amygdala (MeA) and the ventromedial nucleus of the hypothalamus (VMH)[8–11]. Additional studies performed transcriptomic profiling of different brain regions while delineating the effect of circulating hormones on sex differences in gene expression[12–14]. Although these studies provided insights regarding sexually dimorphic behaviors, they did not explore the contribution of specific genes to mediate these behaviors. Additionally, the hormonal regulation in mammalian systems might mask the direct effect of sexual identity (the chromosome complement) on distinct molecular signatures at the level of single neurons.

In the nematode *Caenorhabditis elegans* (*C. elegans*), the genetic sex of each somatic cell, including neurons, is determined in a cell-autonomous manner: the ratio of sex-chromosomes to

autosomes controls a transcriptional cascade which results in the expression of either

hermaphrodite-specific or male-specific genes[15,16]. Recent studies explored the contribution of

the genetic sex to nervous system development, with insights into synaptic connectivity, circuitry,

function, and behavior[7,17–19]. Despite all these efforts, we still lack a complete picture of how

genetic sex shapes gene expression patterns across the nervous system and the different neuronal

classes.

Recently, single-cell RNA sequencing (scRNA-seq) approaches resolved the molecular

heterogeneity of the hermaphrodite nervous system. Its 302 neurons were subdivided into 128

transcriptionally distinct neuron types[20]. The transcriptomic signatures of each neuron class

facilitated the identification of markers, unraveled mechanisms of cell specification, and enabled

deeper characterization of nervous system development and aging[20–29]. Earlier scRNA-seq

focused only on hermaphrodites (with one exception for glia[30]), limiting our knowledge of the

molecular signatures of male neurons and, most importantly, the differences in molecular

signatures between the sexes. To gain a broad perspective on the molecular landscape of the

nervous system in both sexes of *C. elegans*, we generated comparative scRNA-seq transcriptomic

profiles   for   the   sex-shared   neuron   types   in   males   and   hermaphrodites

(https://www.weizmann.ac.il/dimorgena/).

Our research provides a comprehensive atlas for expression patterns of 18,975 genes in the

nervous system of adult males and hermaphrodites. This atlas revealed a variety of neuronal

gene classes with unique dimorphic gene expression patterns, with most of the neuropeptide-

encoding genes demonstrating robust sexually dimorphic gene expression. We focused on the

core, sex-shared nervous system i.e., neurons that exist in both sexes, and uncovered several sex-

shared neurons with novel dimorphic transcriptomic profiles, including the sensory touch

receptor neuron PLM. We harnessed our dataset to predict genes that might regulate

connectivity and revealed novel candidates that operate in both sexes and in a sex-specific

manner to control synaptic wiring.

RESULTS

Single-cell sequencing of adult hermaphrodite and male nervous system

To generate a single-cell transcriptomic profile of the nervous system in both sexes of *C. elegans*

we dissociated young adult males and hermaphrodites expressing TagRFP pan-neuronally into

single cells (Supplementary Figure 1A). To generate pure male populations (XO animals), we used

our recently published protocol that enables large-scale isolation of males with high purity[31]. To

isolate the fluorescently labeled neurons, we dissociated young adult animals into single cells and

sorted the single-cell suspensions by Fluorescence-Activated-Cell-Sorting (FACS) (Figure 1A,

Supplementary Figure 1B-C, *See Methods*). We confirmed the neuronal enrichment and the intact

morphology of RFP+ FACS-purified cells (Supplementary Figure 1D-G, *See Methods*). Single-cell

RNA sequencing was done using a microfluidic-based 10x genomics platform. We corrected for

ambient RNA, removed low-quality and damaged cells, and performed cell-quality filtering (*See*

Methods). This yielded a median of 925 unique molecular identifiers (UMIs) and 358 genes per

cell after filtering. To filter neurons from non-neuronal cells, we used a cutoff based on known

molecular landmarks (563 neuronal and 114 non-neuronal genes, Supplemental Table 1). After

the removal of cells containing more than 20% of UMIs originating from mitochondrial genes, we

obtained 13015 single-cell transcriptomes for hermaphrodites and 19454 single-cell

transcriptomes for males, taken for further analysis and cluster annotation.

To simplify cluster annotation, we merged our datasets with the previously annotated CenGEN

dataset[20]. This combined dataset was first segregated into 72 partitions (Supplementary Figure

2A). Sub-clustering of partition 1 (41219 cells) and partitions 2-72 (61546 cells) yielded 118

clusters (Supplementary Figure 2B-C). A further iteration included manual assignment of varying

identities, producing an overall of 156 clusters (details in Methods). 16 clusters were only present

in males and 13 were classified as "unknown ID" due to lack of previous data. We annotated

clusters using previously assigned cell identities[20] and manually assigned neuronal identities to

88 sex-shared clusters (with at least 5 cells per cluster in both sexes, Figure 1B-C). Despite the

underrepresentation of some neuron classes (e.g., ASH and PHB) due to limitations of cell sorting,

62.7% of the 118 anatomically defined neuron classes in the mature hermaphrodite nervous

system were captured. Interestingly, UMAP representations of sex-shared clusters of

hermaphrodites (adult and L4[20]) and adult males showed a large overlap (Figure 1C), with a strong

Spearman correlation (adult and L4, Supplementary Figure 3). When comparing UMAPs of

hermaphrodites (YA vs L4), we observed a subtle shift likely corresponding to the difference in

their developmental stage[31]. This analysis provides a framework for studying sexual dimorphism

in nervous system function.

Single-cell profiles of the nervous system capture dimorphic gene expression

In this work, we focused on the sex-shared clusters, and we did not further cluster the male-

specific neurons (Supplementary Table 2). For gene expression analysis, we calculated the fraction

of cells expressing a gene in each cluster, the average expression, and markers by using Seurat v2

with default parameters (*See Methods*, https://www.weizmann.ac.il/dimorgena/, Supplementary

Table 2). We also calculated expression thresholds as previously described ([20], *See Methods*) and

combined this information with the marker analysis. Predictive expression patterns across all

neurons in adult males and hermaphrodites can be mined using one of four thresholds of

increased stringency, as observed in Supplementary Table 2 (Figure 1D-E)**.** We characterized the

expression of genes with a known sex-biased expression. The Zn finger transcription factor LIN-

29 promotes a fate switch from a glutamatergic (hermaphrodite) to a cholinergic (male) identity

in the AIM neuron[32,33] (Figure 1F). Indeed, AIM cell clusters showed expression of *lin-29*, the

Choline Acetyltransferase *cho-1*/CHOT, and the vesicular Ach transporter *unc-17/VACHT* in male

clusters, while hermaphrodite clusters expressed the vesicular glutamate transporter *eat-*

4/VGLUT (Figure 1G)*.* We also observed sex-biased expression of *ins-39* in male AFD neurons and

of *mab-3* in male SMD neurons, as previously reported (Figure 1H-I)[31,32,34].

Figure 1: Single-cell RNA sequencing in both sexes of *C. elegans*. (A) Schematic of experimental design used in this study. Hermaphrodite and male *C. elegans* harboring the pan-neuronal marker *rab-3p(prom1)::2xNLS::TagRFP* were subjected to cell dissociation. To purify RFP+ cells, the cell suspension obtained was sorted by FACS. The single-cell RNA transcriptomes were obtained using single-cell droplet microfluidic technologies (10X Genomics, https://www.10xgenomics.com). (B) UMAP representation of all cell-clusters identified with their neuronal identity. Previously published[20] (CenGEN) datasets or manual assignments were both used to determine the IDs of the neurons. (C) UMAP representation of cell clusters with a shifted overlay to illustrate the overlap. Data sets are YA hermaphrodites (magenta), YA males (green), and CenGEN L4 hermaphrodites (yellow). (D) Receiver-Operator Characteristic (ROC) curve showing True Positive Rate (TPR) versus False Positive Rate (FPR) for hermaphrodites, males, hermaphrodites-males together, and CenGEN compared to ground truth expression data using the 65 cell types with 9 or more cells in hermaphrodite and male clusters. (E) Precision-Recall (PR) Curve of Recall (TPR) versus Precision [1 – False Discovery Rate (FDR)] curve. Red dots represent thresholds 1-4. Grey shading represents 95% confidence intervals. (F) Schematic of AIM neurons neurotransmitter switch from glutamatergic identity (in hermaphrodite) to cholinergic identity (in male). (G) AIM representation in the UMAP projection of all cells (inset marked as orange). UMAP projection depicting expression of *unc-17*, *cho-1* cholinergic identity genes, *eat-4* glutamatergic identity gene, and transcriptional factor *lin-29* gene within AIM cluster. Heatmap represents expression levels. (H) High expression of *ins-39* in males. AFD representation in the UMAP projection of all cells (inset marked as orange). UMAP projection depicting expression of *ins-39* gene within AFD cluster. Heatmap represents expression level. (I) Dimorphic expression of *mab-3*. Left, SMD representation in the UMAP projection of all cells (inset marked as orange). UMAP projection depicting expression of *mab-3* gene within SMD cluster. Heatmap represents expression level.

Comparative analysis of sex-shared neuronal clusters reveals novel dimorphisms

To uncover to what extent the transcriptomic profiles of the sex-shared nervous system diverge between males and hermaphrodites, we took a comparative approach, ranking the sex-shared neuronal clusters according to the number of differentially expressed genes (DEGs) (*See Methods*). For ranking purposes, we filtered out clusters with more than one identity, used a stricter cut-off of 9 cells per cluster in both sexes, and proceeded with 62 sex-shared clusters (*See Methods*). This analysis captured known dimorphic sex-shared neurons, including PHC, AFD, AIM,

and AVG[17,32–35] (Figure 2A). Strikingly, it also revealed many sex-shared neurons that were not

previously classified as dimorphic, such as the six touch-receptor-neurons (TRNs), BDU, and PVR

interneurons (Figure 2A).

Gene ontology (GO) enrichment analysis for the 2216 DEGs revealed a significant proportion of

enriched terms related to neuronal function, suggesting their potential relevance to neuronal

activities (Figure 2B). Additional GO terms such as synaptic function, heteromeric G protein, and

calcium signaling underscore the representation of genes that may play a role in shaping a

dimorphic nervous system (Figure 2B). By exploring neuronal DEGs based on their gene family,

we found that while multiple gene families differ between the sexes, neuropeptides,

neuropeptide receptors, and neuropeptide metabolism-related genes are overrepresented with

a significant male-biased expression of neuropeptides and neuropeptides metabolism-related

genes (58.5%, 92 of the 157 neuropeptide genes are DEGs) (Figure 2C-D, Figure 3A,

Supplementary Figure 5A-B).

We validated the predicted sexual dimorphism in the expression pattern of two broadly expressed

neuropeptide genes: *nlp-11* and *flp-20,* focusing on clusters within the head or tail regions. *nlp-*

11 showed dimorphic expression in 12 out of 16 predicted sex-shared neurons and in several

male-specific neurons (Figure 3B-F, Supplementary Figure 4A-B). The remaining four neurons

were either located in the midbody or difficult to identify. Similarly, *flp-20* showed sexually

dimorphic expression in four sex-shared neurons and in the male-specific CEMs (Supplementary

Figure 4C-G). Taken together, we discovered novel sexually dimorphic neurons exhibiting sex

differences in gene expression of various gene families, with significant prominence for the

neuropeptide family, suggesting that the peptidergic system is regulated in a highly sex-specific

manner.

Figure 2. Ranking of sex-shared neuronal clusters highlights novel sexually dimorphic neurons.
(A) The number of differentially expressed genes (DEGs) in the sex-shared neuronal clusters. Clusters are divided to sensory (triangle), interneuron (hexagon), and motor (circle) neurons. A gene was considered as DE according to the following criteria: average log fold change > 1, number of cells in a cluster ≥ 9 in both sexes, p-value < 0.05, and a threshold > 2 in at least one of the sexes (*See Methods*). (B) Gene ontology (GO) term analysis for the 2216 DEGs in both sexes and in hermaphrodite and males separately (only genes meeting the criteria of log fold change > 1, number of cells in a cluster ≥ 9 in both sexes, p-value < 0.05, threshold ≥ 2 in at least one of the sexes are included) reveals specific enrichment for synaptic and signaling gene expression. Wormcat p-values are determined by one-sided Fisher test with FDR correction. (C) Percentage of DEGs from each gene family as a fraction of the number of genes in the specific family. (D) The number of DEGs in the gene families presented in (C) divided according to DEGs that are enriched in males (green) and hermaphrodites (magenta). A DEG that is observed in more than one cluster

206 was counted accordingly (*See Methods*). For (D) we performed a two-sided Fisher's exact test
207 followed by p-value Bonferroni correction for multiple comparisons.
208

A variety of neuronal gene families are differentially expressed between the sexes

In addition to the neuropeptide gene family, other gene families could point towards sex

differences in neuronal function (Supplementary Figure 5-6). Synaptic-related genes and ion

channels included only a limited number of DEGs (Figure 2C, Supplementary Figure 5C-I). Several

transcription factors (TFs) are cell-specifically differentially expressed between the sexes

(Supplementary Figure 6). TFs known to drive sex-specific maturation, such as *tra-1* and *mab-3,*

were differentially expressed in only a few neurons (Supplementary Figure 6A-B) suggesting they

may be expressed only transiently at the transition between L4 and adulthood. The TF *tbx-32* was

differentially expressed in the largest number of clusters with a male bias, suggesting a previously

uncharacterized role for this gene in the development of the male nervous system

(Supplementary Figure 6A). DEGs also included TFs that are predicted to be bound by TRA-1

[15,36,37], supporting possible multiple unexplored roles for TFs in sex-specific neuronal regulation

(Supplementary Figure 6B).

Figure 3. Sexual dimorphism of neuropeptides across the nervous system. (A) Bubble plot of differentially expressed neuropeptide genes between the sexes. Bubble size corresponds to the average fold change in gene expression (only genes meeting the criteria of log fold change > 1, number of cells in a cluster ≥ 9, p-value < 0.05, threshold > 2 in at least one of the sexes are included), and bubble color indicates enrichment in either sex (green for male enrichment, magenta for hermaphrodite enrichment). Rows represent neuron types, grouped by sensory neurons, interneurons, and motor neurons. Columns represent individual genes. (B) UMAP projection of AUA, AVE, RMH, DVC, AVB, AIB, PDB, URA, M2, DVB, SAB, PLM, SIB and AVM clusters showing expression of *nlp-11* gene in both sexes. Heatmap represents the expression levels. Bubble plot representation of *nlp-11* gene expression in AUA, AVE, RMH, DVC, AVB, AIB, PDB, URA, M2, DVB, SAB, PLM, SIB and AVM clusters of the two sexes, ranked from lowest to highest p-value. Bubble size represents average log fold change, bubble color (green for males and magenta for hermaphrodites) represents enrichment in each sex. (C, E) Representative confocal micrographs showing the distinctive multi-colored neuronal nuclei in the NeuroPAL strain otIs669 in the head (C) and tail (E), a strain in which every neuron class can be uniquely identified based on its colour-based barcode[38,39]. These images were utilized to identify neurons expressing the *nlp-11(syb4759[nlp-11::SL2::GFP::H2B])* reporter in young adult hermaphrodites and males. Scale bars represent 10μm. (D) Quantification of *nlp-11* fluorescent intensity from (C) in head neurons AUA, AVE, AIB URAD, URAV, M2, SABV, SIB AVM. (F) Quantification of *nlp-11* fluorescent intensity from (E) in tail neurons DVC, PDB, PLML, PLMR, and DVB neurons in both sexes at YA stage. For each neuron group comparison, n represents the number of animals in each group in which neurons were identified and recorded. AUA, n = 14, 14 ; AVE, n = 14, 14; AIB, n = 12, 12: URAD, n = 14, 14 : URAV, n = 14, 14; M2, n = 12, 13; SABV, n = 13, 13; SIB, n = 9, 9; AVM, n = 13, 13; PDB, n = 14, 14; PLML, n = 14, 14; PLMR, n = 14, 14; DVB, n = 14, 12; DVC, n = 14, 13, for hermaphrodite and males, respectively. In the box-and-whiskers graph, the center line in the box denotes the median, while the box contains the 25th to 75th percentiles of the dataset, whiskers define the minimum and maximum value with dots showing all points. We performed a two-sided Mann-Whitney test for each comparison.

Neurotransmitter identities are mostly maintained in sex-shared neurons

Among the unique functional features that define neuron types are the neurotransmitters they

use[33,40–43]. Since our analysis revealed numerous sex-shared neurons with a sexually dimorphic

gene expression profile, we wondered whether these neurons maintained their neurotransmitter

identity across the sexes. For this purpose, we examined the expression of neurotransmitter-

related genes in hermaphrodite and male sex-shared neuronal clusters (same clusters shown

Figure 2A, Figure 4A). While the known AIM neurotransmitter switch is fully captured in our data

set (Figure 1F-G, Figure 4A), we did not observe any other similar cases in the other sex-shared

neurons (Figure 4A). However, when examining the differential expression of these

neurotransmitter-related genes, we found that some of them, mostly cholinergic-related and

glutamatergic-related genes, exhibited biased expression in several sex-shared clusters (Figure

4B), suggesting sexual dimorphism may be more prominent in neurotransmitter release rather

than identity.

With this in mind, we explored the relationship between dimorphic neurotransmitter release and

dimorphic synaptic connectivity. The complete synaptic wiring diagram of the nervous system is

available in *C. elegans* for both sexes[35], enabling a comparison of the number of synaptic

connections to neurotransmitter-related DEGs. We hypothesized that neurons that show higher

expression of neurotransmitter-related genes in one sex, might also have more outgoing

connections. To test this, we correlated the difference in connectivity between the sexes in

outgoing, incoming, and all synaptic connections (chemical connections)[35] (Figure 4C*, See*

Methods) with the average log-fold change of neurotransmitter-related genes presented in Figure

4B. Strikingly, Pearson correlation coefficient was significant only for outgoing connections but

not for incoming or all synaptic connections (Figure 4D). These results indicate that to some

extent, neurotransmitter release of a neuron is correlated with its outgoing connectivity.

Figure 4. Neurotransmitter identity is preserved across sex-shared neuronal clusters. (A) Left and right bubble plots representing the mean expression values of neuronal identity genes in hermaphrodites and males, respectively. Neuronal clusters and genes are color-coded according to their neurotransmitter identity. (B) Bubble plot representing differentially expressed neurotransmitter identity genes between the sexes. Only genes with significant log fold change are shown. Bubble size in (A) corresponds to gene expression and in (B) to the average log fold change in gene expression. Bubble color indicates enrichment in either sex (green, male; magenta, hermaphrodite). Rows represent neuron types, grouped color coding for neurotransmitter identity. Columns represent individual genes. (C) Bar plot representing the difference in outgoing connectivity between males and hermaphrodites in each cluster. Numbers are represented as an absolute value (*See Methods* for more details). (D) Two-tailed Pearson correlation analysis between the difference in connectivity (outgoing connections, all synaptic connections, and incoming connections) and the average log-fold change of neuronal identity genes (*See Methods* for more details). For (D) we performed a two-sided Pearson correlation test.

Sexually dimorphic gene expression and function of the PLM neuron underlie sex-specific

mechanosensory responses

Among the sex-shared neuronal clusters with a high number of DEGs, we observed all the touch receptor neurons (TRNs): ALM, AVM, PVM, and PLM (Figure 2A), required for mechanosensory responses to gentle touch[44]. We turned to explore this dimorphic molecular repertoire, focusing

specifically on PLM, a TRN that mediates the posterior gentle touch response[45] and is the most

sexually dimorphic among TRNs (Figure 2A, 5A)[44,46]. We validated the differential expression of

multiple neuronal-related genes (Figure 5B-C) using transcriptional and translational reporters

(Figure 5D-G), or single molecule Fluorescent In Situ Hybridization (smFISH) (Figure 5H). We

tested gentle touch responses of hermaphrodites and males. We found that hermaphrodites

showed reduced responses to posterior but not to anterior gentle touch (Figure 5I). As a control,

we tested harsh touch-tail responses in the same animals[7], which were not dimorphic (Figure 5I).

While characterizing the development and growth of PLM, we found sex differences in the length

of the posterior processes at the adult stage, with fewer males having posterior processes

(Supplementary Figure 7A-D). Analysis of PLM connectivity showed minor differences between

the sexes, with males making slightly more gap junctions[35] (Supplementary Figure 7E). Thus, we

speculated that the behavioral dimorphism we observed in posterior gentle touch could stem

from sex differences in PLM neuronal activity. Indeed, when measuring calcium traces of PLM

basal activity, we observed higher activity in males in frequencies up to 2 Hz (Figure 5J-M), further

supporting a sexually dimorphic role for PLM. In addition, optogenetic activation of the tail TRNs,

which includes PLM, resulted in male behavioral responses that were statistically significantly

distinct from hermaphrodites, supporting PLM dimorphism (Supplementary Figure 8). We did not

observe significant changes in the expression of major touch-response genes such as *mec-4* and

mec-10[47,48] (Supplementary Table 2). To test the involvement of specific DEGs in PLM function, we

knocked down the neuropeptide *nlp-11* (higher in PLMs of males) or the RNA-binding protein

mec-8 (higher in hermaphrodites) and assayed the RNAi-silenced animals for posterior gentle

touch responses in both sexes. While *nlp-11* did not show an effect on gentle touch responses in

either of the sexes, *mec-8* was required only in hermaphrodites (Figure 5N). Importantly, over-

expressing MEC-8, specifically in PLM, restored the behavioral responses of hermaphrodites

(Figure 5O). Altogether, our results establish PLM as a sexually dimorphic neuron with a

differential transcriptomic profile and identify a dimorphic gene expression critical for a sex-

specific mechanosensory response.

Figure 5: Sexually dimorphic properties of the touch receptor neuron PLM. (A) Illustration depicting the positioning of the PLM sensory neuron, as outlined in WormAtlas (https://www.wormatlas.org/) (B) Bubble plot representation of *nlp-11, flp-8, dop-3, sng-1,* and *mec-8* genes expression in PLM clusters of the two sexes. Bubble size represents average log fold change, bubble color (green: males and magenta: hermaphrodites) represents enrichment in either sex. Genes with average log fold change <1 but p-value <0.05 are included. (C) Table lists the established functions of *nlp-11, flp-8, dop-3, sng-1,* and *mec-8* genes. (D) Representative confocal micrographs showing expression of *nlp-11::SL2::GFP::H2B* in PLM neurons (white dotted circle) (left). Quantification of *nlp-11::SL2::GFP::H2B* (right). n = 16 hermaphrodites and 15 males. (E) Representative confocal micrographs showing expression of *flp-8p::GFP* in PLM neurons (white dotted circle) (left). Quantification of *flp-8::GFP* (right). n = 15 animals per group. (F) Representative confocal micrographs showing expression of *dop-3::RFP* in PLM neurons (white dotted circle) (left). Quantification of *dop-3::RFP* (right). n = 19 animals per group. (G) Representative confocal micrographs showing expression of *sng-1::GFP* in PLM neurons (white dotted circle) (left). Quantification of *sng-1::GFP* expression in PLM neurite measured along 100 μm from the soma (right). n = 10 animals per group. (H) Representative confocal micrographs showing expression of *mec-8* CAL Fluor Red 590nm smFISH probe in PLM neurons labeled with *mec-4::gfp* and DAPI (white dotted circle) (left). Quantification of *mec-8* smFISH expression (right). n = 27 hermaphrodites and 26 males. a.u., arbitrary units. Scale bars represent 10 μm. (I) Left: Illustration depicting the gentle touch and harsh touch in the anterior and posterior sides of *C. elegans.* Created with Biorender.com. Middle: posterior gentle touch responses of hermaphrodites and males and harsh tail-touch responses tested on the same animals (n = 20 animals per group). Right: anterior and posterior gentle touch responses of hermaphrodites and males (n=15 animals per group). (J) PLM normalized and color-coded GCaMP6s calcium responses of hermaphrodites and males (n = 13 recordings from 13 animals). Heatmaps represent the calcium levels of individual worms. (K) Mean and SEM traces of PLM calcium responses of hermaphrodites and males. (L) Mean traces showing the normalized amplitudes of PLM, zoomed in at frequencies 0-0.2 Hz. (M) Statistical comparisons of the AUC (area under the curve) measured from PLM amplitudes in the two sexes, for chosen frequencies (0-2 Hz) (*See Methods*). (N) Posterior gentle tail-touch responses of hermaphrodites (left) and males (right) after RNAi knock down of control (n=15 each group), *nlp-11* RNAi (n= 14 for hermaphrodites and n= 15 males), and *mec-8* RNAi (n= 15 for hermaphrodites and n= 14 males). We performed a Kruskal-Wallis test followed by a Dunn's multiple comparison test. (O) Posterior gentle tail-touch responses of hermaphrodites in wild-type (n= 17), *mec-8 (ok2043)* mutant (n=18), and *mec-8* rescue in PLM (n=18). We performed a Kruskal-Wallis test followed by a Dunn's multiple comparison test. In the box-and-whiskers graphs, the center line in the box denotes the median, while the box contains the 25th to 75th percentiles of the dataset, whiskers define the minimum and maximum value with dots showing all points. In (D, E, F, G, H, I, M) we performed a two-sided Mann-Whitney test for each comparison.

The cell adhesion molecule LAD-2 is required for proper neurite extension in male but not hermaphrodite PHC neurons

Our study offers molecular candidates to explain the mechanistic basis for previously recognized

sexual dimorphism of neurons such as PHC, AFD, AIM and AVG (Figure 2A). For example, we

validated two differentially expressed neuropeptides, *flp-7* and *flp-22*, with higher expression in

hermaphrodite AVG neurons, suggesting a potential role in modulating AVG's circuit, specifically

in hermaphrodites (Supplementary Figure 9)[7]. We next focused on PHC tail sensory neurons,

which undergo massive remodeling in males compared to hermaphrodites, with males exhibiting

neurite extensions that are twice as long[34]. This elongation is presumably required to

accommodate the increased synaptic connectivity with male-specific neurons[34,35] (Figure 6A,

Figure 4D). The TF DMD-3 was shown to modify the function, connectivity, and molecular features

of the male PHC neurons. We observed increased expression of *dmd-3*, *eat-4/VGLUT* and the

neuropeptide *flp-11* in the male PHC cluster[32,34] (Figure 6B). Our ranking analysis identified 209

DEGs for PHC (Figure 2A). Supporting the dimorphic connectivity and morphology of PHC, these

DEGs were enriched in the GO-terms "synaptic function" and "transmembrane transport" (Figure

6C). We validated a subset of neuronal DEGs predicted by our scRNA-seq using transcriptional

and translational reporters, including a neuropeptide gene, GABA and dopamine receptors, and

the cell adhesion molecule *lad-2* (Figure 6D-I). Since *lad*-2 has been shown to mediate axon

guidance[49], we imaged the neurite extension of PHC in both sexes in *lad-2* mutants. We revealed

that *lad-2* is required only for the male extension, consistent with its higher expression in males

(Figure 6I-K). Taken together, our data set confirms the dimorphic transcriptomic profile for

known dimorphic sex-shared neurons and allows the identification and exploration of novel

molecular mechanisms within these neurons.

Figure 6: Expression of novel DEGs in PHC. (A) Illustration depicting the location of PHC sensory neurons and their neurite length in both sexes, as outlined in WormAtlas. (B) Validation of previously reported DEGs. Left, PHC representation in the UMAP projection of all cells (inset marked as orange). UMAP projection depicting expression of *dmd-3*, *eat-4*, and *flp-11* gene within PHC cluster. Heatmap represents expression level. (C) GO term enrichments for DEGs in PHC

cluster along with the number of genes in each GO term. (D) Bubble plot representation of *capa-*
1, F32B4.5, dop-3, and *lad-2* differential genes expression in PHC cluster. Bubble size represents
average log fold change, and bubble color (green: males and magenta: hermaphrodites)
represents enrichment in each sex. (E) Table lists the established functions for *capa-1, F32B4.5,*
dop-3, and *lad-2* genes. (F) Representative confocal micrographs showing expression of *capa-*
1p::GFP in PHC neurons (white dotted circle) (left). Quantification of *capa-1p::GFP* (right).
Hermaphrodites, n = 14; males, n= 16. (G) Representative confocal micrographs of *F32B4.5p::GFP*
expression in PHC neurons (white dashed circle) (left). Quantification of *F32B4.5p::GFP* (right). n
= 10 animals per group. (H) Representative confocal micrographs of *dop-3p::RFP* expression in
PHC neurons (white dashed circle) (left). Quantification of *dop-3p::RFP* (right). n = 14 animals per
group. (I) Representative confocal micrographs of *lad-2p::GFP* expression in PHC neurons (white
dashed circle) (left). Quantification of *lad-2p::GFP* (right). n = 15 animals per group. (J-K)
Representative confocal micrographs of *eat-4p11::GFP* expression in PHC neurons in wild-type
and *lad-2 (tm3056)* mutant males, n = 16 animals per group (J) and hermaphrodites, n = 10
animals per group (K). Scale bars represent 10 μm. In the box-and-whiskers graph, the center line
in the box denotes the median, while the box contains the 25th to 75th percentiles of the dataset,
whiskers define the minimum and maximum value with dots showing all points. In (F, G, H, I, J, K)
We performed a two-sided Mann-Whitney test for each comparison.
412

Correlation analysis predicts novel regulators of synaptic connectivity

Can we predict which genes regulate synaptic connectivity based on the relationship between

their expression pattern and wiring maps? The relationship between gene expression and

synaptic connectivity is a compelling notion in neurobiology. By integrating transcriptomic profiles

with connectome data, we can generate predictions on the identity of genes that regulate

synaptic wiring. This approach has been previously used, but these studies focused only on

hermaphrodites and used binary gene expression data[50,51]. As the connectome of *C. elegans* is

available for both sexes, we further utilized our non-binary single-cell gene expression dataset of

both sexes and the connectome of *C. elegans* to answer this key question. For this purpose, we

calculated the correlation between the number of outgoing connections for each neuron based

on[35] and the average gene expression values of all genes in our dataset, across the 62 clusters

ranked in Figure 2A, analyzing hermaphrodites and males separately (*See Methods*, Figure 7A-

B)[35]. We chose outgoing connectivity because of the availability of pre-synaptic *in-vivo*

markers[52,53] for further validation of candidates. To identify genes that might regulate the

outward connectivity of neurons, we performed statistical analysis accounting for multiple

comparisons, which resulted in an adjusted p-value for each correlation value (*See Methods*). We

then searched for significantly correlated genes that overlap between hermaphrodite and male

correlation profiles (Figure 7C). We discovered 54 genes with significant correlation in both sexes,

with more than half still uncharacterized. Among the neuronal genes exhibiting a significant

positive correlation, are key presynaptic genes such as the active zone scaffold *syd-2* involved in

presynaptic assembly [54,55], but also postsynaptic genes such as *acc-2,* encoding acetylcholine-

gated chloride channel[56], and *gar-3* encoding G-protein-linked acetylcholine receptor[57] (Figure

7C). This finding highlights numerous uncharacterized genes with potential synaptic function and

suggests that some synaptic regulatory pathways are shared between the two sexes. To gain a

better perspective into dimorphic synaptic regulation, we looked specifically for neuronal genes

that are either significantly correlated with outgoing connectivity in hermaphrodites, in males or

in both sexes (Figure 7D). While exploring the expression of these genes in our data set, we

uncovered the sexually dimorphic expression patterns of some of them (Figure 7E). Importantly,

these genes included *ttx-7,* which encodes a myo-inositol monophosphatase (IMPase), already

shown to mediate proper synaptic localization in the interneuron RIA[58] and *zig-1* predicted to

encode a cell adhesion molecule whose function is unexplored[59,60]. We found that *ttx-7* is

significantly correlated in both sexes (Figure 7C). Based on *ttx-7* expression profile, we

hypothesized that it might regulate synaptic connectivity of RIA neurons in both sexes (since both

express it equally high), but its absence would have a greater effect on the synaptic connectivity

of hermaphrodites CEP neurons (since it is highly expressed in hermaphrodites) (Figure 7E, red

squares, and 7F). We indeed found that in *ttx-7* mutant animals, RIA pre-synapses were mostly

mislocalized in both sexes (Figure 7G), while the number of pre-synapses in CEP neurons was

reduced only in hermaphrodites (Figure 7I). *zig-1* mutant animals showed lower pre-synaptic

signal only in males, in concordance with the gene being significantly correlated only in males

(Figure 7D, Figure 7H). We conclude that combining our dataset together with connectivity maps

uncovers novel genes regulating synaptic wiring, both in non-dimorphic and dimorphic contexts.

Figure 7. Correlation analysis identifies novel genes that regulate synaptic connectivity. (A) Schematic of the correlation analysis. Rows represent the number of outgoing synaptic connections calculated according to[35] (*See Methods*) across neuronal clusters, and columns represent genes and their mean expression values across neuronal clusters. Pearson correlation is calculated between connectivity numbers and mean gene expression for each gene. This analysis was done separately for the hermaphrodite and male data sets. (B) an example of the correlation output. Scatter plots representing the mean expression of the significantly correlated gene *VM106R.1* across the number of outgoing connections in males (right) and hermaphrodites (left). Each dot in the scatter plot represents a neuronal cluster. (C) Venn diagram showing significantly correlated genes in hermaphrodites (magenta) and in males (green) and the number of neuronal genes within these two groups (brown) (top). Heatmap of the Pearson correlation coefficients of the significantly correlated and overlapping genes between hermaphrodites and males (bottom). (D) Heatmaps of the Pearson correlation coefficient of significantly correlated neuronal genes in hermaphrodites (top) and in males (bottom). Common genes in both data sets are marked with a gray square. (E) Bubble plot representation of differential gene expression across neuronal clusters of the neuronal genes from (D). Bubble size represents average log fold change, bubble color (green: males and magenta: hermaphrodites) represents enrichment in each sex. Only significant DEGs are shown. Red square refers to genes and cells analyzed in (G-I). (F) Scatter plots representing the mean expression of *zig-1* (right) and *ttx-7* (left) across the number of outgoing connections in hermaphrodites (magenta dots) and males (green dots). Each dot in the scatter plot represents a neuron. (G) Representative confocal micrographs of *glr-3p::SNB-1::venus* expression in RIA neurons in wild-type and *ttx-7(nj40)* mutant hermaphrodites and males (left). Percentage of animals showing mislocalization signal of SNB-1 in RIA (right). Wild-type hermaphrodites n = 15, Wild-type males n = 14, *ttx-7(nj40)* n = 15 per group. (H) Representative confocal micrographs of *glr-3p::SNB-1::venus* expression in RIA neurons in wild-type and *zig-1(ok784)* mutant hermaphrodites and males (left). Quantification of venus signal in RIA (right). a.u., arbitrary units. Wild-type hermaphrodites n = 15, Wild-type males n = 14, *zig-1(ok784)* n = 15 per group. (I) Representative confocal micrographs of *dat-1::novogfp::CLA-1* expression in CEP neurons in wild-type and *ttx-7(nj40)* mutant hermaphrodites and males (left). Quantification of CLA-1 puncta in CEP (right). n = 15 per group. In the box-and-whiskers graphs, the center line in the box denotes the median, while the box contains the 25th to 75th percentiles of the dataset, whiskers define the minimum and maximum value with dots showing all points. For (C) we performed Pearson correlation analysis with Benjamini Hochberg correction for multiple comparisons (*See Methods*). For (G) we performed a two-sided Fisher's exact for each comparison. For (H-I) we performed a two-sided Mann-Whitney test for each comparison.

DISCUSSION

In this work, we established a comprehensive transcriptome map of the sex-shared nervous

system in both sexes and uncovered novel sexually dimorphic neurons, including the posterior

light touch neurons PLM. In addition to sex-shared neurons, *C. elegans* males have an additional

93 male-specific neurons. Even though these can be subdivided into neuronal subtypes[42,61], for

most male specific-neurons molecular markers that define individual identities haven't been

described. Thus, male-specific clusters have been left out of this analysis, until further tools

become available[62]. Our ranking analysis reveals that the most prominent dimorphic gene family

is neuropeptides, with a male-biased expression, in accordance with our previously published

developmental comparison between the sexes[31]. Neuropeptides are short protein ligands that

modulate synaptic activity mostly through the binding to G protein-coupled receptors (GPCRs),

playing a critical role in regulating functions of the nervous system functions[63]. A

neuropeptidergic connectome has recently been constructed, revealing a dense network with

unique sex-shared neurons that serve as "peptidergic hubs"[64]. This suggests that a significant

rewiring of the neuropeptidergic connectome occurs in addition to the rewiring of the physical

connectome between sexes. Future studies exploring the entire neuropeptidergic connectome in

sexually dimorphic context would shed more light on how neuropeptide signaling has diverged to

drive nervous system function and behavior. [31,65].

Our functional analysis of sex-shared neurons uncovers a novel sexual dimorphism in

mechanosensory posterior gentle touch responses, with males displaying higher sensitivity

compared to hermaphrodites. This difference is likely mediated by the highly dimorphic touch

receptor neuron PLM, which exhibits significant differences in basal neuronal activity between

the sexes. While PLM activity decreases when courtship begins[66], the heightened

mechanosensory responses in males may be essential for mating, as was already shown for

turning behavior[68]. Hermaphrodites, on the other hand, do not display such behavior and show

lower sensitivity. In contrast, both sexes exhibit similar responses to harsh tail touch[7], suggesting

that hermaphrodites maintain robust responses to dangerous stimuli for survival. Our focus on

mechanosensation in this and previous work[7] provides insights into how different touch

modalities (gentle and harsh) can result in distinct outcomes, potentially reflecting different

evolutionary pressures.

We identified the RNA binding protein MEC-8 as a hermaphrodite-enriched gene in PLM and

shown it to be critical for their PLM mechanosensory responses, suggesting that different

molecular mechanisms mediate mechanosensation in the two sexes.

Our single-cell data set identified the cell adhesion molecule LAD-2 as a male-specific regulator

of PHC neurite extension[34], though which specific isoform is responsible remains unclear[69]. Future

screens are required to uncover genes governing PHC neurite structure in hermaphrodites.

We observed that neurotransmitter identity is largely preserved across sexes, with the notable

exception of AIM neurons. Slight changes in expression levels for neurotransmitter genes

between the sexes correlate with changes in outgoing connectivity, although this correlation was

relatively low (r=0.37), suggesting other factors might underlie the differences in outgoing

connectivity, such as synaptic-related genes, transcription factors, or a combination between

different genes. This led us to explore the correlation between gene expression across our data

set and connectivity. Based on this, we were able to predict a large number of candidate genes in

both sexes that might regulate outgoing connectivity, validating two of them. While this

correlation analysis was sufficient to identify gene-to-connectivity matches for specific neurons,

it could not be extended to the entire worm's nervous system, as our analysis did not capture all

sex-shared neurons (62 clusters that have passed all the filters in ranking analysis) and some of

the sex-shared neurons are under-represented in our data set. Further studies profiling the rest

of the sex-shared neurons will possibly allow for a more accurate prediction of the way gene

expression regulates connectivity. Even with a full transcriptomic representation of the nervous

system, predictions can be complicated due to interactions existing between genes and to

differences between connectivity and single-cell RNA sequencing data pooling (small sample size

in EM sections versus pooling large number of worms for single-cell).

Overall, our integrative approach, combining molecular, cellular, behavioral, and computational

analyses, provides significant insights into sexual dimorphisms within the sex-shared nervous

system. This work lays the ground for future studies aimed at uncovering novel molecular

programs that regulate these neuronal dimorphisms, with implications for understanding similar

processes in higher organisms and the development of gender-specific therapeutic strategies.

METHODS

***C. elegans* strains and maintenance**

All *C. elegans* strains used in this study were cultivated according to standard methods[70]. Wild-

type strains were Bristol, N2. *him-5(e1490)* or *him-8(e1489)* were treated as controls for strains

with these alleles in their background. Worms were grown on nematode growth media (NGM)

plates seeded with *E. coli* OP50/NA22 bacteria as a food source. Worms were raised at 20°C,

unless noted otherwise. The sex and age of the animals used in each experiment are noted in the

associated figures and legends. All transgenic strains used in this study are listed in the key

resources table.

Isolation of *C. elegans* pure male population

To generate pure male populations, we followed our previously described protocol[31]. Briefly, *C.*

elegans strain with the genotype *rab-3p(prom1)::2xNLS::TagRFP] IV; dpy-28(y1) III; him-8(e1489)*

IV was employed. This strain harbors a temperature-sensitive mutation (*y1*) in the dosage

compensation gene, *dpy-28* along with *him-8(e1489)* mutation and a pan-neuronal RFP. Worms

were cultured at 15°C on 15 cm NGM plates seeded with NA22 bacteria. Embryos were isolated

by hypochlorite treatment and were age-synchronised 14-16 hours on a foodless NGM plate at

25°C. L1 larvae were collected and were placed on a food NGM plates approximately 1.5 cm from

the food on NGM plates and were allowed to crawl towards food for 1-2 hours at RT. Using a

scalpel, the area where L1/embryos were initially placed was removed. L1 worms were collected

and washed with M9 buffer and were incubated at 20°C until adulthood. This resulted in male

population of ≥98% purity. For each experiment the percentage of males in the population was

recorded.

Worm single cell dissociation

Worms of the desired genotype were grown at 15°C on 15 cm NGM plates seeded with NA22

bacteria until the plates were populated. Gravid adult worms were collected in M9 buffer and

embryos were isolated by hypochlorite treatment. The obtained embryos were hatched for 14-

16 hours on a foodless NGM plate at a restrictive temperature of 25°C. The number of L1 larvae

were counted and were plated (40,000-50,000 animals per plate) on a fresh NA22 seeded

bacterial plates and were grown to young adult stage at 20°C. Young adult worms were washed

10 times at 1300 rpm for 2 minutes with M9 buffer to get rid of any adhering bacteria. The worms

were then washed with ddH$_2$O. The worm pellets were then subjected to single cell dissociation

using protocols described earlier with slight modifications[20,71–74]. For every 0.5 ml of worm pellet

1 ml of freshly thawed sterile SDS-DTT solution (200 mM DTT, 0.25% SDS, 20 mM HEPES, 3%

sucrose, pH 8.0, thawed freshly) was added and incubated at room temperature for 4 min with

gentle mixing every 1 minute. Immediately after SDS-DTT treatment, worms were pelleted at

3,000 rpm for 1 min and washed 5 times with 10 ml egg buffer (118 mM NaCl, 48 mM KCl, 2 mM

$CaCl_2$, 2 mM $MgCl_2$, 25 mM HEPES, pH 7.3). 0.5 ml aliquots of worm pellet were distributed to 1.6

mL microcentrifuge tubes. 1 mL freshly prepared 15 mg/mL Pronase-E was added to each tube

and flicked to mix and then rotated at 20°C for 5 minutes. Using a glass pasteur pipette, worm

suspensions were combined from multiple tubes and were transferred to a 7 mL dounce

homogenizer (pestle clearance 0.02–0.056 mm, Kimble Chase). Pronase E was added to a final

ratio of 4:1 relative to the worm pellet. If the worm pellet was smaller than 0.5 mL a 2mL glass

homogenizer was used. The animals were disintegrated to single cells using type "B" plunger for

4-5 minutes at room temperature and then continued on ice gently. We note that homogenizing

on ice increased cell survival. The status of dissociation was monitored under a dissecting

microscope at 5-minute intervals. Dissociation was stopped when most of the animals were

dissociated but a few whole or fragmented animals were still present. Using a glass pasteur

pipette the cell suspension was aliquoted into 1.6 mL microcentrifuge tubes that were centrifuged

at 5,000 x g for 5 minutes at 4 °C. Supernatant was removed and the cell pellet was washed 3

times with 1 mL ice-cold egg buffer per wash at 5,000 x g for 5 minutes at 4 °C. The cells pellet

was resuspended in 1 ml ice-cold egg buffer and was passed through 5 µm syringe filter gently.

Cell suspension was taken on ice to FACS sorting immediately.

FACS based sorting

Single cell suspension from the desired strain were subjected to Fluorescence Activated Cell

Sorting (FACS) on a BD FACSAria™ III Cell Sorter using a 70-micron nozzle. eFlour 450 dye (final

concentration of 1 μl/mL) was used to discriminate live and dead cells. The cells were gated for

side scatter (SSC)-Area vs forward scatter (FSC)-Area, then single cells were selected from doublet

using FSC-Height vs FSC-Area. Finally, RFP+ cells were gated and sorted under the "4-way Purity"

mask from this subpopulation. Wild-type *him-5(e1490)* without any RPF+ cells was used as

control for gating. Sorted cells were collected in 1.6 mL microcentrifuge tube containing 500 μl L-

15-20 (L-15 medium containing 20% fetal bovine serum). The suspension was centrifuged at 5,000

x g for 10 minutes at 4 °C, resuspended in 50-80 μl of L-15-20 and counted using a

hemocytometer.

RT-PCR

For carrying RT-PCR experiments a previously described protocol was used[31]. YA hermaphrodite

worms were dissociated into single cells and the RFP+ and RFP- cells were FACS sorted in 500 μL

Trizol, flash frozen in liquid nitrogen and stored at -80°C until further processing. Total RNA was

extracted using the TRIzol LS (Invitrogen) protocol until the isopropanol precipitation step then

was re-suspended in the extraction buffer from RNA isolation kit (PicoPure, Arcturus). The

manufacturer's recommendations were followed for further isolation. Using the SuperScript™ IV

first-strand synthesis system (Invitrogen, Catalog # 18091050) and random hexamers, the

extracted total RNA was then converted to cDNA. Fast SYBR™ Green Master Mix (Applied

Biosystems, Catalog # 4385612) was used to carry out qPCR as per manufacturer protocol. The

experiments were carried out in 3 biological replicates. *act-1, pmp-3* and *tba-1* genes served as

internal controls. Ct values were obtained, and relative quantification of the expression of the

target genes was performed utilizing 2−ΔΔCt method.

Cell imaging (imaging flow cytometer)

Cell imaging experiments were carried out using an ImageStreamX Flow Imager system to

evaluate the morphology and viability of cells after single cell dissociation. Single cells were

dissociated from a *C. elegans* strain expressing pan-neuronal TagRFP as described above and were

resuspended in PBS with eFlour 450 dye (Live/Dead) and DRAQ5 dye (DNA binding) to a final

concentration of 1 µl/mL. The positive samples containing all fluorochromes were used to set up

each channel and the laser intensities. Gating was conducted based on Area vs RFP intensity,

followed by the selection of RFP+ cells and by discriminating fluorophore using Area vs intensity.

Cells derived from Wild-type N2 without RFP+ expression served as a control for gating. Images

were acquired for 20,000 cells.

Single-cell RNA sequencing

Depending upon the cell concentration, the single-cell suspensions were diluted to reach the

optimal range as recommended by 10x genomics manufacturer's protocol (approx. 800-1000

cells/µl). Each lane in the 10x genomics chip was targeted for 8,000 or 10,000 cells. Libraries were

prepared with 10x Genomics v3 Chemistry according to the manufacturer's protocol. The libraries

were sequenced using the Illumina NovaSeq 6000 with 150 bp paired end reads, except for one

hermaphrodite sample which was sequenced using the Illumina Nextseq 500 with a 75 cycle high

output kit and paired end sequencing. Detailed experimental information is found in

Supplementary Table 3. Real-Time Analysis software (RTA, version 4.0.0; Illumina) was used for

base calling and analysis was completed using 10X Genomics Cell Ranger software (4.0.0).

Detailed experimental information is found in Supplementary Table 3.

Single-cell RNA-Seq Mapping

Sequenced reads from each experiment were mapped to the *C. elegans* reference transcriptome,

version WBcel235 release WS276, using the Cell Ranger pipeline version 4.0.0. We extended the

3' UTR of each gene by 500 base pairs for additional mapping of reads to the 3' extremity of the

gene body since the 3' end is poorly annotated in *C. elegans*[26].

Background correction and filtering procedure

For each experimental batch (6 hermaphrodite and 6 male samples), the empty droplets were

identified and background correction was performed using SoupX[75] as previously described[20].

Subsequently, a cell filtering step on each batch was performed followed by clustering cells in

UMAP space using Seurat using 50 PCs (Supplementary Figure 10A). Clusters were defined by

computing Shared Nearest Neighbor using only the first two UMAP dimensions and the original

Louvain algorithm with a resolution of 1, using FindNeighbors and FindCluster functions where

all other parameters were set to default. Clusters with more than 0.5% neuronal markers and less

than 1.5% non-neuronal markers were selected for further analysis (Supplementary Figure 10B)

using a list of 563 neuronal and 114 non-neuronal marker genes based on Supplementary Table

1. To ensure no loss of neurons through this procedure, a second round of selection was

performed on cells that did not pass the first filtering step. For the second round, subclustering

was performed on the remaining clusters using same parameters as before and the same filtering

of clusters using neuronal and non-neuronal markers was applied. Finally, cells containing more

than 20% of UMIs originating from mitochondrial genes were removed (Supplementary Figure

10D-E).

Dimensionality reduction, batch correction and cluster Identification

Monocle 3 was used to combine the male and female batches with the CenGEN[20] L4

hermaphrodite neuronal data. Counts were imported to Monocle 3 and batches were aligned

using Batchelor using 135 PCs and an alignment k of 5, followed by UMAP dimensional reduction

using a minimal distance of 0.3 and 75 neighbors, and clustering using Leiden algorithm with a

resolution of 3e-3, resulting in 72 partitions. This UMAP projection was kept to display the final

clustering and annotation analysis. Two iterative rounds of subclustering were performed after

the first one. For the second iteration 2 subclustering steps were performed on partition 1 and on

the merge of partitions 2 to 72. The same parameters were used for both subclusterings: 135 PCs

and an alignment k of 20, followed by UMAP dimensional reduction using a minimal distance of

0.1 and 15 neighbors, and clustering using Leiden algorithm with a resolution of 3e-4 resulting in

36 and 82 new partitions, coming from former partitions 1 and 2-72, respectively. This totaled

118 partitions that were inspected to assign identities or to continue with a third round of

sublcustering. The third iteration of subclustering was only applied to partitions where identity

was not well defined after two iterations. We used partition definition in Monocle 3 to define

clusters that were well segregated and with identities well defined based on the content of

CenGEN annotated cells. If these partitions contained mixed CenGEN identities in some cases we

opted for the cluster definition if the segregation of cells matched well with a single neuronal

identity. In other cases, a third round of dimensionality reduction and clustering was performed.

In total, 27 partitions from the second iteration underwent a third round of subclustering,

producing a total number of 156 cell partitions/clusters. Finally, the CenGEN annotated cells

included in the clusters were used to annotate the clusters.

Pairwise comparison of clusters

To compare the similarity of gene expression between our hermaphrodite dataset and the

hermaphrodite L4 dataset generated by CenGEN, we used the 5000 most variable genes in our

dataset to calculate the Spearman correlation of the cell cluster using their average gene

expression. The results were plotted in a clustered heat map where the complete-linkage

clustering method was calculated for rows and columns using euclidean distances.

Differentially expressed gene analysis

Several metrics were extracted for all genes and all cells, including expression fraction and

average expression (Supplementary Tables 6-7). Differentially expressed genes between

hermaphrodite and male neurons for each cluster was performed using Seurat's "FindMarkers"

function with default parameters on log-normalized dataset. The thresholds were set to include

testing only genes detected in a minimum fraction of cells in either of the two populations, with

a default value of 0.01, and limiting testing to genes that show, on average, at least a 0.1-fold

difference (log-scale) between the two groups of cells, which is also the default value of

FindMarkers function. Finally, this step provided natural log fold changes, percentage of cells

where gene is detected for each sex and p value statistics provided by Wilcoxon Rank Sum test

(Supplementary Table 2).

Thresholding of gene expression

In order to define gene expression across neuron types we used the same thresholding strategy

as Taylor, S. R. *et al.* [20]. First, we defined how many cells should a cluster have in order to poses

similar predictive capabilities. For this, we generated Receiver-Operator Characteristic (ROC)

curve of True Positive Rate (TPR) vs False Positive Rate (FPR) by comparing the single-cell data to

the ground truth. As the ground truth dataset, we used the expression patterns of 160 neuronal

genes determined with high confidence in adult hermaphrodite, as previously described[20,38,76–79].

For this purpose, we assumed expression when the fraction of cells expressing was bigger than 0

and we found that using clusters with 9 cells or more in both sexes (65 cell types) give similar

predictive power when compared to CenGEN dataset (Figure 1D). With this in mind, we used

these 65 cell types to define the thresholding of the whole dataset following the same steps as

described in Taylor, S. R. *et al.*[20]. We used the proportion of cells in which a gene was detected to

define thresholds where genes identified in at least 1% of the cells within each neuron cluster

were regarded as expressed across all types of neurons, whereas transcripts detected in no more

than 2% of the cells within each neuron cluster were regarded as non-neuronal. In the case of

other genes, we implemented the percentile thresholding on a gene-by-gene basis, determining

thresholds as a fraction of the highest cell proportion observed for each gene. Using the R package

boot[80–82], we produced 5,000 stratified bootstraps of the ground truth genes for each threshold

percentile. Subsequently, we determined the TPR, FPR, and False Discovery Rate (FDR) for the 57

clusters and for each distinct neuron type (Figure 1D-E). We used the same 4 thresholds of

increasing stringency and in these 57 clusters, we observed similar performance as reported

before for the whole L4 dataset (Figure 1E). Finally, this thresholding was used for other clusters

even if they contained 10 cells or less.

Ranking analysis

To rank the sex-shared neuronal clusters according to the number of differentially expressed

genes (DEGs) between the sexes, we counted the number of DEGs in each cluster using Python

3.12. Clusters with mixed or unknown identities were removed from this analysis because of

inability to calculate connectivity numbers for these clusters for the analysis below. Clusters were

included only if there were more than 9 cells in a cluster for both sexes. This yielded a list of 62

sex-shared clusters we continued analyzing (Supplementary Table 4). A gene was considered as

differentially expressed if the following criteria were met: average log fold change > 1, p-value <

0.05, threshold ≥ 2 in at least one sex (in order not to exclude genes with no expression in one

sex and high expression in the other). Then, the number of DEGs received after filtering (2216)

was used to generate gene ontology analysis, calculate the fraction of DEGs per gene family and

the number of sex-biased DEGs according to gene family (Figure 2B-D) and bubble plots

representing differential expression of specific gene families (Figure 3A, Supplementary Figure 5,

Supplementary Figure 6) were generated using GraphPad prism 10.

Sex-bias of differentially expressed genes across gene families

Using Python 3.12, we calculated the number of DEGs in males and in herms separately according

to gene families (see gene families in Figure 2D) for all the DEGs between the sexes that passed

the filter (Figure 2A). If a gene appeared in multiple clusters as a DEG, it was counted according

to the number of clusters it appeared. For example, *flp-7* is DE in the clusters AVG, RMF and URX,

so it was counted three times. For statistical analysis, we performed Fisher's exact test for each

gene family (subtracting the number of each count in each sex from the total count of genes in

all families included in the analysis) and then corrected with Bonferroni by multiplying each p-

value by the number of comparisons.

Neurotransmitter identity analysis

To analyze the expression of neurotransmitter genes in our data set, we used Python 3.12,

generating expression bubble plots for the two sexes separately (Figure 4A-B, using

Supplementary Tables 6-7, mean expression values of hermaphrodites and males respectively

across sex-shared clusters) and for dimorphic expression of neuronal identity genes (Figure 4C).

The clusters used for this analysis are listed in Supplementary Table 4. Then, to calculate the

connectivity of the sex-shared neuronal clusters we used Supplementary information 3 from

Cook, SJ. *et al.*[35]. This calculation included only connectivity observed by the electron microscopy

(EM) reconstructions[35]. For each cluster, we calculated using Python 3.12 for each sex the number

of outgoing connections (when the neuron is pre-synaptic), the number of incoming connections

(when the neuron is post-synaptic) and the sum of these two calculations (all synaptic

connections, chemical connections) (Supplementary Table 5). For our analysis we combined left

and right connectivity numbers for each neuron. Then, we calculated the difference between the

number of male connections and the number of hermaphrodite connections in each cluster, for

each synaptic category. These numbers are represented in Figure 4C as an absolute number. To

correlate the difference in connectivity with average log fold change of neurotransmitter identity

genes (Supplementary Table 2), we used two-tailed Pearson correlation analysis in GraphPad

Prism 10. We did this analysis for every synapse category (outgoing, incoming and all). Clusters

with two neurotransmitter identity genes showing dimorphic expression were included twice in

the correlation analysis. SIA cluster was discarded because the two neurotransmitter identity

genes were showing different sex biases (*cha-1* higher in hermaphrodites and *unc-17* higher in

males). AIM was also discarded because its neurotransmitter identity is known to be dimorphic,

being glutamatergic in hermaphrodites and cholinergic in males in the adult stage[33].

Microscopy

Confocal images were acquired using Zeiss LSM 880 confocal microscope using 20x, 40x or 63x oil

immersion objectives. Worms were anaesthetized in sodium azide (NaN3, final concentration 200

mM, prepared in M9 buffer) and mounted on a 5 % agarose pad (freshly prepared) on a glass

slide. The images were processed using ImageJ/Fiji software version 2.3.0/1.53q[83] or Zen

software. All images of the same group were imaged using identical microscope settings. Figures

were prepared using Adobe Illustrator v24.2.1. For imaging NeuroPAL worms, channels were

pseudo-colored in accordance with ref[38,39]. For single neuron expression quantification, ROIs were

chosen manually at the z-plane with the strongest signal using the respective fluorescent marker.

For *sng-1::GFP* analysis we measured the fluorescence intensity of the signal in PLM neurite

(observed by *mec-4p::*BFP) measured along 100 μm from the soma. For PLM development

analysis we measured the length of the anterior and posterior neurites using *mec-4p::BFP*

labeling. For Figure 6J-K we measured the length of PHC neurite from the anus which is visible

through *eat-4p11::GFP*. For Supplementary Figure 9 we identified AVG based on its anatomic

position and shape. For Figure 7G we calculated the percentage of animals with mislocalized

signal of *SNB-1::Venus* and for Figure 7H we measured *SNB::Venus* signal by manually selecting

ROI and normalizing the mean intensity value received to its area for each animal. For Figure 7I

we counted the number of CLA-1 puncta that surround CEP neurons.

Neuron identification using NeuroPAL

For neuronal identification the relevant CRISPR GFP reporters were crossed with NeuroPAL

landmark strain OH15262 harboring *otIs669* transgene. For imaging NeuroPAL worms, the

protocol followed are in accordance with ref[38,39]. For CRISPR GFP neuronal identity, colocalization

with the NeuroPAL landmark strain OH15262 harboring *otIs669* transgene was used to determine

the identity of all neuronal expression. Using ImageJ version 1.52p, the z-plane with the strongest

signal was used to measure the fluorescence intensity of individual neurons that were identified

by reporter/stain expression.

Gene ontology analysis

The WormCat gene set enrichment analysis tool was used for functional annotation of gene

ontology terms that are over-represented in differentially expressed genes combined or for each

sex[84].

Molecular cloning

To generate *mec-4p::BFP* and *eat-4p11::BFP* for PLM and PHC visualization we used PCR fusion[85].

mec-4p was amplified from *mec-4p::GCaMP3::tagRFP* kindly provided by Alon Zaslaver, while *eat-*

4p11 was amplified from an eat-4p11 containing plasmid. *BFP* fragment was amplified from a

plasmid containing *flp-6::tagBFP* kindly provided by Hannes Bulow. To generate PLM specific

rescue of MEC-8, *F10A3.11* promoter (460 bp upstream to *F10A3.11* sequence) amplified from

genomic DNA was fused to *mec-8* cDNA amplified from a N2 mixed-stage cDNA library and to

SL2::GFP amplified from pMO46[17] using PCR fusion. *F10A3.11* is predicted by the CenGEN and

our dataset to be specific to PLM in hermaphrodites, and in males it is predicted to be expressed

in PLM and RIV. Our observations show that its expression is specific in the tail, but it is expressed

in 2-3 additional neurons in the head. *capa-1p* (2154 bp upstream to *capa-1* sequence) and

*F32B3.5*p (2805 bp upstream to *F32B4.5* sequence) were amplified from genomic DNA and fused

to *GFP::H2B* amplified from genomic DNA of *ins-39::GFP::H2B* knock-in animals[31] using PCR

fusion. For *dat-1p::novoGFP::cla-1* generation we amplified *dat-1* promoter from a *dat-1p::IRE-1*

plasmid[86] and *novoGFP::cla-1* from a plasmid kindly provided by Ithai Rabinowitch and Peri

Kurshan.

Touch assays

Gentle touch assays were based on[87]. In brief, L4 animals were isolated the day before the

experiment and stored at 20°C overnight. On the day of the experiment, single 1-day adult

animals were transferred into NGM plates freshly seeded with 30 µl OP50. After 30 minutes of

habituation, animals were tested by applying a touch to the posterior part of the animal

(anteriorly to the anus) with an eyelash. Touch was applied when worms were not moving or

moved very slowly. Each worm was tested five times with intervals of at least 10 seconds between

each trial. The tested worm was given a score of one if it moved forward in response to the touch,

and zero if it did not move forward. The response index was then calculated as the average of the

forward responses. For anterior gentle touch responses, the anterior part of the animal was

touched with an eyelash and a reverse movement response was scored. For harsh tail touch

assays, animals were tested by applying a touch to the tail with a flattened platinum wire pick as

previously described[88] and their forward movements were scored similarly to animals subjected

to posterior gentle touch.

Single Molecule Fluorescent in Situ Hybridization

mec-8 CAL Fluor Red 590 nm custom probe was designed using Stellaris designer

(https://www.biosearchtech.com/stellaris-designer). All the reagents were ordered from

Biosearch Technologies and most of the preparations are based on their protocol (Stellaris RNA

FISH – protocol for *C. elegans*), with minor adjustments. Synchronized L4 worm population

expressing *mec-4p::GFP (zdIs5)* was washed three times with M9 to remove bacterial residues.

Then, after aspirating the M9, 1 mL of fixation buffer (4% formaldehyde in 1XPBS) was added to

the worms followed by 45 minutes incubation at room temperature. After fixation, worms were

washed twice with 1XPBS and supplemented with 70% ethanol. Worms were stored at 4°C

overnight. The next day, larvae pellet was centrifuged, and the ethanol was discarded. Worms

were then added with 1 mL Wash Buffer A (2 mL Stellaris RNA FISH Wash Buffer A (Biosearch

Technologies Cat# SMF-WA1-60), 7 mL Nuclease-free water, 1 mL Deionized Formamide) and

incubated at room temperature for 5 minutes. After discarding Wash Buffer A (spin down), worms

were dispensed with 100 µL of the Hybridization Buffer containing probe (2 µL of probe set (12.5

µM in TE buffer)) into a mixture of 100 µL Hybridization Buffer (450 µL Stellaris RNA FISH

Hybridization Buffer, Biosearch Technologies Cat# SMF-HB1-10 and 50 µL Deionized Formamide)

to achieve 250 nM probe solution and incubated at 37°C in the dark overnight. In the morning,

worms were washed once with Wash Buffer A and incubated with Wash Buffer A at 37°C in the

dark for 30 minutes. Then, Wash Buffer A was discarded, and worms were added 1 mL DAPI (4',6-

diamidino-2-phenylindole, Wash Buffer A consisting of 5 ng/mL DAPI) to stain the nuclei and

incubated at 37°C in the dark for 30 minutes. Then, DAPI was discarded, and worms were added

with 1 mL of Wash Buffer B (Biosearch Technologies Cat# SMF-WB1-20) and incubated for 5

minutes at room temperature. After discarding Wash Buffer B, 30 µL Vectashield® Mounting

Medium (Vector Laboratories Cat #H-1000) were added to the worms for imaging. Imaging was

done using Zeiss LSM 880 confocal microscope with 63x oil immersion objective. To image large

sample size, we stored samples at 4°C for a few weeks with no signal loss. To differentiate between

hermaphrodites and male at all larval stages, we used *ins-39*::SL2::GFP::H2B CRISPR knock-in

worms as described previously[31]. For quantification, we followed a protocol based on[89]. We

performed Gauss blur filter of sigma = 1 for every image. Then, we performed for each image

"Max Entropy" thresholding in Fiji that was selected based on trial-and-error process done on 2-

3 representative images. Then, we carried particle analysis for the image in the ROI determined

by the GFP channel. This gives a count for the number of particles in a specific ROI. We also

performed manual counting of smFISH puncta and got the same results.

Calcium imaging

To image spontaneous calcium traces in PLM, we used animals expressing *mec-4p::GCaMP3.35*.

Using a drop of S-basal with 10 mM levamisole that was filtered with 0.22 um filter, adult worms

were inserted in a dual olfactory chip that was a kind gift from Michael Krieg lab, fabricated as

previously described[90,91]. The flow rate in the chip was 0.005 ml/min. Imaging was done with a

Zeiss LSM 880 confocal microscope using a 40x magnification water objective. When the worm

was properly located inside the chip, PLM was imaged for 1150 frames (2.8 minutes), with

imaging rate of 6.667 Hz. For analysis, the GCaMP fluorescence intensity was measured using FIJI.

ROIs of PLM soma were marked manually to extract mean intensity values. Data analysis was

performed using Python 3.12. For each worm, the baseline fluorescent level (F0) was calculated

by averaging the mean values of 100 frames in the beginning of each recording. Then, for each

frame, ΔF was calculated by subtracting F0 from the value of that time point, and the result was

divided by F0, to normalize the differences in the fluorescence baseline levels between individuals

(ΔF/F0). To calculate the differences between the sexes in PLM basal activity, we performed

Fourier transform on the ΔF/F0 values, then the amplitudes were normalized for each worm to

its own maximum value. Finally, AUC (area under the curve) was measured using the trapz

function in Python, using frequencies up to 2 Hz.

RNAi-Induced Gene Silencing

Gene silencing using RNA interference was carried out as described previously using the feeding

protocol on agar plates[92]. Hermaphrodites at L4 stage were fed on HT115 bacteria expressing

dsRNA (source Ahringer library) for the targeted genes or on control bacteria harboring empty

RNAi vector, L4440. Posterior (gentle) touch responses at 1-day adult from their progeny were

scored. The RNAi efficiency was measured using *pos-1* RNAi as a positive control. For every RNAi

experiment, a new *pos-1* control was prepared.

Optogenetic stimulation assay

To measure the response of mechanosensory stimulus to the head and tail of hermaphrodite and

male *C. elegans* worms, we used AML470[93] strain that expresses light-gated ion channel Chrimson

and a fluorescent reporter mCherry in all the six gentle touch neurons under the control of *mec-4*

promoter. L4 males and hermaphrodites were isolated a day prior to the day of experiments. Males

were separated from hermaphrodites to make sure that they are not mated. The worms were

transferred to freshly seeded plates consisting of 1 ml of 0.5 mM all-trans-retinal (Sigma-Aldrich Cat.

R2500) mixed with *E. Coli* (OP50) as food source and stored in the dark at 20°C until the next day.

Day 1 adult male and hermaphrodite worms were used in the optogenetic behavior assay. On the

day of experiment 20-30 worms were transferred to the unseeded agarose plates to carry out

behavior assay while delivering optogenetic stimulation. To measure the response of

mechanosensory stimulus to head and tail of male and hermaphrodites, we delivered optogenetic

stimulus to the worms using a high throughput optogenetic delivery system[93,94]. Optogenetic

stimulation protocol was same as described in the previous work[93]. Briefly, a 400-μm diameter red

light stimulation was delivered to either the head, tail, or both simultaneously to each tracked worm

on the agarose plate. The illumination spot was centered on the tip of the head or tail of the worm.

Optogenetic stimulus of 630 nm wavelength was delivered through a computer-controlled projector

for 1 second with an inter trigger stimulus interval of 30 seconds. The stimulus intensity was randomly

drawn independently for the head and the tail from the set of 0 and 80 $\mu W/mm^2$ intensities. Thus,

there were four stimulus combinations: 1) Head: 0 $\mu W/mm^2$, Tail 0 $\mu W/mm^2$ 2) Head: 0 $\mu W/mm^2$,

Tail 80 $\mu W/mm^2$, 3) Head: 80 $\mu W/mm^2$, Tail 0 $\mu W/mm^2$, and 4) Head: 80 $\mu W/mm^2$, Tail 80 $\mu W/mm^2$.

Each behavior assay lasted for 30 minutes. The analysis of camera frames of the behavior assay was

same as described in the previous studies[93,94]. Briefly, two independent sets of behavior mapping

algorithms were employed, one for real-time tracking of worms and optogenetic stimulus delivery,

and the other for post processing analysis. The real-time algorithm in LabVIEW tracked all the worms

on the agarose plate and calculated various behavior parameters of each worm such as velocity, body

curvature, centerline, etc. The first and last index of the centerline were assigned as head and tail

respectively. Every 30 seconds, the real-time algorithm signals the computer-controlled projector to

deliver optogenetic stimulus to the head, tail, or both simultaneously for every tracked worm with

one of the four stimulus intensity combinations. In the post-processing algorithm, we correct for any

error in the assignment of head and tail of the worm. This algorithm also classifies behavior of the

worm by classifying pose dynamics into a behavior map of forward, reverse, and turns[94,95]. Same

analysis steps were used as in the previous study to determine the change in velocity[93]. The velocity

of the worm two seconds prior to stimulus onset was defined as the initial velocity of the worm. The

velocity of the worm at the end of one second stimulus was defined as the final velocity. Change in

velocity for each stimulus event for a worm was defined as the difference between the final velocity

and the initial velocity.

Correlation analysis

The entire analysis was done using Python 3.12. Using the number of outgoing connections

calculated for Figure 4 (See Supplementary Table 5), we generated for each sex a matrix including

the number of outgoing connections and mean expression values of all genes taken from

Supplementary Tables 6-7 across the 62 sex-shared neuronal clusters (Supplementary Table 4).

Genes that were not expressed in any of the 62 clusters were removed from the analysis. Then, we

performed Pearson correlation calculation with Benjamini-Hochberg False Discovery Rate (FDR)

correction for each comparison to account for multiple comparisons, providing correlation coefficient

for each gene in the data set of hermaphrodites and male separately. We refer in the text to

significantly correlated genes to genes with p-valued adjusted < 0.05.

Quantification and statistical analysis

Each figure legend and relevant method section include information regarding quantification,

statistical testing, and sample size.

Data availability

All data generated or analyzed in this manuscript are provided in the supplementary information

files. The Supplementary Table 2,6,7 contains the RNA-seq datasets related to this article. The

sequence data of this article have been submitted to NCBI and are available under BioProject

XXXX.

ACKNOWLEDGEMENTS

We thank members of the Oren-Suissa lab and Laurent lab for their critical insights regarding the

manuscript. We thank Dr. Joseph Dicken and Dr. Efrat Hagai at Flow Cytometry Unit, Life Science

Core Facility, Weizmann Institute of Science for helping with FACS sorting. scRNA-seq library

preparation was done with critical advice from Dr. Hadas Keren-Shaul, Dr. Merav Kedmi and Dr.

David Pilzer from the Genomics Sandbox unit at the Life Science Core Facility of Weizmann

Institute of Science. We thank Sapir Sela from the Oren-Suissa lab for cloning the *dat-*

1p::novogfp::cla-1 plasmid. We are grateful to Oliver Hobert lab for sharing strains. Some strains

used in this study were obtained from *Caenorhabditis* Genetics Center (CGC), which is funded by

the NIH Office of Research Infrastructure Programs (P40 OD010440). We thank WormAtlas which

is supported by NIH OD 010943. We are thankful to Alon Zaslaver, Ithai Rabinowitch, Peri Kurshan,

Hannes Buelow labs for providing plasmids. We thank Michael Krieg lab for providing the dual-

olfactory chips. We thank WormBase, an online biological database for *C. elegans*, which is

supported by Grant U41 HG002223 from the National Human Genome Research Institute at the

NIH, the UK Medical Research Council, and the UK Biotechnology and Biological Sciences

Research Council. MOS acknowledges financial support from the European Research Council ERC-

2019-STG 850784, Israel Science Foundation grant 961/21, Dr. Barry Sherman Institute for

Medicinal Chemistry, Sagol Weizmann-MIT Bridge Program and the Azrieli Foundation. MOS is

the incumbent of the Jenna and Julia Birnbach Family Career Development Chair. MOS and PL are

supported by HORIZON TMA MSCA Doctoral Networks grant # 101119745.

AUTHOR CONTRIBUTIONS

R.H. conducted and generated scRNA-seq data and carried out NeuroPAL imaging and analysis.

R.L. and P.L. performed the scRNA-seq data analysis; assigned neuron identities; analyzed

expression. G.S. helped with scRNA-seq initial datasets generation. RR helped with thresholding.

H.S. performed the ranking analysis, neurotransmitter identity with correlation analysis, the

functional analysis of PLM, AVG and PHC neurons including imaging gene expression, calcium

imaging and behavioral experiments and the correlation analysis and imaging presented in Figure

7. G.G. and E.S. contributed to the conceptualization and performance of correlation analysis. Y.S.

generated and conducted *capa-1* reporter expression. S.N.H. contributed to the *lad-2* mutant

experiments. S.K. performed and analyzed the optogenetic experiments A.L. supervised the

optogenetic experiments. M.O.S. supervised and designed the experiments. R.H., H.S., R.L., P.L.

and M.O.S. wrote the paper.

DECLARATION OF INTERESTS

The authors declare no competing interests.

RESOURCE AVAILABILITY

Lead Contact

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the Lead Contact, Meital Oren-Suissa (meital.oren@weizmann.ac.il).

Material Availability

Unique strains generated in this study have been deposited at the *Caenorhabditis* Genetics

Center. Requests for other strains and plasmids should be directed to the Lead Contact.

REFERENCES

1.   Gater, R. *et al.* Sex differences in the prevalence and detection of depressive and anxiety
disorders in general health care settings: report from the World Health Organization
Collaborative Study on Psychological Problems in General Health Care. *Arch Gen*
Psychiatry **55**, 405–413 (1998).

2.   Cerri, S., Mus, L. & Blandini, F. Parkinson's Disease in Women and Men: What's the
Difference? *J Parkinsons Dis* **9**, 501–515 (2019).

3.   Aleman, A., Kahn, R. S. & Selten, J. P. Sex differences in the risk of schizophrenia: evidence
from meta-analysis. *Arch Gen Psychiatry* **60**, 565–571 (2003).

4.   Fang, S. *et al.* Sexually dimorphic control of affective state processing and empathic
behaviors. *Neuron* **112**, 1498-1517.e8 (2024).

5.   Chiu, H. *et al.* Erratum: A circuit logic for sexually shared and dimorphic aggressive
behaviors in Drosophila (Cell (2021) 184(2) (507–520.e16), (S0092867420316202),
(10.1016/j.cell.2020.11.048)). *Cell* **184**, 847 (2021).

6.   Pechuk, V. *et al.* Reprogramming the topology of the nociceptive circuit in C. elegans
reshapes sexual behavior. *Current Biology* **32**, 4372-4385.e7 (2022).

7.   Setty, H. *et al.* Sexually dimorphic architecture and function of a mechanosensory circuit
in C. elegans. *Nat Commun* **13**, (2022).

8.   Moffitt, J. R. *et al.* Molecular, spatial, and functional single-cell profiling of the
hypothalamic preoptic region. *Science* **362**, (2018).

9.   Chen, P. B. *et al.* Sexually Dimorphic Control of Parenting Behavior by the Medial
Amygdala. *Cell* **176**, 1206-1221.e18 (2019).

10.   Welch, J. D. *et al.* Single-Cell Multi-omic Integration Compares and Contrasts Features of
Brain Cell Identity. *Cell* **177**, 1873-1887.e17 (2019).

11.   Kim, D. W. *et al.* Multimodal Analysis of Cell Types in a Hypothalamic Node Controlling
Social Behavior. *Cell* **179**, 713-728.e17 (2019).

12.   Gegenhuber, B., Wu, M. V., Bronstein, R. & Tollkuhn, J. Gene regulation by gonadal
hormone receptors underlies brain sex differences. *Nature 2022 606:7912* **606**, 153–159
(2022).

13.   Knoedler, J. R. *et al.* A functional cellular framework for sex and estrous cycle-dependent
gene expression and behavior. *Cell* **185**, 654-671.e22 (2022).

14.   Li, F. *et al.* Sex differences orchestrated by androgens at single-cell resolution. *Nature*
2024 629:8010 **629**, 193–200 (2024).

15.   Zarkower, D. Somatic sex determination. *WormBook* 1–12 (2006)
doi:10.1895/WORMBOOK.1.84.1.

16.   Barr, M. M., García, L. R. & Portman, D. S. Sexual Dimorphism and Sex Differences in
Caenorhabditis elegans Neuronal Development and Behavior. *Genetics* **208**, 909–935
(2018).

17.   Oren-Suissa, M., Bayer, E. A. & Hobert, O. Sex-specific pruning of neuronal synapses in
Caenorhabditis elegans. *Nature* **533**, 206–211 (2016).

18.   Fagan, K. A. *et al.* A Single-Neuron Chemosensory Switch Determines the Valence of a
Sexually Dimorphic Sensory Behavior. *Current Biology* **28**, 902-914.e5 (2018).

19.   Lawson, H. N., Wexler, L. R., Wnuk, H. K. & Portman, D. S. Dynamic, Non-binary
Specification of Sexual State in the C. elegans Nervous System. *Curr Biol* **30**, 3617-3623.e3
(2020).

20.   Taylor, S. R. *et al.* Molecular topography of an entire nervous system. *Cell* **184**, 4329-
4347.e23 (2021).

21.   Roux, A. E. *et al.* Individual cell types in C. elegans age differently and activate distinct
cell-protective responses. *Cell Rep* **42**, 112902 (2023).

22.   Ghaddar, A. *et al.* Whole-body gene expression atlas of an adult metazoan. *Sci Adv* **9**,
eadg0506 (2023).
23.   Ma, F. & Zheng, C. Transcriptome age of individual cell types in Caenorhabditis elegans.
Proc Natl Acad Sci U S A **120**, e2216351120 (2023).
24.   Lorenzo, R., Onizuka, M., Defrance, M. & Laurent, P. Combining single-cell RNA-
sequencing with a molecular atlas unveils new markers for Caenorhabditis elegans
neuron classes. *Nucleic Acids Res* **48**, 7119–7134 (2020).
25.   Truong, L. *et al.* Single-nucleus resolution mapping of the adult C. elegans and its
application to elucidate inter- and trans-generational response to alcohol. *Cell Rep* **42**,
(2023).
26.   Packer, J. S. *et al.* A lineage-resolved molecular atlas of C. elegans embryogenesis at
single-cell resolution. *Science* **365**, (2019).
27.   Cao, J. *et al.* Comprehensive single-cell transcriptional profiling of a multicellular
organism. *Science* **357**, 661–667 (2017).
28.   Hashimshony, T., Wagner, F., Sher, N. & Yanai, I. CEL-Seq: Single-Cell RNA-Seq by
Multiplexed Linear Amplification. *Cell Rep* **2**, 666–673 (2012).
29.   Schumacher, B., Gallrein, C. & Meyer, D. Neuron-type specific aging-rate reveals age
decelerating interventions preventing neurodegeneration. (2024) doi:10.21203/RS.3.RS-
4360587/V1.
30.   Purice, M. D. *et al.* Molecular heterogeneity of C. elegans glia across sexes. *bioRxiv* (2023)
doi:10.1101/2023.03.21.533668.
31.   Haque, R. *et al.* Sex-specific developmental gene expression atlas unveils dimorphic gene
networks in C. elegans. *Nature Communications 2024 15:1* **15**, 1–18 (2024).
32.   Pereira, L. *et al.* Timing mechanism of sexually dimorphic nervous system differentiation.
Elife **8**, (2019).
33.   Pereira, L. *et al.* A cellular and regulatory map of the cholinergic nervous system of C.
elegans. *Elife* **4**, (2015).
34.   Serrano-Saiz, E., Oren-Suissa, M., Bayer, E. A. & Hobert, O. Sexually Dimorphic
Differentiation of a C. elegans Hub Neuron Is Cell Autonomously Controlled by a
Conserved Transcription Factor. *Current Biology* **27**, 199–209 (2017).
35.   Cook, S. J. *et al.* Whole-animal connectomes of both Caenorhabditis elegans sexes.
Nature **571**, 63–71 (2019).
36.   Hunter, C. P. & Wood, W. B. The tra-1 gene determines sexual phenotype cell-
autonomously in C. elegans. *Cell* **63**, 1193–1204 (1990).
37.   Berkseth, M., Ikegami, K., Arur, S., Lieb, J. D. & Zarkower, D. TRA-1 ChIP-seq reèeals
regulators of sexual differentiation and multileèel feedback in nematode sex
determination. *Proc Natl Acad Sci U S A* **110**, 16033–16038 (2013).
38.   Yemini, E. *et al.* NeuroPAL: A Multicolor Atlas for Whole-Brain Neuronal Identification in
C. elegans. *Cell* **184**, 272-288.e11 (2021).
39.   Tekieli, T. *et al.* Visualizing the organization and differentiation of the male-specific
nervous system of C. elegans. *Development* **148**, (2021).
40.   McIntire, S. L., Jorgensen, E. & Horvitz, H. R. Genes required for GABA function in
Caenorhabditis elegans. *Nature 1993 364:6435* **364**, 334–337 (1993).

41. Sulston, J., Dew, M. & Brenner, S. Dopaminergic neurons in the nematode Caenorhabditis
elegans. *Journal of Comparative Neurology* **163**, 215–226 (1975).

42. Wang, C. *et al.* A neurotransmitter atlas of the nervous system of C. elegans males and
hermaphrodites. *Elife* **13**, (2024).

43. Hobert, O. Terminal Selectors of Neuronal Identity. *Curr Top Dev Biol* **116**, 455–475
(2016).

44. Chalfie, M. & Sulston, J. Developmental genetics of the mechanosensory neurons of
Caenorhabditis elegans. *Dev Biol* **82**, 358–370 (1981).

45. Chalfie, M. & Sulston, J. Developmental genetics of the mechanosensory neurons of
Caenorhabditis elegans. *Dev Biol* **82**, 358–370 (1981).

46. Chen, X. & Chalfie, M. Regulation of Mechanosensation in C. elegans through
Ubiquitination of the MEC-4 Mechanotransduction Channel. *The Journal of Neuroscience*
35, 2200 (2015).

47. O'Hagan, R., Chalfie, M. & Goodman, M. B. The MEC-4 DEG/ENaC channel of
Caenorhabditis elegans touch receptor neurons transduces mechanical signals. *Nature*
Neuroscience 2004 8:1 **8**, 43–50 (2004).

48. Stober, W., Fink, A., Bohn, E., Chalfie, M. & Au, M. Genetic control of differentiation of the
caenorhabditis elegans touch receptor neurons. *Science (1979)* **243**, 1027–1033 (1989).

49. Wang, X. *et al.* The C. elegans L1CAM homologue LAD-2 functions as a coreceptor in
MAB-20/Sema2–mediated axon guidance. *J Cell Biol* **180**, 233 (2008).

50. Arnatkevičiūtė, A., Fulcher, B. D., Pocock, R. & Fornito, A. Hub connectivity, neuronal
diversity, and gene expression in the Caenorhabditis elegans connectome. *PLoS Comput*
Biol **14**, e1005989 (2018).

51. Baruch, L., Itzkovitz, S., Golan-Mashiach, M., Shapiro, E. & Segal, E. Using Expression
Profiles of Caenorhabditis elegans Neurons To Identify Genes That Mediate Synaptic
Connectivity. *PLoS Comput Biol* **4**, e1000120 (2008).

52. Nonet, M. L. Visualization of synaptic specializations in live C. elegans with synaptic
vesicle protein-GFP fusions. *J Neurosci Methods* **89**, 33–40 (1999).

53. Xuan, Z. *et al.* Clarinet (CLA-1), a novel active zone protein required for synaptic vesicle
clustering and release. *Elife* **6**, (2017).

54. Zhen, M. & Jin, Y. The liprin protein SYD-2 regulates the differentiation of presynaptic
termini in C. elegans. *Nature 1999 401:6751* **401**, 371–375 (1999).

55. McDonald, N. A., Tao, L., Dong, M. Q. & Shen, K. SAD-1 kinase controls presynaptic phase
separation by relieving SYD-2/Liprin-α autoinhibition. *PLoS Biol* **21**, e3002421 (2023).

56. Putrenko, I., Zakikhani, M. & Dent, J. A. A Family of Acetylcholine-gated Chloride Channel
Subunits in Caenorhabditis elegans. *Journal of Biological Chemistry* **280**, 6392–6398
(2005).

57. Steger, K. A. & Avery, L. The GAR-3 Muscarinic Receptor Cooperates With Calcium Signals
to Regulate Muscle Contraction in the Caenorhabditis elegans Pharynx. *Genetics* **167**,
633–643 (2004).

58. Tanizawa, Y. *et al.* Inositol monophosphatase regulates localization of synaptic
components and behavior in the mature nervous system of C. elegans. *Genes Dev* **20**,
3296 (2006).

59.   Bénard, C., Tjoe, N., Boulin, T., Recio, J. & Hobert, O. The Small, Secreted Immunoglobulin
Protein ZIG-3 Maintains Axon Position in Caenorhabditis elegans. *Genetics* **183**, 917
(2009).
60.   Aurelio, O., Hall, D. H. & Hobert, O. Immunoglobulin-domain proteins required for
maintenance of ventral nerve cord organization. *Science (1979)* **295**, 686–690 (2002).
61.   Serrano-Saiz, E. *et al.* A Neurotransmitter Atlas of the Caenorhabditis elegans Male
Nervous System Reveals Sexually Dimorphic Neurotransmitter Usage. *Genetics* **206**,
1251–1269 (2017).
62.   Hammarlund, M., Hobert, O., Miller, D. M. & Sestan, N. The CeNGEN Project: The
Complete Gene Expression Map of an Entire Nervous System. *Neuron* **99**, 430–433
(2018).
63.   Qian, T., Wang, H., Xia, X. & Li, Y. Current and emerging methods for probing
neuropeptide transmission. *Curr Opin Neurobiol* **81**, 102751 (2023).
64.   Ripoll-Sánchez, L. *et al.* The neuropeptidergic connectome of C. elegans. *Neuron* **111**,
3570-3589.e5 (2023).
65.   Peedikayil-Kurien, S., Haque, R., Gat, A. & Oren-Suissa, M. Modulation by NPYR underlies
experience-dependent, sexually dimorphic learning. *bioRxiv* 2023.10.19.563073 (2023)
doi:10.1101/2023.10.19.563073.
66.   Susoy, V. *et al.* Natural sensory context drives diverse brain-wide activity during C. elegans
mating. *Cell* **184**, 5122-5137.e17 (2021).
67.   Lipton, J., Kleemann, G., Ghosh, R., Lints, R. & Emmons, S. W. Mate Searching in
Caenorhabditis elegans: A Genetic Model for Sex Drive in a Simple Invertebrate. *Journal
of Neuroscience* **24**, 7427–7434 (2004).
68.   Liu, T., Kim, K., Li, C. & Barr, M. M. FMRFamide-like neuropeptides and mechanosensory
touch receptor neurons regulate male sexual turning behavior in Caenorhabditis elegans.
J Neurosci **27**, 7174–7182 (2007).
69.   Wang, X. *et al.* The C. elegans L1CAM homologue LAD-2 functions as a coreceptor in
MAB-20/Sema2–mediated axon guidance. *J Cell Biol* **180**, 233 (2008).
70.   Brenner, S. THE GENETICS OF CAENORHABDITIS ELEGANS. *Genetics* **77**, 71–94 (1974).
71.   Zhang, S., Banerjee, D. & Kuhn, J. R. Isolation and Culture of Larval Cells from C. elegans.
PLoS One **6**, e19505 (2011).
72.   Spencer, W. C. *et al.* Isolation of Specific Neurons from C. elegans Larvae for Gene
Expression Profiling. *PLoS One* **9**, e112102 (2014).
73.   Kaletsky, R. *et al.* The C. elegans adult neuronal IIS/FOXO transcriptome reveals adult
phenotype regulators. *Nature 2015 529:7584* **529**, 92–96 (2015).
74.   Katz, M. *et al.* Glutamate spillover in C. elegans triggers repetitive behavior through
presynaptic activation of MGL-2/mGluR5. *Nature Communications 2019 10:1* **10**, 1–13
(2019).
75.   Young, M. D. & Behjati, S. SoupX removes ambient RNA contamination from droplet-
based single-cell RNA sequencing data. *Gigascience* **9**, 1–10 (2020).
76.   Bhattacharya, A., Aghayeva, U., Berghoff, E. G. & Hobert, O. Plasticity of the Electrical
Connectome of C. elegans. *Cell* **176**, 1174-1189.e16 (2019).
77.   Stefanakis, N., Carrera, I. & Hobert, O. Regulatory Logic of Pan-Neuronal Gene Expression
in C. elegans. *Neuron* **87**, 733–750 (2015).

78. Reilly, M. B., Cros, C., Varol, E., Yemini, E. & Hobert, O. Unique homeobox codes delineate
all the neuron classes of C. elegans. *Nature 2020 584:7822* **584**, 595–601 (2020).
79. Harris, T. W. *et al.* WormBase: a modern Model Organism Information Resource. *Nucleic*
Acids Res **48**, D762–D767 (2020).
80. Davison, A. C. & Hinkley, D. V. Bootstrap Methods and their Application. *Bootstrap*
Methods and their Application (1997) doi:10.1017/CBO9780511802843.
81. Canty, A. & Ripley, B. D. boot: Bootstrap R (S-Plus) Functions. Preprint at (2024).
82. Davison, A. C. & Hinkley, D. V. Bootstrap Methods and their Application. *Bootstrap*
Methods and their Application (1997) doi:10.1017/CBO9780511802843.
83. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat*
Methods **9**, 676–682 (2012).
84. Holdorf, A. D. *et al.* WormCat: An Online Tool for Annotation and Visualization of
Caenorhabditis elegans Genome-Scale Data. *Genetics* **214**, 279–294 (2020).
85. Hobert, O. PCR fusion-based approach to create reporter Gene constructs for expression
analysis in transgenic C. elegans. *Biotechniques* **32**, 728–730 (2002).
86. Levi-Ferber, M. *et al.* Neuronal regulated ire- 1-dependent mRNA decay controls germline
differentiation in Caenorhabditis elegans. *Elife* **10**, (2021).
87. Chalfie, M., Hart, A. C., Rankin, C. H. & Goodman, M. B. Assaying mechanosensation.
WormBook (2014) doi:10.1895/WORMBOOK.1.172.1.
88. Li, W., Kang, L., Piggott, B. J., Feng, Z. & Xu, X. Z. S. The neural circuits and sensory
channels mediating harsh touch sensation in Caenorhabditis elegans. *Nature*
Communications 2011 2:1 **2**, 1–9 (2011).
89. Ji, N. & van Oudenaarden, A. Single molecule fluorescent in situ hybridization (smFISH) of
C. elegans worms and embryos. *WormBook* 1–16 (2012)
doi:10.1895/WORMBOOK.1.153.1.
90. Karimi, S., Gat, A., Agazzi, C., Oren-Suissa, M. & Krieg, M. Automated dual olfactory
device for studying head/tail chemosensation in Caenorhabditis elegans. *APL Bioeng* **8**,
(2024).
91. Gat, A. *et al.* Integration of spatially opposing cues by a single interneuron guides
decision-making in C. elegans. *Cell Rep* **42**, (2023).
92. Ahringer, J. Reverse genetics. *WormBook* (2006) doi:10.1895/WORMBOOK.1.47.1.
93. Liu, M., Kumar, S., Sharma, A. K. & Leifer, A. M. A high-throughput method to deliver
targeted optogenetic stimulation to moving C. elegans populations. *PLoS Biol* **20**,
e3001524 (2022).
94. Kumar, S., Sharma, A. K., Tran, A., Liu, M. & Leifer, A. M. Inhibitory feedback from the
motor circuit gates mechanosensory processing in Caenorhabditis elegans. *PLoS Biol* **21**,
e3002280 (2023).
95. Liu, M., Sharma, A. K., Shaevitz, J. W. & Leifer, A. M. Temporal processing and context
dependency in caenorhabditis elegans response to mechanosensation. *Elife* **7**, (2018).
1207