The RNA-binding protein HuR impairs adipose tissue anabolism in pancreatic cancer cachexia

Paige C. Arneson-Wissink[1,2], Katherine Pelz[1,3], Beth Worley[1], Heike Mendez [1,2], Peter Pham [1,2], Grace McCarthy [1,3], Alex Chitsazan[4], Jonathan R. Brody [1,3,5], Aaron J. Grossberg*[1,2,4,5]

[1]Brenden-Colson Center for Pancreatic Care, Oregon Health & Science University, Portland, OR

[2]Department of Radiation Medicine, Oregon Health & Science University, Portland, OR

[3]Department of Surgery, Oregon Health & Science University, Portland, OR

[4]Cancer Early Detection Advanced Research Center, Oregon Health & Science University, Portland, OR

[5]Department of Cell, Developmental, and Cancer Biology, Oregon Health & Science University, Portland, OR

Corresponding Author: Aaron J. Grossberg, 3181 SW Sam Jackson Park Rd Mail Code L481, Oregon Health and Science University, Portland, OR, 97239, 503-494-9945, grossber@ohsu.edu

ABSTRACT

Background: Cachexia is defined by chronic loss of fat and muscle, is a frequent complication of pancreatic ductal adenocarcinoma (PDAC), and negatively impacts patient outcomes. Nutritional supplementation cannot fully reverse tissue wasting, and the mechanisms underlying this phenotype are unclear. This work aims to define the relative contributions of catabolism and anabolism to adipose wasting in PDAC-bearing mice. Human antigen R (HuR) is an RNA-binding protein recently shown to suppress adipogenesis. We hypothesize that fat wasting results from a loss of adipose anabolism driven by increased HuR activity in adipocytes of PDAC-bearing mice.

Methods: Adult C57BL/6J mice received orthotopic PDAC cell ($Kras^{G12D}; p53^{R172H/+}; Pdx1\text{-}cre$) (OT-PDAC) or PBS (sham) injections. Mice exhibiting moderate cachexia (9 days after injection) were fasted for 24h, or fasted 24h and refed 24h before euthanasia. A separate cohort of PDAC mice were treated with an established HuR inhibitor (KH-3, 100 mg/kg) and subjected to the fast/refeed paradigm. We analyzed body mass, gross fat pad mass, and adipose tissue mRNA expression. We quantified lipolytic rate as the normalized quantity of glycerol released from 3T3-L1 adipocytes *in vitro*, and gonadal fat pads (gWAT) *ex vivo*.

Results: 3T3-L1 adipocytes treated with PDAC cell conditioned media (CM) liberated less triglyceride into the culture media than control-treated adipocytes (-28.1%) and had lower expression of lipolysis and lipogenesis genes than control cells. PDAC gWAT cultured *ex vivo* displayed decreased lipolysis compared to sham gWAT (-54.7%). PDAC and sham mice lost equivalent fat mass after a 24h fast, however, PDAC mice could not restore inguinal fat pads (iWAT) (-40.5%) or gWAT (-31.8%) mass after refeeding. RNAseq revealed 572 differentially expressed genes in gWAT from PDAC compared to sham mice. Downregulated genes

(n=126) were associated with adipogenesis (adj p=0.05), and expression of adipogenesis

master regulators *Pparg* and *Cebpa* were reduced in gWAT from PDAC mice.

Immunohistochemistry revealed increased HuR staining in gWAT (+74.9%) and iWAT

(+41.2%) from PDAC mice. Inhibiting HuR binding restored lipogenesis in refed animals with a

concomitant increase in iWAT mass (+131.7%) and genes regulating adipogenesis (*Pparγ*,

Cebpa, *Retn*, *Adipoq*, *Fasn*).

Conclusions: Our work highlights deficient adipose anabolism as a driver of wasting in 3T3-

L1 adipocytes treated with PDAC conditioned media and OT-PDAC mice. The small molecule

KH3, which disrupts HuR binding, was sufficient to restore adipogenic and lipogenic gene

expression and prevent adipose wasting. This highlights HuR as a potentially targetable

regulatory node for adipose anabolism in cancer cachexia.

Key words: pancreatic ductal adenocarcinoma, cachexia, adipose, adipogenesis, HuR

INTRODUCTION

Cancer-associated cachexia is a wasting condition characterized by systemic

inflammation, progressive weight loss, and atrophy of white adipose tissue (WAT) and skeletal

muscle[1-3]. In addition to physical deterioration, individuals with cachexia also exhibit fatigue,

anorexia, and cognitive decline[1-4], which contribute significantly to reductions in quality of

life, ability to tolerate chemotherapy or surgery, and patient mortality[5-9]. Cachexia is

estimated to be the direct cause of death in 20-30% of cancer patients[5, 6], and among all

malignancies, pancreatic ductal adenocarcinoma (PDAC) is the most highly associated with

cachexia, with an estimated 83% of patients suffering from this condition[10-12]. Despite much

of cachexia research focusing on improving skeletal muscle mass, a retrospective study of

patients with PDAC revealed that fat loss alone is associated with equally poor outcomes as

combined muscle and fat mass loss [13]. Additionally, current cachexia clinical trials center

around weight maintenance or gain as a primary endpoint. This highlights the importance of

understanding the drivers of adipose tissue loss in cachexia.

Cachexia arises when energy catabolism exceeds anabolism, leading to unsustainable

levels of fat mobilization and muscle depletion. Multiple factors are known to enhance

catabolism, including decreased secretion of anabolic hormones, and altered metabolism of

protein, carbohydrate, and lipid substrates[14]. Current work suggests that inflammation drives

metabolic abnormalities in cachexia[10, 15]. Pro-inflammatory cytokine activity increases

during cancer progression[16, 17] and systemic inflammation can contribute to wasting by

inducing hypercatabolism in muscle and adipose tissue [14, 18-20]. Existing literature

highlights elevated rates of lipolysis and adipose browning as the primary forces underlying

adipose tissue wasting [21, 22]. Browning, in particular, appears to exert a double effect in

cachexia by both reducing lipid stores and increasing energy expenditure[22]. However,

impaired anabolic processes, like adipogenesis and lipogenesis, also contribute to adipose

tissue loss [23]. Targeting peroxisome proliferator-activated receptor gamma (PPARG), a

transcriptional control point of adipogenesis, with the agonist rosiglitazone was sufficient to

improve fat and muscle mass retention in mice with lung cancer [24]. Tumor-derived factors

are also capable of impairing adipogenesis in cultured 3T3L-1 adipocytes [25, 26].

RNA-binding proteins (RBPs) are essential in governing biogenesis, stabilization,

translation, and decay of mRNA transcripts[27, 28]. Several RBPs regulate alternative splicing

in adipocytes[29-31] and promote white adipose tissue browning [32] by inhibiting the

translation efficiency of mitochondrial mRNAs such as UCP1[33]. Until recently, the function of

most RBPs in adipocytes was largely unexplored. Recent work demonstrated that the RBP

human antigen R (HuR), encoded by the embryonic lethal abnormal vision-like 1 (*ELAVL1*)

gene, which regulates the expression of genes involved in inflammation, stress response, and

apoptosis, is also a repressor of adipogenesis in both white and brown adipose tissues[34, 35].

Our goal was to elucidate the relative contributions of enhanced catabolism and impaired

anabolism on fat wasting by investigating adipose tissue response to different nutritional

contexts and HuR inhibition in cachectic mice.

METHODS

Cell culture

KPC PDAC cells

KPC cells expressing pancreas-specific conditional alleles (*Kras^{G12D}; p53$^{R172H/+}$; Pdx1-*

cre)[36] were stored in liquid nitrogen until use and then maintained in RMPI 1640

supplemented with 10% FBS, 1mM sodium pyruvate, and 50 U/mL penicillin/streptomycin

(Gibco, Gaithersburg, MD) at 37C and 5% CO_2. Conditioned media (CM) was collected from

confluent KPC cells grown in DMEM, 1% pen/strep (to accommodate later culturing of 3T3-L1

cells) after 24 hours incubation, centrifuged at 1,200 g for 10 min, filtered with a 0.2-μm syringe

filter, and used immediately or stored at −80°C.

3T3-L1 adipocytes

3T3-L1 adipocytes (ATCC CL-173) were purchased from ATCC and stored in liquid

nitrogen until use. Cells were first treated with preadipocyte expansion media (DMEM, 10%

bovine calf serum, 1% pen/strep) for 3-4 days. Then, cells were plated at a density of $2x10^5$

cells/6-well dish, or $6.7 x10^3$ cells/96well dish and maintained in preadipocyte expansion media

for 2 days after reaching 100% confluency. Media was changed to adipocyte differentiation

media (DMEM, 10% FBS, 1% pen/strep, 1 uM dexamethasone, 0.5 mM IBMX, 1 ug/mL bovine

insulin) for 2 days before switching to adipocyte maintenance media (DMEM, 10% FBS, 1%

pen/strep, 1 ug/mL bovine insulin) for the duration of the experiment. Cells were cultured for up

to 6 days after start of maintenance media. CM was added in place of adipocyte maintenance

media and was supplemented with FBS and insulin.

Oil red O staining

Diluted Oil Red O stock solution was prepared by mixing concentrated Oil Red O (0.5%

Oil Red O in 100% isopropyl alcohol) 6:4 with deionized water. This solution was allowed to

stand for 10 minutes at 4C and then filtered immediately before use through a 0.22 um filter.

3T3-L1 cells were washed with PBS, then incubated in 4% paraformaldehyde for 30 minutes at

room temperature, then washed with deionized water twice. Cells were incubated in 60%

isopropyl alcohol for 5 minutes, then treated with diluted Oil Red O solution for 15 minutes.

Cells were washed 5 times with deionized water, then red signal was quantified at 490 nm in a

plate reader (BioTek).

Triglyceride quantification

3T3-L1 mature adipocytes were treated with KPC CM or control media for 24 hours,

then media was collected off of 3T3-L1 cells for triglyceride quantification according to

manufacturer instructions (Stanbio Liquicolor Triglycerides Assay #2100, Fisher Scientific).

Absorbance was read at 500 nm on a plate reader (BioTek).

Mouse studies

Wildtype C57BL/6J mice (JAX catalog number 000664) were purchased from The

Jackson Laboratory (Bar Harbor, ME) and maintained in standard rodent housing at 26°C with

12h light/12h dark cycles. Animals used for experimentation were 12-15 weeks of age. Mice

were individually housed for acclimation for 7 days prior to tumor implantation and provided *ad*

libitum access to water and food (Rodent diet 5001; Purina Mills, St. Louis, MO, USA). Food

intake was measured daily. Tumor-bearing animals were euthanized by cardiac puncture

under deep isoflurane anesthesia. Studies were conducted in accordance with the U.S.

National Institutes of Health Guide for the Care and Use of Laboratory Animals and approved

by the Institutional Animal Care and Use Committee of Oregon Health & Science University.

Study-specific manipulations

For fasting studies, mice were transferred to clean cages without food for 24 hours prior

to euthanasia or refeeding. Pair-feeding was used on the study days indicated in the figure

legends by feeding sham mice the average of the food consumed by PDAC mice the day prior.

PDAC mice treated with the HuR inhibitor KH3, were injected intraperitoneally (100 mg/kg) at

6-, 8-, and 10-days post-implantation.

Orthotopic PDAC implantation

Wildtype C57BL/6J mice aged 12-15 weeks received orthotopic PDAC tumor injections ($Kras^{G12D}$; $p53^{R172H/+}$; $Pdx1$-cre) or sham injections. PDAC mice were injected with $1x10^6$ KPC cells into the tail of the pancreas parenchyma in a volume of 23ul while sham animals were treated with an equal volume of PBS. Animals were euthanized 10-11 days after tumor implantation. N and sex are defined on a per-study basis in the figure legends.

Echo magnetic resonance imaging body composition

Lean mass, fat mass, total body water, and free water were measured using whole-body magnetic resonance imaging (MRI) (EchoMRI, Houston, TX). Measurements were taken pre-implantation, pre-fasting and at euthanasia in tumor and sham groups to assess body composition.

Tissue collection and histology

Tissues collected at necropsy were weighed and flash frozen in liquid nitrogen prior to storage at -80C. Tissues for HuR histology were fixed with 4% paraformaldehyde overnight and then transferred to 70% ethanol. Tissues were paraffin-embedded, sectioned, incubated with anti-HuR (1:300 #sc-5261, Santa Cruz Biotechnology Inc, Dallas, TX) and stained using horseradish peroxidase-conjugated secondary antibody and incubation in 3,3'-diaminobenzidine by the Histopathology Shared Resource Core at OHSU. Whole tissue sections were scanned by the Advanced Light Microscopy Core at OHSU. ZEN Digital Imaging for Light Microscopy (RRID:SCR_013672) was then used to obtain at least five 10x images per tissue. ImageJ software was used to quantify the percent staining for each section (56). All staining was quantified by running color deconvolution on 10x images, applying a standard

intensity threshold on the corresponding images, and measuring the percent area covered by

the staining.

Ex vivo lipolysis

Ex vivo lipolysis assays of adipose explants were performed as previously

described[37]. Briefly, gonadal white adipose tissue (gWAT) tissue was collected and cut into

approximately 100 mg samples, minced, then incubated in phenol red-free DMEM containing

2% fatty acid free bovine serum albumin at 37°C for 1 hour. Media was collected and snap-

frozen. The tissue was transferred to a new dish containing media only or media with 10 uM

isoproterenol and incubated for 2 hours at 37C. Media was collected and snap-frozen. Media

aliquots were thawed and analyzed for glycerol using Sigma Glycerol Assay Kit (MAK117)

according to manufacturer's directions.

Quantitative qRT-PCR

Total RNA was extracted from cell pellets or tissues with the E.Z.N.A. Total RNA Kit II

(Omega Bio-Tek Inc., Norcross, GA). cDNA was transcribed with the High-Capacity cDNA

Reverse Transcription Kit (Applied Biosystems, Waltham, MA). Quantitative real-time

polymerase chain reaction (qPCR) was run on the ABI 7300 (Applied Biosystems) using

TaqMan Fast Advanced PCR Master Mix (Applied Biosystems) or SYBR Green Master Mix

(Applied Biosystems). The relative expression was calculated using the $\Delta\Delta C_t$ method with

gene expression relative to beta actin or 18S.

RNA sequencing

RNA sequencing was performed on total RNA isolated as described above. RNA

libraries were prepared and sequenced using the Illumina Nova Seq and HiSeq platform

according to the Illumina Tru-Seq protocol (Novogene, Sacramento, CA). Reference genome

and gene model annotation files were downloaded from genome website browser

(NCBI/UCSC/Ensembl) directly. FASTQ files obtained from sequencing were put through fastp

to trim adapters and then fastqc for quality analysis. STAR 2.7.1a was then used to align to

reference genome (mm10) to trimmed reads after which htseq was used to obtain gene

counts. Raw reads were normalized using DESeq2 (1.38.3). Genes with an adjusted *P*-value

<0.05 found by DESeq2 were assigned as differentially expressed. Gene set enrichment

analysis (GSEA) was performed through the Broad Institute GUI. Normalized counts from

DESeq2 from gWAT were input and analyzed with relationship to the Molecular Signature

Database (msigdb). Significant Hallmark genes (p <0.05) were graph from both positive and

negative normalized enrichment scores (NES). Log2 fold change, p-values and p-adj from

differential expression data obtained through deseq2 were loaded into Qiagen's Ingenuity

Pathway Analysis (IPA) to look at upstream and downstream regulators. Volcano plots created

from differential expression data of pairwise comparison with sham groups being baseline

using ggplot2 (3.5). Significant genes from adipogenesis pathway are labeled. Heatmaps were

created of genes in the adipogenesis pathway regardless of significance using

pheatmap(1.0.12).

Statistical Analysis

Specific statistical tests and sample size for each study is indicated in the figure

legends. Error bars in figures show SEM. Statistical analyses were performed using GraphPad

Prism (version 9; GraphPad Software Inc), and graphs were built using GraphPad Prism

(GraphPad Software Inc) statistical analysis software or R studio (4.2.3) using ggplot (3.5.1). P

values are 2 sided with values less than 0.05 regarded as statistically significant.

Data Availability

Further information and resources, including KPC cells and raw data will be shared

upon reasonable request to Aaron J. Grossberg (grossber@ohsu.edu).

RESULTS

Adipocytes suppress lipogenesis in response to PDAC-derived factors

Given a large body of literature suggesting that PDAC cachexia is partly driven by

increased lipolysis in the adipose tissue, we tested the effect of PDAC cell conditioned media

(KPC CM) on lipolysis *in vitro* [21, 22, 38]. We treated differentiated 3T3-L1 cells with control

maintenance media, or with KPC CM for 6 days. At this time, we observed decreased lipid

droplet accumulation by brightfield microscopy, which was confirmed with Oil Red O (**Figure**

1A-B). Liberated triglyceride (TG) levels, a measure of lipolysis, were significantly lower in the

media of KPC CM-treated cells (**Figure 1C**). Correspondingly, mRNA levels of lipolysis

enzymes adipose triglyceride lipase (*Atgl*) and lipase E (*Lipe*) were significantly decreased in

KPC CM-treated 3T3-L1 cells (**Figure 1D**). To explain the apparent decrease in lipid

accumulation without increased lipolysis, we measured the expression of genes associated

with lipogenesis: fatty acid synthase (*Fasn*), nuclear receptor subfamily 1 group H member 3

(*Nr1h3*), solute carrier family 2 member 4 (*Slc2a4*), and lipoprotein lipase (*Lpl*). All of these

genes were suppressed after KPC CM treatment, indicating that the decreased lipid content in

3T3-L1 cells was due to impaired anabolic activity rather than increased catabolic activity

(**Figure 1E**).

PDAC is associated with decreased fat pad mass *in vivo*

We first sought to assess the effects of pancreatic cancer on metabolism by

characterizing tissue physiology in a murine orthotopic PDAC model. 12-week-old C57BL/6J

mice with PDAC or sham implantations were fed *ad libitum* with or without a 24h fast once

mice reached a moderate cachexia burden (9 days after tumor implantation). Body mass

maintained stable among all groups prior to fasting (**Figure 2A**), whereas PDAC mice

exhibited a small reduction in food intake (**Figure 2B**) compared to sham controls. EchoMRI

body composition analysis of fat mass demonstrated significant decreases in overall adiposity

in PDAC animals from baseline to pre-fast due to cachexia progression and reduction in food

intake (**Figure 2C**). As expected, between pre- and post-fast, both PDAC and sham animals

lost significant overall adiposity (**Figure 2D**). Terminal fat pad mass in fasted PDAC animals

was not significantly altered compared to fasted sham mice (**Figure 2E**), indicating that both

groups exhibit similar rates of fasting-induced lipolysis. Fasting caused a significant reduction

in tumor mass compared to *ad libitum*-fed mice (**Figure 2F**).

PDAC impairs lipolysis and adipogenesis

Based on our findings *in vitro*, we hypothesized that PDAC mice would not exhibit

increased lipolysis, which is characteristic of other cachexia models [21, 38]. To measure the

rate of adipose tissue catabolism, we performed an *ex vivo* lipolysis study, using gonadal white

adipose tissue (gWAT) collected from sham and PDAC mice. We assessed fat pads both in

the presence and absence of the beta-adrenergic agonist isoproterenol (10 uM) to determine

baseline and stimulated lipolysis[39]. Baseline and isoproterenol-stimulated glycerol release

were significantly decreased in PDAC gWAT (**Figure 3A**), demonstrating that PDAC

suppresses, rather than enhances, lipolytic rate and capacity. To further characterize the

molecular biology of adipocytes during PDAC, we assessed mRNA expression of genes

regulating lipolysis and browning in adipose tissue. PDAC mice failed to increase gWAT and

inguinal white adipose tissue (iWAT) *Atgl* and *Lipe* expression in response to fasting (**Figure**

3B). We observed similar decreases in browning-associated gene expression, including

peroxisome proliferator-activated receptor gamma coactivator 1 alpha (*Pgc1a),* protein domain

containing 16 (*Prdm16*), and cell death inducing DFFA like effector a (*Cidea*) in PDAC mice in

both *ad libitum* fed and fasted conditions. Uncoupling protein 1 (*Ucp1*), which is also

associated with browning, was not significantly increased in gWAT from PDAC mice (**Figure**

3B-C). Therefore, neither enhanced lipolysis nor browning are significant contributors to

adipose loss in this model.

PDAC downregulates pathways associated with adipogenesis

To gain further insights into PDAC metabolism from global gene expression analysis,

we performed bulk RNA sequencing (RNAseq) on gWAT from sham and PDAC animals. . In

gWAT, we identified a total of 572 differentially expressed genes (DEGs) in PDAC versus

sham mice, of which 446 were enriched, and 126 were depleted (**Figure 4A, S1**). Pathway

analysis referencing the molecular signatures database (MSigDB) hallmark gene set collection

revealed increased expression of gene sets associated with cell cycle control (e.g., E2F

targets and G2M checkpoints) and inflammation (e.g. TNFa and JAK/STAT3 signaling) and

decreased expression of genes linked to adipogenesis, oxidative phosphorylation, and fatty

acid metabolism (**Figure 4B**)[40]. Because inhibition of adipogenesis could provide an

alternative mechanism for adipose wasting, we plotted all significant DEGs in the adipogenesis

gene set and observed nearly universal depletion of these transcripts in PDAC mice (**Figure**

4C). We then validated our RNAseq data by performing qPCR on selected adipogenesis-

associated DEGs, revealing consistent downregulation of both adipogenesis and lipogenesis

genes (**Figure 4D**). We then repeated qPCR in iWAT to determine whether subcutaneous

adipose exhibited the same transcriptomic changes. Transcriptional changes in iWAT were

less dramatic than in gWAT and did not reach significance, with the exception of *Lpl*, which

was significantly decreased in PDAC tissue (**Figure 4E**). Together, these results indicate that

suppressed transcriptomic programs associated with adipogenesis could account for

decreased adipose tissue during cachexia, in the absence of elevated lipolysis.

Adipose tissue anabolism is impaired in orthotopic PDAC mice after refeeding

Based on our results demonstrating downregulation of adipogenic and lipogenic genes

in PDAC mice, we next wanted to confirm whether adipose tissue anabolism is functionally

impaired in mice implanted with orthotopic PDAC. To do this, wildtype C57BL/6J mice with

PDAC or sham implantations were fed *ad libitum* for 9 days, an established timepoint of active

cachexia[41]*,* fasted 24h, and then terminated or allowed to re-feed for 24 h. To control for

differences in caloric intake, we used a pair-fed refeeding scheme in which sham mice were

refed with the average food consumption of the refed PDAC group. Challenging mice with a

24h fast depletes adipose mass equally in PDAC and sham mice (**Figure 2D**), enabling us to

assess anabolic restoration of fat mass during a 24 h refeeding period. We observed

measured cumulative food intake and daily body mass for the duration of the study (**Figure**

5A-B). Fasting and refeeding did not impact pancreas tumor mass (**Figure S2**). There were no

differences in iWAT or gWAT mass in fasted PDAC versus sham mice (**Figure 5C-D**).

However, while sham mice regain significant amounts of gWAT and iWAT mass after

refeeding, gWAT and iWAT masses in PDAC mice remain equivalent before and after

refeeding (**Figure 5C-D**). Refeeding sham mice caused increased gWAT expression of

lipogenic *Fasn* and adipogenic genes *Lep* and *Retn*, while these genes remained suppressed

in PDAC gWAT (**Figure 5E**). Refed sham iWAT showed increased *Fasn,* but not adipogenic

genes, while PDAC iWAT had suppressed expression of both adipogenic and lipogenic genes

(**Figure 5F**). These results confirm that downregulation in pathways mediating adipogenesis

are indeed accompanied by impaired adipose tissue anabolism in orthotopic PDAC mice after

refeeding.

HuR is overexpressed in orthotopic PDAC fat tissue

Following our observation that adipose tissue anabolism is impaired in PDAC animals,

we next sought to understand the mechanism of this phenomenon. We used Ingenuity

Pathway Analysis upstream regulator analysis to identify potential candidates that could drive

impaired anabolism in PDAC adipose tissue. From the DEGs identified in gWAT of PDAC

mice, we found that canonical regulators of adipogenesis, such as troglitazone, fenofibrate,

PPARA, and PGC1A, were predicted to be inhibited. Inflammatory cytokines associated with

PDAC and cachexia, such as MYD88, IL-6, TNFA, IFNG, and IL-1B, were predicted to be

activated. Among other targets that were predicted to be activated but were not typically

associated with cachexia was human antigen R (HuR, *ELAVL1*), an RNA-binding protein

recently established as a suppressor of adipogenesis[34] (**Figure 6A**). Given the predicted

activation of HuR in gWAT and existing literature, we next asked if HuR was more abundant in

adipose tissue from PDAC mice. We performed immunohistochemical staining of HuR in

formalin-fixed, paraffin-embedded sham and PDAC gWAT, iWAT, pancreas, and muscle

tissue. HuR staining was significantly increased in both the gWAT and iWAT of PDAC mice

versus sham controls (**Figure 6C-D, S4).** We also observed increased HuR staining in the

pancreas, but not skeletal muscle from PDAC mice, as compared to sham controls (**Figure**

6B, S3).

HuR inhibition improves metabolic deficits and restores adipogenesis *in vivo*

Since HuR has been established as driving pro-survival pathways in PDAC, we next

wanted to determine if preventing HuR binding to target mRNAs could reverse adipose wasting

in PDAC cachexia[35]. To do this, we treated PDAC mice with a selective HuR inhibitor, KH-3,

which prevents binding of HuR to its target mRNA sequence[42]. All mice received orthotopic

PDAC tumor injections and were fed *ad libitum* then fasted 24h with or without a 24h refeed at

mid-cachexia (9 days after injection). In each feeding group, mice were treated with the HuR

inhibitor KH-3 (100 mg/kg) or vehicle at 6-, 8-, and 10-days post-injection. Cummulative food

intake between vehicle and KH3 groups was not statistically significant prior to fast. KH3

treated PDAC mice gained significantly more weight during refeeding compared to vehicle-

treated mice (**Figure 7A-B**). Refeeding was associated with larger tumors in both vehicle and

KH-3 groups, likely due to both log growth and the propensity for mice to become dehydrated

during fasting (**Figure 7C**). KH-3 treatment itself had no effect on tumor growth over this time

interval. KH-3 treatment caused a nonsignificant increase in gWAT mass and significantly

increased iWAT mass in refed animals (**Figure 7D-F**). In contrast, vehicle-treated mice did not

significantly increase fat pad mass during refeeding. This effect is specific to adipose tissue, as

gastrocnemius muscle mass was not changed between fast/refeed, or vehicle/KH-3 treatment

(**Figure 7F**). To molecularly test the impact of HuR inhibition on adipogenesis and lipogenesis

gene expression, we analyzed gWAT and iWAT tissues by qPCR. KH-3 treatment successfully

restored expression of adipogenic (*Cebpa, Adipoq, Lep*, *Retn*) and lipogenic (*Fasn*) genes

(**Figure 8A-B, D-E**). HuR inhibition did not impact lipolysis gene expression (*Atgl* and *Lipe*)

(**Figure 8C, F**). Thus, we show a role for HuR in adipose tissue metabolism during PDAC-

associated cachexia, whereby inhibiting HuR successfully ameliorates adipose tissue wasting

in vivo by restoring the expression of genes associated with adipogenesis and lipogenesis.

DISCUSSION

This work evaluates the relative contributions of enhanced catabolism and impaired anabolism on fat wasting by investigating adipose tissue response to different nutritional contexts and HuR inhibition in cachectic mice. We demonstrate that our murine model of PDAC cachexia exhibits a near-complete deficit in adipose tissue anabolism in the context of reduced lipolysis, which corroborates a growing set of literature that challenges the dogma that fat wasting occurs as a result of only enhanced catabolism[24-26]. We further identify HuR as a potential molecular mediator of adipogenesis during pancreatic cancer progression. Collectively, these studies support other work describing heterogeneity within cancer-associated cachexia that could have major implications on therapeutic strategies moving forward [13, 43]. In the future, clinical trials may see more therapeutic success if they employ patient selection based on patients' underlying mechanisms of wasting.

Our study implicates the RNA-binding protein HuR in the inhibition of adipogenesis and lipogenesis in adipose tissue. HuR can impact the transcriptome through both RNA binding and HuR-dependent splicing [35]. One documented mechanism of HuR-mediated adipogenesis regulation is via HuR binding to adipogenesis upstream regulator *Pparg*, which suppresses *Adipoq* protein expression in a non-transcriptionally dependent manner [44]. In our RNA sequencing, we did not observe increased HuR expression, although others have documented increased HuR expression in response to nuclear factor-kappa B (NF-κB) signaling[45]. Alternatively, we identified HuR as an upstream regulator based on adipose transcriptomic changes associated with PDAC. In this context, HuR is more likely controlled post-translationally, although further investigation is needed to identify the specific PDAC-associated factors that may activate HuR stability and nuclear translocation [46].

Small molecule antagonists of HuR, such as KH-3, are currently in development and

have been found to have anti-tumor effects[35, 47]. In our studies, KH-3 ameliorated anti-fat

wasting but also induced severe hemolytic anemia in all treated mice. Tumor size was not

impacted by KH3 treatment, likely because of the short period of dosing. However, studies

aimed to evaluate tumor growth over 5 weeks documented slower tumor growth and fewer

metastases in KH-3-treated mice [48]. In our studies, the anti-cachectic effects of KH-3 were

specific to adipose tissue and did not preserve skeletal muscle mass. Although prior studies

link muscle and adipose wasting, these models are all characterized by enhanced lipolysis and

browning [38]. In our model, adipose and muscle wasting appear to be driven by independent

mechanisms. Continued improvement of HuR antagonists could lead to therapeutics that

restore anabolic potential in adipose tissue, providing a novel approach to alleviating one

symptom of cachexia, while also providing anti-tumor benefits.

In addition to impaired anabolic potential in adipose tissue, our PDAC model also

presents suppressed lipolysis. Suppressed lipolysis could contribute to systemic defects in

energy utilization in cancer cachexia. Prior work from our group shows that PDAC impairs

hepatic lipid oxidation but that this does not result in lipid accumulation or fatty liver [41]. While

hepatic metabolic deficits were not HuR dependent, it is possible that impaired lipid

mobilization contributes to global metabolic disruption. Further interrogation is needed to

understand how improved fat retention through HuR inhibition might impact lipid mobilization

and systemic metabolism.

ACKNOWLEDGEMENTS

We thank all members of the Aaron Grossberg and Jonathan Brody labs for their helpful

discussion and suggestions. We also would like to thank Dr. Laing Xu (Department of

Molecular Biosciences, University of Kansas, Lawrence, Kansas) for generously providing us

with the KH3 compound. Author contributions are: Conceptualization, KP, AJG. Methodology,

KP, HM, PP, GM, AC, AJG. Validation, PCAW, KP, HM, BW, PP. Formal Analysis, PCAW, KP,

HM, BW, PP, GM, AC, AJG. Investigation, PCAW, KP, HM, BW, PP, GM, AC. Writing—

Original Draft, PCAW, BLW. Writing – Review and Editing, PCAW, PP, JB, AJG. Visualization,

PCAW, KP, HM, BW, PP, GM, AC. Supervision, JB, AJG. Project Administration, JB, AJG.

Funding Acquisition, JB, AJG. All authors approved this manuscript.

Funding: This work was supported by National Cancer Institute grants K99CA286709

(PCAW), R37CA280692 (AJG), R01264133 (AJG), and K08245188 AJG), R01 CA212600

(JRB), U01CA224012-03 (JRB), R21 CA263996 (JRB), AACR Grant-15-90-25-BROD (JRB),

the Hirshberg Foundation (JRB), and support from the Brenden Colson Center for Pancreatic

Care (AJG, JRB, and PCAW). This work is also supported by the Histopathology Shared

Resource for pathology studies (University Shared Resource Program at Oregon Health and

Sciences University and the Knight Cancer Institute (P30 CA069533 and P30 CA069533

13S5)).

Conflict of interest: The authors do not declare any conflicts of interest.

Ethical standards: The authors of this manuscript certify that they comply with the ethical

guidelines for authorship and publishing in the Journal of Cachexia, Sarcopenia and

Muscle.[49] All human and animal studies were approved by the appropriate ethics committees

and were therefore performed in accordance with the ethical standards laid down in the 1964

Declaration of Helsinki and its later amendments. All human subjects provided informed

consent and any identifying information of individual patients has been omitted.

FIGURE LEGENDS

Figure 1: Adipocytes suppress lipogenesis in response to pancreatic ductal adenocarcinoma-conditioned media. Differentiated 3T3-L1 adipocytes were cultured for 6 days in control or KPC CM, with the exception of the TG assay. **(A)** Brightfield images of differentiated 3T3-L1 cells treated with control media (right) or KPC CM (left). Scale bars represent 200 um. Representative of 3 wells per condition. **(B)** Quantification of Oil Red O staining. N = 4 wells per condition. **(C)** Media triglyceride levels after 24 hours of exposure to control or KPC CM. N=3 wells per condition. **(D)** mRNA expression lipolysis genes N= 3 wells per condition, normalized to 18S expression. **(E)** mRNA expression of lipogenesis genes. N= 3 wells per condition, normalized to 18S expression. Pairwise analyses tested with t-test. p<0.05, **p<0.01, ***p<0.001, and ****p<0.0001.

Figure 2: Pancreatic ductal adenocarcinoma is associated with decreased fat pad mass *in vivo*. Wildtype C57BL/6J mice with PDAC or sham implantations were fed *ad libitum* or fasted 24h at mid-cachexia (9 days after injection). Animals were euthanized 10 or 11 days after tumor implantation following the 24h fast. N= 5 sham, 6 PDAC male mice per feeding condition. **(A)** Daily body mass. Statistically tested with 3-way ANOVA p<0.0001 for time, time x fast; p=0.0073 for fast; p=0.0097 for time x tumor status; p=0.0167 for tumor status. **(B)** Cumulative food intake. Statistically tested with 3-way ANOVA p<0.0001 for time, time x fast; p=0.0035 for time x tumor status. **(C-D)** Body composition changes in total adiposity were characterized at baseline to pre-fast between sham and PDAC mice **(C)**, and at pre- to post-fast **(D)**. MRI analyses were statistically tested with mixed effects model with Fisher's LSD test. **(E)** Terminal gWAT mass. **(F)** Terminal pancreas/tumor mass. 2x2 analyses were statistically tested with two-way ANOVA with Tukey multiple comparisons. *p<0.05, **p<0.01, and ****p<0.0001.

Figure 3: Pancreatic ductal adenocarcinoma impairs lipolysis and adipogenesis *in vivo*. **(A)** Measurement of lipolysis (glycerol release) in explants derived from gWAT collected 10 days post tumor implantation. Lipolysis was induced by the agonist isoproterenol (10 µM). N= 6 male mice per group. Statistically tested with two-way ANOVA with uncorrected Fisher's LSD test to compare groups. **(B)** mRNA expression lipolysis genes in gWAT collected 10 days post-implantation from 24h fasted and ad lib mice, normalized to 18S expression. N=5 male mice sham fasted; 3 male, 3 female mice sham ad lib; 6 male mice PDAC fasted; 3 male, 2 female mice PDAC ad lib. **(C)** mRNA expression of lipolysis genes in iWAT collected 11 days post-implantation from ad lib mice, normalized to beta actin expression. N= 4 male mice per group. 2x2 data were statistically tested with two-way ANOVA with Tukey multiple comparisons comparing tumor experimental groups to sham controls. Pairwise comparisons were made with unpaired t test. *p<0.05, **p<0.01, ***p<0.001, and ****p<0.0001.

Figure 4: PDAC downregulates pathways associated with adipogenesis. RNAseq on gWAT from ad lib fed PDAC and sham mice collected 14 days post tumor implantation. N = 2 female, 3 male PDAC; 3 female, 2 male sham. **(A)** Volcano plot of differentially expressed genes in gWAT. Blue dots represent significantly downregulated genes (log2FC < -1 and adj p<0.05). while red dots represent significantly upregulated genes (log2FC >1 and adj p<0.05). Grey dots are insignificant. Labels denote significant differentially expressed genes from the

Hallmark Adipogenesis pathway. **(B)** Broad Institute GSEA significantly enriched pathways in
PDAC gWAT relative to sham gWAT (pathway pval <0.05). Blue is downregulated, red is
upregulated. **(C)** Heat map of Hallmark Adipogenesis pathways constituents in PDAC and
Sham gWAT. Scale represents row normalized raw counts. **(D)** qPCR validation of
adipogenesis and lipolysis genes in in gWAT collected 10 days post-implantation from 24h
fasted and ad lib mice. N=5 male mice sham fasted; 3 male, 3 female mice sham ad lib; 6 male
mice PDAC fasted; 3 male, 2 female mice PDAC ad lib. Statistically tested with two-way
ANOVA with Tukey multiple comparisons comparing tumor experimental groups to sham
controls. **(E)** qPCR validation of adipogenesis and lipolysis genes in iWAT from ad lib fed mice
collected 11 days post implantation. N = 4 male mice per group. Statistically tested with
unpaired t test. *p<0.05, **p<0.01, ***p<0.001, and ****p<0.0001.
479
Figure 5: Adipose tissue anabolism is impaired in orthotopic PDAC mice after refeeding.
Wildtype C57BL/6J mice with PDAC were fed *ad libitum* then fasted 24h or fasted 24h followed
by a refeed at mid-cachexia (9 days after injection). Sham mice were pair-fed to PDAC
average food intake from 2 days post-implantation until study end. Animals were euthanized 9
or 10 days after tumor implantation following the 24h fast or *ad libitum* refeed. N= 3 male and 3
female mice per group. **(A)** Cumulative food intake. Statistically tested with 3 way ANOVA.
P<0.0001 for time, p=0.0344 for time x tumor status. **(B)** Daily body mass. Statistically tested
with 3 way ANOVA. P<0.0001 for time, p=0.002 for time x tumor status. **(C)** iWAT mass
normalized to initial body mass, normalized to beta actin expression. **(D)** gWAT mass
normalized to initial body mass, normalized to beta actin expression. **(E)** mRNA expression of
anabolic genes in gWAT. **(F)** mRNA expression of anabolic genes in iWAT. Data represented
in C-F were statistically tested with two-way ANOVA with Tukey multiple comparisons
comparing tumor experimental groups to sham controls. *p<0.05, and ****p<0.0001.
493
Figure 6: HuR is overexpressed in PDAC orthotopic pancreas and fat tissue. (A)
Selected upstream regulator predictions from Ingenuity Pathway Analysis of differentially
expressed genes in gWAT. **(B-D)** Representative images and quantification of
immunohistochemical detection of HuR in tissue from PDAC and sham mice. **(B)** Pancreas
N=4 male and 4 female mice per group. **(C)** gWAT N=4 male and 4 female mice per group. **(D)**
iWAT N= 3 female, 2 male sham and 2 female, 3 male PDAC. Scale bars represent 100 um.
Statistically tested with t test. *p<0.05, **p<0.01, ***p<0.001, and ****p<0.0001.
501
Figure 7: HuR inhibition prevents adipose tissue loss *in vivo*. Wildtype C57BL/6J mice
with PDAC were fed *ad libitum* then fasted 24h or fasted 24h followed by a refeed at mid-
cachexia (9 days after injection). In each feeding paradigm, groups were treated with either
vehicle or the HuR antagonist, KH-3, then euthanized 10 or 11 days after tumor implantation
following the 24h fast or *ad libitum* refeed. Vehicle-treated mice were pair-fed to the KH-3
treated group's average food intake from 8 days post-implantation until study end. N = 7 male
mice per group. **(A)** Cumulative food intake, statistically tested as an unpaired t test of vehicle
vs KH3 treatment at day 9 (prior to fast) p =0.1121. **(B)** Change in body weight after refeeding,
statistically tested with unpaired *t* test. **(C)** Terminal pancreas mass. **(D)** Terminal gWAT mass.
(E) Terminal iWAT mass. **(F)** Terminal gastrocnemius mass. 2x2 analyses were statistically
tested with two-way ANOVA with Tukey multiple comparisons*p<0.05, and **p<0.01.
513

Figure 8: HuR inhibition partially restores adipogenesis and lipogenesis *in vivo*. **(A)** mRNA expression of Adipogenesis genes in gWAT, normalized to beta actin expression. **(B)** mRNA expression of lipogenesis genes in gWAT. **(C)** mRNA expression of lipolysis genes in gWAT. **(D)** mRNA expression of Adipogenesis genes in iWAT, normalized to beta actin expression. **(E)** mRNA expression of lipogenesis genes in iWAT. **(F)** mRNA expression of lipolysis genes in iWAT. Data were statistically tested with two-way ANOVA with Tukey multiple comparisons correction. N = 7 male mice per group. $*p<0.05$, $**p<0.01$, $***p<0.001$, and $****p<0.0001$.

REFERENCES

1.   Tisdale MJ. Biology of cachexia. J Natl Cancer Inst. 1997;89:1763-73.

doi:10.1093/jnci/89.23.1763

2.   Grossberg AJ, Scarlett JM, Marks DL. Hypothalamic mechanisms in cachexia. Physiol

Behav. 2010;100:478-89. doi:10.1016/j.physbeh.2010.03.011

3.   Baracos VE, Martin L, Korc M, Guttridge DC, Fearon KCH. Cancer-associated

cachexia. Nat Rev Dis Primers. 2018;4:17105. doi:10.1038/nrdp.2017.105

4.   Olson B, Marks DL. Pretreatment Cancer-Related Cognitive Impairment-Mechanisms

and Outlook. Cancers (Basel). 2019;11:doi:10.3390/cancers11050687

5.   von Haehling S, Anker MS, Anker SD. Prevalence and clinical impact of cachexia in

chronic illness in Europe, USA, and Japan: facts and numbers update 2016. J Cachexia

Sarcopenia Muscle. 2016;7:507-9. doi:10.1002/jcsm.12167

6.   von Haehling S, Anker SD. Cachexia as a major underestimated and unmet medical

need: facts and numbers. J Cachexia Sarcopenia Muscle. 2010;1:1-5. doi:10.1007/s13539-

010-0002-6

7.   Fearon K, Strasser F, Anker SD, Bosaeus I, Bruera E, Fainsinger RL, et al. Definition

and classification of cancer cachexia: an international consensus. Lancet Oncol. 2011;12:489-

95. doi:10.1016/S1470-2045(10)70218-7

8.   Fearon KC, Glass DJ, Guttridge DC. Cancer cachexia: mediators, signaling, and

metabolic pathways. Cell Metab. 2012;16:153-66. doi:10.1016/j.cmet.2012.06.011

9.   Tisdale MJ. Cachexia in cancer patients. Nat Rev Cancer. 2002;2:862-71.

doi:10.1038/nrc927

10. Aoyagi T, Terracina KP, Raza A, Matsubara H, Takabe K. Cancer cachexia, mechanism

and treatment. World J Gastrointest Oncol. 2015;7:17-29. doi:10.4251/wjgo.v7.i4.17

11. Fearon K, Arends J, Baracos V. Understanding the mechanisms and treatment options

in cancer cachexia. Nat Rev Clin Oncol. 2013;10:90-9. doi:10.1038/nrclinonc.2012.209

12. Mueller TC, Burmeister MA, Bachmann J, Martignoni ME. Cachexia and pancreatic

cancer: are there treatment options? World J Gastroenterol. 2014;20:9361-73.

doi:10.3748/wjg.v20.i28.9361

13. Kays JK, Shahda S, Stanley M, Bell TM, O'Neill BH, Kohli MD, et al. Three cachexia

phenotypes and the impact of fat-only loss on survival in FOLFIRINOX therapy for pancreatic

cancer. Journal of cachexia, sarcopenia and muscle. 2018;9:673-84.

14. Mantovani G, Maccio A, Lai P, Massa E, Ghiani M, Santona MC. Cytokine activity in

cancer-related anorexia/cachexia: role of megestrol acetate and medroxyprogesterone

acetate. Semin Oncol. 1998;25:45-52.

15. Straub RH, Cutolo M, Buttgereit F, Pongratz G. Energy regulation and neuroendocrine-

immune control in chronic inflammatory diseases. J Intern Med. 2010;267:543-60.

doi:10.1111/j.1365-2796.2010.02218.x

16. Argiles JM, Busquets S, Toledo M, Lopez-Soriano FJ. The role of cytokines in cancer

cachexia. Curr Opin Support Palliat Care. 2009;3:263-8. doi:10.1097/SPC.0b013e3283311d09

17. MacDonald N, Easson AM, Mazurak VC, Dunn GP, Baracos VE. Understanding and

managing cancer cachexia. J Am Coll Surg. 2003;197:143-61. doi:10.1016/S1072-

7515(03)00382-X

18.   Carson JL, Hernandez M, Jaspers I, Mills K, Brighton L, Zhou H, et al. Interleukin-13

stimulates production of nitric oxide in cultured human nasal epithelium. In Vitro Cell Dev Biol

Anim. 2018;54:200-4. doi:10.1007/s11626-018-0233-y

19.   Haslett PA. Anticytokine approaches to the treatment of anorexia and cachexia. Semin

Oncol. 1998;25:53-7.

20.   Moldawer LL, Copeland EM, 3rd. Proinflammatory cytokines, nutritional support, and

the cachexia syndrome: interactions and therapeutic options. Cancer. 1997;79:1828-39.

21.   Das SK, Eder S, Schauer S, Diwoky C, Temmel H, Guertl B, et al. Adipose triglyceride

lipase contributes to cancer-associated cachexia. Science. 2011;333:233-8.

22.   Petruzzelli M, Schweiger M, Schreiber R, Campos-Olivas R, Tsoli M, Allen J, et al. A

switch from white to brown fat increases energy expenditure in cancer-associated cachexia.

Cell metabolism. 2014;20:433-47.

23.   Taylor J, Uhl L, Moll I, Hasan SS, Wiedmann L, Morgenstern J, et al. Endothelial Notch1

signaling in white adipose tissue promotes cancer cachexia. Nature Cancer. 2023;4:1544-60.

24.   Langer HT, Ramsamooj S, Dantas E, Murthy A, Ahmed M, Ahmed T, et al. Restoring

adiponectin via rosiglitazone ameliorates tissue wasting in mice with lung cancer. Acta

Physiologica. 2024;e14167.

25.   Jang HJ, Kim HY, Lyu JH, Muthamil S, Shin UC, Kim HS, et al. Bee (Apis mellifera L.

1758) wax restores adipogenesis and lipid accumulation of 3T3-L1 cells in cancer-associated

cachexia condition. Food Science & Nutrition. 2024;

26.   Tien S-C, Chang C-C, Huang C-H, Peng H-Y, Chang Y-T, Chang M-C, et al. Exosomal

miRNA 16-5p/29a-3p from pancreatic cancer induce adipose atrophy by inhibiting

adipogenesis and promoting lipolysis. Iscience. 2024;27:

27.   Cook KB, Kazan H, Zuberi K, Morris Q, Hughes TR. RBPDB: a database of RNA-

binding specificities. Nucleic Acids Res. 2011;39:D301-8. doi:10.1093/nar/gkq1069

28.   Gerstberger S, Hafner M, Tuschl T. A census of human RNA-binding proteins. Nat Rev

Genet. 2014;15:829-45. doi:10.1038/nrg3813

29.   Pihlajamaki J, Lerin C, Itkonen P, Boes T, Floss T, Schroeder J, et al. Expression of the

splicing factor gene SFRS10 is reduced in human obesity and contributes to enhanced

lipogenesis. Cell Metab. 2011;14:208-18. doi:10.1016/j.cmet.2011.06.007

30.   Brosch M, von Schonfels W, Ahrens M, Nothnagel M, Krawczak M, Laudes M, et al.

SFRS10--a splicing factor gene reduced in human obesity? Cell Metab. 2012;15:265-6; author

reply 7-9. doi:10.1016/j.cmet.2012.02.002

31.   Huot ME, Vogel G, Zabarauskas A, Ngo CT, Coulombe-Huntington J, Majewski J, et al.

The Sam68 STAR RNA-binding protein regulates mTOR alternative splicing during

adipogenesis. Mol Cell. 2012;46:187-99. doi:10.1016/j.molcel.2012.02.007

32.   Chou CF, Lin YY, Wang HK, Zhu X, Giovarelli M, Briata P, et al. KSRP ablation

enhances brown fat gene program in white adipose tissue through reduced miR-150

expression. Diabetes. 2014;63:2949-61. doi:10.2337/db13-1901

33.   Dai N, Zhao L, Wrighting D, Kramer D, Majithia A, Wang Y, et al. IGF2BP2/IMP2-

Deficient mice resist obesity through enhanced translation of Ucp1 mRNA and Other mRNAs

encoding mitochondrial proteins. Cell Metab. 2015;21:609-21. doi:10.1016/j.cmet.2015.03.006

34.   Siang DTC, Lim YC, Kyaw AMM, Win KN, Chia SY, Degirmenci U, et al. The RNA-

binding protein HuR is a negative regulator in adipogenesis. Nat Commun. 2020;11:213.

doi:10.1038/s41467-019-14001-8

35.   Finan JM, Sutton TL, Dixon DA, Brody JR. Targeting the RNA-binding protein HuR in

cancer. Cancer Research. 2023;83:3507-16.

36.   Michaelis KA, Zhu X, Burfeind KG, Krasnow SM, Levasseur PR, Morgan TK, et al.

Establishment and characterization of a novel murine model of pancreatic cancer cachexia. J

Cachexia Sarcopenia Muscle. 2017;8:824-38. doi:10.1002/jcsm.12225

37.   Schweiger M, Eichmann TO, Taschler U, Zimmermann R, Zechner R, Lass A.

Measurement of lipolysis. Methods Enzymol. 2014;538:171-93. doi:10.1016/B978-0-12-

800280-3.00010-4

38.   Rupert JE, Narasimhan A, Jengelley DH, Jiang Y, Liu J, Au E, et al. Tumor-derived IL-6

and trans-signaling among tumor, fat, and muscle mediate pancreatic cancer cachexia.

Journal of Experimental Medicine. 2021;218:e20190450.

39.   Schweiger M, Eichmann TO, Taschler U, Zimmermann R, Zechner R, Lass A.

Measurement of lipolysis. In: Elsevier; 2014. pp. 171-93.

40.   Liberzon A, Birger C, Thorvaldsdóttir H, Ghandi M, Mesirov JP, Tamayo P. The

molecular signatures database hallmark gene set collection. Cell systems. 2015;1:417-25.

41.   Arneson-Wissink PC, Mendez H, Pelz K, Dickie J, Bartlett AQ, Worley BL, et al. Hepatic

signal transducer and activator of transcription-3 signalling drives early-stage pancreatic

cancer cachexia via suppressed ketogenesis. J Cachexia Sarcopenia Muscle. 2024;15:975-88.

doi:10.1002/jcsm.13466

42.   Wu X, Gardashova G, Lan L, Han S, Zhong C, Marquez RT, et al. Targeting the

interaction between RNA-binding protein HuR and FOXQ1 suppresses breast cancer invasion

and metastasis. Commun Biol. 2020;3:193. doi:10.1038/s42003-020-0933-1

43.   Fearon Kenneth CH, Glass David J, Guttridge Denis C. Cancer Cachexia: Mediators,

Signaling, and Metabolic Pathways. Cell Metabolism. 2012;16:153-66.

doi:10.1016/j.cmet.2012.06.011

44.   Hwang JS, Lee WJ, Hur J, Lee HG, Kim E, Lee GH, et al. Rosiglitazone-dependent

dissociation of HuR from PPAR-γ regulates adiponectin expression at the posttranscriptional

level. The FASEB Journal. 2019;33:7707-20. doi:10.1096/fj.201802643r

45.   Kang MJ, Ryu BK, Lee MG, Han J, Lee JH, Ha TK, et al. NF-κB Activates

Transcription of the RNA-Binding Factor HuR, via PI3K-AKT Signaling, to Promote Gastric

Tumorigenesis. Gastroenterology. 2008;135:2030-42.e3. doi:10.1053/j.gastro.2008.08.009

46.   Grammatikakis I, Abdelmohsen K, Gorospe M. Posttranslational control of HuR

function. Wiley Interdisciplinary Reviews: RNA. 2017;8:e1372.

47.   Huang Z, Liu S, Tang A, Wu X, Aube J, Xu L, et al. Targeting RNA-binding protein HuR

to inhibit the progression of renal tubular fibrosis. Journal of Translational Medicine.

2023;21:428.

48.   Dong R, Chen P, Polireddy K, Wu X, Wang T, Ramesh R, et al. An RNA-Binding

Protein, Hu-antigen R, in Pancreatic Cancer Epithelial to Mesenchymal Transition, Metastasis,

and Cancer Stem Cells. Mol Cancer Ther. 2020;19:2267-77. doi:10.1158/1535-7163.Mct-19-

0822

49.   von Haehling S, Coats AJ, Anker SD. Ethical guidelines for publishing in the Journal of

Cachexia, Sarcopenia and Muscle: update 2021. Wiley Online Library; 2021. pp. 2259-61.

A 3T3-L1: 6 days post treatment

Control

KPC CM

B Oil Red O

C 3T3-L1 Media TG Levels

D Lipolysis

E Lipogenesis

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 7

Figure 8