1 **Stick together: Isolation and characterization of exopolysaccharides producing bacteria**
2 **from degraded permafrost soils**

3 Muhammad Waqas[1*], Simona Zelenková[1], Cordula Vogel[2], Broder Rühmann[3], Milan Varsadiya[1,4],
4 Patrick Liebmann[5], Haitao Wang[6], Olga Shibistova[5], Tim Urich[6], Georg Guggenberger[5] and Jiří
5 Bárta[1,7]

6
1 University of South Bohemia, Department of Ecosystem Biology, České Budějovice, Czech Republic
2 Technical University Dresden, Institute of Soil Science and Site Ecology, Chair of Soil Resources and Land
Use, Tharandt, Germany
3 Chair of Chemistry of Biogenic Resources, Technical University of Munich, Campus for Biotechnology and
Sustainability, Straubing, Germany
4 University of Bayreuth, Ecological Microbiology, Bayreuth Center of Ecology and Environmental Research,
Germany
5 Leibniz University Hannover, Institute of Earth System Sciences, Section Soil Science, Hannover, Germany
6 University of Greifswald, Institute of Microbiology, Greifswald, Germany
7 University of South Bohemia, Faculty of Science, Centre for Polar Ecology, České Budějovice, Czech
Republic
18
19 *__Corresponding author:__ Department of Ecosystem Biology, Faculty of Science, University of South Bohemia
20 in České Budějovice, Branišovská 31, 37005, Czech Republic. Tel: +420 608264752; E-mail:
21 waqasm00@prf.jcu.cz
22
23

ABSTRACT

Bacterial exopolysaccharides (EPSs) are the high-molecular-weight polymers secreted into their
surrounding that play a crucial role in bacterial survival, environmental adaptation, biofilms formation,
and interaction with surrounding matrices. EPSs may also contribute to soil structure development by
enhancing soil aggregate stability, promoting soil cohesion, and interacting with soil particles through
their diverse functional group constituents and conditioning film characteristics. This study aimed to
isolate and characterize potential EPSs-producing bacteria from the active layer of two different
hydrological landscape of degraded permafrost soils, and from undisturbed intact permafrost soil. A
total of 54 bacterial isolates were obtained, representing three phyla: Firmicutes, Actinomycetota, and
Pseudomonadota. EPSs production was assessed by determining the polysaccharide content measured
as glucose equivalent, and 26 isolates were identified as potential EPSs producers. Among the isolates,
Curtobacterium oceanosedimentum, Frigoribacterium faeni, *Streptomyces* strains, *Neobacillus*
bataviensis and *Mesobacillus subterraneus* exhibited the highest polysaccharide yield. The carbohydrate
content of the extracted EPSs varied in both composition and quantity across the different isolates.
Uronic acids such as glucuronic acid was found in the EPSs produced by the isolates closely related to
Curtobacterium oceanosedimentum and *Neobacillus bataviensis*, while the amino hexoses were
identified in EPSs extracted from various isolates, including those affiliated with *Bacillus*, *Streptomyces*,
Luteimonas, and *Phyllobacterium*. The potential EPSs producing bacteria were also found inhabiting in
the different horizons of both degraded permafrost soil and undisturbed intact permafrost soil, by

determining their relative proportion within the total bacterial community based on 16S rRNA gene
sequences similarities.

Key Words

46 Permafrost; Exopolysaccharides (EPSs); Carbohydrates; Bacteria; Soil aggregates; Soil organic
47 matter; Isolation, Sequencing, DNA

INTRODUCTION

Exopolysaccharides (EPSs) are the main part of the complex mixture of organic biopolymers produced
and secreted by a wide range of bacteria into their surroundings (Martino 2018). EPSs are categorized
into two major types based on the biosynthesis pathway: capsular EPSs, which are tightly associated
with the cell surface forming a capsule, and slime EPSs, which are loosely attached to the cell or
completely diffuse into the extracellular environment (Li & Yang, 2007; Sheng et al., 2010). EPSs are
high-molecular-weight polysaccharides that can be broadly classified into homopolysaccharides and
heteropolysaccharides, depending on their monosaccharide composition (Alvarez-Lorenzo et al., 2013;
Monsan et al., 2001). Homopolysaccharides consist of a single type of monosaccharide and include
examples such as dextran, curdlan, and cellulose. In contrast, heteropolysaccharides are more
structurally complex, incorporating diverse monosaccharides such as glucose, galactose, rhamnose, and
glucuronic acids (De Vuyst & Degeest, 1999; Oleńska et al., 2021; Wingender et al., 2001). Their
structural properties are influenced by specific sugar linkages, branching patterns, and the presence of
functional substituents like pyruvates, succinates, and acetates (More et al., 2014; Navarini et al., 1997).

The diverse monosaccharide composition of EPSs, including glucose, galactose, rhamnose, and
mannose, contributes to polymer flexibility, enabling the formation of cross-linked networks with the
surrounding charge surfaces through covalent bonds, ionic interactions, and hydrogen bonding (Huang
et al., 2022). EPSs are rich in distinct functional groups such as hydroxyl, carboxyl, sulfate, phosphate
and amino groups, which play a crucial role in cell attachment, and can make complexes with
surrounding particles. These functional groups contribute to the anionic or cationic nature of EPSs
molecules, enabling electrostatic interactions and hydrogen bonding with various surfaces, including
minerals and organic matter (Flemming et al., 1996; Ha et al., 2010). The structurally diverse
composition of EPSs along with functional groups, highlights the ecological significance of EPSs-
producing bacteria in soil, where they serve a multitude of purposes primarily associated with the biofilm
formation and maintenance (Balducci et al., 2023), protecting bacterial cells under various
environmental stress, adhesion of the cells to each other, and biofilm attachment to the surface
(Wingender et al. 1999; Sandhya and Ali 2015) cryoprotective potentials (Dubey & Jeevaratnam, 2015).

In harsh environments, such as permafrost soils that remain frozen for two or more consecutive years, the unique properties of EPSs are particularly vital, as they facilitate bacterial survival under extreme cold, nutrient scarcity, and fluctuating water availability. Along with it, EPSs are also considered as the main component in the formation and stabilization of soil structures due to their chemical interactions and gelling characteristics(Costa et al., 2018). Therefore, EPSs released by bacteria may have role in protection of soil aggregates under the degradation of permafrost. However, the composition and activity of bacterial communities in permafrost soils are significantly influenced by permafrost degradation (Deng et al., 2015; Jansson & Taş, 2014; Liu et al., 2019; Steven et al., 2008). When rising in Arctic temperatures triggers an increase in the depth of above ground so called active layer, leading to the degradation of permafrost soils and creating conducive environment for bacterial activity (Knoblauch et al. 2013; Koven et al. 2013). This degradation results in several emerging landscape types from which two distinct hydrological conditions, such as, wet soils, e.g. in areas with high moisture, due lower slope position and poor drainage, and dry soils with better drainage and higher evapotranspiration (Jorgenson and Osterkamp 2005; Jorgenson et al. 2013; Liebmann et al. 2024). These landscapes exert different influences on bacterial community compositions and activities, primarily by promoting the decomposition of exposed organic carbon that was previously preserved in frozen ground (Koven et al. 2013). This exposure enhances nutrient availability, which may facilitate bacterial growth, development, and protection through the high production of EPSs under these two different conditions. One way to gain a better understanding of the physiology and ecology of EPSs-producing bacteria is to isolate them from their natural habitats, determine their abundance within the complex soil microbiome, and characterize their EPSs-producing capabilities under laboratory condition. In this study we aimed 1) to isolate and identify the EPSs-producing bacteria from the undisturbed permafrost soil and two degraded permafrost landscapes under lower and higher moisture regimes, respectively, 2) to determine the produced polysaccharides from individual bacterial isolates and 3) to determine the proportion of these EPSs-producing isolates in the total bacterial community.

MATERIALS AND METHODS

Soil sample collection

Soil samples were collected from two different landscapes with degraded permafrost and from an undisturbed intact permafrost soil (hereafter, dry, wet and intact sites) located near Fairbanks, Alaska, USA. The detailed descriptions of the studied sites are reported in Liebmann *et al.* (2024). For each site, a soil profile was prepared with a depth of 100 cm for the dry and wet sites, while for the intact site, the profile was excavated until the permafrost table was reached in 45-55 cm soil depth. Soil samples were collected per horizon including organic layer, topsoil and subsoil by taking the material at various depth within each horizon. All samples were constantly kept at -20°C before processing. For the genetic

material extraction, approx. three g of soil sample were immersed with two volumes of LifeGuard Soil Preservation Solution (Qiagen, Hilden, Germany) immediately after sampling. All Lifeguard-treated samples were constantly kept at 4°C until processing.

Isolation of bacteria

For the isolation of bacteria, one g of frozen soil from the topsoil of each site was serially diluted in sterile saline solution (0.9% NaCl) in a total volume of 1 ml. 100 µl of 10^{-2} to 10^{-4} dilutions were plated on three distinct culture media. Standard I (1 g glucose L^{-1}, 15 g peptone L^{-1}, 3 g yeast extract L^{-1}, and 6 g NaCl L^{-1}), Reasoner´s 2A agar (Reasoner & Geldreich, 1985) and Nutrient agar (2 g Yeast extract, 5 g Peptone L^{-1}, 1g Lab-Lemco L^{-1} and 5 g NaCl L^{-1}) were utilized to ensure the recovery of a diverse range of bacteria, including general heterotrophic, slow-growing oligotrophic bacteria, and actinomycetes. The plates were incubated at 22°C for 48–72 hours. The 22°C was chosen based on higher CFU (data not shown). After the incubation, well-grown single colonies were picked from each diluted agar plate and streaked onto corresponding fresh agar plates, followed by incubation. The streaking process was repeated a minimum of three times to ensure the isolation of pure colonies. The pure colonies were subsequently grown in the Nutrient liquid medium for DNA isolation.

Screening soil isolates for EPSs production

For screening of EPSs production, soil isolates were streaked on agar medium (EPS medium) specifically designed exopolysaccharides production described in Rühmann et al. (2015). The plates were incubated at 22°C for 48–72 hours. The isolates that were able to grow on EPS medium and formed a mucoidal/slimy colony were considered as potential EPSs-producers. They were further screened for exopolysaccharides production via automated analytical system involves various polysaccharide detection modules, including determination of total glucose content using phenol-sulfuric-acid transformation, and carbohydrate content by High-throughput-1-phenyl-3-methyl-5-pyrazolone (HT-PMP)-derivatization of the carbohydrates coupled to Electrospray Ionization Mass Spectrometry. as described in Rühmann et al. (2016). The production of polysaccharides measured as glucose equivalent by soil isolates exceeding 100 mg L^{-1} were considered as the greater EPSs-producers. This threshold was set to distinguish between isolates with high and low EPSs production, making it easier to focus on those with significant activity.

DNA extraction and Polymerase Chain Reaction

The extracted DNA from soil samples and isolates were quantified by Quantus fluorometer (Promega BioSystems, Sunnyvale, California, USA). For this, 2 µl of DNA sample was added to to 98 µl 1x TE buffer and 100 µl QuntiFluor dye (QuntiFluor dsDNA dye, Promega Corporation, Madison, USA) in a 1 ml clean Eppendorf tube. A blank and standard were also prepared by mixing 100 ul of 1x TE buffer with 100 µl of QuantiFluor dye or standard DNA provided by manufacturer, respectively. The tubes

were incubated in the dark for 10 minutes and the fluorimeter was calibrated by standard and blank

samples before quantifying the DNA samples.

Genomic DNA extracted from isolates was utilized to amplify the 16S rRNA gene using master mix

composed of DNA, ddH$_2$O, PCR buffer, BSA and primers: 9bfm (forward primer: 5'-

GAGTTTGATYHTGGCTCAG-3') and 1512uR (reverse primer: 5'-ACGHTACCTTGTTACGACTT-

3'). For a positive control DNA of *Escherichia coli* (strain ATCC 9637) and for a negative control H$_2$O

were used. The PCR amplification was performed using the PCR thermal cycler (SensoQuest GmbH,

Germany) with the following protocol: initial denaturation for three minutes at 95°C, followed by 30

cycles of each denaturation for one minute at 95°C, annealing for one minute at 52°C and elongation

for 90 seconds at 72°C. The final elongation was for 10 minutes at 72°C. After the amplification, 4 µl

of PCR product was determined by electrophoresis on 1% agarose gel and visualized by trans

illuminator (Azure 200, Azure Biosystem, Inc, US). The length of the band was compared to 1 kb DNA

ladder and the correct PCR products were then sent to SEQme sequencing company for Sanger

sequencing. The obtained sequence data underwent quality trimming and the forward and reverse reads

were joined using Geneious Prime bioinformatics software (Java Version 11.0.18+10 (64 bit). The

nucleotide sequences were then searched in 16S rRNA gene NCBI database by standard nucleotide

BLAST.

Determination of proportion of EPSs isolates in total community

The proportion of the EPSs-producing soil isolates in the total relative proportion of bacterial community

from different horizons of each landscape was determined. For this, 16S rRNA gene sequences from the

EPSs-producing soil isolates were used as queries to search against a local 16S rRNA gene database

constructed from the representative sequences of zero-radius operational taxonomic unit (zOTUs) from

the total bacterial community using BLAST (Figure S1, Supporting Information). Afterward, a subset

of assigned zOTUs sharing sequences with more than 97% similarity with 16S rRNA sequence of pure

isolates was used to calculate their proportion within the total bacterial community across the different

horizons within all the three studied sites.

The relative proportion of the bacterial community was calculated by dividing the count of each zOTUs

by the sum of counts within each sample and further summarized across the sites and horizons. The data

were filtered to include only the phylum level and the mean relative proportions were then determined.

The proportion of EPSs-producing soil isolates among the total bacterial community was calculated by

taking the subset of assigned zOTUs with their relative proportions. The data were then filtered at genus

level and computed to mean relative proportion. All the figures were generated using the ggplot v 3.4.4

package (H, 2016).

RESULTS

Isolation and characterization of potential EPSs-producing bacteria

The bacterial isolation was carried out using three distinct culture media, resulting in the isolation of 17, 21, and 16 bacterial isolates from soil of the dry, wet and intact sites, respectively (Table 1, TableS1, Supporting Information). These isolates were grown on specific polysaccharide production medium for the formation of mucoidal "sticky" colonies, which served as an indicator of their potential to produce EPSs. The majority of soil isolates exhibited growth in the form of mucoidal colonies on medium and signifying their potential as EPSs producers (Table S1, Supporting Information). Out of 54 isolates, 48 were successfully sequenced and classified into three phyla: Firmicutes, Actinomycetota, and Pseudomonadota. Within the Firmicutes, potential EPS producing isolates obtained from all three sites were taxonomically assigned to several genera, including *Bacillus, Peribacillus, Neobacillus,* and *Mesobacillus*. Six isolates were identified as *Bacillus mycoides*, while other six isolates from the wet site were assigned to *Bacillus subtilis*, *Peribacillus simplex*, *Neobacillus bataviensis* and *Mesobacillus subterraneus* (Table 1). Among the Actinomycetota, five isolates from the intact site were assigned to *Frigoribacterium faeni*, *Curtobacterium oceanosedimentum* and *Pseudarthrobacter sp.* and three isolates from the dry site were related to *Microbacterium flavescens* and *Streptomyces sp*. One isolate from the wet site was assigned to *Streptomyces luozhongensis*. Within the Pseudomonadota, isolates were taxonomically classified to *Phyllobacterium sp.* and *Luteimonas arsenica* (Table 1). We found only 5.6% overlapping with one similar isolate such as *Bacillus mycoides* between soil of the dry, wet, and intact sites and another one common isolate such as *Curtobacterium oceanosedimentum* found in soil of both wet and intact sites (Fig. S2, Supporting Information).

Differences in EPSs production by soil isolates

In general, isolates with production of polysaccharide measured as glucose equivalent exceeding 100 mg L^{-1} were considered as potential EPSs-producing isolates (Table 1 and Fig. 1). Two isolates showing similarity to *Curtobacterium oceanosedimentum* and *Neobacillus bataviensis* yielded the highest polysaccharide more than 1000 mg L^{-1} such as 1309 mg L^{-1} and 1149 mg L^{-1} respectively. This was followed by the *Frigoribacterium faeni* and *Mesobacillus subterraneus* exhibited polysaccharide above 900 mg L^{-1}. Other isolates belonged to Firmicutes, such as *Bacillus subtilis*, *Peribacillus simplex*, and *Bacillus mycoides* yielded polysaccharide upto 200 mg L^{-1}, except for the one isolate related to *Bacillus mycoides* (A24) accounted 515 mg L^{-1} polysaccharide content. Similarly, isolates assigned to *Streptomyces sp.*, *Pseudarthrobacter sulfonivorans* and *Microbacterium flavescens* yielded in range of 100 to 350 mg L^{-1}. From the phylum Pseudomonadota, isolate of family Enterobacteriaceae exhibited the polysaccharide upto 667 mg L^{-1}, *Luteimonas arsenica* 227 mg L^{-1} and *Phyllobacterium loti* 177 mg L^{-1}. In total, five isolates from soil of the dry site, 12 isolates from soil of the wet site, and nine isolates

from soil of the intact site were identified as potential high EPSs-producers, as their polysaccharide

equivalent as glucose surpassed 100 mg L^{-1}.

Carbohydrate contents of EPSs produced by soil isolates

The soil isolates identified through 16S rRNA sequencing revealed a diverse array of EPSs enriched

with various carbohydrate derivatives determined by High-throughput-1-phenyl-3-methyl-5-pyrazolone

(HT-PMP)-derivatization of the carbohydrates coupled to Electrospray Ionization Mass Spectrometry

(Table 2). Among the Firmicutes isolates, *Bacillus mycoides* exhibited a substantial presence of hexoses,

particularly glucose (Glc) (up to 65 mg L^{-1}), mannose (Man) (upto 5 mg L^{-1}) and galactose (Gal) (upto

102 mg L^{-1}), along with different levels of amino hexoses, such as glucosamine (GlcN) (up to 62 mg L^{-1}

^{1}), Galactosamine (GalN) (upto 10 mg L^{-1}) and N-acetylglucosamine (GlcNAc) (upto 12 mg L^{-1}). One

isolate of *Bacillus mycoide* (A24) with exception of carbohydrate derivatives in their EPSs were

composed of only Gal (102 mg L^{-1}) and Man (3 mg L^{-1}). The sum of the sugar content in EPSs produced

by isolates of *Bacillus mycoide* was varying 52 to 155 mg L^{-1} (Fig. 2). The isolate *Neobacillus*

bataviensis and *Mesobacillus subterraneus*, that exhibited the highest EPSs among the Firmicutes

showed distinct carbohydrate profiles. For examples, the EPSs produced by *Neobacillus bataviensis* was

composed of Glc (103 mg L^{-1}) and glucuronic acid (GlcUA) (21 mg L^{-1}), along with smaller amounts of

Gal (2 mg L^{-1}), rhamnose (Rha) (6 mg L^{-1}), and GlcN (4 mg L^{-1}). The EPSs produced by *Mesobacillus*

subterraneus was characterized by high Glc content (203 mg L^{-1}) and Man (29%), Rha (12 mg L^{-1}), and

GlcN (6 mg L^{-1}). The sum of the carbohydrate content was higher for the *Mesobacillus subterraneus*

(249 mg L^{-1}) than the *Neobacillus bataviensis* (135 mg L^{-1}). The carbohydrate contents of the EPSs

produced by isolates belonged to Actinomycetota were high in the amount of Glc followed by the Man

and GlcN. The highest EPSs producers such as *Curtobacterium oceanosedimentum* showed a

carbohydrate profile dominated by Glc (114 mg L^{-1}) and glucuronic acid (GlcUA) (33 mg L^{-1}), with

minor amounts of Gal (1 mg L^{-1}) and Rha (7 mg L^{-1}). GlcUA was only found in the EPSs of

Curtobacterium oceanosedimentum, and was not detected in the EPSs of other isolates of

Actinomycetota. *Microbacterium flavescens* on the other hand, Man (12 mg L^{-1}) and Rha (7 mg L^{-1})

were only detected in their produced EPSs. Similarly, the sum of the carbohydrate derivatives was higher

for the isolates of *Frigoribacterium sp.*, followed by the *Curtobacterium oceanosedimentum*, and the

least was detected in the EPSs of *Microbacterium flavescens*. In the Pseudomonadota group, EPSs from

Luteimonas arsenica was of only hexoses, in which the Gal was the highest in amount (125 mg L^{-1}).

While the EPSs of other isolates of Pseudomonadota were more enriched with Glc, GlcN and GlcNAc.

The sum of carbohydrate content was about 120 to 190 mg L^{-1} for the isolates of Pseudomonadota.

Proportion of EPSs-producing isolates in total bacterial community

The 16S rRNA gene sequence of 24 EPSs-producing isolates showed similarities higher than 97% to
the sequence of 13 zOTUs (Table S2). The proportion of assigned zOTUs within the total bacterial
community was different among the three sites as well as within the horizon (Fig. 3). The potential
EPSs-producing isolates represented approx. 2% of total bacterial community in soil of the dry site,
~0.5% in soil of the wet site, but below 0.36% in the soil of the intact site. Among the Firmicutes,
Bacillus subtilis (zOTU3) was predominantly detected in highest proportion in both topsoil (~0.67%)
and subsoil (~0.39%) of the dry site. However, it was almost undetected in the horizons of both the other
sites (intact and wet sites), with levels below 0.01%. In contrast, *Bacillus mycoides* (zOTU2010) was
only found with proportion of 0.03% in the topsoil of the wet site. *Peribacillus simplex* (zOTU2176)
was the second after *Bacillus subtilis* found in high proportion of 0.24% in the topsoil of dry site, while
it was not detectable in the subsoil. Additionally, their proportion in the topsoil of the wet site was 0.03%
and was also detected to have less proportion in the organic layer and subsoil of both intact and wet site.
Similarly, *Neobacillus bataviensis* and *Neobacillus pocheonensis* assigned to zOTU702 and
Mesobacillus subterraneus to zOTU3623 were observed in proportions of 0.14% and 0.02%
respectively, in the topsoil of the wet site, contrasting with their proportions of less than 0.001% in other
horizons across all three sites (Fig. 3).From the Actinomycetota, two assigned zOUTUs, *Streptomyces*
luozhongensis (zOTU1852) and *Streptomyces brevispora* (zOTU1848) were found in high proportions
in the organic layer of the dry site of 0.36% and 0.1%, respectively, while the proportion of
Curtobacterium oceanosedimentum (zOTU2238) was high in the organic layer of the intact site with
0.14%. Moreover, *Frigoribacterium faeni* (zOTU1136), *Pseudarthrobacter sp.* (zOTU1072) and
Microbacterium flavescens (zOTU1055) shared almost similar proportion across the topsoil and subsoil
in all three sites, with the exception of *Pseudarthrobacter sp.*, which was also found in the organic layer
of the dry site (Fig. 3).The *Phyllobacterium endophyticum* of Pseudomonadota assigned to zOTU3875
exhibited a range of high proportion of 0.03% in the topsoil to a low proportion of 0.001% in subsoil of
the wet site, while it was not fund in any horizons of the dry and intact sites. Lastly, *Luteimonas arsenica*
(zOTU2611) was also found in all horizons except organic layer of dry site and subsoil of intact site.

DISCUSSION

Bacterial exopolysaccharides and their carbohydrate content

Bacterial EPSs have long been recognized for their structural diversity, which provides a wide range of
physicochemical and biological properties. The structural properties of EPSs serve various roles,
including concealing the bacterial surface, acting as adhesives for interactions with other bacterial
surfaces or substrates, stabilizing biofilm structures, functioning as signaling molecules and providing
protection against environmental stresses (Flemming & Wingender, 2010; Wingender et al., 1999). We
utilized the cultural base approach together with further screening for exopolysaccharide production and

carbohydrate contents using an automated analytical system. Based on it, 24 bacterial strains were selected as potential EPSs producers out of 54 isolates obtained from the dry and wet permafrost degradation landscapes and undisturbed intact permafrost soil. Two isolates, related to *Curtobacterium oceanosedimentum* and *Neobacillus bataviensis*, were the only strains to exhibit high EPSs production containing uronic acids, distinguishing them from the other isolates. EPSs typically contain neutral carbohydrates, such as hexose and pentose, along with uronic acids like glucuronic and galacturonic acids (More et al., 2014). The functional anionic groups in these uronic acids provide a high cation exchange capacity and enable strong complexation with metal ions (Guibaud et al., 2003; Ha et al., 2010). This binding of EPSs to divalent cations like Ca^{2+} and Mg^{2+} promotes the structural integrity of bacterial aggregates, a key interaction in biofilm formation and stability (Mayer et al., 1999). Additionally, amino hexoses component of EPSs such as glucosamine and galactosamine contain amino groups that interact with surrounding molecules, such as metals and organic compounds (Lahiri et al., 2022; B.-B. Wang et al., 2018). These amino hexoses were identified in EPSs extracted from various isolates, including those affiliated with *Bacillus*, *Streptomyces*, *Luteimonas*, and *Phyllobacterium* (Table 2).

In environments like permafrost soils, where nutrient availability is limited and bacterial survival is challenged by extreme cold and low water activity, the uronic acids and amino hexoses content in EPSs may play important role in microbial adaptation by stabilizing with surrounding charged particles. Soil particle composition, including clay, silt, and organic matter, may directly influence the interaction of EPSs amino sugars, and uronic acids. Mineral components like aluminum and iron oxides, along with organic matter, provide surfaces for metal ion binding through the formation of form charged sites on their surfaces (Kleber et al., 2015; Wagai et al., 2020), which then complexes with amino sugars like glucosamine and galactosamine, and the uronic acids in EPSs. These interactions enhance bacterial stability, biofilm formation, and nutrient retention within the soil matrix (More et al., 2014), and may also enhances the soil's overall porosity and aggregate stability (Zhang et al., 2024) and conserving the soil organic carbon pool from degradation and reducing soil dispersion (Kong et al., 2011). However, the production of EPSs with varying compositions is influenced by external environmental factors such as nutrient availability, moisture, pH and temperature (More et al., 2014), in contrast to the production observed with pure strains under controlled laboratory conditions.

Proportion of potential EPSs-producing isolate in total community

We employed a bioinformatics approach to assess the proportion of potential EPSs-producing soil isolates within the proportion of total bacterial community present in the different horizons of intact and degraded permafrost soil. The utilization of high-throughput sequencing and bioinformatic tools allowed us to discern the taxonomic affiliations and relative abundance of potential EPSs-producing isolates within the overall microbial community structure (Fig. 3 and Table S2, Supporting Information). The

proportion of potential EPSs-producing isolates within total community varied among the three sites, with the community structure exhibited greater diversity in soil of the wet and intact sites compared to the one in the dry site (Fig. 3). However, some of the assigned genera such as *Streptomyces* (zOTU1852 and zOTU1848), *Bacillus* (zOTU3) and *Peribacillus* (zOTU2176) to EPSs producing soil isolates found only or with higher proportion in the different horizon of the dry site (Fig 3). This may be due to the hydrophilic nature of carbohydrates in EPSs due to their hydroxyl groups plays a key role in maintaining bacterial hydration and preventing desiccation and supporting the transport of nutrients in the cold and water-limited conditions of permafrost soils, or even during the freezing period (Ali et al., 2020). Strains of *Streptomyces, Bacillus* and *Peribacillus* are known for their ability to produce EPSs as potential response to low soil moisture level (Wang et al. 2003; Marvasi et al. 2010; Homero et al. 2021) and were also investigated for their role in soil aggregation (Aspiras et al., 1971; Raliya et al., 2014). Furthermore, On the other hand, the assigned genera *Mesobacillus*, *Neobacillus*, *Microbacterium and Pseudarthrobacter* were higher in topsoil and subsoil of the intact and wet sites compared to the dry site, whereas some of them were only found more prominent in the subsoil. Similarly, *Curtobacterium oceanosedimentum* (zOTU2238) was found in wet subsoil and intact throughout all the horizon. The presence of these genera in the wet and intact sites suggests their acclimation to high moisture content and low temperatures through the EPSs production in their surroundings. In permafrost soils, where nutrient is often limited, the hydrophobic regions in certain carbohydrates of EPSs may play a critical role in bacterial survival. These hydrophobic regions help EPSs adsorb organic compounds, including aromatics and aliphatics, that are released from decomposed organic matter (Huang et al., 2022; Ibrahim et al., 2022). By enhancing the binding potential of EPSs, these hydrophobic interactions may assist bacteria in retaining nutrients in environments where organic substrates are scarce or undergoing slow decomposition due to the low temperatures and high temperature.

EPS production, with its varying components can enables bacteria to survive and thrive under diverse environmental conditions. In addition to their role in growth and nutrient retention, EPSs may also contribute to soil structure stability and aggregate formation through the interaction of functional groups and the gelling properties of the EPSs, which could play a key role in maintaining soil cohesion and enhancing aggregate development (Kaci et al. 2005; Vardharajula and Sk 2014; Deka et al. 2019).

CONCLUSION

Bacteria have adopted unique mechanisms to cope with and survive in cold habitats experiencing environmental fluctuations. One of these adaptations includes the production of cryoprotective agents in the form of EPSs (Krembs et al., 2002). These agents also act as a gluing substance, physically promoting soil aggregation (Costa et al. 2018). The current study provides information about the bacterial strains isolated from two degrading permafrost soils and an undisturbed intact permafrost soil. These isolates are particularly important for EPSs production in order to survive in extreme conditions

characterized by low temperatures, varying moisture levels and limited nutrients. We observed a notable

tendency for high polysaccharide production with different carbohydrates derivatives by the soil

isolates, indicating that these are potentially high EPSs producers. They were also observed in varying

proportion within the total bacterial community in both degraded permafrost soils and undisturbed intact

permafrost soil. The results of this study suggest that the presence of bacteria in such environment may

be due to their production of EPSs, that provide protection from environmental stress. This EPSs

production in their nature environment may also influence the soil aggregate formation in different

degradation landscapes of permafrost soils.

FUNDING

The work was financially supported by the joint German-Czech Project "CRYOVULCAN –

Vulnerability of carbon in Cryosols", with the individual grants GU 406/35-1, UR 198/4-1, VO

2111/6-1, GACR project n. 20-21259J

DATA AVAILABILITY

367 **DATA AVAILABILITY**

368 Raw fastq 16S rRNA gene sequences of the whole microbiome were uploaded to European nucleotide

369 archive (ENA) under the study n. PRJEB79237. Sequences of the pure strains were uploaded also to

370 ENA under the project n PRJEB79239.

AUTHOR CONTRIBUTIONS

AUTHOR CONTRIBUTIONS

CV, JB, PL, TU and GG completed fieldwork and collected samples from Fairbanks, Alaska, USA.

MW and SZ screened soil samples for bacterial isolation. MW, SZ performed molecular analysis and

MW JB and MV analyzed the sequencing data. BR screened the isolates for EPSs production, MW

and JB wrote the manuscript, and CV, GG, HW and OS contributed to and have approved the final

manuscript.

REFERENCES

Ali, P., Shah, A. A., Hasan, F., Hertkorn, N., Gonsior, M., Sajjad, W., & Chen, F. (2020). A Glacier
Bacterium Produces High Yield of Cryoprotective Exopolysaccharide. *Frontiers in*
Microbiology, *10*. https://doi.org/10.3389/fmicb.2019.03096

Alvarez-Lorenzo, C., Blanco-Fernandez, B., Puga, A. M., & Concheiro, A. (2013). Crosslinked ionic
polysaccharides for stimuli-sensitive drug delivery. *Advanced Drug Delivery Reviews*, *65*(9),
1148–1171. https://doi.org/10.1016/j.addr.2013.04.016

Aspiras, R. B., Allen, O. N., Harris, R. F., & Chesters, G. (1971). The role of microorganisms in the stabilization of soil aggregates. *Soil Biology and Biochemistry*, *3*(4), 347–353. https://doi.org/10.1016/0038-0717(71)90045-9

Balducci, E., Papi, F., Capialbi, D. E., & Del Bino, L. (2023). Polysaccharides' Structures and Functions in Biofilm Architecture of Antimicrobial-Resistant (AMR) Pathogens. *International Journal of Molecular Sciences*, *24*(4), 4030. https://doi.org/10.3390/ijms24044030

Costa, O. Y. A., Raaijmakers, J. M., & Kuramae, E. E. (2018a). Microbial Extracellular Polymeric Substances: Ecological Function and Impact on Soil Aggregation. *Frontiers in Microbiology*, *9*. https://doi.org/10.3389/fmicb.2018.01636

De Vuyst, L., & Degeest, B. (1999). Heteropolysaccharides from lactic acid bacteria. *FEMS Microbiology Reviews*, *23*(2), 153–177. https://doi.org/10.1111/j.1574-6976.1999.tb00395.x

Deka, P., Goswami, G., Das, P., Gautom, T., Chowdhury, N., Boro, R. C., & Barooah, M. (2019). Bacterial exopolysaccharide promotes acid tolerance in Bacillus amyloliquefaciens and improves soil aggregation. *Molecular Biology Reports*, *46*(1), 1079–1091. https://doi.org/10.1007/s11033-018-4566-0

Deng, J., Gu, Y., Zhang, J., Xue, K., Qin, Y., Yuan, M., Yin, H., He, Z., Wu, L., Schuur, E. A. G., Tiedje, J. M., & Zhou, J. (2015). Shifts of tundra bacterial and archaeal communities along a permafrost thaw gradient in <scp>A</scp> laska. *Molecular Ecology*, *24*(1), 222–234. https://doi.org/10.1111/mec.13015

Di Martino, P. (2018). Extracellular polymeric substances, a key element in understanding biofilm phenotype. *AIMS Microbiology*, *4*(2), 274–288. https://doi.org/10.3934/microbiol.2018.2.274

Dubey, A. K., & Jeevaratnam, K. (2015). Structural characterization and functional evaluation of an exopolysaccharide produced by Weissella confusa AJ53, an isolate from fermented Uttapam batter supplemented with Piper betle L. leaves. *Food Science and Biotechnology*, *24*(6), 2117–2124. https://doi.org/10.1007/s10068-015-0281-y

Flemming, H.-C., Schmitt, J., & Marshall, K. C. (1996). Sorption Properties of Biofilms. In *Sediments and Toxic Substances* (pp. 115–157). Springer Berlin Heidelberg. https://doi.org/10.1007/978-3-642-79890-0_5

Flemming, H.-C., & Wingender, J. (2010). The biofilm matrix. *Nature Reviews Microbiology*, *8*(9), 623–633. https://doi.org/10.1038/nrmicro2415

Guibaud, G., Tixier, N., Bouju, A., & Baudu, M. (2003). Relation between extracellular polymers' composition and its ability to complex Cd, Cu and Pb. *Chemosphere*, *52*(10), 1701–1710. https://doi.org/10.1016/S0045-6535(03)00355-2

H, W. (2016). *Package ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.

Ha, J., Gélabert, A., Spormann, A. M., & Brown, G. E. (2010). Role of extracellular polymeric substances in metal ion complexation on Shewanella oneidensis: Batch uptake, thermodynamic modeling, ATR-FTIR, and EXAFS study. *Geochimica et Cosmochimica Acta*, *74*(1), 1–15. https://doi.org/10.1016/j.gca.2009.06.031

Homero, U., Tortella, G., Sandoval, E., & Cuozzo, S. A. (2021). Extracellular Polymeric Substances (EPS) produced by Streptomyces sp. biofilms: Chemical composition and anticancer properties. *Microbiological Research*, *253*, 126877. https://doi.org/10.1016/j.micres.2021.126877

Huang, L., Jin, Y., Zhou, D., Liu, L., Huang, S., Zhao, Y., & Chen, Y. (2022a). A Review of the Role of Extracellular Polymeric Substances (EPS) in Wastewater Treatment Systems. *International*

Journal of Environmental Research and Public Health, *19*(19), 12191.
https://doi.org/10.3390/ijerph191912191

Ibrahim, H. A. H., Abou Elhassayeb, H. E., & El-Sayed, W. M. M. (2022). Potential functions and
applications of diverse microbial exopolysaccharides in marine environments. *Journal of Genetic
Engineering and Biotechnology*, *20*(1), 151. https://doi.org/10.1186/s43141-022-00432-2

Jansson, J. K., & Taş, N. (2014). The microbial ecology of permafrost. *Nature Reviews Microbiology*,
12(6), 414–425. https://doi.org/10.1038/nrmicro3262

Jorgenson, M. T., & Osterkamp, T. E. (2005). Response of boreal ecosystems to varying modes of
permafrost degradation. *Canadian Journal of Forest Research*, *35*(9), 2100–2111.
https://doi.org/10.1139/x05-153

Kaci, Y., Heyraud, A., Barakat, M., & Heulin, T. (2005). Isolation and identification of an EPS-
producing Rhizobium strain from arid soil (Algeria): characterization of its EPS and the effect of
inoculation on wheat rhizosphere soil structure. *Research in Microbiology*, *156*(4), 522–531.
https://doi.org/10.1016/j.resmic.2005.01.012

Kleber, M., Eusterhues, K., Keiluweit, M., Mikutta, C., Mikutta, R., & Nico, P. S. (2015). *Mineral–
Organic Associations: Formation, Properties, and Relevance in Soil Environments* (pp. 1–140).
https://doi.org/10.1016/bs.agron.2014.10.005

Knoblauch, C., Beer, C., Sosnin, A., Wagner, D., & Pfeiffer, E.-M. (2013). Predicting long-term
carbon mineralization and trace gas production from thawing permafrost of Northeast Siberia.
Global Change Biology, *19*(4), 1160–1172. https://doi.org/10.1111/gcb.12116

Kong, A. Y. Y., Scow, K. M., Córdova-Kreylos, A. L., Holmes, W. E., & Six, J. (2011). Microbial
community composition and carbon cycling within soil microenvironments of conventional, low-
input, and organic cropping systems. *Soil Biology and Biochemistry*, *43*(1), 20–30.
https://doi.org/10.1016/j.soilbio.2010.09.005

Koven, C. D., Riley, W. J., & Stern, A. (2013). Analysis of Permafrost Thermal Dynamics and
Response to Climate Change in the CMIP5 Earth System Models. *Journal of Climate*, *26*(6),
1877–1900. https://doi.org/10.1175/JCLI-D-12-00228.1

Krembs, C., Eicken, H., Junge, K., & Deming, J. W. (2002). High concentrations of exopolymeric
substances in Arctic winter sea ice: implications for the polar ocean carbon cycle and
cryoprotection of diatoms. *Deep Sea Research Part I: Oceanographic Research Papers*, *49*(12),
2163–2181. https://doi.org/10.1016/S0967-0637(02)00122-X

Lahiri, D., Nag, M., Dutta, B., Dey, A., & Ray, R. R. (2022). Bacterial extracellular polysaccharides in
biofilm formation and function. In *Application of Biofilms in Applied Microbiology* (pp. 1–23).
Elsevier. https://doi.org/10.1016/B978-0-323-90513-8.00003-0

Li, X. Y., & Yang, S. F. (2007). Influence of loosely bound extracellular polymeric substances (EPS)
on the flocculation, sedimentation and dewaterability of activated sludge. *Water Research*, *41*(5),
1022–1030. https://doi.org/10.1016/j.watres.2006.06.037

Liebmann, P., Bárta, J., Vogel, C., Urich, T., Kholodov, A., Varsadiya, M., Mewes, O., Dultz, S.,
Waqas, M., Wang, H., Shibistova, O., & Guggenberger, G. (2024). Permafrost degradation and
its consequences for carbon storage in soils of Interior Alaska. *Biogeochemistry*, *167*(3), 199–
223. https://doi.org/10.1007/s10533-024-01132-4

Liu, F., Kou, D., Abbott, B. W., Mao, C., Chen, Y., Chen, L., & Yang, Y. (2019). Disentangling the
Effects of Climate, Vegetation, Soil and Related Substrate Properties on the Biodegradability of

Permafrost-Derived Dissolved Organic Carbon. *Journal of Geophysical Research: Biogeosciences*, *124*(11), 3377–3389. https://doi.org/10.1029/2018JG004944

Marvasi, M., Visscher, P. T., & Casillas Martinez, L. (2010). Exopolymeric substances (EPS) from Bacillus subtilis : polymers and genes encoding their synthesis. *FEMS Microbiology Letters*, *313*(1), 1–9. https://doi.org/10.1111/j.1574-6968.2010.02085.x

Mayer, C., Moritz, R., Kirschner, C., Borchard, W., Maibaum, R., Wingender, J., & Flemming, H.-C. (1999). The role of intermolecular interactions: studies on model systems for bacterial biofilms. *International Journal of Biological Macromolecules*, *26*(1), 3–16. https://doi.org/10.1016/S0141-8130(99)00057-4

Monsan, P., Bozonnet, S., Albenne, C., Joucla, G., Willemot, R.-M., & Remaud-Siméon, M. (2001). Homopolysaccharides from lactic acid bacteria. *International Dairy Journal*, *11*(9), 675–685. https://doi.org/10.1016/S0958-6946(01)00113-3

More, T. T., Yadav, J. S. S., Yan, S., Tyagi, R. D., & Surampalli, R. Y. (2014). Extracellular polymeric substances of bacteria and their potential environmental applications. *Journal of Environmental Management*, *144*, 1–25. https://doi.org/10.1016/j.jenvman.2014.05.010

Navarini, L., Stredansky, M., Matulova, M., & Bertocchi, C. (1997). Production and characterization of an exopolysaccharide from Rhizobium hedysari HCNT 1. *Biotechnology Letters*, *19*(12), 1231–1234. https://doi.org/10.1023/A:1018498106893

Oleńska, E., Małek, W., Kotowska, U., Wydrych, J., Polińska, W., Swiecicka, I., Thijs, S., & Vangronsveld, J. (2021). Exopolysaccharide Carbohydrate Structure and Biofilm Formation by Rhizobium leguminosarum bv. trifolii Strains Inhabiting Nodules of Trifoliumrepens Growing on an Old Zn–Pb–Cd-Polluted Waste Heap Area. *International Journal of Molecular Sciences*, *22*(6), 2808. https://doi.org/10.3390/ijms22062808

Raliya, R., Tarafdar, J. C., Mahawar, H., Kumar, R., Gupta, P., Mathur, T., Kaul, R. K., Praveen-Kumar, Kalia, A., Gautam, R., Singh, S. K., & Gehlot, H. S. (2014). ZnO nanoparticles induced exopolysaccharide production by B. subtilis strain JCT1 for arid soil applications. *International Journal of Biological Macromolecules*, *65*, 362–368. https://doi.org/10.1016/j.ijbiomac.2014.01.060

Reasoner, D. J., & Geldreich, E. E. (1985). A new medium for the enumeration and subculture of bacteria from potable water. *Applied and Environmental Microbiology*, *49*(1), 1–7. https://doi.org/10.1128/aem.49.1.1-7.1985

Rühmann, B., Schmid, J., & Sieber, V. (2015). High throughput exopolysaccharide screening platform: From strain cultivation to monosaccharide composition and carbohydrate fingerprinting in one day. *Carbohydrate Polymers*, *122*, 212–220. https://doi.org/10.1016/j.carbpol.2014.12.021

Rühmann, B., Schmid, J., & Sieber, V. (2016). Automated Modular High Throughput Exopolysaccharide Screening Platform Coupled with Highly Sensitive Carbohydrate Fingerprint Analysis. *Journal of Visualized Experiments*, *110*. https://doi.org/10.3791/53249

Sandhya, V., & Ali, Sk. Z. (2015). The production of exopolysaccharide by Pseudomonas putida GAP-P45 under various abiotic stress conditions and its role in soil aggregation. *Microbiology*, *84*(4), 512–519. https://doi.org/10.1134/S0026261715040153

Sheng, G.-P., Yu, H.-Q., & Li, X.-Y. (2010). Extracellular polymeric substances (EPS) of microbial aggregates in biological wastewater treatment systems: A review. *Biotechnology Advances*, *28*(6), 882–894. https://doi.org/10.1016/j.biotechadv.2010.08.001

Steven, B., Pollard, W. H., Greer, C. W., & Whyte, L. G. (2008). Microbial diversity and activity
through a permafrost/ground ice core profile from the Canadian high Arctic. *Environmental*
Microbiology, *10*(12), 3388–3403. https://doi.org/10.1111/j.1462-2920.2008.01746.x

Torre Jorgenson, M., Harden, J., Kanevskiy, M., O'Donnell, J., Wickland, K., Ewing, S., Manies, K.,
Zhuang, Q., Shur, Y., Striegl, R., & Koch, J. (2013). Reorganization of vegetation, hydrology
and soil carbon after permafrost degradation across heterogeneous boreal landscapes.
Environmental Research Letters, *8*(3), 035017. https://doi.org/10.1088/1748-9326/8/3/035017

Vardharajula, S., & Sk Z, A. (2014). Exopolysaccharide production by drought tolerant bacillus spp.
and effect on soil aggregation under drought stress. *Journal of Microbiology, Biotechnology and*
Food Sciences, *4*(1), 51–57. https://doi.org/10.15414/jmbfs.2014.4.1.51-57

Wagai, R., Kajiura, M., & Asano, M. (2020). Iron and aluminum association with microbially
processed organic matter via meso-density aggregate formation across soils: organo-metallic glue
hypothesis. *SOIL*, *6*(2), 597–627. https://doi.org/10.5194/soil-6-597-2020

Wang, B.-B., Liu, X.-T., Chen, J.-M., Peng, D.-C., & He, F. (2018). Composition and functional group
characterization of extracellular polymeric substances (EPS) in activated sludge: the impacts of
polymerization degree of proteinaceous substrates. *Water Research*, *129*, 133–142.
https://doi.org/10.1016/j.watres.2017.11.008

Wang, L., Li, S., & Li, Y. (2003). Identification and characterization of a new exopolysaccharide
biosynthesis gene cluster from *Streptomyces*. *FEMS Microbiology Letters*, *220*(1), 21–27.
https://doi.org/10.1016/S0378-1097(03)00044-2

Wingender, J., Neu, T. R., & Flemming, H.-C. (1999). What are Bacterial Extracellular Polymeric
Substances? In *Microbial Extracellular Polymeric Substances* (pp. 1–19). Springer Berlin
Heidelberg. https://doi.org/10.1007/978-3-642-60147-7_1

Wingender, J., Strathmann, M., Rode, A., Leis, A., & Flemming, H.-C. (2001). *[25] Isolation and*
biochemical characterization of extracellular polymeric substances from Pseudomonas
aeruginosa (pp. 302–314). https://doi.org/10.1016/S0076-6879(01)36597-7

Zhang, H., Wang, K., Liu, X., Yao, L., Chen, Z., & Han, H. (2024). Exopolysaccharide-Producing
Bacteria Regulate Soil Aggregates and Bacterial Communities to Inhibit the Uptake of Cadmium
and Lead by Lettuce. *Microorganisms*, *12*(11), 2112.
https://doi.org/10.3390/microorganisms12112112

546
547

548 **Table 1.** Taxonomic classification of soil isolates identified as potential EPSs producers, that produced more than
549 100 mg L^{-1}. Isolates sharing more than 97% sequence similarity are presented at the species level and lower than
550 97% are presented at family level. Polysaccharide contents are listed for isolates exceeding 100 mg L^{-1} production

Phylum	ID	Closest Blast Hit 16s rRNA sequencing	% Identity	Locality	Polysaccharide content (mg L^{-1})
Firmicutes					
	S29	*Bacillus mycoides* strain ATCC 6462	99.9	Intact	145.34
	S31	*Bacillus mycoides* strain ATCC 6462	100	Intact	180.69
	S37	*Bacillus mycoides* strain ATCC 6462	97.6	Intact	150.05

A6	*Bacillus proteolyticus* strain MCCC 1A00365	98.0	Intact	170.17
A13	*Bacillus mycoides* strain ATCC 6462	100	Dry	212.37
S28	*Bacillus mycoides* strain ATCC 6462	100	Dry	140.79
A24	*Bacillus mycoides* strain ATCC 6462	100	Wet	515.24
S38	*Bacillus subtilis* strain SBMP4	99.2	Wet	171.48
S39	*Bacillus subtilis* strain SBMP4	98.0	Wet	164.54
S3	*Peribacillus simplex* strain LMG 11160	96.5	Wet	100.0
A25	*Neobacillus bataviensis* strain NBRC 102449	99.0	Wet	1148.6
S19	*Neobacillus pocheonensis* strain Gsoil 420	99.1	Wet	146.0
S6	*Mesobacillus subterraneus* strain COOI3B	98.2	Wet	911.44
Actinomycetota				
S17	*Frigoribacterium faeni* strain 801	99.4	Intact	1309.14
S18	*Frigoribacterium faeni* strain 801	99.5	Intact	735.79
A17	*Curtobacterium oceanosedimentum* strain ATCC 31317	99.3	Intact	1309.14
S15	*Pseudarthrobacter sulfonivorans* strain ALL	98.6	Intact	115.63
S23	Micrococcaceae	91.7	Intact	208.21
A11	*Streptomyces brevispora* strain BK160	99.5	Dry	342.92
A2	*Microbacterium flavescens* strain 401	98.3	Dry	136.85
S9	*Streptomyces luozhongensis* strain TRM49605	99.5	Wet	118.63
S7	*Curtobacterium oceanosedimentum* strain ATCC 31317	98.8	Wet	216.69
Pseudomonadota				
A12	*Luteimonas arsenica* strain 26-35	99.8	Dry	227.26
A22	*Phyllobacterium loti* strain S658	99.0	Wet	177.28
A23	*Phyllobacterium endophyticum* strain PEPV15	99.5	Wet	124.15
S5	Enterobacteriaceae	91.3	Wet	667.23

554 **Table 2:** Sugar derivatives of EPSs produced by soil isolates measured in mg L^{-1}

Phylum	ID	Closest Blast Hit 16s rRNA sequencing	Hexoses			Uronic acids		Deoxy hexoses		Amino hexoses		
			Glc	Man	Gal	GlcUA	GalUA	Rha	Fuc	GlcN	GalN	GlcNAc
Firmicutes	S29	*Bacillus mycoides* strain ATCC 6462	45	4	6	0	0	0	0	38	7	11
	S31	*Bacillus mycoides* strain ATCC 6462	0	3	11	0	0	4	0	25	5	0
	S37	*Bacillus mycoides* strain ATCC 6462	53	5	0	0	0	0	0	43	8	11
	A13	*Bacillus mycoides* strain ATCC 6462	65	5	1	0	0	0	0	62	10	12
	S28	*Bacillus mycoides* strain ATCC 6462	5	5	6	0	0	0	0	42	10	8
	A24	*Bacillus mycoides* strain ATCC 6462	0	3	102	0	0	0	0	0	0	0
	S38	*Bacillus subtilis* strain SBMP4	0	0	11	0	0	0	0	23	5	0
	S39	*Bacillus subtilis* strain SBMP4	0	5	0	0	0	0	0	7	0	0
	A6	*Bacillus proteolyticus* strain MCCC 1A00365	0	6	13	0	0	4	0	32	6	0
	S3	*Peribacillus simplex* strain LMG 11160	0	4	0	0	0	0	0	0	0	0
	A25	*Neobacillus bataviensis* strain NBRC 102449	103	0	2	21	0	6	0	4	0	0
	S19	*Neobacillus pocheonensis* strain Gsoil 420	5	3	11	0	0	4	0	30	0	11
	S6	*Mesobacillus subterraneus* strain COOI3B	203	29	0	0	0	12	0	6	0	0
Actinomycetota	S17	*Frigoribacterium faeni* strain 801	68	67	2	0	0	7	0	4	0	0
	S18	*Frigoribacterium faeni* strain 801	136	46	0	0	0	9	0	6	0	0
	A17	*Curtobacterium oceanosedimentum* strain ATCC 31317	114	0	1	33	0	7	0	0	0	0
	S15	*Pseudarthrobacter sulfonivorans* strain ALL	11	7	0	0	0	0	0	7	0	0
	S23	*Pseudarthrobacter sulfonivorans* strain ALL	103	3	0	0	0	0	0	8	0	0
	A11	*Streptomyces brevispora* strain BK160	101	6	0	0	0	0	0	41	7	9
	A2	*Microbacterium flavescens* strain 401	0	12	0	0	0	7	0	0	0	0
	S9	*Streptomyces luozhongensis* strain TRM49605	75	5	0	0	0	0	0	3	0	0
	S7	*Curtobacterium oceanosedimentum* strain ATCC 31317	10	4	14	0	0	5	0	32	5	13
Pseudomonadota	A12	*Luteimonas arsenica* strain 26-35	52	8	0	0	0	0	0	48	8	14
	A22	*Phyllobacterium loti* strain S658	87	3	8	0	0	0	0	30	6	9
	A23	*Phyllobacterium endophyticum* strain PEPV15	39	4	0	0	0	0	0	46	8	15

S5	*Leclercia adecarboxylata* ATCC 23216 = NBRC 102595 strain LMG 2803	63	4	125	0	0	0	0	0	0	0

555   Glc: Glucose, Man: Mannose, Gal: Galactose, GlcUA: Glucuronic Acid, GalUA: Galacturonic Acid, Rha: Rhamnose, Fuc: Fucose, GlcN: Glucosamine,

556   GalN: Galactosamine, GlcNAc: N-Acetylglucosamine.

559 **Figure 1.** A: Polysaccharide production measured as glucose equivalent in mg L^{-1}.

561 **Figure 2:** Sum of the sugar derivatives determined from the extracted EPSs.

Figure 3. Relative proportion of assigned zOTUs for EPSs-producing soil isolates at genera level in total bacterial community of two sites with degraded permafrost soil (dry and wet site) and site with undisturbed permafrost soil (Intact site).

563
564
565