Structural complexity shapes the global distribution of ecosystems and people

Authors: James Cant[1]*, Nina Schiettekatte[1], Elizabeth M. P. Madin[2], Arie C. Seijmonsbergen[3],
Kenneth F. Rijsdijk[3], Joshua S. Madin[2] & Maria Dornelas[1,4]

Affiliations:

[1]Centro de Ciências do Mar e do Ambiente (MARE), Faculdade de Ciências, Universidade de Lisboa; Lisboa,
Portugal.

[2]Hawai'i Institute of Marine Biology, University of Hawai'i at Manoa; Kaneohe, Hawai'i, USA.

[3]Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam; Amsterdam, The Netherlands.

[4]Centre for Biological Diversity, University of St Andrews; St Andrews, Fife, UK.

*Corresponding author: jicant@ciencias.ulisboa.pt [ORCID: 0000-0002-5529-1752].

Abstract: The three-dimensional structure of land- and seascapes, links biotic and abiotic
processes to shape how organisms use their environments. Much effort has been dedicated to
understanding this relationship between structural complexity and biodiversity. However, tailored,
system-specific approaches to quantifying structural complexity preclude cross-system
comparisons. Expanding a geometric framework developed for quantifying the structural
complexity of local-scale surface environments, we generate high-resolution, global maps of the
key complexity measures of height range, rugosity, and fractal dimension. We illustrate how
patterns in these complexity attributes across the globe reflect the distribution of ecosystems, land
cover types, and the presence of people. Using these tools to understand how structural complexity
determines the arrangement of natural and socio-ecosystems, we can better predict their responses
to ongoing global change.

Main Text:

The physical structure of natural environments sets the stage on which ecological, geological, and climatic processes shape global biodiversity and socio-ecosystems (*1*, *2*). Climatic regimes are a major determinant of global species richness trends (*3*). However, abiotic topography modulates the spatio-temporal configuration of climate systems, thereby mediating local climate exposure, resource availability, and resource access (*4–6*). For instance, the structural complexity of natural landscapes influences the taxonomic and functional diversity of the forest ecosystems they support (*7*). Indeed, some conservation approaches, such as *Conserving Nature's Stage*, a global initiative aimed at incorporating landscape features in conservation planning (*8*), argue that protecting the enduring physical structure of natural habitats is a more effective use of conservation resources than safeguarding more transitory biotic assemblages (*9–13*). The structure of natural environments also contributes to human wellbeing, provides key cultural benefits, and supports agricultural diversification (*2*, *14*). Despite this ecological and socioeconomic value, the term complexity is poorly defined, prompting a lack of consensus for how it is quantified and measured (*15*, *16*). As a result, insight into the role of structural complexity in determining the configuration of natural systems is typically tailored to specific organisms or ecosystems such as terrestrial mammals, inshore reef fish, rocky reefs and mussel beds (*17–20*). Identifying scalable metrics applicable across systems is thus essential for quantifying the general effect of structural complexity in shaping ecological systems across the globe.

Interwoven with long established theory linking changes in area to patterns in species diversity (e.g. (*21–23*)), increases in structural complexity are hypothesised to promote biodiversity (*24*). More structurally complex environments are thought to accommodate a greater diversity of available habitat types and, therefore, support higher species richness (*25*, *26*). Yet, the relationship between complexity and habitat diversity is scale dependent with the greater habitat variety brought about by enhanced complexity likely reducing the size of available habitats (the area-heterogeneity trade-off) (*27*). Thus, increasing structural complexity can also reduce population sizes and increase the likelihood of local extirpation (*28–31*). The importance of evaluating the nuanced effects of structural complexity on the assembly of biological systems has given rise to concepts such as *geodiversity* (*11*). An abiotic analogue of the term biodiversity, geodiversity combines measures of the richness of structural features (e.g. sediment types and landforms) and processes (e.g. riverbed erosion) comprising different environments (*32*). By synthesising the chemical, geological, and physical features comprising natural environments, geodiversity offers a proxy measure of habitat diversity that has been associated with the maintenance of ecosystem function and functional diversity (*33*, *34*). However, for identifying mechanistic drivers, there is a tension between synthesising the characteristics of underlying abiotic variables and evaluating their individual attributes. To establish the association between abiotic structure and the functioning of biotic systems, we must disentangle how the structural complexity of global landscapes influences the natural assemblages they can support (*33*, *35*).

Here, we present a quantitative global-scale assessment of structural complexity patterns using a geometric framework originally developed to quantify the structural complexity of coral

reef ecosystems (*36*) and forest canopies (*7, 37*). Complementing efforts to classify global
ecosystem types (*5*) and map the distribution of land use practices (*38*) and human occupancy (*39*),
we evaluate how ecosystem and land cover classifications correspond with global patterns in the
structural complexity attributes of *height range* (*ΔH*), *rugosity,* (*R*) and *fractal dimension* (*D*).
Describing the shape of surfaces, *ΔH* captures the variation between maximum and minimum
elevation, *R* represents a measure of true surface area, and *D* details the shape-filling capacity of a
surface (*i.e.*, increases in *D* correspond with an increase in contortion) (*40, 41*). Specifically, we
set out to test two key hypotheses. Firstly, (H1) we expect that similar ecosystem types will exhibit
similar structural attributes, with structural characteristics refining local climate exposures and
thus driving the establishment of similar biotic communities (*42, 43*). Secondly, (H2) we
hypothesise that human activities will be concentrated in less complex environments, since
anthropogenic pressures are often attributed to simplifying natural landscapes (*44–46*). Overall,
we demonstrate the value of using geometric descriptors of structural complexity to detect effects
of abiotic structure influencing the global configuration of ecosystems, land use, and socio-
ecosystems.
78
Results
Quantifying global complexity patterns
Using a global terrain model sourced from the General Bathymetric Chart of the Oceans (GEBCO)
(*47*) we derived estimates of height range (*ΔH*), rugosity (*R*), and fractal dimension (*D*) for the
Earth's surface at a resolution of 1.87 km, reflecting ~185 million grid cells. Across the globe, we
report that estimates of *ΔH* and *R* align with patterns of topographic relief (Fig. 1A & B). Higher
ΔH and *R* estimates have been shown to reflect rougher benthic surfaces in both rocky intertidal
and coral reef environments (*36, 48*). At the global scale, this relationship is maintained, with
higher *ΔH* and *R* estimates corresponding with regions characterised by frequent tectonic activity
such as the Himalayan, Andean, and Mid- Atlantic mountain ranges, and the Pacific Ring of Fire
(a high density of plate boundaries around the coastal Pacific region associated with increased
volcanic and tectonic activity (*49*)). However, unlike *ΔH*, *R* also provides a representation of the
space made available by increasing roughness and offers insight into finer-scale variation across
surface gradients (*1, 36*). Indeed, focusing on selected locations within the Amazon, the Mariana
Trench, and the Sahara Desert (Fig. 1) illustrates how it is possible to distinguish the
topographically complex canyon at the entrance to the Mariana Trench and the topographic
channels associated with the Amazon River and Saharan sand dunes using just their associated
rugosity values.
Areas of high fractal dimension (*D*) occur in the arctic tundras of North America and
Siberia, the Amazonian and Patagonian regions of South America, and in central Africa (Fig 1C).
A notable difference between the fractal properties of marine and terrestrial environments is also
evident, with the global seabed typically having lower fractal dimension than the terrestrial
landscape (Fig. 1C; see Appendix S2 for further details). Fractal dimension relates to the intricacy
of a surface, relative to its elevation, with higher *D* suggesting increased convolution (*36*).

Accordingly, differences between the fractal dimension of terrestrial and marine landscapes are
consistent with the prominence of sedimentation throughout deep sea habitats (*50*), which, in areas
of low tectonic activity and high sediment influx, likely buries bathymetric lows and the more
intricate features of marine landscapes (*51*). Geometric constraints ensure that the intricacy of a
surface comes at the expense of its topographic complexity; such that increases in D correspond
with declines in ΔH and R (when either R or ΔH are held constant) (*36*) (Appendix S1).
Characterised by regions of geologically younger and less rugged terrain, rainforest river basins
such as the Amazon are therefore capable of possessing high fractal dimension (Fig. 1).
Alternatively, when viewed through the lens of its fractal dimension, the main canyon of the
Mariana Trench almost disappears, as its surface complexity is small relative to its associated large
changes in elevation (Fig. 1). Intriguingly, with higher fractal dimension often associated with an
increase in habitat diversity and resource partitioning (*7, 25, 52, 53*), perhaps it is not coincidental
that highly diverse rainforest ecosystems are affiliated with environments characterised by higher
fractal dimension, allowing for greater habitat variety, and lower elevational changes, minimising
the development of topographic barriers.
118
Structural complexity across ecosystems and land cover types
The geometric attributes of differing environments position global ecosystem types along a
spectrum of structural complexity (Fig. 2). To delineate the structural signatures associated with
differing global ecosystems we quantified R and D across the distributions of different IUCN
ecosystem functional groups (EFGs; *sensu* (*5*)). Despite rugosity and fractal dimension estimates
varying considerably across global EFGs, we highlight an association between particular
complexity signatures and particular ecosystem types (*H1*) (Figs. 2A, S4 & S5). Marine
ecosystems stand out as being characterised by a higher prevalence of less fractal and less rugose
environments (Fig. 2A). Meanwhile, whilst transitional coastal ecosystems possess the lower
rugosities of their fully marine counterparts (*54*), they exhibit higher fractal dimension (Fig 2A).
Across the terrestrial realm, Polar and Alpine landscapes and forest and woodland ecosystems are
typically more rugose environments, whilst desert and grassland ecosystems have lower height
variation. Wetland ecosystems also comprise lower height variation but possess very high fractal
dimension. Our spectrum of ecosystem complexities also illustrates how, irrespective of their
associated realm, anthropogenically-modified habitats are associated with lower rugosities
compared to corresponding natural environments (Fig. 2A).
Cross-system similarities in ecosystem complexities offer insights into the causal
mechanisms and processes involved in generating and maintaining different ecosystem types.
Coastal and freshwater ecosystems typically possess comparatively high fractal dimension (Fig.
2B). One exception to this pattern are permanent salt lakes (e.g. the Atacama Desert), which
display low structural complexity compared to freshwater ecosystems, and instead possess similar
geometric attributes to sclerophyll desert ecosystems (Fig 2B). Both permanent salt lakes and
sclerophyll deserts are found in semi-arid plains (*55, 56*), the key difference between them being
the proximity of salt lake catchments to sources of periodic water inflow (*57*). Cross-system

similarities in ecosystem complexities are also evident between permanent marshlands and river
deltas (Fig. 2B). Aside from the association of river deltas with saline conditions, these ecosystem
types are largely indistinguishable and possess the shared characteristics of a high deposition of
silt and clay-based sediments, constant or periodic inundation, and high turbidity (*58*). Indeed,
permanent marshlands and river deltas are often components of the same environments (e.g. the
Okavango Delta). We also demonstrate similar attributes of rugosity and fractal dimension across
oceanic temperate rainforests and polar tundra ecosystems (Fig 2B). Presently, these ecosystems
support very different biological communities due to their differing climate exposures. However,
during the mid-Cretaceous (the Turonian–Santonian age, circa 92 to 83 million years ago), when
a far warmer global climate maintained polar temperatures > 14ºC (*59*), the structural complexity
associated with contemporary polar tundra ecosystems supported diverse oceanic temperate
rainforests (*60*). Together these highlighted examples evidence the mediatory role structural
complexity plays in delineating global ecosystems through time and space.

We illustrate how patterns in rugosity and height range correlate with patterns in the
recently established measure of geodiversity (Fig. 3), which represents the combined richness of
geological, geomorphological and hydrological features comprising different environments.
However, fractal dimension and geodiversity exhibit no apparent relationship (Fig. 3), suggesting
that they are orthogonal descriptors of environments. To further distinguish the mechanisms
through which the structure of natural environments modulates the biotic assemblages they
support, we compared the different but complementary perspectives of structural complexity and
geodiversity across the terrestrial landscape. Aligning global maps of terrestrial geodiversity with
our global maps of structural complexity, we observed how environments comprising a greater
richness of different geological and hydrological features typically have a more rugged topography
(Fig 3A & B). Alternatively, geodiversity and fractal dimension offer different perspectives on the
structural regimes of natural environments (Fig 3C). Although increases in both geodiversity and
fractal dimension have been reported to increase biodiversity (*25, 61–63*), often their effects on
diversity are complex and non-linear (*64*). Consequently, it would be worthwhile exploring how
the interaction between a system's abiotic richness and its fractal dimension mediates resource
availability and how organisms exploit them.

Finally, our evaluation of global structural complexity patterns continually emphasises
evidence in support of our hypothesis that (*H2*) human societies are typically associated with more
lowland environments with reduced elevational variation (Figs. 4). We quantified the rugosity and
fractal dimension patterns associated with the global distributions of different land cover
classifications obtained from the Copernicus Climate Change Service (C3S) land cover survey
(*38*). Although clustering rugosity and fractal dimension estimates by associated land cover
categories represents a coarse filter, it is apparent how land cover corresponding with increased
human modification (*i.e.*, urban and cropland environments) are associated with lower estimates
in both R and D, compared to more 'natural' land cover (Fig. 4A). Our findings also suggest that
human occupancy, but not density, is strongly associated with the structural complexity of natural
environments. By overlaying gridded human population densities sourced from NASA (*39*) onto

our maps of global complexity patterns we evaluated how human population densities worldwide
correspond with patterns in ΔH, R, and D. Specifically, the likelihood of observing human
population counts > 0 decreases with increasing ΔH, R, and D (Fig 4B). However, we only observe
a directional association between fractal dimension and human population density, with the density
of human populations decreasing with increasing fractal dimension (Fig 4B[iii]).
188
Discussion
We demonstrate here that using geometric measures to quantify cross-system patterns in global
structural complexity can inform our understanding of the processes through which structural
complexity delineates abiotic and biotic regimes. Understanding the drivers of biodiversity, and
how natural communities respond to ongoing global change, requires an appreciation for the
intrinsic association between biodiversity and abiotic structural complexity (*65*). Colonisation,
extinction, and speciation are fundamental drivers of global patterns in species richness (*66*, *67*),
and are acutely linked to the structural shape of global landscapes over time (*6*). Moreover,
landscape morphology and structure mediate whether, how, and when, organisms can access
available resources (*6*). Indeed, by shaping dispersal pathways and niche partitioning trends,
historic shifts in landscape structure are thought to have influenced the contemporary arrangement
of global biodiversity more so than prevailing climatic regimes (*65*, *68–70*).
Emphasising the value of considering both climatic regimes and structural complexity, our
findings offer numerous quantitative insights supporting the notion that the interaction between
structural complexity and climate shapes biodiversity (*1*). Increases in the fractal dimension of
natural landscapes correspond to increasing habitat diversity, thereby supporting species
coexistence through enhanced resource partitioning (*52*, *71*, *72*). Increases in fractal dimension
could, therefore, be expected to increase speciation by fostering elevated rates of diversification.
Indeed, we have shown how the Amazonian Basin, considered amongst the planet's most
biodiverse ecosystems (*73*, *74*), is an environment dominated by the fractal dimension of its
complexity signature. Nevertheless, the complexity signature of the Sahara Desert, a
comparatively more inhospitable environment, is similarly dominated by its fractal dimension. Yet
rather than weaken the argument that structural complexity plays a role in configuring biodiversity,
this scenario elevates the importance of considering the interaction between local climate regimes
and abiotic structural complexity (*4*). Moreover, alongside illustrating similarities across the
structural regimes associated with ecosystems supporting similar biotic communities, we highlight
how the association between biodiversity and structural complexity is not static through time. The
biotic components of natural ecosystems are typically more transient through time than their
geological counterparts (*2*). Despite being associated with similar structural regimes, oceanic
temperate rainforests and the polar tundra represent very different ecosystems. Yet, the historical
biotic communities associated with polar tundra ecosystems under a warmer climate were similar
to the contemporary biotic communities of oceanic temperate rainforests. Likewise, although
presently comparatively species poor, under differing past climate regimes, the Sahara Desert

supported vibrant grassland communities (*75, 76*). Overall, these examples highlight the key
underlying role of abiotic structural complexity in shaping global biodiversity over time.

The relationship between species diversity and area has long been central to our
understanding of biodiversity theory (*77*), yet environmental context underpins how the
relationship actually manifests (*78, 79*). Applying our quantification of structural complexity to
the assessment of avian species-area relationships observed across global island systems highlights
the nuanced insights structural measures can offer into the mechanisms mediating the
configuration of global species and functional diversity patterns (Box 1). Area is important for
providing the space necessary for species to coexist (*24, 67, 80*). Whether changes in available
area constitute increases or decreases in rugosity, height range, or fractal dimension will determine
whether these changes actually benefit or limit species diversity. Using the geometric measures of
structural complexity it is possible to quantify and detect the mechanisms (e.g. changes in habitat
diversity, niche size and/or species isolation) through which changes in area influence the
configuration of biotic communities. Accordingly, using these measures we can reconcile
understanding of the nuanced effects of area on species diversity patterns across ecosystems and
taxa.

Human presence is negatively correlated with increasing landscape complexity. Beyond its
effects on biodiversity, the complexity of natural landscapes is known to have profound impacts
on multiple aspects of human society and its development (e.g. tourism, agriculture, mobility, and
wellbeing) (*2, 14*). For instance, with mountain ranges and high alpine regions typically associated
with a reduction in the space and arable land needed for agriculture and food production, the
development of human settlements has often gravitated towards flatter, more lowland
environments (*81*). Accordingly, environments exposed to the greatest levels of anthropogenic
pressure and exploitation are those that typically possess reduced structural complexity. However,
this association is unlikely to be solely due to the selective bias of humans having historically
selected less complex environments for development (*14, 82*). Instead, increased anthropogenic
pressure being associated with reduced structural complexity is also indicative of human activities
simplifying natural landscapes through the active removal of geometric and compositional
structure (*44–46, 83, 84*). While we are unable to distinguish between cause and effect with our
static layers, the two processes are not mutually exclusive, and both instances are likely true across
the global landscape.
253

Conclusions
Our global-scale assessment suggests that geometric structural complexity plays a substantial role
in mediating the configuration of natural and anthropogenic communities. Having long appreciated
the association between diversity and area (*7, 77*), a recent focus of ecological research has been
to quantify the mechanistic drivers linking biodiversity and the physical structure of natural
environments (*10, 33, 66*). Nature also does not exist in a vacuum, with the majority of global
ecosystems facing some form of anthropogenic pressure (*85–87*). How we interact with and use
our local environment is mediated by patterns in their structural complexities. So, too, is how these

natural environments respond to our activities. The use of geometric theory to quantify and isolate patterns in the structural complexity of global environments can provide continuous quantitative variables for delineating ecotones based on their shared structural characteristics. Moreover, we have demonstrated the value of including measures of geometric structural complexity within assessments into the drivers of spatial biodiversity patterns. Overall, our approach offers insight into the future configuration of global biodiversity by serving as a basis for evaluating how structural complexity helps to shape biotic communities, and how this association will be influenced by anthropogenic pressures and ongoing global change.

References and Notes:

1. J. S. Madin, M. Asbury, N. Schiettekatte, M. Dornelas, O. Pizarro, J. Reichert, D. Torres-Pulliza, A word on habitat complexity. *Ecol. Lett.* **26**, 1021–1024 (2023).

2. J. Alahuhta, H. Tukiainen, M. Toivanen, T. Ala-Hulkko, J. Hjort, T. Lankila, T. Maliniemi, S. Puhakka, H. Salminen, M. Seppänen, R. Korpelainen, D. Ding, Acknowledging geodiversity in safeguarding biodiversity and human health. *Lancet Planet Health* **6**, 987–92 (2022).

3. G. Pecl, M. B. Araujo, J. D. Bell, J. Blanchard, T. C. Bonebrake, Biodiversity redistribution under climate change: Impacts on ecosystems and human well-being Publication Details. *Science* **355**, 1–9 (2017).

4. M. T. P. Coelho, E. Barreto, T. F. Rangel, J. A. F. Diniz-Filho, R. O. Wüest, W. Bach, A. Skeels, I. R. McFadden, D. W. Roberts, L. Pellissier, N. E. Zimmermann, C. H. Graham, The geography of climate and the global patterns of species diversity. *Nature* **622**, 537–544 (2023).

5. D. A. Keith, J. R. Ferrer-Paris, E. Nicholson, M. J. Bishop, B. A. Polidoro, E. Ramirez-Llodra, M. G. Tozer, J. L. Nel, R. Mac Nally, E. J. Gregr, K. E. Watermeyer, F. Essl, D. Faber-Langendoen, J. Franklin, C. E. R. Lehmann, A. Etter, D. J. Roux, J. S. Stark, J. A. Rowland, N. A. Brummitt, U. C. Fernandez-Arcaya, I. M. Suthers, S. K. Wiser, I. Donohue, L. J. Jackson, R. T. Pennington, T. M. Iliffe, V. Gerovasileiou, P. Giller, B. J. Robson, N. Pettorelli, A. Andrade, A. Lindgaard, T. Tahvanainen, A. Terauds, M. A. Chadwick, N. J. Murray, J. Moat, P. Pliscoff, I. Zager, R. T. Kingsford, A function-based typology for Earth's ecosystems. *Nature* **610**, 513–518 (2022).

6. C. Badgley, T. M. Smiley, R. Terry, E. B. Davis, L. R. G. DeSantis, D. L. Fox, S. S. B. Hopkins, T. Jezkova, M. D. Matocq, N. Matzke, J. L. McGuire, A. Mulch, B. R. Riddle, V. L. Roth, J. X. Samuels, C. A. E. Strömberg, B. J. Yanites, Biodiversity and Topographic Complexity: Modern and Geohistorical Perspectives. *Trends Ecol. Evol.* **32**, 211–226 (2017).

7. M. Ehrbrecht, D. Seidel, P. Annighöfer, H. Kreft, M. Köhler, D. C. Zemp, K. Puettmann, R. Nilus, F. Babweteera, K. Willim, M. Stiers, D. Soto, H. J. Boehmer, N. Fisichelli, M. Burnett, G. Juday, S. L. Stephens, C. Ammer, Global patterns and climatic controls of forest structural complexity. *Nat. Commun.* **12**, 519 (2021).

8.   P. Beier, M. L. Hunter, M. Anderson, Special Section: Conserving Nature's Stage. *Conserv.*
Biol. **29**, 613–617 (2015).

9.   J. J. Lawler, D. D. Ackerly, C. M. Albano, M. G. Anderson, S. Z. Dobrowski, J. L. Gill, N.
E. Heller, R. L. Pressey, E. W. Sanderson, S. B. Weiss, The theory behind, and the
challenges of, conserving nature's stage in a time of rapid change. *Conserv. Biol.* **29**, 618–
629 (2015).

10.   F. Schrodt, J. J. Bailey, W. D. Kissling, K. F. Rijsdijk, A. C. Seijmonsbergen, D. Van Ree,
J. Hjort, R. S. Lawley, C. N. Williams, M. G. Anderson, P. Beier, P. Van Beukering, D. S.
Boyd, J. Jos′, J. Brilha, L. Carcavilla, K. M. Dahlin, J. C. Gill, J. E. Gordon, M. Gray, M.
Grundy, M. L. Hunter, J. J. Lawler, M. Monge-Ganuzas, K. R. Royse, I. Stewart, S. Record,
W. Turner, P. L. Zarnetske, R. Field, To advance sustainable stewardship, we must
document not only biodiversity but geodiversity. *Proc. Natl. Acad. Sci.* **116**, 16155–16158
(2019).

11.   F. Schrodt, G. Vernham, J. Bailey, R. Field, J. E. Gordon, M. Gray, J. Hjort, C. Hoorn, M.
L. Hunter, J. Larwood, A. Lausch, M. Monge-Ganuzas, S. Miller, D. van Ree, A. C.
Seijmonsbergen, P. L. Zarnetske, W. D. Kissling, The status and future of essential
geodiversity variables. *Philos. Trans. R. Soc. A* **382**, 1–18 (2024).

12.   P. Cienciala, A case for stronger integration of physical landscape processes in conservation
science and practice. *Conserv. Biol.* **38**, e14229 (2024).

13.   M. L. Hunter, V. Acuña, D. M. Bauer, K. P. Bell, A. J. K. Calhoun, M. R. Felipe-Lucia, J.
A. Fitzsimons, E. González, M. Kinnison, D. Lindenmayer, C. J. Lundquist, R. A. Medellin,
E. J. Nelson, P. Poschlod, Conserving small natural features with large ecological roles: A
synthetic overview. *Biol. Conserv.* **211**, 88–95 (2017).

14.   K. S. Nelson, E. K. Burchfield, Landscape complexity and US crop production. *Nat. Food*
2, 330–338 (2021).

15.   L. H. L. Loke, R. A. Chisholm, Measuring habitat complexity and spatial heterogeneity in
ecology. *Ecol. Lett.* **25**, 2269–2288 (2022).

16.   A. Stein, H. Kreft, Terminology and quantification of environmental heterogeneity in
species-richness research. *Biol. Rev.* **90**, 815–836 (2015).

17.   A. E. Hall, M. J. Kingsford, Habitat type and complexity drive fish assemblages in a tropical
seascape. *J. Fish Biol.* **99**, 1364–1379 (2021).

18.   K. E. Kovalenko, S. M. Thomaz, D. M. Warfe, Habitat complexity: Approaches and future
directions. *Hydrobiologia* **685**, 1–17 (2012).

19.   A. Stein, J. Beck, C. Meyer, E. Waldmann, P. Weigelt, H. Kreft, Differential effects of
environmental heterogeneity on global mammal species richness. *Glob. Ecol. Biogeogr.* **24**,
1072–1083 (2015).

20. J. Sola, J. N. Griffin, Resolving the context-dependency of local heterogeneity–diversity relationships across rocky reefs worldwide. *Proc. R. Soc. B Biol. Sci.* **292**, 20242723 (2025).

21. R. M. May, "Patterns of species abundance and diversity" in *M. L. Cody and J. M. Diamond, Editors. Ecology and Evolution of Communities* (Harvard University Press, Cambridge, Massachusetts, USA, 1975), pp. 81–120.

22. O. Arrhenius, Species and Area. *J. Ecol.* **9**, 95–99 (1921).

23. M. L. Rosenzweig, *Species Diversity in Space and Time* (Cambridge University Press, Cambridge, UK, 1995; https://cir.nii.ac.jp/crid/1361418519057201536).

24. R. H. MacArthur, E. O. Wilson, *The Theory of Island Biogeography* (Princeton University Press, 1967; http://www.jstor.org/stable/j.ctt19cc1t2).

25. A. Stein, K. Gerstner, H. Kreft, Environmental heterogeneity as a universal driver of species richness across taxa, biomes and spatial scales. *Ecol. Lett.* **17**, 866–880 (2014).

26. M. K. Borregaard, T. J. Matthews, R. J. Whittaker, The general dynamic model: towards a unified theory of island biogeography? *Glob. Ecol. Biogeogr.* **25**, 805–816 (2016).

27. O. Allouche, M. Kalyuzhny, G. Moreno-Rueda, M. Pizarro, R. Kadmon, Area–heterogeneity tradeoff and the diversity of ecological communities. *Proc. Natl. Acad. Sci.* **109**, 17495–17500 (2012).

28. R. Kadmon, O. Allouche, Integrating the Effects of Area, Isolation, and Habitat Heterogeneity on Species Diversity: A Unification of Island Biogeography and Niche Theory. *Am. Nat.* **170**, 443–454 (2007).

29. E. Ben-Hur, R. Kadmon, Heterogeneity–diversity relationships in sessile organisms: a unified framework. *Ecol. Lett.* **23**, 193–207 (2020).

30. E. Ben-Hur, R. Kadmon, An experimental test of the area–heterogeneity tradeoff. *Proc. Natl. Acad. Sci.* **117**, 4815–4822 (2020).

31. R. Chocron, C. H. Flather, R. Kadmon, Bird diversity and environmental heterogeneity in North America: a test of the area–heterogeneity trade-off. *Glob. Ecol. Biogeogr.* **24**, 1225–1235 (2015).

32. M. Gray, *Geodiversity: Valuing and Conserving Abiotic Nature* (Wiley-Blackwell, 2013).

33. G. Vernham, J. J. Bailey, J. M. Chase, J. Hjort, R. Field, F. Schrodt, Understanding trait diversity: the role of geodiversity. *Trends Ecol. Evol.*, doi: 10.1016/j.tree.2023.02.010 (2023).

34. G. Vernham, J. J. Bailey, R. Field, F. Schrodt, What is the Relationship Between Plant Trait Diversity and Geodiversity? A Plot-Based, Pan-European Analysis. *Glob. Ecol. Biogeogr.* **33**, e13904 (2024).

35.   P. L. Zarnetske, B. Baiser, A. Strecker, S. Record, J. Belmaker, M.-N. Tuanmu, The
Interplay Between Landscape Structure and Biotic Interactions. *Curr. Landsc. Ecol. Rep.* **2**,
12–29 (2017).

36.   D. Torres-Pulliza, M. A. Dornelas, O. Pizarro, M. Bewley, S. A. Blowes, N. Boutros, V.
Brambilla, T. J. Chase, G. Frank, A. Friedman, M. O. Hoogenboom, S. Williams, K. J. A.
Zawada, J. S. Madin, A geometric basis for surface habitat complexity and biodiversity.
Nat. Ecol. Evol. **4**, 1495–1501 (2020).

37.   G. Arseniou, D. W. MacFarlane, Fractal dimension of tree crowns explains species
functional-trait responses to urban environments at different scales. *Ecol. Appl.* **31**, e02297
(2021).

38.   Copernicus Climate Change Service, Land cover classification gridded maps from 1992 to
present derived from satellite observation (2019).

39.   Center for International Earth Science Information Network (CIESIN), Gridded Population
of the World, Version 4 (GPWv4): Population Count, Revision 11. (2018).

40.   D. M. Warfe, L. A. Barmuta, S. Wotherspoon, Quantifying Habitat Structure: Surface
Convolution and Living Space for Species in Complex Environments. *Oikos* **117**, 1764–
1773 (2008).

41.   B. B. Mandelbrot, Self-Affine Fractals and Fractal Dimension. *Phys. Scr.* **32**, 257 (1985).

42.   B. T. Trew, I. M. D. Maclean, Vulnerability of global biodiversity hotspots to climate
change. *Glob. Ecol. Biogeogr.* **30**, 768–783 (2021).

43.   S. Z. Dobrowski, A climatic basis for microrefugia: the influence of terrain on climate. *Glob.*
Change Biol. **17**, 1022–1035 (2011).

44.   T. D. Meehan, B. P. Werling, D. A. Landis, C. Gratton, Agricultural landscape
simplification and insecticide use in the Midwestern United States. *Proc. Natl. Acad. Sci.*
108, 11500–11505 (2011).

45.   S. J. Norder, R. F. de Lima, L. de Nascimento, J. Y. Lim, J. M. Fernández-Palacios, M. M.
Romeiras, R. B. Elias, F. J. Cabezas, L. Catarino, L. M. P. Ceríaco, A. Castilla-Beltrán, R.
Gabriel, M. M. de Sequeira, K. F. Rijsdijk, S. Nogué, W. D. Kissling, E. E. van Loon, M.
Hall, M. Matos, P. A. V. Borges, Global change in microcosms: Environmental and societal
predictors of land cover change on the Atlantic Ocean Islands. *Anthropocene* **30**, 100242
(2020).

46.   S. J. Norder, A. C. Seijmonsbergen, S. D. D. V. Rughooputh, E. E. van Loon, V. Tatayah,
A. T. Kamminga, K. F. Rijsdijk, Assessing temporal couplings in social–ecological island
systems: historical deforestation and soil loss on Mauritius (Indian Ocean). *Ecol. Soc.* **22**
(2017).

47.   GEBCO Compilation Group, GEBCO 2023 Grid (2023).

48. N. J. Frost, M. T. Burrows, M. P. Johnson, M. E. Hanley, S. J. Hawkins, Measuring surface
complexity in ecological studies. *Limnol. Oceanogr. Methods* **3**, 203–210 (2005).

49. S. C. Pal, A. Saha, I. Chowdhuri, D. Ruidas, R. Chakrabortty, P. Roy, M. Shit, Earthquake
hotspot and coldspot: Where, why and how? *Geosystems Geoenvironment* **2**, 100130
(2023).

50. L. A. Levin, R. J. Etter, M. A. Rex, A. J. Gooday, C. R. Smith, J. Pineda, C. T. Stuart, R. R.
Hessler, D. Pawson, Environmental Influences on Regional Deep-Sea Species Diversity.
Annu. Rev. Ecol. Syst. **32**, 51–93 (2001).

51. B. Smith, D. Sandwell, Accuracy and resolution of shuttle radar topography mission data.
Geophys. Res. Lett. **30** (2003).

52. M. W. Palmer, The Coexistence of Species in Fractal Landscapes. *Am. Nat.* **139**, 375–397
(1992).

53. V. E. Kostylev, J. Erlandsson, M. Y. Ming, G. A. Williams, The relative importance of
habitat complexity and surface area in assessing biodiversity: Fractal application on rocky
shores. *Ecol. Complex.* **2**, 272–286 (2005).

54. P. T. Harris, M. Macmillan-Lawler, J. Rupp, E. K. Baker, Geomorphology of the oceans.
Mar. Geol. **352**, 4–24 (2014).

55. M. Saccò, N. E. White, C. Harrod, G. Salazar, P. Aguilar, C. F. Cubillos, K. Meredith, B.
K. Baxter, A. Oren, E. Anufriieva, N. Shadrin, Y. Marambio-Alfaro, V. Bravo-Naranjo, M.
E. Allentoft, Salt to conserve: a review on the ecology and preservation of hypersaline
ecosystems. *Biol. Rev.* **96**, 2828–2850 (2021).

56. P. Jara-Arancio, M. T. K. Arroyo, P. C. Guerrero, L. F. Hinojosa, G. Arancio, M. A.
Méndez, Phylogenetic perspectives on biome shifts in eucocoryne (Alliaceae) in relation to
climatic niche evolution in western South America. *J. Biogeogr.* **41**, 328–338 (2014).

57. Kingsford, R. T., Hollibaugh, J. T., Harper, R., Keith, D. A., "F2.6 Permanent salt and soda
lakes." in *Keith, D.A., Ferrer-Paris, J.R., Nicholson, E. et al. (2022). A Function-Based*
Typology for Earth's Ecosystems – Appendix S4. The IUCN Global Ecosystem Typology
v2.1: Descriptive Profiles for Biomes and Ecosystem Functional Groups. (2022)*Nature*.

58. C. Zavala, M. Arcuri, M. Di Meglio, A. Zorzano, G. Otharán, A. Irastorza, A. Torresi,
Deltas: a new classification expanding Bates's concepts. *J. Palaeogeogr.* **10**, 23 (2021).

59. J. A. Tarduno, D. B. Brinkman, P. R. Renne, R. D. Cottrell, H. Scher, P. Castillo, Evidence
for Extreme Climatic Warmth from Late Cretaceous Arctic Vertebrates. *Science* **282**, 2241–
2243 (1998).

60. J. P. Klages, U. Salzmann, T. Bickert, C.-D. Hillenbrand, K. Gohl, G. Kuhn, S. M. Bohaty,
J. Titschack, J. Müller, T. Frederichs, T. Bauersachs, W. Ehrmann, T. van de Flierdt, P. S.
Pereira, R. D. Larter, G. Lohmann, I. Niezgodzki, G. Uenzelmann-Neben, M. Zundel, C.

Spiegel, C. Mark, D. Chew, J. E. Francis, G. Nehrke, F. Schwarz, J. A. Smith, T. Freudenthal, O. Esper, H. Pälike, T. A. Ronge, R. Dziadek, Temperate rainforests near the South Pole during peak Cretaceous warmth. *Nature* **580**, 81–86 (2020).

61. J. Hjort, J. E. Gordon, M. Gray, M. L. Hunter JR., Why geodiversity matters in valuing nature's stage. *Conserv. Biol.* **29**, 630–639 (2015).

62. H. Tukiainen, J. Alahuhta, R. Field, T. Ala-Hulkko, R. Lampinen, J. Hjort, Spatial relationship between biodiversity and geodiversity across a gradient of land-use intensity in high-latitude landscapes. *Landsc. Ecol.* **32**, 1049–1063 (2017).

63. J. A. Dijkstra, L. G. Harris, K. Mello, A. Litterer, C. Wells, C. Ware, Invasive seaweeds transform habitat structure and increase biodiversity of associated species. *J. Ecol.* **105**, 1668–1678 (2017).

64. Y. Ren, Y. Lü, J. Hu, L. Yin, Geodiversity underpins biodiversity but the relations can be complex: Implications from two biodiversity proxies. *Glob. Ecol. Conserv.* **31**, e01830 (2021).

65. J. Igea, A. J. Tanentzap, Global topographic uplift has elevated speciation in mammals and birds over the last 3 million years. *Nat. Ecol. Evol.* **5**, 1530–1535 (2021).

66. L. Valente, A. B. Phillimore, M. Melo, B. H. Warren, S. M. Clegg, K. Havenstein, R. Tiedemann, J. C. Illera, C. Thébaud, T. Aschenbach, R. S. Etienne, A simple dynamic model explains the diversity of island birds worldwide. *Nature* **579**, 92–96 (2020).

67. R. J. Whittaker, K. A. Triantis, R. J. Ladle, A General Dynamic Theory of Oceanic Island Biogeography. *J. Biogeogr.* **35**, 977–994 (2008).

68. S. Huang, M. J. M. Meijers, A. Eyres, A. Mulch, S. A. Fritz, Unravelling the history of biodiversity in mountain ranges through integrating geology and biogeography. *J. Biogeogr.* **46**, 1777–1791 (2019).

69. J. Fjeldså, R. C. K. Bowie, C. Rahbek, The role of mountain ranges in the diversification of birds. *Annu. Rev. Ecol. Evol. Syst.* **43**, 249–265 (2012).

70. A. C. Seijmonsbergen, J. Guldenaar, K. F. Rijsdijk, Exploring Hawaiian long-term insular geodiversity dynamics. *Landf. Anal.* **35**, 31–43 (2017).

71. M. Tokeshi, S. Arakaki, Habitat complexity in aquatic systems: Fractals and beyond. *Hydrobiologia* **685**, 27–47 (2012).

72. M. P. Johnson, N. J. Frost, M. W. J. Mosley, M. F. Roberts, S. J. Hawkins, The area-independent effects of habitat complexity on biodiversity vary between regions. *Ecol. Lett.* **6**, 126–132 (2003).

73. M. S. Bass, M. Finer, C. N. Jenkins, H. Kreft, D. F. Cisneros-Heredia, S. F. McCracken, N. C. A. Pitman, P. H. English, K. Swing, G. Villa, A. Di Fiore, C. C. Voigt, T. H. Kunz,

Global Conservation Significance of Ecuador's Yasuní National Park. *PLoS ONE* **5**, e8767
(2010).

74.   Science Panel for the Amazon, *Amazon Assessment Report 2021* (United Nations
Sustainable Development Solutions Network, New York, USA, 2021;
https://www.theamazonwewant.org/spa-reports/. DOI: 10.55161/RWSX6527).

75.   M. Claussen, V. Gayler, The Greening of the Sahara during the Mid-Holocene: Results of
an Interactive Atmosphere-Biome Model. *Glob. Ecol. Biogeogr. Lett.* **6**, 369–377 (1997).

76.   S. Kröpelin, D. Verschuren, A.-M. Lézine, H. Eggermont, C. Cocquyt, P. Francus, J.-P.
Cazet, M. Fagot, B. Rumes, J. M. Russell, F. Darius, D. J. Conley, M. Schuster, H. von
Suchodoletz, D. R. Engstrom, Climate-Driven Ecosystem Succession in the Sahara: The
Past 6000 Years. *Science* **320**, 765–768 (2008).

77.   M. V. Lomolino, Ecology's most general, yet protean pattern: the species-area relationship.
J. Biogeogr. **27**, 17–26 (2000).

78.   T. J. Matthews, F. Guilhaumon, K. A. Triantis, M. K. Borregaard, R. J. Whittaker, On the
form of species–area relationships in habitat islands and true islands. *Glob. Ecol. Biogeogr.*
25, 847–858 (2016).

79.   T. E. Martin, Species-Area Slopes and Coefficients: A Caution on Their Interpretation. *Am.*
Nat. **118**, 823–837 (1981).

80.   R. H. MacArthur, E. O. Wilson, An Equilibrium Theory of Insular Zoogeography. *Evolution*
17, 373–387 (1963).

81.   T. Shi, X. Li, L. Xin, X. Xu, The spatial distribution of farmland abandonment and its
influential factors at the township level: A case study in the mountainous area of China.
Land Use Policy **70**, 510–520 (2018).

82.   D. G. Green, N. Klomp, G. Rimmington, S. Sadedin, *Complexity in Landscape Ecology*
(Springer, The Netherlands, 2006)vol. 4 of *Landscape Series*.

83.   L. Fahrig, J. Baudry, L. Brotons, F. G. Burel, T. O. Crist, R. J. Fuller, C. Sirami, G. M.
Siriwardena, J.-L. Martin, Functional landscape heterogeneity and animal biodiversity in
agricultural landscapes. *Ecol. Lett.* **14**, 101–112 (2011).

84.   T. Wang, I. Belle, U. Hassler, Modelling of Singapore's topographic transformation based
on DEMs. *Geomorphology* **231**, 367–375 (2015).

85.   E. C. Ellis, N. Gauthier, K. Klein Goldewijk, R. Bliege Bird, N. Boivin, S. Díaz, D. Q.
Fuller, J. L. Gill, J. O. Kaplan, N. Kingston, H. Locke, C. N. H. McMichael, D. Ranco, T.
C. Rick, M. R. Shaw, L. Stephens, J.-C. Svenning, J. E. M. Watson, People have shaped
most of terrestrial nature for at least 12,000 years. *Proc. Natl. Acad. Sci.* **118**, e2023483118
(2021).

86.   A. P. Sullivan, D. W. Bird, G. H. Perry, Human behaviour as a long-term ecological driver
of non-human evolution. *Nat. Ecol. Evol.* **1**, 1–11 (2017).

87.   B. M. Flores, C. Levis, Human-food feedback in tropical forests. *Science* **372**, 1146–1147
(2021).

88.   E. M. N. Polman, A. C. Seijmonsbergen, H. Versteegh, W. D. Kissling, Global geodiversity
components are not equally represented in UNESCO Global Geoparks. *Philos. Trans. R.*
Soc. A **382**, 1–18 (2024).

89.   C. Badgley, Tectonics, topography, and mammalian diversity. *Ecography* **33**, 220–231
(2010).

90.   M. J. Steinbauer, R. Field, J.-A. Grytnes, P. Trigas, C. Ah-Peng, F. Attorre, H. J. B. Birks,
P. A. V. Borges, P. Cardoso, C.-H. Chou, M. De Sanctis, M. M. de Sequeira, M. C. Duarte,
R. B. Elias, J. M. Fernández-Palacios, R. Gabriel, R. E. Gereau, R. G. Gillespie, J. Greimler,
D. E. V. Harter, T.-J. Huang, S. D. H. Irl, D. Jeanmonod, A. Jentsch, A. S. Jump, C. Kueffer,
S. Nogué, R. Otto, J. Price, M. M. Romeiras, D. Strasberg, T. Stuessy, J.-C. Svenning, O.
R. Vetaas, C. Beierkuhnlein, Topography-driven isolation, speciation and a global increase
of endemism with elevation. *Glob. Ecol. Biogeogr.* **25**, 1097–1107 (2016).

91.   C. Rahbek, M. K. Borregaard, R. K. Colwell, B. Dalsgaard, B. G. Holt, N. Morueta-Holme,
D. Nogues-Bravo, R. J. Whittaker, J. Fjeldså, Humboldt's enigma: What causes global
patterns of mountain biodiversity? *Science* **365**, 1108–1113 (2019).

92.   C. Körner, Why are there global gradients in species richness? mountains might hold the
answer. *Trends Ecol. Evol.* **15**, 513–514 (2000).

93.   R. G. Davies, C. D. L. Orme, D. Storch, V. A. Olson, G. H. Thomas, S. G. Ross, T.-S. Ding,
P. C. Rasmussen, P. M. Bennett, I. P. F. Owens, T. M. Blackburn, K. J. Gaston, Topography,
energy and the global distribution of bird species richness. *Proc. R. Soc. B Biol. Sci.* **274**,
1189–1197 (2007).

94.   R Core Team, R: A language and environment for statistical computing., R Foundation for
Statistical Computing (2022); https://www.R-project.org/.

95.   QGIS.org, QGIS Geographic Information System., QGIS Association (2023).

96.   J. Madin, N. Schiettekatte, habtools: Tools and Metrics for 3D Surfaces and Objects, (2024).

97.   Microsoft Corporation, S. Weston, foreach: Provides Foreach Looping Construct, (2022).

98.   Microsoft Corporation, S. Weston, doParallel: Foreach Parallel Adaptor for the "parallel"
Package, (2022).

99.   D. A. Keith, R. J. Williams, "T4.3 Hummock savannas." in *Keith, D.A., Ferrer-Paris, J.R.,*
Nicholson, E. et al. (2022). A Function-Based Typology for Earth's Ecosystems – Appendix

S4. The IUCN Global Ecosystem Typology v2.1: Descriptive Profiles for Biomes and Ecosystem Functional Groups (2022)*Nature.*

100. D. A. Keith, D. Faber-Langendoen, J. Franklin, "T4.4 Temperate woodlands." in *Keith, D.A., Ferrer-Paris, J.R., Nicholson, E. et al. (2022). A Function-Based Typology for Earth's Ecosystems – Appendix S4. The IUCN Global Ecosystem Typology v2.1: Descriptive Profiles for Biomes and Ecosystem Functional Groups* (2022)*Nature.*

101. C. E. R. Lehmann, A. Etter, K. R. Young, D. A. Keith, "T4.2 Pyric tussock savannas." in *Keith, D.A., Ferrer-Paris, J.R., Nicholson, E. et al. (2022). A Function-Based Typology for Earth's Ecosystems – Appendix S4. The IUCN Global Ecosystem Typology v2.1: Descriptive Profiles for Biomes and Ecosystem Functional Groups.* (2022)*Nature.*

102. C. E. R. Lehmann, D. A. Keith, "T4.1 Trophic savannas" in *Keith, D.A., Ferrer-Paris, J.R., Nicholson, E. et al. (2022). A Function-Based Typology for Earth's Ecosystems – Appendix S4. The IUCN Global Ecosystem Typology v2.1: Descriptive Profiles for Biomes and Ecosystem Functional Groups.* (2022)*Nature.*

103. D. A. Keith, F. Essl, D. J. Gibson, "T4.5 Temperate subhumid grasslands." in *Keith, D.A., Ferrer-Paris, J.R., Nicholson, E. et al. (2022). A Function-Based Typology for Earth's Ecosystems – Appendix S4. The IUCN Global Ecosystem Typology v2.1: Descriptive Profiles for Biomes and Ecosystem Functional Groups* (2022)*Nature.*

104. D. A. Keith, J. R. Ferrer-Paris, E. Nicholson, R. T. Kingsford, Indicative distribution maps for Ecosystem Functional Groups - Level 3 of IUCN Global Ecosystem Typology (2023). https://doi.org/10.5281/zenodo.10081251.

105. R. J. Hijmans, raster: Geographic data analysis and modeling, (2020); https://cran.r-project.org/package=raster.

106. R. J. Hijmans, terra: Spatial Data Analysis, (2023).

107. S. N. Wood, *Generalized Additive Models* (Chapman and Hall/CRC, New York, 2017).

108. Bürkner, P., brms: An R Package for Bayesian Multilevel Models Using Stan. *J. Stat. Softw.* **80**, 1–28 (2017).

109. T. J. Matthews, J. P. Wayman, R. J. Whittaker, P. Cardoso, J. P. Hume, F. Sayol, K. Proios, T. E. Martin, B. Baiser, P. A. V. Borges, Y. Kubota, L. dos Anjos, J. A. Tobias, F. C. Soares, X. Si, P. Ding, C. D. Mendenhall, Y. C. K. Sin, F. E. Rheindt, K. A. Triantis, F. Guilhaumon, D. M. Watson, L. Brotons, C. Battisti, O. Chu, F. Rigal, A global analysis of avian island diversity–area relationships in the Anthropocene. *Ecol. Lett.*, doi: 10.1111/ele.14203 (2023).

110. T. M. Blackburn, P. Cassey, R. P. Duncan, K. L. Evans, K. J. Gaston, Avian extinction and mammalian introductions on oceanic islands. *Science* **305**, 1955–1958 (2004).

111. J. A. Tobias, C. Sheard, A. L. Pigot, A. J. M. Devenish, J. Yang, F. Sayol, M. H. C. Neate-Clegg, N. Alioravainen, T. L. Weeks, R. A. Barber, P. A. Walkden, H. E. A. MacGregor, S. E. I. Jones, C. Vincent, A. G. Phillips, N. M. Marples, F. A. Montaño-Centellas, V. Leandro-Silva, S. Claramunt, B. Darski, B. G. Freeman, T. P. Bregman, C. R. Cooney, E. C. Hughes, E. J. R. Capp, Z. K. Varley, N. R. Friedman, H. Korntheuer, A. Corrales-Vargas, C. H. Trisos, B. C. Weeks, D. M. Hanz, T. Töpfer, G. A. Bravo, V. Remeš, L. Nowak, L. S. Carneiro, A. J. Moncada R., B. Matysioková, D. T. Baldassarre, A. Martínez-Salinas, J. D. Wolfe, P. M. Chapman, B. G. Daly, M. C. Sorensen, A. Neu, M. A. Ford, R. J. Mayhew, L. Fabio Silveira, D. J. Kelly, N. N. D. Annorbah, H. S. Pollock, A. M. Grabowska-Zhang, J. P. McEntee, J. Carlos T. Gonzalez, C. G. Meneses, M. C. Muñoz, L. L. Powell, G. A. Jamie, T. J. Matthews, O. Johnson, G. R. R. Brito, K. Zyskowski, R. Crates, M. G. Harvey, M. Jurado Zevallos, P. A. Hosner, T. Bradfer-Lawrence, J. M. Maley, F. G. Stiles, H. S. Lima, K. L. Provost, M. Chibesa, M. Mashao, J. T. Howard, E. Mlamba, M. A. H. Chua, B. Li, M. I. Gómez, N. C. García, M. Päckert, J. Fuchs, J. R. Ali, E. P. Derryberry, M. L. Carlson, R. C. Urriza, K. E. Brzeski, D. M. Prawiradilaga, M. J. Rayner, E. T. Miller, R. C. K. Bowie, R. M. Lafontaine, R. P. Scofield, Y. Lou, L. Somarathna, D. Lepage, M. Illif, E. L. Neuschulz, M. Templin, D. M. Dehling, J. C. Cooper, O. S. G. Pauwels, K. Analuddin, J. Fjeldså, N. Seddon, P. R. Sweet, F. A. J. DeClerck, L. N. Naka, J. D. Brawn, A. Aleixo, K. Böhning-Gaese, C. Rahbek, S. A. Fritz, G. H. Thomas, M. Schleuning, AVONET: morphological, ecological and geographical data for all birds. *Ecol. Lett.* **25**, 581–597 (2022).

112. C. Magneville, N. Loiseau, C. Albouy, N. Casajus, T. Claverie, A. Escalas, F. Leprieur, E. Maire, D. Mouillot, S. Villéger, mFD: an R package to compute and illustrate the multiple facets of functional diversity. *Ecography* **2022**, 1–15 (2022).

113. R. Sayre, S. Noble, S. Hamann, R. Smith, D. Wright, S. Breyer, K. Butler, K. Van Graafeiland, C. Frye, D. Karagulle, D. Hopkins, D. Stephens, K. Kelly, Z. Basher, D. Burton, J. Cress, K. Atkins, D. P. Van Sistine, B. Friesen, R. Allee, T. Allen, P. Aniello, I. Asaad, M. J. Costello, K. Goodin, P. Harris, M. Kavanaugh, H. Lillis, E. Manca, F. Muller-Karger, B. Nyberg, R. Parsons, J. Saarinen, J. Steiner, A. Reed, A new 30 meter resolution global shoreline vector and associated global islands database for the development of standardized ecological coastal units. *J. Oper. Oceanogr.* **12**, S47–S56 (2019).

114. R. D. Morey, J. N. Rouder, BayesFactor: Computation of Bayes Factors for Common Designs, (2024).

115. C. Brockmann, M. Boettcher, G. Kirches, "Tutorial: CDR Land Cover (brokered from CCI Land Cover)" (Copernicus Climate Change Service, 2018).

Acknowledgments:

Funding:

Marie Sklodowska-Curie-European Postdoctoral Fellowship through Horizon Europe, ReefCoRe 101204139 (JC).

National Science Foundation – Natural Environment Research Council Biological Oceanography Grant 1948946 (MD, JSM, ESPM).

ERC Consolidator Grant through the European Union, CoralINT 101044975 (MD).

Author Contributions:

Conceptualization: JC, NS, EMPM, JSM, MD

Methodology: JC, NS, ACS, KFR, JSM, MD

Investigation: JC

Visualization: JC, JSM, MD

Funding acquisition: JC, JSM, EMPM, MD

Project administration: MD, JSM

Supervision: MD

Writing – original draft: JC, MD

Writing – review & editing: JC, NS, EMPM, ACS, KFR, JSM, MD

Competing interests: The authors declare that there are no conflicts of interest associated with this piece of work.

Data and materials availability: All the R scripts used in developing these maps and implementing all analyses discussed throughout this manuscript have been uploaded to a public repository on GitHub (https://github.com/CantJ/Global-Geometric-Complexity). This code repository and the global maps of structural complexity presented in this manuscript have also been archived open access on Zenodo (https://doi.org/10.5281/zenodo.18153610).

Supplementary Materials:

Materials and Methods [2028 words]

Supplementary Text

Figs. S1 to S6

Tables S1 to S3

References (94 - 115)

A.

i. Amazon Rainforest

ii. Mariana Trench

iii. Sahara Desert

Longitude

Height Range $\log_{10}(\Delta H)$

B.

Rugosity $\log_{10}(R)$

C.

Fractal Dimension (D)

i. Amazon Rainforest

ii. Mariana Trench

iii. Sahara Desert

Longitude

Fig. 1. Global patterns in geometric structural complexity. High resolution maps (1.87×1.87 km) depicting variation in the structural complexity measures of (A) Height Range (ΔH), (B) Rugosity (R), and (C) Fractal Dimension (D), across all terrestrial and marine environments. Global maps are displayed in a global Mollweide equal area projection and comprise ~146 million estimates of ΔH, R, and D, with white pixels reflecting missing or omitted entries. Comparing measures of fractal dimension and rugosity can offer differing insights into the complexity characteristics of natural environments, with the close-up panels highlighting the fractal dimension (*left*) and rugosity (*right*) of the (i) Amazonian Basin, (ii) Mariana Trench, and (iii) Sahara Desert. Each panel shows a representative 935×935 km section of the selected locations and is displayed in a WGS84 projection.

A.

B.

Fig. 2. The relative structural characteristics of differing environments delineates the differing complexity regimes underlying ecosystems worldwide. (A) Ridgeline plots illustrating the distribution of rugosity (R) and fractal dimension (D) values associated with ecosystem types across different IUCN biome classifications (see Figs. S4 & S5 for a within biome breakdown of rugosities and fractal dimension across ecosystem types). Global distributions of different biomes and ecosystem types obtained from the IUCN function-based ecosystem typology (*5*). Fill colour used to delineate the realm types associated with each biome class. **(B)** Mean fractal dimension and rugosity estimates derived for differing ecosystem types. Point colouring used to cluster ecosystem types based on their associated realms, with panel colouring used to highlight regions of comparatively high and low rugosity and/or fractal dimension. Red inset text highlights the corresponding pattern in height range (*ΔH*) across this parameter space, with representative ecosystem types shown for reference. Fit line shows modelled relationship between mean ecosystem rugosity and fractal dimension estimates (r^2: 0.26 [95% CI. 0.13, 0.38]) with error displayed as 95% CI.

A.

$r^2 = 0.362$ [95% CI: 0.361, 0.364]

B.

$r^2 = 0.377$ [0.375, 0.379]

C.

$r^2 = 0.098$ [0.096, 0.099]

Geodiversity

Fig. 3. Comparing the geometry and structural richness of natural environments allows for disentangling the mechanisms through which structural complexity mediates the configuration of global biodiversity. The alignment of the structural complexity measures of **(A)** Height Range (*ΔH*), **(B)** Rugosity (*R*), and **(C)** Fractal Dimension (*D*) with estimates of geodiversity (geology, geomorphology and hydrology combined). Geodiversity estimates were extracted from a high resolution (1.87 × 1.87 km) global map depicting patterns in the richness of structural features comprising an environment (*88*). The solid line across each plot represents the predicted relationship between each geodiversity and each complexity measure, with the thickness of the line corresponding with the 95% confidence intervals.

Fig. 4. Global patterns in structural complexity influence how humans interact with and use natural environments worldwide. (A) Ridgeline plots illustrating the distribution of rugosity (R) and fractal dimension (D) estimates associated with different land cover classifications. Land cover categories and their spatial distributions obtained from the Copernicus Climate Change Service (C3S) land cover survey (*38*). **(B)** The alignment of the structural complexity measures of **(i)** Height Range (ΔH), **(ii)** Rugosity (R), and **(iii)** Fractal Dimension (D) with patterns in human population density. Human population density patterns were sourced from the Gridded Population of the World (*39*), with each plot showing the predicted relationships (including 95% CI) derived from two-part hurdle models separately exploring patterns in the probability of greater than zero population densities (red) and recorded population count densities (blue). Population density estimates were transformed using a power transformation ($y^{0.05}$).

Box 1. *Case-study: Disentangling drivers of island biodiversity*

To demonstrate how metrics of structural complexity can help us understand the intrinsic association between biodiversity and abiotic habitat structure, we separately modelled the relationships between the complexity variables of height range (ΔH), rugosity (R), and fractal dimension (D), and measures of native (*i.e.*, non-introduced) island bird species and functional richness. Consistent with established theory, island area has a positive effect on both island bird species richness (Fig. 5A) and functional richness (Fig. S6A). Underlying this effect of island area, however, we evidence how the effects of structural complexity on species and functional richness patterns contribute to the multifaceted and heterogeneous relationship between species diversity and area. Overall, we illustrate that whilst increases in island rugosity and height range have little effect on bird species richness, they have a positive effect on functional diversity (Fig. 5B). Alternatively, fractal dimension has an overall negative effect on bird species richness but little effect on bird functional richness (Fig. 5B). Yet, these overall effects mask important non-linear relationships between structural complexity and species diversity. Large elevational gradients (*i.e.*, high R and/or ΔH), such as those associated with mountainous regions, generate geographical barriers and support high habitat and climatic heterogeneity, thus promoting high rates of speciation and diversification (*6*, *69*, *89*). Areas of high elevational complexity also support high endemism due to the specialisation required for inhabiting their associated ecosystems (*90*, *91*). Hence, at lower levels of rugosity and height range, increases in these complexity measures correlate with slight increases in island bird species richness (Fig. 5C[i & ii]) and large increases in island bird functional richness (Fig. S6B & C). However, this positive effect of rugosity and height range tapers off as complexity increases. Alongside a potential reduction in available niche space, the specialisation needed to initially access and colonise island systems coupled with the specialisation needed to tolerate the climate regimes associated with areas of high elevational and topographical complexity (*92*) may explain why, beyond a particular threshold, increases in ΔH and R diminish the effect of area on bird diversity. Meanwhile, at lower levels, changes in fractal dimension have little effect on either island bird species or functional richness (Fig. 5C[iii] & S6D). However, at higher levels, increases in island fractal dimension have a negative effect on island bird species and functional diversity. Higher fractal dimension across natural landscapes is attributed with promoting speciation through diversification by increasing habitat diversity and enabling more effective resource partitioning (*52*, *71*, *72*). However, the increased habitat diversity associated with higher fractal dimension likely reduces the availability of realised niche space; eliciting an increase in local extirpation rates (*27*). Our assessment here focused on avian fauna whose distributions are more defined by coarse-scale topographic features than fine-scale patterns in the shape of surface environments (*93*). However, the patterns we highlight here testament the necessity for continued exploration into the relationship between structural complexity and species diversity across taxa, as the necessary data becomes available.

Fig. 5. Habitat structural complexity influences how changes in available space influence biotic configurations. (A) The conditional effect of Island area (km^2) on island bird species richness when accounting for the nested effect of archipelago and the fixed effects of island latitude, distance to nearest mainland, and human population density ($r^2 = 0.844$ [95% CI: 0.829, 0.857]). **(B)** Posterior distributions of the first polynomial slope coefficients extracted from models evaluating the effects of island area, rugosity, height range, and fractal dimension, on island bird species and functional richness. **(C)** The conditional effects of **(i)** rugosity (R), **(ii)** height range (ΔH), and **(iii)** fractal dimension (D) on island bird species richness (r^2: $R = 0.795$ [0.775, 0.811]; $\Delta H = 0.794$ [0.775, 0.810]; $D = 0.795$ [0.776, 0.812]). All models account for the nested effects of archipelago and the fixed effect of island area, island latitude, distance to nearest mainland, and human population density. Error displayed as 95% CI across all panels.