CLCNet: a contrastive learning and chromosome-aware network for genomic prediction in plants

Jiangwei Huang[1,4], Zhihan Yang[1,2], Mou Yin[1,2], Chao Li[1,4], Jinmin Li[1,4], Yu Wang[1,4], Lu Huang[1,4], Fei He[1,2], Chengzhi Liang[1,2], Miaomiao Li[1,3], Rongcheng Han[1,4,*], and Yuqiang Jiang[1,4,*]

[1]*State Key Laboratory of Molecular Developmental Biology, Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, Beijing, China*

[2]*State Key Laboratory of Plant Genomics, Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, Beijing, China*

[3]*State Key Laboratory of Plant Cell and Chromosome Engineering, Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, Beijing, China*

[4]*University of Chinese Academy of Sciences, Beijing, China*

* Correspondence (Tel: +86-10-64806367; Email: hanrch@genetics.ac.cn (R.H.); Tel: +86-1064806362; Email: yqjiang@genetics.ac.cn (Y.J.))

Keywords: Genomic Prediction, Deep Learning, Contrastive Learning, Complex Traits

Abstract

Genomic selection (GS) is a sophisticated breeding technology that utilizes genomic markers in conjunction with phenotypic data to predict breeding values and phenotypes of candidate populations. The foundation of GS is the construction of precise genomic prediction (GP) models. Traditional GP models, which are typically linear in nature, encounter difficulties in capturing non-linear relationships within genetic data, thereby constraining their capacity to adequately characterize complex genetic architectures. To address this limitation, we have developed a novel deep learning framework, namely contrastive learning and chromosome-aware network (CLCNet), which has been specifically tailored for plant GP. The architecture of CLCNet comprises two key modules: a contrastive learning module designed to capture phenotypic variations, and a chromosome-aware module that accounts for the effects of linkage, as well as local and global epistasis. The performance of CLCNet was evaluated across eight datasets representing seven plant species: maize (*Zea mays*), rice (*Oryza sativa*), cotton (*Gossypium hirsutum*), millet (*Setaria italica*), chickpea (*Cicer arietinum*), rapeseed (*Brassica napus*), and soybean (*Glycine max*). In comparative assessments against three classical models (rrBLUP, Bayesian Ridge, Bayesian Lasso), two machine learning approaches (LightGBM, SVR), and two other deep learning models (DNNGP, DeepGS), CLCNet exhibited superior performance, achieving higher Pearson correlation coefficients (PCC) and lower root mean squared errors (RMSE). Due to its superior prediction accuracy, generalization capability and scalability in high-dimensional datasets, CLCNet is expected to be a powerful tool for future GS applications in the field of plant breeding.

Introduction

As an advanced plant and animal breeding technology, genomic selection (GS) utilizes genome-wide markers and phenotypic data to predict genomic estimated breeding values (GEBVs) (Meuwissen et al., 2001). In contrast to the marker-assisted selection (MAS) method, GS captures the collective effects of numerous loci, including those with minor contributions, enabling earlier and more efficient selection (Crossa et al., 2017). A prevalent issue in GS is the discrepancy between the large number of markers and the relatively small number of phenotypic observations. To address this, various models and algorithms, such as GBLUP, Bayesian approaches, and machine learning techniques, have been developed to enhance prediction accuracy (Habier et al., 2007; Maenhout et al., 2007; Crossa et al., 2010; Heslot et al., 2012; Wang et al., 2018). Nevertheless, the efficacy of these methodologies is often impeded by their inability to adequately capture nonlinear interactions and manage noise in genomic data.

Deep learning (DL) has emerged as a powerful alternative, capable of modelling complex, high-dimensional datasets through multi-layer neural networks for genomics (Liu et al., 2020,). The efficacy of DL in capturing non-linear genotype-phenotype relationships without the necessity for predefined assumptions renders it a particularly advantageous approach for complex traits (Abdollahi-Arpanahi et al., 2020; Wang et al., 2020). Furthermore, DL supports a range of modelling paradigms, encompassing single-task, multi-task, multi-modal, and transfer learning (Eraslan et al., 2019). These paradigms enhance performance by focusing on specific phenotypes, leveraging shared trait information, integrating multi-omics data, and utilizing pre-trained models for data-scarce scenarios.

DL-based models have demonstrated notable success in genomic prediction (GP).

For instance, DeepGS, which employed convolutional layers, outperformed RR-BLUP

in wheat predictions (Ma et al., 2018). Furthermore, DLGWAS has been shown to

enhance soybean phenotypic predictions through the utilization of residual and parallel

convolutional layers (Liu et al., 2019). Addressing the issues of denoising and

dimensionality, DNNGP combined PCA with DL for robust predictions across wheat,

maize, and tomato (Wang et al., 2023). SoyDNGP further innovated by transforming

genomic matrices into RGB-like channels for soybean GS (Gao et al., 2023).

Multi-modal DL models have also advanced GS by integrating multi-omics data. For

instance, DEM combined genomic, transcriptomic, and epigenomic data using self-

attention mechanisms, thereby improving predictions for various species (Ren et al.,

2024). DeepCCR enhanced rice by incorporating SNP and environmental data with a

Bi-LSTM module (Ma et al., 2024). Transfer learning, as demonstrated by TrG2P,

employed pre-trained models on non-yield traits to improve yield predictions,

leveraging pleiotropic relationships (Li et al., 2024).

Despite these above advances, multi-task learning in GS remains underexplored. A

multi-task learning based GS strategy that leverages shared information across traits

has the potential to improve generalization and robustness, particularly in data-limited

scenarios (Eraslan et al., 2019). However, challenges such as inadequate modelling of

inter-task correlations, variability in phenotype data quality leading to negative transfer,

and doubled complexity in model training limit its application. The advancement of GS

through the implementation of innovative multi-task architectures and optimization

strategies represents a promising avenue for future research endeavors. Consequently,

there is an urgent necessity to develop innovative multitasking architectures and

optimization strategies to propel further advancements in this domain.

Herein, we report a novel deep learning framework, named as contrastive learning

and chromosome-aware network (CLCNet), which has been specifically tailored for plant GP. The architecture of CLCNet comprises two fundamental modules: a contrastive learning module for capturing phenotypic variations, and a chromosome-aware module that accounts for linkage, as well as both local and global epistasis. The efficacy of CLCNet was evaluated across eight datasets from seven plant species (maize (*Zea mays*), rice (*Oryza sativa*), cotton (*Gossypium hirsutum*), millet (*Setaria italica*), chickpea (*Cicer arietinum*), rapeseed (*Brassica napus*), and soybean (*Glycine max*)). The results indicated that CLCNet outperformed three classical models (rrBLUP, Bayesian Ridge, Bayesian Lasso), two machine learning approaches (LightGBM, SVR), and two deep learning models (DNNGP, DeepGS), achieving higher Pearson correlation coefficients (PCC) and lower root mean squared errors (RMSE). These results underscore the superior accuracy, generalization, and scalability of CLCNet, indicating it as a powerful tool for future GS applications in plant breeding.

Results

1. The architectural design of CLCNet

As shown in Figure 1a, CLCNet is a multi-task model that uses single-nucleotide polymorphism (SNP) data to predict phenotypic values as the primary task, whilst concurrently generating low-dimensional feature representations as an auxiliary task. The model incorporates a dual-pathway backbone, comprising one pathway with three non-linear layers and another with a linear layer, combined through a residual structure. Outputs are processed through separate branches for the primary and auxiliary tasks.

Figure 1. CLCNet model framework. (a) CLCNet model architecture. CLCNet is a multi-task
model using SNP data to predict phenotypic values while generating low-dimensional feature
representations. It features a dual-pathway backbone with non-linear and linear layers
combined via a residual structure, supporting both primary and auxiliary tasks. (b)
Chromosome-aware module. SNPs are grouped by chromosome, with LightGBM used for
feature selection. Local selection identifies key SNPs within chromosomes, while global
selection captures genome-wide key SNPs. Only SNPs with positive information gain (IG > 0)
to LightGBM are retained. The selected local and global SNPs are then concatenated to form
the input features. (c) Comparative learning module. Contrastive learning module employs a
random pairing strategy per epoch, processing two samples simultaneously to predict
phenotypic values and feature vectors. Optimization uses three loss functions: MSE for
phenotypic predictions and a contrastive loss linking feature vector distances to phenotypic
differences.

As demonstrated in Figure 1b, the framework incorporates two innovations. Firstly,

it includes a chromosome-aware module for the preprocessing of SNPs. Secondly, it

incorporates a contrastive learning module for the training process. SNPs are organized

by chromosome, and key loci are selected both locally and globally using a feature

selection algorithm that efficiently identifies important loci. These selected features are

then concatenated to form the model input. The contrastive learning module

incorporates a random pairing strategy during training to optimize the model. This

module minimizes a combined loss function that includes both phenotypic prediction

loss and contrastive loss, leveraging pairwise relationships between samples (Figure

1c).

2. Performance of CLCNet and comparison with other methods

CLCNet demonstrated notable advantages in the domain of phenotypic prediction tasks.

Out of 27 tasks, CLCNet significantly outperformed all other methods in 21 instances.

This superiority was characterized by an enhanced Pearson's correlation coefficient

(PCC) in 21 tasks, as demonstrated in Figure 2a, and a reduction in root mean square

error (RMSE) in 20 tasks, as shown in Figure 2b. Notably, for one phenotype (DTT in

GSTP001), CLCNet showed a significantly higher PCC, while the RMSE difference

was not significant. This indicated that CLCNet's performance for DTT remains robust

despite dataset limitations, as its PCC consistently exceeded 95% across other methods

(Supplementary Figures 1).

Figure 2. CLCNet overall prediction performance. (a) Prediction accuracy performance of CLCNet on eight datasets. (b) Prediction error performance of CLCNet on eight datasets. ΔPCC and ΔRMSE measure the improvement of CLCNet over the best baseline model specific to each task. Significant differences between CLCNet and the best baseline model are indicated by asterisks, and non-significant differences are indicated by ns. Significant level, *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$, ****: $p < 0.0001$.

However, for six phenotypes, CLCNet did not show performance improvements,

with both PCC and RMSE being statistically comparable to the best baseline method.

These phenotypes included fiber elongation (Felo), fiber length (Flen), micronaire

value (FMic), fiber strength (FStr), hundred seed weight (HSW), and grain weight per

plant (PGW), which were distributed across the GSTP010 (cotton) and GSTP011

(millet) datasets (Figure 2a; Figure 2b; Supplementary Figures 4-5). Among the 21 tasks

where CLCNet achieved significant PCC improvements, it outperformed rrBLUP 16

times, SVR 4 times, and Bayesian Lasso once (Supplementary Figures 1-8). These

results underscore the superiority of CLCNet in predictive accuracy and robustness

across diverse phenotypic traits.

CLCNet exhibited outstanding performance in PCC improvements across multiple

datasets (Table 1, Supplementary Table 1). In the GSTP012 (chickpea) dataset, the

ΔPCC_{mean} increase reached 36.015%, with the largest improvement observed in yield

per plant (YPP) at 54.603% (Supplementary Table 1). In GSTP013 (rapeseed), the

ΔPCC_{mean} increase was 31.866%, with protein content (PC) showing the greatest

improvement at 50.389% (Supplementary Table 1). Similarly, in GSTP014 (soybean),

the ΔPCC_{mean} improvement was 16.375%, with protein content (PC) again exhibiting

the largest increase at 25.651% (Supplementary Table 1). In contrast, ΔPCC_{mean} in

GSTP001 (hybrid maize) was relatively small, at only 0.599%. This modest

improvement may be attributed to the performance plateau of the dataset, where the

prediction accuracy for traits such as days to tassel (DTT), ear weight (EW), and plant

height (PH) had already exceeded 95%, 80%, and 94%, respectively, with the other

seven methods (Supplementary Figures1).

Table 1. Benchmark dataset

ID	Species	Sample type	chromosomes	Sample number	Raw SNP	SNP filter	SNP number	Traits	ΔPCC$_{mean}$	ΔRMSE$_{mean}$
GSTP001	Maize	Hybrid Line	2n = 2x = 20	8,652	4,549,828	LD	125,820	EW, DTT, PH	0.599%	0.008
GSTP003	Maize	Inbred Line	2n = 2x = 20	1,458	4,549,828	LD	132,584	EW, DTT, PH	12.897%	-0.114
GSTP007	Rice	Hybrid Line	2n = 2x = 24	1,495	1,651,507	LD	87,363	YPP_HZ, YPP_SY, GW, PN	12.779%	-0.073
GSTP010	Cotton	Inbred Line	2n = 4x = 52	1,245	1,122,352	LD; MAF	14,201	FElo, FLen, FMic, FStr	-0.600%	0.007
GSTP011	Millet	Inbred Line	2n = 2x = 18	827	161,562	LD	20,860	MSW, HSW, PGW	3.387%	-0.028
GSTP012	Chickpea	Inbred Line	2n = 2x = 16	2,921	2,418,866	LD; MAF	289,653	HSW, YPP, SPP	36.015%	-0.243
GSTP013	Rapeseed	Inbred Line	2n = 4x = 38	991	4,286,896	LD	666,221	FT, OC, PC	31.866%	-0.385
GSTP014	Soybean	Inbred Line	2n = 2x = 40	2,795	29,599,017	LD	173,511	SL, VBN, HSW, PC	16.375%	-0.127

The LD and MAF commands are *--indep-pairphase 100 10 0.2* and *--maf 0.05*, respectively. EW, DTT, PH, YPP_HZ, YPP_SY, GW, PN, Felo, Flen, Fmic, FStr, MSW, HSW, PGW, YPP, SPP, FT, OC, PC, SL, and VBN denote ear weight, days to tasseling, plant height, yield per plant in Hangzhou and Sanya, grain weight, panicle number, fiber elongation rate, fiber length, fiber micronaire, fiber strength, main stem width, hundred seed weight, per plant grain weight, yield per plant, seeds per pod, flowering time, oil content, protein content, seed length, and valid branch number, respectively. ΔPCC and ΔRMSE indicate the improvement of CLCNet over the task-specific best baseline model, while ΔPCC mean and ΔRMSE mean represent the average ΔPCC and ΔRMSE across all tasks in one dataset.

Regarding ΔRMSE $_{mean}$, CLCNet performed best in GSTP013 (rapeseed), GSTP012 (chickpea), and GSTP014 (soybean), with reductions of -0.385, -0.243, and -0.127, respectively. Conversely, the smallest ΔRMSE reduction was observed in GSTP001 (hybrid maize), at just -0.008.

Collectively, CLCNet exhibited significant predictive advantages across most phenotypic prediction tasks, underscoring its superior potential value and applicability.

3. SNP analysis of the chromosome-aware module

An analysis was conducted of the characteristics of SNPs in the chromosome-aware module across eight datasets. The following proportions were calculated: (a) the proportion of SNPs specific to the global chromosome, (b) the proportion of SNPs specific to the local chromosome, (c) the proportion of SNPs that overlap, and (d) the proportion of SNPs in quantity. These metrics were defined based on global SNPs and local SNPs. The proportion of global-specific SNPs represents the percentage of global SNPs not included in the local SNP set, relative to the total global SNPs. The proportion

of local-specific SNPs is defined as the percentage of local SNPs not included in the

global SNP set, relative to the total local SNPs. The proportion of overlapping SNPs

represents the percentage of overlapping SNPs in the global and local SNP sets relative

to their union minus the overlap. Finally, the proportion of SNP quantity represents the

incremental count of local SNPs compared to global SNPs, relative to their union minus

the overlap.

Figure 3. Analysis of SNP selected by the chromosome-aware module for each of the eight
datasets. (a) Proportion of global-specific SNP for each of the eight datasets. It calculated as
the percentage of global SNPs not included in the local SNP set relative to the total global SNPs.
(b) Proportion of local-specific SNP for each of the eight datasets. It the percentage of local
SNPs not included in the global SNP set relative to the total local SNPs. (c) Proportion of
overlapping SNP for each of the eight datasets. It calculated as the percentage of shared SNPs
between global and local SNP sets relative to their union minus the overlap. (d) Proportion of
number of SNP for each of the eight datasets. It representing the incremental count of local
SNPs compared to global SNPs relative to their union minus the overlap.

The results (Figures 3a, 3b, 3c and 3d) demonstrated that GSTP010 (cotton) and

GSTP011 (millet) exhibited significant outlier characteristics in all four proportions.

Specifically, they had the lowest proportion of global-specific SNPs, proportion of

local-specific SNPs, and proportion of SNP quantity, but the highest proportion of

overlapping SNPs. Despite differences in SNP counts, sample sizes, and chromosome

numbers across datasets, the outlier features of GSTP010 and GSTP011 aligned with

their SNP counts, which were the lowest among all datasets (Supplementary Figures

9a-c).

Notably, ΔPCC and ΔRMSE for these two datasets showed no significant differences

(Figure 2, Table 1, Supplementary Figures 4 and 5). These findings suggest that the

quantity of SNPs may have a substantial impact on the predictive performance of

CLCNet.

4. Impact of SNP number on CLCNet's predictive performance

The LD pruning and MAF filtering steps were removed from the preprocessing of the

GSTP010 (cotton) and GSTP011 (millet) datasets. Consequently, the number of SNPs

for GSTP010 increased substantially from 14,201 to 1,122,352, and for GSTP011 from

20,860 to 161,562. Subsequent SNP analysis through the chromosome-aware module

revealed increases in the proportions of global-specific SNPs, local-specific SNPs, and

overlapping SNPs, while the proportion of SNP quantity decreased (Supplementary

Figures 9d-g).

Subsequently, a comparison was made of the performance of CLCNet and the other

seven methods on the GSTP010 (cotton) and GSTP011 (millet) datasets after increasing

the SNP count. For GSTP010, among its four traits, three traits that initially showed no

significant difference between CLCNet and the best-performing method demonstrated

significant differences after the SNP count was increased, while fiber length (FElo)

remained non-significant (Supplementary Figures 4 and 10). In GSTP011, the PCC for

hundred seed weight (HSW) remained non-significant in comparison to other methods;

however, all RMSE improved from non-significant to significant. For plant grain

weight (PGW), both PCC and RMSE shifted from non-significant to significant

differences. Additionally, for main stem width (MSW), the difference improved from

significant to highly significant (Supplementary Figures 5 and 11).

Figure 4. Analysis of GSTP10 and GSTP011 before and after increasing the number of SNPs. (a) ΔPCC of GSTP10 and GSTP011 before and after increasing the number of SNPs. (b) ΔRMSE of GSTP10 and GSTP011 before and after increasing the number of SNPs. ΔPCC and ΔRMSE reflect the improvement of CLCNet over the best-performing baseline model for each specific task.

Further analysis revealed that both ΔPCC and ΔRMSE were significantly better after the SNP count increased (Figure 4a; Figure 4b), indicating that the performance improvement of CLCNet surpassed that of other methods. These results demonstrate that increasing SNP count can substantially enhance the predictive performance of CLCNet.

5. Analysis of the Importance of CLCNet's Core Module

To ascertain the impact of the two core modules of CLCNet on the model's performance, a series of modified models were constructed in which the two core modules of CLCNet were removed. The performance of these models was then compared with that of the complete CLCNet model. This comparison was made across 27 phenotypic prediction tasks in terms of PCC and RMSE. Upon the removal of the contrastive learning module, the model transitioned from multi-task learning to single-task learning, and this transition is referred to as CLCNet_1. The third model, named as CLCNet_2, was characterized by the omission of single-chromosome feature selection, whilst retaining feature selection for the entire chromosome group. The architecture of the three models is detailed in Supplementary Figure 12.

Figure 5. Comparison of CLCNet and its variants across 27 phenotype prediction tasks from eight datasets. (a) ΔPCC comparison between CLCNet and its variants. (b) ΔRMSE comparison between CLCNet and its variants. ΔPCC and ΔRMSE represent the performance improvement of CLCNet relative to the best-performing baseline model for each individual task.

The results indicate that, with respect to the ΔPCC, CLCNet demonstrated the optimal performance, with CLCNet_1 and CLCNet_2 exhibiting the lowest predictive accuracy. The ΔPCC for CLCNet_2 decreased to a negative value, indicating that its performance inferior to that of traditional methods (Figure 5a). A parallel tendency was observed in terms of the ΔRMSE (Figure 5b). Specifically, CLCNet demonstrated the lowest error, followed by CLCNet_1, while CLCNet_2 exhibited the highest error, exceeding the baseline, indicating that its error was greater than that of traditional methods (Figure 5b).

The comprehensive analysis of this study indicates that the chromosome-aware module is crucial for improving prediction accuracy and reducing error, and its removal results in a significant decline in model performance. The contrastive learning module validates the effectiveness of multi-task modelling, optimizing errors and enhancing accuracy through weight sharing and loss-driven mechanisms. It is only when both modules are used in combination that the model performs optimally. This emphasizes

the key role of the chromosome-aware module and the contrastive learning module and

their synergistic effect in complex phenotypic prediction tasks.

6. Comparison of algorithm performance under SNP position perturbation

As shown in Figure 1b, the chromosome-aware module employs a two-tiered approach

for the selection of representative SNPs, categorizing them into local and global groups.

Local SNPs are identified through local selection, while global SNPs are obtained

through genome-wide selection. This methodology accounts for both local and global

epistasis, *i.e.* non-linear SNP relationships within chromosome groups and across the

genome. It is important to note that local and global selections are distinct processes;

local-specific SNPs are not identifiable through global selection, and vice versa. The

efficacy of the chromosome-aware module is evidenced by its integration of both local

and global selections, a feature absented in CLCNet_2, which relies solely on global

epistasis selection. This finding serves to confirm the limitations of global-only SNP

selection (Figures 5a and 5b).

Figure 6. Analysis of SNP selected by the chromosome-aware module for each of the eight

datasets before and after position disruption and linkage disequilibrium decay analysis for

each of the eight datasets. (a) Proportion of global-specific SNP during positional shuffling. (b)

Proportion of local-specific SNP during positional shuffling. (c) Proportion of overlapping

SNP during position shuffling. (d) Proportion of numbers of SNP during position shuffling. (e)

Linkage disequilibrium decay analysis.

These proportions were previously defined and calculated based on the global and local SNPs

for each dataset, to compared before and after the disruption of SNP positions.

It is hypothesized that the chromosome-aware module is responsible for the linkage

effects observed within chromosomes, thereby enhancing the model's performance. To

verify this hypothesis, an experiment was conducted in which the positions of SNP were

randomized across the genome. The performance of the model was then compared

before and after the randomization. This disruption of linkage effects introduced more

redundant SNPs, leading to a subsequent decrease in prediction accuracy. The analysis

revealed that in most datasets (GSTP001 for hybrid maize, GSTP003 for inbred maize,

GSTP007 for rice, and GSTP014 for soybean), the shuffle increased the proportion of

both global-specific and local-specific SNPs. Global-specific SNPs increased by an

average of 20%, while local-specific SNPs increased by only 2%. Furthermore,

overlapping SNPs decreased, thereby supporting our hypothesis that linkage dispersion

causes the model to favour global-specific SNPs (Figure 6a-d). It is hypothesised that

GSTP012 (chickpea) and GSTP013 (rapeseed) exhibit smaller linkage effects, a

hypothesis that is confirmed by linkage decay distance calculations showing lower

linkage effects in these datasets (Figure 6e).

Figure 7. Comparison of PCC and RMSE of six models on six datasets before and after

position shuffling.

(a-f) Performance changes in PCC and RMSE on each dataset (GSTP001, GSTP003,

GSTP007, GSTP012, GSTP013, and GSTP014) before and after SNP position shuffling.

For datasets from hybrid maize (GSTP001), inbred maize (GSTP003), rice

(GSTP007), and soybean (GSTP014), following the shuffling of the SNP positions, the

PCC of CLCNet significantly decreased and the RMSE significantly increased. In

contrast, methods such as rrBLUP, Bayesian Ridge, Bayesian Lasso, LightGBM, and

DNNGP demonstrated no substantial alterations in performance, suggesting that they

are unable to account for linkage effects. Furthermore, for the chickpea (GSTP012) and

rapeseed (GSTP013) dataset, CLCNet's PCC and RMSE did not exhibit significant

changes before and after the shuffle, likely attributable to the reduced linkage effects

observed in these species (Figure 6e). Collectively, these findings validate the

hypothesis that the process of shuffling the positions of SNPs leads to the dispersal of

linkage effects, thereby introducing redundant SNPs that result in a substantial decline

in model performance.

Discussion

CLCNet employs a multi-task learning framework, a distinction that sets it apart from

single-task, multi-modal, and transfer learning approaches. In comparison to single-task

models, multi-task models leverage auxiliary tasks' losses to guide the learning process,

often leading to superior performance (Eraslan et al., 2019). In this study, we have

integrated contrastive learning within the multi-task framework of CLCNet. Within this

framework, the primary task is the prediction of phenotypic value, while the prediction

of low-dimensional SNP representation serving as the auxiliary task. The contrastive

loss is calculated by comparing the Euclidean distance between the low-dimensional

SNP representations of two randomly paired samples with the absolute difference in

their phenotypic values. This design has been shown to strengthen inter-task

correlations, ensure diverse information sources for shared learning, as well as facilitate

continuous model weight updates.

To the best of our knowledge, the application of a multitasking framework based on

comparative learning to genomic prediction (GP) models is the first of its kind, although

there is potential for further optimization. One such potential enhancement could be

achieved with the random pairing method, which could be supplemented by exhaustive

pairing. Specifically, in each training iteration, CLCNet randomly pairs a sample with

another from the dataset to compute contrastive loss. While this random pairing method

has proven to be effective, it typically results in a smaller number of unique pairings

(e.g., with 100 samples, fewer than 100 unique pairings was made out of a theoretical

4950 possible combinations). The implementation of exhaustive pairing could enhance

model performance by leveraging a broader diversity of sample combinations. To

balance computational efficiency and performance, we propose increasing the sampling

rate of pairwise combinations moderately.

The chromosome-aware module we designed in this study is an effective feature

selection method. In the context of genomic data, where sample sizes are limited and

feature dimensions are substantial, dimensionality reduction is paramount for achieving

computational efficiency. Feature selection facilitates dimensionality reduction and

noise reduction by identifying representative SNPs. While traditional methods

combining GWAS and GS are useful for feature selection, GWAS typically employs

linear models to assess associations between individual SNPs and phenotypes,

overlooking SNP interactions (Liu et al., 2019). In contrast, LightGBM accounts for

SNP interactions during feature selection, offering greater sensitivity and accuracy in

identifying relevant SNPs (Yan et al., 2021). In our approach, we employ LightGBM to

capture SNP interactions, applying chromosome grouping for local SNPs and genome-

wide selection for global SNPs. The selected local and global SNPs are then combined

to form the final set of representative SNPs. These local and global SNPs complement

each other and cannot replace one another.

In addition, the chromosome-aware module, another distinctive feature of CLCNet,

accounts for linkage effects, a special form of local interaction in genomic data. The

integration of local, global, and linkage effects enhances the model's performance,

setting CLCNet apart from other models. Furthermore, the chromosome-aware module

is flexible and can be conveniently introduced to a variety of GP models, including

linear methods, machine learning approaches, and deep learning frameworks. This

versatility positions the chromosome-aware module as a potent instrument for data

preprocessing across a range of GP tasks.

CLCNet demonstrated no substantial enhancement in performance when compared

to the optimal approach for GSTP10 (cotton) and GSTP11 (millet), primarily due to the

restricted number of SNPs, which was impacted by LD pruning and MAF preprocessing.

Following the elimination of these preprocessing steps, there was a substantial increase

in the SNP number (Table 1). Upon re-evaluation, CLCNet exhibited superior

performance, surpassing the most effective baseline methods (Figure 4a; Figure 4b). A

similar observation was made in majority of tasks that had hitherto exhibited no

improvement, which demonstrated significant gains, while those with significant

improvements exhibited even higher levels of enhancement (Supplementary Figures 4

and 10 for GSTP10; Supplementary Figures 5 and 11 for GSTP11). Notably, this was

accomplished without altering sample sizes, highlighting the strong potential of

CLCNet for cost-effective planting scenarios by increasing sequencing depth.

Conclusion

We developed a novel deep learning model, the CLCNet, for plant GP. The integration

of chromosome-aware and contrastive learning modules has resulted in CLCNet

demonstrating superior performance in terms of prediction accuracy when compared to

traditional linear models, machine learning models, and other deep learning approaches.

The model effectively captures linkage effects, local and global epistasis, and optimizes

molecular marker selection, thereby reducing data noise and improving overall

performance. Furthermore, the multi-task modelling strategy of CLCNet, which

incorporates the prediction of phenotypic difference as an auxiliary task, facilitates

enhanced inter-task information sharing and a consequent increase in prediction

accuracy. In summary, CLCNet is a reliable, efficient, and powerful tool for GS,

excelling in high-dimensional datasets and holding significant potential for plant

breeding applications.

Materials and methods

Data description and preprocessing

This study utilized the CropGS-Hub database, which consolidates genotypic and

phenotypic data from 14 populations of 7 major crops, comprising over 30,000

individuals and 224 billion genotypic records (Chen et al., 2024). From these, 8 datasets

were selected as benchmarks based on sample size, SNP loci, and population type. Data

dimensionality was reduced using PLINK (v1.9) for linkage disequilibrium (LD)

pruning (*--indep-pairphase 100 10 0.2*) and minor allele frequency (MAF) filtering (*--*

maf 0.05 for GSTP012 and GSTP014). The final datasets, representing maize (*Zea*

mays), rice (*Oryza sativa*), cotton (*Gossypium hirsutum*), foxtail millet (*Setaria italica*),

chickpea (*Cicer arietinum*), rapeseed (*Brassica napus*), and soybean (*Glycine max*),

include 27 complex traits. Preprocessed datasets were summarized in Table 1.

The architecture of GenomicCNN

We developed CLCNet using PyTorch (v2.1.0, https://pytorch.org/). CLCNet is a multi-

task deep learning model that takes genomic single-nucleotide polymorphism (SNP)

data as input and outputs phenotypic predictions (primary task) alongside low-

dimensional feature vectors (auxiliary task). The input receives an n × m matrix,

where n represents the sample number and m represents the number of markers. The

0/1/2 genotype encoding was transformed into one-hot encoding. The model's backbone

network employs a dual-pathway architecture: one pathway comprises three non-linear

fully connected layers (ReLU activation; dimensions: 4096→2048→1024), while the

other consists of a single linear fully connected layer (dimension: 1024). The outputs

of the two pathways are summed at the 1024-dimensional level and subsequently passed

to the primary and auxiliary task branches. Each branch is composed of three non-linear

fully connected layers, with dimensions mapped as 4096→2048→1 for the primary

task and 4096→2048→1024 for the auxiliary task.

The chromosome-aware module is essential for preprocessing genomic data. SNPs are

grouped into chromosomal categories based on their genomic locations, followed by

local feature selection within each group to identify key SNPs (local SNPs). Global

feature selection is then applied across the entire genome to capture comprehensive

features (global SNPs). Local and global SNPs are concatenated to form the input

features for the model. Feature selection is performed using the LightGBM algorithm,

a tree-based approach to model non-linear relationships among SNPs, facilitating the

identification of key loci (Yan et al., 2021). It identifies key loci by retaining SNPs with

positive information gain (IG > 0) and discarding those with IG = 0.

The contrastive learning module introduces a customized training strategy. During each

epoch, when loading each sample, a random pairing mechanism selects another sample

for pairing. Both samples are fed into the model simultaneously, and their phenotypic

values and low-dimensional feature vectors are predicted. Model optimization is guided

by three loss functions: the mean squared error (MSE) between the predicted and true

phenotypic values for each of the two input samples (equation 1), and a custom

contrastive loss based on the Euclidean distance between the feature vectors of the two

samples and the difference in their phenotypic values (equation 2). The model aims to

minimize the total loss, which is the sum of the MSE for both samples and the

contrastive loss. The server was equipped with two Xeon Xeon(R) Gold 6330 CPU,

512 GB of memory, and an NVIDIA GeForce RTX3090 GPU.

equation 1 $$Loss_{mse} = \frac{1}{N}\sum_{i=1}^{N}(y_i - \hat{y}_i)^2$$

Here, N is the number of samples. y_i is the true value for the i-th observation. $\hat{y}_i$ is

the predicted value for the i-th observation.

equation 2 $$Loss_{contrastive} = \frac{1}{N}\sum_{i=1}^{N}(\|embedding_1^i - embedding_2^i\| - diff^i)^2$$

Here, N is the number of samples. $\|embedding_1^i - embedding_2^i\|$ denotes the

Euclidean distance between the i-th pair of embeddings. $diff^i$ is the phenotypic

difference for the i-th pair of samples.

Model comparison and cross-validation

The predictive performance of CLCNet was evaluated against seven baseline models

on benchmark datasets. These included three classical statistical methods (Ridge

Regression Best Linear Unbiased Prediction, rrBLUP; Bayesian Ridge; Bayesian

Lasso), two machine learning models (Light Gradient Boosting Machine, LightGBM;

Support Vector Regression, SVR), and two deep learning models (Deep Neural

Network Genomic Prediction, DNNGP; Deep Learning Genomic Selection, DeepGS).

Classical methods were implemented using the R packages rrBLUP (v4.6.2) and BGLR

(v1.1.3). For machine learning, LightGBM was implemented with Python's lightgbm

package (v3.3.5), while SVR utilized Python's scikit-learn library (v0.24.2).

DeepGS utilized a convolutional neural network (CNN) architecture comprising input,

convolutional, max-pooling, dropout, fully connected, and output layers, with dropout

rates of 0.2, 0.1, and 0.05 applied at various stages to mitigate overfitting. The ReLU

activation function was employed throughout the architecture (Ma et al., 2018).

Similarly, DNNGP adopted a featuring three convolutional layers, batch normalization,

dropout, a flattening layer, a dense layer, and an output layer. Principal components

accounting for over 98% of the variance were extracted from the genomic data using

PLINK (v1.9) as input features (Wang et al., 2023).

Predictive accuracy was measured using the Pearson Correlation Coefficient (PCC),

and prediction error was assessed using the Root Mean Squared Error (RMSE). To

compare CLCNet with the best-performing baseline models, we calculated the PCC

$_{CLCNet}$ and PCC $_{Best}$ (PCC values for CLCNet and the best baseline, respectively),

RMSE $_{CLCNet}$ and RMSE $_{Best}$ (RMSE values for CLCNet and the best baseline,

respectively), and ΔPCC and ΔRMSE (differences in PCC and RMSE between CLCNet

and the best baseline).

The significance of performance differences was assessed using paired t-tests under a

10-fold cross-validation. Phenotype values of test set were z-normalized using the mean

and variance from the training set to prevent data leakage. All the models were trained

on the training set, and their performance was evaluated on the test set.

SNP position perturbation

The input is an n ×m matrix, where n represents the number of samples and m denotes

the number of markers, as previously described and defined. The order of m reflected

the genomic positions of the markers. To perform SNP position perturbation, the order

of m was randomly shuffled across all n samples, disrupting the original SNP positions.

After shuffling, the input matrix shape remains unchanged. Subsequently, CLCNet and

other methods were used to calculate the changes in PCC and RMSE before and after

the SNP position shuffle.

Linkage disequilibrium decay analysis

SNPs with a minor allele frequency (MAF) < 0.05 and a missing rate > 10% were

filtered using VCFtools (v0.1.15). Pairwise linkage disequilibrium (LD) was calculated

as r² for the remaining SNPs within a 1 MB window.

Acknowledgements

We would like to express our gratitude to Dr. Jiachao Xu, a postdoctoral researcher

from the Institute of Genetics and Developmental Biology of the Chinese Academy of

Sciences, for his professional guidance on server construction. We also appreciate Prof.

Fei Lu (the Institute of Genetics and Developmental Biology of the Chinese Academy

of Sciences) for the insightful discussions. We thanks to Dr. Shihao Shao from Peking

University Health Science Center for his valuable teaching and constructive

suggestions on deep learning.

Code availability

The scripts are available on GitHub (https://github.com/SuppurNewer/CLCNet).

Conflict of interest

The authors declare no conflicts of interest.

Authors' contributions

Y.J., and R.H. conceived the study and supervised the research; J.H. developed the

model and drafted the manuscript; J.H., Z. Y., M.Y., C.L., and J.L analyzed data. R.H.

Y.W., L.W., F.H., C.L., and M.L. discussed the results and revised the manuscript. All

authors read and approved the final manuscript.

Supplemental information

Supplemental Figure 1

Performance comparison of eight models on the GSTP001 dataset

Supplemental Figure 2

Performance comparison of eight models on the GSTP003 dataset

Supplemental Figure 3

Performance comparison of eight models on the GSTP007 dataset

Supplemental Figure 4

Performance comparison of eight models on the GSTP010 dataset

Supplemental Figure 5

Performance comparison of eight models on the GSTP011 dataset

Supplemental Figure 6

Performance comparison of eight models on the GSTP012 dataset

Supplemental Figure 7

Performance comparison of eight models on the GSTP013 dataset

Supplemental Figure 8

Performance comparison of eight models on the GSTP014 dataset

Supplemental Figure 9

he relationship between dataset characteristics and SNPs selected by the chromosome-

aware module

Supplemental Figure 10

Performance comparison of eight models on the GSTP010 dataset after increasing the

number of SNPs

Supplemental Figure 11

Performance comparison of eight models on the GSTP011 dataset after increasing

number of SNPs

Supplemental Figure 12

The CLCNet and its variant model architectures

Supplemental Table S1

A comparison of the results obtained by the CLCNet in relation to 27 trait predictions,

with the results of the most effective method among the other methods.

References

Abdollahi-Arpanahi, R., Gianola, D., and Pen˜ agaricano, F. (2020).Deep learning
versus parametric and ensemble methods for genomic prediction of complex
phenotypes. Genet. Sel. Evol. 52:1-15.

Crossa, J., de los Campos, G., Pérez, P., Gianola, D., Burgueño, J., Araus, J. L.,
Makumbi, D., Singh, R. P., Dreisigacker, S., Yan, J., et al. (2010). Prediction of
genetic values of quantitative traits in plant breeding using pedigree and molecular
markers. Genetics, 186(2), 713-724.

Crossa, J., Pérez-Rodríguez, P., Cuevas, J., Montesinos-López, O., Jarquín, D., de los
Campos, G., Burgueño, J., González-Camacho, J. M., Pérez-Elizalde, S., Beyene,
Y., Dreisigacker, S., Singh, R., Zhang, X., Gowda, M., Roorkiwal, M., Rutkoski,
J., and Varshney, R. K. (2017). Genomic selection in plant breeding: Methods,
models, and perspectives. Trends Plant Sci. 22: 961-975.

Eraslan, G., Avsec, Ž., Gagneur, J., and Theis, F.J. (2019). Deep learning: new
computational modelling techniques for genomics. Nat. Rev. Genet. 20, 389-403.

Gao, P., Zhao, H., Luo, Z., Lin, Y., Feng, W., Li, Y., Kong, F., Li, X., Fang, C., and
Wang, X. (2023). SoyDNGP: A web-accessible deep learning framework for
genomic prediction in soybean breeding. Brief. Bioinform. 24(6): 1-12.

Habier, D., Fernando, R. L., and Dekkers, J. C. M. (2007). The impact of genetic
relationship information on genome-assisted breeding values. Genetics, 177(4),
2389-2397.

Heslot, N., Yang, H.-P., Sorrells, M. E., and Jannink, J.-L. (2012). Genomic selection
in plant breeding: A comparison of models. Crop Science, 52(1), 146-160.

Li, J., Zhang, D., Yang, F., Yang, J., Wang, K., and Zhao, C. (2024). TrG2P: A transfer-
learning-based tool integrating multi-trait data for accurate prediction of crop yield.
Plant Commun. 5: 100975.

Liu, H. J., and Yan, J. (2019). Crop genome-wide association study: A harvest of
biological relevance. Plant J. 97: 8-18.

Liu, J., Li, J., Wang, H., and Yan, J. (2020). Application of deep learning in genomics.
Science China Life Sciences, 63, 1860-1878.

Liu, Y., Wang, D., He, F., Wang, J., Joshi, T., and Xu, D. (2019). Phenotype prediction
and genome-wide association study using deep convolutional neural network of
soybean. Front. Genet. 10: 1091.

Ma, W., Qiu, Z., Song, J., Li, J., Cheng, Q., Zhai, J., and Ma, C. (2018). A deep
convolutional neural network approach for predicting phenotypes from genotypes.
Planta 248: 1307-1318.

Ma, X., Wang, H., Wu, S., Han, B., Cui, D., Liu, J., Zhang, Q., Xia, X., Song, P., Tang,
C., Geng, L., Yang, Y., Yan, S., Zhou, K., and Han, L. (2024). DeepCCR: Large-
scale genomics-based deep learning method for improving rice breeding. Plant
Biotechnol. J. 1-3.

Maenhout, S., De Baets, B., Haesaert, G., and Van Bockstaele, E. (2007). Support
vector machine regression for the prediction of maize hybrid performance.
Theoretical and Applied Genetics, 115(7), 1003–1013.

Meuwissen, T.H.E., Hayes, B.J., and Goddard, M.E. (2001). Prediction of Total Genetic
Value Using Genome-Wide Dense Marker Maps. Genetics 157:1819-1829.

Ren, Y., Wu, C., Zhou, H., Hu, X., and Miao, Z. (2024). Dual-extraction modeling: A
multi-modal deep-learning architecture for phenotypic prediction and functional
gene mining of complex traits. Plant Commun. 5: 100102.

Wang, H., Cimen, E., Singh, N., and Buckler, E. (2020). Deep learning for plant
genomics and crop improvement. Curr. Opin. Plant Biol. 54: 34–41.

Wang, K., Abid, M.A., Rasheed, A., Crossa, J., Hearne, S., and Li, H. (2023). DNNGP,

a deep neural network-based method for genomic prediction using multi-omics

data in plants. Mol. Plant 16: 279-293.

Wang, X., Xu, Y., Hu, Z., and Xu, C. (2018). Genomic selection methods for crop

improvement: Current status and prospects. Crop J. 6: 330-340.

Yan, J., Xu, Y., Cheng, Q., Jiang, S., Wang, Q., Xiao, Y., Ma, C., Yan, J., and Wang, X.

(2021). LightGBM: Accelerated genomically designed crop breeding through

ensemble learning. Genome Biology, 22, 271.

625